1

Prefrontal neuronal dynamics in the absence of task execution

Shusen Pu[1,2+], Wenhao Dang[1+], Xue-Lian Qi[3], Christos Constantinidis[1,4,5]

1. Department of Biomedical Engineering, Vanderbilt University, Nashville TN 37235, USA.
2. Department of Mathematics and Statistics, University of West Florida, Pensacola, FL 32514, USA.
3. Department of Neurobiology & Anatomy, Wake Forest School of Medicine, Winston-Salem, NC 27157, USA.
4. Neuroscience Program, Vanderbilt University, Nashville TN 37235, USA.
5. Department of Ophthalmology and Visual Sciences, Vanderbilt University Medical Center, Nashville TN 37232, USA.

+These authors contributed equally to this work

Keywords: monkey; population coding; prefrontal cortex; working memory

Lead Contact / Corresponding Author:
Christos.Constantinidis.1@vanderbilt.edu

SUMMARY

Activity across prefrontal cortical populations represents stimuli in a low-dimensional

manifold that often undergoes transformations during the execution of cognitive tasks.

These dynamics are commonly thought to be determined by whatever cognitive operation

is being performed at the time, for example, rotation of stimulus representation in

population activity during working memory is thought to reduce interference by

subsequent distractors. The assumption that cognitive operations dictate stimulus

transformations has not yet been tested experimentally. Here we show that the activity

space of prefrontal neurons in naïve monkeys exhibits significant rotations, some aspects

of which remain remarkably unchanged after training to perform working memory tasks.

Training does induce changes in population dynamics, which further distinguish correct

and error trials during task execution. Our results reveal automatic functions of prefrontal

neural circuits that form the substrate of cognitive flexibility.

INTRODUCTION

Prefrontal neurons represent sensory stimuli in a dynamic, task-specific manner [1, 2, 3]. Analysis of population activity with methods of dimensionality reduction reveals that stimuli can be represented in a low dimensional space, or manifold, with the majority of the firing rate variance captured by a few dimensions [4, 5]. Furthermore, the representation of stimuli in the reduced space is not static but changes dynamically during the course of a trial, as the subjects perform a cognitive operation in the context of a task [6, 7, 8]. The stimulus manifold may therefore be projected to a different subspace, rotated, or otherwise geometrically transformed [6, 7, 8].

Transformations of stimuli in state space have been thought to be specific for the cognitive operations being performed. Orthogonal rotation of a stimulus representation has been proposed as a possible mechanism for reducing interference between sensory and memory representations, protecting the memory of an initial stimulus from the interference of a subsequent stimulus presentation [9]. The same stimuli are represented in different subspaces when used in the context of different tasks [10] and errors are characterized by changes in stimulus representation geometry [11].

This evidence appears to link the geometric transformation of stimulus information in neuronal populations to cognitive operations [12], however, a critical assumption that task execution is causal to such transformations has not been tested experimentally. It is thus unknown how learning to perform a task in the first place changes the representation of stimuli. Here we test this idea by analyzing population of prefrontal neurons in subjects before and after they were trained to perform cognitive tasks involving identical stimuli. We checked multiple aspects of the activity space,

including subspace alignment, geometrical similarity, and dynamics, to determine the

relative contribution of training and regional specificity in the formation of the observed

low dimensional geometry.

RESULTS

Neurophysiological recordings were collected from the lateral PFC of six monkeys

before and after they were trained to perform working memory tasks [13, 14]. Once fully

trained, the monkeys viewed two stimuli appearing in sequence with intervening delay

periods between them and reported whether the second stimulus was the same as the first

(Fig. 1A-B). The stimulus sets used in these experiments varied in terms of spatial

location or shape (Fig. 1A-B). Recordings were also obtained from these animals,

viewing the same stimuli presented with the same timing prior to any training in the task.

A total of 1164 neurons from six monkeys in five prefrontal subdivisions (Fig. 1C) were

recorded during the passive, pre-training viewing of spatial stimuli; 847 neurons were

recorded during the passive, pre-training viewing of feature stimuli. Additionally, 1031

neurons from three monkeys were obtained after training, when the animals performed

actively the spatial working memory task; 793 neurons were obtained during the active

feature working memory task.

Spatial stimuli are geometrically arranged in a low dimensional subspace

We applied Principal Component Analysis (PCA) on the covariance matrix of the zero-

centered data, and selected eigenvectors sorted in decreasing order in terms of explained

variance. In agreement with previous studies that have suggested that prefrontal neurons

represent stimuli in a low-dimensional manifold [4, 5], we found that the first three

principal components generally explain more than 50% of the variance (Fig. S1) across

brain regions and training phases, validating the approach of visualizing and measuring

geometry structure in low dimensional space.

Previous studies also revealed that location information is arranged in a well-

structured geometry resembling the physical appearance of visual stimuli in working

memory [6, 15], however, it is not clear if this representation is learned through task training.

Visualizing in the subspace defined by the first two principal components confirmed that

spatial stimuli roughly fell on a ring with the same relative positioning in the low-

dimensional subspace (Fig. 2A), and this geometry did not depend on training to perform

a task. We calculated the unexplained variance between the empirical geometry and the

'standard' geometry that maximizes the distance between diametric locations, and found

that spatial representations were well fitted both before and after training using this

approach (Fig. 2B). Across prefrontal subdivisions, the mid-dorsal region had the best

match for the hypothetical arrangement, indicated by the overall lowest unexplained

variance.

Some subspace transformations are task independent

Low-dimensional stimulus representations in neuronal population activity are

transformed during cognitive operations [7, 8, 9] and we therefore sought to test how such

transformations differ systematically before and after training to perform a cognitive task.

We quantified transformations based on the rotation of subspaces in different task epochs

or task conditions.

Orthogonality of stimulus representations at different task intervals has been

described previously in the context of behavioral tasks [9], and we sought to compare how

training affects such transformations in prefrontal areas. Strikingly, representations of the

same stimuli during the cue and match intervals already exhibited significant angles of

rotation in multiple brain areas for both the spatial and feature sets before any task

training (Fig. 3A-B). We also found that the pattern of transformation between the cue

and match stimuli was highly area specific, that is, similar angles of plane rotation were

seen across different prefrontal areas before and after training (Fig. 3B). The Pearson

correlation coefficient between pre- and post-training rotation angles was r=0.78

(p=0.023, Fig. S2 left). To further confirm the subspace orthogonality observed in the

angle measure, we calculated the variance accounted for (VAF) ratio between the cue and

the match period. The results qualitatively agree with the angle measure (Fig. 3C), in that

high angle measurements always correspond to low VAF. Once again, we found that the

relative order of values across subdivisions matched between the pre- and post-training

phases. This result indicates that populations of neurons in different subregions of the

prefrontal cortex transform matching stimuli in a stereotypical fashion, independent of

training and task execution.

Some representation transformations were only observed after training, for

example involving the representation of stimuli in match and nonmatch conditions. In the

spatial task, and specifically in the most selective, mid-dorsal region of PFC[16], the angle

of rotation between the match and nonmatch representations (both during the second

stimulus presentation period) changed considerably after training (Fig. 4A). Across areas,

there was little correlation between the plane angles observed before and after training

across areas, and the correlation coefficient was r=-0.29 (p=0.491, Fig. S2 middle). The

VAF ratio measurement agreed with angle measurement in this case too, evidenced by a

major left shift of the distribution for the mid-dorsal region in the spatial task. A similar

effect of training was observed for the subspace angle between the cue and the first delay

period, with anterior subregions of the PFC exhibiting increased rotation for the spatial

task, and posterior subdivisions exhibiting increased rotation for the feature task (Fig.

S3).

Task dependent and independent representation geometry

In addition to subspace transformations, we also examined the within-subspace geometry.

Two indexes were calculated to represent relative representational amplitude and

structure similarity between task epochs or conditions: the unexplained variance, which

quantified geometrical structure similarity after controlling the scale and orientation

differences of representation in respective subspaces, and the scale factor, which was the

ratio of within-subspace to total variance. We additionally used a within epoch measure

as control (i.e. comparing the geometry of data from the same dataset to itself, see

Methods). For the unexplained variance, the more similar the two conditions are

compared to each other, the closer the measurement should be to 0; for the scale factor,

more similar measurements correspond to values close to 1.

In the geometry similarity analysis, we found that the effect of training for the

most selective subdivision, the mid-dorsal PFC, was to decrease the similarity between

epochs. This was true for the comparison of the two stimulus presentation epochs (cue vs

match): the spatial unexplained variance increased from a pre-training value of

0.097±0.012, to a post-training value of 0.25±0.01; the feature pre-training unexplained

variance similarly increased from 0.11±0.003, to a post-training value of 0.24±0.014

(Fig. 3C). The same relationship held between the cue stimulus and first delay period

epochs: the spatial pre-training unexplained variance increased from 0.1±0.003 to a post-

training value of 0.22±0.007 (Fig. S3C).

The representation scale measured across the cue and sample epochs aggregated

around 1, both before and after training (Fig. 3D and Fig. 4C). This means that the

stimulus representation generally spanned roughly the same amount of area in the

respective subspace, either when compared between the cue and sample (first and second

stimulus presentation) period, or between match and nonmatch conditions within the

same period, thus indicating a stable magnitude of population responses. For the cue vs.

delay comparison, the scale factor measure fell below 1 (Fig. S3C) due in part to the fact

that, even with persistent delay activity, the overall average firing rate in the delay is

lower compared to the stimulus presentation epochs. The stability of the scale factor

during the stimulus presentation periods suggests that on the population level, changes in

the correlation structure induced by training did not result in an overall increase of

stimulus-driven firing rate changes; whereas the amplitude of stimulus representation was

held fairly constant.

Subspace rotation and dynamics correlates with behavior performance

We also compared the dynamic trajectory of population activity before and after training.

As expected, pre-training activity was largely confined to a narrow subspace during the

delay period, while post-training activity became increasingly dynamic in an expanded

space (Fig. S4A, B). To further quantify this change, we randomly sampled half of the

trials from all periods and calculated its PCA space as the base to project the remaining

half trials (see Methods). The mean ratios of the decoding space area between fixation,

first delay and second delay period (using the fixation interval as reference) were 1 : 1.07

: 1.26 for the pre-training data and 1 : 8.62 : 1.83 for the post-training data (Fig. S4C-E).

We also sought to test whether the post-training population rotation in the match

relative to the nonmatch condition was linked to improved task performance. We thus

compared the subspace rotation angle in correct and error trials in a subset of neurons

with sufficient error trials across conditions (n=295 in the spatial working memory task,

and n=201 in the feature working memory task), pooling data across all areas (see

Methods). In correct trials (Fig. 5A, C), match and nonmatch trials were nearly

orthogonal to each other (79.9±5.4 and 78.0±14.5 degrees for the spatial and feature

stimuli, respectively), consistent with the post-training findings of the full population

(Fig. 3). In error trials, this rotation was significantly reduced (Fig. 5B, D,19.7±9.0 and

18.8±14.5 respectively). To determine how population responses evolved across the

length of the trial, we investigated the dynamics of representations for match and

nonmatch trials in the reduced PCA space. Example trajectories for a single stimulus

condition are plotted for the mid-dorsal area in Fig. 5E. These, too, resembled the

trajectories in state space described previously for neuronal activity in animals trained to

perform behavioral tasks involving perceptual decisions [8]. In correct trials, as expected,

match and nonmatch trajectories stayed close to each other until the sample period and

then diverged during the sample presentation and the delay period that followed it. In

error trials, an abnormal rotation was already present in the cue period and match and

nonmatch trajectories were less-distinguishable (Fig. S5). The result suggests that

incomplete rotation of the stimulus subspace is more likely to result in errors. However,

this failure of population activity to follow the desired trajectory is present early in the

trial and does not depend specifically on orthogonalization of the nonmatch stimulus,

which, as we showed here, was present even in naïve animals.

DISCUSSION

Computation in neural activity is characterized by the transformation of representations

across stages of processing [12]. Our results demonstrate that neuronal populations perform

stimulus transformations and exhibit dynamics that are currently thought to represent

cognitive operations [5, 8, 9, 10], even in the absence of task execution, in animals naïve to

any such training. Our results in the same dataset from trained animals corroborated

previous findings: sequential cue and match stimuli from the same trials were represented

in orthogonal planes; match and nonmatch stimuli exhibited large state space separations;

and error trials exhibited smaller plane rotations than correct ones. These positive results

only accentuate the finding that naïve animals, viewing passively the same stimuli,

produced similar rotations as trained animals for the cue vs. match comparison. The

result indicates that some dynamic transformations of population-based stimulus

representations are not caused by task operations requiring explicit training.

Neural basis of cognitive flexibility

Natural plasticity allows for significant training-induced improvements of working

memory, particularly through changes in PFC activity [17, 18, 19, 20, 21]. Representation of

stimuli in neural activity has also been shown to be dynamic at much faster timescales,

during the time course of a single behavioral session [22] or, depending on trial events, on a

moment-by-moment basis [1]. The transformation of stimulus representations has

previously been considered indicative of flexible task execution, e.g. rotating

representations may minimize interference between multiple stimuli, thus allowing

additional stimuli to be encoded with greater efficiency [9]. Our current results suggest that

the prefrontal cortex may perform these operations automatically, even in the absence of

task training or execution. Furthermore, the scale of stimulus representation was largely

unchanged after training. The idea of stable population information under shifting neural

responses within the range of days to weeks is an increasingly appreciated phenomenon

in studies involving post-learning plasiticity[23, 24]. Our results here showed a similar

phenomenon after learning a new task.

Regardless of transformations, data across prefrontal areas and task conditions

were generally well fit by low dimensional representations. Low dimensionality in

working memory representations is considered a hallmark of generalizability, while high

dimensionality is correlated with better discriminability [25]. In our experiment, stimuli

were highly discriminable from each other from the outset, and task training involved

incorporating them into the context of a new task rather than learning fine features of the

stimuli [26], suggesting that an increase in dimensionality was not required to perform the

task after training.

Regional Specialization

Prior studies have debated whether different types of information are represented across

the dorsal-ventral and anterior-posterior axes of the prefrontal cortex has been

contentious [27, 28, 29, 30]. Converging evidence, however, suggests that anterior subdivisions

of the prefrontal cortex generally represent more abstract information, and their activation

depends to a greater extent on the task subjects have been trained to perform [31].

Consistent with this idea, we observed that anterior areas consistently exhibited high

rotation values for match and nonmatch stimuli relative to the cue (and relative to each

other), both before and after training. Our results raise the possibility that this innate

stimulus transformation endows anterior areas of the prefrontal cortex with greater

capacity for plasticity.

METHODS

Data obtained from six male rhesus monkeys (*Macaca mulatta*), ages 5–9 years old, weighing 5–12 kg, as previously documented [14], were analyzed in this study. None of these animals had any prior experimentation experience at the onset of our study. Monkeys were either single-housed or pair-housed in communal rooms with sensory interactions with other monkeys. All experimental procedures followed guidelines set by the U.S. Public Health Service Policy on Humane Care and Use of Laboratory Animals and the National Research Council's Guide for the Care and Use of Laboratory Animals and were reviewed and approved by the Wake Forest University Institutional Animal Care and Use Committee.

Monkeys sat with their heads fixed in a primate chair while viewing a monitor positioned 68 cm away from their eyes with dim ambient illumination and were required to fixate on a 0.2° white square appearing in the center of the screen. During each trial, the animals were required to maintain fixation on a 0.2° white square appearing in the center of the screen while visual stimuli were presented either at a peripheral location or over the fovea, in order to receive a liquid reward (typically fruit juice). Any break of fixation immediately terminated the trial and no reward was given. Eye position was monitored throughout the trial using a non-invasive, infrared eye position scanning system (model RK-716; ISCAN, Burlington, MA). The system achieved a $< 0.3°$ resolution around the center of vision. Eye position was sampled at 240 Hz, digitized and recorded. The visual stimulus display, monitoring of eye position, and synchronization of stimuli with neurophysiological data was performed with in-house software implemented

in the MATLAB environment (Mathworks, Natick, MA), utilizing the Psychophysics

Toolbox [32].

Pre-training task

Following a brief period of fixation training and acclimation to the stimuli, monkeys were

required to fixate on a center position while stimuli were displayed on the screen. The

stimuli shown in the pre-training, passive, spatial task consisted of white 2° squares,

presented in one of nine possible locations arranged in a 3×3 grid with 10° of distance

between adjacent stimuli. Only the eight peripheral locations are analyzed here, as the

center location never appeared as a nonmatch. The stimuli shown in the pre-training

passive feature task consisted of white 2° geometric shapes drawn from a set comprising

a circle, a diamond, the letter H, the hashtag symbol, the plus sign, a square, a triangle,

and an inverted Y-letter. These stimuli could be presented in one of the possible locations

of the spatial grid.

288   The presentation began with a fixation interval of 1 s where only the fixation

point was displayed, followed by 500 ms of stimulus presentation (referred to hereafter as

cue), followed by a 1.5 s "delay" interval where, again, only the fixation point was

displayed. A second stimulus (referred to hereafter as sample) was subsequently shown

for 500 ms. In the spatial task, this second stimulus would be either identical in location

to the initial stimulus, or diametrically opposite to the first stimulus. In the feature task,

this second stimulus would appear in the same location as the initial stimulus and would

either be an identical shape or the corresponding non-match shape (each shape was paired

with one non-match shape). Only one nonmatch stimulus was paired with each cue, so

that the number of match and nonmatch trials were balanced in each set. In both the

spatial and feature task, this second stimulus display was followed by another "delay"

period of 1.5 s where only the fixation point was displayed. The location and identity of

stimuli were of no behavioral relevance to the monkeys during the "pre-training" phase,

as fixation was the only necessary action for obtaining rewards.

Post-training task

Three monkeys, data of which are analyzed here, were trained to perform active working

memory tasks that involved the presentation of identical stimuli as the spatial and feature

tasks during the "pre-training" phase. Monkeys were required to remember the spatial

location and/or shape of the first presented stimulus, and report whether the second

stimulus was identical to the first or not, via saccading to one of two target stimuli (green

for matching stimuli, blue for non-matching). Each target stimulus could appear at one of

two locations orthogonal to the cue/sample stimuli, pseudo-randomized in each trial. In

the spatial task, the second stimulus could be either identical in location to the initial

stimulus, or diametrically opposite to it. In the feature task, this second stimulus would

appear in the same location as the initial stimulus and would either be an identical shape

or the corresponding non-match shape.

Surgery and neurophysiology

A 20 mm diameter craniotomy was performed over the PFC and a recording cylinder was

implanted over the site. The location of the cylinder was visualized through anatomical

magnetic resonance imaging (MRI) and stereotaxic coordinates post-surgery. For two of

the four monkeys that were trained to complete active spatial and feature working

memory tasks, the recording cylinder was moved after an initial round of recordings in

the post-training phase to sample an additional surface of the PFC.

Anatomical localization

Each monkey underwent an MRI scan prior to neurophysiological recordings. Electrode

penetrations were mapped onto the cortical surface. We identified six lateral PFC

regions: a posterior-dorsal region that included area 8A, a mid-dorsal region that included

areas 8B and 9/ 46, an anterior-dorsal region that included area 9 and area 46, a posterior-

ventral region that included area 45, an anterior-ventral region that included area 47/12,

and a frontopolar region that included area 10. However, the frontopolar region was not

sampled sufficiently to be included in the present analyses.

Neuronal recordings

Neural recordings were carried out in the aforementioned areas of the PFC both before

and after training in each WM task. Extracellular recordings were performed with

multiple microelectrodes that were either glass- or epoxylite-coated tungsten, with a 250

μm diameter and 1–4 MΩ impedance at 1 kHz (Alpha-Omega Engineering, Nazareth,

Israel). A Microdrive system (EPS drive, Alpha- Omega Engineering) advanced arrays of

up to 8 microelectrodes, spaced 0.2–1.5 mm apart, through the dura and into the PFC.

The signal from each electrode was amplified and band-pass filtered between 500 Hz and

8 kHz while being recorded with a modular data acquisition system (APM system, FHC,

Bowdoin, ME). Waveforms that exceeded a user-defined threshold were sampled at 25 μs

resolution, digitized, and stored for offline analysis. Neurons were sampled in an

unbiased fashion, collecting data from all units isolated from our electrodes, with no

regard to the response properties of the isolated neurons. A semi-automated cluster

analysis relied on the KlustaKwik algorithm, which applied principal component analysis

of the waveforms to sort recorded spike waveforms into separate units. To ensure a stable

firing rate in the analyzed recordings, we identified recordings in which a significant

effect of trial sequence was evident at the baseline firing rate (ANOVA, $p < 0.05$), e.g.,

due to a neuron disappearing or appearing during a run, as we were collecting data from

multiple electrodes. Data from these sessions were truncated so that analysis was only

performed on a range of trials with stable firing rates. Less than 10% of neurons were

corrected in this way. Identical data collection procedures, recording equipment, and

spike sorting algorithms were used before and after training in order to prevent any

analytical confounds.

Dimensionality reduction and rotation of subspaces

We applied principal components analysis (PCA) to visualize the neural population

activity manifolds during the spatial and feature tasks. PCA was performed on the mean

firing rate of neurons across different prefrontal regions, for data collected both before

and after training. We examined the submanifolds of neural activity as a function of the

spatial and feature stimulus sets.

In the spatial task, we collected data from five regions during five epochs

(fixation, cue, delay1, sample, delay2) before and after the monkeys were trained to

perform the task. In each region, trials were collected when the cue stimuli appeared at

$L = 8$ locations. We then selected the recordings with more than 16 trials at each

location. For each neuron, the average firing rate of the match trials (and nonmatch

trials) during a given period at the eight locations formed a column entry (8x1 vector) for

the population activity matrix A_1 (A_2 for nonmatch trials). During each phase of the

experiment (fixation, cue, delay1, sample, delay2) we defined the corresponding activity

matrix A_1 (A_2) for each prefrontal region as an $8 \times N$ matrix, where 8 is the number of

locations where the cue occurred and N is the number of neurons. To find the rotation

between two periods, we aligned the activity matrix into one single matrix B, which is a

$16 \times N$ matrix with the first eight rows containing the activities in one period and the rest

containing for another period. Then we normalized B by subtracting the mean across

each column to guarantee the matrix was zero-centered. The PCA was applied on the

centered data using singular value decomposition (SVD). We selected the first three

eigenvectors of the covariance matrix of the zero-centered data and sorted them by

decreasing order in terms of explaining the variance. The first three principal components

explained an average of 73% of the response matrix variance across all examined periods

and locations. Similar procedures were applied to visualize the responses in the feature

task, where we grouped the data according to the shape of cue stimulus. Specifically,

there were eight different shapes in the feature task. Therefore, the corresponding activity

matrix had a dimension of $8 \times N$, in which N is the number of neurons and each row

stands for a shape in the feature task (see Figure1 B). For each pair of comparison for the

rotations, we aligned the data in the same way as described above.

To visualize the population representations, we projected the centered activity

matrix into a three-dimensional (3D) PCA space. After projection, the population

representations of the eight spatial locations (and eight shapes in the feature task) formed

eight points in the reduced 3D space. The eight projected points seemed to lie on the

same plane in both the spatial and feature tasks. Following a similar procedure, we

further applied PCA on the eight points in the reduced plane, with a dimension of 8×3,

to find its best-fit plane. The high proportion of variance explained by the fitted plane

validated our hypothesis that the reduced space could be presented in 2D.

We defined the angles between different conditions based on the best-fit planes

that we constructed above. Given the vectors spanning the best-fit plane (P_1) for one

period are $\vec{v}_1$ and $\vec{v}_2$ (obtained from the PCs of the reduced eight points), and the best-fit

plane (P_2) for another period in the same region are $\vec{v}_3$ and $\vec{v}_4$, the angle of between P_1

and P_2 was calculated as:

$$\langle P_1, P_2 \rangle = \cos^{-1}\left(\frac{|(\vec{v}_1 \times \vec{v}_2) \cdot (\vec{v}_3 \times \vec{v}_4)|}{|\vec{v}_1 \times \vec{v}_2| \cdot |\vec{v}_3 \times \vec{v}_4|}\right),$$

where "$\langle P_1, P_2 \rangle$" denotes the angle between P_1 and P_2, $\cos^{-1}$ is the inverse cosine

function, and "$\vec{v}_1 \times \vec{v}_2$" is the cross product that finds the vector that is perpendicular to

the plane spanned by $\vec{v}_1$ and $\vec{v}_2$, the "·" sign stands for the dot product, and "|x|" returns

the absolute value (for a scaler) or length (for a vector) of x.

To distinguish between a true large rotation of stimulus representation and a

spurious rotation coming about in a population of neurons exhibiting low selectivity and

high variability, we created a control condition for every rotation-angle measurement.

Specifically, we first individually fitted a Poisson distribution for each cell and each

stimulus class (e.g. the circle stimulus in the shape working memory task), using the

same number of trials as in the empirical data angle calculation. Then we randomly

sampled the same number of trials from the simulated distribution. This controlled

dataset has the same level of variability and selectivity statistics, and thus provides a

baseline rotation within each epoch. Only cells with at least 6 match and 6 nonmatch

trials for each stimulus category were included in the rotation angle analysis. Variability

was estimated by randomly drawing 80% of the cells from each prefrontal subdivision

over 100 iterations.

Variance accounted for ratio (VAF)

To cross-check the rotation angle between the 2D subspace of different task epochs, we

measured the Variance Accounted For (VAF) ratio for each angle measurement[15]. The

VAF ratio for epoch subspace pair (1,2) was defined as

$$VAF_{1,2} = \frac{Var(v_2 v_2^T v_1 L_1^T)}{Var(v_1 L_1^T)}$$

Where v_i $(i = 1,2)$ is a 3×2 matrix representing two subspace axes, and L_1 represents the

stimulus projection in the first subspace. VAF ranges from 0 to 1; large values indicate

better alignment of subspaces, while values close to 0 suggest orthogonality. Only cells

with at least 6 match and 6 nonmatch trials for each stimulus category were included in

the VAF analysis.

Geometry of low-dimensional representation

A matrix L of size 8×2 was used to represent the projection of 8 locations or shapes on

the 2D plane. To quantify the amplitude of stimuli representations in two conditions, we

used a scale factor, which is simply the variance ratio between two geometrical

arrangement calculated as follows

$$Scale\ factor = \frac{\sqrt{\sum_p \sum_q C_{second}^2(p,q)}}{\sqrt{\sum_p \sum_q C_{first}^2(p,q)}}$$

$$C_{(p,q)} = L_{(p,q)} - \bar{L}$$

Where C_{first} and C_{second} represent mean-subtracted projections on a 2D plane of the first

and second condition in a comparison, respectively, and where $p=1, \dots ,8$ and $q=1,2$,

representing two coordinates for eight shapes in a 2D subspace. To quantify the

arrangement consistency between experimental conditions, and between experiment data

and hypothetical predictions, we first used the Kabsch algorithm [33] to find the optimal

rotation that aligns the low-dimensional representation in different subspaces. Then we

calculated the covariance matrix $H = (Scale\ factor * C_1)^T C_2$, and determined the SVD

of H as $H = UDV^T$, where D is a rectangular diagonal matrix with non-negative real

values. Thereafter, the optimal rotation matrix R was calculated

$$R = V \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & d \end{bmatrix} U^T$$

$$d = sign(\det(VU^T))$$

441   Lastly, the difference between two low-dimensional manifolds after controlling for scale

442   was quantified as

$$Unexplained\ variance = \frac{2 \times \sqrt{\sum \sum (F_{first} - C_{second})^2}}{\sqrt{\sum_p \sum_q F_{first}^2(p,q)} + \sqrt{\sum_p \sum_q C_{second}^2(p,q)}}$$

Where F_{first} = R(Scale factor*C_{first}) represents the low-dimensional geometry of the first

condition after scaling, translation and rotation.

When comparing the empirical geometry with a hypothetical geometry (Fig. 2),

we assumed that the projection of eight stimuli should fall approximately on a ring, and

that the distance between each cue and its nonmatch pair should be maximized. For the

spatial task, the match-nonmatch structure is symmetric (e.g. when A is cue, B is

nonmatch; and when B is cue A is nonmatch). These two assumptions should lead to a

predicted arrangement similar to the physical appearance of the spatial stimuli, where the

nonmatch pair always perfectly falls on two ends of a diagonal line.

Only cells with at least 6 match and 6 nonmatch trials for each stimulus category

were included in the geometrical similarity analysis. Standard error was measured by

random drawing 80% of cells in each subdivision in 100 iterations.

Time course of subspace rotation

In addition to visualizing the rotation between different phases of the WM task, we

investigated the dynamics of presentations in the reduced PCA space. To construct the

trajectory, we discretized the whole task with a timestep of $\Delta_t = 50\ ms$ and an interval of

$l_t = 250\ ms$. For each timestep k, we recorded the mean firing rates for the match and

nonmatch trials in the interval $[t_0 + (k-1)\Delta_t - 250, t_0 + (k-1)\Delta_t]$ as the entries for

population activity matrix A_1 and A_2 (defined in the above section). Similar as the

procedure of constructing the PCA space, we projected the neural responses for each

condition into their first three principal components. To plot the trajectories, we started

from 250 milliseconds after the start of fixation, i.e., set the initial time $t_0 = 250\ ms$.

Moreover, we use the first three PCAs in the match trials during the cue period as the

common basis to calculate the coordinates. In other words, we projected the activity

matrix into the same common subspace spanned by the PCA space of the match trials.

Note that we can pick up any common subspaces to project the data, and the similar

geometry relations in Figure 3 should hold for all bases. As illustrated in Figure 3, we

explored the dynamics of representation for each location (shape) in the spatial (feature)

task, where we plot every six points from the dynamics, i.e., we set an increment of

$dt = 300 \, ms$ for the purpose of visualization.

To further investigate the changes in decoding spaces, we randomly sampled half

of the trials from all periods (fixation, cue, delay1, sample and delay2) and calculated its

PCA space as the base to project the remaining half trials. The area of the projection for a

given period indicates how much information the state space contains from all eight

locations, which we defined as the decoding subspace (S_t). Take the spatial task for

example, in the reduced space formed by PC1 and PC2, the eight locations were

presented by eight points. To quantify the size of decoding subspace, we first found the

boundaries of the eight points, which was a set of points representing a single conforming

2-D boundary around the eight points (see Figure S4C and D). Then we calculated the

areas of the polygons defined by the boundary points, which we defined as the areas of

the decoding space (A_t). We calculated the ratios of the areas of decoding spaces based

on the fixation period (A_{fix}). For example (as in Figure S4C), the ratios (R) between

fixation, delay1 and delay2 were defined as

487
$$R = \frac{A_{fix}}{A_{fix}} : \frac{A_{delay1}}{A_{fix}} : \frac{A_{delay2}}{A_{fix}} \, .$$

488   Note that we divided the area at fixation period to normalize the data so that we could

489   compare the ratios from different samples. We repeated the random sampling across

490   10,000 iterations, and the statistics of the ratios were reported in the results section.

Significance testing of rotations with bootstrap

Our previous analysis was based on all data recorded from neurons that had more than 16

trials for all cue conditions and in each region in Figure 1C. To test the significance and

robustness of the rotations in the representations, we applied a bootstrap method to

sample 100 iterations from the original data. On each iteration, 80% of the cells were

drawn with replacement. The angles were calculated following the same procedure as

above. Mean angles from 100 iterations obtained from bootstrap plus/minus standard

errors of the mean were superimposed as error bars in figures.

Analysis of error trials

In the post-training spatial task, most cells did not have an error trial. Therefore, the

corresponding activity matrix (as defined above) for the error trials was very sparse. In

the spatial task, there were 295 neurons where error trials presented at more than half of

the eight locations, i.e., 295 rows of the population activity matrix had more than four

non-empty entries. We selected those 295 neurons for the error trial analysis in the spatial

task. Following the same criterion, we selected 201 neurons containing error trials in the

feature task. We first aligned the activity matrix for both cases in the same order, i.e., the

same neuron occupied the same row in both matrices. In most cases, for a given stimulus,

there were more correct trials than error trials, which led to a larger number of samples in

most entries of the activity matrix. Each entry of the activity matrix was calculated from

the mean activity across all trials. We thus matched the number of trials in each entry to

allow comparison between correct and error trials. Specifically, for each empty entry in

the error matrix, we removed the corresponding entry in the correct matrix. Additionally,

we selected the same number (minimum of the corresponding entries) of trials in both

cases to ensure the same number of samples in both the correct and error trials.

Dimensionality reduction was performed on the error trials using PCA. Note that

there are several possible versions of PCA for a sparse matrix; we relied on a

probabilistic principal component (PPCA) with variational Bayesian learning [34] as this

method worked best for our neural data.

Acknowledgments

Research reported in this paper was supported by the National Eye Institute of the

National Institutes of Health under award number R01 EY017077. We wish to thank Rye

Jaffe, Junda Zhu, and Zhengyang Wang for helpful comments on the manuscript.

Author Contributions

C.C. conceived and designed the experiments. C.C. and X. L. Q. performed experiments.

S.P, W.D., and C.C. performed data analysis. C.C., S. P. and W.D. wrote the manuscript

with input from all authors.

Declaration of Interests.

The authors declare no competing interest

REFERENCES

1. Mante V, Sussillo D, Shenoy KV, Newsome WT. Context-dependent computation by recurrent dynamics in prefrontal cortex. *Nature* **503**, 78-84 (2013).

2. Rainer G, Asaad WF, Miller EK. Selective representation of relevant information by neurons in the primate prefrontal cortex. *Nature* **393**, 577-579 (1998).

3. Rigotti M*, et al.* The importance of mixed selectivity in complex cognitive tasks. *Nature* **497**, 585-590 (2013).

4. Ebitz RB, Hayden BY. The population doctrine in cognitive neuroscience. *Neuron*, (2021).

5. Cueva CJ*, et al.* Low-dimensional dynamics for working memory and time encoding. *Proc Natl Acad Sci U S A* **117**, 23021-23032 (2020).

6. Murray JD, Bernacchia A, Roy NA, Constantinidis C, Romo R, Wang XJ. Stable population coding for working memory coexists with heterogeneous neural dynamics in prefrontal cortex. *Proc Natl Acad Sci U S A* **114**, 394-399 (2017).

7. Tang C, Herikstad R, Parthasarathy A, Libedinsky C, Yen SC. Minimally dependent activity subspaces for working memory and motor preparation in the lateral prefrontal cortex. *Elife* **9**, (2020).

8. Okazawa G, Hatch CE, Mancoo A, Machens CK, Kiani R. Representational geometry of perceptual decisions in the monkey parietal cortex. *Cell* **184**, 3748-3761 e3718 (2021).

9. Libby A, Buschman TJ. Rotational dynamics reduce interference between sensory and memory representations. *Nat Neurosci* **24**, 715-726 (2021).

10. Minxha J, Adolphs R, Fusi S, Mamelak AN, Rutishauser U. Flexible recruitment of memory-based choice representations by the human medial frontal cortex. *Science* **368**, (2020).

11. Panichello MF, Buschman TJ. Shared mechanisms underlie the control of working memory and attention. *Nature* **592**, 601-605 (2021).

12. Kriegeskorte N, Wei XX. Neural tuning and representational geometry. *Nat Rev Neurosci* **22**, 703-718 (2021).

575 13. Meyer T, Qi XL, Stanford TR, Constantinidis C. Stimulus selectivity in dorsal and
576   ventral prefrontal cortex after training in working memory tasks. *J Neurosci* **31**,
577   6266-6276 (2011).
578

579 14. Riley MR, Qi XL, Zhou X, Constantinidis C. Anterior-posterior gradient of plasticity
580   in primate prefrontal cortex. *Nat Commun* **9**, 3790 (2018).
581

582 15. Xie Y*, et al.* Geometry of sequence working memory in macaque prefrontal
583   cortex. *Science* **375**, 632-639 (2022).
584

585 16. Dang W, Jaffe RJ, Qi XL, Constantinidis C. Emergence of Nonlinear Mixed
586   Selectivity in Prefrontal Cortex after Training. *J Neurosci* **41**, 7420-7434 (2021).
587

588 17. Dux PE, Tombu MN, Harrison S, Rogers BP, Tong F, Marois R. Training improves
589   multitasking performance by increasing the speed of information processing in
590   human prefrontal cortex. *Neuron* **63**, 127-138 (2009).
591

592 18. Klingberg T*, et al.* Computerized Training of Working Memory in Children with
593   ADHD - a Randomized, Controlled Trial. *J Am Acad Child Adolesc Psychiatry* **44**,
594   177-186 (2005).
595

596 19. Jaeggi SM, Buschkuehl M, Jonides J, Perrig WJ. Improving fluid intelligence with
597   training on working memory. *ProcNatlAcadSciUSA* **105**, 6829-6833 (2008).
598

599 20. Constantinidis C, Klingberg T. The neuroscience of working memory capacity and
600   training. *Nat Rev Neurosci* **17**, 438-449 (2016).
601

602 21. Qi XL, Constantinidis C. Neural changes after training to perform cognitive tasks.
603   *Behav Brain Res* **241**, 235-243 (2013).
604

605 22. Asaad WF, Rainer G, Miller EK. Neural activity in the primate prefrontal cortex
606   during associative learning. *Neuron* **21**, 1399-1407 (1998).
607

608 23. Rule ME, Loback AR, Raman DV, Driscoll LN, Harvey CD, O'Leary T. Stable task
609   information from an unstable neural population. *Elife* **9**, (2020).
610

611 24. Rule ME, O'Leary T. Self-healing codes: How stable neural populations can track
612   continually reconfiguring neural representations. *Proc Natl Acad Sci U S A* **119**,
613   (2022).
614

615 25. Bernardi S, Benna MK, Rigotti M, Munuera J, Fusi S, Salzman CD. The Geometry
616   of Abstraction in the Hippocampus and Prefrontal Cortex. *Cell* **183**, 954-967 e921
617   (2020).
618

619   26.   Meyers EM, Qi XL, Constantinidis C. Incorporation of new information into
620   prefrontal cortical activity after learning working memory tasks. *Proc Natl Acad*
621   *Sci U S A* **109**, 4651-4656 (2012).
622
623   27.   Wilson FA, Scalaidhe SP, Goldman-Rakic PS. Dissociation of object and spatial
624   processing domains in primate prefrontal cortex. *Science* **260**, 1955-1958 (1993).
625
626   28.   Badre D, D'Esposito M. Is the rostro-caudal axis of the frontal lobe hierarchical?
627   *Nat Rev Neurosci* **10**, 659-669 (2009).
628
629   29.   Rao SC, Rainer G, Miller EK. Integration of what and where in the primate
630   prefrontal cortex. *Science* **276**, 821-824 (1997).
631
632   30.   Owen AM, Stern CE, Look RB, Tracey I, Rosen BR, Petrides M. Functional
633   organization of spatial and nonspatial working memory processing within the
634   human lateral frontal cortex. *Proc Natl Acad Sci U S A* **95**, 7721-7726 (1998).
635
636   31.   Constantinidis C, Qi XL. Representation of Spatial and Feature Information in the
637   Monkey Dorsal and Ventral Prefrontal Cortex. *Front Integr Neurosci* **12**, 31
638   (2018).
639
640   32.   Meyer T, Constantinidis C. A software solution for the control of visual
641   behavioral experimentation. *J Neurosci Methods* **142**, 27-34. (2005).
642
643   33.   Kabsch W. A solution for the best rotation to relate two sets of vectors. *Acta*
644   *Crystallogr* **A32**, 922 (1976).
645
646   34.   Ilin A, Raiko T. Practical Approaches to Principal Component Analysis in the
647   Presence of Missing Values. *Journal of Machine Learning Research*, 1957-2000
648   (2010).
649
650

FIGURES

Figure 1. **Task structure and recording areas**. The animals were required to maintain
center fixation throughout both active and passive task trials. At the end of active tasks
trials, monkeys were required to make a saccade to a green target if the stimuli matched
or to a blue target if the stimuli did not match. (A) Spatial match-to-sample task; the eight
cue locations analyzed are shown in the inset. (B) Shape feature match-to-sample task;
eight possible shapes in a session shown are shown in the inset. (C). Neurons were
recorded from 5 subdivisions of lateral PFC, which we refer to as Anterior Dorsal (AD),
Anterior Ventral (AV), Mid-Dorsal (MD), Posterior Dorsal (PD), and Posterior Ventral
(PV).

665
Figure 2. **Comparison of within-subspace geometry for both tasks.** (A) A graphical
summary of how representational geometry was compared. Geometries were calculated
from cue period activity of the spatial task, pre- and post-training. The geometries were
compared to a hypothetical geometry that maximizes the distance between match and
nonmatch stimulus pairs (see Methods). (B) Geometrical similarity to the hypothetical
configuration for both tasks in the cue and the first delay period. Filled bars represent
empirical data, and dashed bars represent similarity measure after shuffling shape labels
as a control. The bar height represents deviation from both the circle configuration, and
from the 'standard' order of stimuli on the circle. A smaller difference between the filled
and dashed bars shows a higher similarity between the representational geometry and the
'standard' hypothetical geometry. The y-axis units are arbitrary.
677
678

679
Figure 3. Subspace rotation between the cue period and the sample period in match
trials. (A) Illustration of subspace rotations. Average neural response of different
conditions were projected onto a 3D PCA-space, where the rotation and geometrical
similarity were calculated (see Methods for details). (B) Filled boxes: angles between the
cue and the sample-match condition in different PFC subdivisions for the spatial and the
feature task. Unfilled boxes: angles from the control experiments obtained by randomly

drawing data from a distribution fitted to the empirical data for each task epoch. The dot
and horizontal line in the box represent the mean and median, respectively. Whiskers are
five percentile minimum and maximum values. (C) Variance accounted for (VAF)
between the cue and the sample-match condition in difference PFC subdivisions for the
spatial and the feature task. (D) Left: a graphic summary of subspace geometrical
similarity. The scale factor was the ratio between within-subspace total variance. The
unexplained variance quantifies geometrical structure similarity after controlling the scale
and orientation differences of representation in respective subspaces. Right: Geometrical
similarity between cue and sample-match. Gray boxes represent results from the cue-
match dataset, boxes with oblique lines are results from the within cue period control
group.
697

698
Figure 4. Subspace rotation between the match and nonmatch conditions in the
sample period. (A) Primary angles between the match and non-match in difference PFC
subdivisions for the spatial and the feature task. Primary angles between the cue and the
sample-match condition in different PFC subdivisions for the spatial and the feature task.
As a control we randomly draw data from a distribution fitting to empirical data from a
particular task epoch (unfilled boxes). In this figure control is within the match condition
of the sample period. (B) Variance accounted for (VAF) between the match and non-
match conditions in difference PFC subdivisions for the spatial and the feature task. (C)
Geometrical similarity between match and nonmatch in the sample period. Gray boxes
represent results from match-nonmatch dataset, while boxes with oblique lines are results
from the within match period control group.
710

711
712
Figure 5. Population representations in correct and error trials. (A) Representation
of spatial stimuli during the match and nonmatch period in correct trials (n=295 neurons).
Rotation φ=76°. (B) Representation of spatial stimuli during the match and nonmatch
period in the error trials (same population as in panel A). Rotation φ=15°. (C)
Representation of feature stimuli during the match and nonmatch period in correct trials
(n=201 neurons). Rotation φ=84°. (D) Representation of feature stimuli during the match
and nonmatch period in error trials (same population as in panel C). Rotation φ=7°. (E)
Dynamics of stimulus representations in correct and error trials, for one example spatial
location. Solid black: match in correct trials, solid gray: nonmatch in correct trials,
dashed red: match in error trials, dashed magenta: nonmatch in error trials. Always same
cells are plotted (n=295 neurons).
724
725

727
728 **Supplemental Figure S1.** Explained variance distribution for both tasks pre- and post-
729 training in the PCA analysis. In general, fewer dimensions were needed for characterizing
730 population activity in the spatial task, but under most conditions, three dimensions were
731 enough to explain more than 50% of the variance. The training process has a
732 heterogenous effect on the distribution of variance in different subdivisions of LFPC.
733

734
735

736
737
738 **Supplemental Figure S2.** Scatter plot of subspace rotation angle for the spatial and the
739 feature task. Each dot represents the mean angle measure for one PFC subdivision, blue
740 color dots indicate spatial task, and red dots indicate feature task.

741
Supplemental Figure S3. Subspace rotation between the cue and the first delay. (A)
Filled boxes: primary angles between the cue and the first delay in difference PFC
subdivisions for the spatial and the feature task. Unfilled boxes: control group obtained
by randomly drawing data from a distribution fitting to empirical data from a particular
task epoch. In this figure control is within the first delay period. (B) Variance accounted
for (VAF) between the cue and the first delay in difference PFC subdivisions for the
spatial and the feature task**.** (C) Geometrical similarity between the cue and the firs delay
period. Gray boxes: results from cue-delay dataset; boxes with oblique lines: results from
the within first delay control group.

751
752
Supplemental Figure S4. Complete population activity trajectories before and after training. (A) Trajectory of population representation in state space prior to training, for the spatial stimuli in the mid dorsal area (n=453). Solid lines: match trials, dashed lines: nonmatch trials. **(B)** As in A, for neuronal activity in the mid-dorsal area after training (n=374). **(C)** Area covered by projection of spatial stimuli prior to training, in three task epochs: fixation, delay1, and delay2. **(D)** As in C, for data obtained after training. **(E)**. Boxplot of surface areas: summarizing statistics of panels C and D with bootstrapping sampling, where each sample contained 80% of cells. Area values have been normalized by that of the fixation period.

763
764

765
766
767
768 **Supplemental Figure S5. Population representations in correct and error trials. (A)**
769 Representation of spatial stimuli during the fixation period in correct and error trials
770 (n=295 neurons). Rotation φ=45°. (B). As in panel A, for the cue period. Rotation φ=81°.
771 (C) As in panel A for the first delay period. Rotation φ=82°.