Genetic, Clinical Underpinnings of Brain Change Along Two Neuroanatomical Dimensions of Clinically-defined Alzheimer's Disease

Junhao Wen[1,2*], Zhijian Yang[2], Ilya M. Nasrallah[2], Yuhan Cui[2], Guray Erus[2], Dhivya Srinivasan[2], Ahmed Abdulkadir[2,3], Elizabeth Mamourian[2], Ioanna Skampardoni[2], Gyujoon Hwang[2], Ashish Singh[2], Mark Bergman[2], Jingxuan Bao[4], Erdem Varol[5], Zhen Zhou[2], Aleix Boquet-Pujadas[6], Jiong Chen[2], Arthur W. Toga[7], Andrew J. Saykin[8], Timothy J. Hohman[9], Paul M. Thompson[10], Sylvia Villeneuve[11], Randy Gollub[12], Aristeidis Sotiras[13], Katharina Wittfeld[14], Hans J. Grabe[14], Duygu Tosun[15], Murat Bilgel[16], Yang An[16], Daniel S. Marcus[17], Pamela LaMontagne[17], Tammie L. Benzinger[17], Susan R. Heckbert[18], Thomas R. Austin[18], Lenore J. Launer[19], Mark Espeland[20], Colin L Masters[21], Paul Maruff[21], Jurgen Fripp[22], Sterling C. Johnson[23], John C. Morris[24], Marilyn S. Albert[25], R. Nick Bryan[26], Susan M. Resnick[16], Luigi Ferrucci[27], Yong Fan[2], Mohamad Habes[28], David Wolk[2, 29], Li Shen[4], Haochang Shou[2,30], Christos Davatzikos[2*], for the iSTAGING, the AI4AD, and the ADSP phenotypic harmonization consortia, the BLSA, the PREVENT-AD, and the ADNI studies

[1]Laboratory of AI and Biomedical Science (LABS), Stevens Neuroimaging and Informatics Institute, Keck School of Medicine of USC, University of Southern California, Los Angeles, California, USA.
[2]Artificial Intelligence in Biomedical Imaging Laboratory (AIBIL), Center for AI and Data Science for Integrated Diagnostics (AI^{2}D), Perelman School of Medicine, University of Pennsylvania, Philadelphia, USA.
[3]Research Lab in Neuroimaging of the Department of Clinical Neurosciences at Lausanne University Hospital, Lausanne, Switzerland
[4]Department of Biostatistics, Epidemiology and Informatics University of Pennsylvania Perelman School of Medicine, Philadelphia, USA
[5]Department of Statistics, Center for Theoretical Neuroscience, Zuckerman Institute, Columbia University, New York, New York
[6]Biomedical Imaging Group, EPFL, Lausanne, Switzerland
[7]Laboratory of Neuro Imaging, Stevens Neuroimaging and Informatics Institute, Keck School of Medicine of USC, University of Southern California, Marina del Rey, California
[8]Radiology and Imaging Sciences, Center for Neuroimaging, Department of Radiology and Imaging Sciences, Indiana Alzheimer's Disease Research Center and the Melvin and Bren Simon Cancer Center, Indiana University School of Medicine, Indianapolis
[9]Vanderbilt Memory and Alzheimer's Center, Vanderbilt Genetics Institute, Department of Neurology, Vanderbilt University Medical Center, Nashville, TN USA
[10]Imaging Genetics Center, Mark and Mary Stevens Neuroimaging and Informatics Institute, Keck School of Medicine of USC, University of Southern California, Marina del Rey, California
[11]Douglas Mental Health University Institute, McGill University, Montréal, Québec, Canada
[12]Athinoula A. Martinos Center for Biomedical Imaging, Massachusetts General Hospital, Harvard Medical School, Charlestown, Massachusetts
[13]Department of Radiology and Institute for Informatics, Washington University School of Medicine, St. Louis, USA
[14]Department of Psychiatry and Psychotherapy, University Medicine Greifswald, Greifswald, Germany

47 [15]Department of Radiology and Biomedical Imaging, University of California, San Francisco,
48 CA, USA
49 [16]Laboratory of Behavioral Neuroscience, National Institute on Aging, NIH, USA
50 [17]Department of Radiology, Washington University School of Medicine, St. Louis, Missouri,
51 USA
52 [18]Cardiovascular Health Research Unit and Department of Epidemiology, University of
53 Washington, Seattle, WA, USA
54 [19]Neuroepidemiology Section, Intramural Research Program, National Institute on Aging,
55 Bethesda, Maryland, USA
56 [20]Sticht Center for Healthy Aging and Alzheimer's Prevention, Wake Forest School of
57 Medicine, Winston-Salem, North Carolina, USA
58 [21]Florey Institute of Neuroscience and Mental Health, The University of Melbourne,
59 Parkville, VIC, Australia
60 [22]CSIRO Health and Biosecurity, Australian e-Health Research Centre CSIRO, Brisbane,
61 Queensland, Australia
62 [23]Wisconsin Alzheimer's Institute, University of Wisconsin School of Medicine and Public
63 Health, Madison, Wisconsin, USA
64 [24]Knight Alzheimer Disease Research Center, Washington University in St. Louis, St. Louis,
65 MO, USA
66 [25]Department of Neurology, Johns Hopkins University School of Medicine, USA
67 [26]Department of Radiology, University of Pennsylvania, Philadelphia, USA.
68 [27]Translational Gerontology Branch, Longitudinal Studies Section, National Institute on
69 Aging, National Institutes of Health, MedStar Harbor Hospital, 3001 S. Hanover Street,
70 Baltimore, MD, 21225, USA
71 [28]Glenn Biggs Institute for Alzheimer's & Neurodegenerative Diseases, University of Texas
72 Health Science Center at San Antonio, San Antonio, USA
73 [29]Department of Neurology and Penn Memory Center, University of Pennsylvania,
74 Philadelphia, USA
75 [30]Penn Statistics in Imaging and Visualization Center, Department of Biostatistics,
76 Epidemiology, and Informatics, Perelman School of Medicine, University of Pennsylvania,
77 Philadelphia, USA
78
79 [*]Correspondence to:
80 Junhao Wen, Ph.D. – junhaowe@usc.edu
81 2025 Zonal Ave, Los Angeles, CA 90033, United States
82 Christos Davatzikos, Ph.D. – Christos.Davatzikos@pennmedicine.upenn.edu
83 3700 Hamilton Walk, 7th Floor, Philadelphia, PA 19104

85 **Running title**: Two AI Dimensions of Alzheimer's Disease
86 **Keywords:** Alzheimer's disease; dimensions; imaging-genomics; deep learning; semi-
87 supervised representation learning.

Abstract

Alzheimer's disease (AD) is associated with heterogeneous atrophy patterns. We employed a semi-supervised clustering technique known as Surreal-GAN, through which we identified two dominant dimensions of brain atrophy in symptomatic mild cognitive impairment (MCI) and AD patients: the "diffuse-AD" (R1) dimension shows widespread brain atrophy, and the "MTL-AD" (R2) dimension displays focal medial temporal lobe (MTL) atrophy. Critically, only R2 was associated with widely known sporadic AD genetic risk factors (e.g., *APOE ε4*) in MCI and AD patients at baseline. We then independently detected the presence of the two dimensions in the early stages by deploying the trained model in the general population and two cognitively unimpaired cohorts of asymptomatic participants. In the general population, genome-wide association studies found 77 genes unrelated to *APOE* differentially associated with R1 and R2. Functional analyses revealed that these genes were overrepresented in differentially expressed gene sets in organs beyond the brain (R1 and R2), including the heart (R1) and the pituitary gland, muscle, and kidney (R2). These genes were enriched in biological pathways implicated in dendritic cells (R2), macrophage functions (R1), and cancer (R1 and R2). Several of them were "druggable genes" for cancer (R1), inflammation (R1), cardiovascular diseases (R1), and diseases of the nervous system (R2). The longitudinal progression showed that *APOE ε4*, amyloid, and tau were associated with R2 at early asymptomatic stages, but this longitudinal association occurs only at late symptomatic stages in R1. Our findings deepen our understanding of the multifaceted pathogenesis of AD beyond the brain. In early asymptomatic stages, the two dimensions are associated with diverse pathological mechanisms, including cardiovascular diseases, inflammation, and hormonal dysfunction – driven by genes different from *APOE* – which may collectively contribute to the early pathogenesis of AD.

Introduction

Alzheimer's disease (AD) is the most common cause of dementia in older adults and remains incurable despite many pharmacotherapeutic clinical trials, including anti-amyloid drugs[1,2] and anti-tau drugs.[3] This is largely due to the complexity and multifaceted nature of the underlying neuropathological processes leading to dementia. The research community has embraced several mechanistic hypotheses to elucidate AD pathogenesis.[4–6] Among these, the amyloid hypothesis has been dominant over the past decades and has proposed a dynamic biomarker chain: extracellular beta-amyloid (Aβ) triggers a cascade that leads to subsequent intracellular neurofibrillary tangles, including hyperphosphorylated tau protein (tau and p-tau), neurodegeneration, including medial temporal lobe atrophy, and cognitive decline.[7,8] However, the amyloid hypothesis has been reexamined and revised due to substantial evidence which questions its current form[8–10]. While amyloid remains critical to AD development, the amyloid cascade model has been continually refined as other biological factors are discovered to influence the pathway from its accumulation to cell death.

Cardiovascular dysfunction has been widely associated with an increased risk for AD.[11] There is also growing evidence that inflammatory[10–12] and neuroendocrine processes[5,13] influence pathways of amyloid accumulation and neuronal death. The inflammation hypothesis claims that microglia and astrocytes release pro-inflammatory cytokines as drivers, byproducts, or beneficial responses associated with AD progression and severity.[12–14] The neuroendocrine hypothesis, first introduced in the context of aging[15], has been extended to AD[16], where it proposes that neurohormones secreted by the pituitary and other essential endocrine glands can affect the central nervous system (CNS), which subsequently contribute to developing AD. For example, Xiong and colleagues[17] recently found that blocking the action of follicle-stimulating hormone in mice abrogates the AD-like phenotype (e.g., cognitive decline) by inhibiting the neuronal C/EBPβ–δ-secretase pathway. These findings emphasize the need to further elucidate early brain and body changes well before they lead to irreversible clinical progression.[18]

Recent advances in artificial intelligence (AI), especially deep learning (DL), applied to magnetic resonance imaging (MRI), showed great promise in biomedical applications[19,20]. DL models discover complex non-linear relationships between phenotypic and genetic

features and clinical outcomes, thereby providing informative imaging-derived
endophenotypes[21]. In particular, AI has been applied to MRI to disentangle the
neuroanatomical heterogeneity of AD with categorical disease subtypes.[22–26] The genetic
underpinnings[27,28] of this neuroanatomical heterogeneity in AD are also complex and
heterogeneous. The most recent large-scale genome-wide association study[27] (GWAS:
111,326 AD vs. 677,633 controls) has identified 75 genomic loci, including *APOE* genes,
associated with AD. However, such case-control group comparisons conceal genetic factors
that might contribute differentially to different dimensions of AD-related brain change. More
importantly, the genetic variants that contribute to the initiation and early progression of brain
change in younger and asymptomatic individuals are poorly understood.

In this study, we utilize a novel semi-supervised deep learning approach, Surreal-
GAN, to characterize the neuroanatomical heterogeneity of the disease. Unlike our previous
model, Smile-GAN[22], which categorized subtypes, Surreal-GAN generates multiple
continuous dimensional scores, simultaneously accounting for spatial and temporal disease
heterogeneity, similar to what was accomplished in a previous unsupervised clustering model
known as Sustain. These multi-dimensional scores reflect the co-expression level of
respective brain atrophy dimensions in the same patient. Refer to the method (**Surreal-GAN**
for disease heterogeneity) and **Supplementary eMethod 1** for methodological details of
Surreal-GAN, comparisons to other subtyping methods, and strengths of semi-supervised
representation learning. We hypothesized that genetic variants, potentially unrelated to *APOE*
genes, contribute to early manifestations of multiple dimensions of brain atrophy in early
asymptomatic stages. We first trained the Surreal-GAN model to define the AD dimensions
to test this hypothesis in the late symptomatic stages. We then examined their expression
back to early asymptomatic stages. In our previous study,[29] we derived two neuroanatomical
dimensions (R1 and R2) by applying Surreal-GAN to the MCI/AD participants (target
population) and cognitively unimpaired (CU) participants (reference population) from the
Alzheimer's Disease Neuroimaging Initiative study (ADNI[30]). Herein, we applied the trained
model to three asymptomatic populations and one symptomatic population: the *general*
population (*N*=39,575; age: 64.12±7.54 years) from the UK Biobank (UKBB[31]) excluding
demented individuals; the *cognitively unimpaired population* (*N*=1658; age: 65.75±10.90
years) from ADNI and the Baltimore Longitudinal Study of Aging study (BLSA[32]); the
cognitively unimpaired population with a family risk (*N*=343; age: 63.63±5.05 years) from
the Pre-symptomatic Evaluation of Experimental or Novel Treatments for Alzheimer's

Disease (PREVENT-AD[33]); the *MCI/AD population* (N=1534; age: 73.45±7.69 years) from

ADNI and BLSA. Refer to the method (**Study design and populations**) and **Table 1** for

details of the definition of these populations.

Materials and methods

Study design and populations

The current study consists of four main populations (**Table 1**), which were jointly consolidated by the iSTAGING[34] and the AI4AD consortia (http://ai4ad.org/): the iSTAGING consortium consolidated all imaging and clinical data; imputed genotyping data were downloaded from UKBB; the AI4AD consortium consolidated the whole-genome sequencing (WGS) data for the ADNI study. The definition of each population and inclusion criteria are detailed in **Supplementary eMethod 2**.

Image preprocessing

All T1w-weighted MR images were first corrected for magnetic field intensity inhomogeneity.[35] A deep learning-based skull stripping algorithm was applied for the removal of extra-cranial material. In total, 145 anatomical regions of interest (ROIs) were generated in gray matter (GM, 119 ROIs), white matter (WM, 20 ROIs), and ventricles (6 ROIs) using a multi-atlas label fusion method[36] (**Supplementary eMethod 3**). The 119 ROIs were statistically harmonized by an extensively validated approach, i.e., ComBat-GAM[37], using the entire imaging data of iSTAGING. **Supplementary eFigure 1** demonstrates the normality check for the MUSE ROI (right accumbent area) before and after statistical harmonization, illustrating that our statistical harmonization enhanced the normality of ROIs across various studies.

Surreal-GAN for disease heterogeneity

Surreal-GAN[29] dissects underlying disease-related heterogeneity via a deep representation learning approach under the principle of semi-supervised learning.[22,23] At a high level, its most fundamental novelty is that it provides a continuous representation of the presence of multiple, non-exclusive abnormal brain patterns in each individual, rather than clustering individuals into one of many clusters, i.e. disease subtypes. More specifically, several methodological advancements were considered compared to its predecessor, Smile-GAN[22]. First, Surreal-GAN is to model neuroanatomical heterogeneity by considering both spatial and temporal (i.e., disease severity) variation using only cross-sectional MRI data. Secondly, Surreal-GAN disentangles the neuroanatomical heterogeneity of AD by enabling patients to

simultaneously exhibit multiple distinct imaging patterns (i.e., high scores for expressing all
these patterns), resulting in high-dimensional scores across multiple dimensions. Lastly, in
contrast to prior probability-based clustering methods like Smile-GAN, Surreal-GAN
operates without the constraint that all dimensional scores must sum to 1. This allows for a
more normal distribution of dimensional scores suited for GWAS (**Supplementary eFigure**
2). Further methodological details are elaborated upon in **Supplementary eMethod 1**.

Alternative clustering techniques, such as Sustain[24] and Bayesian latent[38] methods, are
available for deciphering the neuroanatomical heterogeneity in AD[22–26,39]. Surreal-GAN
distinguishes itself from these approaches based on fundamental methodological distinctions,
such as its utilization of semi-supervised deep learning compared to the unsupervised
approach of others[40]. Additionally, Surreal-GAN generates continuous dimensions associated
with distinct phenotypic outcomes, allowing the simultaneous co-expression of multiple
patterns instead of categorizing patients into a single dominant subtype or stage, as seen in
other methods. Notably, the two dimensions, R1 and R2, displayed correlations with the four
subtypes generated by Smile-GAN, particularly R2 exhibited a correlation with P3 (reflecting
medial temporal lobe atrophy), and both R1 and R2 displayed correlations with P4
(representing global atrophy), as depicted in **Supplementary eFigure 3**.

Brain and clinical variable associations

We performed brain-wide associations for the 119 GM ROIs. For baseline brain-wide
associations, linear regression models were fitted with R1 and R2 dimensions as independent
variables, with each ROI as the dependent variable, controlling for age, sex, intracranial
volume (ICV), and/or diagnosis as covariates.

We performed a two-step linear regression for longitudinal brain-wide associations.
First, we derived the individual-level age change rate using a linear mixed-effects model. To
this end, we included a participant-specific random slope for age and random intercept; age
and sex were treated as fixed effects. Secondly, the same linear regression model as in
baseline brain associations was fitted with the age change rate as the independent variable.

We also performed clinical variable association for all clinical variables and
neuropsychological testing available for each population, using the same model in the
baseline brain-wide associations. Bonferroni correction of 119 GM ROIs was performed to
adjust for the multiple comparisons.

Genetic analyses

Genetic analyses were performed for the WGS data from ADNI and the imputed genotype data from UKBB. Our quality check protocol is detailed in **Supplementary eMethod 4**. This resulted in 1487 participants and 24,194,338 SNPs in ADNI WGS data. For UKBB, we limited our analysis to European ancestry participants, resulting in 33,541 participants and 8,469,833 SNPs.

Using UKBB data, we first estimated the SNP-based heritability using GCTA-GREML,[41] controlling for confounders of age (at imaging), age-squared, sex, age-sex interaction, age-squared-sex interaction, ICV, and the first 40 genetic principal components, following a previous pioneer study.[42] In GWAS, we performed a linear regression for each neuroanatomical dimension and included the same covariates as in the heritability estimates. We adopted the genome-wide P-value threshold (5×10^{-8}) in all GWAS. The annotation of genomic loci (displayed by its top lead SNP) and gene mappings, prioritized gene set enrichment, and tissue specificity analyses were performed using FUMA (https://fuma.ctglab.nl/, version: v1.3.8) (**Supplementary eMethod 5** and **6**). A two-step procedure (**Supplementary eMethod 7**) was performed to determine if an annotated genomic locus or gene was associated with AD-related clinical traits. We calculated the polygenic risk scores (PRS)[43] using both ADNI and UKBB genetic data (**Supplementary eMethod 8**). Finally, we constructed a target-drug-disease network for these genes associated with R1 and R2 to identify these "druggable genes" (**Supplementary eMethod 9**).

Results

Two dominant dimensions of brain atrophy found in MCI and AD

In MCI/AD patients, the "diffuse-AD" dimension (R1) showed widespread brain atrophy without an exclusive focus on the medial temporal lobe (**Fig. 1A** and **Supplementary eTable 1** for P-values and effect sizes). In contrast, the "MTL-AD" dimension (R2) displayed more focal medial temporal lobe atrophy, prominent in the bilateral parahippocampal gyrus, hippocampus, and entorhinal cortex (**Fig. 1A**). All results, including P-values and effect sizes (Pearson's correlation coefficient r), are presented in **Supplementary eTable 1**. The atrophy patterns of the two dimensions defined in the symptomatic MCI/AD population (**Fig. 1A**) were present in the asymptomatic populations, albeit with a smaller magnitude of r. (**Supplementary eTable 1**, **4**, and **8**). We presented the age distribution (**Supplementary eFigure 4A**), as well as the expression of R1 and R2, along with the population-level difference for the four populations in **Supplementary eFigure 4B-D**.

At baseline, the R1-dominant group had 25.72 % AD patients (N=222 out of 863); the R2-dominant group consisted of 30.10% AD patients (202 out of 671). Within a 7-year follow-up period, MCI participants from both the R1-dominant and R2-dominant groups progressed to AD, with the R2-dominant group exhibiting a higher proportion of AD patients (40% vs. 25%) (**Supplementary eFigure 5 A-B**); the two dominant dimensions developed independently throughout the 7-year follow-up period (**Supplementary eFigure 5 C-D**).

APOE genes are associated with R2 but not with R1 in the MCI/AD population

In GWAS, the R2 dimension, but not R1, was associated with well-established AD genomic loci (_rs429358_, chromosome: 19, 45411941; minor allele: C, P-value: 1.05×10^{-11}) and genes (_APOE_, _PVRL2_, _TOMM40_, and _APOC1_) (**Fig. 1B**). The details of the identified genomic locus and annotated genes are presented in **Supplementary eTable 2**. The PRS of AD showed a slightly stronger positive association with the R2 dimension [r=0.11, -$\log_{10}$(P-value)=3.14] than with the R1 dimension [r=0.09, -$\log_{10}$(P-value)=2.31, **Supplementary eFigure 6**]. The QQ plots of the baseline GWAS are presented in **Supplementary eFigure 7**.

Clinical profiles of the R1 and R2 dimensions in the MCI/AD population

Clinical association studies correlated the two dimensions with 45 clinical variables and

biomarkers. Compared to the R1 dimension, R2 showed associations, to a larger extent than

R1, with SPARE-AD and the Rey Auditory Verbal Learning Test (RAVLT). SPARE-AD

quantifies the presence of a typical imaging signature of AD-related brain atrophy, which has

been previously shown to predict clinical progression in both CU and MCI individuals.[44]

RAVLT measures episodic memory, a reliable neuropsychological phenotype in AD, which

is also correlated with medial temporal lobe atrophy.[45,46] The R1 dimension was associated to

a greater extent with 1) SPARE-BA, which captures the individualized expression of

advanced brain age from MRI[47]; 2) white matter lesions (WML), which are commonly

associated with vascular risk factors and cognitive decline[48], and 3) whole-brain uptake of

18F-fluorodeoxyglucose (FDG) PET, which is a biomarker of brain metabolic function and

atrophy. Both dimensions were positively associated with cerebrospinal fluid (CSF) levels of

tau and p-tau and negatively associated with the CSF level of Aβ42[49] (**Fig. 1C**), as well as the

whole-brain standardized uptake value ratio of 18F-AV-45 PET (**Supplementary eTable 3**).

Results for all 45 clinical variables, including cognitive scores, modifiable risk factors, CSF

biomarkers, disease/condition labels, demographic variables, and imaging-derived

phenotypes, are presented in **Supplementary eTable 3** for P-values and effect sizes (i.e., beta

coefficients).

Clinical profiles of the R1 and R2 dimensions in the general population

Brain association studies confirmed the presence of the two atrophy patterns in the general population (**Fig. 2A** and **Supplementary eTable 4** for P-values and effect sizes). In clinical association studies, the R1 dimension was significantly associated, to a larger extent than R2, with cardiovascular (e.g., triglycerides) and diabetes factors (e.g., Hba1c and glucose), executive function (TMT-B), intelligence, physical measures (e.g., diastolic blood pressure), SPARE-BA [$-\log_{10}$(P-value) = 236.89 for R1 and -46.35 for R2] and WML [$-\log_{10}$(P-value) = 120.24 for R1 and 2.06 for R2]. In contrast, the R2 dimension was more significantly associated with SPARE-AD [$-\log_{10}$(P-value) = 136.01 for R1 and 250.41 for R2] and prospective memory (**Fig. 2B**). Results for all 61 clinical variables, including cardiovascular factors, diabetic blood markers, social demographics, lifestyle, physical measures, cognitive scores, and imaging-derived phenotypes, are presented in **Supplementary eTable 5** for P-values and effect sizes.

Twenty-four genomic loci and seventy-seven genes unrelated to *APOE* are associated with the R1 and R2 dimensions in the general population

GWAS identified 24 genomic loci, 14 of which are novel (not previously associated with any traits in GWAS Catalog), and 77 positionally and functionally mapped genes unrelated to *APOE* associated with R1 or R2. In particular, the R1 dimension was significantly associated with 11 genomic loci and 49 genes. Eight genes (blue-colored genes in **Fig. 2D**) were previously associated with AD-related traits; 12 novel loci/genes have not been previously associated with any clinical traits. The R2 dimension was significantly associated with 13 genomic loci and 40 annotated genes. 13 genes (red-colored genes in **Fig. 2D**) were associated with AD-related traits; 8 loci/genes were novel (**Fig. 2C**, **Supplementary eTable 6**). These genomic loci and genes were also associated with many clinical traits in the literature from the GWAS Catalog.[50] These included hormones (e.g., sex hormone-binding globulin measurement vs. *CCKN2C*), inflammatory factors (e.g., macrophage inflammatory protein 1b measurement vs. *CDC25A*), imaging-derived phenotypes (e.g., cerebellar volume measurement from MRIs vs. *DMRTA2*), and psychiatric disorders (e.g., unipolar depression vs. *ASTN2*) (**Fig. 2D**). Details of the GWAS Catalog results are presented in **Supplementary eFile 1**. The Manhatton and QQ plots of the baseline GWAS are presented in **Supplementary eFigure 8**. The LDSC[51] interpcept of the two GWASs was close to 1,

indicating no substantial genomic inflation (R1=1.0032±0.0084; R2=1.023±0.0084).

Furthermore, our main GWASs using European ancestry were robust in three sensitivity

check analyses: split-sample, sex-stratified, and mixed-effect[52] linear model analyses.

Detailed results are presented in **Supplementary eText 1** and **Supplementary eFile2-4**.

The two dimensions were significantly heritable in the general population based on

the SNP-based heritability estimates (R1: $h^2 = 0.49 \pm 0.02$; R2: $h^2 = 0.55 \pm 0.02$). The PRS

of AD showed a marginally positive association with the R2 dimension [-$\log_{10}$(P-

value)=1.42], but not with the R1 dimension [-$\log_{10}$(P-value)=0.47<1.31] in this population.

Genes associated with the R1 and R2 dimensions are overrepresented in organs beyond

the brain in the general population

Tissue specificity analyses test whether the mapped genes are overrepresented in

differentially expressed gene sets (DEG) in one organ/tissue compared to all other

organs/tissues using different gene expression data[53]. The genes associated with the R1

dimension were overrepresented in caudate, hippocampus, putamen, amygdala, substantia

nigra, liver, heart, and pancreas; the genes associated with the R2 dimension were

overrepresented in caudate, hippocampus, putamen, amygdala, anterior cingulate, pituitary,

liver, muscle, kidney, and pancreas (**Fig. 3A** and **Supplementary eFigure 9**). Genes in DEG

over-expressed in the heart were only associated with R1, while those in DEG over-expressed

in the pituitary gland, muscle, and kidney were unique in R2. The expression values of every

single gene for all tissues are presented in **Supplementary eFigure 10**.

Genes associated with the R1 and R2 dimensions are enriched in key biological

pathways in the general population

Genes associated with the two dimensions were enriched in different biological pathways.

Genes associated with the two dimensions were implicated in several types of cancer,

including up-regulation of fibroblast, breast cancer, and neuroblastoma tumors (**Fig. 3B**),

which indicate a certain extent of genetic overlaps and shared pathways that may explain the

intriguing inverse relationship between AD and cancer.[54] Genes associated with the R1

dimension were implicated in pathways involved in the down-regulation of macrophages

(**Fig. 3B**), which are involved in the initiation and progression of various inflammatory

processes, including neuroinflammation and AD.[13] Inflammation is also known to be

associated with vascular compromise and dysfunction. This further concurs with the stronger

cardiovascular profile of R1, especially with increased WML and predominant SPARE-BA

increases. Genes associated with the R2 dimensions were enriched in pathways involved in

AD onset, hippocampus-related brain volumes, and dendritic cells (**Fig. 3B**). In particular,

dendritic cells may regulate amyloid-β-specific T-cell entry into the brain,[55] as well as the

inflammatory status of the brain.[56] The gene set enrichment analysis results are presented in

Supplementary eTable 7.

Genes associated with the R1 and R2 dimensions show potential for drug discovery and

repurposing

We queried whether these 77 genes associated with R1 and R2 are "druggable genes" from

the constructed target-drug-disease network – the target genes express proteins to bind drug-

like molecules, and the drug is at any stage of the clinical trial. For the 49 R1-annotated

genes, 9 genes were targets for 15 drugs and drug-like molecules, treating various cancer,

inflammation, and cardiovascular dysfunctions. For the 40 R2-annotated genes, 6 genes were

targets for 7 drugs developed for diseases of the nervous system, such as Parkinson's (**Fig.**

3C). The pharmacological mechanisms targeted by these identified drugs are largely related

to the pathogenesis of AD in previous literature. For example, FDA-approved Niacin [R1;

target gene: *NNMT*; Anatomical Therapeutic Chemical (ATC) code: C10AD02] is a B

vitamin used to treat various deficiencies and diseases in the cardiovascular system, including

myocardial infarctions,[57] hyperlipidemia,[58] and coronary artery disease.[59] Interestingly, a

recent study[60] showed that Niacin detained AD progression in a 5xFAD mice model. The

niacin receptor HCAR2 modulates microglial response to amyloid deposition, ultimately

alleviating neuronal loss and cognitive decline. Other drugs for potential drug repurposing of

AD are the FDA-approved Docetaxel (R1; target gene: *MAP4*; ATC: L01CD02) and

Paclitaxel (R1; target gene: *MAP4*; ATC: L01CD01), which both target various cancers,

including breast cancer and metastatic prostate cancer. The intriguing inverse relationship

between AD and cancer has long been established, but the underlying shared etiology

remains unclear.[54,61] One hypothesis was that microtubule-associated protein tau – a

pathological biomarker of AD – was associated with resistance to Docetaxel in certain cancer

treatments.[62] In addition, Docetaxel impacted the blood-brain barrier function of breast

cancer brain metastases.[63] Another drug called KM-819 (R2; target gene: *FAF1*) is currently

in Phase 1 for a clinical trial of Parkinson's disease,[64] which aims to suppress α-synuclein-

induced mitochondrial dysfunction,[65] consistent with the mitochondrial hypothesis[66] of AD.

To sum up, R1 and R2 show distinct landscapes of the "druggable genome"[67] on drug

discovery and repurposing[68] for future clinical translation.

The longitudinal rate of change in the R2 dimension, but not R1, is marginally associated with the *APOE ε4* allele, tau in cognitively unimpaired individuals

Using cognitively unimpaired participants from ADNI and BLSA, longitudinal brain association studies showed that the rate of change in the R1 dimension was associated with the change of brain volume in widespread brain regions. In contrast, the rate of change in the R2 dimension was associated with the change of brain volume in the focal medial temporal lobe (**Fig. 4A** and **Supplementary eTable 8** for P-values and effect sizes). This further indicates that the two dominant patterns discovered cross-sectionally also progress in consistent directions longitudinally. The two dimensions were not associated with CSF biomarkers (Aβ42, tau, and p-tau) and the *APOE ε4* allele (*rs429358*) at baseline [$-\log_{10}$(P-value) < 1.31)]. The rate of change of the R2 dimension, but not R1, was marginally [nominal threshold: $-\log_{10}$(P-value) > 1.31] associated with the *APOE ε4* allele, the CSF level of tau, and p-tau (**Fig. 4B** and **Supplementary eTable 9** for P-values and effect sizes), but they did not survive the Bonferroni correction [$-\log_{10}$(P-value) = 2.95]. The longitudinal rate of change of both dimensions was negatively associated [$-\log_{10}$(P-value) > 2.95] with the total CSF level of Aβ42.

We tested these associations using cognitively unimpaired individuals with a high risk of AD based on their family history from the PREVENT-AD cohort. Similarly, at baseline, the two dimensions were not associated with CSF biomarkers or the *APOE ε4* allele (*rs429358*). The longitudinal rate of change in the R2 dimension, but not R1, was marginally [nominal threshold: $-\log_{10}$(P-value) > 1.31] associated with the *APOE ε4* allele [$-\log_{10}$(P-value) = 1.92], the CSF level of tau [$-\log_{10}$(P-value) = 1.65], and p-tau [$-\log_{10}$(P-value) = 1.66].

Longitudinal brain association studies also confirmed the longitudinal progression of the two dimensions in the MCI/AD population (**Supplementary eFigure 11A**). The rates of change in the two dimensions were both associated with *APOE ε4* [$-\log_{10}$(P-value) = 12.54 for R1 and 9.05 for R2] in GWAS (**Supplementary eFigure 11B**), and related to CSF levels of tau [$-\log_{10}$(P-value) = 16.47 for R1 and 9.73 for R2], p-tau [$-\log_{10}$(P-value) = 19.13 for R1 and 10.81 for R2], and Aβ42 [$-\log_{10}$(P-value) = 13.64 for R1 and 13.55 for R2] (**Supplementary eFigure 11C).**

Discussion

The current study leveraged a deep semi-supervised representation learning method to establish two predominant dimensions in the symptomatic MCI/AD population, which were independently found to be expressed, to a lesser degree, in three asymptomatic populations. In particular, the R1 dimension represented a "diffuse-AD" atrophy pattern: varying degrees of brain atrophy throughout the entire brain. In contrast, the R2 dimension showed an "MTL-AD" atrophy pattern: brain atrophy predominantly concentrated in the medial temporal lobe (**Fig. 1A**). Importantly, only R2 was found to be significantly associated with genetic variants of the *APOE* genes in MCI/AD patients. Furthermore, our study examined early manifestations of the R1 and R2 dimensions in asymptomatic populations with varying levels of AD risks and their associations with genetics, amyloid plaques and tau tangles, biological pathways, and body organs. We identified that 24 genomic loci, 14 of which are novel, and 77 annotated genes contribute to early manifestations of the two dimensions. Functional analyses showed that genes unrelated to *APOE* were overrepresented in DEG sets in organs beyond the brain (R1 and R2), including the heart (R1) and the pituitary gland (R2), and enriched in several biological pathways involved in dendritic cells (R2), macrophage functions (R1), and cancer (R1 and R2). Several of these genes were "druggable genes" for cancer (R1), inflammation (R1), cardiovascular diseases (R1), and diseases of the nervous system (R2). Longitudinal findings in the cognitively unimpaired populations showed that the rate of change of the R2 dimension, but not R1, was marginally associated with the *APOE ε4* allele, the CSF level of tau, and Aβ42 (R1 and R2). Our findings suggested that diverse pathologic processes, including cardiovascular risk factors, neurohormone dysfunction, and inflammation, might occur in the early asymptomatic stages, supporting and expanding the current amyloid cascade.[7,8]

AD has been regarded as a CNS disorder. However, increasing evidence has indicated

that the origins or facilitators of the pathogenesis of AD might involve processes outside the

brain.[6] For example, recent findings revealed that gut microbiota disturbances might

influence the brain through the immune and endocrine system and the bacteria-derived

metabolites.[69,70] Our findings support the view that multiple pathological processes might

contribute to early AD pathogenesis and identify non-*APOE* genes in the two dimensions

overrepresented in tissues beyond the brain (e.g., the heart, pituitary gland, muscle, and

kidney). Pathological processes may be involved in different cells, molecular functions, and

biological pathways, exaggerating amyloid plaque and tau tangle accumulation and leading to

the downstream manifestation of neurodegeneration and cognitive decline.

The genetic and clinical underpinnings of the R1 dimension support inflammation, as

well as cardiovascular diseases, as a core pathology contributing to AD.[12,71,72] Genes

associated with the R1 dimension were previously associated with various inflammation-

related clinical traits (**Fig. 2D**), and enriched in biological pathways involved in

immunological response (e.g., up-regulation in macrophages[73], **Fig. 3B**). In addition, genes in

this dimension were overrepresented in DEG sets in the heart (**Fig. 3A**). Previous literature

indicated that inflammation is likely an early step that initiates the amyloidogenic pathway –

the expression of inflammatory cytokines leads to the production of β-amyloid plaques.[13]

Several markers of inflammation are also present in serum and CSF before any indications of

Aβ or tau tangles.[74] For example, clusterin, a glycoprotein involved in many processes and

conditions (e.g., inflammation, proliferation, and AD) induced by tumor necrosis factor

(TNF), was present ten years earlier than Aβ deposition.[75] In addition, the R1 dimension was

also strongly associated with cardiovascular and diabetes biomarkers (**Fig. 2B**). Inflammatory

processes have been critical, well-established risk factors for compromised cardiovascular

function,[76] such as coronary artery disease and the breakdown of the blood-brain barrier. Our

results corroborated the close relationships between AD, cardiovascular diseases, and

inflammation.

The genetic and clinical underpinnings of the R2 dimension support that

neuroendocrine dysfunction might be an early event contributing to the pathogenesis of

AD.[16,17] Genes in the R2 dimension were previously associated with different hormone and

pancreas-related traits from GWAS Catalog (**Fig. 2D**); they were also overrepresented in

DEG in the pituitary and pancreas glands, muscle and kidney (**Fig. 3A**), which are master

glands or key organs in the endocrine system.[77] Previous literature suggested that

neuroendocrine dysfunction might contribute to AD development by secreting
neurohormonal analogs and affecting CNS function.[16] For example, luteinizing hormone-
releasing hormone and follicle-stimulating hormone in serum or neurons were associated
with the accumulation of Aβ plaques in the brain.[17,78],[79] However, early experimental studies
on antagonists of Luteinizing hormone-releasing hormone and growth hormone-releasing
hormone in animal models of AD have shown promising but not entirely convincing
evidence.[16] Taken together, neurodegeneration in the R2 dimension represents an AD-
specific phenotype that might be driven by hormonal dysfunction, leading to rapid
accumulation of amyloid plaques, and was potentially accelerated by the *APOE ε4* allele –
the rate of change in R2, but not R1, was associated with the *APOE ε4* allele in cognitively
unimpaired individuals (**Fig. 4B**).

The hypothesized implications above of the R1 and R2 dimensions on inflammation,
cardiovascular functions, and neuroendocrine dysfunctions are not mutually exclusive and
may collectively contribute to AD pathogenesis. It has been shown that dysregulation of the
hypothalamic-pituitary-gonadal axis is associated with dyotic signaling, modulating the
expression of TNF and related cytokines in systemic inflammation, and the induction of
downstream neurodegenerative cascades within the brain.[80,81] These studies hypothesized that
the neuroendocrine dysfunction and the inflammation mechanism might be the upstream and
downstream neuropathological processes along the disease course of AD.[16] That is, the loss
of sex steroids and the elevation of gonadotropins might lead to a higher level of
inflammatory factors in the brain. Finally, other competing hypotheses may also play a role in
developing AD in early asymptomatic stages, including the mitochondrial hypothesis,[66] the
metabolic hypothesis,[82] and the tau hypothesis.[3]

The NIA-AA framework[83] claims that AD is a continuum in which AD pathogenesis
is initiated in early asymptomatic cognitively unimpaired stages and progresses to amyloid-
positive and tau-positive (A+T+) in late symptomatic stages.[83] Our findings are consistent
with this framework and elucidate the cross-sectional and longitudinal associations of the two
dimensions with genetic and clinical markers from early asymptomatic to late symptomatic
stages. In early asymptomatic stages, the rates of change in the two dimensions are both
associated with amyloid. However, only the R2 dimension, not R1, is marginally associated
with the *APOE ε4* allele and the CSF level of tau (**Fig. 4B**). In contrast, in late symptomatic
stages, the rates of change in the two dimensions are both associated with the *APOE ε4* allele,
CSF levels of tau, p-tau, and amyloid (**Supplementary eFigure 11**). Our findings suggest

that comorbidities or normal aging in R1 may alter the rate or trajectory of neurodegeneration
at early asymptomatic stages, but *APOE*-related genes might play a more pronounced role in
the acceleration and progression during late symptomatic stages for both dimensions (**Fig. 5**).

Several recent studies[84–88], as detailed in an insightful overview by Luo et al.,
collectively provide a comprehensive transcriptomics and epigenomics atlas depicting AD
progression at the single-cell level[89]. These studies highlight the involvement of microglia-
related inflammation, lipid metabolism, and mitochondrial dysfunction. This substantiates the
primary hypothesis in our study: the two dimensions are linked to diverse pathological
mechanisms, encompassing cardiovascular diseases, inflammation, and hormonal
dysfunction, potentially driven by genes beyond *APOE*.

Limitations

This study has several limitations. Firstly, there is a need for longitudinal data from the
general population, as exemplified by the UK Biobank, to provide further validation for the
hypotheses proposed to cover the entire AD spectrum in the same population. Secondly, it is
essential to extend the generalization of the current GWAS findings to include
underrepresented ethnic groups, going beyond the European ancestry populations. Lastly,
substantial methodological advancements are imperative, particularly in the integration of
imaging and genetic data, to effectively capture the full spectrum of AD heterogeneity.

Outlook

In conclusion, the current study used a novel deep semi-supervised representation
learning method to establish two AD dimensions. Our findings support that those diverse
pathological mechanisms, including cardiovascular diseases, inflammation, hormonal
dysfunction, and involving multiple organs,[90–92] collectively affect AD pathogenesis in
asymptomatic stages. These novel biomarkers may serve as instrumental variables to guide
future treatments in the early asymptomatic stages of AD, targeting multi-organ dysfunction
beyond the brain.

Data availability

The GWAS summary statistics corresponding to this study are publicly available on the MEDICINE knowledge portal (http://labs.loni.usc.edu/medicine/) and the BRIDGEPORT web portal (https://www.cbica.upenn.edu/bridgeport/).

Acknowledgements

This research has been conducted using the UK Biobank Resource under Application Number 35148. Data used in the preparation of this article were in part obtained from the Alzheimer's Disease Neuroimaging Initiative (ADNI) database (adni.loni.usc.edu). As such, the investigators within the ADNI contributed to the design and implementation of ADNI and/or provided data but did not participate in the analysis or writing of this report. A complete listing of ADNI investigators can be found at: http://adni.loni.usc.edu/wpcontent/uploads/how_to_apply/ADNI_Acknowledgement_List.pdf. ADNI is funded by the National Institute on Aging, the National Institute of Biomedical Imaging and Bioengineering, and through generous contributions from the following: AbbVie, Alzheimer's Association; Alzheimer's Drug Discovery Foundation; Araclon Biotech; BioClinica, Inc.; Biogen; Bristol-Myers Squibb Company; CereSpir, Inc.; Cogstate; Eisai Inc.; Elan Pharmaceuticals, Inc.; Eli Lilly and Company; EuroImmun; F. Hoffmann-La Roche Ltd and its affiliated company Genentech, Inc.; Fujirebio; GE Healthcare; IXICO Ltd.; Janssen Alzheimer Immunotherapy Research & Development, LLC.; Johnson & Johnson Pharmaceutical Research & Development LLC.; Lumosity; Lundbeck; Merck & Co., Inc.; Meso Scale Diagnostics, LLC.; NeuroRx Research; Neurotrack Technologies; Novartis Pharmaceuticals Corporation; Pfizer Inc.; Piramal Imaging; Servier; Takeda Pharmaceutical Company; and Transition Therapeutics. The Canadian Institutes of Health Research is providing funds to support ADNI clinical sites in Canada. Private sector contributions are facilitated by the Foundation for the National Institutes of Health (www.fnih.org). The grantee organization is the Northern California Institute for Research and Education, and the study is coordinated by the Alzheimer's Therapeutic Research Institute at the University of Southern California. ADNI data are disseminated by the Laboratory for Neuro Imaging at the University of Southern California.

Funding

The primary funding support for this present study is from the initial funding package as an Assistant Professor of Neurology for WJ, provided by Stevens Neuroimaging and Informatics Institute, Keck School of Medicine of USC, University of Southern California. The iSTAGING consortium is a multi-institutional effort funded by NIA by RF1 AG054409 for DC. The Baltimore Longitudinal Study of Aging neuroimaging study is funded by the Intramural Research Program, National Institute on Aging, National Institutes of Health and by HHSN271201600059C. Other supporting grants include 5U01AG068057-02, 1U24AG074855-01. SRH is Supported by multiple grants and contracts from NIH.

Authors' contributions

Dr. Wen has full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Study concept and design: Wen, Davatzikos

Acquisition, analysis, or interpretation of data: Wen, Davatzikos

Drafting of the manuscript: Wen

Critical revision of the manuscript for important intellectual content: all authors

Statistical and genetic analysis: Wen

Competing interests

DAW served as Site PI for studies by Biogen, Merck, and Eli Lilly/Avid. He has received consulting fees from GE Healthcare and Neuronix. He is on the DSMB for a trial sponsored by Functional Neuromodulation. AJS receives support from NIH (P30 AG010133, P30 AG072976, R01 AG019771, R01 AG057739, U01 AG024904, R01 LM013463, R01 AG068193, T32 AG071444, and U01 AG068057 and U01 AG072177). He has also received support from Avid Radiopharmaceuticals, a subsidiary of Eli Lilly (in-kind contribution of PET tracer precursor); Bayer Oncology (Scientific Advisory Board); Eisai (Scientific Advisory Board); Siemens Medical Solutions USA, Inc. (Dementia Advisory Board); Springer-Nature Publishing (Editorial Office Support as Editor-in-Chief, Brain Imaging, and Behavior). ME receives support from multiple NIH grants, the Alzheimer's Association, and the Alzheimer's Therapeutic Research Institute. HJG has received travel grants and speaker honoraria from Fresenius Medical Care, Neuraxpharm, Servier, and Janssen Cilag and research funding from Fresenius Medical Care. HJG had personal contracts approved by the university administration for speaker honoraria and one IIT with Fresenius Medical Care.

Supplementary material

Supplementary material is available at *Brain* online.

References

1. Guthrie, H. *et al.* Safety, Tolerability, and Pharmacokinetics of Crenezumab in Patients with Mild-to-Moderate Alzheimer's Disease Treated with Escalating Doses for up to 133 Weeks. *J Alzheimers Dis* **76**, 967–979 (2020).

2. Sevigny, J. *et al.* The antibody aducanumab reduces Aβ plaques in Alzheimer's disease. *Nature* **537**, 50–56 (2016).

3. Congdon, E. E. & Sigurdsson, E. M. Tau-targeting therapies for Alzheimer disease. *Nat Rev Neurol* **14**, 399–415 (2018).

4. Hardy, J. & Selkoe, D. J. The Amyloid Hypothesis of Alzheimer's Disease: Progress and Problems on the Road to Therapeutics. *Science* **297**, 353–356 (2002).

5. Brinkmalm, G. & Zetterberg, H. The phosphorylation cascade hypothesis of Alzheimer's disease. *Nat Aging* **1**, 498–499 (2021).

6. Du, X., Wang, X. & Geng, M. Alzheimer's disease hypothesis and related therapies. *Translational Neurodegeneration* **7**, 2 (2018).

7. Jack, C. R. *et al.* Tracking pathophysiological processes in Alzheimer's disease: an updated hypothetical model of dynamic biomarkers. *Lancet Neurol* **12**, 207–216 (2013).

8. Frisoni, G. B. *et al.* The probabilistic model of Alzheimer disease: the amyloid hypothesis revised. *Nat Rev Neurosci* **23**, 53–66 (2022).

9. Makin, S. The amyloid hypothesis on trial. *Nature* **559**, S4–S7 (2018).

10. Herrup, K. The case for rejecting the amyloid cascade hypothesis. *Nat Neurosci* **18**, 794–799 (2015).

11. Barisano, G. *et al.* Blood–brain barrier link to human cognitive impairment and Alzheimer's disease. *Nat Cardiovasc Res* **1**, 108–115 (2022).

12. Wyss-Coray, T. Inflammation in Alzheimer disease: driving force, bystander or

beneficial response? *Nat Med* **12**, 1005–1015 (2006).

13. Heneka, M. T. *et al.* Neuroinflammation in Alzheimer's Disease. *Lancet Neurol* **14**, 388–

405 (2015).

14. Leng, F. & Edison, P. Neuroinflammation and microglial activation in Alzheimer disease:

where do we go from here? *Nat Rev Neurol* **17**, 157–172 (2021).

15. Dean, D. W. Neuroendocrine Theory of Aging: Chapter 1. (2012).

16. Schally, A. V. Endocrine approaches to treatment of Alzheimer's disease and other

neurological conditions: Part I: Some recollections of my association with Dr. Abba

Kastin: A tale of successful collaboration. *Peptides* **72**, 154–163 (2015).

17. Xiong, J. *et al.* FSH blockade improves cognition in mice with Alzheimer's disease.

Nature **603**, 470–476 (2022).

18. Dubois, B. *et al.* Preclinical Alzheimer's disease: Definition, natural history, and

diagnostic criteria. *Alzheimers Dement* **12**, 292–323 (2016).

19. Rajpurkar, P., Chen, E., Banerjee, O. & Topol, E. J. AI in health and medicine. *Nat Med*

28, 31–38 (2022).

20. Wen, J. *et al.* Convolutional neural networks for classification of Alzheimer's disease:

Overview and reproducible evaluation. *Medical Image Analysis* **63**, 101694 (2020).

21. Kendler, K. & Neale, M. Endophenotype: a conceptual analysis. *Mol Psychiatry* **15**, 789–

797 (2010).

22. Yang, Z. *et al.* A deep learning framework identifies dimensional representations of

Alzheimer's Disease from brain structure. *Nat Commun* **12**, 7065 (2021).

23. Wen, J. *et al.* Multi-scale semi-supervised clustering of brain images: Deriving disease

subtypes. *Med Image Anal* **75**, 102304 (2021).

24. Young, A. L. *et al.* Uncovering the heterogeneity and temporal complexity of

neurodegenerative diseases with Subtype and Stage Inference. *Nat Commun* **9**, 4273

(2018).

25. Zhang, X. *et al.* Bayesian model reveals latent atrophy factors with dissociable cognitive

trajectories in Alzheimer's disease. *COGNITIVE SCIENCES* 10.

26. Vogel, J. W. *et al.* Four distinct trajectories of tau deposition identified in Alzheimer's

disease. *Nat Med* **27**, 871–881 (2021).

27. Bellenguez, C. *et al.* New insights into the genetic etiology of Alzheimer's disease and

related dementias. *Nat Genet* **54**, 412–436 (2022).

28. Lambert, J.-C. *et al.* Meta-analysis of 74,046 individuals identifies 11 new susceptibility

loci for Alzheimer's disease. *Nat Genet* **45**, 1452–1458 (2013).

29. Yang, Z., Wen, J. & Davatzikos, C. Surreal-GAN:Semi-Supervised Representation

Learning via GAN for uncovering heterogeneous disease-related imaging patterns. *ICLR*

(2021).

30. Petersen, R. C. *et al.* Alzheimer's Disease Neuroimaging Initiative (ADNI): clinical

characterization. *Neurology* **74**, 201–209 (2010).

31. Bycroft, C. *et al.* The UK Biobank resource with deep phenotyping and genomic data.

Nature **562**, 203–209 (2018).

32. Resnick, S. M. *et al.* One-year age changes in MRI brain volumes in older adults. *Cereb*

Cortex **10**, 464–472 (2000).

33. Breitner, J. C. S., Poirier, J., Etienne, P. E. & Leoutsakos, J. M. Rationale and Structure
for a New Center for Studies on Prevention of Alzheimer's Disease (StoP-AD). *J Prev*
Alzheimers Dis **3**, 236–242 (2016).

34. Habes, M. *et al.* The Brain Chart of Aging: Machine-learning analytics reveals links
between brain aging, white matter disease, amyloid burden, and cognition in the
iSTAGING consortium of 10,216 harmonized MR scans. *Alzheimer's & Dementia* **17**,
89–102 (2021).

35. Tustison, N. J. *et al.* N4ITK: improved N3 bias correction. *IEEE Trans. Med. Imaging*
29, 1310–1320 (2010).

36. Doshi, J. *et al.* MUSE: MUlti-atlas region Segmentation utilizing Ensembles of
registration algorithms and parameters, and locally optimal atlas selection. *Neuroimage*
127, 186–195 (2016).

37. Pomponio, R. *et al.* Harmonization of large MRI datasets for the analysis of brain
imaging patterns throughout the lifespan. *Neuroimage* **208**, 116450 (2020).

38. Zhang, X. *et al.* Bayesian model reveals latent atrophy factors with dissociable cognitive
trajectories in Alzheimer's disease. *Proc Natl Acad Sci USA* **113**, E6535–E6544 (2016).

39. Ferreira, D., Nordberg, A. & Westman, E. Biological subtypes of Alzheimer disease: A
systematic review and meta-analysis. *Neurology* **94**, 436–448 (2020).

40. Wen, J. *et al.* Subtyping Brain Diseases from Imaging Data. in *Machine Learning for*
Brain Disorders (ed. Colliot, O.) 491–510 (Springer US, 2023). doi:10.1007/978-1-0716-
3195-9_16.

41. Yang, J., Lee, S. H., Wray, N. R., Goddard, M. E. & Visscher, P. M. GCTA-GREML
accounts for linkage disequilibrium when estimating genetic variance from genome-wide
SNPs. *PNAS* **113**, E4579–E4580 (2016).

42. Zhao, B. *et al.* Genome-wide association analysis of 19,629 individuals identifies variants

influencing regional brain volumes and refines their genetic co-architecture with

cognitive and mental health traits. *Nat Genet* **51**, 1637–1644 (2019).

43. Choi, S. W., Mak, T. S.-H. & O'Reilly, P. F. Tutorial: a guide to performing polygenic

risk score analyses. *Nat Protoc* **15**, 2759–2772 (2020).

44. Davatzikos, C., Xu, F., An, Y., Fan, Y. & Resnick, S. M. Longitudinal progression of

Alzheimer's-like patterns of atrophy in normal older adults: the SPARE-AD index. *Brain*

132, 2026–2035 (2009).

45. Moradi, E., Hallikainen, I., Hänninen, T. & Tohka, J. Rey's Auditory Verbal Learning

Test scores can be predicted from whole brain MRI in Alzheimer's disease. *NeuroImage:*

Clinical **13**, 415–427 (2017).

46. Squire, L. R., Stark, C. E. L. & Clark, R. E. The medial temporal lobe. *Annu Rev*

Neurosci **27**, 279–306 (2004).

47. Bashyam, V. M. *et al.* MRI signatures of brain age and disease over the lifespan based on

a deep brain network and 14 468 individuals worldwide. *Brain* **143**, 2312–2324 (2020).

48. Prins, N. D. & Scheltens, P. White matter hyperintensities, cognitive impairment and

dementia: an update. *Nat Rev Neurol* **11**, 157–165 (2015).

49. Andreasen, N. *et al.* Evaluation of CSF-tau and CSF-Aβ42 as Diagnostic Markers for

Alzheimer Disease in Clinical Practice. *Archives of Neurology* **58**, 373–379 (2001).

50. Buniello, A. *et al.* The NHGRI-EBI GWAS Catalog of published genome-wide

association studies, targeted arrays and summary statistics 2019. *Nucleic Acids Res* **47**,

D1005–D1012 (2019).

51. Bulik-Sullivan, B. K. *et al.* LD Score regression distinguishes confounding from
polygenicity in genome-wide association studies. *Nat Genet* **47**, 291–295 (2015).

52. Jiang, L., Zheng, Z., Fang, H. & Yang, J. A generalized linear mixed model association
tool for biobank-scale data. *Nat Genet* **53**, 1616–1621 (2021).

53. Watanabe, K., Taskesen, E., van Bochoven, A. & Posthuma, D. Functional mapping and
annotation of genetic associations with FUMA. *Nat Commun* **8**, 1826 (2017).

54. Lanni, C., Masi, M., Racchi, M. & Govoni, S. Cancer and Alzheimer's disease inverse
relationship: an age-associated diverging derailment of shared pathways. *Mol Psychiatry*
26, 280–295 (2021).

55. Fisher, Y., Nemirovsky, A., Baron, R. & Monsonego, A. Dendritic cells regulate
amyloid-β-specific T-cell entry into the brain: the role of perivascular amyloid-β. *J*
Alzheimers Dis **27**, 99–111 (2011).

56. Brezovakova, V., Valachova, B., Hanes, J., Novak, M. & Jadhav, S. Dendritic Cells as an
Alternate Approach for Treatment of Neurodegenerative Disorders. *Cell Mol Neurobiol*
38, 1207–1214 (2018).

57. Kong, D. *et al.* Niacin Promotes Cardiac Healing after Myocardial Infarction through
Activation of the Myeloid Prostaglandin D2 Receptor Subtype 1. *J Pharmacol Exp Ther*
360, 435–444 (2017).

58. Crouse, J. R. New developments in the use of niacin for treatment of hyperlipidemia: new
considerations in the use of an old drug. *Coron Artery Dis* **7**, 321–326 (1996).

59. Duggal, J. K. *et al.* Effect of niacin therapy on cardiovascular outcomes in patients with
coronary artery disease. *J Cardiovasc Pharmacol Ther* **15**, 158–166 (2010).

60. Moutinho, M. *et al.* The niacin receptor HCAR2 modulates microglial response and limits disease progression in a mouse model of Alzheimer's disease. *Sci Transl Med* **14**, eabl7634 (2022).

61. Wen, J. *et al.* Novel genomic loci and pathways influence patterns of structural covariance in the human brain. 2022.07.20.22277727 Preprint at https://doi.org/10.1101/2022.07.20.22277727 (2022).

62. Yang, J. *et al.* Microtubule-associated protein tau is associated with the resistance to docetaxel in prostate cancer cell lines. *Res Rep Urol* **9**, 71–77 (2017).

63. Bernatz, S. *et al.* Impact of Docetaxel on blood-brain barrier function and formation of breast cancer brain metastases. *J Exp Clin Cancer Res* **38**, 434 (2019).

64. Shin, W. *et al.* A first-in-human study to investigate the safety, tolerability, pharmacokinetics, and pharmacodynamics of KM-819 (FAS-associated factor 1 inhibitor), a drug for Parkinson's disease, in healthy volunteers. *Drug Des Devel Ther* **13**, 1011–1022 (2019).

65. Kim, B.-S. *et al.* Pharmacological Intervention Targeting FAF1 Restores Autophagic Flux for α-Synuclein Degradation in the Brain of a Parkinson's Disease Mouse Model. *ACS Chem Neurosci* **13**, 806–817 (2022).

66. Swerdlow, R. H., Burns, J. M. & Khan, S. M. The Alzheimer's disease mitochondrial cascade hypothesis: progress and perspectives. *Biochim Biophys Acta* **1842**, 1219–1231 (2014).

67. Hopkins, A. L. & Groom, C. R. The druggable genome. *Nat Rev Drug Discov* **1**, 727–730 (2002).

68. Pushpakom, S. *et al.* Drug repurposing: progress, challenges and recommendations. *Nat Rev Drug Discov* **18**, 41–58 (2019).

69. Jiang, C., Li, G., Huang, P., Liu, Z. & Zhao, B. The Gut Microbiota and Alzheimer's

Disease. *J Alzheimers Dis* **58**, 1–15 (2017).

70. Seo, D., Boros, B. D. & Holtzman, D. M. The microbiome: A target for Alzheimer

disease? *Cell Res* **29**, 779–780 (2019).

71. Heppner, F. L., Ransohoff, R. M. & Becher, B. Immune attack: the role of inflammation

in Alzheimer disease. *Nat Rev Neurosci* **16**, 358–372 (2015).

72. Kinney, J. W. *et al.* Inflammation as a central mechanism in Alzheimer's disease.

Alzheimers Dement (N Y) **4**, 575–590 (2018).

73. Fujiwara, N. & Kobayashi, K. Macrophages in inflammation. *Curr Drug Targets*

Inflamm Allergy **4**, 281–286 (2005).

74. Laurin, D., David Curb, J., Masaki, K. H., White, L. R. & Launer, L. J. Midlife C-

reactive protein and risk of cognitive decline: a 31-year follow-up. *Neurobiol Aging* **30**,

1724–1727 (2009).

75. Thambisetty, M. *et al.* Association of Plasma Clusterin Concentration With Severity,

Pathology, and Progression in Alzheimer Disease. *Archives of General Psychiatry* **67**,

739–748 (2010).

76. Ruparelia, N., Chai, J. T., Fisher, E. A. & Choudhury, R. P. Inflammatory processes in

cardiovascular disease: a route to targeted therapies. *Nat Rev Cardiol* **14**, 133–144

(2017).

77. Vitale, G., Salvioli, S. & Franceschi, C. Oxidative stress and the ageing endocrine

system. *Nat Rev Endocrinol* **9**, 228–240 (2013).

78. Strittmatter, W. J. Old Drug, New Hope for Alzheimer's Disease. *Science* **335**, 1447–

1448 (2012).

79. Strittmatter, W. J. Medicine. Old drug, new hope for Alzheimer's disease. *Science* **335**,

1447–1448 (2012).

80. Clark, I. A. & Atwood, C. S. Is TNF a Link between Aging-Related Reproductive

Endocrine Dyscrasia and Alzheimer's Disease? *JAD* **27**, 691–699 (2011).

81. Clark, I. A., Alleva, L. M. & Vissel, B. The roles of TNF in brain dysfunction and

disease. *Pharmacol Ther* **128**, 519–548 (2010).

82. Demetrius, L. A. & Driver, J. Alzheimer's as a metabolic disease. *Biogerontology* **14**,

641–649 (2013).

83. Jack, C. R. *et al.* NIA-AA Research Framework: Toward a biological definition of

Alzheimer's disease. *Alzheimers Dement* **14**, 535–562 (2018).

84. Gazestani, V. *et al.* Early Alzheimer's disease pathology in human cortex involves

transient cell states. *Cell* **186**, 4438-4453.e23 (2023).

85. Xiong, X. *et al.* Epigenomic dissection of Alzheimer's disease pinpoints causal variants

and reveals epigenome erosion. *Cell* **186**, 4422-4437.e21 (2023).

86. Sun, N. *et al.* Human microglial state dynamics in Alzheimer's disease progression. *Cell*

186, 4386-4403.e29 (2023).

87. Dileep, V. *et al.* Neuronal DNA double-strand breaks lead to genome structural variations

and 3D genome disruption in neurodegeneration. *Cell* **186**, 4404-4421.e20 (2023).

88. Mathys, H. *et al.* Single-cell atlas reveals correlates of high cognitive function, dementia,

and resilience to Alzheimer's disease pathology. *Cell* **186**, 4365-4385.e27 (2023).

89. Luo, W., Qu, W. & Gan, L. The AD odyssey 2023: Tales of single cell. *Cell* **186**, 4257–

4259 (2023).

90. Tian, Y. E. *et al.* Biological aging of human body and brain systems. 2022.09.03.22279337 Preprint at https://doi.org/10.1101/2022.09.03.22279337 (2022).

91. Zhao, B. *et al.* Heart-brain connections: Phenotypic and genetic insights from magnetic resonance images. *Science* **380**, abn6598 (2023).

92. Wen, J. *et al.* The Genetic Architecture of Biological Age in Nine Human Organ Systems. *medRxiv* 2023.06.08.23291168 (2023) doi:10.1101/2023.06.08.23291168.

93. Liberzon, A. *et al.* Molecular signatures database (MSigDB) 3.0. *Bioinformatics* **27**, 1739–1740 (2011).

94. Wishart, D. S. *et al.* DrugBank 5.0: a major update to the DrugBank database for 2018. *Nucleic Acids Res* **46**, D1074–D1082 (2018).

95. Zhou, Y. *et al.* Therapeutic target database update 2022: facilitating drug discovery with enriched comparative data of targeted agents. *Nucleic Acids Res* **50**, D1398–D1407 (2022).

Figure legends

Figure 1: **The manifestation of the R1 and R2 dimensions of brain atrophy in the MCI/AD population.**

A) Brain association studies reveal two dominant brain atrophy dimensions. A linear regression model was fit to the 119 GM ROIs at baseline for the R1 and R2 dimensions. The $-\log_{10}$(P-value) of each significant ROI (Bonferroni correction for the number of 119 ROIs: $-\log_{10}$(P-value) > 3.38) is shown. A negative value denotes brain atrophy with a negative coefficient in the linear regression model. P-value and effect sizes (r, Pearson's correlation coefficient) are presented in **Supplementary eTable 1**. The range of r for each dimension is also shown. Of note, the sample size (N) for R1 and R2 is the same for each ROI. **B)** Genome-wide association studies demonstrate that the R2, but not R1, dimension is associated with variants related to *APOE* genes (genome-wide P-value threshold with the red line: $-\log_{10}$(P-value) > 7.30). We associated each common variant with R1 and R2 using the whole-genome sequencing data from ADNI. Gene annotations were performed via positional, expression quantitative trait loci, and chromatin interaction mappings using FUMA.[53] We then manually queried whether they were previously associated with AD-related traits in the GWAS Catalog.[50] Red-colored loci/genes indicate variants associated with AD-related traits in previous literature. **C)** Clinical association studies show that the R2 dimension is associated to a larger extent with AD-specific biomarkers, including SPARE-AD[44], an imaging surrogate to AD atrophy patterns, and *APOE ε4,* the well-established risk allele in sporadic AD. The R1 dimension is associated to a larger extent with aging (e.g., SPARE-BA,[47] an imaging surrogate for brain aging) and vascular-related biomarkers (e.g., WML, white matter lesion). The same linear regression model was used to associate the R1 and R2 dimensions with the 45 clinical variables, including cognitive scores, modifiable risk factors, CSF biomarkers, disease/condition labels, demographic variables, and imaging-derived phenotypes. The radar plot shows representative clinical variables; results for all 45 clinical

variables are presented in **Supplementary eTable 3**. The SPARE-AD and SPARE-BA scores
are rescaled for visualization purposes. The gray-colored circle lines indicate the P-value
threshold in both directions (Bonferroni correction for the 45 variables: $-\log_{10}$(P-value) >
2.95). A positive/negative $-\log_{10}$(P-value) value indicates a positive/negative correlation
(beta). The transparent dots represent the associations that do not pass the Bonferroni
correction; the blue-colored dots and red-colored dots indicate significant associations for the
R1 and R2 dimensions, respectively.

Figure 2: **The expression of the R1 and R2 dimensions in the general population. A)**

Brain association studies confirm the presence of the two dimensions in the general
population: the R1 dimension shows widespread brain atrophy, whereas the R2 dimension
displays focal medial temporal lobe atrophy. P-value and effect sizes (*r*, Pearson's correlation
coefficient) are presented in **Supplementary eTable 4**. The range of *r* for each dimension is
also shown. Of note, the sample size (*N*) for R1 and R2 is the same for each ROI. **B)** Clinical
association studies further show that the R2 dimension is associated with prospective

memory, and the R1 dimension is associated with several cognitive dysfunctions, cardiovascular risk factors (e.g., triglycerides), and diabetes (e.g., HbA1c). The same linear regression models were used to associate the R1 and R2 dimensions with the 61 clinical variables, including cardiovascular factors, diabetic blood markers, social demographics, lifestyle, physical measures, cognitive scores, and imaging-derived phenotypes. The radar plot shows representative clinical variables; all other results are presented in **Supplementary eTable 5**. The gray circle lines indicate the P-value threshold in both directions (Bonferroni correction for the 61 variables: $-\log_{10}$(P-value) > 3.08). A positive/negative $-\log_{10}$(P-value) value indicates a positive/negative correlation (beta). Transparent dots represent the associations that do not pass the Bonferroni correction; the blue-colored dots and red-colored dots indicate significant associations for the R1 and R2 dimensions, respectively. **C)** Genome-wide association studies demonstrate that the R2 dimension is associated to a larger extent with genomic loci and genes previously associated with AD-related traits in the literature (genome-wide P-value threshold with the red line: $-\log_{10}$(P-value) > 7.30). Each genomic locus is represented by its top lead SNP. The R1 dimension identified 8 (blue-colored in bold) out of the 49 mapped genes associated with AD-related traits. The R2 dimension identified 13 (red-colored in bold) out of 40 mapped genes associated with AD-related traits. Gene annotations were performed via positional, expression quantitative trait loci, and chromatin interaction mappings using FUMA (**Supplementary eTable 6** for all mapped genes).[53] The genomic loci and mapped genes were manually queried in the GWAS Catalog[50] to determine whether they were previously associated with AD (novel or not). * denotes that the genomic locus is novel. **D)** Besides AD-related traits, the genes and genomic loci in the two dimensions were also associated with other clinical traits, including inflammation, neurohormones, and imaging-derived phenotypes, in the literature from the GWAS Catalog.[50] The flowchart first maps the genomic loci and genes (left) identified in the two dimensions onto the human genome (middle). It then links these variants to any clinical traits identified in previous literature from the GWAS Catalog (right). In the middle of the human genome, we show chromosomes 1 to 22 (above to below); the blue and red-colored genes are AD-related for the R1 and R2 dimensions, respectively. The black-colored genes (Fig. C) are not annotated. INF: inflammation; PD: psychiatric disorder; PM: physical measure; Novel (pink-colored in bold, corresponding to the novel loci/genes in Fig C) indicates that the locus or gene was not associated with any traits in the literature.

Figure 3: **Tissue specificity and biological pathway enrichment analysis of the R1 and R2 dimensions in the general population.**

A) Tissue specificity analyses show that genes associated with the two dimensions of neurodegeneration are overrepresented in organs/tissues beyond the human brain (R1 and R2). The unique overrepresentation of genes in differentially expressed gene sets (DEG) in the heart (R1) and the pituitary gland, muscle, and kidney (R2) may imply the involvement of inflammation[12,71,72] and neurohormone dysfunction[15–17], respectively. The *GENE2FUNC*[53]

pipeline from FUMA was performed to examine the overrepresentation of prioritized genes
(Fig. 2C) in pre-defined DEGs (up-regulated, down-regulated, and both-side DEGs) from
different gene expression data. The input genes (**Fig. 2C**) were tested against each DEG using
the hypergeometric test. We present only the organs/tissues that passed the Bonferroni
correction for multiple comparisons. **B)** Gene set enrichment analysis shows that genes
associated with the two dimensions are enriched in different biological pathways. For
example, genes associated with the R1 dimension are implicated in down-regulated
macrophage functions, which have been shown to be associated with inflammation.[13] In
contrast, the R2 dimension is enriched in AD hallmarks (e.g., hippocampus), AD-related gene
sets, and the pathway involved in dendritic cells, which may regulate amyloid-β-specific T-
cell entry into the brain.[55] Both dimensions are enriched in gene sets involved in cancer,
which may indicate overlapped genetic underpinnings between AD and cancer.[54] The
GENE2FUNC[53] pipeline from FUMA was performed to examine the enrichment of
prioritized genes (**Fig. 2C**) in pre-defined gene sets. Hypergeometric tests were performed to
test whether the input genes were overrepresented in any pre-defined gene sets. Gene sets
were obtained from different sources, including MsigDB[93] and GWAS Catalog.[50] We show
the significant results from gene sets defined in the GWAS Catalog, curated gene sets, and
immunologic signature gene sets. All results are shown in **Supplementary eTable 7. C)** The
target-drug-disease network for R1 and R2-associated genes provides great potential for drug
discovery and repurposing. R1-annotated "druggable genes" were developed for
cardiovascular diseases, various cancers, and inflammation, whereas R2-annotated
"druggable genes" were developed for diseases of the nervous system (e.g., Parkinson's
disease). For the target-drug-disease network, the 5th level of the Anatomical Therapeutic
Chemical (ATC) code is displayed for the DrugBank database,[94] and the disease name
defined by the International Classification of Diseases (ICD-11) code is showed for the
Therapeutic Target Database.[95]

970
971

Figure 4: **The longitudinal rate of change in R1 and R2 in the cognitively unimpaired**
population.

A) Longitudinal brain association studies show that the R1 dimension exhibits longitudinal
brain volume decrease in widespread brain regions, whereas the R2 dimension displays
longitudinal brain volume decrease in the focal medial temporal lobe. We first derived the
rate of change of the 119 GM ROIs and the R1 and R2 dimensions using a linear mixed effect
model; a linear regression model was then fit to the rate of change of the ROIs, R1, and R2 to
derive the beta coefficient value of each ROI. A negative value denotes longitudinal brain
changes with a negative coefficient of the rate of change in the linear regression model. P-
value and effect sizes (r, Pearson's correlation coefficient) are presented in **Supplementary**
eTable 8. The range of r for each dimension is also shown. Of note, the sample size (N) for
R1 and R2 is the same for each ROI. **B)** The rate of change, not the baseline measurement, in
the two dimensions is negatively associated with the CSF level of Aβ42 (Bonferroni
correction for the 45 variables: $-\log_{10}$(P-value) > 2.95). The rate of change in the R2
dimension, not the R1 dimension, was marginally ($-\log_{10}$(P-value) > 1.31) associated with
the CSF level of tau and p-tau, and *APOE ε4*. All other clinical associations are presented in
Supplementary eTable 9. The gray-colored circle lines indicate different P-value thresholds
in both directions (Bonferroni correction for the 45 variables: $-\log_{10}$(P-value) > 2.95 and the
nominal P-value threshold: $-\log_{10}$(P-value) > 1.31). A positive/negative $-\log_{10}$(P-value) value
indicates a positive/negative correlation (beta). Transparent dots represent the associations
that do not pass the nominal P-value threshold [$\log_{10}$(P-value) = 1.31]; the blue-colored dots
and red-colored dots indicate significant associations [$\log_{10}$(P-value) > 1.31] for the R1 and
R2 dimensions, respectively.

996
997

Figure 5: Genes unrelated to *APOE* influence early manifestations of R1 and R2.

Genes unrelated to *APOE* and overrepresented in organs beyond the human brain are
associated with early manifestations of the R1 (diffuse-AD) and R2 (MTL-AD) dimensions,
which capture the heterogeneity of AD-related brain atrophy. For visualization purposes, we
display the two genes with the highest expression values in the tissue specificity analyses for
each organ/tissue. The black arrow line emulates the longitudinal progression trajectory along
these two dimensions. The positions of beta-amyloid, tau, and *APOE* (increasing *APOE*-
mediated progression) indicate the time point when they are associated with the two
dimensions. The blue/red gradient-color background indicates a higher influence of *APOE*-
related genes (left to right; early to late stages). The brain atrophy patterns are presented in
the 3D view. In early asymptomatic stages, the R1-related genes are implicated in
cardiovascular diseases and inflammation; the R2-related genes are involved in hormone-
related dysfunction. Critically, longitudinal progression of the dimension demonstrates an
impact of the *APOE* genes in early asymptomatic stages in R2, but this longitudinal effect
occurs only in late symptomatic stages in R1. These results suggest that comorbidities (e.g.,
cardiovascular conditions) or normal aging in R1 may alter or delay the trajectory of
neurodegeneration in early asymptomatic stages; *APOE*-related genes may play a pronounced
role in the acceleration and progression in late symptomatic stages for both dimensions. Of
note, the underlying pathological processes that initiate and drive the progression of the two
dimensions are not mutually exclusive. Hence, both R1 and R2 can be co-expressed in the
same individual. In addition, the two dimensions can also be affected by other AD
hypotheses, such as the mitochondrial hypothesis[66] and the metabolic hypothesis.[82] MTL:
medial temporal lobe.

1022

1023