Abnormal intrinsic neural timescale in Parkinson's disease

Yarui Wei,[1, 3-8#*] Chunyan Zhang,[1, 3-8#] Yuanyuan Peng,[1, 3-9] Chen Chen,[1, 3-8] Shaoqiang Han,[1, 3-8] Weijian Wang,[1, 3-8] Yong Zhang,[1, 3-8] Hong Lu,[2*] Jingliang Cheng[1, 3-8*]

[1]Department of Magnetic Resonance Imaging, The First Affiliated Hospital of Zhengzhou University, Zhengzhou 450052, China

[2]Department of Neurology, The First Affiliated Hospital of Zhengzhou University, Zhengzhou 450052, China

[3]Key Laboratory for Functional Magnetic Resonance Imaging and Molecular Imaging of Henan Province, Zhengzhou 450052, China

[4]Engineering Technology Research Center for Detection and Application of Brain Function of Henan Province, Zhengzhou 450052, China

[5]Engineering Research Center of Medical Imaging Intelligent Diagnosis and Treatment of Henan Province, Zhengzhou 450052, China

[6]Key Laboratory of Magnetic Resonance and Brain Function of Henan Province, Zhengzhou 450052, China

[7]Key Laboratory of Brain Function and Cognitive Magnetic Resonance Imaging of Zhengzhou, Zhengzhou 450052, China

[8]Key Laboratory of Imaging Intelligence Research Medicine of Henan Province, Zhengzhou 450052, China

[9]Imaging Center, Luoyang Orthopedic-Traumato Logical Hospital of Henan Province, Henan Provincal Orthopedic Hospital, Luoyang 471002, China

[#] **These authors have contributed equally to this work.**

[*]**Correspondence to:**

Yarui Wei

Department of Magnetic Resonance Imaging, The First Affiliated Hospital of Zhengzhou University, Zhengzhou 450052, China

E-mail: yarui_wei@163.com

Jingliang Cheng

Department of Magnetic Resonance Imaging, The First Affiliated Hospital of Zhengzhou University, Zhengzhou 450052, China

E-mail: fccchengjl@zzu.edu.cn

Hong Lu

Department of Neurology, The First Affiliated Hospital of Zhengzhou University

Zhengzhou 450052, China

E-mail: zzu_luhong@163.com

Abstract

Background: The intrinsic neural timescale in a local brain area has been specialized, and the sensory and prefrontal areas have shorter and longer timescales, respectively. Parkinson's disease (PD) is a progressive neurological disorder that also exhibits with abnormal neural activity in sensorimotor, visual, and cognitive-related areas. However, it was unclear whether PD patients exhibited abnormal intrinsic timescales and whether PD patients exhibited different temporal feature at different stages.

Methods: Here, we estimated the intrinsic timescales using the magnitude of the autocorrelation of intrinsic neural signals by resting state functional magnetic

resonance imaging data in 74 PD patients, including 44 patients in the early stage (PD-ES) and 30 patients in the late stage (PD-LS), and 73 healthy controls (HC).

Results: We identified that the PD group had shorter timescales in bilateral lingual and calcarine gyri, bilateral postcentral and precentral gyri, and the right middle cingulum gyrus than the HC group, which correlated with the age, but not in the HC group. Moreover, shorter timescales in the PD group also correlated with the symptom severity and reduced grey matter volume (GMV). Longer timescale in the right middle frontal gyrus was also found in the PD group than that in the HC group. Increasingly, compared to the HC group, the PD-ES group had longer timescales in the anterior cortical regions, including the left superior and inferior frontal gyri, and the right middle frontal gyrus, whereas the PD-LS group had shorter timescales in the posterior cortical regions, including bilateral calcarine and lingual gyri, which accompanied with more serious cognitive impairments and reduced GMV.

Conclusions: Our findings suggest that PD patients exhibit abnormal intrinsic timescales in visual, sensorimotor, and cognitive systems, which might provide new insights for the neural substrate of PD. Distinct patterns of intrinsic timescales in cerebral cortex between the PD-LS and PD-ES groups indicate that intrinsic timescale might be a new neuroimaging biomarker across disease stage in PD.

Keywords: intrinsic timescale; Parkinson's disease; sensorimotor cortex; visual system; cognitive system

Introduction

Specialization and hierarchy are organizing principles for primate cortex, and cortical areas are also specialized in the temporal domain. The neural timescale in a local brain area reflects the function of that area in many trials of a task[1,2] or even in the absence of direct stimulus processing[3,4]. Moreover, the intrinsic neural timescales at resting state could predict the neural activity of the brain region in a task,[5-7] and the

sensory and prefrontal areas exhibit shorter and longer timescales at resting state, respectively[3]. Recently, the intrinsic timescales were also verified by using multiple neuroimaging methods, such as single-neuron level[5], electroencephalography (EEG)[8], magnetoencephalogram (MEG),[7] and functional magnetic resonance imaging (fMRI)[4,9-11]. Abnormal intrinsic timescales were also found in some neurological or mental diseases, such as schizophrenia,[12,13] autism,[10] epilepsy,[14] and depression,[15] which showed that intrinsic timescales might bring new insights about abnormal local neural dynamics on the brain of patients with some diseases. However, it's unclear whether abnormal intrinsic timescales were exhibited in patients with Parkinson's disease (PD).

PD is a progressive neurological disorder characterized by tremor, rigidity, and slowness of movements and abnormal sensorimotor integration.[16,17] Nonmotor symptoms, such as visual dysfunction (retinal deficits, facial emotion recognition deficits, and visual hallucinations)[18,19] and cognitive impairment[20,21], also occur frequently in PD. Resting state fMRI studies also showed abnormal neural activity or functional connection in motor[22-25] and nonmotor[26-29] areas. Moreover, studies with somatosensory evoked potentials (SEPs),[30] prepulse inhibition,[31] and event-related potentials[32] support the hypothesis of central abnormalities of sensorimotor integration in PD. Importantly, numerous studies showed that patients with PD with cognitive impairments exhibited changes of brain structure and function.[26,27,29,33] In addition, visual-related areas also showed abnormal neural activation, and structural and functional changes in PD.[34-36] However, whether abnormal neural information stored in sensorimotor, visual, and cognitive-related areas was also unclear.

Here, we assessed whether abnormal intrinsic timescales were exhibited in the brain of PD patients. In line with the previous studies,[10,11] the present study also estimated the intrinsic timescales by using the magnitude of the autocorrelation of intrinsic neural signals in PD patients. To my best knowledge, no study has investigated the intrinsic timescales of neural activity in PD. Moreover, we also explored the correlation between abnormal intrinsic timescales in PD patients and clinical symptoms. Because the previous studies found abnormal intrinsic timescales

confirmed by age and significant correlation between abnormal intrinsic timescales and the structural atypicality in autism,[10] and the factor of age[37-39] and structural changes of the brain[40-42] in PD are also important. Therefore, we also explored the correlations between abnormal intrinsic timescales and the age/structure information in PD patients. Finally, because PD is a progressive neurodegenerative disease and imaging biomarkers in PD are increasingly important for monitoring progression in clinical trials,[43,44] we also explored the differences of intrinsic timescales among patients with PD in the early stage (PD-ES) and the late stage (PD-LS) and healthy controls (HC).

Results

Demographic and head-motion data

For the PD and HC groups, no significant between-group difference in age (Z = -0.240, P = 0.810) or sex (X^2 = 0.819, P = 0.365) was found (Table 1). For the PD-ES and PD-LS groups, significant between-group differences in age (Z = -2.282, P = 0.023), sex (X^2 = 29.480, P = 3.968 × 10^{-7}), and the scores of UPDRS (Z = -5.788, P = 7.120 × 10^{-9}), HY scale (Z = -7.420, P = 1.173 × 10^{-13}), MoCA (Z = 2.166, P = 0.030) and MMSE (Z = 2.811, P = 0.005) were found (Supplementary table 1).

Abnormal intrinsic timescales between the PD and HC groups

The mean intrinsic timescale maps in the PD and HC groups were presented in Fig. 1. Both PD and HC groups had a similar whole-brain pattern of intrinsic timescales: longer timescales in frontal and parietal cortices and shorter timescales in sensorimotor, visual, and auditory areas (Fig. 1, *upper*).

Two-sample t test revealed that the PD group had shorter timescales in the bilateral lingual gyri, bilateral postcentral gyri, and right middle cingulum gyrus compared with the HC group (Supplementary table 2, Fig. 2). The upper first two

clusters also included bilateral calcarine gyri and bilateral precentral gyri. Longer timescale in the right middle frontal gyrus was also found in the PD group than the HC group (Supplementary table 2, Fig. 2).

Correlations among intrinsic timescale, symptom severity, and age in the PD and HC groups

The intrinsic timescale of the bilateral lingual gyri in the PD group negatively correlated with the scores of the Hoehn and Yahr scale (Spearman's rho = -0.265, P = 0.022, Fig. 3). Moreover, the intrinsic timescales in bilateral lingual gyri (Spearman's rho = -0.331, P = 0.004), the right postcentral gyrus (Spearman's rho = -0.262, P = 0.024), and the right middle cingulum gyrus (Spearman's rho = -0.338, P = 0.003) negatively correlated the age of the PD group, respectively; the older the age, the shorter the intrinsic timescale (Fig. 4, *upper*). In HC group, no significant correlation result was found between the intrinsic timescale and the age in any cluster from corrected statistical intrinsic timescale map. And we also found that the age in the PD group negatively correlated with the scores of MoCA (Spearman's rho = -0.372, P = 0.004), and MMSE (Spearman's rho = -0.371, P = 0.001) (Fig. 4, *lower*). However, this significance did not remain after Bonferroni correction (P < 0.05/106 = 0.0005).

Grey matter volume and intrinsic timescale in the PD and HC groups

No significant difference of GMV was found between the PD and HC groups. However, we extracted the mean GMV values of all voxels within each cluster from corrected statistical intrinsic timescale map and found positively significant correlation between the intrinsic timescale and the GMV in the right postcentral gyrus (Spearman's rho = 0.254, P = 0.029, Fig. 5). However, this significance did not remain after Bonferroni correction (P < 0.05/106 = 0.0005). In HC group, no significant correlation results were found between the intrinsic timescale and the GMV.

Abnormal intrinsic timescales in PD at different stages

The mean intrinsic timescale maps for the PD-ES and PD-LS groups were also presented in Fig. 1. Both PD-ES and PD-LS groups also showed a similar whole-brain pattern of intrinsic timescales like the PD and HC groups (Fig. 1, *lower*).

The one-sample ANOVA revealed significant between-group differences in intrinsic timescales of bilateral lingual gyri, the left precuneus, and the right middle cingulum gyrus (Supplementary table 3, Fig. 6). The upper first two clusters also included bilateral calcarine gyri. Post hoc comparisons using Bonferroni's test showed that the PD-LS group had shorter timescale in the left precuneus than that in the PD-ES group and shorter timescales in bilateral calcarine and lingual gyri than that in the HC group (Supplementary table 3, Fig. 6). Increasingly, longer timescales in the left superior frontal gyrus, the left inferior frontal gyrus, the right middle frontal gyrus were also found in the PD-ES group than the HC group (Supplementary table 3, Fig. 6). However, no significant correlation was found among the intrinsic timescale, symptom severity, and age in the PD-ES, PD-LS, or HC group.

Grey matter volume and intrinsic timescale in PD at different stages

No significant difference of GMV was found among the PD-ES, PD-LS, and HC groups. However, we extracted the mean GMV values of all voxels within each cluster from corrected statistical intrinsic timescale maps and found significant differences of mean GMV values between groups in all the upper regions (Table 2). Then Dunn's test performed, and we found that: (1) compared to the PD-ES and HC groups, the PD-LS group had reduced GMV in the bilateral lingual and calcarine gyri, and the left precuneus (Table 2). Moreover, compared to the PD-ES group, the PD-LS group also had reduced GMV in the right middle cingulum gyrus, the left superior frontal gyrus, the left inferior frontal gyrus, the right middle frontal gyrus (Table 2). However, no significant correlation was found between the intrinsic timescale and the GMV in the PD-ES, PD-LS, or HC group.

Discussion

In this study, we explored the intrinsic timescales, which relates to the functional hierarchy of the brain, in PD patients. By analyzing the time-dependent magnitude of the autocorrelation function in resting state fMRI data, we found that the PD group had shorter timescales in bilateral lingual and calcarine gyri, bilateral postcentral and precentral gyri, and the right middle cingulum gyrus, which was correlated with the age. Moreover, shorter timescale in the PD group correlated with the symptom severity and reduced GMV. Longer timescale in the right middle frontal gyrus were also found in the PD group. Increasingly, compared to the HC group, the PD-ES group had longer timescales in the anterior cortical regions, whereas the PD-LS group had shorter timescales in the posterior cortical regions, which accompanied with more serious cognitive impairment and reduced local GMV. The findings indicate that patients with PD exhibited abnormal intrinsic timescales in visual, sensorimotor, and cognitive systems and distinct patterns of intrinsic timescales in cerebral cortex in PD at different stages, which might provide new insights for the neural substrate of PD.

Our findings showed abnormal temporal property of local neural activity in the visual, sensorimotor, and cognitive systems in PD, which was consistent with the previous studies. For example, the previous fMRI studies that calculated the variability of brain signal across time found abnormal variance in visual[47-49], sensorimotor[47,50], and frontal[51,52] regions in PD. Some studies also found that the symptom severity of PD significantly correlated with such abnormal variability in the visual and sensorimotor regions.[49,50] We speculate that if abnormal signal variability in these regions indicates more random activity of the brain and then result in weaker autocorrelation, which can be interpreted the current findings. Moreover, abnormal intrinsic timescales in multisystem also refer to abnormal functional hierarchy of the brain in individuals with PD. Therefore, the intrinsic timescale might be a suitable index to explore the neural substrate of PD.

Furthermore, using the index of the intrinsic timescale, we also found that distinct patterns of intrinsic timescales in cerebral cortex in PD at different stages. The

cognitive impairment of the PD-LS group was more severity than that of the PD-ES group. Distinct patterns of local cerebral glucose metabolism were also found in PD with and without mild cognitive impairment (PDNC and PDMCI): the PDNC patients had limited areas of hypometabolism in the frontal and occipital cortices, and the PDMCI patients had extensive areas of hypometabolism in the posterior cortical regions, including the temporo-parieto-occipital junction, medial parietal, and inferior temporal cortices.[53] Hirano *et al.*[54] suggested that the frontal cortex was associated with impulse control disorders and that posterior brain areas were related to cognitive decline in PD. Distinct patterns of intrinsic timescales between the PD-ES and PD-LS groups also indicate that intrinsic timescale might be a new neuroimaging biomarker across disease stage in PD, which might have the potential to improve clinical care and management.

It is reasonable that significant correlations between cognitive impairments and the age were found in PD, because PD is one of the most common age-related neurodegenerative disorders[55]. It's well known that ageing remains the biggest risk factor for developing idiopathic PD and that researches[56,57] into the mechanisms leading to cell death in Parkinson's disease has shed light on many aspects of the pathogenesis of PD. Reeve *et al.*[55] explained why advancing age is the biggest risk factor in PD, and suggested that changes in protein degradation pathways and mitochondrial DNA mutations might result in neural loss. Moreover, we also found that significantly negative association between the age and the intrinsic timescales in the lingual gyrus, the postcentral gyrus, and the middle cingulate gyrus, which also indicate that the intrinsic timescale may act as another biomarker of PD. The present study helps us to understand that age-related neural factors predispose some individuals to develop this common neurodegenerative disease.

The GMV and intrinsic timescales might be independent and complementary measures for PD. On the one hand, reduced GMV of bilateral lingual and calcarine gyri, the left precuneus, the right middle cingulate gyrus, and bilateral frontal gyri were found in the PD-LS group, but not in the PD-ES group or in the combined group including all patients at different stages, which could explain the result of no

significant difference of GMV between the PD and HC groups. However, the PD-LS group had shorter intrinsic timescales in bilateral lingual and calcarine gyri, and the left precuneus, and the PD-ES group had higher intrinsic timescales in bilateral frontal gyri. Therefore, changes of the GMV in the upper brain regions were consistent and stable for PD patients, and changes of intrinsic timescales were different for PD patients, which might indicate abnormal functional hierarchy in PD. On the other hand, shorter intrinsic timescale in the lingual gyrus correlated with reduced GMV in PD patients, the latter might be the structure base of abnormal temporal pattern. Therefore, a combination of the GMV and intrinsic timescale approaches may provide more evidence for the neural substrate of PD.

Several limitations should be noted when interpreting our findings. First, although we totally recruited 74 PD patients and 73 HC, the sample size is small. Maybe this is the reason why no significant correlation between abnormal intrinsic timescales and symptom/age/GMV was found in PD at different stages. Abnormal intrinsic timescales in the sensorimotor regions were not found in PD at different stages, which might also originate from the small sample size. The future study should increase the sample size to make the findings more reliable. Second, since abnormal intrinsic timescales of visual regions were obvious in PD, especially in the PD-LS group, detailed information about vision of the participants should have been collected to ensure whether temporal feature in visual regions associated with visual dysfunction. Abnormal neural activity of the visual regions and the symptoms of visual hallucination and other visual dysfunction were also found in PD.[34,58] Third, significant difference of sex was found between the PD-ES and PD-LS groups. However, PD patients were not always ready for recruitment, and their availability would have been reduced even further if the factor of the sex had been applied. The further study should match the factor of the sex between the PD-ES and PD-LS groups, which makes the findings more reliable.

Overall, our findings suggest that PD patients exhibit abnormal intrinsic timescales in visual, sensorimotor, and cognitive systems, which might provide new insights for the neural substrate of PD. Distinct patterns of intrinsic timescales in

cerebral cortex in PD patients at early and late stages indicate that intrinsic timescale might be a new neuroimaging biomarker across disease stage in PD.

Materials and methods

Participants

A total of 74 PD patients (41 females) and 73 age- and sex-matched HC (35 females) were included in the study (Table 1). The PD patients were all from Department of Neurology in the First Affiliated Hospital of Zhengzhou University. The clinical diagnosis of patients with PD was made by two experienced neurologists based on the clinical diagnostic criteria for PD introduced by the International Movement Disorder Society (MDS),[45] and all patients were diagnosed as idiopathic PD. PD patients were assessed the severity of motor symptoms and cognitive impairments by using the unified Parkinson's disease rating scale (UPDRS), a revised severity classification Hoehn-Yahr (HY) scale, Montreal cognitive assessment (MoCA), and mini-mental state examination (MMSE). Based on HY scale, 74 patients with PD were split into 44 patients with PD-ES (the scores of the HY scale were ≤2) and 30 patients with PD-LS (the scores of the HY scale were >2) (Supplementary table 1). Moreover, we collected UPDRS data for 72 of the patients, and collected HY scale data for all patients, and collected MoCA data for 57 of the patients, and collected MMSE data for 71 of the patients. The scores of MoCA and MMSE lower than 25 considered cognitive impairment.[46] Exclusion criteria for the patients were as follows: (1) secondary or atypical parkinsonism; (2) history of other neurological diseases (cerebrovascular disease, seizures, brain surgery, brain tumors, moderate to severe head trauma, or hydrocephalus); (3) history of psychiatric illness; (4) use of antipsychotics or other medicines that may affect clinical assessment; (5) alcohol or drug abuse; (6) contraindications for MRI. The HC were healthy middle-aged and elderly people recruited from across Henan Province. Exclusion criteria for HC were as follows: (1) history of neurological or psychiatric illness; (2) alcohol or drug abuse;

(3) contraindications for MRI. The present research was approved by the Ethics Committee of the First Affiliated Hospital of Zhengzhou University. All subjects or their guardians signed the informed consent forms according to the Declaration of Helsinki. Because we performed the Shapiro-Wilk test to check the normality of demographic and clinical data for each group, and the results showed that the factor of demographic and clinical data was not normally distributed for each group. Therefore, Wilcoxon signed-rank test were used to compare demographic and clinical data between groups and X^2 test were used to compare sex data between groups with SPSS software, respectively.

Data acquisition

All subjects were scanned using one of two 3T MAGNETOM Prisma scanners (Siemens Healthcare, Erlangen, Germany), with a 64-channel receiver array head coil. Head motion and scanner noise were reduced using foam paddings and earplugs, respectively. All participants were asked to remain alert, not to move, not to sleep with eyes closed, not to think systematically. Structural images were acquired using a 3D prototypic T1 magnetization prepared rapid acquisition gradient echo (MPRAGE) sequence with the following settings: repetition time (TR)/echo time (TE) = 2300/2.9 ms, slice number = 176, slice thickness = 1.2 mm, slice gap = 0 mm, flip angle = 9º, field of view (FOV) = 24.0 × 25.6 cm^2, number of averages = 1, matrix size = 240 × 256, voxel size = 1 × 1 × 1.2 mm^3, scan time = 5.2 mins. Functional images were acquired transversely with echo planar imaging (EPI) sequence with the following settings: TR/TE = 1000/30 ms, slice number = 52, slice thickness = 2.2 mm, slice gap = 2.64 mm, flip angle = 70º, FOV = 22 × 22 cm^2, number of averages = 1, matrix size = 64 × 64, voxel size = 3.4375 × 3.4375 × 2.2 mm^3. A total of 400 volumes were collected, resulting in a total scan time of 6.67 mins.

Data preprocessing

The functional images were preprocessed using Data Processing Assistant for

Resting-State fMRI (DPARSF) programs, which are based on Statistical Parametric Mapping (SPM12, http://www.fil.ion.ucl.ac.uk/spm) and MATLAB (MathWorks). The first 5 volumes were discarded due to unsteady magnetization. Slice-timing and realignment were performed. Then, data were spatially normalized to the Montreal Neurological Institute (MNI) template (resampling voxel size = $3 \times 3 \times 3$ mm^3), detrended, and filtered (0.01-0.08 Hz). Image volumes with framewise displacement (FD) >0.2 mm were scrubbed to reduce the effect of head motion using spline interpolation. Because we performed the Shapiro-Wilk test to check the normality of FD for each group and found that FD was not normally distributed for each group. No significant between-group difference in FD was found for the PD and HC groups (Z = -1.604, P = 0.109) and for the PD-ES and PD-LS groups (Z = -0.011, P = 0.991).

Grey matter volume (GMV) was calculated from structural MRI data using SPM12 and DPABI as follows: firstly, we segmented the structural images into grey matter, white matter, and cerebrospinal fluid; then, segmented images were also spatially normalized to the MNI template (resampling voxel size = $2 \times 2 \times 2$ mm^3) and modulated; finally, the modulated gray matter maps were smoothed using an isotropic Gaussian kernel [full width at half maximum (FWHM) = 6 mm].

Intrinsic timescale map

Based on the previous study[10], we used the preprocessed fMRI data to evaluate the intrinsic neural timescale for each voxel in the whole brain of each participant as follows. First, we estimated an autocorrelation function (ACF) of the fMRI signal of each voxel (time bin = TR), and then calculated the sum of ACF values in the initial period where the ACF showed positive values. The upper limit of this period was set at the point where the ACF hits zero for the first time. After repeating this procedure for every voxel, we also spatially smoothed the brain map (FWHM = 6 mm) to improve the signal-to-noise ratio and then obtained Z-transformed brain map to eliminate individual differences within group. We used this whole-brain map as an intrinsic timescale map in which the value at each voxel is equal to the intrinsic neural

timescale of the brain region. Two-sample t test was performed to compare the group differences in intrinsic timescales between the PD and HC groups, with age, sex, and mean FD as covariates. One-way ANOVA was also performed to compare the group differences in intrinsic timescales between the PD-ES, PD-LS, and HC groups, with age, sex, and mean FD as covariates. Then, post hoc comparisons using Bonferroni's test were performed, and pair-wise different Z-value maps corresponding to between-group intrinsic timescale differences were computed. The statistically significant threshold was set at voxel-wise $P < 0.001$, cluster-wise $P < 0.05$, two-tailed, and the minimum cluster size of 22 voxels after Gaussian random field (GRF) correction.

Correlations among intrinsic timescale, symptom severity, and age

To investigate the correlations among abnormal intrinsic timescale, symptom severity and age, we extracted the mean intrinsic timescale values of all voxels within each cluster from corrected statistical map and calculated the Spearman rank correlation coefficients for clinical measures and age with significant results between groups. Multiple comparisons were corrected using the Bonferroni method ($P < 0.05/106 = 0.0005$).

Grey matter volume and intrinsic timescale

We designed to investigate the neuroanatomical bases for intrinsic timescale by comparing GMV to the temporal property of neural signals. Two-sample t test was performed to compare the group differences in GMV between the PD and HC groups, and one-way ANOVA was also performed to compare the group differences in intrinsic timescales between the PD-ES, PD-LS, and HC groups, with age and sex as covariates. The statistically significant threshold was set at voxel-wise $P < 0.001$, cluster-wise $P < 0.05$, two-tailed, and the minimum cluster size of 58 voxels after GRF correction. We also extracted the mean GMV value of all voxels within each

significant cluster from corrected statistical intrinsic timescale maps, performed Mann-Whitney U test to compare the group differences of GMV between the PD and HC groups, and performed Kruskal-Wallis' H test with Dunn's test as the post hoc test compare the group differences of GMV among the PD-ES, PD-LS, and HC groups. To investigate the correlations between the intrinsic timescale and the GMV within each cluster from corrected statistical intrinsic timescale maps and calculated the Spearman rank correlation coefficients. Multiple comparisons were corrected using the Bonferroni method ($P < 0.05/106 = 0.0005$).

Acknowledgements

This research study was supported by the National Natural Science Foundation of China (81601467, 81871327), Medical Science and Technology Co-construction Project of Henan Province (LHGJ20220404), and Industrial Technology Foundation Public Service Platform Project of China (CEIEC-2020-ZM02-0103/03).

Competing interests

The authors report no competing interests.

References

1.   Hasson U, Yang E, Vallines I, Heeger DJ, Rubin N. A hierarchy of temporal receptive windows in human cortex. *The Journal of neuroscience : the official journal of the Society for Neuroscience*. Mar 5 2008;28(10):2539-50.

2.   Honey CJ, Thesen T, Donner TH, *et al*. Slow cortical dynamics and the accumulation of information over long timescales. *Neuron*. 2012;76(2):423-434.

3.   Murray JD, Bernacchia A, Freedman DJ, *et al*. A hierarchy of intrinsic timescales across primate cortex. *Nature neuroscience*. Dec 2014;17(12):1661-3.

4.   Wengler K, Goldberg AT, Chahine G, Horga G. Distinct hierarchical alterations of intrinsic neural timescales account for different manifestations of psychosis. *eLife*. Oct 27 2020;9:56151.

5.   Cirillo R, Fascianelli V, Ferrucci L, Genovesio A. Neural Intrinsic Timescales in the Macaque Dorsal Premotor Cortex Predict the Strength of Spatial Response Coding. *iScience*. Dec 21 2018;10:203-210.

6.   Golesorkhi M, Gomez-Pilar J, Tumati S, Fraser M, Northoff G. Temporal hierarchy of intrinsic neural timescales converges with spatial core-periphery organization. *Communications biology*. Mar 4 2021;4(1):277.

7.   Wolff A, Berberian N, Golesorkhi M, Gomez-Pilar J, Zilio F, Northoff G. Intrinsic neural timescales: temporal integration and segregation. *Trends in cognitive sciences*. Feb 2022;26(2):159-173.

8.   Zilio F, Gomez-Pilar J, Cao S, *et al*. Are intrinsic neural timescales related to sensory processing? Evidence from abnormal behavioral states. *NeuroImage*. Feb 1 2021;226:117579.

9.   Manea AMG, Zilverstand A, Ugurbil K, Heilbronner SR, Zimmermann J. Intrinsic timescales as an organizational principle of neural processing across the whole rhesus macaque brain. *eLife*. Mar 2 2022;11:75540.

10.   Watanabe T, Rees G, Masuda N. Atypical intrinsic neural timescale in autism. *eLife*. Feb 5 2019;8:42256.

11.   Raut RV, Snyder AZ, Raichle ME. Hierarchical dynamics as a macroscopic organizing principle of the human brain. *Proceedings of the National Academy of Sciences of the United States of America*. Aug 25 2020;117(34):20890-20897.

12.   Uscatescu LC, Said-Yurekli S, Kronbichler L, *et al*. Reduced intrinsic neural timescales in schizophrenia along posterior parietal and occipital areas. *NPJ schizophrenia*. Nov 22 2021;7(1):55.

13.   Northoff G, Sandsten KE, Nordgaard J, Kjaer TW, Parnas J. The Self and Its Prolonged Intrinsic Neural Timescale in Schizophrenia. *Schizophrenia bulletin*. Jan 23 2021;47(1):170-179.

14.   Wang K, Zhang X, Song C, *et al*. Decreased Intrinsic Neural Timescales in Mesial Temporal Lobe Epilepsy. *Frontiers in human neuroscience*. 2021;15:772365.

15.   Han S, Zheng R, Li S, *et al*. Integrative Functional, Molecular, and Transcriptomic Analyses of Altered Intrinsic Timescale Gradient in Depression. *Frontiers in neuroscience*. 2022;16:826609.

16.   Abbruzzese G, Berardelli A. Sensorimotor integration in movement disorders. *Movement disorders : official journal of the Movement Disorder Society*. Mar 2003;18(3):231-240.

17.   Lewis GN, Byblow WD. Altered sensorimotor integration in Parkinson's disease. *Brain : a journal of neurology*. Sep 2002;125(Pt 9):2089-99.

18. Weil RS, Schrag AE, Warren JD, Crutch SJ, Lees AJ, Morris HR. Visual dysfunction in Parkinson's disease. *Brain : a journal of neurology*. Nov 1 2016;139(11):2827-2843.

19. Ciccarelli N, Anzuino I, Pepe F, Magni E, Traficante D, Silveri MC. The facial emotion recognition deficit in Parkinson's disease: Implications of a visual scanning strategy. *Neuropsychology*. May 2022;36(4):279-287.

20. Litvan I, Aarsland D, Adler CH*, et al*. MDS Task Force on mild cognitive impairment in Parkinson's disease: critical review of PD-MCI. *Movement disorders : official journal of the Movement Disorder Society*. Aug 15 2011;26(10):1814-24.

21. Petrou M, Dwamena BA, Foerster BR*, et al*. Amyloid deposition in Parkinson's disease and cognitive impairment: a systematic review. *Movement disorders : official journal of the Movement Disorder Society*. Jun 2015;30(7):928-35.

22. Bell PT, Gilat M, O'Callaghan C*, et al*. Dopaminergic basis for impairments in functional connectivity across subdivisions of the striatum in Parkinson's disease. *Human brain mapping*. Apr 2015;36(4):1278-91.

23. Manza P, Zhang S, Li CS, Leung HC. Resting-state functional connectivity of the striatum in early-stage Parkinson's disease: Cognitive decline and motor symptomatology. *Human brain mapping*. Feb 2016;37(2):648-62.

24. Zang Z, Song T, Li J*, et al*. Simultaneous PET/fMRI revealed increased motor area input to subthalamic nucleus in Parkinson's disease. *Cerebral cortex*. Feb 23 2022;Advance online publication

25. Li J, Liao H, Wang T*, et al*. Alterations of Regional Homogeneity in the Mild and Moderate Stages of Parkinson's Disease. *Frontiers in aging neuroscience*. 2021;13:676899.

26. Ruppert MC, Greuel A, Freigang J*, et al*. The default mode network and cognition in Parkinson's disease: A multimodal resting-state network approach. *Human brain mapping*. Jun 1 2021;42(8):2623-2641.

27. Baggio HC, Sala-Llonch R, Segura B*, et al*. Functional brain networks and cognitive deficits in Parkinson's disease. *Human brain mapping*. Sep 2014;35(9):4620-34.

28. Sun HH, Pan PL, Hu JB, Chen J, Wang XY, Liu CF. Alterations of regional homogeneity in Parkinson's disease with "pure" apathy: A resting-state fMRI study. *Journal of affective disorders*. Sep 1 2020;274:792-798.

29. Lopes R, Delmaire C, Defebvre L*, et al*. Cognitive phenotypes in parkinson's disease differ in

terms of brain-network organization and connectivity. *Human brain mapping*. Mar 2017;38(3):1604-1621.

30. Rossini PM, Babiloni F, Bernardi G, *et al*. Abnormalities of short-latency somatosensory evoked potentials in parkinsonian patients. *Electroencephalography and clinical neurophysiology*. Jul-Aug 1989;74(4):277-89.

31. Nakashima K, Shimoyama R, Yokoyama Y, Takahashi K. Auditory effects on the electrically elicited blink reflex in patients with Parkinson's disease. *Electroencephalography and clinical neurophysiology*. Apr 1993;89(2):108-12.

32. Gerschlager W, Alesch F, Cunnington R, *et al*. Bilateral subthalamic nucleus stimulation improves frontal cortex function in Parkinson's disease. An electrophysiological study of the contingent negative variation. *Brain : a journal of neurology*. Dec 1999;122 (Pt 12):2365-73.

33. Ibarretxe-Bilbao N, Tolosa E, Junque C, Marti MJ. MRI and cognitive impairment in Parkinson's disease. *Movement disorders : official journal of the Movement Disorder Society*. 2009;24 Suppl 2:S748-53.

34. Cardoso EF, Fregni F, Maia FM, *et al*. Abnormal visual activation in Parkinson's disease patients. *Movement disorders : official journal of the Movement Disorder Society*. Aug 15 2010;25(11):1590-6.

35. Vignando M, Ffytche D, Lewis SJG, *et al*. Mapping brain structural differences and neuroreceptor correlates in Parkinson's disease visual hallucinations. *Nature communications*. Jan 26 2022;13(1):519.

36. Li N, Lei D, Peng J, *et al*. Brain network topology and future development of freezing of gait in Parkinson's disease: a longitudinal study. *Journal of neurology*. May 2022;269(5):2503-2512.

37. Pederzoli M, Girotti F, Scigliano G, Aiello G, Carella F, Caraceni T. L-dopa long-term treatment in Parkinson's disease: age-related side effects. *Neurology*. Nov 1983;33(11):1518-22.

38. Gai WP, Blumbergs PC, Geffen LB, Blessing WW. Age-related loss of dorsal vagal neurons in Parkinson's disease. *Neurology*. Nov 1992;42(11):2106-11.

39. Pihlstrom L, Fan CC, Frei O, *et al*. Genetic Stratification of Age-Dependent Parkinson's Disease Risk by Polygenic Hazard Score. *Movement disorders : official journal of the Movement Disorder Society*. Jan 2022;37(1):62-69.

40. Yang Y, Ye C, Sun J, *et al*. Alteration of brain structural connectivity in progression of Parkinson's disease: A connectome-wide network analysis. *NeuroImage Clinical*. 2021;31:102715.

41. Starr P, Little S. A space-time continuum in DBS: structural and functional advances in

Parkinson's disease. *Brain : a journal of neurology*. Mar 3 2021;144(2):357-359.

42.   Sarasso E, Agosta F, Piramide N, Filippi M. Progression of grey and white matter brain damage in Parkinson's disease: a critical review of structural MRI literature. *Journal of neurology*. Sep 2021;268(9):3144-3179.

43.   Mitchell T, Lehericy S, Chiu SY, Strafella AP, Stoessl AJ, Vaillancourt DE. Emerging Neuroimaging Biomarkers Across Disease Stage in Parkinson Disease: A Review. *JAMA neurology*. Oct 1 2021;78(10):1262-1272.

44.   Filippi M, Basaia S, Sarasso E*, et al*. Longitudinal brain connectivity changes and clinical evolution in Parkinson's disease. *Molecular psychiatry*. Sep 2021;26(9):5429-5440.

45.   Postuma RB, Berg D, Stern M*, et al*. MDS clinical diagnostic criteria for Parkinson's disease. *Movement disorders : official journal of the Movement Disorder Society*. Oct 2015;30(12):1591-601.

46.   Nasreddine ZS, Phillips NA, Bedirian V*, et al*. The Montreal Cognitive Assessment, MoCA: a brief screening tool for mild cognitive impairment. *Journal of the American Geriatrics Society*. Apr 2005;53(4):695-9.

47.   Shang S, Ye J, Wu J*, et al*. Early disturbance of dynamic synchronization and neurovascular coupling in cognitively normal Parkinson's disease. *Journal of cerebral blood flow and metabolism : official journal of the International Society of Cerebral Blood Flow and Metabolism*. Apr 26 2022;42(9):1719-1731.

48.   Bellot E, Kauffmann L, Coizet V, Meoni S, Moro E, Dojat M. Effective connectivity in subcortical visual structures in de novo Patients with Parkinson's Disease. *NeuroImage Clinical*. 2022;33:102906.

49.   Nackaerts E, Nieuwboer A, Broeder S, Swinnen S, Vandenberghe W, Heremans E. Altered effective connectivity contributes to micrographia in patients with Parkinson's disease and freezing of gait. *Journal of neurology*. Feb 2018;265(2):336-347.

50.   Tian Y, Chen HB, Ma XX*, et al*. Aberrant Volume-Wise and Voxel-Wise Concordance Among Dynamic Intrinsic Brain Activity Indices in Parkinson's Disease: A Resting-State fMRI Study. *Frontiers in aging neuroscience*. 2022;14:814893.

51.   Cai W, Young CB, Yuan R*, et al*. Dopaminergic medication normalizes aberrant cognitive control circuit signalling in Parkinson's disease. *Brain : a journal of neurology*. Mar 31 2022;Advance online publication

52. Xu J, Yu M, Wang H, *et al*. Altered Dynamic Functional Connectivity in de novo Parkinson's Disease Patients With Depression. *Frontiers in aging neuroscience*. 2021;13:789785.

53. Hosokai Y, Nishio Y, Hirayama K, *et al*. Distinct patterns of regional cerebral glucose metabolism in Parkinson's disease with and without mild cognitive impairment. *Movement disorders : official journal of the Movement Disorder Society*. Apr 30 2009;24(6):854-62.

54. Hirano S, Shinotoh H, Eidelberg D. Functional brain imaging of cognitive dysfunction in Parkinson's disease. *Journal of neurology, neurosurgery, and psychiatry*. Oct 2012;83(10):963-9.

55. Reeve A, Simcox E, Turnbull D. Ageing and Parkinson's disease: why is advancing age the biggest risk factor? *Ageing research reviews*. Mar 2014;14:19-30.

56. Kang SS, Zhang Z, Liu X, *et al*. TrkB neurotrophic activities are blocked by alpha-synuclein, triggering dopaminergic cell death in Parkinson's disease. *Proceedings of the National Academy of Sciences of the United States of America*. Oct 3 2017;114(40):10773-10778.

57. Martin LJ, Pan Y, Price AC, *et al*. Parkinson's disease alpha-synuclein transgenic mice develop neuronal mitochondrial degeneration and cell death. *The Journal of neuroscience : the official journal of the Society for Neuroscience*. Jan 4 2006;26(1):41-50.

58. Harrington DL, Shen Q, Vincent Filoteo J, *et al*. Abnormal distraction and load-specific connectivity during working memory in cognitively normal Parkinson's disease. *Human brain mapping*. Apr 1 2020;41(5):1195-1211.

Figure legends

Figure 1. The intrinsic timescales of the whole brain in all PD patients and the HC group, and in the PD patients at different stages. In each group, we found longer intrinsic timescales in frontopartial regions and shorter intrinsic timescales in sensorimotor, visual, and auditory regions. ES, early stage; HC, healthy controls; L, left; LS, late stage; PD, Parkinson's disease; R, right.

Figure 2. Voxel-wise intrinsic timescales between the PD and HC groups. The PD group had shorter timescales in bilateral postcentral and precentral gyri, bilateral lingual and calcarine gyri, and the right middle cingulum gyrus, and longer timescales in the right middle frontal gyrus than the HC group. HC, healthy controls; L, left; PD, Parkinson's disease; R, right.

Figure 3. Correlation between the intrinsic timescale and the symptom severity of PD. We found negative correlation between the intrinsic timescale of the lingual gyrus and the scores of the Hoehn and Yahr (HY) scale in patients with Parkinson's disease (PD).

Figure 4. Correlations between the intrinsic timescales and the age and between the cognitive-related scales and the age. We found negative correlation between the intrinsic timescales and the age in the lingual gyrus, postcentral gyrus, and the middle cingulum gyrus for the PD group (*Upper*) and negative correlation between the age and the scores of Montreal cognitive assessment (MoCA) and Mini-mental state examination (MMSE) (*Lower*).

Figure 5. Correlations between the intrinsic timescale and the grey matter volume. Shorter intrinsic timescale in the right postcentral gyrus correlated with its reduced grey matter volume.

Figure 6. Voxel-wise intrinsic timescales in PD at different stages. One-sample analysis of variance (ANOVA) revealed significant between-group differences in intrinsic timescales of bilateral lingual and calcarine gyri, the left precuneus, and the right middle cingulum gyrus (*Upper*). Post hoc comparisons using Bonferroni's test showed that the PD-LS group had shorter timescale in the left precuneus than the PD-ES group and shorter timescales in bilateral lingual and calcarine gyri than that in the HC group (*Lower*), and that the PD-ES group had longer timescales in the left superior frontal gyrus, the left inferior frontal gyrus, and the right middle frontal gyrus than the HC group (*Lower*). ES, early stage; HC, healthy controls; L, left; LS, late stage; PD, Parkinson's disease; R, right.

Table 1. The demographic and clinical data of PD and HC

	PD (n = 74)	HC (n = 73)	Z/X^2 values	P values
Age (SD)	59.03 (9.52)	59.55 (9.36)	-0.240	0.810
Sex (M/F)	33/41	38/35	0.819	0.365
UPDRS (SD, n = 72)	42.86 (19.33)	-	-	-
HY scale (SD, n = 74)	2.09 (0.88)	-	-	-
MoCA (SD, n = 57)	19.81 (5.38)	-	-	-
MMSE (SD, n = 71)	22.99 (3.85)	-	-	-

F, female; HC, healthy control; HY, Hoehn and Yahr; M, male; MMSE, mini-mental state examination; MoCA, Montreal cognitive assessment; PD, Parkinson's disease; UPDRS, unified Parkinson's disease rating scale

Table 2. The differences of average grey matter volumes between patients with PD at different stages and healthy controls in regions from corrected statistical intrinsic timescale maps

Between-group differences	Cluster	Regions	Hemisphere	Peak MNI Coordinate			PD-LS ($n = 44$)	PD-ES ($n = 30$)	HC ($n = 73$)	*P* values
				X	Y	Z				
ANOVA analysis	1	Lingual gyrus	L	-15	-54	0	0.389[ab]	0.429	0.431	0.003
		Calcarine gyrus								
	2	Lingual gyrus	R	9	-48	3	0.360[ab]	0.392	0.395	0.005
		Calcarine gyrus								
	3	Precuneus	L	41	-3	-63	0.481[a]	0.509	0.499	0.011
	4	Middle cingulum gyrus	R	26	3	-30	0.438[a]	0.399	0.426	0.006
PD-ES vs PD-LS	1	Precuneus	L	52	-3	-63	0.474[ab]	0.497	0.490	0.014
PD-LS vs HC	1	Calcarine gyrus	R	150	24	-57	0.355[ab]	0.389	0.390	0.005
		Lingual gyrus								
	2	Lingual gyrus	L	-15	-54	0	0.384[a]	0.423	0.424	0.005
		Calcarine gyrus								
PD-ES vs HC	1	Superior frontal gyrus	L	-6	54	39	0.359[a]	0.380	0.381	0.017
	2	Inferior frontal gyrus	L	-42	42	-6	0.329[a]	0.351	0.330	0.021
	3	Middle frontal gyrus	R	24	54	-12	0.306[a]	0.343	0.338	0.032

| 4 | Middle frontal gyrus | R | 36 | 51 | 24 | 0.220 | 0.248 | 0.244 | 0.046 |

ES, early stage; HC, healthy control; L, left; LS, late stage; MNI, Montreal Neurological Institute; PD, Parkinson's disease; R, right

[a] represents that the local grey matter volume of the PD-LS group is significantly lower than that of the PD-ES group, and [b] represents that the local grey matter volume of the PD-LS group is significantly lower than that of the HC group

PD

L   R

HC

L   R

Intrinsic timescales (s)

1   3.5

PD-ES

L   R

PD-LS

L   R

Postcentral gyrus (right)

rho = 0.254
P = 0.029