PROTEIN KINASE A ACTIVITY IN THE LEADING EDGE OF MIGRATING CELLS IS DEPENDENT ON THE ACTIVITY OF FOCAL ADHESION KINASE

Kathryn V. Svec[1], Mingu Kang[1], and Alan K. Howe[1,2,3*]

[1]Department of Pharmacology, University of Vermont, Burlington, VT, United States,
[2]Department of Molecular Physiology and Biophysics, University of Vermont, Burlington, VT, United States
[3]University of Vermont Cancer Center, University of Vermont, Burlington, VT, United States
*CORRESPONDENCE: Alan K. Howe, alan.howe@med.uvm.edu

ABSTRACT

Protein Kinase A (PKA) is a pleiotropic serine/threonine kinase whose localized and dynamic activity is required for cellular migration. Several cell types have shown robust PKA activity in the leading edge during migration. This activity is regulated by changes in actomyosin contractility, but the mechanism of the mechanochemical regulation of PKA remains unclear. Our ongoing investigation into this mechanism led us to discover a novel relationship between PKA and the non-receptor tyrosine kinase Focal Adhesion Kinase (FAK). Here, we find that while inhibition of actomyosin contractility leads to a relatively slow decrease in total cellular levels of phosphorylated FAK over tens of minutes, it decreases the focal adhesion-associated pool of both phospho-FAK and phospho-paxillin within two minutes, similar to the timing of the effect of inhibition of contractility on leading edge PKA activity. We then show that pharmacologic inhibition of FAK rapidly decreases leading edge PKA activity. Additionally, leading edge PKA activity is significantly reduced in FAK-null mouse embryonic fibroblasts and essentially eliminated by also silencing expression of the closely-related kinase Pyk2 in these cells. Importantly, leading edge PKA activity in FAK-null cells is rescued by re-expression of wild-type but not kinase-dead FAK. Taken together, these observations show that FAK activity is required for localized regulation of PKA during cell migration and, in so doing, establish a novel and unexpected connection between an adhesion-regulated tyrosine kinase and a serine/threonine kinase canonically downstream of G-protein coupled receptor and second-messenger signaling.

INTRODUCTION

Cell migration is essential to many diverse normal and pathological processes alike, including wound healing, development, and cancer cell metastasis. Many decades of research have elucidated myriad signaling pathways involved in cell migration, but in such an intricate process, seemingly discrete signaling nodes must work in concert to coordinate directed motility.

Substantial evidence points to Protein Kinase A (PKA) as an important mediator of cell migration upstream of membrane dynamics, actin organization, and many other processes critical to cell motility [1-3]. PKA is a principle serine/threonine kinase canonically activated downstream of G-protein coupled receptor (GPCR) activation and the activation of adenylyl cyclases [4]. Further, we and others have shown that PKA is active in the leading edge of migrating cells [5-11], alongside innumerable other signaling molecules that guide and coordinate cell migration, such as Cdc42 and RhoA [12, 13]. Our previous work identified a dependence of leading edge PKA activity on cellular contractility as it is rapidly lost upon treatment with myosin II inhibitor, blebbistatin [11]. Further, durotactic stretch can potentiate rapid, local bursts of PKA activity and PKA is required for mechanically gated cell migration, or durotaxis [11]. Thus, PKA activity in the leading edge of migrating cells is coupled to cellular contractility.

However, it remains unclear how PKA activity is regulated by mechanical inputs. Generally, cells sense changes in extracellular forces via focal adhesion complexes and associated dynamic actin structures. Mechanical force is translated into chemical signals via physical deformation of focal adhesion components, allowing for the confluence of different combinations of proteins and modified signaling [14-17]. Prominent among mechanically sensitive signaling proteins is Focal Adhesion Kinase (FAK), a nonreceptor tyrosine kinase well-characterized for its role in adhesive signaling, principally through the phosphorylation of paxillin and several other focal adhesion proteins [18, 19]. Further, FAK is involved in the iterative probing of the extracellular environment performed by focal adhesions through fluctuations in force during durotaxis [20] and is a key mediator of signaling to and from contractile forces [21-24]. Yet while FAK mediates mechanochemical signaling and is required for durotaxis[20, 25], it has never been implicated in the regulation of PKA activity, in any context.

Several studies provide evidence of crosstalk between tyrosine kinases and PKA signaling, but almost exclusively involving the regulation of tyrosine phosphorylation by PKA [26-34]. Though it is uncommon, it is not unheard of for a tyrosine kinase to regulate PKA activity. Both receptor tyrosine kinases and Src family kinases have been shown to regulate PKA activity by phosphorylating the PKA catalytic subunit directly [35, 36]. However, this work has been largely conducted *in vitro* and has not yet extended to FAK. Therefore, given the regulation of leading edge PKA activity by mechanical inputs and the central role of FAK as a mechanotransducer, we tested whether FAK plays a role in regulating localized PKA activity in migrating cells.

RESULTS

Disrupting actin polymerization does not lead to an immediate loss of leading edge PKA activity

Our prior research identified leading edge PKA activity as being rapidly modulated by mechanical inputs, both applied and intrinsic. Transduction of mechanically-gated signals during migration depends on actin polymerization [37, 38]. Therefore, we examined PKA activity in migrating cells before and after treatment with inhibitors of actin polymerization. Specifically, PKA activity was monitored in migrating SKOV-3 ovarian cancer cells expressing a membrane targeted PKA biosensor (pmAKAR3) before and after treatment with either latrunculin A (100nM) or cytochalasin D (50nM). As expected, leading edge dynamics were rapidly inhibited (within 1 min) after addition of either drug (Supplemental Videos 1 and 2). However, neither actin polymerization inhibitor led to a significant decrease in the mass of relative leading edge PKA activity within 10 minutes of treatment (Supplemental Figure 1; Supplemental Videos 1 and 2). These observations, along with a previous study which showed that leading edge PKA activity in α4 integrin-expressing CHO cells gradually decreased 9-20 minutes after latrunculin A treatment [7], are in contrast to the rapid (within 2 min) and complete inhibition of PKA upon loss of contractility after treatment with blebbistatin and the even more rapid (within 30 sec) activation of PKA upon cell stretch [11]. Taken together, these data suggest that leading-edge PKA activity is not immediately or directly regulated by actin dynamics *per se* and suggest that PKA is coupled to changes in actomyosin contractility by means other than through actin polymerization.

Disruption of cellular traction forces rapidly and transiently inhibits FAK activity within focal adhesions

The rapid effect of cell-ECM traction on PKA activity led us to consider other adhesion- and/or migration-related signaling pathways that are known to be mechanosensitive. Focal adhesion kinase (FAK) is a centrally important mediator of integrin-mediated adhesive signaling, regulator of cell migration, and component of mechanotransduction pathways [18, 19, 22, 23]. Thus, given that PKA activity is stimulated by integrin-mediated adhesion [29, 39], important for efficient cell migration [1, 3, 5, 9], and regulated by mechanical signaling [3, 40], FAK is well-positioned to be a regulator of leading edge PKA activity. Moreover, like PKA activity, FAK activity (as measured by autophosphorylation on Y397) also decreases upon inhibition of actomyosin contractility [41, 42]. Importantly, however, the effect of loss of contractility on FAK activity and/or autophosphorylation has, to date, only been observed after a period of 30-60 minutes, at which point focal adhesion morphology is dramatically altered [40-42]. In contrast, PKA activity decreases much faster after loss of tension - within 2 minutes - rapidly following the loss of cellular traction force and significantly preceding the structural dissolution of focal adhesions [11]. We therefore investigated whether there was an effect of blebbistatin on FAK activity at early timepoints that might correspond to the effect on PKA activity. Specifically, migrating SKOV-3 cells were left untreated or were treated with blebbistatin for varying amounts of time (from 2 to 60 min) then harvested for immunoblotting or processed for immunofluorescence with antibodies against pY397 FAK (the autophosphorylation site). In agreement with published observations [41, 42], inhbition of contractility led to a decrease in total cellular FAK autophosphorylation by 30 min but had no apparent effect on autophosphorylation before 5 min (Figure 1A). However, when the phospho-FAK content within focal adhesions was assessed by immunofluorescence, there was a significant decrease in pY397-FAK (normalized to total focal adhesion area demarcated by vinculin staining as described in Figure 1B) within 2 minutes of blebbistatin treatment, (Figure 1C

and D). This rapid decrease in FAK autophosphorylation was paralleled by a similar decrease in phosphorylation of paxillin at Y118 paxillin, a major substrate of FAK (Figure 1C and E). These data suggest that a pool of active FAK is rapidly dephosphorylated within focal adhesions before the observable dissolution of the focal adhesion structures themselves. To our knowledge, this is the most immediate effect of myosin II inhibition on FAK phosphorylation described to date. Interestingly, the initial blebbistatin-mediated decrease in both pY397 FAK and pY118 paxillin levels was followed by a recovery by 5 min and a slight but steady and significant increase at later time points, becoming even higher than control levels as the focal adhesions dissolved and diminished in size (Figure 1C-E). This recovery in pY397-FAK levels is consistent with earlier observations that while focal adhesion size is greatly reduced 30 minutes after blebbistatin treatment, the level of phosphorylated FAK in remaining adhesive structures is comparable to the level found in mature focal adhesions in untreated cells [40]. This recovery notwithstanding, these observations confirm that FAK can be regulated by actomyosin contractility and establish that at least a portion of FAK activity responds with the same rapid kinetics as the loss of PKA activity to the loss of contractility.

Pharmacologic inhibition of FAK decreases leading edge PKA activity
To directly test whether FAK activity was required for leading edge PKA activity, pmAKAR3-expressing SKOV-3 cells were analyzed by FRET microscopy before and after treatment with PF-562271 (PF271), a potent, ATP-competitive FAK inhibitor. Treatment with 250nM PF271 led to a rapid decrease in membrane-proximal leading edge PKA activity within 10 minutes of treatment (Figure 2A-D; Supplemental Video 3). This effect did not change the ability of PKA to respond to increases in cAMP using a saturating dose of forskolin (to activate adenylyl cyclases) and IBMX (to inhibit phosphodiesterases; Supplemental Figure 2), though the measurement of this effect was limited by the dynamic range of the biosensor. Importantly, the concentration used here was 2.5-fold lower than the reported *in vitro* IC_{50} of PF271 against PKA directly [43]. Nonetheless, to ensure that the decrease in PKA activity was not an off-target effect of PF271, the assay was repeated with two additional FAK inhibitors - PND1186 and the lesser known, but structurally dissimilar, FAK inhibitor 14 (also known as Y15). Treatment with FAK inhibitor 14 caused a rapid and complete reduction of leading edge PKA activity, even more striking than that caused by PF271 (Figure 2D). While we found no data reporting any direct effect of FAK inhibitor 14 on PKA activity, this inhibitor is reported as highly selective for FAK activity, having no measurable effect on even highly similar tyrosine kinases such as Pyk2 or Src [44-46]. Treatment with a third FAK inhibitor, PND1186 (also called VS-4718), led to a distinct effect. Whereas peaks of PKA activity in the leading edge were diminished, as seen with the previous inhibitors, overall PKA activity in other parts of the cell rose (Supplemental Figure 3 A and B). While the reason for this increase is unknown, it might be explained by *in vitro* studies that showed PND1186 can cause a slight direct increase in PKA activity [47]. Taken together, the demonstration that three chemically distinct inhibitors of FAK rapidly decreased leading edge PKA activity solidly suggests a functional connection between these two pathways.

Genetic loss of FAK leads to decreased PKA activity in the leading edge
Despite both the current observations as well as the parallels between FAK and PKA in terms of their regulation by and role in adhesion, migration, and mechanotransduction, there are no reports of FAK acting upstream of PKA activity and no obvious mechanism or target through which FAK might exert its effect. Thus, to eliminate the possibility of off-target effects of pharmacologic

inhibition and to further confirm this novel regulatory connection, leading edge PKA activity was observed in wild-type mouse embryonic fibroblasts (MEFs) as well as MEFs isolated from FAK-null embryos [48]. Similar to SKOV-3 and other cell types previously assessed for PKA activity during migration [1, 3, 5, 7-11, 49], wild-type MEF cells exhibit leading edge PKA events (Figure 3A; Supplemental Video 4). Also as in SKOV-3 cells, this localized PKA activity is rapidly decreased upon pharmacologic inhibition of FAK by PF271 (Figure 3B; Supplemental Video 4). However, FAK-null MEFs showed significantly lower peak and average PKA activity in the leading edge compared to wild-type MEFs (Figure 3C, D, and G; Supplemental Video 5). Of note, the whole-cell mean FRET ratios of the FAK-null cells was not significantly different than that of wild-type MEFs and a minimal level of membrane-proximal PKA activity was retained in protrusive regions (Figure 3C, F, and G).

Interestingly, the low level of leading edge PKA activity observed in FAK-null cells was further diminished by treatment with PF271 (Figure 3D; Supplemental Video 5), raising the question of what might be the target for this inhibitor in the absence of FAK. Pyk2 is a related tyrosine kinase which shares 45% amino acid identity with FAK and is upregulated and highly active in FAK-null cells [50]. Prior studies have shown that this increase in Pyk2 abundance and activity can partially compensate for the loss of FAK in some contexts [50, 51]. Further, Pyk2 is also inhibited by PF271, albeit with a higher IC_{50} than FAK [43]. Thus, we assessed whether loss of Pyk2 would further diminish PKA activity in the FAK-null MEFs. To this end, we assessed PKA activity in FAK-null MEFs stably expressing a short hairpin RNA against Pyk2 to knock down Pyk2 activity (shPyk2). Total PKA activity in the FAK-null/shPyk2 cells was significantly diminished compared to both the wild-type and FAK-null cells, as measured by whole cell FRET ratio (Figure 3E and F; Supplemental Video 6). Also, though the relative mass of leading edge PKA activity in FAK-null/shPyk2 cells was lower than in FAK-null cells, this decrease was not statistically significant, likely due to the already low level of activity in the FAK-nulls and the limited number of FAK-null/shPyk2 cells per experiment that were amenable to analysis due to low transfection efficiency (Figure 3E and G). Taken together, these genetic data corroborate the observations from the pharmacological inhibitors and demonstrate that FAK - and to a lesser extent, Pyk2 - is required for leading edge PKA activity in migrating cells.

Leading edge PKA activity in FAK-null cells is rescued by wild-type but not kinase-dead FAK.
In addition to its tyrosine kinase activity, FAK has been shown to exhibit important cellular effects by acting as a scaffold that physically organizes various signaling intermediates [52]. To further corroborate the pharmacologic inhibitor data and confirm that FAK kinase activity rather its structural and scaffolding functions were required for regulation of leading edge PKA activity, FAK-null MEFs were co-transfected with constructs expressing pmAKAR3 and either mCherry-tagged wild-type or kinase-dead (R454) FAK, or mCherry-paxillin as a control. Re-expression of mCherry tagged wild-type FAK rescued leading edge PKA activity to the level of wildtype cells (Figure 4A and E; Supplemental Video 7). As expected, expression of a control mCh-paxillin construct was unable to rescue leading edge PKA activity above that of the parent FAK-null cell line (Figure 4 B and E; Supplemental Video 8). Similarly, cells expressing the kinase dead mCh-R454 FAK exhibited limited PKA activity in the leading edge, similar to FAK-null cells alone or those expressing mCh-paxillin (Figure 4 C and E; Supplemental Video 9). These data validate the causal connection between FAK function and leading edge PKA activity and demonstrate that

FAK kinase activity, and not its scaffolding or other functions, is required for its regulation of PKA activity in the leading edge of migrating cells.

DISCUSSION

We have identified a novel relationship whereby FAK promotes PKA activity in the leading edge of migrating SKOV-3 ovarian cancer cells and MEF cells. This relationship is sensitive to both pharmacologic and genetic inhibition of FAK and is dependent on FAK kinase activity. It is rather surprising that FAK inhibition has such a marked effect on PKA activity in the leading edge. The canonical pathway leading to activation of PKA, the generation of cAMP downstream of adenylyl cyclases (ACs), heterotrimeric G-proteins, and GPCRs, is well characterized and ostensibly wholly independent of tyrosine phosphorylation. There is no precedent suggesting the mechanism by which this regulation occurs. It is possible that FAK is acting on regulatory targets upstream of PKA, such as heterotrimeric G proteins, ACs, or phosphodiesterases, that control the local availability of cAMP. To our knowledge, however, there are no published reports of FAK activity contributing to the generation of cAMP, in any system. There is evidence to support the regulation of FAK downstream of ACs, both promoting FAK activity downstream of increased AC activity [31] or norepinephrine stimulation [53], and decreasing FAK activity after stimulation of ACs in microdomains [54]. Further, inhibition of PKA has been shown to delay dephosphorylation of FAK upon cellular detachment, putting PKA upstream of FAK in anchorage-dependent signaling [29]. Yet no mechanism for these relationships has been identified, nor has the reciprocal relationship, whereby FAK modulates the activity of adenylyl cyclases, been demonstrated.

While other tyrosine kinases such as EGFR and Fyn have been shown to directly phosphorylate the PKA catalytic domain [35, 36], these modifications lead to only a modest increase in the catalytic efficiency of PKA *in vitro*, and the direct PKA phosphorylation by FAK has not been investigated. Thus, while phosphorylation of PKA subunits by FAK has not been shown, it is possible that FAK, like EGFR and Fyn, may directly modify and thus regulate PKA catalytic subunits. However, if FAK were directly phosphorylating PKA, it seems unlikely that we would see a marked decrease in PKA activity within minutes of FAK inhibition as we see in Figure 2. If, similar to EGFR and Fyn, phosphorylation of PKA by FAK slightly increased PKA catalytic efficiency, then inhibiting FAK would cease ongoing phosphorylation of PKA. A decrease in PKA activity would then rely on swift, overwhelming action by tyrosine phosphatases, which would likely only modestly decrease the catalytic response of PKA to the unchanged availability of cAMP. Therefore, it is more likely that FAK is acting upstream, potentiating PKA through increased availability of cAMP, allowing for more immediate and marked change through signal amplification via second messenger production.

Moreover, using the R454 kinase-dead FAK construct in FAK-null cells, we have demonstrated that it is FAK kinase activity, and not one of the numerous scaffolding functions of FAK, that is responsible for regulation of PKA. FAK is a large protein known for its ability to scaffold higher order signaling complexes through its binding with numerous other proteins—other kinases such as Src, adapter molecules such as p130Cas, and focal adhesion components like talin and paxillin [52, 55-58]. Our results suggest that FAK is acting to influence PKA activity more directly than by scaffolding some components of the signaling pathway in a way that requires the input of its kinase activity.

With these results, it is unclear whether mechanical inputs, PKA, and FAK exist in a linear hierarchy where FAK acts as an intermediate signaling node. Prior experiments suggested that FAK inhibition does not rapidly change actomyosin contractility [40]. If FAK is acting to transmit mechanical signals to PKA, we would expect that disrupting contractility would lead to rapid decreases in FAK activity. Though we have shown that at least some pools of FAK are rapidly sensitive to contractile forces (within 2 minutes), the data presented do not categorically place FAK activity in the same pathway as mechanical regulation of PKA. While the loss of contractility rapidly inhibited FAK activity in focal adhesions, our prior work has shown that leading edge PKA events are not directly spatially related to focal adhesions [11]. Additionally, FAK functions in distinct pools within the cell, not always localized to focal adhesions [59]. Therefore, further work must be done at higher time resolution to identify whether a subcellular pool of FAK activity is contributing to PKA activity in migrating cells. This can be addressed using subcellular fractionation techniques to sensitize western blotting to small, subcellular changes in FAK activity downstream of myosin II inhibition.

The novel involvement of FAK in leading edge PKA activity is intriguing regardless of whether FAK is involved specifically in its mechanical regulation. PKA is active in the leading edge of numerous cell types, is required for the migration of many different types of cells, and has important downstream targets involved in many aspects of this complex process [3]. FAK is similarly widely required for cell migration, adhesion, and mechanotransduction, yet these two principal kinases are canonically regarded to function separately. The identification of crosstalk between FAK and PKA represents an important finding in the regulation of migration and implies that these ubiquitous but heretofore disparate pathways may crosstalk in other cellular processes. This represents a wholly new aspect of regulation of the PKA pathway and an itinerant new avenue of investigation to identify its extent and underlying mechanism.

MATERIALS AND METHODS
Cell Culture
SKOV-3 cells were obtained from ATCC and maintained in DMEM supplemented with 10% fetal bovine serum below 60% confluency. Cells plated for transfection were allowed to reach 70% confluency and transfected with FuGene 6. After 30 hours, transfected SKOV-3 cells were cryopreserved at -80°C in DMEM 10% fetal bovine serum, 10% DMSO.

MEF WT, FAK KO, and FAK KO shPyk2 cells were obtained from David Schlaepfer and maintained in DMEM supplemented with 10% fetal bovine serum on dishes coated with 0.1% gelatin [48]. Cells were transfected at 70% confluency, replated at 24 hours, and cryopreserved after 48 hours at -80°C in DMEM 10% fetal bovine serum, 10% DMSO.

FRET Microscopy
Imaging dishes (29mm; CellVis) were coated overnight at 4°C with 10µg/ml fibronectin. SKOV-3 cells were thawed and plated at 25,000 cells per dish in DMEM supplemented with 10% fetal bovine serum. Four hours after plating, cells were rinsed and media was exchanged to Ringer's Solution supplemented with 12.5ng/ml epidermal growth factor. Cells were allowed to equilibrate for at least 45 min before imaging. Imaging began 20 minutes before drug treatment and extended 20 minutes beyond treatment to assess the effect of each drug.

MEF cells were thawed and plated at 20,000 cells per dish in DMEM supplemented with 10% fetal bovine serum. Eighteen hours after plating, cells were rinsed and media was exchanged with Ringer's Solution supplemented with 10ng/ml plasma derived growth factor-BB.

CFP and FRET images were acquired simultaneously using a Nikon TiE inverted fluorescent microscope and the Cairn OptoSplitII optical splitter. Temperature was maintained at 37°C. FRET images were acquired using 2X2 binning in Nikon Elements imaging software.

FRET Image Analysis
FRET images were processed using a protocol based on Broussard *et al*. as described previously [11, 60] (Add to citation list). Briefly, a median filter of radius =1 was applied and background was subtracted from each channel using a mean background pixel value for each image. The FRET image was then corrected for CFP bleed-through and division of the corrected FRET image by the CFP image resulted in the FRET ratio image. This image was then masked using the sum of the corrected FRET and CFP images, sharpened with an unsharp mask, radius 10. FRET ratio images were visualized using a custom LUT.

Quantification of relative leading edge PKA activity was calculated by the number of pixels in the leading edge with a FRET ratio value measuring more than 1.5 standard deviations above the mean pixel value of the entire cell during the pre-drug portion of the experiment, divided by the total number of pixels in the leading edge. The remaining leading edge activity after a given treatment was calculated by dividing the relative LE PKA activity 10 minutes after by that 10 minutes before treatment. Whole cell mean FRET ratio calculations were averaged over 7 frames of the movie, absent of any treatment, while the cell migrated.

Immunofluorescence
SKOV-3 cells were plated in DMEM supplemented with 10% fetal bovine serum on coverslips coated with 10μg/mL fibronectin overnight then treated with 25 uM blebbistatin for the indicated periods. Cells were then fixed in 4% PFA in PBS for 15 min at RT, permeabilized in 0.25% triton X100 in PBS for 10 min at RT with gentle shaking, and blocked for at least 1 h in 4% low- IgG BSA + 0.1% tween-20 in PBS. Coverslips were stained for 1 h at RT with antibodies against vinculin (hVin1, Sigma; 1:400) and either (phosphoTyr397-FAK (Cat# 44-625G, ThermoFisher; 1:50) or phosphoTyr118-paxillin (Cat# 44-722G; ThermoFisher; 1:100) diluted in blocking solution. Coverslips were washed 4 times for 7 min in PBS, then incubated in secondary antibodies (Alexa594-ocnjugated donkey anti-mouse and Alexa488-conjugated anti-rabbit, both from Abcam and both diluted 1:250 in block) and washed again as above. Coverslips were mounted on slides using Prolong Glass Antifade (ThermoFisher) and imaged as described elsewhere [40].

Western Blot
SKOV-3 cells were plated in DMEM supplemented with 10% fetal bovine serum on 60 mm tissue culture dishes previously coated in 10μg/mL fibronectin. Following a similar protocol to imaging parameters, at ~70% confluency, cells were stimulated with 12.5ng/mL EGF for 30 minutes prior to drug treatments. Timed treatments with 25μM blebbistatin were ended by a double rinse in ice-cold complete PBS followed by harvest in lysis buffer containing 50mM Tris, pH 8.0, 150mM NaCl, 5mM EDTA, 5% glycerol, 1% Triton-X 100, 25mM NaF, 2mM Na_2VO_4 supplemented with 1x protease inhibitor cocktail (Thermo Scientific). Cells were freeze-thawed twice in liquid nitrogen and whole cell extracts were collected upon centrifugation at 16,000 x g. 10μg of protein from each treatment condition was separated by SDS-PAGE and transferred to PVDF membrane. Membranes were blocked with 5% nonfat dry milk and immunoblotted with antibodies against pY397-FAK (Invitrogen 44-624G) or total FAK (Millipore, 05-537).

Figure Legends

Figure 1: Blebbistatin rapidly but transiently inhibits the focal adhesion pool of FAK. (A) Western blot of whole cell lysates from cells treated with 25μM blebbistatin for 0, 2, 5, 15, 30, or 60 minutes probed for pY397 FAK or total FAK. (B) Schematic describing the method used to calculate and mask relative phospho-FAK and phospho-Pxn in focal adhesions. Briefly, SKOV-3 cells treated with 25 μM blebbistatin for 0, 2, 5, 15, 30, or 60 minutes were probed for vinculin, to demarcate total focal adhesions, and either pY397 FAK or pY118 Pxn. Vinculin signal was used to create a binary focal adhesion mask which was multiplied by the phosphoprotein signal. Phosphoprotein signal was then divided by vinculin signal to give relative phosphoprotein images as shown in B. (C) Pseudocolored relative phosphoprotein images for pY397 FAK (top) or pY118 Pxn (bottom) upon treatment with 25μM Blebbistatin for the indicated time period. (D) Relative phosphoprotein signal in each masked adhesion for each pY397 FAK (left) or pY118 Pxn (right). * p≤0.05, **** p≤0.0001.

Figure 2: Inhibiting FAK activity rapidly decreases leading edge PKA activity. (A) FRET ratio image montage of the leading edge of an SKOV-3 cell expressing pmAKAR3 treated with 250nM PF271 at time 0,scale=10μm. (B) Kymograph of leading edge PKA activity through line in panel A,.(C) Relative peak PKA activity in the leading edge of cells over time treated with vehicle control, 0.1% DMSO (N=3, n=36), and cells treated with 250nM PF271 (N=3, n=36). (D) Peak PKA activity remaining in the leading edge 10 minutes after treatment compared to that 10 minutes before treatment with 0.1% DMSO, 250nM PF271, and 10μM FAKi 14 (N=1, n=10). Kruskal-Wallis with multiple comparisons, * p≤0.05, ** p≤0.01, **** p<0.0001) Comparisons shown are a subset of all comparisons assessed by Kruskal-Wallis test. Comprehensive graph including all test results found in Supplemental Figure 4.

Figure 3: Fibroblasts lacking FAK exhibit decreased leading edge PKA activity.(A) FRET ratio image of a migrating wildtype MEF cell expressing pmAKAR3, PKA activity reporter, scale bar=10μm. (B) Montage of FRET ratio in the leading edge of a wildtype MEF cell upon treatment with 250nM PF271 showing decreased leading edge PKA activity six and twelve minutes after treatment, scale bar=10μm. (C) FRET ratio image of a migrating FAK-null MEF cell expressing pmAKAR3, scale bar=10μm. Note the uniform LUT scaling in panels A and C. (D) Montage of FRET ratio in the leading edge of a FAK-null MEF cell upon treatment with 250nM PF271, showing very slightly decreased leading edge PKA activity six and twelve minutes after treatment, scale bar=10μm. (E) FRET ratio image of a MEF FAK-null cell stably expressing shPyk2 and transiently expressing pmAKAR3, scale bar=10μm. LUT scaling enhanced due to decreased mean FRET ratios in shPyk2 cells. (F) Whole cell PKA activity as assessed by mean FRET ratio values for entire cell in wildtype MEF (N=5, n=30), FAK-null MEF (N=3, n=27), and FAK-null/shPyk2 MEF (N=5, n=24) cells. Comparisons shown are a subset of all comparisons assessed by Kruskal-Wallis test. Comprehensive graph including all test results found in Supplemental Figure 5A. (G) Relative leading edge PKA activity in wildtype MEF (N=5, n=19), FAK-null MEF (N=3, n=26), and FAK-null/shPyk2 MEF (N=4, n=14) cells. Comparisons shown are a subset of all comparisons assessed by Kruskal-Wallis test. Comprehensive graph including all test results found in Supplemental Figure 5B. * p≤0.05, *** p≤0.001, **** p≤0.0001.

Figure 4: Leading edge PKA activity is rescued by wild-type but not kinase-dead FAK. (A) FRET ratio image of a migrating FAK-null MEF cell expressing mCh-FAK and pmAKAR3, scale bar=10μm. (B) FRET ratio image of a migrating FAK-null MEF cell expressing mCh-paxillin and pmAKAR3, scale bar=10μm. (C) FRET ratio image of a MEF FAK-null cell expressing mCh-

R454 FAK and pmAKAR3, scale bar=10μm. Note the LUT scaling is the same for A-C. (D) Whole cell PKA activity as assessed by mean FRET ratio values for entire cell. mCh-FAK (N=3, n=42), mCh-Paxillin (N=3, n=22), or mCh-R454 FAK (N=3, n=30). Comparisons shown are a subset of all comparisons assessed by Kruskal-Wallis test. Comprehensive graph including all test results found in Supplemental Figure 5A. (E) Relative leading edge PKA activity in FAK-null MEF cells expressing mCh-FAK (N=3, n=42), mCh-Paxillin (N=3, n=22), or mCh-R454 FAK (N=3, n=30). Comparisons shown are a subset of all comparisons assessed by Kruskal-Wallis test. Comprehensive graph including all test results found in Supplemental Figure 5B. * p≤0.05, ** p≤0.01, *** p≤0.001, **** p≤0.0001.

Supplemental Figure Legends
Supplemental Figure 1: Inhibiting actin polymerization does not rapidly decrease leading edge PKA activity (A) FRET ratio image montage of the leading edge of an SKOV-3 cell expressing pmAKAR3 treated with 50nM Cytochalasin D at time 0, scale=10μm. (B) Kymograph of pmAKAR3 raw YFP emission (greyscale, intentionally over-brightened for ease of visualization; *top*) and leading edge PKA activity (pseudocolored; *bottom*) through the linear region of interest (ROI) in panel A. (C) FRET ratio image montage of the leading edge of an SKOV-3 cell expressing pmAKAR3 treated with 100nM Latrunculin A at time 0. Scale=10μm (D) Kymograph of pmAKAR3 raw YFP emission and leading edge PKA activity through the ROI in panel C. (E) Relative peak PKA activity in the leading edge of cells over time treated with vehicle control, 0.1% DMSO (N=3, n=36), and cells treated with 50nM Cytochalasin D (N=3, n=34) or 100nM Latrunculin A (N=3, n=28). (D) Peak PKA activity remaining in the leading edge 10 minutes after treatment compared to that 10 minutes before treatment with 0.1% DMSO, 50nM Cytochalasin D, or 100nM Latrunculin A. Kruskal-Wallis with multiple comparisons. Comparisons shown are a subset of all comparisons assessed by Kruskal-Wallis test. Comprehensive graph including all test results found in Supplemental Figure 4.

Supplemental Figure 2: Whole cell FRET ratio to Forskolin and IBMX in SKOV-3 cells after treatment with PF271. After treatment with PF271 or DMSO control, cells from Figure 2 were treated with Forskolin and IBMX to increase intracellular cAMP and assess maximum PKA signal. No significant difference was found between the cell means.

Supplemental Figure 3: Inhibiting FAK with PND1186 leads to decrease in leading edge PKA activity, but slight whole cell increase. (A) FRET ratio image montage of the leading edge of an SKOV-3 cell expressing pmAKAR3 treated with 10μM at time 0, scale=10μm (B) Peak PKA activity remaining in the leading edge 10 minutes after treatment compared to that 10 minutes before treatment with 0.1% DMSO (N=3, n=36) or 10μM PND1186 (N=2, n=18). Kruskal-Wallis with multiple comparisons. Comparisons shown are a subset of all comparisons assessed by Kruskal-Wallis test. Comprehensive graph including all test results found in Supplemental Figure 4.

Supplemental Figure 4: Comprehensive graph including all test results from Kruskal-Wallis assessment of the treatment of SKOV-3 cells with pharmacologic inhibitors. Comparisons shown in Figure 2 and Supplemental Figures 1-3 are a subset of all comparisons assessed by Kruskal-Wallis test, the complete results of which are shown here. * p≤0.05, ** p≤0.01, *** p≤0.001, **** p≤0.0001.

Supplemental Figure 5: Comprehensive graph including all test results from Kruskal-Wallis assessment of MEF cell lines and MEF FAK-null cells expressing mCherry tagged addback constructs. Comparisons shown in Figures 3 and 4 are a subset of all comparisons assessed by Kruskal-Wallis test, the complete results of which are shown here. (A) Mean PKA activity as assessed by mean FRET ratio. (B) Relative leading edge PKA activity * $p \leq 0.05$, ** $p \leq 0.01$, *** $p \leq 0.001$, **** $p \leq 0.0001$.

References

1. Howe AK. Regulation of actin-based cell migration by cAMP/PKA. Biochimica et biophysica acta. 2004;1692(2-3):159-74. Epub 2004/07/13. doi: 10.1016/j.bbamcr.2004.03.005. PubMed PMID: 15246685.
2. Diviani D, Scott JD. AKAP signaling complexes at the cytoskeleton. J Cell Sci. 2001;114(Pt 8):1431-7. Epub 2001/04/03. doi: 10.1242/jcs.114.8.1431. PubMed PMID: 11282019.
3. Svec K, Howe A. Protein Kinase A in Cellular Migration – Niche Signaling of a Ubiquitous Kinase. Frontiers in Molecular Biosciences. 2022;9. doi: 10.3389/fmolb.2022.953093.
4. Lodish HF. Molecular cell biology. Eighth ed. New York: W.H. Freeman-Macmillan Learning; 2016.
5. Howe AK, Baldor LC, Hogan BP. Spatial regulation of the cAMP-dependent protein kinase during chemotactic cell migration. Proceedings of the National Academy of Sciences of the United States of America. 2005;102(40):14320-5. Epub 2005/09/24. doi: 10.1073/pnas.0507072102. PubMed PMID: 16176981; PubMed Central PMCID: PMCPMC1242330.
6. Lim CJ, Han J, Yousefi N, Ma Y, Amieux PS, McKnight GS, et al. Alpha4 integrins are type I cAMP-dependent protein kinase-anchoring proteins. Nature cell biology. 2007;9(4):415-21. Epub 2007/03/21. doi: 10.1038/ncb1561. PubMed PMID: 17369818.
7. Lim CJ, Kain KH, Tkachenko E, Goldfinger LE, Gutierrez E, Allen MD, et al. Integrin-mediated protein kinase A activation at the leading edge of migrating cells. Molecular biology of the cell. 2008;19(11):4930-41. Epub 2008/09/12. doi: 10.1091/mbc.e08-06-0564. PubMed PMID: 18784251; PubMed Central PMCID: PMCPMC2575143.
8. Paulucci-Holthauzen AA, Vergara LA, Bellot LJ, Canton D, Scott JD, O'Connor KL. Spatial distribution of protein kinase A activity during cell migration is mediated by A-kinase anchoring protein AKAP Lbc. The Journal of biological chemistry. 2009;284(9):5956-67. Epub 2008/12/25. doi: 10.1074/jbc.M805606200. PubMed PMID: 19106088; PubMed Central PMCID: PMCPMC2645839.
9. McKenzie AJ, Campbell SL, Howe AK. Protein kinase A activity and anchoring are required for ovarian cancer cell migration and invasion. PLoS One. 2011;6(10):e26552. Epub 2011/10/27. doi: 10.1371/journal.pone.0026552. PubMed PMID: 22028904; PubMed Central PMCID: PMCPMC3197526.
10. Tkachenko E, Sabouri-Ghomi M, Pertz O, Kim C, Gutierrez E, Machacek M, et al. Protein kinase A governs a RhoA-RhoGDI protrusion-retraction pacemaker in migrating cells. Nature cell biology. 2011;13(6):660-7. Epub 2011/05/17. doi: 10.1038/ncb2231. PubMed PMID: 21572420; PubMed Central PMCID: PMCPMC3746034.
11. McKenzie AJ, Svec KV, Williams TF, Howe AK. Protein kinase A activity is regulated by actomyosin contractility during cell migration and is required for durotaxis. Molecular biology of the cell. 2020;31(1):45-58. Epub 2019/11/14. doi: 10.1091/mbc.E19-03-0131. PubMed PMID: 31721649; PubMed Central PMCID: PMCPMC6938270.
12. Machacek M, Hodgson L, Welch C, Elliott H, Pertz O, Nalbant P, et al. Coordination of Rho GTPase activities during cell protrusion. Nature. 2009;461(7260):99-103. Epub 2009/08/21. doi: 10.1038/nature08242. PubMed PMID: 19693013; PubMed Central PMCID: PMCPMC2885353.
13. Ridley AJ. Life at the leading edge. Cell. 2011;145(7):1012-22. Epub 2011/06/28. doi: 10.1016/j.cell.2011.06.010. PubMed PMID: 21703446.

14. Pelham RJ, Jr., Wang YL. Cell locomotion and focal adhesions are regulated by the mechanical properties of the substrate. Biol Bull. 1998;194(3):348-9; discussion 9-50. Epub 2001/09/07. doi: 10.2307/1543109. PubMed PMID: 11536880.

15. Discher DE, Janmey P, Wang YL. Tissue cells feel and respond to the stiffness of their substrate. Science. 2005;310(5751):1139-43. Epub 2005/11/19. doi: 10.1126/science.1116995. PubMed PMID: 16293750.

16. Bershadsky AD, Balaban NQ, Geiger B. Adhesion-dependent cell mechanosensitivity. Annu Rev Cell Dev Biol. 2003;19:677-95. Epub 2003/10/23. doi: 10.1146/annurev.cellbio.19.111301.153011. PubMed PMID: 14570586.

17. Bershadsky AD, Kozlov M, Geiger B. Adhesion-mediated mechanosensitivity: a time to experiment, and a time to theorize. Curr Opin Cell Biol. 2006;18(5):472-81. Epub 2006/08/26. doi: 10.1016/j.ceb.2006.08.012. PubMed PMID: 16930976.

18. Mitra SK, Hanson DA, Schlaepfer DD. Focal adhesion kinase: in command and control of cell motility. Nat Rev Mol Cell Biol. 2005;6(1):56-68. Epub 2005/02/03. doi: 10.1038/nrm1549. PubMed PMID: 15688067.

19. Schaller MD. Cellular functions of FAK kinases: insight into molecular mechanisms and novel functions. J Cell Sci. 2010;123(Pt 7):1007-13. Epub 2010/03/25. doi: 10.1242/jcs.045112. PubMed PMID: 20332118.

20. Plotnikov SV, Pasapera AM, Sabass B, Waterman CM. Force fluctuations within focal adhesions mediate ECM-rigidity sensing to guide directed cell migration. Cell. 2012;151(7):1513-27. Epub 2012/12/25. doi: 10.1016/j.cell.2012.11.034. PubMed PMID: 23260139; PubMed Central PMCID: PMCPMC3821979.

21. Tilghman RW, Parsons JT. Focal adhesion kinase as a regulator of cell tension in the progression of cancer. Semin Cancer Biol. 2008;18(1):45-52. Epub 2007/10/12. doi: 10.1016/j.semcancer.2007.08.002. PubMed PMID: 17928235; PubMed Central PMCID: PMCPMC2267763.

22. Mierke CT. The role of focal adhesion kinase in the regulation of cellular mechanical properties. Phys Biol. 2013;10(6):065005. Epub 2013/12/07. doi: 10.1088/1478-3975/10/6/065005. PubMed PMID: 24304934.

23. Tomakidi P, Schulz S, Proksch S, Weber W, Steinberg T. Focal adhesion kinase (FAK) perspectives in mechanobiology: implications for cell behaviour. Cell Tissue Res. 2014;357(3):515-26. Epub 2014/07/06. doi: 10.1007/s00441-014-1945-2. PubMed PMID: 24988914.

24. Urciuoli E, Peruzzi B. Involvement of the FAK Network in Pathologies Related to Altered Mechanotransduction. Int J Mol Sci. 2020;21(24). Epub 2020/12/17. doi: 10.3390/ijms21249426. PubMed PMID: 33322030; PubMed Central PMCID: PMCPMC7764271.

25. Wang HB, Dembo M, Hanks SK, Wang Y. Focal adhesion kinase is involved in mechanosensing during fibroblast migration. Proceedings of the National Academy of Sciences of the United States of America. 2001;98(20):11295-300. Epub 2001/09/27. doi: 10.1073/pnas.201201198. PubMed PMID: 11572981; PubMed Central PMCID: PMCPMC58723.

26. Armaiz-Pena GN, Allen JK, Cruz A, Stone RL, Nick AM, Lin YG, et al. Src activation by beta-adrenoreceptors is a key switch for tumour metastasis. Nat Commun. 2013;4:1403. Epub 2013/01/31. doi: 10.1038/ncomms2413. PubMed PMID: 23360994; PubMed Central PMCID: PMCPMC3561638.

27. Abrahamsen H, Vang T, Tasken K. Protein kinase A intersects SRC signaling in membrane microdomains. The Journal of biological chemistry. 2003;278(19):17170-7. Epub 2003/02/28. doi: 10.1074/jbc.M211426200. PubMed PMID: 12606547.

28. Beristain AG, Molyneux SD, Joshi PA, Pomroy NC, Di Grappa MA, Chang MC, et al. PKA signaling drives mammary tumorigenesis through Src. Oncogene. 2015;34(9):1160-73. Epub 2014/03/26. doi: 10.1038/onc.2014.41. PubMed PMID: 24662820.

29. Howe AK, Juliano RL. Regulation of anchorage-dependent signal transduction by protein kinase A and p21-activated kinase. Nature cell biology. 2000;2(9):593-600. Epub 2000/09/12. doi: 10.1038/35023536. PubMed PMID: 10980699.

30. Jin H, Garmy-Susini B, Avraamides CJ, Stoletov K, Klemke RL, Varner JA. A PKA-Csk-pp60Src signaling pathway regulates the switch between endothelial cell invasion and cell-cell adhesion during vascular sprouting. Blood. 2010;116(25):5773-83. Epub 2010/09/10. doi: 10.1182/blood-2010-07-296210. PubMed PMID: 20826718; PubMed Central PMCID: PMCPMC3031419.

31. Sanchez-Collado J, Lopez JJ, Jardin I, Camello PJ, Falcon D, Regodon S, et al. Adenylyl Cyclase Type 8 Overexpression Impairs Phosphorylation-Dependent Orai1 Inactivation and Promotes Migration in MDA-MB-231 Breast Cancer Cells. Cancers (Basel). 2019;11(11). Epub 2019/10/28. doi: 10.3390/cancers11111624. PubMed PMID: 31652779; PubMed Central PMCID: PMCPMC6893434.

32. Schmitt JM, Stork PJ. PKA phosphorylation of Src mediates cAMP's inhibition of cell growth via Rap1. Mol Cell. 2002;9(1):85-94. Epub 2002/01/24. doi: 10.1016/s1097-2765(01)00432-4. PubMed PMID: 11804588.

33. Vang T, Torgersen KM, Sundvold V, Saxena M, Levy FO, Skalhegg BS, et al. Activation of the COOH-terminal Src kinase (Csk) by cAMP-dependent protein kinase inhibits signaling through the T cell receptor. J Exp Med. 2001;193(4):497-507. Epub 2001/02/22. doi: 10.1084/jem.193.4.497. PubMed PMID: 11181701; PubMed Central PMCID: PMCPMC2195911.

34. Yeo MG, Oh HJ, Cho HS, Chun JS, Marcantonio EE, Song WK. Phosphorylation of Ser 21 in Fyn regulates its kinase activity, focal adhesion targeting, and is required for cell migration. J Cell Physiol. 2011;226(1):236-47. Epub 2010/07/27. doi: 10.1002/jcp.22335. PubMed PMID: 20658524.

35. Schmoker AM, Barritt SA, Weir ME, Mann JE, Hogan TC, Ballif BA, et al. Fyn Regulates Binding Partners of Cyclic-AMP Dependent Protein Kinase A. Proteomes. 2018;6(4). Epub 2018/10/03. doi: 10.3390/proteomes6040037. PubMed PMID: 30274258; PubMed Central PMCID: PMCPMC6313912.

36. Caldwell GB, Howe AK, Nickl CK, Dostmann WR, Ballif BA, Deming PB. Direct modulation of the protein kinase A catalytic subunit α by growth factor receptor tyrosine kinases. Journal of cellular biochemistry. 2012;113(1):39-48. Epub 2011/08/26. doi: 10.1002/jcb.23325. PubMed PMID: 21866565; PubMed Central PMCID: PMCPMC3245341.

37. Puleo JI, Parker SS, Roman MR, Watson AW, Eliato KR, Peng L, et al. Mechanosensing during directed cell migration requires dynamic actin polymerization at focal adhesions. J Cell Biol. 2019;218(12):4215-35. Epub 2019/10/09. doi: 10.1083/jcb.201902101. PubMed PMID: 31594807; PubMed Central PMCID: PMCPMC6891092.

38. Romero S, Le Clainche C, Gautreau AM. Actin polymerization downstream of integrins: signaling pathways and mechanotransduction. Biochem J. 2020;477(1):1-21. Epub 2020/01/09. doi: 10.1042/BCJ20170719. PubMed PMID: 31913455.

39. Howe AK, Hogan BP, Juliano RL. Regulation of vasodilator-stimulated phosphoprotein phosphorylation and interaction with Abl by protein kinase A and cell adhesion. The Journal of biological chemistry. 2002;277(41):38121-6. Epub 2002/06/28. doi: 10.1074/jbc.M205379200. PubMed PMID: 12087107.

40. McKenzie AJ, Hicks SR, Svec KV, Naughton H, Edmunds ZL, Howe AK. The mechanical microenvironment regulates ovarian cancer cell morphology, migration, and spheroid disaggregation. Sci Rep. 2018;8(1):7228. Epub 2018/05/10. doi: 10.1038/s41598-018-25589-0. PubMed PMID: 29740072; PubMed Central PMCID: PMCPMC5940803.

41. Dumbauld DW, Shin H, Gallant ND, Michael KE, Radhakrishna H, Garcia AJ. Contractility modulates cell adhesion strengthening through focal adhesion kinase and assembly of vinculin-containing focal adhesions. J Cell Physiol. 2010;223(3):746-56. Epub 2010/03/06. doi: 10.1002/jcp.22084. PubMed PMID: 20205236; PubMed Central PMCID: PMCPMC2874193.

42. Pasapera AM, Schneider IC, Rericha E, Schlaepfer DD, Waterman CM. Myosin II activity regulates vinculin recruitment to focal adhesions through FAK-mediated paxillin phosphorylation. J Cell Biol. 2010;188(6):877-90. Epub 2010/03/24. doi: 10.1083/jcb.200906012. PubMed PMID: 20308429; PubMed Central PMCID: PMCPMC2845065.

43. Roberts WG, Ung E, Whalen P, Cooper B, Hulford C, Autry C, et al. Antitumor activity and pharmacology of a selective focal adhesion kinase inhibitor, PF-562,271. Cancer Res. 2008;68(6):1935-44. Epub 2008/03/15. doi: 10.1158/0008-5472.CAN-07-5155. PubMed PMID: 18339875.

44. Golubovskaya VM, Nyberg C, Zheng M, Kweh F, Magis A, Ostrov D, et al. A small molecule inhibitor, 1,2,4,5-benzenetetraamine tetrahydrochloride, targeting the y397 site of focal adhesion kinase decreases tumor growth. J Med Chem. 2008;51(23):7405-16. Epub 2008/11/08. doi: 10.1021/jm800483v. PubMed PMID: 18989950; PubMed Central PMCID: PMCPMC2662449.

45. Hochwald SN, Nyberg C, Zheng M, Zheng D, Wood C, Massoll NA, et al. A novel small molecule inhibitor of FAK decreases growth of human pancreatic cancer. Cell Cycle. 2009;8(15):2435-43. Epub 2009/07/03. doi: 10.4161/cc.8.15.9145. PubMed PMID: 19571674; PubMed Central PMCID: PMCPMC4824314.

46. O'Brien S, Golubovskaya VM, Conroy J, Liu S, Wang D, Liu B, et al. FAK inhibition with small molecule inhibitor Y15 decreases viability, clonogenicity, and cell attachment in thyroid cancer cell lines and synergizes with targeted therapeutics. Oncotarget. 2014;5(17):7945-59. Epub 2014/10/04. doi: 10.18632/oncotarget.2381. PubMed PMID: 25277206; PubMed Central PMCID: PMCPMC4202172.

47. Tanjoni IW, C.; Uryu, S.; Tomar, A.; Nam, J.; Mielgo, A.; Lim, S.; Liang, C.; Koenig, M.; Sun, C.; Patel, N.; Kwok, C.; McMahon, G.; Stupack, D.G.; Schlaepfer, D.D. PND-1186 FAK inhibitor selectively promotes tumor cell apoptosis in three-dimensional environments Cancer Biol Ther. 2010;9(10):764-77. PubMed Central PMCID: PMC PMC2933317

48. Sieg DJ, Hauck CR, Schlaepfer DD. Required role of focal adhesion kinase (FAK) for integrin-stimulated cell migration. J Cell Sci. 1999;112 (Pt 16):2677-91. Epub 1999/07/22. PubMed PMID: 10413676.

49. Hung WC, Yang JR, Yankaskas CL, Wong BS, Wu PH, Pardo-Pastor C, et al. Confinement Sensing and Signal Optimization via Piezo1/PKA and Myosin II Pathways. Cell Rep. 2016;15(7):1430-41. Epub 2016/05/11. doi: 10.1016/j.celrep.2016.04.035. PubMed PMID: 27160899; PubMed Central PMCID: PMCPMC5341576.

50. Sieg DJ, Ilic D, Jones KC, Damsky CH, Hunter T, Schlaepfer DD. Pyk2 and Src-family protein-tyrosine kinases compensate for the loss of FAK in fibronectin-stimulated signaling events but Pyk2 does not fully function to enhance FAK- cell migration. EMBO J. 1998;17(20):5933-47. Epub 1998/10/17. doi: 10.1093/emboj/17.20.5933. PubMed PMID: 9774338; PubMed Central PMCID: PMCPMC1170921.

51. Weis SM, Lim ST, Lutu-Fuga KM, Barnes LA, Chen XL, Göthert JR, et al. Compensatory role for Pyk2 during angiogenesis in adult mice lacking endothelial cell FAK. J Cell Biol. 2008;181(1):43-50. Epub 2008/04/09. doi: 10.1083/jcb.200710038. PubMed PMID: 18391070; PubMed Central PMCID: PMCPMC2287283.

52. Parsons JT. Focal adhesion kinase: the first ten years. J Cell Sci. 2003;116(Pt 8):1409-16. Epub 2003/03/18. doi: 10.1242/jcs.00373. PubMed PMID: 12640026.

53. Sood AK, Armaiz-Pena GN, Halder J, Nick AM, Stone RL, Hu W, et al. Adrenergic modulation of focal adhesion kinase protects human ovarian cancer cells from anoikis. J Clin Invest. 2010;120(5):1515-23. Epub 2010/04/15. doi: 10.1172/JCI40802. PubMed PMID: 20389021; PubMed Central PMCID: PMCPMC2860925.

54. Swaney JS, Patel HH, Yokoyama U, Head BP, Roth DM, Insel PA. Focal adhesions in (myo)fibroblasts scaffold adenylyl cyclase with phosphorylated caveolin. The Journal of biological chemistry. 2006;281(25):17173-9. Epub 2006/04/19. doi: 10.1074/jbc.M513097200. PubMed PMID: 16618703.

55. Schlaepfer DD, Hauck CR, Sieg DJ. Signaling through focal adhesion kinase. Prog Biophys Mol Biol. 1999;71(3-4):435-78. Epub 1999/06/04. doi: 10.1016/s0079-6107(98)00052-2. PubMed PMID: 10354709.

56. Schaller MD. Biochemical signals and biological responses elicited by the focal adhesion kinase. Biochimica et biophysica acta. 2001;1540(1):1-21. Epub 2001/07/31. doi: 10.1016/s0167-4889(01)00123-9. PubMed PMID: 11476890.

57. Guan JL. Integrin signaling through FAK in the regulation of mammary stem cells and breast cancer. IUBMB Life. 2010;62(4):268-76. Epub 2010/01/27. doi: 10.1002/iub.303. PubMed PMID: 20101634; PubMed Central PMCID: PMCPMC2848709.

58. Golubovskaya VM, Cance WG. Focal adhesion kinase and p53 signaling in cancer cells. Int Rev Cytol. 2007;263:103-53. Epub 2007/08/30. doi: 10.1016/S0074-7696(07)63003-4. PubMed PMID: 17725966.

59. Kleinschmidt EG, Schlaepfer DD. Focal adhesion kinase signaling in unexpected places. Curr Opin Cell Biol. 2017;45:24-30. Epub 2017/02/19. doi: 10.1016/j.ceb.2017.01.003. PubMed PMID: 28213315; PubMed Central PMCID: PMCPMC5482783.

60. Broussard JA, Rappaz B, Webb DJ, Brown CM. Fluorescence resonance energy transfer microscopy as demonstrated by measuring the activation of the serine/threonine kinase Akt. Nat Protoc. 2013;8(2):265-81. Epub 2013/01/12. doi: 10.1038/nprot.2012.147. PubMed PMID: 23306460; PubMed Central PMCID: PMCPMC3756929.

Figure 1

Figure 2

Figure 3

Figure 4

Supp. Figure 1

Supp. Figure 2

Supp. Figure 3

Supp. Figure 4

A

Whole Cell PKA Activity

B

Leading Edge PKA Activity

Supp. Figure 5