TransCistor reveals the landscape of *cis*-regulatory long noncoding RNAs

Authors

Panagiotis Chouvardas[123], Marc Zimmerli[#12], Daniel Hanhart[#23], Mario Moser[#12], Hugo Guillen-Ramirez[1245], Sanat Mishra[6], Roberta Esposito[12], Taisia Polidori[12], Maro Widmer[12], Raquel García-Pérez[7], Marianna Kruithof-de Julio[23], Dmitri Pervouchine[8], Marta Melé[7], Rory Johnson*[1245]

1. Department of Medical Oncology, Inselspital, Bern University Hospital, University of Bern, 3010 Bern, Switzerland.

2. Department for BioMedical Research, University of Bern, 3008 Bern, Switzerland.

3. Department of Urology, Inselspital, Bern University Hospital, Bern, Switzerland.

4. School of Biology and Environmental Science, University College Dublin, Dublin D04 V1W8, Ireland.

5. Conway Institute for Biomolecular and Biomedical Research, University College Dublin, Dublin D04 V1W8, Ireland.

6. Indian Institute of Science Education and Research, Mohali, India.

7. Life Sciences Department, Barcelona Supercomputing Centre, 08034, Barcelona, Spain.

8. Center of Life Sciences, Skolkovo Institute of Science and Technology, Moscow, Russia.

# Equal contribution.

* Correspondence to rory.johnson@dbmr.unibe.ch

Abstract

Long noncoding RNAs (lncRNAs) can positively and negatively regulate expression of target genes encoded in *cis*. However, the extent, characteristics and mechanisms of such *cis*-regulatory lncRNAs (*cis*-lncRNAs) remain obscure. Until now, they have been defined using inconsistent, *ad hoc* criteria that can result in false-positive predictions. Here, we introduce *TransCistor*, a framework for defining and identifying *cis*-lncRNAs based on enrichment of targets amongst proximal genes. Using transcriptome-wide perturbation experiments for 190 human and 133 mouse lncRNAs, we provide the first large-scale view of *cis*-lncRNAs. Our results ascribe *cis*-regulatory activity to only a small fraction (~10%) of lncRNAs, with a prevalence of activators over repressors. *Cis*-lncRNAs are detected at similar rates by RNA interference (RNAi) and antisense oligonucleotide (ASO) perturbations. We leverage this cis-lncRNA catalogue to evaluate mechanistic models for *cis*-lncRNAs involving enhancers and chromatin folding. Thus, TransCistor places *cis*-regulatory lncRNAs on a quantitative foundation for the first time.

Main

The first characterised long noncoding RNAs (lncRNAs), *H19* and *XIST,* were both found to have *cis*-regulatory activity: their perturbation by loss-of-function (LOF) led to increased expression of protein-coding genes encoded "in *cis*" - i.e. within a relatively short linear distance on the same chromosome[1,2]. Genes whose expression responds to lncRNA LOF are considered "targets" of that lncRNA, while the direction of this change (up or down) defines the lncRNA as a "repressor" or "activator", respectively. Since then, numerous more *cis*-regulatory lncRNAs have been reported [3,4]. Conversely, other lncRNAs have no apparent positional preference for their targets, and are termed *trans*-lncRNAs [5]. This *cis/trans* duality provides a fundamental framework for understanding regulatory lncRNAs[6], yet the global prevalence of *cis*- and *trans*-regulatory lncRNAs remains poorly defined.

Within reported *cis*-lncRNAs there appears to be great diversity, in terms of regulatory activity (activators and repressors), distance of the target (ranging from one hundred basepairs[4] to hundreds of kilobases[7]) and number of targets (one[4] to many[8]). Two overarching molecular mechanisms have been proposed for *cis*-lncRNAs: enhancer elements and chromatin folding[9]. Some *cis*-activating lncRNAs, termed "enhancer lncRNAs" (e-lncRNAs), have been found to overlap DNA-encoded enhancer elements[10–12], similar to lncRNAs more generally[13]. The expression and splicing of the e-lncRNA transcripts correlate with enhancer activity, implying that RNA processing somehow promotes target gene activation. Similarly, it has been proposed that *cis*-lncRNAs find their targets via spatial proximity, determined by chromatin looping or within the confines of local topologically-associating domains (TADs)[6]. An attractive corollary of these models is that *cis*-regulatory lncRNAs may act via non-sequence dependent mechanisms, perhaps involving phase separation[14] and local concentration gradients[15]. It has recently been posited that lncRNAs proceed through an evolutionary trajectory commencing with fortuitous *cis*-regulatory activity before acquiring targeting capabilities and graduating to *trans*-regulation[16]. Nonetheless, these conclusions are drawn from piecemeal studies of individual lncRNAs, and a holistic view of *cis* and *trans* lncRNAs, the features that distinguish them, and resulting clues to their molecular mechanisms and biological significance, await a comprehensive catalogue of *cis*-lncRNAs.

Despite their importance, we lack a rigorous and agreed definition for *cis*-lncRNAs. Until now,
they have been defined by the existence of ≥1 proximal targets. Targets are defined as those
whose expression changes in response to lncRNA LOF, as measured using single-gene (RT-
PCR) or whole-transcriptome (RNA-seq, CAGE, microarray) techniques[3,5,17]. "Proximity" is
defined on a case-by-case basis, using a wide range of windows spanning 10^2 to 10^5 bp[7]. A single
proximal target is usually considered sufficient. The problem with this approach is that, as the total
number of targets and/or *cis*-window size increase, so will the chance of observing ≥1 *cis*-target
genes by random chance.

In this study, we consider *cis*-lncRNAs from a quantitative perspective. We show that conventional
definitions are prone to high false positive rates. We introduce statistical methods for definition of
cis-lncRNAs at controlled false discovery rates, and use them to classify regulatory lncRNAs
across hundreds of perturbation datasets. Our results enable us to estimate, for the first time, the
prevalence of *cis*-acting lncRNAs and evaluate hypotheses regarding their molecular
mechanisms of action.

Results

A quantitative definition of *cis*-lncRNAs

We began by investigating the usefulness of the present "naïve" definition of *cis*-lncRNAs, based
on the existence of ≥1 target gene within a local genomic window. To create a dataset of lncRNA-
target relationships, we collected 382 lncRNA LOF experiments targeting 188 human lncRNAs
from a mixture of sources (including the recently published dataset of ASO knockdowns in human
dermal fibroblasts from the FANTOM consortium[18]) and 137 experiments for 131 lncRNAs in
mouse (Figure 1A). To this we added 6 hand-curated LOF experiments targeting previously
reported *cis*-acting lncRNAs (*UMLILO*, *XIST x2*, *Chaserr*, *Paupar* and *Dali*). We employed a
functional definition of "targets", as genes whose steady-state levels significantly change in
response to a given lncRNA's LOF (Figure 1B). We further define targets as activated or
repressed, where they decrease / increase in response to lncRNA LOF, respectively. Finally, we
define genes to be proximal / distal, if their annotated transcription start site (TSS) lies inside /
outside a defined distance window of the lncRNA's TSS, respectively. In the dataset, 126 lncRNAs
were represented by ≥1 independent experiments, and the median number of target genes
identified per experiment was 65 (Supplementary Figure S1).

Using a range of *cis*-window sizes from 50 kb to 1 Mb centred on the lncRNAs' TSSs, we
evaluated the fraction of lncRNAs that would be defined as *cis*-lncRNAs under the naïve definition.
This approach defines ~2 to 12% of lncRNAs as *cis*-regulators (Figure 1C, line). To test whether
this rate is greater than random chance, we shuffled the target / non-target labels of all protein-
coding genes and repeated this analysis. The rate of *cis*-lncRNA predictions in this random data
overlapped the true rates in all windows (Figure 1C, boxplots), suggesting that the conventional
definition of *cis*-lncRNAs yields high rates of false-positive predictions.

To overcome this issue, we adopted the following definition for *cis*-lncRNAs: cis-*lncRNAs are*
those whose targets are significantly enriched amongst proximal genes. This definition has the
advantage of being quantitative and statistically testable. Below we incorporate this definition into
two alternative methods for identifying *cis*-lncRNAs.

TransCistor: digital and analogue identification of *cis*-lncRNAs

The first method employs a simple definition of proximal genes, being those whose TSS falls within a defined window centred on the lncRNA TSS. We developed a pipeline, *TransCistor-digital*, which takes as input a processed whole-transcriptome list of target genes ("regulation file"), and tests for statistical enrichment in proximal genes using the hypergeometric distribution (Figure 1D). Although in principle any sized window may be used, we reasoned that the most biologically-meaningful would be the local TAD, in line with previous studies[19]. Chromatin folding varies to an extent between cell types[20]. Therefore, TransCistor-digital calculates enrichment across a set of experimentally-defined cell-type-specific TADs (40 human, 3 mouse)[21] and aggregates the resulting p-values by their harmonic mean.

The above fixed-window approach is intuitive yet has drawbacks. Several reported *cis*-lncRNAs have individual targets that are not immediately adjacent[7], and might be overlooked by the fixed-window approach. Furthermore, many lncRNAs may have no neighbouring genes in their local TAD, or no identified local TAD. Therefore, we developed an alternative method that dispenses with fixed windows, while still examining proximity biases in targets. This method, *TransCistor-analogue*, defines a distance statistic as the mean TSS-to-TSS distance of all same-chromosome targets of a given lncRNA (Figure 1E). To estimate statistical significance, a null distribution is calculated by randomisation of target labels (Figure 1E). Now, *cis*-lncRNAs are defined as those having a distance statistic that is lower than a majority of simulations.

We sought to test the performance of TransCistor-digital and evaluate the global landscape of *cis*-lncRNAs. After filtering out unusable datasets (having no *cis*-targets or no overlapping TAD), 168 datasets remained. The majority of p-values produced by this analysis follow the null distribution, underlining the conservative statistical behaviour of TransCistor (Figure 2A,B). We discovered 19 *cis*-acting lncRNAs (12 activators, 7 repressors), with a relatively relaxed false discovery rate (FDR) threshold of 0.5, while no *cis*-lncRNAs are simultaneously classified as activator and repressor (Figure 2C). Amongst the top-ranked *cis*-lncRNAs is *UMLILO*, previously described to activate multiple genes in its local genomic neighbourhood[8]. *UMLILO* exhibits a significant enrichment of activated targets amongst proximal genes, which is not observed for repressed targets (Figure 2D,E). Analysis of the entire perturbation dataset by TransCistor-analogue identified 20 *cis*-lncRNAs (15 activators, 5 repressors, FDR≤0.5). Statistical behaviour is good (Figure 2F,G), while *cis*-lncRNAs once again are cleanly split between activators and repressors (Figure 2H).

The usefulness of these methods is supported by their internal and external consistency. Together, the TransCistor approaches correctly identify previously-described *cis*-activators *H19*[22], *JPX*[23]*, Evx1os*[24] and *DA125942*[25] amongst the top ranked *cis*-activators, while *XIST* is amongst the top repressors[26] (Figure 2J). Both human and mouse orthologues of *CHASERR* (ENSG00000272888) are identified as *cis-repressors*[4]. Of the hits, two are concordantly classified by ≥1 independent perturbation experiments (*XIST* & *DNAAF3-AS1* classified as *cis*-repressors based on two separate experiments each) (Figure 2I,J). We observed agreement between the two TransCistor methods, with 6 *cis*-lncRNAs in common (*DA125942, linc1427, RAD51-AS1, H19, Xist, DNAAF3-AS1*) (p-value < 0.05, hypergeometric test) (Figure 3A).

TransCistor predicted *cis*-regulatory activity for a number of known lncRNAs that have never been described as such in prior literature. These include *BANCR* (*cis*-activator), and *SBF2-AS1*,

LASTR, NORAD, *DANCR* (*cis*-repressors). However, the latter two are only identified in one out
of multiple independent perturbation experiments (6/5 for *NORAD/DANCR* respectively). In the
case of *DANCR*, the *cis* definition arises from the repression of two same-strand small RNAs
(has-mir-4449, SNORA26). It is not yet clear if these results reflect false-positive or false-negative
predictions. To investigate this, we merged all hits across experiments and repeated the analysis,
but here we found no *cis* signal, suggesting that they are false-positive predictions. On the other
hand, analysis of an independent dataset for *SBF2-AS1* from different cells (A549 lung
adenocarcinoma) and perturbation (siRNA) yielded concordant *cis*-repressor prediction from
TransCistor-analogue (Supplementary Figure S2). This is strong evidence that *SBF2-AS1* is a
novel *cis*-repressive lncRNA.

Surprisingly, TransCistor failed to find evidence supporting two previously reported *cis*-lncRNAs,
Paupar[27] and *Dali*[28]. Inspection of the originating microarray data revealed that, for neither case,
do the claimed *cis*-target genes pass cutoffs of differential expression (Supplementary Figure S3).

Overall, if we consider lncRNAs where at least one method in one dataset is called as *cis*-acting,
then our data implicates 10% (33/323) of lncRNAs as *cis*-regulators (Figure 3B). When broken
down by direction of regulation, we find that 7% (23) of these are activators and 3% (10) are
repressors, of which none overlap. We henceforth define the remaining 290 tested lncRNAs as
trans-lncRNAs. Together, these findings indicate that TransCistor is capable of identifying known
and novel *cis*-lncRNAs, and a relatively small minority of lncRNAs display significant *cis*-activity.

TransCistor identifies *cis*-lncRNA independently of perturbation technology

The perturbation experiments contained a mixture of RNA interference (RNAi) and antisense
oligonucleotide (ASO) LOF perturbations. While early experiments were performed using the two
RNAi approaches of siRNA and shRNA, it is widely thought that these principally degrade targets
in the cytoplasm[29,30] or ribosome[31]. In contrast, ASOs are becoming the method of choice to knock
down lncRNAs, since they are thought to act on nascent RNA in chromatin[32]. If correct, then one
would expect ASO perturbations to have greater power to discover *cis*-lncRNAs. To test this, we
compared predictions from each perturbation technology (Figure 3C). Surprisingly, we observed
broadly similar rates of *cis*-lncRNA identification between perturbation methods. However, ASO
experiments discover similar rates of activators and repressors, while RNAi perturbations yield an
apparent excess of activators over repressors.

We conclude that TransCistor is capable of discovering *cis*-lncRNAs across perturbation types
While the small numbers preclude statistical confidence, these findings broadly support the use
of RNAi in targeting nuclear lncRNAs and identifying *cis*-lncRNAs, although the possibility for
perturbation-specific biases should be further investigated.

Association of *cis*-lncRNAs with enhancer elements

It has been widely speculated that *cis*-lncRNAs, particularly activators (ie e-lncRNAs), act in
concert with DNA enhancer elements to upregulate target gene expression[3,9,12]. Our catalogue of
cis-lncRNAs represents an opportunity to independently test this. We calculated the rate of
overlap of lncRNAs with enhancers using epigenomics data across human tissues (Figure 4A,
Supplementary Figure S4). Analyses were performed at a variety of epigenome thresholds (the
minimum number of samples required to define a given epigenomic state) and window sizes (the
distance from the lncRNA TSS to the nearest epigenome element).

This analysis revealed several intriguing relationships between *cis*-lncRNAs and enhancer
elements. First, we noted an enrichment of super-enhancers at the TSS of *cis*-activator lncRNAs
(boxed, Figure 4B). Inspection of overlaps at other thresholds and window sizes revealed a similar
effect (Supplementary Figure S4). Second, we observed an enrichment of Enhancer(1) elements
around the TSS of *cis*-repressor lncRNAs (boxed, Figure 4B), which similarly was corroborated
by analyses with a variety of thresholds (Supplementary Figure S4). More broadly, we observed
a generalised enrichment of various enhancer element annotations with *cis*-lncRNAs (Figure 4B,
left column). However, we do not observe a preference for such enrichment in *cis*-activator over
cis-repressor lncRNAs (Figure 4B, right column). Overall, within the limits of statistical power
given our relatively small sample size, these findings are consistent with a relationship between
cis-lncRNAs and enhancer elements.

Some *cis*-lncRNAs are brought into spatial proximity to their targets by chromatin looping

A second key mechanistic model posits that regulatory interactions between *cis*-lncRNAs and
target genes are effected by close spatial proximity, brought about by chromatin looping (Figure
5A). To measure proximity, we utilised published Hi-C interactions from a range of human cell
lines[33]. We evaluated the importance of proximity for regulatory targeting, by combining an
asymptotic regression model to predict an "expected interaction" at a given linear genomic
distance, with a logistic regression model to evaluate whether strong deviations from this
expectation were indicative of targeting (Figure 4C). This approach revealed a significant (p-value
≤ 0.1) contribution of spatial proximity to targeting for two *cis*-activator lncRNAs: *UMLILO* (8 cell
lines) and *DA125942* (1 cell line) (Figure 4D). In both cases, previous studies have implicated
chromatin looping in target identification [8,25]. An excellent example is represented by HUVEC
cells, where *UMLILO* target genes tend to be located in higher proximity (Interaction, *y*-axis),
compared to other non-targets at similar distances in linear DNA (*x*-axis) (Figure 4E). An
alternative inverse square model yielded the same two lncRNAs (Supplementary Figure S5).
Together, this indicates that for a subset of *cis*-lncRNAs, spatial proximity may determine identity
of target genes.

Discussion

We have described *TransCistor*, a modular quantitative method for identification of *cis*-regulatory
lncRNAs. We applied it to a corpus of perturbation datasets to create the first large-scale survey
of *cis*-regulatory RNAs. We evaluated the performance of *TransCistor* in light of the present state-
of-the-art and used the resulting catalogue of *cis*-lncRNAs to address fundamental questions
regarding their prevalence and molecular mechanisms.

TransCistor-digital and -analogue represent practical tools for *cis*-lncRNA discovery. Previous
studies used a "naïve" criterion of ≥1 *cis*-target gene within an arbitrarily-sized window; however,
we show that this method is prone to predominantly false-positive predictions at ≥50 kb windows.
TransCistor improves on this situation by making predictions at a defined false discovery rate
(FDR). The two distinct statistical methods are designed to capture a range of *cis*-activity, from
lncRNAs regulating the most proximal neighbour gene's expression within the local TAD, such as
Chaserr[4], to those regulating a more distal target amongst other non-target genes, such as
CCAT1-L[7]. The value of resulting predictions is supported by good statistical behaviour as judged
by quantile-quantile (QQ) analysis, consistency between methods and datasets, and recall of

numerous known *cis*-lncRNAs, including founding members *H19* and *XIST*. TransCistor is made
available both as a webserver and standalone software. It is compatible with a wide range of input
data, since "regulation" files can be readily generated from any experimental dataset comprising
lncRNA perturbation and global readout of gene expression changes, including two decades of
experiments from microarrays to RNA-sequencing and future parallelised CRISPR LOF methods
such as Perturb-Seq[34].

This work builds on important previous attempts to comprehensively discover *cis*-regulatory
lncRNAs. Basu and Larsson utilised gene expression correlation as a means for inferring
candidate *cis*-regulatory relationships[35]. Very recently, de Hoon and colleagues employed
genome-wide RNA-chromatin and chromatin folding to train a predictive model for *cis*-regulatory
lncRNAs[36]. While these methods are valuable, they infer target genes based on indirect correlates
of *cis*-regulation, which may not reflect causation[37]. Furthermore, we failed to find evidence that
chromatin folding links *cis*-lncRNAs to their target genes in all cases. What distinguishes
TransCistor from these approaches, is its use of LOF perturbations to directly identify gene
targets. We argue that, due to its direct and functional nature, this approach should be considered
the gold standard evidence for defining *cis*-regulatory relationships.

Our results afford important insights into the regulatory lncRNA landscape. Notwithstanding the
caveats discussed above, we provide the first global estimate of *cis*-lncRNA prevalence,
suggesting they represent a modest fraction (10%) of the total, with a slight prevalence of
activators over repressors. These values are certainly impacted by a variety of errors discussed
above, which we hope will be corrected by future, larger-scale studies. The preponderance of *cis*-
activators may be an artefact of RNAi perturbations, which appear to yield an excess of activators
over repressors, with no apparent explanation yet. Our results shed light on *cis*-lncRNAs'
molecular mechanisms, finding evidence supporting their relationship with enhancer elements
and, in some cases, a preference to loop into spatial proximity to targets. Surprisingly, we
observed evidence that enhancers are associated with both activator and repressor lncRNAs.

Finally, it is worth revisiting the assumptions we make when interpreting lncRNA perturbation
experiments. These involve a small oligonucleotide with perfect sequence complementarity to a
lncRNA target in RNA *and* DNA, and assess the outcome in terms of steady state RNA levels.
Two key assumptions are made. Firstly, any change in downstream gene expression is assumed
to occur through changes in the targeted lncRNA transcript. It is well known that small oligos are
not only capable of hybridising to genomic DNA[38], but also to affect local chromatin
modifications[39], raising the possibility of chromatin/DNA-mediated *cis*-regulatory mechanisms.
The second assumption is more fundamental: that, when local gene changes are observed to
occur, such changes reflect the *biological function* of the lncRNA[40,41]. The alternative explanation
is that perturbations of a lncRNA lead to changes to local gene expression, but that this is a by-
product of altering lncRNA expression (e.g. by disrupting local transcription factories), and that
the evolutionarily-selected function of the lncRNA is something quite different. In other words, is
observed *cis*-activity a reflection of genuine, adaptive biological regulatory pathway, or is it merely
a technical artefact without biological relevance? Testing these alternative explanations will be an
interesting challenge for the future, facilitated by the tools provided here.

Methods

TransCistor

TransCistor was developed under the R statistical software (v4.0). Gene locations were extracted from GENCODE annotation file in GTF format (v38 for human, v25 for mouse)[42] and were converted into a matrix. The TransCistor input consists of a "regulation file", containing all genes and a flag indicating their regulation status: 1 (upregulated after perturbation; repressed by the lncRNA), -1 (downregulated after perturbation; activated by the lncRNA) or 0 (not target). Regulation status can be defined by the user, and here is based on differential expression after lncRNA perturbation. The perturbed lncRNA itself is removed from the regulation file to avoid false positive predictions. Results are visualized with ggplot2 (v3.3.5), ggpubr (v0.4), pheatmap (v1.0.12) packages and custom in-house generated scripts.

TransCistor includes two modules; Digital and Analogue. TransCistor-digital defines *cis*-lncRNAs based on statistical overrepresentation of proximal targets, defined as targets in the same topologically associated domain (TAD) as the lncRNA. Membership of a TAD is defined based on a gene's TSS. Digital TransCistor utilizes a collection of TADs for human and mouse cell types accessed via the 3D-Genome Browser[21]. For each cell type, TransCistor identifies the lncRNA TAD, estimates the number of proximal (within TAD) and distal (outside TAD) targets / non-targets (separately for activated and repressed). Then, it tests for overrepresentation of proximal targets over distal targets by the hypergeometric test. The p-values for all the cell types are then integrated by their harmonic mean. P-values are corrected for multiple hypothesis testing using the False Discovery Rate (FDR) method and taking into account the experiments which show at least one proximal target. TransCistor-analogue evaluates whether the mean distance of targets from the same chromosome are closer than random chance. Distance is defined by TSS to TSS. Analysis is performed separately for activated and repressed targets. Then, the random distribution is calculated, by randomly shuffling the regulation flags on genes within the same chromosome, and recalculating the test statistic each time. By default, 1000 simulations are performed. Finally, the empirical p-value is calculated from the proportion of simulations with a statistic less than the true value.

Both modules of TransCistor are available as a standalone R package and along with all regulation files (https://github.com/pchouvardas/TransCistor) and Rshiny webserver (https://transcistor.unibe.ch/). The input comprises metadata about the lncRNA, and a regulation file containing target gene information that can be readily derived from any transcriptome-wide data including RNA-sequencing, CAGE and microarray experiments.

Collecting and processing perturbation datasets

The FANTOM perturbation datasets were downloaded from the Core FANTOM6 repository[18]. The differential expression results were transformed into regulation files by applying an adjusted p-value threshold of 0.05 and using custom bash scripts. The respective metadata were also downloaded from FANTOM6 and were integrated to the GENCODE annotation matrix. 31 perturbation experiments were removed because they target protein coding genes, and an additional 19 were removed because target lncRNAs had no ENSEMBL identifier. The LncRNA2Target datasets were downloaded from the webserver (Version 2.0)[43] and targets were defined by using an adjusted p-value cutoff of 0.05. The lncRNA locations were manually obtained

315 from the website or original publications, when necessary. The rest of the datasets were accessed
316 through the original publications and post-processed to generate the regulation files. All regulation
317 files are available from the project Github repository, linked above.

Analysis of chromatin states

Chromatin states annotations were retrieved from three sources: EpiMap[44], genoSTAN[45], and
dbSUPER[46]. EpiMap consists of 18 chromatin states across 833 samples, genoSTAN identifies
promoter and enhancer regions genome-wide across 127 samples, and dbSUPER aggregates
82234 human superenhancers from 102 cell types/tissues. The annotations were relabelled as
follows: Superenhancer – dbSUPER's superenhancers; Enhancer(1) – genoSTAN's enhancers;
Enhancer(2.1) – EpiMap's Genic enhancer 1; Enhancer(2.2) – EpiMap's Active enhancer 1;
Enhancer(2.3) – EpiMap's Weak enhancer; Promoter(1) – genoSTAN's promoters; and
Promoter(2) – EpiMap's Active TSS.

The human TSS annotations were intersected with chromatin states at several genomic windows
(1 bp, 100 bp, 1000 bp, 10000 bp) and a given state-TSS intersection was counted only if it was
present in more samples than a given threshold (0, 1, 5, or 10 samples). For each pair of genomic
window and filter, a contingency matrix was computed for each pair of predicted labels (*cis*-
activator vs. *cis*-repressor, *cis*-activator vs. *trans*, and *cis*-repressor vs. *trans*) or the grouped label
(*cis* vs. *trans*), counting the number of TSSs falling into each category. Fisher's exact test was
used to compute the p-value of each contingency matrix.

Chromatin folding analysis

HiC interaction data was obtained using the python package "hic-straw" (v1.2.1)
(https://github.com/aidenlab/straw), using human HiC datasets from Aiden laboratory[33]. The gene
coordinates reported by TransCistor were converted to hg19 to match the HiC data. The binning
resolution was set to 25 kb, and interaction scores were normalized by Knight-Ruiz matrix
balancing method. Due to gaps in the HiC matrices, ~7% of lncRNA: (non-/)target interactions
were approximated by using a "next best" pair of bins, for which an interaction score was available,
instead of the correct binning. In 6.8% of cases this only required replacing either one of the ideal
bins by a direct neighbour and for the remaining 0.2% either shifting both genes by one bin or one
of the genes by two bins. An estimate for the expected interaction at a given distance was then
calculated by fitting a regression model to the HiC data with the interaction score as the response
and the TSS distance between the two genes as the explanatory variable. An asymptotic
regression model was chosen for this step ('SSasymp' and 'nls' of the R base package 'stats'
v4.0.3). Due to model limitations only *cis*-lncRNAs identified by TransCistor digital were included
in this analysis. For 2/12 lncRNAs from this subgroup (*RAD51-AS1, NARF-AS2*), model
generation failed for one or more of the cell types. Modelling the interaction as a function of the
inverse square distance was also considered ('glm' also from 'stats'). This model had the
advantage of not failing for either combination of *cis*-lncRNA and cell type, but fit the data less
well and it had a clear bias to underestimate the interaction in close 2D proximity and overestimate
interaction further away (Supplementary Figure S5). The significance of interaction on the
targeting status was then assessed by fitting a logistic regression model to predict whether a gene
is a target of a given lncRNA based on the difference between observed and expected interaction
(again using the 'glm' function).

Figures

Figure 1

<u>Figure 1: TransCistor is a quantitative framework for classifying *cis*- and *trans*-regulatory</u>
<u>lncRNAs.</u>

A) Origin of the perturbation data: The y-axis displays the number of perturbation experiments
(left panel) or individual lncRNA genes (right panel) per model organism (x-axis). Note that the
difference between number of perturbation experiments & lncRNAs arises from the fact that many
lncRNA genes are represented by > 1 experiment. The bars are split/color-coded according to
the origin dataset. The 6 hand curated perturbation experiments are the ones targeting UMLILO,
XIST (2), Chaserr, Paupar and Dali. B) Definition of target genes: A target gene is defined as
one whose expression significantly changes after loss-of-function perturbation of a given lncRNA
(pink). The direction of that change (down/up) defines the target as activated/repressed (green,
orange), respectively. C) Evaluating accuracy of naïve *cis*-lncRNA definition: The plot displays
the number of lncRNAs classified as "*cis*-regulatory" using a definition of ≥1 proximal target genes
(*y*-axis), while varying the size of the genomic window (centred on the lncRNA TSS) within which
a target is defined as "proximal" (*x*-axis). Line: real data from Panel A; Boxplot: Simulations
created by 50 random shuffles of the target labels across all annotated genes. D) TransCistor-
digital method: TransCistor-digital evaluates the enrichment of targets (green) in proximal regions,
defined as those residing within the same topologically associating domain (TAD) as the lncRNA
TSS (pink) (left panel), compared to the background target rate in the rest of the genome ("Distal")
(centre panel). *Cis*-lncRNAs are defined as those having a significantly higher proximal target
rate, defined using hypergeometric test (right panel). E) TransCistor-analogue method: A
distance statistic is defined as the mean genomic distance (bp) of all targets (green) on the same
chromosome as the lncRNA (pink) (left panel). 1000 simulations are performed where target
labels are shuffled across genes within the same chromosome (centre panel). *Cis*-lncRNAs are
defined as those whose real statistic (dashed line) falls below the majority of simulations (right
panel).

Figure 2

A) **Digital − Activator**

B) **Digital − Repressor**

C)

D) **UMLILO:**
[Activator p = 0.004 | Repressor p = 0.905]

E) Chr4:72323028-75514764

F) **Analogue − Activator**

G) **Analogue − Repressor**

H)

I) **DNAAF3−AS1**
ASO-G0267577-01

J) **DNAAF3−AS1**
ASO-G0267577-03

Figure 2: Large scale classification of *cis*-lncRNAs in human and mouse.

A) Quantile-quantile plot displays the random expected (*x*-axis) and observed (*y*-axis) p-values for lncRNAs (points) tested for activated targets by TransCistor-digital. The grey diagonal *y*=*x* line indicates the expectation if no hits were present. B) As for (A), for TransCistor-digital and repressed targets. C) Comparison of activator and repressor activity detected by TransCistor-digital. For each lncRNA (points), their false-discovery rate (FDR)-adjusted significance is plotted on the *x*-axis (activator) and *y*-axis (repressor). Note the absence of lncRNAs that are both activators and repressors. D) *UMLILO*, an example *cis*-activator: The plot shows the number of

genes, divided by targets / non-targets (colour / grey), location (distal/proximal) and regulation
direction (activated/repressed). *UMLILO* is classified as a *cis*-activator, due to the significant
excess (8) of proximal activated targets. Statistical significance (uncorrected) is displayed above.
E) *UMLILO* genomic locus: Vertical bars denote gene TSS. Grey: non-targets; green: activated
targets; pink: *UMLILO*. F) As for (A), for TransCistor-analogue and activated targets. G) As for
(B), for TransCistor-analogue and repressed targets. H) As for (C), for TransCistor-analogue. I)
DNAAF3-AS1, an example *cis*-repressor identified by TransCistor-analogue. Shown is the target
distance statistic (*x*-axis) for real data (vertical bar) and simulations (boxes). The number of
simulations in each distance bin is displayed on the *y*-axis. J) As for (I), for a second perturbation
experiment.

Figure 3

Figure 3: Rate of *cis*-lncRNA across perturbations, datasets and species.

A) Summary of TransCistor results: The values represent numbers of experiments classified in the bins indicated on the two axes, at a cutoff of FDR ≤ 0.5. The names of lncRNA genes are displayed. Previously-described cis-lncRNAs are red. B) The rate of lncRNA genes defined to be *cis*-regulatory based on our analysis. Note that one single experiment is sufficient to label a lncRNA gene as *cis*-regulatory. C) The rate of experiments defined as *cis*-regulatory, broken down by perturbation method.

<u>Figure 4: Intersection of *cis*-lncRNAs with enhancer elements and generation of a chromatin</u>
<u>conformation based model.</u>

A) Method of calculating overlap by enhancer annotations (horizontal purple bars) of lncRNA TSS
(pink bar). Overlaps are considered while varying two important thresholds: numbers of individual
enhancer annotations are considered the minimum necessary (epigenome threshold) and various
sized windows around the TSS of overlap calculation between extended TSS region (span). Only
the TSS spans with overlaps in more samples than a given epigenome threshold are considered.
B) Enrichment results epigenome threshold=1 and span=100 bp. Rows show enrichment for
super-enhancer, enhancer, and promoter states while comparing the TSS according to their
mechanism of action (see Methods). P-values were computed using Fisher's Exact Test. C) A
model for proximity-driven target selection: (Left panel) Chromatin folding brings lncRNA (pink)
into spatial proximity with proximal genes, which are subsequently targeted (green). (Right panel)
Chromatin proximity maps, such as provided by HiC methodology, enable one to evaluate the
spatial proximity (*y*-axis) of targets, while normalising for confounder of linear 2D DNA distance
(*x*-axis). These parameters were modelled using an Asymptotic regression model (right panel,
inset). D) Evaluating the contribution of proximity to target selection in human cells: The model
significance of *cis*-lncRNAs (identified by TransCistor-digital) (*x*-axis) was evaluated across HiC
interaction data from a panel of human cell lines (*y*-axis). Colour scale shows uncorrected p-
values. Green cells indicate cases where target genes tend to be significantly more proximal than
non-targets. No cases of the inverse were observed. E) Example data for *UMLILO* in HUVEC
cells. Note that target genes (green) tend to be more spatially proximal (*y*-axis) than non-target
genes (grey) at a similar TSS-to-TSS genomic distance (*x*-axis).

 Supplementary Figures

Supplementary Figure 1

A

B

Number of detected targets

<u>Supplementary Figure S1: Summary statistics of perturbation datasets.</u>

A) Histogram displaying the numbers of separate perturbation experiments (*x*-axis) available for each lncRNA gene (*y*-axis). B) Histograms displaying the number of significantly changing genes (targets) (*x*-axis) for each perturbation experiment (*y*-axis). Regulated genes represent the union of activated and repressed genes.

Supplementary Figure 2

443 <u>Supplementary Figure S2: Analysis of *cis*-regulation by *SBF2-AS1* in independent datasets.</u>

444 (A) TransCistor-analogue results for FANTOM ASO knockdown targeting *SBF2-AS1* in human
445 dermal fibroblasts. B) As for (A), but for independent data from A549 cells treated with siRNA. C)
446 Numbers indicate the genes in each category, classified by their regulation in the two distinct
447 datasets in (A) and (B).

Supplementary Figure 3

Supplementary Figure S3: Analysis of *Dali* and *Paupar* target genes using public microarray data.

A) Global transcriptome changes upon *Dali* knockdown were obtained from Gene Expression Omnibus (https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE62031). B) Neighbour gene and putative target *Pou3f3* mRNA expression in control and *Dali* knockdown samples. C) Global transcriptome changes upon *Paupar* knockdown were obtained from Gene Expression Omnibus (https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE52569). D) Neighbour gene and putative target *Pax6* mRNA expression in control and *Paupar* knockdown samples.

Supplementary Figure 4

<u>Supplementary Figure S4: Enrichment of enhancers, super-enhancers and promoters in *cis*-</u>
<u>lncRNAs.</u> Each row represents a different enhancer annotation (see Methods). Columns
represent comparisons between indicated pairs of lncRNA classes. Heatmaps display the
enrichment of overlap at different genomic windows around the lncRNA TSSs (span, *x*-axis) and
the minimum number of observed samples required to define an enhancer (epigenome threshold,
y-axis). Statistical significance is indicated where p-value ≤ 0.05 by Fisher's exact test (1-sided),
and not corrected for multiple hypothesis testing.

<u>Supplementary Figure S5: Alternative inverse square model for target gene interaction.</u>

A) and B) are equivalent to main Figure 4 panels D and E respectively but using an inverse square
model instead of an asymptotic regression. C) Comparison of asymptotic (solid line) and inverse
square (dashed line) regression models.

Supplementary Data Files

<u>Data File 1: Information for all lncRNAs studied in this work.</u>

Acknowledgements

We thank the other members of GOLD Lab for insightful feedback and discussions. Hyeonjoo Lee and Jin-Wu Nam (Hanyang University, Republic of Korea) kindly shared *Xist* knockdown RNA-sequencing data. All computation was carried out on the University of Bern Interfaculty Bioinformatics Unit cluster maintained by Rémy Bruggman and Pierre Berthier. We acknowledge administrative and logistical support from Basak Ginsbourger, Rahel Tschudi, Marla Rittiner, Beatrice Stalder, Willy Hofstetter and Patrick Furer (DBMR, University of Bern). HG-R is supported by a Marie Skłodowska-Curie Actions postdoctoral fellowship (R21950). Work in the GOLD Lab is funded by the Swiss National Science Foundation through the National Centre of Competence in Research (NCCR) "RNA & Disease" (51NF40-182880), Project funding "The elements of long noncoding RNA function" (31003A_182337), Sinergia project "Regenerative strategies for heart disease via targeting the long noncoding transcriptome" (173738), by the Medical Faculty of the University and University Hospital of Bern, by the Helmut Horten Stiftung, Swiss Cancer Research Foundation (4534-08-2018), and Science Foundation Ireland through Future Research Leaders award 18/FRL/6194.

References

1. Thorvaldsen, J. L., Duran, K. L. & Bartolomei, M. S. Deletion of the H19 differentially methylated domain results in loss of imprinted expression of H19 and Igf2. *Genes & Development* **12**, 3693 (1998).

2. Brown, C. J. *et al.* A gene from the region of the human X inactivation centre is expressed exclusively from the inactive X chromosome. *Nature* **349**, 38–44 (1991).

3. Ørom, U. A. *et al.* Long Noncoding RNAs with Enhancer-like Function in Human Cells. *Cell* **143**, 46–58 (2010).

4. Rom, A. *et al.* Regulation of CHD2 expression by the Chaserr long noncoding RNA gene is essential for viability. *Nature Communications* **10**, 1–15 (2019).

5. Leucci, E. *et al.* Melanoma addiction to the long non-coding RNA SAMMSON. *Nature* **531**, 518–522 (2016).

6. Gil, N. & Ulitsky, I. Regulation of gene expression by cis-acting long non-coding RNAs. *Nature Reviews Genetics* vol. 21 102–117 Preprint at https://doi.org/10.1038/s41576-019-0184-5 (2020).

7. Xiang, J. F. *et al.* Human colorectal cancer-specific CCAT1-L lncRNA regulates long-range chromatin interactions at the MYC locus. *Cell Research* **24**, 513–531 (2014).

8. Fanucchi, S. *et al.* Immune genes are primed for robust transcription by proximal long noncoding RNAs located in nuclear compartments. *Nature Genetics 2018 51:1* **51**, 138–150 (2018).

9. Gil, N. & Ulitsky, I. Production of spliced long noncoding RNAs specifies regions with increased enhancer activity. *Cell Syst* **7**, 537 (2018).

10. Engreitz, J. M. *et al.* Local regulation of gene expression by lncRNA promoters, transcription and splicing. *Nature* **539**, 452–455 (2016).

11. Ørom, U. A. & Shiekhattar, R. Long non-coding RNAs and enhancers. *Curr Opin Genet Dev* **21**, 194–198 (2011).

12. Tan, J. Y., Biasini, A., Young, R. S. & Marques, A. C. Splicing of enhancer-associated lincRNAs contributes to enhancer activity. *Life Sci Alliance* **3**, (2020).

13. Marques, A. C. *et al.* Chromatin signatures at transcriptional start sites separate two equally populated yet distinct classes of intergenic long noncoding RNAs. *Genome Biology* **14**, R131 (2013).

14. Hnisz, D., Shrinivas, K., Young, R. A., Chakraborty, A. K. & Sharp, P. A. A Phase Separation Model for Transcriptional Control. *Cell* **169**, 13–23 (2017).

15. Wu, M., Yang, L. Z. & Chen, L. L. Long noncoding RNA and protein abundance in lncRNPs. *RNA* **27**, 1427 (2021).

16. Palazzo, A. F. & Koonin, E. V. Functional Long Non-coding RNAs Evolve from Junk Transcripts. *Cell* **183**, 1151–1161 (2020).

17.   Ramilowski, J. A. *et al.* Functional Annotation of Human Long Non-Coding RNAs
via Molecular Phenotyping. *bioRxiv* 700864 (2019) doi:10.1101/700864.

18.   Ramilowski, J. A. *et al.* Functional annotation of human long noncoding RNAs via
molecular phenotyping. *Genome Research* **30**, 1060–1072 (2020).

19.   Tan, J. Y. *et al.* cis -Acting Complex-Trait-Associated lincRNA Expression
Correlates with Modulation of Chromosomal Architecture. *Cell Reports* **18**, 2280–2288
(2017).

20.   Marstrand, T. T. & Storey, J. D. Identifying and mapping cell-type-specific
chromatin programming of gene expression. *Proc Natl Acad Sci U S A* **111**, E645–E654
(2014).

21.   Wang, Y. *et al.* The 3D Genome Browser: A web-based browser for visualizing 3D
genome organization and long-range chromatin interactions. *Genome Biology* **19**, 1–12
(2018).

22.   Forne, T. *et al.* Loss of the maternal H19 gene induces changes in Igf2 methylation
in both cis and trans. *Proc Natl Acad Sci U S A* **94**, 10243 (1997).

23.   Carmona, S., Lin, B., Chou, T., Arroyo, K. & Sun, S. LncRNA Jpx induces Xist
expression in mice using both trans and cis mechanisms. *PLoS Genet* **14**, (2018).

24.   Luo, S. *et al.* Divergent lncRNAs Regulate Gene Expression and Lineage
Differentiation in Pluripotent Cells. *Cell Stem Cell* **18**, 637–652 (2016).

25.   Maass, P. G. *et al.* A misplaced lncRNA causes brachydactyly in humans. *The
Journal of Clinical Investigation* **122**, 3990 (2012).

26.   Engreitz, J. M. *et al.* The Xist lncRNA Exploits Three-Dimensional Genome
Architecture to Spread Across the X Chromosome. *Science (1979)* **341**, 1237973 (2013).

27.   Vance, K. W. *et al.* The long non-coding RNA Paupar regulates the expression of
both local and distal genes. *EMBO J* **33**, 296–311 (2014).

28.   Chalei, V. *et al.* The long non-coding RNA Dali is an epigenetic regulator of neural
differentiation. *Elife* **3**, 1–24 (2014).

29.   Stalder, L. *et al.* The rough endoplasmatic reticulum is a central nucleation site of
siRNA-mediated RNA silencing. *The EMBO Journal* **32**, 1115 (2013).

30.   Zeng, Y. & Cullen, B. R. RNA interference in human cells is restricted to the
cytoplasm. *RNA* **8**, 855–60 (2002).

31.   Biasini, A. *et al.* Translation is required for miRNA-dependent decay of
endogenous transcripts. *The EMBO Journal* **40**, (2021).

32.   Lai, F., Damle, S. S., Ling, K. K. & Rigo, F. Directed RNase H Cleavage of Nascent
Transcripts Causes Transcription Termination. *Molecular Cell* **77**, 1032-1043.e4 (2020).

33.   Durand, N. C. *et al.* Juicer Provides a One-Click System for Analyzing Loop-
Resolution Hi-C Experiments. *Cell Systems* **3**, 95–98 (2016).

34.   Dixit, A. *et al.* Perturb-Seq: Dissecting Molecular Circuits with Scalable Single-Cell
RNA Profiling of Pooled Genetic Screens. *Cell* **167**, 1853-1866.e17 (2016).

35.   Basu, S. & Larsson, E. A Catalogue of Putative cis-Regulatory Interactions
Between Long Non-coding RNAs and Proximal Coding Genes Based on Correlative
Analysis Across Diverse Human Tumors. *G3 Genes|Genomes|Genetics* **8**, 2019–2025
(2018).

36.   Agrawal, S. *et al.* Systematic identification of cis-interacting lncRNAs and their
targets. *bioRxiv* 2021.01.13.426305 (2022) doi:10.1101/2021.01.13.426305.

37.   Goyal, A. *et al.* A cautionary tale of sense-antisense gene pairs: independent
regulation despite inverse correlation of expression. *Nucleic Acids Res* **45**, 12496–12508
(2017).

38.   Scharner, J. *et al.* Hybridization-mediated off-target effects of splice-switching
antisense oligonucleotides. *Nucleic Acids Research* **48**, 802 (2020).

39.   Marasco, L. E. *et al.* Counteracting chromatin effects of a splicing-correcting
antisense oligonucleotide improves its therapeutic efficacy in spinal muscular atrophy. *Cell*
185, 2057-2070.e15 (2022).

40.   Doolittle, W. F. We simply cannot go on being so vague about "function." *Genome*
Biology **19**, 1–3 (2018).

41.   Graur, D. *et al.* On the Immortality of Television Sets: "Function" in the Human
Genome According to the Evolution-Free Gospel of ENCODE. *Genome Biology and*
Evolution **5**, 578–590 (2013).

42.   Frankish, A. *et al.* GENCODE 2021. *Nucleic Acids Res* **49**, D916–D923 (2021).

43.   Cheng, L. *et al.* LncRNA2Target v2.0: a comprehensive database for target genes
of lncRNAs in human and mouse. *Nucleic Acids Research* **47**, (2019).

44.   Hoon, D. S. B., Rahimzadeh, N. & Bustos, M. A. EpiMap: Fine-tuning integrative
epigenomics maps to understand complex human regulatory genomic circuitry. *Signal*
Transduction and Targeted Therapy 2021 6:1 **6**, 1–3 (2021).

45.   Zacher, B. *et al.* Accurate Promoter and Enhancer Identification in 127 ENCODE
and Roadmap Epigenomics Cell Types and Tissues by GenoSTAN. *PLOS ONE* **12**,
e0169249 (2017).

46.   Khan, A. & Zhang, X. dbSUPER: a database of super-enhancers in mouse and
human genome. *Nucleic Acids Research* **44**, D164–D171 (2016).

47.   Ramilowski, J. A. *et al.* Functional annotation of human long noncoding RNAs via
molecular phenotyping. *Genome Research* **30**, 1060–1072 (2020).

48.   Cheng, L. *et al.* LncRNA2Target v2.0: a comprehensive database for target genes
of lncRNAs in human and mouse. *Nucleic Acids Research* **47**, D140 (2019).