Gobies inhabiting natural CO₂ seeps reveal acclimation strategies to long-term acidification

Sneha Suresh[1], Alice Mirasole[2], Timothy Ravasi[3,4], Salvatrice Vizzini[5,6], Celia Schunter[1,7*]

[1]Swire Institute of Marine Science, School of Biological Sciences, The University of Hong Kong, Pokfulam Road, Hong Kong SAR

[2]Department of Integrative Marine Ecology, Ischia Marine Centre, Stazione Zoologica Anton Dohrn, via F. Buonocore 42, 80077 Ischia, Naples, Italy

[3]Marine Climate Change Unit, Okinawa Institute of Science and Technology Graduate University, Onna-son, Japan

[4]Australian Research Council Centre of Excellence for Coral Reef Studies, James Cook University, Townsville, Australia

[5]University of Palermo, Department of Earth and Marine Sciences, via Archirafi 22, 90123 Palermo, Italy

[6]CoNISMa, National Inter-University Consortium for Marine Science, Piazzale Flaminio 9, 00196 Roma, Italy

[7]State Key Laboratory of Marine Pollution, City University of Hong Kong, Hong Kong, Hong Kong SAR, China

*Corresponding author

Correspondence: celiaschunter@gmail.com

Abstract

Ocean acidification (OA) is known to affect the physiology, survival, behaviour, and fitness of various fish species with repercussions at the population, community, and ecosystem levels. Some fish species, however, seem to acclimate rapidly to OA conditions and even thrive in acidified environments. The molecular mechanisms that enable species to successfully inhabit high CO_2 environments has not been fully elucidated especially in wild fish populations. Here, we used the natural CO_2 seep in Vulcano Island, Italy to study the effects of OA on the brain transcriptome of the anemone goby, a species with high population density in the CO_2 seep and investigate their potential for acclimation. When compared to fish from environments with ambient CO_2, gobies living in acidified waters showed differences in expression of transcripts involved in ion transport and pH homeostasis, cellular stress, immune response, circadian rhythm, and metabolism. We also found evidence of potential adaptive mechanisms to restore the functioning of GABAergic pathways, whose activity can be affected by exposure to elevated CO_2 levels. Our findings indicate that gobies living in the CO_2 seep may be capable of mitigating CO_2 induced oxidative stress and maintaining physiological pH while meeting the consequent increased energetic costs. The conspicuous difference in expression of core circadian rhythm transcripts could provide an adaptive advantage by increasing flexibility of physiological processes under acidified conditions thereby facilitating acclimation. Our results show potential molecular processes of acclimation to elevated CO_2 in gobies enabling them to thrive in the acidified waters of Vulcano Island.

Keywords: Ocean acidification, transcriptomics, climate change, brain, anemone goby

Introduction

Global change driven by anthropogenic activities is having profound effects on marine ecosystems with rising carbon dioxide (CO_2) levels being one of the most crucial problems (Doney et al., 2009, 2012; Hofmann et al., 2010; Kroeker et al., 2010). Oceans are major carbon sinks and have absorbed nearly 30% of anthropogenic CO_2 emissions since the 1980s resulting in ocean acidification (OA; Bindoff et al., 2019). Exposure to CO_2 levels predicted to occur by the year 2100 ($\geq$700 µatms) is known to impact the physiology, growth rate, development, survival, behaviour, and reproduction in various species of fish (Domenici et al., 2012; Esbaugh, 2017; Ferrari et al., 2012; S. L. Hamilton et al., 2017; T. J. Hamilton et al., 2014; Rachael M. Heuer & Grosell, 2014; Munday, Pratchett, et al., 2013; Radford et al., 2021; Simpson et al., 2011). This can have far-reaching consequences on species interactions, population sustainability and ultimately impact ecosystem functioning (Fabry et al., 2008; Ferrari et al., 2011; Kroeker et al., 2010; Munday et al., 2010).

Most knowledge about these impacts of ocean acidification (OA) on teleosts, however, comes from short-term laboratory-based studies and mostly only account for short-term effects of OA. Accurate assessment of the future biological impacts of OA on species and populations requires consideration of their potential for acclimation and adaptive evolution (Sunday et al., 2014). The effects of elevated CO_2 are also known to be variable both within and across species. Inter-individual variability in CO_2 tolerance has been reported both in laboratory studies and in the wild (Cattano et al., 2018; Lehmann et al., 2022; Monroe et al., 2021; Munday et al., 2013; Paula et al., 2019; Welch & Munday, 2017). The underlying reason for the observed variable effect is not fully known but could be the result of specific genetic and epigenetic signatures or environmental factors or a combination of those. Environmental factors such as nutrient gradients, species composition, habitat modifications, and predation risk influence the effects of elevated CO_2 on various aspects of fish physiology and behaviour (Mirasole et al.,

2019; Munday et al., 2014; Nagelkerken et al., 2015; Zunino et al., 2019). Studies on wild fish

populations that naturally inhabit high CO_2 environments provide a unique opportunity for

integrating the direct and indirect effects of elevated CO_2 exposure on fish and investigate the

molecular mechanisms enabling acclimation.

Volcanic CO_2 seeps in the ocean, where emission of CO_2-rich fluids locally reduce pH,

have been used as natural analogues to study the effects of long-term exposure of fish to

elevated CO_2 in their natural environment. Shifts in habitat, trophic diversity and fish

assemblages at CO_2 seeps have been reported which could obscure some of the negative effects

of OA on fish (Fabricius et al., 2011; Hall-Spencer et al., 2008; Mirasole et al., 2019;

Nagelkerken et al., 2015; Vizzini et al., 2017). For example, predator recognition ability and

risk behaviour of species living in CO_2 seeps was found to be correlated with predator

abundance at the seeps, with perception of predator threat to be unaffected by elevated CO_2 in

high predator risk environments (Cattano et al., 2017) and low-risk predation environments

offsetting any negative impacts of CO_2 induced impairment in predator recognition (Munday

et al., 2014). Habitat shifts is another factor that can work antagonistically with CO_2 to

influence species behaviour. Gobies from the CO_2 seeps in Vulcano Island, Italy living in more

open habitats showed reduced predator escape behaviour, however, the same was not observed

in gobies living in more sheltered sites indicating the influence of habitat on species response

to OA (Nagelkerken et al., 2015). Similarly, habitat shifts resulting in net resource enrichment

at CO_2 seeps seems to benefit certain fish species by providing sufficient food to meet the

increased energy demands of living in a high CO_2 environment (Connell et al., 2013; Mirasole

et al., 2019; Nagelkerken et al., 2015). It is therefore important to consider the indirect effects

of OA while making inferences about the adaptive potential of species to future OA conditions

and volcanic CO_2 seeps provide the perfect opportunity to do so.

Prolonged exposure to elevated CO_2 at the seeps may enable fish to acclimate through

physiological and behavioural compensation resulting in long-term survival and subsequent

adaptation. Evidence from studies using transcriptomics to determine gene expression patterns

in the brain and gonads of fish from natural CO_2 seeps revealed plasticity and adaptive selection

to elevated CO_2 (Kang et al., 2022; Petit-Marty et al., 2021). These findings suggest that species

that are continuously exposed to elevated CO_2 levels can develop mechanisms to mitigate

negative effects of CO_2 exposure and have the potential to adapt to acidified environments.

While several studies have investigated the effects of elevated CO_2 on fish physiology,

behaviour, and molecular processes, we are only starting to learn about the effects of CO_2 on

brain function in wild fish species.

Here, we sequenced and analysed the brain transcriptome of the anemone goby, *Gobius*

incognitus to identify gene expression patterns underlying acclimation to elevated CO_2 in the

wild population. This benthic fish species has a very limited home range making it an ideal

model to study the effects of OA *in situ*. Previous studies have reported potentially negative

physiological effects such as changes in otolith morphology and skeletal composition in *Gobius*

from the natural CO_2 seeps at Vulcano Island (Mirasole et al., 2017, 2021), however, they still

maintain higher population densities at the seeps (Nagelkerken et al., 2015). This could mean

that gobies have developed alternative strategies/mechanisms to offset the negative effects of

elevated CO_2 exposure. In fact, swimming activity and predator perception of *Gobius*

incognitus living in natural CO_2 seeps in Vulcano Island was not affected by elevated CO_2

indicating higher tolerance of this species to changing ocean conditions (Spatafora et al., 2022).

By examining brain gene expression patterns of the anemone goby from the natural CO_2 seeps

in Vulcano Island, Italy we aim to identify molecular signatures of acclimation that enable this

species to mitigate negative effects of elevated CO_2 and thrive in the acidified waters of

Vulcano Island.

Methods

Study area and sample collection

Gobius incognitus samples were collected by SCUBA diving on 30th September 2018

at a depth of 1-2 meters with small hand nets from two locations in Levante Bay, Vulcano

Island (Italy): a low pH site (38.419 ºN, 14.961 ºE; pH 7.80 ± 0.09) and a control site located

around 600 m from the low pH site (38.420 ºN, 14.965 ºE; pH 8.19 ± 0.03) (Figure 1). The

control site was similar to the low pH site in terms of orientation (south-east), depth and habitat

(sandy with beds of *Cymodocea nodosa* seagrass). A total of eight individuals were sampled at

each site at the same time of day (between 11:00 am and 12:00 pm). The total length of all fish

was measured in the field, immediately after which they were euthanized on ice and the brain

tissues were dissected, preserved in RNAlater, and stored at -80ºC until further processing. The

age of all sixteen individuals was estimated using the Von Bertalanffy function. On average all

the sampled fish were four cm long and two years old (Supplementary Table S1).

Figure 1: (a) Map showing the sampling sites in Levante Bay, Vulcano Island. The star

indicates the location of the primary vent and the circles indicate the low pH site and the control

(ctrl) site. (b) Picture of the study species *Gobius incognitus* in the wild.

RNA extraction and sequencing

Frozen whole brains were homogenized in RLT-lysis buffer (Qiagen) for 30s at 30Hz/s using Qiagen TissueLyzer II with 4-5 single-use silicon beads per sample and total RNA was extracted using the Qiagen RNeasy Micro Kit. On-column DNase digestion was performed using RNase-Free DNase1 (Qiagen) following the manufacturer's instructions, to remove any DNA contamination. RNA concentration and quality was determined using Qubit and Agilent 2100 Bioanalyzer respectively (mean RIN = 7.79) following which RNA-Seq libraries were built using the KAPA Stranded mRNA-Seq Kit with 100 ng total RNA as input from each of the 16 individuals. The samples were subsequently sequenced on an Illumina NovaSeq 6000 to get 151 bp, paired-end reads at the Centre for PanorOmics Sciences, The University of Hong Kong. An average of 33 ± 2.8 million read pairs were obtained from the 16 RNA libraries sequenced (Supplementary Table S2).

Sequence processing and primary *de novo* transcriptome assembly

The quality of all the sequencing reads was determined using FastQC v0.11.5 (Andrews, 2010). To ensure high quality of all reads, we trimmed adapters and sequences with an average Phred-scaled quality score below 30 using fastp v0.20.0 (Chen et al., 2018) with a 4-base sliding window approach. Sequences less than 40 bp in length after trimming were removed. The trimmed sequences were further filtered to remove potential contaminants, ambiguous bases, and rRNA sequences. Potential contaminant sequences were removed using kraken v2.0.8-beta (Wood & Salzberg, 2014) with a confidence score of 0.3 using the bacterial, fungal, and viral databases from NCBI RefSeq as reference. For sequences containing ambiguous base calls (Ns), the longest sub-sequence of at least 40 bp without any ambiguous bases was extracted and retained if it was at least half the original sequence length and had an average Phred-scaled quality score above 30 using the 'filter_illumina' script from DRAP (Cabau et al.,

2017). Reads originating from ribosomal RNA (rRNA) were identified and removed by

mapping the sequences to the SSUParc (SILVA_138_SSUParc_tax_silva.full_metadata.gz)

and LSUParc datasets (SILVA_132_LSUParc.full_metadata.gz) from SILVA (Quast et al.,

2013) using Bowtie2 v2.4.1 (Langmead & Salzberg, 2012) in the very sensitive, local mode.

An average of 26 ± 2.5 million adapter free, filtered, quality-trimmed, and decontaminated

paired reads were finally retained after the stringent filtering process from all 16 samples

(Supplementary Table S2). These high quality reads were subsequently merged and assembled

de novo using Trinity v2.8.4 (Grabherr et al., 2013) within DRAP v1.92 (parameters: -c '-x'; --

no-trim; -m bwa; --no-rate) using the pooled-assembly strategy resulting in a primary

transcriptome assembly consisting of 633,517 contigs.

De novo transcriptome assembly optimization

Candidate coding regions (open reading frames (ORFs)) within the transcript contigs

assembled by DRAP were identified using TransDecoder v5.5.0 (Haas et al., 2013) and only

contigs with ORFs of at least 100 amino acids long were retained to prevent retention of false

positive ORFs as done in previous studies on CO_2 seeps (Kang et al., 2022; Petit-Marty et al.,

2021). These contigs were then blasted against the UniProtKB/Swiss-Prot Release 2021_01

database (Apweiler, 2004) using the BLASTX algorithm (Blast+ v2.6.0) (Altschul et al., 1990)

with an e-value cut-off 1e-05 and retaining only one blast hit per contig (--max_target_seqs =

1). For each assembled contig only the best ORF was retained prioritized first based on blast

homology and then based on ORF length (--retain_blastp_hits; --single_best_only) after which

113,895 contigs were retained. The quality of the selected contigs was further assessed using

the runAssesment module within DRAP v1.92 and 68 putative chimeric contigs were removed.

The high-quality RNA-seq reads were then re-mapped to the filtered transcriptome assembly

using Bowtie2 v2.4.1 in the sensitive mode and disallowing any discordant and unpaired

alignments, but without any upper limit on the number of alignments. To identify and cluster

individually assembled contigs belonging to the same gene, the alignment results were used by

Corset v1.09 (Davidson & Oshlack, 2014) to hierarchically cluster contigs based on the

proportion of multi-mapped reads and the expression patterns. For each of the Corset generated

clusters the longest transcript sequence was extracted using the fetchClusterSeqs.py script from

Corset-tools (https://github.com/Adamtaranto/Corset-tools) and those clusters with no aligned

reads were removed from the *de novo* transcriptome assembly resulting in 44,614 high quality

contigs retained.

To further validate the transcriptome assembly, blast search (TBLASTX, e-value = 1e-

05, max_target_seqs = 1) was performed against the complete cDNA sequence of *Neogobius*

melanostomus (round goby) obtained from Ensembl (release-104) (Howe et al., 2021), the

phylogenetically closest species with a sequenced genome to *Gobius incognitus* (Thacker &

Roje, 2011). To further reduce transcript redundancy and identify potential alternative

transcripts that failed to cluster in the previous steps, the blast results were filtered to retain

only the longest transcript for each blast hit to get the final *de novo* transcriptome assembly.

Transcripts with no blast hits to *Neogobius melanostomus* were also retained in the final

assembly as these could potentially be transcripts that are specific to our target species, *Gobius*

incognitus. The final transcriptome assembly consisted of 43,349 non-redundant transcripts.

The basic statistics of the transcriptome assembly determined using rnaQUAST v2.1.0

(Bushmanova et al., 2016), 'stats.sh' script from BBMap v38.87 (Bushnell, 2014) and

'contig_ExN50_statistic.pl'   from   Trinity   accessory   scripts

(https://github.com/trinityrnaseq/trinityrnaseq/tree/master/util/misc),   are   shown   in

Supplementary Table S3 and Supplementary Figure S1.

Transcriptome annotation and evaluation of assembly completeness

The *de novo* assembled transcriptome was compared against the UniProtKB/Swiss-Prot

Release 2021_01 database using BLASTX (e-value = 1e-05 and HSP length cut-off = 33)

resulting in 43.6% (24,454) of the assembled transcripts with blast hits to known proteins. Gene

Ontology (GO) mapping and annotation was then carried out within OmicsBox v1.4.11 (Gö

Tz et al., 2008). InterProScan was used to identify the protein domains within the transcript

sequences and the InterProScan GOs were merged with the GO annotations resulting in a total

of 24,875 (42.6%) annotated transcripts. For contigs that did not blast to the Swiss-Prot

database, the Ensembl gene ID obtained by blasting against *Neogobius melanostomus* was

retained. Combining the annotation results from OmicsBox and the additional 9,362 transcripts

with blast hits to the cDNA sequence of *Neogobius melanostomus*, 78.9% (34,237) of the

transcripts in the final *de novo* assembly has a functional annotation (Supplementary Table S3

and S4).

To incorporate the transcript expression levels in evaluating the contiguity of the *de*

novo assembly, the ExN50 (expression-informed N50) metric was determined. ExN50

(expression-informed N50) represents the N50 value for the most highly expressed transcripts

representing x% of the total normalized expression data and is more appropriate for

transcriptome assemblies than the conventional N50 statistic (Galachyants et al., 2019). A peak

of the ExN50 value at higher x percentiles indicates the effectiveness of the de novo assembly

pipeline to assemble longer transcripts (Sahraeian et al., 2017). The expression levels of the

transcripts were quantified using RSEM v1.3.1 (Li & Dewey, 2014) and the ExN50 value was

calculated across different expression percentiles using the TMM normalized expression values.

The *Gobius incognitus reference* transcriptome assembled here has a sufficiently high coverage

of the longer transcripts indicated by a peak of ExN50 value at E86 (Supplementary Table S5,

Supplementary Figure S2). The level of completeness of the transcriptome assembly was

determined using BUSCO v4.1.3 (Manni et al., 2021) using the Actinopterygii_odb10 and

Eukaryota_odb10 sets of Benchmarking Universal Single-Copy Orthologs (BUSCO) which

revealed high representation of both the Actinopterygian core genes and the Eukaryota core

genes. Specifically, 76.4% and 91.4% of Actinopterygii core genes and Eukaryota core genes

respectively, were recovered completely. An additional 7.7% of Actinopterygii core genes and

7.1% of Eukaryota core genes were recovered partially (Supplementary Figure S3) indicating

sufficiently good assembly and annotation completeness, similar to other *de novo* assemblies

(Pomianowski et al., 2021; Wong et al., 2019).

Differential expression analysis

The high-quality, filtered RNA-seq reads were re-mapped to the final transcriptome

assembly using Bowtie2 v2.4.1 in the sensitive mode, disallowing any discordant and unpaired

alignments and using additional parameters (--dpad 0 --gbar 99999999 --mp 1,1 --np 1 --score-

min L,0,-0.1) to ensure the resulting alignment files are compatible with RSEM, which was

used to quantify the transcript expression levels. On average 72% of the RNA-Seq reads

mapped back to the assembled transcriptome (Supplementary Table S6) indicating sufficiently

high quality of the *de novo* transcriptome. Principal component analysis (PCA) was performed

to determine the overall transcript expression patterns and identify any outlier samples in R

v4.1.1 (R Core Team, 2021) following which differential expression analysis was done using

DESeq2 v1.32.0 (Love et al., 2014). The transcripts were considered to be significantly

differentially expressed between the control and CO_2 seep sites if the FDR corrected p-value

(padj) was less than 0.05. Additionally, to remove further false positives, we set other filters to

absolute log2 fold change (abs log2FC) greater than 0.3 and baseMean greater than 10 as done

in previous studies on brain fish transcriptomics (Schunter et al., 2016, 2018). Enrichment of

GO functional categories among the significant DE genes were determined using Fisher's

Exact Test in OmicsBox v1.4.11 (*OmicsBox – Bioinformatics Made Easy, BioBam*

Bioinformatics, March 3, 2019) with an FDR threshold of 0.05, using the option "reduce to

most specific" to reduce the redundancy in the enriched GO categories identified. The heatmap

showing relative transcript expression patterns across samples was created using the pheatmap

package v1.0.12 (where transcripts with similar expression levels were clustered together) and

all other figures were created using the ggplot2 package v3.3.6 in R v4.2.1.

SNP calling and detection of outlier loci

SNP calling and outlier detection was performed to determine if there were any genetic

differences between the samples from the CO_2 seep and control sites. The previously obtained

BAM files were marked for duplicate reads using MarkDuplicates (Picard) and reads

containing Ns in their cigar string were split using SplitNCigarReads within the Genome

Analysis Toolkit (GATK v4.3.0.0) (Depristo et al., 2011). SNPs within the samples were called

using HaplotypeCaller within GATK and filtered using the VariantFiltration tool (parameters:

-window 35 -cluster 3 --filter-name FS -filter "FS>30.0" --filter-name QD -filter "QD<2.0").

The resulting VCF files from all 16 individuals were merged and further filtered to remove all

indels and retain only SNPs with a Phred- scaled quality score of 30 and minor allele frequency

(MAF) ≥ 0.2. Additionally, the minimum and maximum depth cut-off was set to 10 and 70

respectively per site and per genotype to account for possible mapping/assembly errors and all

sites where over 25% of the individuals are missing a genotype were removed. Discriminant

Analysis of Principal Components (DAPC; Jombart et al., 2010) was used to determine genetic

structuring between the samples and to evaluate if particular variants were potentially under

selection, outlier loci detection was performed in BayeScan v2.0 software (Foll & Gaggiotti,

2008) with the assumption that the individuals from the control and CO_2 seep sites belong to

two different populations.

Results

Brain transcriptome analysis

A total of 43,301 transcripts were expressed across all 16 samples with fish from the

CO_2 seep site showing more variability in transcript expression among individuals compared

to fish from the control site (Figure 2). There was no significant difference in the total body

length (Welch's t-test, $t(19.48) = 0.43$, $p = 0.67$) and age (Welch's t-test, $t(19.64) = 0.47$, $p =$

0.64) of the fish sampled from the low pH site and control site. Therefore none of the measured

biological factors drive the observed transcript expression patterns. When comparing transcript

expression levels between the control and CO_2 seep sites, 996 transcripts were identified to be

significantly differentially expressed (DE) (padj < 0.05, abs log2FC > 0.3, baseMean > 10;

Supplementary Table S7). Majority of these (68%) were expressed at higher levels in fish from

the CO_2 seep site compared to the control site (Supplementary Figure S4).

Figure 2: Overall transcript expression pattern across the 16 samples from the natural CO_2

seep and control site at Vulcano Island. Principal component analysis (PCA) was performed

using the regularized log transformed (rlog) counts of all expressed transcripts. The circles

represent the individuals from the control site and the triangles represent the individuals from

the low pH site. The blue and grey region is the 95% confidence interval of the samples from

the CO_2 seep and control sites respectively.

Functional analysis of DE transcripts

Functional enrichment analysis identified 434 GO terms to be enriched among the DE

transcripts (Supplementary Table S8). The transcripts associated with each enriched GO term

was further categorised into six broader functional groups (acid-base and ion homeostasis,

neural activity and related effects, circadian rhythm, metabolism, cellular stress response, and

immune response) based on their functional description from the NCBI gene database

(https://www.ncbi.nlm.nih.gov/gene/)   and   UniProt   knowledgebase   (UniProtKB;

https://www.uniprot.org/) (Figure 3, Supplementary Table S9).

Figure 3: Functional groups associated with the differentially expressed transcripts. Each

circle represents one transcript, and the colour represents the FDR corrected p-value (padj)

from the differential expression analysis.

The five most significant DE transcripts (with the lowest padj values) were all

associated with the circadian rhythm. These include two core circadian genes Period circadian

protein homolog 3 (*PER3*) and Orphan nuclear receptor RVR (*NR1D2*), two nuclear hormone

receptor genes involved in transcriptional regulation of core circadian genes Nuclear receptor

ROR-alpha A (*RORA*) and Nuclear receptor ROR-beta (*RORB*), and a transcriptional regulator

known to regulate circadian rhythm genes Nuclear factor interleukin-3-regulated protein

(*NFIL3*) (Supplementary Figure S4). In fact, all core clock transcripts were DE between the

control and CO_2 seep site; specifically, there was an overall downregulation of repressors of

the circadian rhythm and upregulation of the activators (Figure 4). Furthermore, *NOCT* a

circadian deadenylase that functions as a post transcriptional modulator in circadian control of

metabolic genes was upregulated in individuals from the CO_2 seep (Supplementary Table S9).

Figure 4: Relative expression levels of transcripts with enriched functions of fish from control and CO_2 seep sites. Each row represents a transcript, and the row z-score is calculated from the log2 transformed transcripts per million (TPM) values for each transcript. The transcripts are clustered based on expression levels.

Various transcripts involved in cellular stress response was also DE in our study. In fact

this group had the highest number of DE transcripts majority of which were upregulated in

individuals from the CO_2 seep (Figure 4). This included transcripts known to play a role in the

production of reactive oxygen species (ROS), transcripts involved in phosphatidylinositol 3-

kinase (PI3K) activity and signalling, DNA damage repair, unfolded protein response and

ubiquitination, molecular chaperones including heat shock proteins (HSPs), cell cycle control,

apoptosis, various members of the Ras superfamily of small GTPases and transcripts associated

with redox-sensitive signalling pathways such as the NF-kB and MAPK signalling. Notably,

there was also an overall increase in the expression of transcripts associated with cellular

detoxification and redox regulation (such as glutathione transferase, glutathione peroxidase,

heme oxygenase, isocitrate dehydrogenase, and succinate dehydrogenase), cellular response to

hypoxia, immune response and immune system function in individuals from the CO_2 seep site.

Of particular interest was the *galactose-specific lectin nattectin*, a C-type lectin which is

involved in innate immune responses as it had the highest fold-change in expression among all

DE transcripts (Supplementary Table S9).

Ion channels and transporters were another major functional group that was

differentially regulated between fish from the CO_2 seep and control sites. Notably, transcripts

encoding the Na^+/K^+ transporting ATPase (*ATP1A1*) and carbonic anhydrase 10 (*CA10*), which

play critical roles in ion-regulation and maintaining pH homeostasis, were upregulated in fish

from the CO_2 seep site (Figure 3). Additionally, various transcripts involved in actin

cytoskeleton organisation were also DE (Supplementary Table S9, Figure 4). This could

indirectly affect membrane transport as the cytoskeleton closely interacts with the plasma

membrane. Changes in ionic composition across neuronal membranes can affect the activity of

neurotransmitters such as GABA and glutamate and we found transcripts involved in

degradation of GABA (*SLC6A13* and *GABAT*) and a transcript encoding the metabotropic

glutamate receptor 4 (*GRM4*) to be differentially expressed in our study (Supplementary Table

S9, Figure 4). Additionally, transcripts encoding chloride transporters and claudins, whose

expression is affected by neural activity, were also DE (Supplementary Table S9, Figure 4).

Several transcripts involved in metabolism were also found to be differentially

regulated between fish from control and CO_2 seep sites. Transcripts involved in key metabolic

processes that result in increased ATP production such as glycolysis, gluconeogenesis, TCA

cycle, mitochondrial beta-oxidation, mitochondrial respiratory chain, lipid and amino acid

metabolism were expressed at higher levels in gobies from the CO_2 seep site. Interestingly,

transcripts involved in fatty acid synthesis, an ATP consuming process were downregulated in

individuals from the CO_2 seep (Supplementary Table S9, Figure 4).

Outlier loci detection

We evaluated the genetic background of the fish living in CO_2 seep and control site. A

total of 8,629 high-quality SNPs were obtained across all 16 individuals from the CO_2 seep and

control site. There was no genetic structuring found among the analyzed samples and no outlier

was detected (Supplementary Figure S5). Therefore there was no evidence of genetic

divergence between the individuals from the control and CO_2 seep site. Gobies have a pelagic

larval phase and hence the individuals from the CO_2 seep and control sites come from a

heterogeneous pool of larvae and there is no evidence of post-settlement processes between the

sites.

Discussion

Natural CO_2 seeps provide an excellent opportunity to study the effects of long-term

exposure of fish to elevated CO_2 in their natural environment and determine their adaptive

potential while also accounting for various indirect effects of ocean acidification (OA) such as

habitat shifts and changes in species assemblages. Here, we investigated changes in the brain

transcriptome of the anemone goby from a natural CO_2 seep in Vulcano Island to identify

molecular processes enabling acclimation to elevated CO_2 in the wild. Previous studies at

Vulcano Island have found limited behavioural effects of CO_2 on this fish species and increased

population size at the seep suggesting that this species has higher tolerance to OA conditions

possibly through differential regulation of gene expression enabling physiological and

behavioural adjustments (Nagelkerken et al., 2015; Spatafora et al., 2022).

We found large differences in overall brain transcript expression levels between

individuals from CO_2 seep and ambient control sites. The expression patterns were more

variable among individuals from the CO_2 seep compared to those from the control site which

is not due to the age or length of the fish. Several previous studies have reported variability in

species response to elevated CO_2 (Cattano et al., 2018; Monroe et al., 2021; Munday et al.,

2013; Paula et al., 2019; Welch & Munday, 2017), which could be due to variation in

acclimatization strategies or genetic and epigenetic differences. Further research is needed to

fully understand the underlying causes of individual variation in CO_2 tolerance especially since

intraspecific variability can be heritable (Welch & Munday, 2017) and hence is a key factor in

influencing adaptive capacity of populations to future global change (Lehmann et al., 2022;

Munday et al., 2013).

The majority of differentially expressed (DE) transcripts exhibited elevated expression

in fish from the CO_2 seep similar to what has been observed in gonads of the common triplefin

from the CO_2 seep in New Zealand, another species known to have increased population size

in acidified environments (Petit-Marty et al., 2021). The gobies seem to be able to elicit

acclimation responses to elevated CO_2 by differentially regulating transcripts associated with

crucial functions which potentially enables them to maintain cellular homeostasis and thrive in

the acidified waters of Vulcano Island. Specifically, 2.3% of the brain transcriptome involved

in key functions such as acid-base and ion homeostasis, neurological function, circadian

rhythm, metabolism, cellular stress response, and immune response were differentially

expressed (DE) between fish from the CO_2 seep and control sites.

Cellular Stress Response was the most gene rich response in gobies at the CO_2 seep.

We observed DE of various transcripts associated with redox sensitive pathways such as NF-

*k*B and MAPK signalling as well as transcripts known to induce an increase in the intracellular

concentration of reactive oxygen species (ROS). Therefore, gobies inhabiting the CO_2 seep

could experience changes in cellular redox balance resulting in oxidative stress (Poli et al.,

2012; Zhang et al., 2016). They however also seem to be capable of counteracting stress-

induced damage by upregulating transcripts involved in cellular detoxification and redox

regulation. Oxidative stress in fish can trigger cell damage and apoptosis (Pizzino et al., 2017;

Strader et al., 2020) and also cause conformational changes and oxidative damage to all

macromolecules including DNA, proteins, and lipids thereby affecting transmission of genetic

information and resulting in intracellular accumulation of unfolded and misfolded proteins

(Juan et al., 2021) which is cytotoxic (Hetz, 2012). Differential regulation of transcripts

involved in DNA damage response and DNA repair, unfolded protein response (UPR),

chaperones such as the heat shock protein (HSP) family observed here suggest that gobies are

capable of minimising oxidative damage on macromolecules caused by ROS. High constitutive

expression of several universally conserved genes in the cellular stress response machinery in

fish from the CO_2 seep indicate an adaptive response to increase cellular resistance to oxidative

stress induced cell damage.

Transcripts involved in innate and adaptive immune response and the complement

system were found to be up-regulated in gobies from the CO_2 seep site. Elevated levels of

cellular stress can result in increased infections from opportunistic pathogens and hence

activation of immune responses could be a protective mechanism to prevent potential infections

in the face of increased physiological stress under conditions of OA (de Souza et al., 2016).

Immune response genes are commonly differentially regulated in fish exposed to elevated CO_2

(de Souza et al., 2016; De Souza et al., 2014; Machado et al., 2020) revelaing it as an important

function that is regulated under OA conditions. Additionally, two hypoxia-inducible transcripts,

HIF1A and *EGLN2* were upregulated in individuals from the CO_2 seep. While the primary

function of these transcripts is to regulate cellular response to hypoxic conditions, their

expression could also be enhanced in response to cellular stress. Increased mRNA expression

of *HIF1A* in response to hypercapnia has been previously reported under stable oxygen levels

(Machado et al., 2020). These results indicate that gobies living in the naturally acidified waters

of Vulcano Island are able to elicit physiological compensatory mechanisms to counter the

cellular stress induced by prolonged exposure to elevated CO_2 levels.

Living in CO_2 seeps also induced expression differences in various transcripts

associated with ion transport and conserving pH balance. In particular, carbonic anhydrase

(CA), an enzyme which plays a key role in CO_2 excretion, acid-base balance, and ion-

regulation was upregulated in gobies from the CO_2 seep, which could facilitate increased CO_2

excretion thereby preventing acidosis and also enhance transport of O_2 to the brain (Rummer

& Brauner, 2011). Additionally, transcripts encoding potassium channel transporters (*KCNAB3*,

KCNK10 and *KCNJ10*), and a Na^+/K^+ transporting ATPase (*ATP1A1*) were differentially

expressed (DE) in the gobies. Acid-base balance and ion homeostasis in fish are closely linked

processes as transport of H^+ and HCO_3^- ions are coupled to K^+, Na^+ and Cl^- transport (Gilmour

& Perry, 2009). Regulation of ion transport is therefore key to ensure proper cellular

functioning, maintain pH homeostasis and prevent respiratory acidosis which is a common

effect of elevated CO_2 exposure in fish (Heuer & Grosell, 2014; Ishimatsu et al., 2005). DE of

transcripts encoding potassium channel proteins in our study could potentially be involved in

enabling gobies living in the CO_2 seep to sense ambient CO_2 levels and in turn trigger

downstream homeostatic responses to restore internal acid-base status as potassium channels

are involved in CO_2 chemoreception and sensing acid-base disturbance (Qin et al., 2010).

Additionally, the increased expression of *ATP1A1* in fish from the CO_2 seep may play an

important role in pH homeostasis as the sodium and chloride gradient created by this enzyme

affects the activity of other key secondary transporters involved in pH regulation (Casey et al.,

2010; Singh Jaggi et al., 2015; Tresguerres et al., 2020). Transcripts involved in actin

cytoskeleton organisation was also DE which combined with DE of Ras GTPases, which are

involved in remodelling the cytoskeleton, could imply changes in membrane-cytoskeleton

interactions in individuals from the CO_2 seep, in turn affecting membrane transport (de Curtis

& Meldolesi, 2012). Therefore, gobies inhabiting the naturally acidified waters of Vulcano

Island can efficiently regulate various ion transport mechanisms to buffer any intracellular pH

changes caused by elevated environmental CO_2 and thereby maintain cellular homeostasis.

We found subtle evidence of regulation of GABAergic signalling in gobies from the

CO_2 seep which could be a consequence of changes in ion channel activity as neural signalling

pathways are reliant on ion transport and stable ion gradients. Two transcripts, *SLC6A13* and

GABAT regulating the turnover rate of extracellular γ-aminobutyric acid (GABA), which is a

major neurotransmitter in the brain, were upregulated in gobies from the seep. Given that

GABA is not prone to enzymatic degradation, increased expression of these transcripts could

be an adaptive response to regulate GABAergic signalling which is known to be altered in high

CO$_2$ environments as a result of pH buffering processes (Heuer et al., 2016; Hübner & Holthoff,

2013; Kang et al., 2022; Nilsson et al., 2012; Schunter et al., 2019). Furthermore, upregulation

of the glutamate metabotropic receptor 4 (*GRM4*) in fish from the CO$_2$ seep could be a

compensatory response to reduce presynaptic GABA release (Bobeck et al., 2014; Flor et al.,

1995). Additionally, transcripts involved in the transport of chloride ions, whose concentration

gradient influences GABA activity, were DE, which could potentially regulate GABAergic

neurotransmission (Jentsch et al., 2002). Particularly, the downregulation of *SLC12A1* in the

gobies could be an adaptive response to restore any changes in GABA function (Grosell, 2011;

Hübner & Holthoff, 2013) similar to what has been observed with the potassium-chloride co-

transporter 2 (KCC2) gene in the damselfish *Acanthochromis polyacanthus* exposed to

elevated CO$_2$ (Schunter et al., 2018). Therefore, gobies inhabiting the naturally acidified waters

of Vulcano Island possibly use alternative strategies to ensure proper functioning of neural

signalling pathways reliant on maintenance of stable ionic gradients which are altered by pH

buffering processes.

The circadian rhythm is a key pathway regulating nearly all physiological processes

and all core transcripts in this pathway showed differential expression. CO$_2$ induced changes

in the circadian rhythm have been repeatedly observed across various species and are possibly

linked to changes in GABA receptor activity (Lee et al., 2021). The circadian pathway enables

organisms to anticipate and respond to environmental changes which is evolutionarily

advantageous (Ayyar & Sukumaran, 2021; Dmitriev & Mangel, 2000; Prokkola & Nikinmaa,

2018). Changes in the expression of circadian genes in response to elevated CO$_2$, have been

suggested to result in an amplitude or phase shift of the circadian clock providing an adaptive

advantage by increasing flexibility of physiological processes (Kang et al., 2022; Lee et al.,

2021; Schunter et al., 2016, 2021). We found evidence of potential circadian regulation of

metabolism in gobies in response to elevated environmental CO$_2$. In addition, *NOCT,* an

immediate early gene with deadenylase activity was upregulated in gobies from CO_2 seep.

NOCT functions as a mediator of post-transcriptional circadian regulation of metabolic

processes and can also respond to changes in the external environment (Stubblefield et al.,

2012). Increased expression of *NOCT* in individuals from the CO_2 seep combined with DE of

all core circadian genes suggest rhythmic regulation of energy metabolism potentially enabling

gobies to meet the increased energetic demands of living in an acidified environment. Given

the ubiquity of the circadian clock, there is a need for further research to understand the

biological implications of the effect of elevated CO_2 on the circadian rhythm and its impacts

on neuronal signal transduction and other downstream physiological processes.

In addition to the above-mentioned rhythmic regulation of metabolism, we found an

overall increase in expression of transcripts involved in ATP production in gobies from the

CO_2 seep. Interestingly, transcripts involved in fatty-acid biosynthesis, an anabolic process was

downregulated. The constant need to maintain acid-base balance and homeostasis while

minimising oxidative stress in a high CO_2 environment can be energetically expensive (Cattano

et al., 2018; Kelly & Hofmann, 2013; Strader et al., 2020). Decreased expression of transcripts

involved in fatty acid synthesis could be an alternative strategy of gobies living in the seep to

facilitate allocation of acetyl-CoA, a key metabolite that intersects with glycolysis, TCA cycle,

fatty acid beta-oxidation and fatty acid synthesis (Currais et al., 2019), for use in energy

producing metabolic pathways. This in addition to up-regulation of *SLC16A1* and *SLC25A22*,

which are also involved in transport of key metabolic intermediates, suggests that gobies

selectively regulate metabolic pathways to ensure increased energy production. Additionally,

all transcripts encoding hexose/glucose transporters, with the exception of *SLC45A1* were also

upregulated in gobies from the CO_2 seep which could facilitate increased availability of glucose

in the brain. This is particularly important as glucose is the primary substrate for energy

production in the brain. Despite being a glucose transporter, *SLC45A1* was selectively

downregulated probably because it is a sugar-H^+ symporter (Shimokawa et al., 2002) and hence

facilitates H^+ import into neuronal cells thereby increasing intracellular pH which could be

detrimental. Therefore, gobies have potentially acclimated to a high CO_2 environment by

selective regulation of genes involved in metabolism to increased energy production. Changes

in the energy budgets of fish living in acidified waters could result in physiological trade-offs

(Kelly & Hofmann, 2013). However, the increased metabolic capacity of gobies observed here

combined with CO_2 driven habitat shifts resulting in enrichment of key trophic resources at the

CO_2 seep (Nagelkerken et al., 2015) potentially enables them to meet increased energy

requirements. This aids in maintaining pH homeostasis and mitigating any potential cellular

damage due to elevated CO_2 levels without a significant trade-off in the energy available for

other biological processes. In fact, the CO_2 vents have a higher abundance of gobies compared

to nearby control sites (Mirasole et al., 2019; Nagelkerken et al., 2015; Spatafora et al., 2022).

At other CO_2 seeps at White Island, New Zealand, a similar habitat shift with net resource

enrichment can be found and the common triplefin *F. lapillum* also exhibit a higher abundance

with a similar increase in metabolic capacity (Nagelkerken et al., 2015; Petit-Marty et al., 2021).

Therefore, gobies similar to the common triplefin, might benefit from the habitat shifts at CO_2

seeps and show an adaptive response to prolonged CO_2 exposure with an overall increase in

metabolism.

Our findings show molecular signatures of acclimation in the anemone goby to the

acidified waters of Vulcano Island. The gobies effectively regulate disturbances in acid-base

balance, maintain cellular homeostasis and mitigate OA induced oxidative stress. Gobies also

showed potential adaptive mechanisms to maintain proper functioning neurological signalling

pathways, which could be disrupted as a consequence of altered ionic composition due to pH

buffering processes. Although these processes can be energetically expensive potentially

resulting in trade-offs, the overall increase in metabolism seen in gobies from the CO_2 seep

suggests that they meet the increased energy demands associated with living in an acidified

environment. This could be further facilitated by net trophic resource enrichment at the CO_2

seep. The increased population density of gobies at the CO_2 seep in Vulcano Island reported in

previous studies combined with evidence of acclimation to OA found in our study indicates

that gobies can effectively mitigate the behavioural effects of elevated CO_2. If gobies undergo

adaptive selection like what has been seen in other species inhabiting CO_2 seeps, they might

be able to endure future ocean acidification conditions.

Author contributions

AM conceived the project with input from CS and TR. AM conducted the fieldwork and dissected the fish. SV provided the funding for fieldwork. SS and CS carried out the RNA extractions, designed the RNA-Seq analysis pipeline and drafted the manuscript. All authors contributed to data interpretation and editing the manuscript. All authors read and approved the manuscript.

Acknowledgments

We thank Dr. Juan Diego Gaitan-Espitia and Dr. Kang Jingliang for providing guidance and feedback on the *de novo* transcriptome assembly.

Ethics

All sample collection were carried out in accordance with the institutional and national law guidelines concerning the use of animals in research.

Competing financial interests

All authors declare they have no competing interests.

Funding

This project was funded by the start-up funds from The University of Hong Kong to CS.

Data accessibility

RNA-seq raw sequences and the *de novo* assembled transcriptome assemblies have been deposited in NCBI under BioProject PRJNA869880.

References

Altschul, S. F., Gish, W., Miller, W., Myers, E. W., & Lipman, D. J. (1990). Basic local

alignment search tool. *Journal of Molecular Biology*, *215*(3), 403–410.

https://doi.org/10.1016/S0022-2836(05)80360-2

Andrews, S. (2010). FastQC: a quality control tool for high throughput sequence data.

Apweiler, R. (2004). UniProt: the Universal Protein knowledgebase. *Nucleic Acids Research*,

32(90001), 115D – 119. https://doi.org/10.1093/nar/gkh131

Ayyar, V. S., & Sukumaran, S. (2021). Circadian rhythms: influence on physiology,

pharmacology, and therapeutic interventions. *Journal of Pharmacokinetics and*

Pharmacodynamics, *48*(3), 321–338. https://doi.org/10.1007/S10928-021-09751-2

Bindoff, N. L., Cheung, W. W. L., Kairo, J. G., Arístegui, J., Guinder, V. A., Hallberg, R.,

Hilmi, N., Jiao, N., Karim, M. saiful, Levin, L., Sean, O., Purca, S. R., Baruch, R., Suga,

T., Tagliabue, A., & Williamson, P. (2019). Chapter 5: Changing Ocean, Marine

Ecosystems, and Dependent Communities. *Intergovernmental Panel of Climate Change.*

IPCC Special Report on the Ocean and Cryosphere in a Changing Climate.

https://www.ipcc.ch/srocc/chapter/chapter-5/

Bobeck, E. N., Chen, Q., Morgan, M. M., & Ingram, S. L. (2014). Contribution of Adenylyl

Cyclase Modulation of Pre- and Postsynaptic GABA Neurotransmission to Morphine

Antinociception and Tolerance. *Neuropsychopharmacology 2014 39:9*, *39*(9), 2142–

2152. https://doi.org/10.1038/npp.2014.62

Bresolin de Souza, K., Asker, N., Jönsson, E., Förlin, L., & Sturve, J. (2016). Increased

activity of lysozyme and complement system in Atlantic halibut exposed to elevated

CO_2 at six different temperatures. *Marine Environmental Research*, *122*, 143–147.

https://doi.org/10.1016/J.MARENVRES.2016.09.005

Bushmanova, E., Antipov, D., Lapidus, A., Suvorov, V., & Prjibelski, A. D. (2016).

RnaQUAST: A quality assessment tool for de novo transcriptome assemblies.

Bioinformatics, *32*(14), 2210–2212. https://doi.org/10.1093/bioinformatics/btw218

Bushnell, B. (2014). BBMap: A Fast, Accurate, Splice-Aware Aligner. (No. LBNL-7065E).

Lawrence Berkeley National Lab.(LBNL), Berkeley, CA (United States).

https://www.osti.gov/servlets/purl/1241166

Cabau, C., Escudié, F., Djari, A., Guiguen, Y., Bobe, J., & Klopp, C. (2017). Compacting and

correcting Trinity and Oases RNA-Seq de novo assemblies. *PeerJ*, *2017*(2).

https://doi.org/10.7717/peerj.2988

Casey, J. R., Grinstein, S., & Orlowski, J. (2009). Sensors and regulators of intracellular pH.

Nature reviews Molecular cell biology, 11(1), 50-61.https://doi.org/10.1038/nrm2820

Cattano, C., Calò, A., Di Franco, A., Firmamento, R., Quattrocchi, F., Sdiri, K., Guidetti, P.,

& Milazzo, M. (2017). Ocean acidification does not impair predator recognition but

increases juvenile growth in a temperate wrasse off CO_2 seeps. *Marine Environmental*

Research, *132*(August), 33–40. https://doi.org/10.1016/j.marenvres.2017.10.013

Cattano, C., Claudet, J., Domenici, P., & Milazzo, M. (2018). Living in a high CO_2 world: a

global meta-analysis shows multiple trait-mediated fish responses to ocean acidification.

Ecological Monographs, *88*(3), 320–335. https://doi.org/10.1002/ECM.1297

Chen, S., Zhou, Y., Chen, Y., & Gu, J. (2018). Fastp: An ultra-fast all-in-one FASTQ

preprocessor. *Bioinformatics*, *34*(17), i884–i890.

https://doi.org/10.1093/bioinformatics/bty560

Connell, S. D., Kroeker, K. J., Fabricius, K. E., Kline, D. I., & Russell, B. D. (2013). The

other ocean acidification problem: CO_2 as a resource among competitors for ecosystem

dominance. *Philosophical Transactions of the Royal Society B: Biological Sciences*,

368(1627). https://doi.org/10.1098/RSTB.2012.0442

Currais, A., Huang, L., Goldberg, J., Petrascheck, M., Ates, G., Pinto-Duarte, A., Shokhirev,

M. N., Schubert, D., & Maher, P. (2019). Elevating acetyl-CoA levels reduces aspects of

brain aging. *ELife*, *8*. https://doi.org/10.7554/ELIFE.47866

Davidson, N. M., & Oshlack, A. (2014). Corset: Enabling differential gene expression

analysis for de novo assembled transcriptomes. *Genome Biology*, *15*(7), 410.

https://doi.org/10.1186/s13059-014-0410-6

de Curtis, I., & Meldolesi, J. (2012). Cell surface dynamics - how Rho GTPases orchestrate

the interplay between the plasma membrane and the cortical cytoskeleton. *Journal of*

Cell Science, *125*(Pt 19), 4435–4444. https://doi.org/10.1242/JCS.108266

De Souza, K. B., Jutfelt, F., Kling, P., Förlin, L., & Sturve, J. (2014). Effects of Increased

CO_2 on Fish Gill and Plasma Proteome. *PLOS ONE*, *9*(7), e102901.

https://doi.org/10.1371/JOURNAL.PONE.0102901

Dmitriev, A. V., & Mangel, S. C. (2000). A circadian clock regulates the pH of the fish

retina. *The Journal of Physiology*, *522*(Pt 1), 77. https://doi.org/10.1111/J.1469-

7793.2000.0077M.X

Domenici, P., Allan, B., McCormick, M. I., & Munday, P. L. (2012). Elevated carbon dioxide

affects behavioural lateralization in a coral reef fish. *Biology Letters*, *8*(1), 78–81.

https://doi.org/10.1098/RSBL.2011.0591

Doney, S. C., Fabry, V. J., Feely, R. A., & Kleypas, J. A. (2009). Ocean acidification: The

other CO_2 problem. *Annual Review of Marine Science*, *1*, 169–192.

https://doi.org/10.1146/annurev.marine.010908.163834

Doney, S. C., Ruckelshaus, M., Emmett Duffy, J., Barry, J. P., Chan, F., English, C. A.,

Galindo, H. M., Grebmeier, J. M., Hollowed, A. B., Knowlton, N., Polovina, J.,

Rabalais, N. N., Sydeman, W. J., & Talley, L. D. (2012). Climate change impacts on

marine ecosystems. *Annual Review of Marine Science*, *4*, 11–37.

https://doi.org/10.1146/annurev-marine-041911-111611

Esbaugh, A. J. (2017). Physiological implications of ocean acidification for marine fish:

emerging patterns and new insights. *Journal of Comparative Physiology B 2017 188:1*,

188(1), 1–13. https://doi.org/10.1007/S00360-017-1105-6

Fabricius, K. E., Langdon, C., Uthicke, S., Humphrey, C., Noonan, S., De'ath, G., Okazaki,

R., Muehllehner, N., Glas, M. S., & Lough, J. M. (2011). Losers and winners in coral

reefs acclimatized to elevated carbon dioxide concentrations. *Nature Climate Change*

2011 1:3, *1*(3), 165–169. https://doi.org/10.1038/NCLIMATE1122

Fabry, V. J., Seibel, B. A., Feely, R. A., & Orr Fabry, J. C. (2008). Impacts of ocean

acidification on marine fauna and ecosystem processes. *ICES Journal of Marine*

Science, 65(3), 414-432. https://academic.oup.com/icesjms/article/65/3/414/789605

Ferrari, M. C. O., Dixson, D. L., Munday, P. L., Mccormick, M. I., Meekan, M. G., Sih, A.,

& Chivers, D. P. (2011). Intrageneric variation in antipredator responses of coral reef

fishes affected by ocean acidification: implications for climate change projections on

marine communities. *Global Change Biology*, *17*(9), 2980–2986.

https://doi.org/10.1111/J.1365-2486.2011.02439.X

Ferrari, M. C. O., Mccormick, M. I., Munday, P. L., Meekan, M. G., Dixson, D. L.,

Lönnstedt, O., & Chivers, D. P. (2012). Effects of ocean acidification on visual risk

assessment in coral reef fishes. *Functional Ecology*, *26*(3), 553–558.

https://doi.org/10.1111/j.1365-2435.2011.01951.x

Flor, P. J., Lukic, S., Rüegg, D., Leonhardt, T., Knöpfel, T., & Kühn, R. (1995). Molecular

cloning, functional expression and pharmacological characterization of the human

metabotropic glutamate receptor type 4. *Neuropharmacology*, *34*(2), 149–155.

https://doi.org/10.1016/0028-3908(94)00149-M

Foll, M., & Gaggiotti, O. (2008). A genome-scan method to identify selected loci appropriate

for both dominant and codominant markers: a Bayesian perspective. *Genetics*, *180*(2),

977–993. https://doi.org/10.1534/GENETICS.108.092221

Galachyants, Y. P., Zakharova, Y. R., Volokitina, N. A., Morozov, A. A., Likhoshway, Y. V.,

& Grachev, M. A. (2019). De novo transcriptome assembly and analysis of the

freshwater araphid diatom Fragilaria radians, Lake Baikal. *Scientific Data*, *6*(1), 1–11.

https://doi.org/10.1038/s41597-019-0191-6

Gilmour, K. M., & Perry, S. F. (2009). Carbonic anhydrase and acid-base regulation in fish.

Journal of Experimental Biology, *212*(11), 1647–1661.

https://doi.org/10.1242/jeb.029181

Gö Tz, S., García-Gó Mez, J. M., Terol, J., Williams, T. D., Nagaraj, S. H., Nueda, M. J.,

Robles, M., Taló, M., Dopazo, J., & Conesa, A. (2008). High-throughput functional

annotation and data mining with the Blast2GO suite. *Nucleic Acids Research*, *36*(10),

3420–3435. https://doi.org/10.1093/nar/gkn176

Grabherr, M. G. ., Brian J. Haas, Moran Yassour Joshua Z. Levin, Dawn A. Thompson, Ido

Amit, Xian Adiconis, Lin Fan, Raktima Raychowdhury, Qiandong Zeng, Zehua Chen,

Evan Mauceli, Nir Hacohen, Andreas Gnirke, Nicholas Rhind, Federica di Palma, Bruce

W., N., & Friedman, and A. R. (2013). Trinity: reconstructing a full-length

transcriptome without a genome from RNA-Seq data. *Nature Biotechnology*, *29*(7),

644–652. https://doi.org/10.1038/nbt.1883.Trinity

Grosell, M. (2011). Intestinal anion exchange in marine teleosts is involved in

osmoregulation and contributes to the oceanic inorganic carbon cycle. *Acta Physiologica*

(Oxford, England), *202*(3), 421–434. https://doi.org/10.1111/j.1748-1716.2010.02241.x

Haas, B. J., Papanicolaou, A., Yassour, M., Grabherr, M., Philip, D., Bowden, J., Couger, M.

B., Eccles, D., Li, B., Macmanes, M. D., Ott, M., Orvis, J., Pochet, N., Strozzi, F.,

Weeks, N., Westerman, R., William, T., Dewey, C. N., Henschel, R., LeDuc, R. D.,

Friedman, N., Regev, A. (2013). De novo transcript sequence recostruction from RNA-

Seq: reference generation and analysis with Trinity. In *Nature protocols* (Vol. 8, Issue

8). https://doi.org/10.1038/nprot.2013.084.De

Hall-Spencer, J. M., Rodolfo-Metalpa, R., Martin, S., Ransome, E., Fine, M., Turner, S. M.,

Rowley, S. J., Tedesco, D., & Buia, M. C. (2008). Volcanic carbon dioxide vents show

ecosystem effects of ocean acidification. *Nature 2008 454:7200*, *454*(7200), 96–99.

https://doi.org/10.1038/NATURE07051

Hamilton, S. L., Logan, C. A., Fennie, H. W., Sogard, S. M., Barry, J. P., Makukhov, A. D.,

Tobosa, L. R., Boyer, K., Lovera, C. F., & Bernardi, G. (2017). Species-Specific

Responses of Juvenile Rockfish to Elevated pCO_2: From Behavior to Genomics. *PLOS*

ONE, *12*(1), e0169670. https://doi.org/10.1371/JOURNAL.PONE.0169670

Hamilton, T. J., Holcombe, A., & Tresguerres, M. (2014). CO_2-induced ocean acidification

increases anxiety in Rockfish via alteration of GABAA receptor functioning.

Proceedings of the Royal Society B: Biological Sciences, *281*(1775).

https://doi.org/10.1098/RSPB.2013.2509

Hetz, C. (2012). The unfolded protein response: controlling cell fate decisions under ER

stress and beyond. *Nature Reviews. Molecular Cell Biology*, *13*(2), 89–102.

https://doi.org/10.1038/NRM3270

Heuer, R M, Welch, M. J., Rummer, J. L., Munday, P. L., & Grosell, & M. (2016). Altered

brain ion gradients following compensation for elevated CO_2 are linked to behavioural

alterations in a coral reef fish. *Nature Publishing Group*.

https://doi.org/10.1038/srep33216

Heuer, Rachael M., & Grosell, M. (2014). Physiological impacts of elevated carbon dioxide

and ocean acidification on fish. *American Journal of Physiology - Regulatory*

Integrative and Comparative Physiology, *307*(9), R1061–R1084.

https://doi.org/10.1152/AJPREGU.00064.2014/SUPPL_FILE/TABLES1.PDF

Hofmann, G. E., Barry, J. P., Edmunds, P. J., Gates, R. D., Hutchins, D. A., Klinger, T., &

Sewell, M. A. (2010). The effect of Ocean acidification on calcifying organisms in

marine ecosystems: An organism-to-ecosystem perspective. *Annual Review of Ecology,*

Evolution, and Systematics, *41*, 127–147.

https://doi.org/10.1146/annurev.ecolsys.110308.120227

Howe, K. L., Achuthan, P., Allen, J., Allen, J., Alvarez-Jarreta, J., Amode, M. R., Armean, I.

M., Azov, A. G., Bennett, R., Bhai, J., Billis, K., Boddu, S., Charkhchi, M., Cummins,

C., Da, L., Fioretto, R., Davidson, C., Dodiya, K., Houdaigui, B. El, … Flicek, P.

(2021). Ensembl 2021. *Nucleic Acids Research*, *49*(2).

https://doi.org/10.1093/nar/gkaa942

Hübner, C. A., & Holthoff, K. (2013). Anion transport and GABA signaling. *Frontiers in*

Cellular Neuroscience, *0*(OCT), 177.

https://doi.org/10.3389/FNCEL.2013.00177/BIBTEX

Ishimatsu, A., Hayashi, M., Lee, K. S., Kikkawa, T., & Kita, J. (2005). Physiological effects

on fishes in a high-CO_2 world. *Journal of Geophysical Research: Oceans*, *110*(C9), 1–8.

https://doi.org/10.1029/2004JC002564

Jentsch, T. J., Stein, V., Weinreich, F., & Zdebik, A. A. (2002). Molecular structure and

physiological function of chloride channels. *Physiological Reviews*, *82*(2), 503–568.

https://doi.org/10.1152/physrev.00029.2001

Juan, C. A., de la Lastra, J. M. P., Plou, F. J., & Pérez-Lebeña, E. (2021). The Chemistry of

Reactive Oxygen Species (ROS) Revisited: Outlining Their Role in Biological

Macromolecules (DNA, Lipids and Proteins) and Induced Pathologies. *International*

Journal of Molecular Sciences, *22*(9). https://doi.org/10.3390/IJMS22094642

Kang, J., Nagelkerken, I., Rummer, J. L., Rodolfo-Metalpa, R., Munday, P. L., Ravasi, T., &

Schunter, C. (2022). Rapid evolution fuels transcriptional plasticity to ocean

acidification. *Global Change Biology*. https://doi.org/10.1111/GCB.16119

Kelly, M. W., & Hofmann, G. E. (2013). Adaptation and the physiology of ocean

acidification. *Functional Ecology*, *27*(4), 980–990. https://doi.org/10.1111/J.1365-

2435.2012.02061.X

Kolosov, D., Bui, P., Chasiotis, H., & Kelly, S. P. (2013). Claudins in teleost fishes. *Tissue*

Barriers, *1*(3), e25391. https://doi.org/10.4161/TISB.25391

Kroeker, K. J., Kordas, R. L., Crim, R. N., & Singh, G. G. (2010). Meta-analysis reveals

negative yet variable effects of ocean acidification on marine organisms. *Ecology*

Letters, *13*(11), 1419–1434. https://doi.org/10.1111/j.1461-0248.2010.01518.x

Langmead, B., & Salzberg, S. L. (2012). Fast gapped-read alignment with Bowtie 2. *Nature*

Methods, *9*(4), 357–359. https://doi.org/10.1038/nmeth.1923

Lee, D.-W., Song, J. A., Park, H.-S., Cheol, ·, & Choi, Y. (2021). Circadian Rhythm

Disturbances Due to Exposure to Acidified Conditions and Different Photoperiods in

Juvenile Olive Flounder (Paralichthys olivaceus). *Ocean Science Journal*, *56*, 198–206.

https://doi.org/10.1007/s12601-021-00018-y

Li, B., & Dewey, C. N. (2014). RSEM: Accurate transcript quantification from RNA-seq data

with or without a reference genome. In *Bioinformatics: The Impact of Accurate*

Quantification on Proteomic and Genetic Analysis and Research (pp. 41–74). Apple

Academic Press. https://doi.org/10.1201/b16589

Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. *Genome Biology*, *15*(12), 1–21.

https://doi.org/10.1186/s13059-014-0550-8

Machado, M., Arenas, F., Svendsen, J. C., Azeredo, R., Pfeifer, L. J., Wilson, J. M., &

Costas, B. (2020). Effects of Water Acidification on Senegalese Sole Solea senegalensis

Health Status and Metabolic Rate: Implications for Immune Responses and Energy Use.

Frontiers in Physiology, *11*, 26. https://doi.org/10.3389/FPHYS.2020.00026/BIBTEX

Manni, M., Berkeley, M. R., Seppey, M., Simão, F. A., & Zdobnov, E. M. (2021). BUSCO

Update: Novel and Streamlined Workflows along with Broader and Deeper

Phylogenetic Coverage for Scoring of Eukaryotic, Prokaryotic, and Viral Genomes.

Molecular Biology and Evolution, *38*(10), 4647–4654.

https://doi.org/10.1093/MOLBEV/MSAB199

Mirasole, A., Gillanders, B. M., Reis-Santos, P., Grassa, F., Capasso, G., Scopelliti, G.,

Mazzola, A., & Vizzini, S. (2017). The influence of high pCO_2 on otolith shape,

chemical and carbon isotope composition of six coastal fish species in a Mediterranean

shallow CO_2 vent. *Marine Biology*, *164*(9). https://doi.org/10.1007/s00227-017-3221-y

Mirasole, A., Scopelliti, G., Tramati, C., Signa, G., Mazzola, A., & Vizzini, S. (2021).

Evidences on alterations in skeleton composition and mineralization in a site-attached

fish under naturally acidified conditions in a shallow CO_2 vent. *Science of The Total*

Environment, *761*, 143309. https://doi.org/10.1016/J.SCITOTENV.2020.143309

Mirasole, A., Signa, G., Gianguzza, P., Bonaviri, C., Mazzola, A., & Vizzini, S. (2019). Fish

assemblages cope with ocean acidification in a shallow volcanic CO_2 vent benefiting

from an adjacent recovery area. *Marine Environmental Research*.

https://doi.org/10.1016/j.marenvres.2019.104851

Mittermayer, H., Stiasny, M. H., Puvanendran, V., Chierici, M., Jentoft, S., & Reusch, B. H.

(2019). Transcriptome profiling reveals exposure to predicted end-of-century ocean

acidification as a stealth stressor for Atlantic cod larvae. *Scientific Reports*, *9(1)*, 1–11.

https://doi.org/10.1038/s41598-019-52628-1

Monroe, A. A., Schunter, C., Welch, M. J., Munday, P. L., & Ravasi, T. (2021). Molecular

basis of parental contributions to the behavioural tolerance of elevated pCO_2 in a coral

reef fish. *Proceedings of the Royal Society B: Biological Sciences*, *288*(1964).

https://doi.org/10.1098/rspb.2021.1931

Munday, P. L., Cheal, A. J., Dixson, D. L., Rummer, J. L., & Fabricius, K. E. (2014).

Behavioural impairment in reef fishes caused by ocean acidification at CO_2 seeps.

Nature Climate Change, *4*(6), 487–492. https://doi.org/10.1038/nclimate2195

Munday, P. L., Dixson, D. L., McCormick, M. I., Meekan, M., Ferrari, M. C. O., & Chivers,

D. P. (2010). Replenishment of fish populations is threatened by ocean acidification.

Proceedings of the National Academy of Sciences of the United States of America,

107(29), 12930–12934.

https://doi.org/10.1073/PNAS.1004519107/SUPPL_FILE/PNAS.201004519SI.PDF

Munday, P. L., McCormick, M. I., Meekan, M., Dixson, D. L., Watson, S.-A., Chivers, D. P.,

& Ferrari, M. C. O. (2013). Selective mortality associated with variation in CO_2

tolerance in a marine fish. *Ocean Acidification*, *1*, 1–5. https://doi.org/10.2478/OAC-

2012-0001

Munday, P. L., Pratchett, M. S., Dixson, D. L., Donelson, J. M., Endo, G. G. K., Reynolds,

A. D., & Knuckey, R. (2013). Elevated CO_2 affects the behavior of an ecologically and

economically important coral reef fish. *Marine Biology*, *160*(8), 2137–2144.

https://doi.org/10.1007/S00227-012-2111-6/FIGURES/2

Nagelkerken, I., Russell, B. D., Gillanders, B. M., Connell, S. D., & Kong SAR, H. (2015).

Ocean acidification alters fish populations indirectly through habitat modification.

Nature Climate Change, 6(1), 89-93. https://doi.org/10.1038/NCLIMATE2757

Nilsson, G. E., Dixson, D. L., Domenici, P., McCormick, M. I., Sørensen, C., Watson, S. A.,

& Munday, P. L. (2012). Near-future carbon dioxide levels alter fish behaviour by

interfering with neurotransmitter function. *Nature Climate Change*, *2*(3), 201–204.

https://doi.org/10.1038/nclimate1352

OmicsBox – Bioinformatics Made Easy, BioBam Bioinformatics, March 3. (2019).

Paula, J. R., Baptista, M., Carvalho, F., Repolho, T., Bshary, R., & Rosa, R. (2019). The past,

present and future of cleaner fish cognitive performance as a function of CO_2 levels.

Biology Letters, *15*(12). https://doi.org/10.1098/RSBL.2019.0618

Petit-Marty, N., Nagelkerken, I., Connell, S. D., & Schunter, C. (2021). Natural CO_2 seeps

reveal adaptive potential to ocean acidification in fish. *Evolutionary Applications*, *14*(7),

1794–1806. https://doi.org/10.1111/eva.13239

Pizzino, G., Irrera, N., Cucinotta, M., Pallio, G., Mannino, F., Arcoraci, V., Squadrito, F.,

Altavilla, D., & Bitto, A. (2017). Oxidative Stress: Harms and Benefits for Human

Health. *Oxidative Medicine and Cellular Longevity*, *2017*.

https://doi.org/10.1155/2017/8416763

Poli, G., Leonarduzzi, G., Biasi, F., & Chiarpotto, E. (2012). Oxidative Stress and Cell

Signalling. *Current Medicinal Chemistry*, *11*(9), 1163–1182.

https://doi.org/10.2174/0929867043365323

Prokkola, J. M., & Nikinmaa, M. (2018). Circadian rhythms and environmental disturbances

- Underexplored interactions. *Journal of Experimental Biology*, *221*(16).

https://doi.org/10.1242/JEB.179267/19339

Qin, Z., Lewis, J. E., & Perry, S. F. (2010). Zebrafish (Danio rerio) gill neuroepithelial cells

are sensitive chemoreceptors for environmental CO_2. *The Journal of Physiology*, *588*(5),

861–872. https://doi.org/10.1113/JPHYSIOL.2009.184739

Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., & Glöckner,

F. O. (2013). The SILVA ribosomal RNA gene database project: Improved data

processing and web-based tools. *Nucleic Acids Research*, *41*(D1), 590–596.

https://doi.org/10.1093/nar/gks1219

R Core Team. (2021). R: A language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna, Austria. https://www.r-project.org/.

Radford, C. A., Collins, S. P., Munday, P. L., & Parsons, D. (2021). Ocean acidification

effects on fish hearing. *Proceedings of the Royal Society B: Biological Sciences*,

288(1946). https://doi.org/10.1098/rspb.2020.2754

Rummer, J. L., & Brauner, C. J. (2011). Plasma-accessible carbonic anhydrase at the tissue of

a teleost fish may greatly enhance oxygen delivery: in vitro evidence in rainbow trout,

Oncorhynchus mykiss. *Journal of Experimental Biology*, *214*(14), 2319–2328.

https://doi.org/10.1242/JEB.054049

Sahraeian, S. M. E., Mohiyuddin, M., Sebra, R., Tilgner, H., Afshar, P. T., Au, K. F., Bani

Asadi, N., Gerstein, M. B., Wong, W. H., Snyder, M. P., Schadt, E., & Lam, H. Y. K.

(2017). Gaining comprehensive biological insight into the transcriptome by performing

a broad-spectrum RNA-seq analysis. *Nature Communications*, *8*(1), 1–14.

https://doi.org/10.1038/s41467-017-00050-4

Schunter, C., Jarrold, M. D., Munday, P. L., & Ravasi, T. (2021). Diel pCO_2 fluctuations alter

the molecular response of coral reef fishes to ocean acidification conditions. *Molecular*

Ecology, *30*(20), 5105–5118. https://doi.org/10.1111/mec.16124

Schunter, C., Ravasi, T., Munday, P. L., & Nilsson, G. E. (2019). Neural effects of elevated

CO_2 in fish may be amplified by a vicious cycle. *Conservation Physiology*, *7*(1).

https://doi.org/10.1093/CONPHYS/COZ100

Schunter, C., Welch, M. J., Nilsson, G. E., Rummer, J. L., Munday, P. L., & Ravasi, T.

(2018). An interplay between plasticity and parental phenotype determines impacts of

ocean acidification on a reef fish. *Nature Ecology & Evolution*, *2*(2), 334–342.

https://doi.org/10.1038/s41559-017-0428-8

Schunter, C., Welch, M. J., Ryu, T., Zhang, H., Berumen, M. L., Nilsson, G. E., Munday, P.

L., & Ravasi, T. (2016). Molecular signatures of transgenerational response to ocean

acidification in a species of reef fish. *Nature Climate Change*, *6*(11), 1014–1018.

https://doi.org/10.1038/nclimate3087

Shimokawa, N., Okada, J., Haglund, K., Dikic, I., Koibuchi, N., & Miura, M. (2002). Past-A,

a novel proton-associated sugar transporter, regulates glucose homeostasis in the brain.

The Journal of Neuroscience : The Official Journal of the Society for Neuroscience,

22(21), 9160–9165. https://doi.org/10.1523/JNEUROSCI.22-21-09160.2002

Simpson, S. D., Munday, P. L., Wittenrich, M. L., Manassa, R., Dixson, D. L., Gagliano, M.,

& Yan, H. Y. (2011). Ocean acidification erodes crucial auditory behaviour in a marine

fish. *Biology Letters*, *7*(6), 917–920. https://doi.org/10.1098/RSBL.2011.0293

Singh Jaggi, A., Kaur, A., Bali, A., & Singh, N. (2015). Expanding Spectrum of Sodium

Potassium Chloride Co-transporters in the Pathophysiology of Diseases. *Current*

Neuropharmacology, 13(3), 369-388.

Spatafora, D., Cattano, C., Aglieri, G., Quattrocchi, F., Turco, G., Quartararo, G.,

Dudemaine, J., Calosi, P., & Milazzo, M. (2022). Limited behavioural effects of ocean

acidification on a Mediterranean anemone goby (*Gobius incognitus*) chronically

exposed to elevated CO_2 levels. *Marine Environmental Research*, *181*, 105758.

https://doi.org/10.1016/J.MARENVRES.2022.105758

Strader, M. E., Wong, J. M., & Hofmann, G. E. (2020). Ocean acidification promotes broad

transcriptomic responses in marine metazoans: A literature survey. *Frontiers in Zoology*,

17(1), 1–23. https://doi.org/10.1186/s12983-020-0350-9

Stubblefield, J. J., Terrien, J., & Green, C. B. (2012). Nocturnin: At the crossroads of clocks

and metabolism. *Trends in Endocrinology and Metabolism*, *23*(7), 326–333.

https://doi.org/10.1016/j.tem.2012.03.007

Sunday, J. M., Calosi, P., Dupont, S., Munday, P. L., Stillman, J. H., & Reusch, T. B. H.

(2014). Evolution in an acidifying ocean. *Trends in Ecology & Evolution*, *29*(2), 117–

125. https://doi.org/10.1016/J.TREE.2013.11.001

Thacker, C. E., & Roje, D. M. (2011). Phylogeny of Gobiidae and identification of gobiid

lineages. *Systematics and Biodiversity*, *9*(4), 329–347.

https://doi.org/10.1080/14772000.2011.629011

Tresguerres, M., Clifford, A. M., Harter, T. S., Roa, J. N., Thies, A. B., Yee, D. P., &

Brauner, C. J. (2020). Evolutionary links between intra-and extracellular acid-base

regulation in fish and other aquatic animals. *J. Exp. Zool*, *333*, 449–465.

https://doi.org/10.1002/jez.2367

Tresguerres, M., & Hamilton, T. J. (2017). Acid–base physiology, neurobiology and

behaviour in relation to CO_2-induced ocean acidification. *Journal of Experimental*

Biology, *220*(12), 2136–2148. https://doi.org/10.1242/JEB.144113

Vizzini, S., Martínez-Crego, B., Andolina, C., Massa-Gallucci, A., Connell, S. D., & Gambi,

M. C. (2017). Ocean acidification as a driver of community simplification via the

collapse of higher-order and rise of lower-order consumers. *Scientific Reports 2017 7:1*,

7(1), 1–10. https://doi.org/10.1038/s41598-017-03802-w

Welch, M. J., & Munday, P. L. (2017). Heritability of behavioural tolerance to high CO_2 in a

coral reef fish is masked by nonadaptive phenotypic plasticity. *Evolutionary*

Applications, *10*(7), 682–693. https://doi.org/10.1111/eva.12483

Wood, D. E., & Salzberg, S. L. (2014). Kraken: Ultrafast metagenomic sequence

classification using exact alignments. *Genome Biology*, *15*(3).

https://doi.org/10.1186/gb-2014-15-3-r46

Zhang, J., Wang, X., Vikash, V., Ye, Q., Wu, D., Liu, Y., & Dong, W. (2016). ROS and

ROS-Mediated Cellular Signaling. *Oxidative Medicine and Cellular Longevity*, *2016*.

https://doi.org/10.1155/2016/4350965

Zunino, S., Canu, D. M., Zupo, V., & Solidoro, C. (2019). Direct and indirect impacts of

marine acidification on the ecosystem services provided by coralligenous reefs and

seagrass systems. *Global Ecology and Conservation*, *18*, e00625.

https://doi.org/10.1016/J.GECCO.2019.E00625

928