Title: Medial accumbens shell spiny projection neurons encode relative reward preference

Authors: Christian E. Pedersen[1,5], Daniel C. Castro[2,3,5], Madelyn M. Gray[3,5], Zhe C. Zhou[5], Sean C. Piantadosi[3,5], Raajaram Gowrishankar[3,5], Shane A. Kan[5], Patrick J. Murphy[5], Patrick R. O'Neill[2], Michael R. Bruchas[1,2,3,4,5*]

[1] Department of Bioengineering, University of Washington, Seattle, WA, 98195
[2] Departments of Anesthesiology, Neuroscience and Psychiatry, and Washington University Pain Center, Washington University School of Medicine, St. Louis, MO 63110
[3] Department of Anesthesiology and Pain Medicine, University of Washington, Seattle, Washington, 98195
[4] Department of Pharmacology, University of Washington, Seattle, Washington 98195
[5] Center for Neurobiology of Addiction, Pain, and Emotion, University of Washington, Seattle, Washington, 98195

*Corresponding Authors:
Michael R. Bruchas: mbruchas@uw.edu

Abstract

Medial nucleus accumbens shell (mNAcSh) is a critical brain region for driving motivated behaviors. Despite this well-established role, the underlying reward processing of individual neurons, circuits and cell-types within mNAcSh remains largely unknown. Here, we leverage deep brain 2-photon calcium imaging through endoscopic lenses to record mNAcSh spiny projection neuron (SPN) ensemble responses to rewards of different preference and to reward-predictive cues across cue-reward learning. Reward responses were found to be heterogeneous and particularly differentiated based on reward preference and cell type. A large subpopulation of reward-excited enkephalinergic SPNs were found to be specifically recruited during consumption of unpreferred rewards. A major enkephalinergic efferent projection from mNAcSh to ventral pallidum (VP) was also found to be recruited to unpreferred rewards and to causally drive low positive reward preference. Enkephalin and dynorphinergic SPNs in mNAcSh distinctly represent rewards of different preference and propagate distinct signals through efferent projections to drive consummatory behavior.

The brain must compute and represent distinct identities of rewards to facilitate reward

preference and to drive focused reward acquisition. The nucleus accumbens (NAc) is

established as a major brain region for driving reward seeking and consumption (*1*, *2*). The NAc

has many glutamatergic afferents such as basolateral amygdala, prefrontal cortex, ventral

pallidum (VP) and ventral hippocampus and dopaminergic innervation from ventral tegmental

area (VTA) (*3*). These NAc afferents have been shown to be highly reward modulated, and in

many cases, even reflect the quantity and value/preference of rewards (*4–9*).

The NAc is primarily composed of two dissociable types of spiny projection neurons

(SPN) that are canonically defined by their expression of either dopamine 1 (D1R) or dopamine

2 (D2R) type receptors. D1R-SPNs also largely co-express the endogenous opioid peptide

dynorphin, while D2R-SPNs primarily express enkephalin (*10*, *11*). Previous studies have

shown NAc SPNs to be modulated by rewards and reward-predictive cues but have not

resolved the molecular identity of the recorded neurons or tracked changes in activity of

individual neurons across multiple days of cue-reward learning (*12*, *13*). Most reports have also

primarily focused on NAc core and have not given specific attention to the reward-responsivity

of the medial shell subregion (mNAcSh). mNAcSh has been shown to be very distinct from NAc

core in its connectivity, function and dopaminergic innervation (*3*, *14–18*). Even within mNAcSh,

functional distinctions have been isolated across anterior-posterior and dorsal-ventral axes (*2*,

11).

Here, we sought to characterize the responses of individual mNAcSh SPNs to rewards

of different identities and across cue-reward learning with particular focus on the molecular

identity and relative anatomical position of the record cells. Recent experimental advances have

allowed for the simultaneous recording of dozens of genetically defined neurons in deep brain

structures using 2-photon calcium imaging through endoscopic lenses (*19*, *20*). The low noise-

floor of 2-photon microscopy also allows for the resolution of individual cell morphology and

permits high fidelity tracking of individual neurons across multiple days (*21*). In the current

study, we leverage this technique to conduct longitudinal studies on how mNAcSh SPNs

respond to multiple reward identities and change across cue-reward learning. This high

resolution approach revealed a distinct subpopulation of enkephalinergic SPNs (Penk-SPN) that

are selectively activated during the consumption of less preferred reward. We then isolate a

Penk-SPN circuit from mNAcSh to VP is selectively activated during less preferred reward to

causally drive a low, positive reward preference. These results provide foundational insight into

how relative reward preference information is integrated and separated across mNAcSh

neuronal ensembles and how these SPNs contribute to driving consummatory behavior.

In order to characterize differences in how rewards of varying preference are processed

in mNAcSh, we presented mice with 4 distinct reward conditions that had differing concentration

but similar taste: 0% (water), 3%, 10% and 30% sucrose. To test the relative preference of the 4

sucrose reward conditions to water-restricted mice, 2-bottle choice experiments were conducted

in which the animal was given access to 2 of the 4 conditions at a time and had to choose the

preferred reward (**Extended Data Fig. 1a**). Previous studies have consistently found that mice

and rats prefer sucrose concentrations between 10-15% and that the distribution of sucrose

preferences is shifted toward lower concentrations during water deprivation (*22–24*). Mice also

tend to drink smaller volumes of higher concentration sucrose (20%+) due to satiety (*25*). To

test the relative preference of the 4 sucrose reward conditions while minimizing effects from

satiety, we tested mice in an adapted 2-bottle choice assay where the animals were only given

brief access (20 s) to sucrose sippers with a variable intertrial interval between access periods.

It was found that water-restricted C57 mice preferred 10% sucrose over all other conditions and

30% sucrose less than all other conditions (**Extended Data Fig. 1e**).

To measure the activity of mNAcSh SPNs during the consumption of varying reward

preference conditions, we imaged the calcium dynamics of hundreds of individual SPNs using

2-photon fluorescence microscopy through endoscopic GRIN lenses. Preprodynorphin-cre

(Pdyn-cre) and proenkephalin-cre (Penk-cre) animals were stereotaxically injected with 500 nl of

AAV-DJ-Ef1α-DIO-GCaMP6s virus and implanted with a 600 um diameter cylindrical GRIN lens

in mNAcSh (**Fig. 1a**). After 4-6 weeks of recovery from surgery, animals were water restricted

and trained to sip 10% sucrose reward at variable intervals while reversibly immobilized and

head-restrained (**Fig. 1b**). All animals displayed consistent licking behavior while immobilized

after 4 days of lick training and were then tested in 30 minute, 20 trial calcium imaging sessions

each day for 4 consecutive days. In each session, animals consumed 1 of 4 possible sucrose

reward conditions (0%, 3%, 10% or 30% sucrose in water), with the order of condition days

pseudorandomized and counterbalanced across animals (**Fig. 1c**). Animals immediately licked

the presented sucrose reward, with an average licking bout duration of 7-8 seconds (**Fig. 1d**).

There was no significant difference between reward conditions in the average amount of licking

for each reward due to the small 10 ul reward volumes (**Fig. 1e**). This uniformity in lick behavior

allowed for accurate comparison of SPN activity in response to the content of the different

reward identities with minimal variation in behavioral output.

The resolution and low noise floor of 2-photon calcium imaging allowed for visualization

of individual cell morphology and enabled high fidelity tracking of 715 individual SPNs across

multiple days and conditions (**Fig. 1g-h, Extended Data Fig. 2a**). Individually tracked Pdyn-

SPNs and Penk-SPNs displayed reward responses that were consistent for a given sucrose

concentration, but that varied between the different sucrose concentrations (**Extended Data**

Fig. 2b-c). Groups of individual SPNs were also correlated in their selectivity for specific

sucrose concentrations. In order to group individual neurons together that had similar response

profiles across the 4 sucrose conditions, k-means clustering was performed across all tracked

SPNs. This unbiased approach to grouping neurons on the basis of similar activity revealed that

there were 4 distinct clusters of Penk-SPNs and 5 distinct clusters of Pdyn-SPNs (**Fig. 1o-r,**

Extended Data Fig. 2d-g). Notably, every cluster was composed of neurons from nearly all, if

not all, of the mice that were tested, demonstrating that these functionally-defined mNAcSh

clusters are generalizable across animals (**Extended Data Fig. 2b-c**). 32% of tracked Penk-

SPNs and 41% of tracked Pdyn-SPNs were not consistently modulated by any of the reward

conditions (**Fig. 1i**). A greater proportion of Penk-SPNs (32%) were inhibited in response to all 4

sucrose conditions than for Pdyn-SPNs (18%) (**Fig. 1j**). A large cluster of Penk-SPNs and Pdyn-

SPNs were found to be active in response to the preferred 0%, 3% and 10% conditions, but

were relatively quiescent in response to the unpreferred (see **Extended Data Fig. 1**) 30%

sucrose condition (**Fig. 1k**). Nearly half of all reward-excited Penk-SPNs were selectively active

far more for the unpreferred 30% sucrose condition (**Fig. 1l**). The presence of 3 robust and

complimentary SPN clusters that are differently activated at the 30% condition indicates that

there is a dramatic shift of mNAcSh SPN recruitment during unpreferred rewards. A large 30%

sucrose selective population of Penk-SPNs is recruited only for the unpreferred condition and

this occurs simultaneously as otherwise reward-excited SPNs are systematically silent (**Fig. 1k-**

l).

A distinct cluster of Pdyn-SPNs was also found to be active to all sucrose reward

conditions (3%, 10% and 30%) but much less active to water (0% sucrose) (**Fig. 1m**). A small

minority of Pdyn-SPNs (7%) were also found to be highly excited by every reward condition

(**Fig. 1n**). To further quantify the specificity of mNAcSh SPN activity to the 4 reward preference

conditions, a linear Support Vector Classifier (SVC) model was created. This SVC model was

trained to predict which of the 4 sucrose conditions an animal consumed on a given trial only by

considering the mean activity from each SPN during the reward epoch (**Fig. 1s**). The SVC

performed with high accuracy and specificity for all 4 reward conditions for both Pdyn-SPNs and

Penk-SPNs (**Fig. 1v**). The SVC model coefficients were greater for SPNs in the Preferred and

Unpreferred clusters for discriminating between 30% sucrose and the other reward conditions

(**Fig. 1t-u**). This reaffirms that the primary differences in recruitment of reward-excited SPNs

were occurring between the preferred and unpreferred reward conditions. Altogether, these

mNAcSh calcium imaging experiments across multiple reward preference conditions revealed

substantial heterogeneity in the reward-evoked activity of individual SPNs. There are distinct

mNAcSh SPN populations that are selective for preferred reward conditions and populations

that are selective for unpreferred reward conditions. Reward-excited Penk-SPNs were highly

selective in their reward response on the basis of reward preference, while reward-excited

Pdyn-SPNs were less selective for particular reward preference conditions. Consistent

recruitment of distinct mNAcSh SPN subpopulations to rewards of different preference suggests

that these neurons encode specific reward preference information.

Since we observed heterogeneity in the activity of mNAcSh SPNs to rewards of different

preference, we then sought to characterize potential differences in how individual mNAcSh

SPNs respond to reward-predictive cues as a function of cue-reward associative learning. The

same animals that were tested in the variable sucrose 2-photon imaging paradigm (see **Fig. 1**)

were also trained to associate an auditory cue to subsequent delivery of 10% sucrose reward

across 7 days of Pavlovian conditioning (**Fig. 2a-c**). Animals again displayed immediate licking

in response to sucrose reward presentation, but initially did not show robust anticipatory licking

to the cue (**Fig. 2d,h**). Animals began to display consistent anticipatory licking after multiple

sessions of Pavlovian conditioning, indicating that the cue-reward association had been learned

(**Fig. 2e,h**).

2-photon calcium imaging of mNAcSh SPNs expressing GCaMP6s occurred

simultaneously to Pavlovian conditioning on training days 1, 3, 5 and 7, and 632 individual

SPNs were successfully tracked across all Pavlovian imaging sessions (**Fig. 2f-g, Extended**

Data Fig. 3a). K-means clustering was performed on all 632 SPNs to stratify groups of neurons

on the basis of similar cue and reward responses. This approach revealed that there were 3

distinct clusters of Penk-SPNs and 4 distinct clusters of Pdyn-SPNs within the context of the

Pavlovian conditioning paradigm (**Fig. 2m-n, Extended Data Fig. 3b-g**). Not all reward-

responsive mNAcSh SPNs (61% of all SPNs) were found to also be cue-responsive (41% of all

SPNs) (**Fig. 2m-n**). There were also no cue-inhibited SPNs observed and all reward-inhibited

SPNs had no consistent cue-response (**Fig. 2j**). There was also a distinct cluster of Pdyn-SPNs

that was reward-excited but not cue-excited (**Fig. 2l**), while all reward-excited Penk-SPNs were

also cue-excited (**Fig. 2k**). The Pdyn-SPN cluster that was only reward-excited also showed

smaller reward-excitation to the 10% sucrose reward in later Pavlovian conditioning days,

suggesting that the expectation of reward diminished the magnitude of reward-excitation (**Fig.**

2l). To test for the persistence of cue and reward-responses during the absence of expected

reward delivery, the latter half (last 10 out of 20 trials) of trials on Pavlovian day 7 were reward

omission trials, where the cue is presented as usual but the expected sucrose reward is not

delivered. The SPNs that were found to be exclusively reward-responsive but not cue-

responsive showed no response during Pavlovian extinction trials when reward was omitted

(**Fig. 2j,l**), further suggesting that the reward-responses were truly specific to the content of the

reward itself. However, cue and reward-excited SPNs displayed a diminished, but persistent cue

and reward-excitation during the 10 extinction trials (**Fig. 2k**).

Surprisingly, the magnitude of cue-excitation of the cue-excited SPN clusters did not

increase across Pavlovian training days (**Fig. 2o**) as is canonically observed in VTA DA neurons

during cue-reward associative learning (*26, 27*). We also found that the magnitude of cue-

excitation was only weakly correlated with amount of anticipatory licking per trial (**Fig. 2p**).

Taken together, these results demonstrate that mNAcSh SPN cue-responses do not robustly

change as a function of cue-reward associative learning and do not reflect reward expectation.

Additionally, Pdyn-SPNs decreased the magnitude of their cue-excitation across trials within an

imaging session, suggesting that the cue-response of Pdyn-SPNs may reflect the salience of

the cue tone rather than its motivational value (**Fig. 2q-r**).

Previous studies have demonstrated stark differences in the molecular identity and

function of SPNs, even within the mNAcSh (*2, 11*). Accordingly, the functionally defined SPN

clusters that were identified in the variable sucrose and Pavlovian conditioning behavioral

paradigms (see **Fig. 1-2**) were analyzed to relate the functional identity of SPNs to their relative

position within the focal plane during in vivo 2-photon calcium imaging. The x and y position of

each tracked SPN in the focal plane was calculated using the ROIs of the identified neuron. The

distribution of x and y positions in the focal plane for each cluster was considered and mapped

to the anterior-posterior and medial-lateral orientation of the animal during the imaging session.

The implanted endoscopic lenses were 600 um in diameter and thus, segregation of functional

SPN clusters within the imaging focal plane represents hundreds of microns of anatomical

separation within mNAcSh. Visualizing the spatial distribution of functionally defined SPN

clusters revealed that reward-inhibited SPNs were specific to anterior portions of the focal

plane, for the variable sucrose and Pavlovian imaging sessions (**Fig. 3a-j**). This strong

divergence in reward response of mNAcSh SPNs across the anterior-posterior axis may

indicate a critical difference in the function of these anatomically separable SPN populations.

Distinct subpopulations of Pdyn-SPNs and Penk-SPNs are recruited in response to

rewards of different preference (see **Fig. 1**). Next, we determined whether the stratification of

reward preference information in mNAcSh is propagated selectively to downstream brain

regions via major mNAcSh efferent projections. We first retrotraced the major output projections

from mNAcSh Penk-SPNs and Pdyn-SPNs using AAVretro viral techniques, retrobeads and

fluorescent in situ hybridization (FISH). In congruence with some previous literature (*28*), Penk-

SPNs were found to densely project to ventral pallidum (VP) and less densely to lateral

hypothalamus (LH) and these circuits had minimal collateralization (13.3%) (**Extended Data**

Fig. 7d-f). It was also found that Pdyn-SPNs densely project to LH and sparsely to VTA with

minimal collateralization (7.3%) (**Extended Data Fig. 7a-c**), as has been previously reported of

D1R expressing SPNs (*29*).

To determine if these major SPN efferent circuits were selectively recruited to particular

reward preference conditions and not others, we conducted simultaneous dual-site fiber

photometry recordings while animals consumed the 4 reward preference conditions (0%, 3%,

10% or 30% sucrose). Penk-cre and Pdyn-cre mice were first injected with AAV-DJ-Ef1α-DIO-

GCaMP6s in the mNAcSh and then implanted 400 um diameter fiber optic cannulas into the VP

and LH (Penk-cre) or LH and VTA (Pdyn-cre) (**Fig. 4e**). Simultaneous *in vivo* recording of

calcium dynamics at the axon terminals of two mNAcSh efferents allowed us to make direct

comparisons of reward preference-related activity within-subject. Photometry animals were

tested in a head-restrained behavior paradigm identical to the 2-photon calcium imaging

experiments (see **Fig. 1**) (**Fig. 4a-b**).

For the Penk-cre mice, both the mNAcSh-VP^{Penk} and mNAcSh-LH^{Penk} circuits had

reward-excitation to all of the 4 sucrose preference conditions. However, the magnitude of

mNAcSh-VP^{Penk} reward-excitation was significantly greater for the unpreferred reward condition

than the preferred reward conditions (**Fig. 4f-g**). This greater excitation to the unpreferred

condition directly parallels the preference selectivity of the Penk-Unpreferred cluster we

identified during 2-photon imaging experiments (see **Fig. 1l**). In contrast, mNAcSh-LH^{Pdyn} was

not significantly selective for particular reward preference conditions (**Fig. 4l-m**). The non-

selective reward-excitation of mNAcSh-LH^{Pdyn} parallels the less selective reward-excitation of

the Pdyn-SPN clusters from 2-photon imaging experiments (see **Fig. 1m-n**). Reward preference

selectivity of some mNAcSh efferents and not others suggests that the heterogeneous reward

preference encoding observed in mNAcSh SPN populations is selectively propagated

downstream through particular efferents.

The dual-site fiber photometry animals were also tested in a 7 day Pavlovian

conditioning paradigm identical to the head-restrained paradigm used during 2-photon calcium

imaging (see **Fig. 2b-c**). mNAcSh-VP^{Penk} and mNAcSh-LH^{Pdyn} were found to be cue-excited

throughout Pavlovian conditioning in addition to having consistent 10% sucrose reward-

excitation (**Extended Data Fig. 8f-j, 8p-t**). Furthermore, the cue-excitation of mNAcSh-VP^{Penk}

persisted even during the 10 extinction trials where the reward was omitted (**Extended Data**

Fig. 8j). The magnitude of mNAcSh-VP^{Penk} and mNAcSh-LH^{Pdyn} reward-excitation was also

significantly lower when animals were expecting the cued 10% sucrose reward than when they

receive uncued 10% sucrose reward (**Extended Data Fig. 8h,r**). This mNAcSh-VP^{Penk} response

profile within the Pavlovian conditioning paradigm parallels the cue and reward-excitation that

was observed in Penk-SPNs cue and reward-excited clusters (see **Fig. 2k, Extended Data Fig.**

5). Given our calcium imaging-based characterization of 355 Penk-SPNs in mNAcSh, this

suggests that the mNAcSh-VP^{Penk} circuit is likely largely composed of the unpreferred reward

selective Penk-SPN subpopulation in mNAcSh. Thus, the separation of distinct reward

preference information streams in mNAcSh is continued through the mNAcSh-VP^{Penk} circuit,

which selectively encodes unpreferred rewards.

mNAcSh-VP^{Penk} was found to have greater reward-excitation to the unpreferred 30%

sucrose reward condition than the preferred conditions. However, it is still unclear whether

mNAcSh-VP^{Penk} reward-related activity causally contributes to consummatory behavior. To test

the role of mNAcSh-VP^{Penk} in casually driving consummatory behavior, we injected Penk-cre

mice with 300 nl of AAV5-EF1α-DIO-parapinopsin(PPO)-Venus (*30*) or AAV5-EF1α-DIO-ChR2-

eYFP bilaterally into mNAcSh and implanted a 200 um diameter fiber optic bilaterally into VP

(**Fig. 5e,i**). After recovering from surgery, animals were water-restricted and trained to lick 10%

sucrose reward while reversibly immobilized and head-restrained (**Fig. 5a**). After 4 days of lick

training, all animals displayed consistent lick behavior in response to variable interval sucrose
presentation and were then put through 2 consecutive test days.

Each test day involved one 30 minute, 20 trial session in which the animals were
presented with either 10% or 30% sucrose reward. The order of sucrose condition days was
pseudorandomized and counterbalanced across animals. For consistency with the imaging and
dual site photometry behavior paradigms, animals were presented with 5 droplets of sucrose
reward per trial with a total aggregate volume of 10 ul and 1 second between droplet delivery
(**Fig. 5b**). Animals immediately licked the presented sucrose reward, with an average licking
bout durations of 7-8 seconds (**Fig. 5c**). On a pseudorandom 50% of the trials within a session,
time-locked optogenetic manipulation of mNAcSh-VP^{Penk} activity would be paired with reward
delivery, via 4 seconds of 10 Hz blue light delivery (**Fig. 5b**). 10Hz optogenetic stimulation
frequency was chosen because it is physiologically relevant level of neural activity for SPNs
during phasic reward-related activity (*12*, *13*). Any differences in consummatory licking between
no laser and laser trials were recorded via a contact licometer with millisecond precision (**Fig.**
5d) and all differences were evaluated within subject to account for the baseline variability in lick
rate between animals. To control for possible behavioral effects from the delivery of the blue
light itself, cre- littermate controls were also injected, implanted, and tested in identical
conditions.

Time-locked photoinhibition (PPO-injected mice) of mNAcSh-VP^{Penk} during consumption
significantly reduced the reward epoch lick rate regardless of whether animals were consuming
10% or 30% sucrose reward (**Fig. 5g-h**). This demonstrates that mNAcSh-VP^{Penk} reward-
excitation is required for normal consummatory licking. Since mNAcSh Penk-SPNs specifically
encode reward preference information and mNAcSh-VP^{Penk} is reward-excited, the reduction in
consummatory licking during mNAcSh-VP^{Penk} photoinhibition is likely due to complete disruption
of reward information in this circuit.

Time-locked photoexcitation of mNAcSh-VP^{Penk} during consumption had an effect on

consummatory licking that was reward-dependent. Time-locked photoexcitation of mNAcSh-

VP^{Penk} reduced the consummatory licking of preferred 10% sucrose reward (**Fig. 5k**) but did not

significantly change the consumption of unpreferred 30% sucrose reward (**Fig. 5l**). The fact that

mNAcSh-VP^{Penk} already has greater endogenous activity during the consumption of 30%

sucrose reward (see **Fig. 4**) suggests that exogenous photoexcitation is not changing the

activity of this circuit as much during consumption of unpreferred reward. While in the preferred

reward condition, mNAcSh-VP^{Penk} has less endogenous activity and 10 Hz exogenous

photoexcitation would more profoundly changes the activity of the circuit.

Notably, mNAcSh-VP^{Penk} exogenous photoexcitation is not disrupting the motor action of

licking itself, as no disruption of licking behavior was observed during the consumption of 30%

sucrose reward (**Fig. 5l**). To isolate any possible effects from motor action impairment, a set of

additional experiments were designed in which mNAcSh-VP^{Penk} was photoexcited or

photoinhibited as animals freely moved through a behavioral arena in red light conditions

(**Extended Data Fig. 11h-i**). Movement velocity was not changed by mNAcSh-VP^{Penk}

photoinhibition (**Extended Data Fig. 11k**), nor by mNAcSh-VP^{Penk} photoexcitation (**Extended**

Data Fig. 11m), further suggesting that manipulation of this circuit during headfixed licking

behavior was influencing the processing of reward information and not disrupting the motor

execution of licking.

We next posited that if mNAcSh-VP^{Penk} is causally driving a low positive reward

preference, then photoexcitation of this circuit alone should be inherently rewarding. To test this

hypothesis, we again tested mice in our brief access 2-bottle choice paradigm (see **Extended**

Data Fig. 1). mNAcSh-VP$^{Penk:ChR2}$ expressing animals were trained to lick 10% sucrose reward

from two sucrose spouts within a behavioral chamber (**Fig. 5m**). After training the mice to

approach and lick the sucrose spouts for 5 days, we put the animals through 6 consecutive test

days. Test days 1, 3, 5 were 'reset' days to neutralize any left versus right biases that resulted

from other test conditions as described before (see **Extended Data Fig. 1**). On test day 2,

animals were given brief access to both a 10% and a 30% sucrose sipper. In accordance with

our earlier results (see **Extended Data Fig. 1**), mNAcSh-VP$^{Penk:ChR2}$ animals showed a

preference for 10% sucrose over 30% sucrose (**Fig. 5n**). On test days 4 and 6, the animals

were again tested for their preference between two sippers. However, one of the two sippers

contained no liquid and licking it triggered 2 seconds of 10 Hz closed loop optogenetic

photoexcitation (**Fig. 5m**). This stimulation duration and frequency was chosen to reasonably

recapitulate mNAcSh-VP^{enk} activity during 30% sucrose reward licking. The other sipper that the

animals could choose instead of the self-stimulation sipper contained either 10% or 30%

sucrose as before. The order of test days in which animals got the 10% or 30% alternative

sipper and the sipper location were pseudorandomized and counterbalanced across animals.

This experiment was designed to test if animals find mNAcSh-VP^{Penk} photoexcitation inherently

rewarding and how the inherent reinforcement of mNAcSh-VP^{Penk} photoexcitation compares to

the 10% and 30% sucrose conditions. It was found that mNAcSh-VP$^{Penk:ChR2}$ animals licked the

empty self-stimulation sipper significantly more than cre- controls, demonstrating that mNAcSh-

VP^{Penk} photoexcitation is inherently reinforcing. Furthermore, cre+ animals still licked the 10%

sucrose sipper significantly more than the self-stimulation sipper (**Fig. 5o**) but had no significant

preference between the 30% sucrose and the self-stimulation sippers (**Fig. 5p**). This allows us

to assess the relative preference of exogenous mNAcSh-VP^{Penk} excitation and to observe that it

is comparable to that of 30% sucrose reward, but measurably lower than the 10% sucrose

reward.

Overall, these complementary optogenetics experiments demonstrate that mNAcSh-

VP^{Penk} causally drives a low positive reward preference. This leads us to conclude that not only

is mNAcSh-VP^{Penk} endogenously recruited during the consumption of unpreferred rewards, but

that this endogenous activity during reward causally contributes to driving perceived reward
preference and reward consumption.

Here we report that distinct mNAcSh SPN ensembles encode relative reward
preference. Separate subpopulations of mNAcSh SPNs are recruited during the consumption of
preferred rewards and unpreferred rewards. We find that this stratification of reward preference
information in mNAcSh is then propagated through efferent projections, with unpreferred reward
signals relayed to VP by Penk-SPNs. Optogenetic activation of the mNAcSh-VP^{Penk} circuit was
found to cause no change in consummatory behavior when the animals were consuming
unpreferred reward but caused reductions in the consumption of preferred reward. Self-
stimulation experiments further revealed that mNAcSh-VP^{Penk} activation itself was inherently
rewarding and elicited consummatory behavior approximately equivalent to the lower preference
natural reward. These findings present a detailed mechanism for how mNAcSh integrates
incoming reward-related information, represents reward information on the basis of relative
preference and coordinates the appropriately motivated consummatory behavior.

It remains unresolved how much separation of reward identity, their subsequent
preference and the underlying relative value attributed to these rewards is present in brain
regions upstream of accumbens and which afferents are supplying this information. There is
recent evidence that neurons in VP encode reward value and project to NAc (*7*, *8*).
Furthermore, reward value signals in VP may even be present earlier than in NAc (*6*). These
previous findings and our current results suggest that some reward value information may be
originally transmitted from VP to mNAcSh, where it is further amplified and propagated.

In addition to the evidence for separation of reward information upstream of mNAcSh,
stratification of this information may be further enhanced locally due to interconnectivity between
SPNs in mNAcSh. Previous studies show a high degree of interconnectivity between nearby
SPNs in NAc (*31*, *32*). The divergence in SPN recruitment on the basis of reward preference

may be further stratified due to local inhibition of preferred reward SPNs by unpreferred reward

SPNs when they are selectively activated.

The observation that reward-inhibited SPNs are anterior to other cue and reward-excited

SPN clusters (see **Fig. 3**) is consistent with recent studies showing that incoming glutamatergic

innervation from basolateral amygdala, ventral hippocampus and medial thalamus to anterior

NAc is decreased during reward consumption and glutamatergic innervation to posterior NAc is

increased during reward consumption (*33*). This finding strongly suggests that differences in

glutamatergic innervation in NAc along the anterior-posterior axis result in differences in SPN

reward processing and that the anterior-posterior division is a continuum of individual neurons

rather than a strict boundary. Divergence in the reward-related activity of NAc populations has

also previously been observed across the mediolateral and dorsoventral axes (*11*, *34*).

Reward and cue-related signaling in NAc core has been found to be largely mediated by

mesolimbic dopamine release (*17*, *35*). However, it remains unclear how the differences in

dopaminergic innervation of NAc core and mNAcSh contribute to differences in reward

processing. Our current results demonstrate that many Penk-SPNs, that largely co-express

inhibitory D2Rs (*2*, *36*), are excited during reward and reward-predictive cues (see **Fig. 1-2**).

This may be attributable to unique dopamine receptor expression in mNAcSh, where D3-type

dopamine receptors are highly enriched (*37*, *38*). However, we also find that mNAcSh SPNs are

cue-excited but do not increase their cue-excitation across training days, and thus, do not

encode reward expectation as is canonically observed in NAc core (*13*, *15*). DA is also likely not

released as robustly to reward-predictive cues in mNAcSh as it is in NAc core (*18*, *39*). Recent

evidence also demonstrates that evoked DA release in mNAcSh does not reinforce

simultaneously presented cues (*17*). Altogether, these converging lines of research and the

present study suggest that DA is not a major mechanism underlying cue information processing

and cue reinforcement in mNAcSh, like it is in NAc core.

Mu opioid receptor (MOPR) activation in the basal forebrain has been thoroughly

characterized for its localized robust induction of reward 'liking' and 'disliking'. In particular,

MOPR activation in anterior VP suppresses reward 'liking' and consumption while MOPR

activation in posterior VP enhances reward 'liking' and consumption (*40, 41*). Penk-SPNs in

mNAcSh likely release enkephalin, which is a high affinity endogenous ligand for MOPR (*42*).

Here we find that mNAcSh Penk-SPNs densely project to VP and are activated by lower

preference rewards. Enkephalinergic activation of MOPR from Penk-SPNs may be a potential

mechanism by which reward preference signals are transduced in VP. This potential opioidergic

mechanism is beyond the scope of the present study but presents an interesting avenue for

further investigation.

Although future studies are necessary to understand the neurotransmitter and circuit-

level mechanisms underlying reward identity, preference and value computation in the NAc-VP

reward system, this work provides critical insight that mNAcSh robustly represents distinct

reward identities, their relative preference and transmits this information through a pallidal

efferent. This critical neural encoding mechanism in NAc plays a central role in how the brain

integrates incoming reward information and uses it to drive motivated and consummatory

behavior.

References

1.   F. Mannella, K. Gurney, G. Baldassarre, The nucleus accumbens as a nexus between
411   values and goals in goal-directed behavior: a review and a new hypothesis. *Frontiers in*
412   *Behavioral Neuroscience*. **7** (2013), doi:10.3389/fnbeh.2013.00135.

413   2.   D. C. Castro, M. R. Bruchas, A Motivational and Neuropeptidergic Hub: Anatomical and
414   Functional Diversity within the Nucleus Accumbens Shell. *Neuron*. **102**, 529–552 (2019).

415   3.   J. P. Britt, F. Benaliouad, R. A. McDevitt, G. D. Stuber, R. A. Wise, A. Bonci, Synaptic and
416   behavioral profile of multiple glutamatergic inputs to the nucleus accumbens. *Neuron*. **76**,
417   790–803 (2012).

4.   G. D. Stuber, D. R. Sparta, A. M. Stamatakis, W. A. van Leeuwen, J. E. Hardjoprajitno, S.
Cho, K. M. Tye, K. A. Kempadoo, F. Zhang, K. Deisseroth, A. Bonci, Excitatory
transmission from the amygdala to nucleus accumbens facilitates reward seeking. *Nature*.
475, 377–380 (2011).

5.   J. M. Otis, V. M. K. Namboodiri, A. M. Matan, E. S. Voets, E. P. Mohorn, O. Kosyk, J. A.
McHenry, J. E. Robinson, S. L. Resendez, M. A. Rossi, G. D. Stuber, Prefrontal cortex
output circuits guide reward seeking through divergent cue encoding. *Nature*. **543**, 103–
107 (2017).

6.   D. Ottenheimer, J. M. Richard, P. H. Janak, Ventral pallidum encodes relative reward value
earlier and more robustly than nucleus accumbens. *Nat Commun*. **9**, 4350 (2018).

7.   J. M. Richard, F. Ambroggi, P. H. Janak, H. L. Fields, Ventral Pallidum Neurons Encode
Incentive Value and Promote Cue-Elicited Instrumental Actions. *Neuron*. **90**, 1165–1173
(2016).

8.   Y. M. Vachez, J. R. Tooley, K. Abiraman, B. Matikainen-Ankney, E. Casey, T. Earnest, L.
M. Ramos, H. Silberberg, E. Godynyuk, O. Uddin, L. Marconi, C. E. Le Pichon, M. C.
Creed, Ventral arkypallidal neurons inhibit accumbal firing to promote reward consumption.
Nat Neurosci. **24**, 379–390 (2021).

9.   W. Schultz, Dopamine reward prediction-error signalling: a two-component response. *Nat
Rev Neurosci*. **17**, 183–195 (2016).

10.   M. J. Besson, A. M. Graybiel, B. Quinn, Co-expression of neuropeptides in the cat's
striatum: an immunohistochemical study of substance P, dynorphin B and enkephalin.
Neuroscience. **39**, 33–58 (1990).

11.   R. Al-Hasani, J. G. McCall, G. Shin, A. M. Gomez, G. P. Schmitz, J. M. Bernardi, C.-O.
Pyo, S. I. Park, C. M. Marcinkiewcz, N. A. Crowley, M. J. Krashes, B. B. Lowell, T. L. Kash,
J. A. Rogers, M. R. Bruchas, Distinct Subpopulations of Nucleus Accumbens Dynorphin
Neurons Drive Aversion and Reward. *Neuron*. **87**, 1063–1077 (2015).

12.   S. A. Taha, Encoding of Palatability and Appetitive Behaviors by Distinct Neuronal
Populations in the Nucleus Accumbens. *Journal of Neuroscience*. **25**, 1193–1202 (2005).

13.   M. F. Roitman, R. A. Wheeler, R. M. Carelli, Nucleus accumbens neurons are innately
tuned for rewarding and aversive taste stimuli, encode their predictors, and are linked to
motor output. *Neuron*. **45**, 587–597 (2005).

14.   H. J. Groenewegen, C. I. Wright, A. V. Beijer, P. Voorn, Convergence and segregation of
ventral striatal inputs and outputs. *Ann N Y Acad Sci*. **877**, 49–63 (1999).

15.   J. J. Day, R. A. Wheeler, M. F. Roitman, R. M. Carelli, Nucleus accumbens neurons
encode Pavlovian approach behaviors: evidence from an autoshaping paradigm. *European
Journal of Neuroscience*. **23**, 1341–1351 (2006).

16.   F. Ambroggi, A. Ghazizadeh, S. M. Nicola, H. L. Fields, Roles of nucleus accumbens core
and shell in incentive-cue responding and behavioral inhibition. *J Neurosci*. **31**, 6820–6830
(2011).

17.   B. T. Saunders, J. M. Richard, E. B. Margolis, P. H. Janak, Dopamine neurons create
458   Pavlovian conditioned stimuli with circuit-defined motivational properties. *Nat Neurosci*. **21**,
459   1072–1083 (2018).

460   18.   J. W. de Jong, S. A. Afjei, I. Pollak Dorocic, J. R. Peck, C. Liu, C. K. Kim, L. Tian, K.
461   Deisseroth, S. Lammel, A Neural Circuit Mechanism for Encoding Aversive Stimuli in the
462   Mesolimbic Dopamine System. *Neuron*. **101**, 133-151.e7 (2019).

463   19.   S. L. Resendez, J. H. Jennings, R. L. Ung, V. M. K. Namboodiri, Z. C. Zhou, J. M. Otis, H.
464   Nomura, J. A. McHenry, O. Kosyk, G. D. Stuber, Visualization of cortical, subcortical and
465   deep brain neural circuit dynamics during naturalistic mammalian behavior with head-
466   mounted microscopes and chronically implanted lenses. *Nat Protoc*. **11**, 566–597 (2016).

467   20.   M. A. Rossi, M. L. Basiri, J. A. McHenry, O. Kosyk, J. M. Otis, H. E. van den Munkhof, J.
468   Bryois, C. Hübel, G. Breen, W. Guo, C. M. Bulik, P. F. Sullivan, G. D. Stuber, Obesity
469   remodels activity and transcriptional state of a lateral hypothalamic brake on feeding.
470   *Science*. **364**, 1271–1274 (2019).

471   21.   V. M. K. Namboodiri, J. M. Otis, K. van Heeswijk, E. S. Voets, R. A. Alghorazi, J.
472   Rodriguez-Romaguera, S. Mihalas, G. D. Stuber, Single-cell activity tracking reveals that
473   orbitofrontal neurons acquire and maintain a long-term memory to guide behavioral
474   adaptation. *Nat Neurosci*. **22**, 1110–1121 (2019).

475   22.   S. Pothion, J.-C. Bizot, F. Trovero, C. Belzung, Strain differences in sucrose preference
476   and in the consequences of unpredictable chronic mild stress. *Behav Brain Res*. **155**, 135–
477   146 (2004).

478   23.   D. L. Wallace, V. Vialou, L. Rios, T. L. Carle-Florence, S. Chakravarty, A. Kumar, D. L.
479   Graham, T. A. Green, A. Kirk, S. D. Iñiguez, L. I. Perrotti, M. Barrot, R. J. DiLeone, E. J.
480   Nestler, C. A. Bolaños-Guzmán, The influence of DeltaFosB in the nucleus accumbens on
481   natural reward-related behavior. *J Neurosci*. **28**, 10272–10277 (2008).

482   24.   D. J. Ottenheimer, K. Wang, X. Tong, K. M. Fraser, J. M. Richard, P. H. Janak, Reward
483   activity in ventral pallidum tracks satiety-sensitive preference and drives choice behavior.
484   *Sci Adv*. **6** (2020), doi:10.1126/sciadv.abc9321.

485   25.   J. D. Davis, The effectiveness of some sugars in stimulating licking behavior in the rat.
486   *Physiol Behav*. **11**, 39–45 (1973).

487   26.   W. Schultz, P. Dayan, P. R. Montague, A neural substrate of prediction and reward.
488   *Science*. **275**, 1593–1599 (1997).

489   27.   K. C. Berridge, T. E. Robinson, What is the role of dopamine in reward: hedonic impact,
490   reward learning, or incentive salience? *Brain Res Brain Res Rev*. **28**, 309–369 (1998).

491   28.   M. Creed, N. R. Ntamati, R. Chandra, M. K. Lobo, C. Lüscher, Convergence of Reinforcing
492   and Anhedonic Cocaine Effects in the Ventral Pallidum. *Neuron*. **92**, 214–226 (2016).

493   29.   S. Thoeni, M. Loureiro, E. C. O'Connor, C. Lüscher, Depression of Accumbal to Lateral
494   Hypothalamic Synapses Gates Overeating. *Neuron*. **107**, 158-172.e4 (2020).

495   30.   B. A. Copits, R. Gowrishankar, P. R. O'Neill, J.-N. Li, K. S. Girven, J. J. Yoo, X. Meshik, K.
496   E. Parker, S. M. Spangler, A. J. Elerding, B. J. Brown, S. E. Shirley, K. K. L. Ma, A. M.
497   Vasquez, M. C. Stander, V. Kalyanaraman, S. K. Vogt, V. K. Samineni, T. Patriarchi, L.
498   Tian, N. Gautam, R. K. Sunahara, R. W. Gereau, M. R. Bruchas, A photoswitchable
499   GPCR-based opsin for presynaptic inhibition. *Neuron*. **109**, 1791-1809.e11 (2021).

500   31.   H. Planert, S. N. Szydlowski, J. J. J. Hjorth, S. Grillner, G. Silberberg, Dynamics of
501   synaptic transmission between fast-spiking interneurons and striatal projection neurons of
502   the direct and indirect pathways. *J Neurosci*. **30**, 3499–3507 (2010).

503   32.   H. A. Tejeda, J. Wu, A. R. Kornspun, M. Pignatelli, V. Kashtelyan, M. J. Krashes, B. B.
504   Lowell, W. A. Carlezon, A. Bonci, Pathway- and Cell-Specific Kappa-Opioid Receptor
505   Modulation of Excitation-Inhibition Balance Differentially Gates D1 and D2 Accumbens
506   Neuron Activity. *Neuron*. **93**, 147–163 (2017).

507   33.   S. J. Reed, C. K. Lafferty, J. A. Mendoza, A. K. Yang, T. J. Davidson, L. Grosenick, K.
508   Deisseroth, J. P. Britt, Coordinated Reductions in Excitatory Input to the Nucleus
509   Accumbens Underlie Food Consumption. *Neuron*. **99**, 1260-1273.e4 (2018).

510   34.   R. Al-Hasani, R. Gowrishankar, G. P. Schmitz, C. E. Pedersen, D. J. Marcus, S. E. Shirley,
511   T. E. Hobbs, A. J. Elerding, S. J. Renaud, M. Jing, Y. Li, V. A. Alvarez, J. C. Lemos, M. R.
512   Bruchas, Ventral tegmental area GABAergic inhibition of cholinergic interneurons in the
513   ventral nucleus accumbens shell promotes reward reinforcement. *Nat Neurosci* (2021),
514   doi:10.1038/s41593-021-00898-2.

515   35.   P. E. M. Phillips, G. D. Stuber, M. L. A. V. Heien, R. M. Wightman, R. M. Carelli,
516   Subsecond dopamine release promotes cocaine seeking. *Nature*. **422**, 614–618 (2003).

517   36.   J. Bertran-Gonzalez, C. Bosch, M. Maroteaux, M. Matamales, D. Hervé, E. Valjent, J.-A.
518   Girault, Opposing patterns of signaling activation in dopamine D1 and D2 receptor-
519   expressing striatal neurons in response to cocaine and haloperidol. *J Neurosci*. **28**, 5671–
520   5685 (2008).

521   37.   E. M. Pich, G. Collo, Pharmacological targeting of dopamine D3 receptors: Possible clinical
522   applications of selective drugs. *Eur Neuropsychopharmacol*. **25**, 1437–1447 (2015).

523   38.   J. D. Berke, What does dopamine mean? *Nat Neurosci*. **21**, 787–793 (2018).

524   39.   M. P. Saddoris, F. Cacciapaglia, R. M. Wightman, R. M. Carelli, Differential Dopamine
525   Release Dynamics in the Nucleus Accumbens Core and Shell Reveal Complementary
526   Signals for Error Prediction and Incentive Motivation. *J. Neurosci*. **35**, 11572–11582
527   (2015).

528   40.   K. S. Smith, K. C. Berridge, Opioid Limbic Circuit for Reward: Interaction between Hedonic
529   Hotspots of Nucleus Accumbens and Ventral Pallidum. *Journal of Neuroscience*. **27**,
530   1594–1605 (2007).

531   41.   D. C. Castro, S. L. Cole, K. C. Berridge, Lateral hypothalamus, nucleus accumbens, and
532   ventral pallidum roles in eating and hunger: interactions between homeostatic and reward
533   circuitry. *Frontiers in Systems Neuroscience*. **9**, 90 (2015).

534   42.   B. A. Jordan, S. Cvejic, L. A. Devi, Opioids and their complicated receptor complexes.
535   *Neuropsychopharmacology.* **23**, S5–S18 (2000).

536   43.   A. François, S. A. Low, E. I. Sypek, A. J. Christensen, C. Sotoudeh, K. T. Beier, C.
537   Ramakrishnan, K. D. Ritola, R. Sharif-Naeini, K. Deisseroth, S. L. Delp, R. C. Malenka, L.
538   Luo, A. W. Hantman, G. Scherrer, A Brainstem-Spinal Cord Inhibitory Circuit for
539   Mechanical Pain Modulation by GABA and Enkephalins. *Neuron.* **93**, 822-839.e6 (2017).

540   44.   K. E. Parker, C. E. Pedersen, A. M. Gomez, S. M. Spangler, M. C. Walicki, S. Y. Feng, S.
541   L. Stewart, J. M. Otis, R. Al-Hasani, J. G. McCall, K. Sakers, D. L. Bhatti, B. A. Copits, R.
542   W. Gereau, T. Jhou, T. J. Kash, J. D. Dougherty, G. D. Stuber, M. R. Bruchas, A
543   Paranigral VTA Nociceptin Circuit that Constrains Motivation for Reward. *Cell.* **178**, 653-
544   671.e19 (2019).

545   45.   P. Kaifosh, J. D. Zaremba, N. B. Danielson, A. Losonczy, SIMA: Python software for
546   analysis of dynamic fluorescence imaging data. *Front. Neuroinform.* **8** (2014),
547   doi:10.3389/fninf.2014.00080.

548   46.   M. Pachitariu, C. Stringer, M. Dipoppa, S. Schröder, L. F. Rossi, H. Dalgleish, M.
549   Carandini, K. D. Harris, "Suite2p: beyond 10,000 neurons with standard two-photon
550   microscopy" (preprint, Neuroscience, 2016), , doi:10.1101/061507.

Figure Legends

Fig. 1. Distinct subpopulations of mNAcSh SPNs are recruited in response to rewards of
different preference.

(a) Coronal section cartoon displaying target brain region, cell-types and lens implant location.

(b) Schematic of 2-photon calcium imaging through endoscopic lens during consumption of
variable sucrose reward.

(c) Time-course of sucrose reward delivery for a given trial. Droplets were either 0%, 3%, 10%
or 30% sucrose solution in water.

(d) Representative lick raster depicting individual licks that were detected by contact licometer
during behavior and imaging session.

(e) Mean lick probability during reward epoch (0-8 seconds from reward onset) for each sucrose
reward condition (n = 18 animals; one-way ANOVA: p = 0.9275).

(f) Representative coronal section showing Penk-SPNs expressing GCaMP6s underneath the
implanted endoscopic GRIN lens in mNAcSh (scale bar = 200 um).

(G and H) Fluorescence time series of individual SPNs that were tracked from (g) day 1 session
through (h) day 4 session.

(i) SPN clusters that were not consistently recruited during any of the reward conditions
(averaged response to 20 trials of each reward condition).

(j) SPN clusters that were consistently transiently inhibited during all reward conditions.

(k) SPN clusters that were consistently transiently excited during the 0%, 3% and 10% conditions, but less for the 30% condition.

(l) Penk-SPN cluster that was consistently transiently excited during 30% condition, moderately for the 0% condition and less for the 3% and 10% condition.

(m) Pdyn-SPN cluster that was consistently transiently excited during the 3%, 10% and 30% sucrose conditions but less for the 0% (water) condition.

(n) Pdyn-SPN cluster that was consistently transiently excited for all reward conditions.

(o) t-sne plot of variance in activity of Penk-SPN clusters during all reward conditions. Each dot represents an individual tracked neuron that composes the cluster.

(p) Fraction of all tracked Penk-SPN neurons that belong to each functional cluster.

(q) t-sne plot of variance in activity of Pdyn-SPN clusters during all reward conditions. Each dot represents an individual tracked neuron that composes the cluster.

(r) fraction of all tracked Pdyn-SPN neurons that belong to each functional cluster.

(s) Cartoon depicting multiclass linear Support Vector Classifier implementation (features = neurons, observations = mean activity during given trial, target = class of reward condition).

(t) Mean of SVC coefficients for all Penk-SPNs (SVC features) of each functional cluster.

(u) Mean of SVC coefficients for all Pdyn-SPNs (SVC features) of each functional cluster.

(v) Linear SVC classification accuracy for every animal using mean SPN activity during reward epoch (0-8s from reward onset) and during control epoch (12-4s before reward onset) (n = 7 Penk-cre and 11 Pdyn-cre animals).

Fig. 2. Most reward-excited mNAcSh SPNS are cue-excited, but do not reflect reward expectation

(a) Coronal section cartoon displaying target brain region, cell-types and lens implant location.

(b) Schematic of 2-photon calcium imaging through endoscopic lens during cue-reward Pavlovian conditioning.

(c) Time-course of auditory cue and subsequent sucrose reward delivery for a given trial.

(d and e) Representative lick raster depicting individual licks that were detected by contact licometer during cue and reward when animals were cue-naïve (d) and after conditioning (e).

(f and g) Fluorescence time series of individual SPNs that were tracked from Pavlovian (f) day 1 session through (g) day 7 session.

(h) Mean lick probability during cue epoch (0-5 seconds from cue-onset) for each day of Pavlovian conditioning and extinction (n = 16 animals; One-way ANOVA: Day 1 vs Day 7, *p = 0.020, Day 7 vs Extinction, **p = 0.002).

(i) SPN clusters that were not consistently recruited during cue or reward throughout 7 days of Pavlovian conditioning (averaged response to 20 trials of each reward condition).

(j) SPN clusters that were consistently transiently inhibited during reward throughout 7 days of Pavlovian conditioning.

(k) SPN clusters that were consistently transiently excited during cue and reward throughout 7 days of Pavlovian conditioning and extinction.

(l) Pdyn-SPN cluster that was consistently transiently excited during reward but not cue throughout 7 days of Pavlovian conditioning.

(m) Fraction of all tracked Pdyn-SPN neurons that belong to each Pavlovian functional cluster.

(n) Fraction of all tracked Penk-SPN neurons that belong to each Pavlovian functional cluster.

(o) Mean cue response (0-5 seconds from cue-onset) of all cue-excited SPNs during each day of Pavlovian conditioning and extinction (n = 7 Penk-cre animals, 155 neurons; Pearson's correlation: $p = 0.105$; n = 9 Pdyn-cre animals, 104 neurons; Pearson's correlation: $p = 0.083$).

(p) Correlation between mean cue response of all cue-excited cluster SPNs and mean cue epoch lick probability for every Pavlovian conditioning trial. Blue and red lines capture linear fit between cue response and cue licking for each cell-type. (Pearson's correlation: ***$p < 0.001$).

(q) Mean lick probability during cue epoch (0-5 seconds from cue-onset) for each trial within a session (Days 1, 3, 5 combined) (n = 16 animals; Pearson's correlation: $p = 0.288$).

(r) Mean cue response of all cue-excited cluster SPNs during each trial within a session (Days 1, 3, 5 combined) (n = 7 Penk-cre animals; Pearson's correlation: $p = 0.254$; n = 9 Pdyn-cre animals; Pearson's correlation: ***$p < 0.001$).

Fig. 3. Reward-inhibited SPN clusters are anatomically anterior to other SPN clusters

(a) Fraction of all tracked Penk-SPNs and Pdyn-SPNs that are part of reward-inhibited clusters.

(b) Mean activity of SPN clusters that were consistently transiently inhibited during all variable sucrose reward conditions.

(c) Position of individual Penk-SPNs in focal plane during in vivo 2-photon calcium imaging sessions; arranged by variable sucrose response cluster (n = 4 clusters, 7 animals; one-way ANOVA: ***$p < 0.001$).

(d) Position of individual Pdyn-SPNs in focal plane during in vivo 2-photon calcium imaging sessions; arranged by variable sucrose response cluster (n = 5 clusters, 11 animals; one-way ANOVA: ***$p < 0.001$).

(e) Relative anatomical location of SPN variable sucrose response clusters.

(f) Fraction of all tracked Penk-SPNs and Pdyn-SPNs that are part of Pavlovian reward-inhibited clusters.

(g) Mean activity of SPN clusters that were consistently transiently inhibited during reward throughout 7 days of Pavlovian conditioning.

(h) Position of individual Penk-SPNs in focal plane during in vivo 2-photon calcium imaging sessions; arranged by Pavlovian conditioning response cluster (n = 3 clusters, 7 animals; one-way ANOVA: ***$p < 0.001$).

(i) Position of individual Pdyn-SPNs in focal plane during in vivo 2-photon calcium imaging sessions; arranged by Pavlovian conditioning response cluster (n = 4 clusters, 9 animals; one-way ANOVA: **$p = 0.004$).

(j) Relative anatomical location of SPN Pavlovian conditioning response clusters.

Fig. 4. mNAcSh-VP^{Penk} parallels unpreferred reward selectivity and cue-excitation of Penk-SPN cluster

(a) Schematic of simultaneous two-site fiber photometry during consumption of variable sucrose
reward.

(b) Time-course of sucrose reward delivery for a given trial. Droplets were either 0%, 3%, 10%
or 30% sucrose in water.

(c) Representative lick raster depicting individual licks that were detected by contact licometer
during behavior and photometry session.

(d) Mean lick rate during reward epoch (0-8 seconds from reward onset) for each sucrose
reward condition (n = 15 animals; one-way ANOVA: p = 0.280).

(e) Sagittal section cartoon displaying target injection region, cell-type and fiber implant
locations.

(f) Mean mNAcSh-VP^{Penk} reward-response for every trial of each variable sucrose reward
condition.

(g) Mean mNAcSh-VP^{Penk} activity over reward response epoch (2-12 seconds from reward-
onset) for each condition (n = 9 animals, 190+ trials; one-way ANOVA: 30% vs 0%, *p = 0.044,
30% vs 10%, **p = 0.009).

(h) Representative coronal section showing Penk-SPN terminals in VP expressing GCaMP6s
underneath the fiber implant (scale bar = 200 um).

(i) Mean mNAcSh-LH^{Penk} reward-response for every trial of each variable sucrose reward
condition.

(j) Mean mNAcSh-LH^{Penk} activity over reward response epoch (2-12 seconds from reward-
onset) for each condition (n = 6 animals, 127+ trials; one-way ANOVA: p = 0.6167).

(k) Sagittal section cartoon displaying target injection region, cell-types and fiber implant
locations.

(l) Mean mNAcSh-LH^{Pdyn} reward-response for every trial of each variable sucrose condition.

(m) Mean mNAcSh-LH^{Pdyn} activity over reward response epoch (2-12 seconds from reward-
onset) for each condition (n = 6 animals, 123+ trials, one-way ANOVA: p = 0.550).

(n) Representative coronal section showing Pdyn-SPN terminals in LH expressing GCaMP6s
underneath the fiber implant (scale bar = 150 um).

(o) Mean mNAcSh-VTA^{Pdyn} reward-response for every trial of each variable sucrose condition.

(p) Mean mNAcSh-VTA^{Pdyn} activity over reward response epoch (2-12 seconds from reward-
onset) for each condition (n = 7 animals, 143+ trials, one-way ANOVA: 3% vs 10%, *p = 0.031).

Fig. 5. mNAcSh-VP^{Penk} causally drives low-positive reward preference

(a) Schematic of timelocked-optogenetic neural activity manipulation during consumption of
variable sucrose reward.

(b) Time-course of sucrose reward delivery for a given trial. 50% of trials also had 4 seconds of
10 Hz optogenetic stimulation concurrent with reward delivery.

(c) Lick rate distribution relative to reward-onset for all conditions. Comparison of lick rate
between conditions was restricted to the indicated reward epoch (0-8 seconds from reward-
onset).

(d) Representative lick raster displaying individual measured licks during trials without
stimulation and trials with stimulation.

(e) Coronal section cartoon displaying target injection region, cell-type and bilateral fiber implant
location.

(f) Representative coronal section showing Penk-SPN terminals in VP expressing PPO
underneath optic fiber implant (scale bar = 200 um).

(g) Within-session comparison of 10% sucrose consumption between trials with and without
reward-paired optogenetic inhibition (n = 7 treatment, 5 control animals; pairwise t-test: ***p <
0.001 , p = 0.418).

(h) Within-session comparison of 30% sucrose consumption between trials with and without
reward-paired optogenetic inhibition (n = 7 treatment, 5 control animals; pairwise t-test: **p =
0.003 , p = 0.619).

(i) Coronal section cartoon displaying target injection region, cell-type and bilateral fiber implant
location.

(j) Representative coronal section showing Penk-SPN terminals in VP expressing ChR2
underneath optic fiber implant (scale bar = 200 um).

(k) Within-session comparison of 10% sucrose consumption between trials with and without
reward-paired optogenetic excitation (n = 9 treatment, 6 control animals; pairwise t-test: **p =
0.003 , p = 0.437).

(l) Within-session comparison of 30% sucrose consumption between trials with and without
reward-paired optogenetic excitation (n = 9 treatment, 6 control animals; pairwise t-test: p =
0.057 , p = 0.327).

(m) Schematic of brief access 2-bottle choice task: animal makes choice between sucrose
reward sipper and lick-triggered 2 second, 10 Hz optogenetic self-stimulation.

(n) Preference between 10% sucrose reward sipper and 30% sucrose reward sipper (n = 15
animals; pairwise t-test: *p = 0.021).

(o) Preference between 10% sucrose reward and optogenetic self-stimulation (n = 9 treatment,
6 control animals; pairwise t-test: **p = 0.001 , **p = 0.004).

(p) Preference between 30% sucrose reward and optogenetic self-stimulation (n = 9 treatment,
6 control animals; pairwise t-test: p = 0.956 , **p = 0.002).

Methods

Animals

Adult C57BL/6 (18–35 g) male and female wildtype, prodynorphin-IRES-Cre (*dyn*-cre) and

preproenkephalin-IRES-Cre (*enk*-cre) mice were group housed, given access to food pellets

and water ad libitum, and maintained on a 12 hr:12 hr light:dark cycle. All mice were kept in a

sound-attenuated, isolated holding facility one week prior to surgery, post-surgery, and

throughout the duration of the behavioral assays to minimize stress. For cell-type specific

optogenetic experiments, we used Cre- cage and littermate controls. Unless otherwise noted,

animals had *ad libitum* access to food and water. The mice were bred at Washington University

in Saint Louis or the University of Washington. All animals were initially test naive, individually

assigned to specific experiments as described, and not involved with other experimental

procedures. Statistical comparisons did not detect any significant differences between male and

female mice and were therefore combined to complete final group sizes. All animals were

monitored for health status daily and before experimentation for the entirety of the study. All

procedures were approved by the Animal Care and Use Committee of Washington University,

Animal Care and Use Committee of the University of Washington and conformed to US National

Institutes of Health guidelines.

Tissue collection after behavioral experiments

Unless otherwise stated, animals were transcardially perfused with 0.1 M phosphate-buffered

saline (PBS) and then 40 mL 4% paraformaldehyde (PFA). Brains were dissected and post-

fixed in 4% PFA overnight and then transferred to 30% sucrose solution for cryoprotection.

Brains were sectioned at 30 mM on a microtome and stored in a 0.01M phosphate buffer at 4⁰C

prior to immunohistochemistry and tracing experiments. For behavioral cohorts, viral expression

and optical fiber placements were confirmed before inclusion in the presented datasets.

RNAscope Fluorescent *In Situ* Hybridization

Following rapid decapitation of wildtype mice, brains were rapidly frozen in 100mL -50°C

isopentane and stored at -80°C. Coronal sections corresponding to the site of interest or

injection plane used in the behavioral experiments were cut at 20uM at -20°C and thaw-

mounted onto SuperFrost Plus slides (Fisher). Slides were stored at -80°C until further

processing. Fluorescent in situ hybridization was performed according to the RNAscope 2.0

Fluorescent Multiplex Kit User Manual for Fresh Frozen Tissue (Advanced Cell Diagnostics,

Inc.). Briefly, sections were fixed in 4% PFA, dehydrated, and treated with pretreatment 4

protease solution. Sections were then incubated for target probes for mouse proenkephalin

(*Penk*, accession number NM_001002927.2, probe region 106 – 1332) or prodynorphin (*Pdyn*,

accession number NM_018863.3, probe region 33 - 700). All target probes consisted of 20 ZZ

oligonucleotides and were obtained from Advanced Cell Diagnostics. Following probe

hybridization, sections underwent a series of probe signal amplification steps followed by

incubation of fluorescently labeled robes designed to target the specific channel associated with

the probes. Slides were counterstained with DAPI, and coverslips were mounted with

Vectashield Hard Set mounting medium (Vector Laboratories). Images were obtained on an

Olympus Fluoview 3000 confocal microscope and analyzed with HALO software. To analyze the

images, each image was opened in the HALO software. DAPI positive cells were then

registered and used as markers for individual cells. An observer blind to the brain tissue origin

and probes used then counted the total number of probe-labeled cells for each channel

separately. A positive cell consisted of an area within the radius of a DAPI nuclear staining that

measured at least 10 total positive pixels for neurotransmitter probes. Two - three separate

slices from the NAc or were used for each animal and that total is presented in the data.

Stereotaxic Surgery

After mice were acclimated to the holding facility for at least seven days, the mice were

anaesthetized in an induction chamber (1%-4% isoflurane) and placed into a stereotaxic frame

(Kopf Instruments, model 1900) where they were mainlined at 1%-2% isoflurane. A small hole

was drilled in the skull above the target brain region. For mice receiving viral injections, a needle

syringe (Hamilton Company, 65458) was used to deliver the vector at a rate of 100 μL/min.

Animals used for 2-photon calcium imaging experiments were injected unilaterally with 500 uL

of AAVdj-ef1a-DIO-GCaMP6s into the mNAcSh (coordinates: AP +1.3, ML +0.5, DV -4.4 from

bregma), implanted with a GRIN lens (Inscopix, 7.3 mm length, 0.6 mm diameter) and had a 1 cm diameter headring horizontally adhered to the top of the skull. Animals used for fiber photometry experiments were injected unilaterally with 400 uL of AAVdj-ef1a-DIO-GCaMP6s into the mNAcSh (coordinates: AP +1.3, ML +0.5, DV -4.4 from bregma), implanted ipsilaterally with fiber optic cannulas (Doric Lenses, 400 um diameter, 6 mm length, 0.48 NA) in VP (coordinates: AP +0.3, ML +1.2, DV -4.7 from bregma), LH (coordinates: AP -1.3, ML +1.1, DV -4.7 from bregma) or VTA (coordinates: AP -3.3, ML +0.4, DV -4.4 from bregma), and had a 1 cm diameter headring horizontally adhered to the top of the skull. Animals used for retrotracing and FISH were injected unilaterally with 200 uL of green or red retrobeads (Lumafluor, Retrobeads IX) in VP (coordinates: AP +0.3, ML +1.2, DV -4.7 from bregma), LH (coordinates: AP -1.3, ML +1.1, DV -4.7 from bregma) or VTA (coordinates: AP -3.3, ML +0.4, DV -4.4 from bregma). Animals used for optogenetics experiments were injected bilaterally with 200 uL of AAV5-DIO-ChR2-EYFP or AAV5-DIO-parapinopsin-Venus into the mNAcSh (coordinates: AP +1.3, ML +/-0.5, DV -4.4 from bregma), implanted bilaterally with custom-built fiber optic cannulas (200 um diameter) in VP (coordinates: AP +0.3, ML +1.2, DV -4.7 from bregma) or LH (coordinates: AP -1.3, ML +1.1, DV -4.7 from bregma), and had a 1 cm diameter headring horizontally adhered to the top of the skull. All implants and headrings were secured to the skull using dental cement (Parkell, C&B Metabond). Animals were injected with slow-release Carprofen (5 mg/kg) prior to surgery to act as a post-surgery analgesic. Animals were monitored daily for 7 days following surgery.

Water Access Restriction prior to behavioral testing

Mice were allowed to recover from surgery for a minimum of 4 weeks prior to water restriction or behavioral testing. To adequately motivate consistent sucrose consumption in behavioral subjects, animals were water-restricted by giving them access to water only once per day as previously described (*5*, *20*, *21*). Starting 5 days prior to behavioral testing animals were given

1.0 ml of water in a 2x2cm paper tray in their home cage each day at approximately 5-6pm,

after behavioral testing. Animals readily consumed the water while supervised by the researcher

to ensure they consumed most of the presented amount. Animals were also weighed routinely

to ensure they stayed at 85-90% of their original body weight while under water-restriction. No

health problems related to dehydration arose throughout the water-restriction period of up to

~16 days. After the conclusion of a behavioral testing schedule, water-restriction was

immediately stopped and animals were given ad-libitum access to water in their home cages

again.

Brief Access Two Bottle Choice

To test the relative preference of the 4 variable sucrose reward conditions (0%, 3%, 10%, 30%

sucrose dissolved in water) while minimizing effects from satiety, we tested mice in an adapted

2-bottle choice assay where the animals were only given brief access (20 s) to sucrose sippers

with a variable intertrial interval (20-40s) between access periods. Brief access 2-bottle choice

assays were conducted within enclosed operant chambers, that had two retractable sucrose

sippers side-by-side (MedAssociates, MED-307A, ENV-352). Discrete licks on the metallic

sippers were recorded by a contact licometer system (MedAssociates, ENV-250). Prior to 2-

bottle choice test days, each animal was trained with free access (2 days) and variable access

(2 days) to both sippers containing 10% sucrose for 30 minute sessions. Each preference test

involved presentation of 2 of the 4 different sucrose conditions and each session had two

phases. In the first phase, animals were given access to either sipper in isolation so that they

could sample them independently. In the second phase, animals were simultaneously presented

with both sippers side-by-side and had to choose where to allocate their sipper access time.

The 6 possible pairings of the 4 sucrose reward conditions were presented in a

pseudorandomized and counterbalanced sequence to minimize effects from condition history

and right versus left sipper location bias. Preference test days were also buffered by 10%

versus 10% sucrose days to neutralize any left-versus-right biases that arose from test days.

As indicated in the Results, some sessions included a variation of this paradigm where one of

the sucrose sippers contained no liquid and instead triggered closed loop optogenetic

stimulation. Licking the empty sucrose spout would register a lick in the MED-PC program and

immediately trigger a TTL sent from the MED-307A to an Arduino Uno. The Arduino Uno

controlled a 470 nm light laser (4 mW intensity at end of patch cable, 10 Hz, 10 ms pulse width,

2 second in duration) that was optically coupled to the fiber implants on the animal's head.

Variable Sucrose Access during Reversible Head-fixation

After 4-6 weeks of recovery from surgery, animals were water restricted (see above) and trained

to sip 10% sucrose reward at variable intervals while reversibly immobilized and head-

restrained. Reversible head-fixation was conducted by scruffing the animals, backing them into

a 50 ml canonical tube and sliding the animals and tube into place in a custom headstage as

has been previously described (*20*, *21*). All animals displayed no signs of physical distress and

consistent licking behavior while immobilized after 4 days of lick training and were then tested in

30 minute, 20 trial sessions each day for 4 consecutive days. Intertrial interval was uniformly

distributed between 70 and 100 seconds. On each trial, animals were presented with 5 droplets

of sucrose reward with a total aggregate volume of 10 ul, with 1 second between droplet

delivery. Reward delivery was controlled by custom code on an Arduino Uno. Animals

immediately licked the presented sucrose reward, with an average licking bout duration of 7-8

seconds. In each session, animals consumed 1 of 4 possible sucrose reward conditions (0%,

3%, 10% or 30% sucrose in water), with the order of condition days pseudorandomized and

counterbalanced across animals. Discrete lick events were detected and recorded with <1 ms

time resolution using a contact licometer circuit as has been previously described (*20*, *21*). In

some cases, 15-17 FPS behavioral video of the animal's face was recorded alongside the

contact licometer or instead of the contact licometer. The presence of sucrose licking in a

particular frame was assessed manually or through automated scoring by a recurrent neural

network model (see below) that was extensively trained and validated using simultaneous

contact licometer information. The behavioral testing was conducted in dark or far-red light

conditions to remove contamination from visual stimuli. 2-photon calcium imaging data or

photometry data was acquired concurrently with this behavioral paradigm on days 1-4.

Pavlovian conditioning during Reversible Head-fixation

The same animals that were tested in the variable sucrose during reversible head-fixation

paradigm (see above) were then also trained to associate an auditory cue to subsequent

delivery of 10% sucrose reward across 7 days of Pavlovian conditioning during reversible head-

fixation. Animals were again tested in 30 minute, 20 trial sessions each day for 7 consecutive

days. Intertrial interval was uniformly distributed between 70 and 100 seconds. On each trial,

animals were presented with 3 seconds of a 4 kHz auditory cue. 2 seconds after cue

termination, 10% sucrose reward was delivered in a series of 5 droplets with total aggregate

volume of 10 ul, with 1 second between droplet delivery. Reward delivery and cues were

controlled by custom code on an Arduino Uno. The latter half of trials (last 10 out of 20 trials) on

Pavlovian day 7 were reward omission trials, where the cue was presented as usual, but the

expected sucrose reward was not delivered. Licking behavior was detected and recorded with

<1 ms time resolution using a contact licometer and/or by 15-17 FPS behavioral video. The

behavioral testing was conducted in dark or far-red light conditions to remove contamination

from visual stimuli. 2-photon calcium imaging data or photometry data was acquired

concurrently with this behavioral paradigm on Pavlovian training days 1, 3, 5 and 7.

Automated Lick Detection from Video using Recurrent Neural Network

In some reversible head-fixed behavior sessions, 15-17 FPS, 720p or 1080p behavioral video

was collected concurrently with or instead of the contact licometer data for lick detection. The

camera was positioned approximately 6-8 cm in front of the animal's face to capture licking

behavior. Far-red or infrared light was used to illuminate the animals face during behavioral

video acquisition. ROIs near the animal's mouth (160x160 pixels for 1080p video, 100x100

pixels for 720p video) were manually selected and cropped from the original video. Pixel data

types were converted from 16-bit RGB to 8-bit integer grayscale. Individual pixels were

normalized by taking the absolute value of the z score, using the mean and standard deviation

of an individual pixel over the duration of the entire video. Normalized frames were then spatially

downsampled to 5x5 pixels. 48 behavioral sessions that had simultaneous contact licometer lick

detection and behavioral video were used to identify which video frames contained spout licks.

10 frame sequences of these ground-truth labelled video frames were then used to train a

custom recurrent neural network (RNN) (10 long-short-term-memory node layer, followed by 10

node dense layer) using standard tools from the Python TensorFlow library. A RNN

implementation was chosen because it considers changes in pixels across sequential frames,

rather than only considering one frame at a time. The RNN model for binary frame classification

performed with high accuracy and specificity with a ROC AUC = 0.91. The model was then used

to predict licking for every frame of 104 30-minute behavioral videos. To further validate the

accuracy of the automated detection approach, the predicted classifications of the model were

compared to manually scored video data where researchers scored 1 second time bins as

containing animal licks or not. The model again reliably matched the compared lick scoring

information, with 82% median overlap between manually scored bins and recurrent neural

network frames (see **Extended Data Fig. 6**). Furthermore, the overall behavioral findings of

how much animals licked for different reward conditions and to reward-predictive cues across

Pavlovian conditioning were consistent between contact licometer, RNN-scored video and

manually-scored video approaches for lick detection. Code for RNN lick detection model is

available at github.com/christianepedersen.

In vivo 2-photon Calcium Imaging Acquisition

Image acquisition was performed simultaneously with reversible head-fixed behavior (see above) using the Olympus Fluoview FVMPE-RS 2-photon microscope. A 20x magnification air objective lens (Olympus, LCPLN20XIR, 0.45 numerical aperture, 8.3mm working distance) was optically coupled to the surgically implanted endoscopic lens (Inscopix, 7.3mm length, 0.6mm diameter GRIN lens) of an animal. 920 nm laser light (SpectraPhysics MaiTai Laser, ~100-femtosecond pulse width) excited GCaMP6s fluorophores during resonant scanning (30-Hz roundtrip frame rate, averaging every 6 frames in real-time). Emitted ~535 nm photons were directed to detectors (Olympus, GaAsP PMT) and attributed to the excited pixel. The same focal plane (<5 um tolerance) was optically accessed for each imaging session with the same animal so we could reliably track the activity of the same neurons across multiple days and conditions. A 200-frame, 40-second image series was initiated 20 seconds before reward delivery on each behavioral trial and was triggered by an Arduino Uno that also controlled reward delivery and cues.

***In vivo* 2-photon Calcium Imaging Data Processing**

The Olympus OIR files collected during imaging through Olympus FluoView (FV1200) were exported as tif files. Tif files for individual behavioral trials were motion corrected using a nonrigid motion correction package (Flatiron Institute: github.com/flatironinstitute/NoRMCorre) and concatenated using custom code in MatLab. These whole-session tif files were then converted to an HDF5 format using a custom code. These HDF5 files were then motion corrected in the x–y plane using an adapted hidden Markov model (SIMA version 1.3.0: losonczylab.org/sima/1.3/api/motion.html) as previously reported (*45*). We had found that the imaging plane showed very little z movement in mNAcSh (<5 μm based on a random sample of sessions). Individual neurons were then identified manually and selected by drawing regions of interest (ROIs) manually using the ImageJ polygon tool on the mean projection of pixel intensity

across all frames in a session. ROIs for different sessions that corresponded to the same

neuron and accurately captured the dynamics of the cell were used to 'track' that neuron across

days and conditions. Individual neurons that did not have consistent ROIs for all sessions were

not 'tracked' and were not used for comparison of responses to different conditions. The

retained ROIs were imported into SIMA and then used for signal extraction.

Custom code added neuropil correction to the SIMA signal extraction as previously described

(*21*). GCaMP6s labelled neurons were fairly sparse in the focal plane, with minimal overlap of

neurons in x, y dimensions. However, changes in the fluorescence of individual neuron ROIs

were still contaminated with slight variation in back fluorescence due to neuropil. Neuropil

correction was done by first calculating a neuropil signal around each ROI. This was done by

calculating a weighted sum of all recorded pixels excluding those falling within a 15-pixel (~17-

µm) radius of all ROIs. The weight for any pixel was calculated using a Gaussian function

centered on the ROI of interest with a radius of 50 pixels (~45µm). These parameters were

obtained after a systematic search of the parameter space in a small subset of sessions and

visually comparing the obtained fluorescence traces against the raw videos in the current

mNAcSh dataset as well as in previous studies (*21*). The results in this manuscript are robust

against large variation in these parameters. Once the neuropil signal was calculated for every

ROI, a correction of this signal was done by subtracting 0.8 multiplied by this signal from the raw

calcium trace of the ROI. The results of this previously published neuropil correction procedure

(*21*) were also found to be highly consistent with results from another prominent background

subtraction approach from the package Suite2p (*46*). After single cell fluorescence time series

were extracted and neuropil-corrected from raw calcium image sequences, they were detrended

by fitting a double exponential curve to the entire session and subtracting it. Changes in

fluorescence from the now consistent baseline were z-score normalized, using the mean and

standard deviation of the entire session.

In Vivo 2-photon Calcium Imaging Data Analysis

Individual neurons that were successfully tracked across days and conditions were further analyzed to understand their responsivity to different behavioral conditions. First, the average response of individual neurons to all variable sucrose and Pavlovian conditions was taken by taking the mean fluorescence of each cell across all 20 trials of a given behavioral session (or 10 rewarded trials or 10 extinction trials for Pavlovian Day 7). Prior to clustering, dimensionality reduction was performed on trial-averaged responses to all reward conditions (0-10 seconds from reward onset for variable sucrose sessions, -5-10 seconds from reward onset for Pavlovian sessions) to capture the most important Euclidean distance measures during subsequent clustering. Specifically, principal component analysis was performed and the number of components retained (6-8 components) was chosen through standard Scree methods. The principal components of reward responses of individual neurons across all variable sucrose or Pavlovian conditions were then used for K-means clustering (number of clusters with lowest silhouette score was chosen, 500x bootstrapping was performed to minimize variability from initial conditions) to group together neurons that had similar responses across these different conditions. K-means clustering and silhouette score calculations were performed in Python using the Scikit-learn library. Several clusters were identified among dyn-SPN and enk-SPN neurons for each behavioral paradigm. The trial-averaged responses for all neurons within a given cluster were then further averaged to generate the average response traces for each cluster to every condition.

To investigate whether the different reward conditions in the variable sucrose paradigm (0%, 3%, 10%, 30%) could be uniquely identified only by the simultaneous activity of mNAcSh SPNs, a multi-class linear support vector classifier (SVC) model (cost = 0.8) was created. The linear SVC model was trained and tested on the data for each animal individually. The class of the reward condition and the magnitude of the reward response (0-8 s from onset of reward

delivery) of every neuron for a given trial was used as input for the model (80 trials per animal,

20 trials per reward class). 80% of the trials were used to train the linear SVC and 20% were

used for cross-validation. The accuracy of the model during cross-validation for each animal is

reported in the Results. The linear SVC model was created in Python using the Scikit-learn

library and feature coefficients were taken directly from the model and averaged by which K-

means cluster the neurons belonged to. To test whether cue-excitation of mNAcSh SPNs was

correlated to Pavlovian conditioning session index, trial index within a session or to anticipatory

licking to the cue, Pearson's correlations were calculated relative to the magnitude of cue-

response (0-5 s from onset of cue) of SPNs that belonged to clusters that were both cue-excited

and reward-excited.

To investigate the relationship between SPN cue and reward-response and relative anatomical

position, we calculated the relative position of tracked SPNs in the focal plane during 2-photon

calcium imaging sessions. The 2-photon resonant scanner generated raw frames that were

512x512 pixels. The x and y pixel position at the center of each ROI that was manually drawn in

ImageJ was considered for each successfully tracked neuron. The relative anatomical position

of each neuron was calculated as the mean x and y position of its ROI centers across all

imaging sessions. Medial-lateral and anterior-posterior axes were inferred based on the

orientation of the animal being orthogonal to the objective lens during the imaging and behavior

session. The depth of the focal plane ventral to the GRIN lens during the imaging session

introduces a complex magnification effect, so the precise ratio between pixels and anatomical

distance in the brain could only be approximated based on the 600 um diameter of the lens and

the width of the focal plane being approximately equivalent.

***In Vivo* Fiber Photometry During Reversible Head-fixation**

Fiber photometry recordings were made throughout the entirety of 20 trial, 30-minute head-fixed

variable sucrose and Pavlovian conditioning sessions using a previously published approach

(*44*). Prior to recording, an optic fiber was attached to the implanted fiber using a ferrule sleeve

(Doric, ZR_2.5). Two LEDs were used to excite GCaMP6s. A 531-Hz sinusoidal LED light

(Thorlabs, LED light: M470F3; LED driver: DC4104) was bandpass filtered (470 ± 20 nm, Doric,

FMC4) to excite GCaMP6s and evoke Ca^{2+}-dependent emission. A 211-Hz sinusoidal LED light

(Thorlabs, LED light: M405FP1; LED driver: DC4104) was bandpass filtered (405 ± 10 nm,

Doric, FMC4) to excite GCaMP6s and evoke Ca^{2+}-independent isosbestic control emission.

Prior to behavior and recording, a 120 s period of GCaMP6s excitation with 405 nm and 470 nm

light was used to remove the majority of baseline drift. Laser intensity for the 470 nm and

405 nm wavelength bands were measured at the tip of the optic fiber and adjusted to ~70 μW

before each day of recording. GCaMP6s fluorescence traveled through the same optic fiber

before being bandpass filtered (525 ± 25 nm, Doric, FMC4), transduced by a femtowatt silicon

photoreceiver (Newport, 2151) and recorded by a real-time processor (TDT, RZ5P). The

envelopes of the 531-Hz and 211-Hz signals were extracted in real-time by the TDT program

Synapse at a sampling rate of 1017.25 Hz.

Photometry Data Analysis

Custom MATLAB scripts were developed for analyzing fiber photometry data in context of

mouse behavior and can be accessed online (github.com/christianepedersen). The isosbestic

405 nm excitation control signal was subtracted from the 470 nm excitation signal to remove

movement artifacts from intracellular Ca^{2+}-dependent GCaMP6s fluorescence (see **Extended**

Data Fig. 10). Baseline drift was evident in the signal due to slow photobleaching artifacts,

particularly during the first several minutes of each 30-minute recording session. A double

exponential curve was fit to the raw trace and subtracted to correct for baseline drift. After

baseline correction, the photometry trace was z-scored relative to the mean and standard

deviation of the entire session. The processed fluorescence timeseries was then aligned to

animal behavior and the average fluorescence relative to the time of behavioral events (sucrose

reward delivery) was visualized. Quantification of peri-event fluorescence was calculated by

taking the average of all samples over a given time window for each trial and comparing

between behavioral conditions.

Optogenetic Stimulation during Head-fixed Sucrose Consumption

After recovering from surgery, animals were water-restricted and trained to lick 10% sucrose

reward while reversibly immobilized and head-restrained. After 4 days of lick training, all animals

displayed consistent lick behavior in response to variable interval sucrose presentation and

were then put through 2 consecutive test days. Each test day involved one 30 minute, 20 trial

session in which the animals were presented with either 10% or 30% sucrose reward. The order

of sucrose condition days was pseudorandomized and counterbalanced across animals. For

consistency with the imaging and photometry behavior paradigms, animals were presented with

5 droplets of sucrose reward per trial with a total aggregate volume of 10 ul and 1 second

between droplet delivery. Animals immediately licked the presented sucrose reward, with

average licking bout durations of 7-8 seconds. On a pseudorandom 50% of the trials within a

session, time-locked optogenetic manipulation of mNAcSh-VP^{enk} or mNAcSh-LH^{dyn} activity

would be paired with reward delivery, via 4 seconds of 470 nm blue laser light delivery (4 mW

for ChR2; 10 mW for PPO measured at end of patch cable, 10 Hz, 10 ms pulse width). 10Hz

optogenetic stimulation frequency was chosen because it is physiologically relevant level of

neural activity for SPNs during phasic reward-related activity (*12*, *13*). Any differences in

consummatory licking between no laser and laser trials were recorded via the contact licometer

circuit with <1 ms time precision. All differences in licking between no laser and laser trials were

evaluated within subject to account for the baseline variability in lick rate between animals. To

control for possible behavioral effects from the delivery of the blue light itself, cre- littermate

controls were also injected, implanted, and tested in identical conditions.

Open-field Locomotion Assay

To determine whether mNAcSh-VP^{Penk} or mNAcSh-LH^{Pdyn} is naturally activated during bouts of

motor activity as they are during reward consumption (see **Fig. 4, Extended Data Fig. 8**),

photometry animals were tested in a locomotion assay. Animals freely roamed a 0.5x0.5 m

arena in far red light conditions in a quiet environment to measure the amount of spontaneous

locomotion in a context that was not anxiogenic. Simultaneous fiber photometry recordings were

conducted during the 30-minute long behavioral sessions. 15 FPS video was captured

throughout the behavioral session by a camera position directly above the behavioral arena.

Automated animal location tracking software (Noldus, Ethovision) was used on the video data

after the session to get the x and y coordinates of the animal in the behavioral arena for every

frame of the video. Changes in animal coordinates between frames were used to calculate the

instantaneous velocity of the animal for each frame. Pearson's correlations were calculated

between the velocity of the animal and z scored photometry fluorescence for every frame. Local

maxima for movement velocity were identified as discrete events by using the MatLab

'findpeaks()' command (peak prominence >1 SD, peak width >4 seconds). Windows of the

fluorescence time series were aligned to velocity peaks to visualize peri-velocity peak averaged

fluorescence. Contrary to the mNAcSh-VP^{Penk} and mNAcSh-LH^{Pdyn} reward-excitations observed

during sucrose consumption, bouts of substantial locomotion did not correlate to circuit

recruitment.

To determine if mNAcSh-VP^{Penk} or mNAcSh-LH^{Pdyn} activity manipulation could causally disrupt

the execution of motor actions, we then performed a similar locomotion assay experiment with

intermittent optogenetic photo-excitation or photo-inhibition. Bilaterally implanted optogenetics

animals (see above) freely roamed a 0.5x0.5 m arena in far red light conditions in a quiet

environment to measure the amount of spontaneous locomotion in a context that was not

anxiogenic. Every minute of the 14 minute behavioral session had 30 seconds of pulsing 470

nm laser light (4 mW for ChR2; 10 mW for PPO measured at end of patch cable, 10 Hz, 10 ms

pulse width) followed by 30 seconds of no laser. 15 FPS behavioral video was captured

throughout the session and was again used to calculate the instantaneous velocity of the animal

for every frame. Movement velocity was compared within subject and within session between

laser on epochs and laser off epochs. Optogenetic photo-excitation or photo-inhibition of

mNAcSh-VP^{enk} or mNAcSh-LH^{dyn} had no significant effect on animal movement velocity. There

was also no cumulative effect on locomotion throughout the session as movement velocity was

found to be consistent throughout the entire duration of the session.

Statistical Approaches and Analysis

All data collected were averaged and expressed as mean ± SEM. Statistical significance was

taken as *$p < 0.05$, **$p < 0.01$, and ***$p < 0.001$, as determined by Pearson's correlation,

Student's t-test or one-way ANOVA followed by Tukey post hoc tests as appropriate. For *in

situ* hybridization data, we used Student's t-test. For 2-photon imaging and photometry

experiments, we used Pearson's correlation and Student's t-tests, as appropriate. For

optogenetic behavioral experiments, we used pairwise t-tests for comparison within subject and

within session. All n values for each experimental group are described in the appropriate figure

legend. For behavioral experiments, group size ranged from $n = 6$ to $n = 10$. For *in

situ* hybridization quantification experiments, slices were collected from 2-3 mice, with data

averaged from 2-3 slices per mouse. Statistical analyses were performed in GraphPad Prism

8.0 (Graphpad, La Jolla, CA), MatLab 2020 (The MathWorks, Natick, MA) or with Python SciPy

library.

Acknowledgements

We thank Dylan Blumenthal, Michelle Chung, Taylor Hobbs, and Carina Pizzano for animal

colony maintenance. We thank the Bruchas lab, Stuber lab and NAPE Center for helpful

discussions. **Funding:** C.E.P. was funded by NIH grant DA051124. D.C.C was funded by NIH

grants NS007205, DA043999, DA049862, DA051489. M.R.B. was funded by NIH grants

R37DA033396, P30DA048736, and the Mallinckrodt Endowed Professorship. **Author Contributions:** C.E.P, D.C.C and M.R.B conceived the study and designed experiments. C.E.P, D.C.C., S.A.K and P.J.M. performed experiments and collected data. C.E.P analyzed data and designed analyses. M.M.G, Z.C.Z. and P.R.O contributed to experimental design and provided technical support for data collection. C.E.P., D.C.C and M.R.B wrote the manuscript. **Competing Interests:** The authors declare no competing interests. **Data and materials availability:** This study did not generate any new/unique plasmids, mouse lines or reagents. Custom MatLab and Python analysis codes were created to appropriately organize, process, and combine photometry and 2-photon recording data with associated behavioral data. Analysis code for photometry and 2-photon imaging data will be made available on github.com/christianepedersen. Data sets supporting the current study are available from the corresponding author upon reasonable request.

Materials and Correspondence

Further information and requests for resources and reagents should be directed to and will be fulfilled by corresponding author Michael R. Bruchas (mbruchas@uw.edu).

Extended Data Figure Legends

Extended Data Figure 1. Sucrose concentration preference of C57 mice under water restriction

(a) Schematic of brief access 2-bottle choice task: animal makes choice between two sippers containing different concentrations of sucrose reward.

(b) Time-course of task and representative lick data throughout session.

(c) Brief access 2-bottle choice task schedule: Alternating between pseudorandomized test days and 10% versus 10% days to neutralize spatial biases that arise during test days.

(d) Preference between left and right sippers containing identical rewards on spatial bias neutralization days (n = 10 animals, pairwise t-test: $p = 0.411$, $p = 0.760$, $p = 0.885$, $p = 0.784$, $p = 0.911$, $p = 0.949$).

(e) Preference between each pairing of the 4 sucrose concentration conditions; graphs rearranged in order of preference (n = 10 animals, pairwise t-test: *$p = 0.031$, **$p = 0.003$, *$p = 0.012$, $p = 0.227$, **$p = 0.001$, $p = *0.026$).

Extended Data Figure 2. Individual SPN tracking and clustering based on response to variable sucrose conditions

(a) Mean pixel values across entire in vivo 2-photon calcium imaging session for a representative Penk-cre animal to demonstrate the ability to resolve individual cell morphology and track neurons reliably across multiple days and conditions.

(b) Mean response (20 trials per condition) of individual Penk-SPNs to all sucrose concentration conditions. Individual tracked cells are arranged by their cluster ID and animal ID to show representation of different animals within the neuronal clusters.

(c) Mean response (20 trials per condition) of individual Pdyn-SPNs to all sucrose concentration conditions. Individual tracked cells are arranged by their cluster ID and animal ID to show representation of different animals within the neuronal clusters.

(d) Silhouette plot shown for optimal k-means clustering of Penk-SPNs.

(e) Silhouette scores for different Penk-SPN cluster counts to determine the optimal number of clusters.

(f) Silhouette plot shown for optimal k-means clustering of Pdyn-SPNs.

(g) Silhouette scores for different Pdyn-SPN cluster counts to determine the optimal number of clusters.

Extended Data Figure 3. Individual SPN tracking and clustering based on response to Pavlovian conditioning

(a) Mean pixel values across entire in vivo 2-photon calcium imaging session for a representative Pdyn-cre animal to demonstrate the ability to resolve individual cell morphology and track neurons reliably across multiple days and conditions.

(b) Mean response (20 trials per condition) of individual Penk-SPNs to cue and reward for all Pavlovian conditioning days. Individual tracked cells are arranged by their cluster ID and animal ID to show representation of different animals within the neuronal clusters.

(c) Mean response (20 trials per condition) of individual Pdyn-SPNs to cue and reward for all Pavlovian conditioning days. Individual tracked cells are arranged by their cluster ID and animal ID to show representation of different animals within the neuronal clusters.

(d) t-sne plot of variance in activity of Penk-SPN clusters during cue and reward for all Pavlovian conditioning days. Each dot represents an individual tracked neuron that composes the cluster.

(e) Silhouette scores for different Penk-SPN Pavlovian response cluster counts to determine the optimal number of clusters.

(f) t-sne plot of variance in activity of Pdyn-SPN clusters during cue and reward for all Pavlovian conditioning days. Each dot represents an individual tracked neuron that composes the cluster.

(g) Silhouette scores for different Pdyn-SPN Pavlovian response cluster counts to determine the optimal number of clusters.

Extended Data Figure 4. GRIN lens placements

(a) Surgical placements of endoscopic GRIN lenses in mNAcSh of Penk-cre animals.

(b) Surgical placements of endoscopic GRIN lenses in mNAcSh of Pdyn-cre animals.

Extended Data Figure 5. Overlap of variable sucrose clusters and Pavlovian conditioning clusters

(a) Fraction of Penk-SPNs in each variable sucrose cluster (outer donut) that were found to be cue or reward-responsive (inner donut) within the context of the Pavlovian conditioning paradigm.

(b) Fraction of Pdyn-SPNs in each variable sucrose cluster (outer donut) that were found to be cue or reward-responsive (inner donut) within the context of the Pavlovian conditioning paradigm.

(c) Number of Penk-SPNs that were tracked across entirety of variable sucrose and Pavlovian conditioning paradigms, organized by the intersection of their variable sucrose cluster and Pavlovian conditioning cluster.

(d) Number of Pdyn-SPNs that were tracked across entirety of variable sucrose and Pavlovian conditioning paradigms, organized by the intersection of their variable sucrose cluster and Pavlovian conditioning cluster.

Extended Data Figure 6. Training and Validation of Recurrent Neural Network for Lick Detection

(a) Representative frame from head-fixed behavior video. ROI near animals mouth cropped from video, spatially downsampled and normalized prior to frame classification.

(b) Cartoon depicting organization of RNN that uses changes in previous and future frames to predict licking in the current frame.

(c) Table depicting which data sets were used to initially train the RNN model using contact licometer information as frame labels and used to manually validate RNN model accuracy.

(d) Representative sequential RNN prediction probabilities for when animal is licking (blue) overlaid with timestamps of animal licking from contact licometer (orange).

(e) ROC curve showing specificity of RNN binary classification on entire 48 video training set.

(f and g) Representative session frame classification by RNN model (f) and manual video scoring (g).

(h) Overlap between manually scored video and RNN model video classification for all 104 behavioral videos.

Extended Data Figure 7. There is limited collateralization of Penk-SPN and Pdyn-SPN efferent circuits

(a) Schematic (left) and representative images (right) of ventral tegmental area (top) or lateral hypothalamus (bottom) of retrobead injection sites.

(b and c) In situ hybridization (b) and quantification (c) of prodynorphin, red, and green retrobeads in mNAcSh (scale bar = 200um). Percentage of DAPI-labelled cells that are also labelled with indicated probe.

(d) Schematic (left) and representative images (right) of ventral pallidum (top) or lateral hypothalamus (bottom) of retrobead injection sites.

1217 (e and f) In situ hybridization (e) and quantification (f) of proenkephalin, red, and green
1218 retrobeads in mNAcSh (scale bar = 200um). Percentage of DAPI-labelled cells that are also
1219 labelled with indicated probe.

1220

Extended Data Figure 8. mNAcSh-LH^{Pdyn} and mNAcSh-VP^{Penk} are mildly cue-responsive

(a) Schematic of simultaneous two-site fiber photometry during Pavlovian cue-reward
conditioning.

(b) Time-course of auditory cue and subsequent sucrose reward delivery for a given trial.

(c and d) Representative lick raster depicting individual licks that were detected by contact
licometer during cue and reward when animals were cue-naïve (m) and after conditioning (n).

(e) Mean lick rate during cue epoch (0-5 seconds from cue-onset) for each day of Pavlovian
conditioning and extinction (n = 15 animals; one-way ANOVA: Day 1 vs Day 5, *p = 0.041, Day
3 vs Ext., *p = 0.011, Day 5 vs Ext., ***p < 0.001, Day 7 vs Ext., **p = 0.005).

(f) Sagittal section cartoon displaying target injection region, cell-types and fiber implant
locations.

(g) Mean mNAcSh-VP^{Penk} reward-response for every uncued 10% sucrose trial and every cued
10% sucrose trial.

(h) Mean mNAcSh-VP^{Penk} activity over reward response epoch (2-12 seconds from reward-
onset) for uncued 10% sucrose, cued 10% sucrose and extinction trials (n = 9 animals, 96+
trials; one-way ANOVA: Uncued vs Cued, *p = 0.014, Cued vs Ext., ***p < 0.001).

(i) Mean mNAcSh-VP^{Penk} reward-response for every trial of day 7 Pavlovian extinction.

(j) Mean mNAcSh-VP^{Penk} activity over cue response epoch (2-5 seconds from cue-onset) for
cued 10% sucrose and extinction trials (n = 9 animals, 96+ trials; two sample t-test: p = 0.909).

(k) Representative coronal section showing Penk-SPN terminals in LH expressing GCaMP6s
underneath the fiber implant (scale bar = 200 um).

(l) Mean mNAcSh-LH^{Penk} reward-response for every uncued 10% sucrose trial and every cued
10% sucrose trial.

(m) Mean mNAcSh-LH^{Penk} activity over reward response epoch (2-12 seconds from reward-
onset) for uncued 10% sucrose, cued 10% sucrose and extinction trials (n = 6 animals, 64+
trials; one-way ANOVA: Uncued vs Cued, ***p < 0.001).

(n) Mean mNAcSh-LH^{Penk} reward-response for every trial of day 7 Pavlovian extinction.

(o) Mean mNAcSh-LH^{Penk} activity over cue response epoch (2-5 seconds from cue-onset) for
cued 10% sucrose and extinction trials (n = 6 animals, 64+ trials; two sample t-test: p = 0.851).

(p) Sagittal section cartoon displaying target injection region, cell-types and fiber implant
locations.

(q) Mean mNAcSh-LH^{Pdyn} reward-response for every uncued 10% sucrose trial and every cued
10% sucrose trial.

(r) Mean mNAcSh-LH^{Pdyn} activity over reward response epoch (2-12 seconds from reward-
onset) for uncued 10% sucrose, cued 10% sucrose and extinction trials (n = 6 animals, 63+
trials, one-way ANOVA: Uncued vs Cued, ***p < 0.001, Cued vs Ext., ***p < 0.001).

(s) Mean mNAcSh-LH^{Pdyn} reward-response for every trial of day 7 Pavlovian extinction.

(t) Mean mNAcSh-LH^{Pdyn} activity over cue response epoch (2-5 seconds from cue-onset) for
cued 10% sucrose and extinction trials (n = 6 animals, 63+ trials, two sample t-test: p = 0.073).

(u) Representative coronal section showing Pdyn-SPN terminals in VTA expressing GCaMP6s
underneath the fiber implant (scale bar = 200 um).

(v) Mean mNAcSh-VTA^{Pdyn} reward-response for every uncued 10% sucrose trial and every
cued 10% sucrose trial.

(w) Mean mNAcSh-VTA^{Pdyn} activity over reward response epoch (2-12 seconds from reward-
onset) for uncued 10% sucrose, cued 10% sucrose and extinction trials (n = 7 animals, 71+
trials, one-way ANOVA: Uncued vs Ext., **p = 0.009, Cued vs Ext., ***p < 0.001).

(x) Mean mNAcSh-VTA^{Pdyn} reward-response for every trial of day 7 Pavlovian extinction.

(y) Mean mNAcSh-VTA^{Pdyn} activity over cue response epoch (2-5 seconds from cue-onset) for
cued 10% sucrose and extinction trials (n = 7 animals, 71+ trials, two sample t-test: p = 0.387).

1270

Extended Data Figure 9. mNAcSh-LH^{Pdyn} may not causally drive sucrose consumption

(a) Schematic of time-locked optogenetic neural activity manipulation during consumption of
variable sucrose reward.

(b) Coronal section cartoon displaying target injection region, cell-type and bilateral fiber implant
location.

(c) Within-session comparison of 10% sucrose consumption between trials with and without
reward-paired optogenetic inhibition (n = 7 treatment, 5 control animals; pairwise t-test: p =
0.776, p = 0.187).

(d) Within-session comparison of 30% sucrose consumption between trials with and without
reward-paired optogenetic inhibition (n = 7 treatment, 5 control animals; pairwise t-test: p =
0.698, p = 0.220).

(e) Time-course of sucrose reward delivery for a given trial. 50% of trials also had 4 seconds of
10 Hz optogenetic stimulation concurrent with reward delivery.

(f) Coronal section cartoon displaying target injection region, cell-type and bilateral fiber implant
location.

(g) Within-session comparison of 10% sucrose consumption between trials with and without
reward-paired optogenetic excitation (n = 10 treatment, 5 control animals; pairwise t-test: **p =
0.002, p = 0.973).

(h) Within-session comparison of 30% sucrose consumption between trials with and without
reward-paired optogenetic excitation (n = 10 treatment, 5 control animals; pairwise t-test: p =
0.174, p = 0.857).

1292

Extended Data Figure 10. Fiber photometry signal preprocessing and implant placements

(a) Demodulated fluorescence timeseries for 405nm and 470 nm GCaMP6s excitation from
same recording session.

(b) Fitting a second exponential curve to the 470nm-405nm subtracted timeseries and
subtracting the fitted curve to 'detrend' the slow drift artifact from the whole session timeseries.

(c) Final normalized trace after calculation of ΔF/F using median of initial 470nm channel for
division.

(d) Mean mNAcSh-VP^{Penk} reward-response for 9 animals across all variable sucrose conditions.

(e) Surgical placements of 400 um diameter photometry fiber implants in VP of Penk-cre
animals.

(f) Surgical placements of 400 um diameter photometry fiber implants in LH of Penk-cre
animals.

(g) Surgical placements of 400 um diameter photometry fiber implants in VTA of Pdyn-cre
animals.

(h) Surgical placements of 400 um diameter photometry fiber implants in LH of Pdyn-cre
animals.

1309

Extended Data Figure 11. mNAcSh-VP^{Penk} and mNAcSh-LH^{Pdyn} are not recruited during
locomotion and cannot disrupt locomotion

(a) Schematic of fiber photometry recording during locomotion in a behavioral arena in red light
conditions.

(b) Simultaneous recording of fiber photometry signal and tracking of animal movement velocity.
Detection of instantaneous velocity peaks was performed post-hoc to identify movement bouts.

(c) Correlation between 1 second time bins of movement velocity and 1 second time bins of
GCaMP6s fluorescence (Pearson's correlation; n = 10 Penk-cre mice, 6 Pdyn-cre mice).

(d) Sagittal section cartoon displaying target injection region, cell-types and photometry
recording location.

(e) Mean mNAcSh-VP^{Penk} activity during every movement velocity peak.

(f) Sagittal section cartoon displaying target injection region, cell-types and photometry
recording location.

(g) Mean mNAcSh-LH^{Pdyn} activity during every movement velocity peak.

(h) Schematic of optogenetic stimulation (30 seconds on, 30 seconds off, repeating) during
locomotion in a behavioral arena in red light conditions.

(i) Representative timeseries of movement velocity throughout behavioral session, with
alternating stimulation schedule displayed.

(j) Mean movement velocity throughout duration of session for all stimulation and control
conditions.

(k) Within session comparison of movement velocity during mNAcSh-VP^{Penk} photo-inhibition via
PPO (n = 7 treatment, 5 control animals; pairwise t-test: p = 0.497, p = 0.378).

(l) Within session comparison of movement velocity during mNAcSh-LH^{Pdyn} photo-inhibition via
PPO (n = 7 treatment, 5 control animals; pairwise t-test: p = 0.903, p = 0.489).

(m) Within session comparison of movement velocity during mNAcSh-VP^{Penk} photo-excitation
via ChR2 (n = 9 treatment, 6 control animals; pairwise t-test: p = 0.910, p = 0.440).

(n) Within session comparison of movement velocity during mNAcSh-LH^{Pdyn} photo-excitation via
ChR2 (n = 8 treatment, 5 control animals; pairwise t-test: p = 0.923, p = 0.480).

FIGURE 1

FIGURE 2

FIGURE 3

Anatomical Separation of Variable Sucrose Clusters

a 355 Penk+ SPN — 112 Cells; 360 Pdyn+ SPN — 64 Cells

b Unpreferred — Preference (water-restricted) — Preferred

Cluster ID: 30% Sucrose, 0% Sucrose, 3% Sucrose, 10% Sucrose

Penk-Inhibited
Pdyn-Inhibited

0.5 SD | 5 s

c Penk-cre — Bregma, Lambda
Anterior → Posterior; Medial → Lateral; Approx. 100 um
Penk-SPN Inhibited Cluster *** Other Penk-SPN Clusters; 10% of FOV

d Pdyn-cre — Bregma, Lambda
Anterior → Posterior; Medial → Lateral; Approx. 100 um
Pdyn-SPN Inhibited Cluster *** Other Pdyn-SPN Clusters; 10% of FOV

e Relative Anatomical Cluster Location

Penk-Null
Penk-Inhibited ***
Penk-Preferred
Penk-Unpreferred

Pdyn-Null
Pdyn-Inhibited ***
Pdyn-Preferred
Pdyn-Sucrose **
Pdyn-AllRewards

Anterior ←→ Posterior; Lateral ←→ Medial
10% of FOV; 10% of FOV

Anatomical Separation of Pavlovian Clusters

f 363 Penk+ SPN — 51 Cells; 269 Pdyn+ SPN — 43 Cells

g Cluster ID: Day 1, Day 3, Day 5, Day 7, Day 7 Extinction
Reward-inhibited Clusters
0.5 SD | 5 s

h Penk-cre — Bregma, Lambda
Anterior → Posterior; Medial → Lateral; Approx. 100 um
Penk-SPN Reward-Inhibited Cluster *** Other Penk-SPN Clusters; 10% of FOV

i Pdyn-cre — Bregma, Lambda
Anterior → Posterior; Medial → Lateral; Approx. 100 um
Pdyn-SPN Reward-Inhibited Cluster ** Other Pdyn-SPN Clusters; 10% of FOV

j Relative Anatomical Cluster Location

Penk-Null
Penk-Rew-inhibited ***
Penk-Rew, cue-excited

Pdyn-Null
Pdyn-Rew-inhibited **
Pdyn-Rew, cue-excited
Pdyn-Rew-excit.

Anterior ←→ Posterior; Lateral ←→ Medial
10% of FOV; 10% of FOV

FIGURE 4

FIGURE 5

SUPPLEMENTAL FIGURE 2

a

b

Sucrose Reward Conditions

0%   3%   10%   30%

Penk-Preferred Cluster

Penk-cre Animals
- 1
- 2
- 3
- 4
- 5
- 6
- 7

Penk-Null Cluster

z score
1.5
1.0
0.5
0
-0.5
-1.0
-1.5

Penk-Unpreferred Cluster

50 neurons

Penk-Inhibited Cluster

20 s

c

Sucrose Reward Conditions

0%   3%   10%   30%

Pdyn-Inhibited Cluster

Pdyn-cre Animals
- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11

Pdyn-Preferred Cluster

z score
1.5
1.0
0.5
0
-0.5
-1.0
-1.5

Pdyn-Null Cluster

Pdyn-All Rewards Cluster

50 neurons

Pdyn-Sucrose Cluster

20 s

d

Penk-Preferred
Penk-Unpreferred
Penk-Null
Penk-Inhibited

Silhouette Coefficient Values
0   0.2   0.4   0.6   0.8   1

e

Penk-SPN Silhouette Scores

Silhouette Score

Number of Clusters
2   3   4   5   6   7   8

f

Pdyn-All Rewards
Pdyn-Preferred
Pdyn-Sucrose
Pdyn-Null
Pdyn-Inhibited

Silhouette Coefficient Values
0   0.2   0.4   0.6   0.8   1

g

Pdyn-SPN Silhouette Scores

Silhouette Score

Number of Clusters
2   3   4   5   6   7   8

a

b

Pavlov Day 1 | Pavlov Day 3 | Pavlov Day 5 | Pavlov Day 7 | Day 7 Extinction

Cue 10% Sucrose

Penk-cre Animals

z score
1.5
1.0
0.5
0
-0.5
-1.0
-1.5

Reward and Cue-Excited Cluster

Reward-Inhibited Cluster

50 neurons

Null Cluster

20 s

c

Pavlov Day 1 | Pavlov Day 3 | Pavlov Day 5 | Pavlov Day 7 | Day 7 Extinction

Cue 10% Sucrose

Pdyn-cre Animals

Reward and Cue-Excited Cluster

z score
1.5
1.0
0.5
0
-0.5
-1.0
-1.5

Null Cluster

Reward-Inhibited Cluster

50 neurons

Reward-Excited Cluster

20 s

d

Null
Reward-inhibited
Reward, cue-excited

t-SNE 1
t-SNE 2

e

Penk-SPN Silhouette Scores

Silhouette Score
0.5
0.4
0.3
0.2
0.1
0

Number of Clusters
2 3 4 5 6 7 8

f

Null
Reward-inhibited
Reward, cue-excited
Reward-excited

t-SNE 1
t-SNE 2

g

Pdyn-SPN Silhouette Scores

Silhouette Score
0.5
0.4
0.3
0.2
0.1
0

Number of Clusters
2 3 4 5 6 7 8

SUPPLEMENTAL FIGURE 4

a

Penk-cre mice
GRIN lens placements

1.42mm

- 152a
- 155a
- 164a
- 167a
- 167e

1.34mm

- 164b
- 164c

1.18mm

b

Pdyn-cre mice
GRIN lens placements

1.42mm

- 20b
- 20c
- 22a
- 22b
- 312c
- 312d

1.34mm

- 29a
- 305a
- 311c
- 251a
- 768a

1.18mm

SUPPLEMENTAL FIGURE 5

a Penk-SPN Variable Sucrose Clusters

- Penk-Null
- Penk-Inhibited
- Penk-Preferred
- Penk-Unpreferred

Pavlovian Clusters

- Cue + reward-excited
- Reward-excited
- Reward-inhibited
- Null

70 Cells
61 Cells
112 Cells
112 Cells

b Pdyn-SPN Variable Sucrose Clusters

- Pdyn-Null
- Pdyn-Inhibited
- Pdyn-Preferred
- Pdyn-Sucrose
- Pdyn-AllRewards

66 Cells
56 Cells
26 Cells
64 Cells
148 Cells

c

8 neurons not tracked	Penk-Null	Penk-Inhibited	Penk-Preferred	Penk-Unpreferred
Null	62	60	19	10
Reward-inhibited	9	39	2	1
Cue + reward-excited	41	13	49	50

d

112 neurons not tracked	Pdyn-Null	Pdyn-Inhibited	Pdyn-Preferred	Pdyn-Sucrose	Pdyn-AllRewards
Null	49	11	4	10	3
Reward-inhibited	22	16	0	4	0
Reward-excited	5	4	4	15	4
Cue + reward-excited	26	1	25	26	19

SUPPLEMENTAL FIGURE 6

a

Frames with contact licometer triggered

Frames without contact licometer triggered

| z-score pixel intensity |

0   1.4

b

Automated lick detection for every frame using Recurrent Neural Network

Predict licking in frame t

Detect changes in sequences of frames

c

		Behavior Video	Simultaneous Contact Licometer	Manual Video Scoring
Supervised Training	48 videos	X	X	
Predicting	104 videos	X		
Manual Validation	104 videos	X		X

Supervised Training

d

● Contact licometer triggered   — RNN prediction probability

e

Receiver Operating Characteristic

AUC = 0.91

Manual Validation

f

g

h

Manual Validation of RNN Accuracy

Percent overlap with manually scored lick bins

SUPPLEMENTAL FIGURE 7

SUPPLEMENTAL FIGURE 9

a

Time-locked
Optogenetic
Manipulation

Sipper

Sucrose

b

Pdyn-cre^{mNAcSh}
AAV5-hSyn-DIO-PPO

c

10% Sucrose

mNAcSh-LH^{Pdyn} cre- control

Mean licks per second

Off   Laser   Off   Laser

d

30% Sucrose

mNAcSh-LH^{Pdyn} cre- control

Mean licks per second

Off   Laser   Off   Laser

e

Reward-Paired
ChR2 Excitation
or PPO Inhibition

Laser on
50% of trials

10 Hz

Random
ITI

1s

● 2 ul Sucrose Delivered
| Consummatory Licks

f

Pdyn-cre^{mNAcSh}
AAV5-hSyn-DIO-ChR2

g

10% Sucrose

mNAcSh-LH^{Pdyn} cre- control

**

Mean licks per second

Off   Laser   Off   Laser

h

30% Sucrose

mNAcSh-LH^{Pdyn} cre- control

Mean licks per second

Off   Laser   Off   Laser

SUPPLEMENTAL FIGURE 10

a Fiber Photometry

b Velocity Peaks

c r = 0.0105 p = 0.1609
r = 0.0113 p = 0.2390

mNAcSh-VP^{Penk}
mNAcSh-LH^{Pdyn}

d VP LH
Penk-cre
AAV-DJ-DIO-GCaMP6s

e Peak velocity
mNAcSh-VP^{Penk}

f LH VTA
Pdyn-cre
AAV-DJ-DIO-GCaMP6s

g Peak velocity
mNAcSh-LH^{Pdyn}

h Optogenetic Excitation or Inhibition
10 Hz

i laser on

j Penk:ChR2   Pdyn:ChR2
Penk:PPO   Pdyn:PPO
cre- controls

k Penk-cre^{mNAcSh}
AAV5-hSyn-DIO-PPO
mNAcSh-VP^{Penk}
cre- control

l Pdyn-cre^{mNAcSh}
AAV5-hSyn-DIO-PPO
mNAcSh-LH^{Pdyn}
cre- control

m Penk-cre^{mNAcSh}
AAV5-hSyn-DIO-ChR2
mNAcSh-VP^{Penk}
cre- control

n Pdyn-cre^{mNAcSh}
AAV5-hSyn-DIO-ChR2
mNAcSh-LH^{Pdyn}
cre- control