1 **Visual processing and collective motion-related decision-making in desert locusts**

2 Itay Bleichman[1], Pratibha Yadav[1,2] and Amir Ayali[1,2*]

3 *[1] School of Zoology, Tel Aviv University, Israel*
4 *[2] Sagol School of Neuroscience, Tel Aviv University, Israel*
5

*Author for correspondence

Abstract

Collectively moving groups of animals rely on the decision-making of locally interacting

individuals in order to maintain swarm cohesion. However, the complex and noisy visual

environment poses a major challenge to the extraction and processing of relevant

information. We addressed this challenge by studying swarming-related decision-making

in desert locust last-instar nymphs. Controlled visual stimuli, in the form of random dot

kinematograms, were presented to tethered locust nymphs in a trackball setup, while

monitoring movement trajectory and walking parameters. In a complementary set of

experiments, the neurophysiological basis of the observed behavioral responses was

explored. Our results suggest that locusts utilize filtering and discrimination upon

encountering multiple stimuli simultaneously. Specifically, we show that locusts are

sensitive to differences in speed at the individual conspecific level, and to movement

coherence at the group level, and may use these to filter out non-relevant stimuli. The

locusts also discriminate and assign different weights to different stimuli, with an observed

interactive effect of stimulus size, relative abundance, and motion direction. Our findings

provide insights into the cognitive abilities of locusts in the domain of decision-making

and visual-based collective motion, and support locusts as a model for investigating

sensory-motor integration and motion-related decision-making in the intricate swarm

environment.

Keywords: *Schistocerca gregaria*, cognition, discrimination, visual processing, swarming

Introduction

A fundamental aspect of all instances of collective motion is that of individual repeated decision-making [1–3]. This, in turn, is both driven by and relies on local interactions among the constituent agents, requiring each agent to obtain information about its surrounding social environment [4]. The consequent formation and maintenance of this distinctive form of synchronized movement is understood to be beneficial to the participating individuals [5–7].

A quintessential example of the above process is displayed by the desert locust, *Schistocerca gregaria* (Acrididae). When in the gregarious phase, they collectively move in huge dense marching swarms ([8][9], figure 1A). Locust swarming is commonly accepted as heavily relying on visual perception [10]: each individual locust, with limited visibility amidst an unpredictable terrain, and an intricate, continuously changing social environment, must engage in repeated and dynamic decision-making to avoid getting derailed, while at the same time sustaining the collective motion. This can be translated into a two-layer process: the continuous extraction of the (unknown) state of the social surroundings from the input received by the sensory system (i.e. the eyes); and sensory-motor integration to facilitate the appropriate motor response. Different approaches, ranging from mathematical modeling to studying synchronization in small groups of locusts in laboratory settings, have been employed in the study of swarming behavior in the desert locust [11–14]. However, our understanding of swarm formation and

maintenance is still far from complete, partly due to a lack of answers to some fundamental

questions regarding decision-making at the individual level.

The swift extraction and processing of relevant information from a changing, complex

sensory environment presents a critical challenge [15], especially in the noisy and cluttered

visual surroundings of a locust swarm. Insects may adopt a range of strategies to increase

the efficiency of information perception and processing by reducing the information load

[16]. Such strategies include filtering relevant stimuli [17], categorizing the targets [18],

and generalizing visual patterns [19]. Filtering relevant visual stimuli, for example by

employing a "matched filter" in the visual modality, can reduce the amount of information

that needs to be processed [16,20]. In dragonflies and hoverflies, for example, small target

detectors are specifically tuned to objects that constitute only a 1–3° angle of the visual

field [21,22]. The filtering may occur at different levels of stimuli processing and vary with

the ecological relevance of the stimuli. The insect's nervous system can then channel its

resources into performing essential computations, even if complex, in order to extract the

task-relevant visual information at low energetic cost [23] . In the case of the desert locust,

we hypothesize that, during collective-motion-related visual processing, the locust

identifies and extracts relevant stimuli – swarming-related visual cues – from the overall

visual scenery, based on a subset of visual features, enabling swift and appropriate

decision-making. It is possible that a matched filter for walking speed is used to recognize

marching conspecifics; while filtering based on the coherence of the moving group, as

inferred from a subset of the swarm, might be used to estimate the overall direction of the

swarm.

An additional difficulty imposed on information gathering can arise from the presence of

multiple relevant competing inputs [24–26]. In this case, reducing the information load can

also be achieved through selective attention – the ability to focus on one type of preferred

stimulus while ignoring other perceivable ones [16,27]. Although selective attention

remains arguable in the context of insects, the much related key capability to discriminate

among different stimuli based on shape, color, and pattern orientation has been observed

in honey bees and bumblebees [28–31]; as well as in fruit flies, which show anticipatory

behavior consistent with selective attention to the tracked visual stimulus [32,33].

Desert locusts exhibit a characteristic pause-and-go motion, with pause duration correlated

with a high probability of turning to change direction [34]. We can thus refer to the locust

collective motion as comprising a series of repeated decisions taken by the individuals in

the group [13]. Additionally, the decision-making process itself can be considered as a

problem of vector selection, including a choice between continued standing or initiating

walking, and a choice of direction. Observed variations in the fraction of time spent

walking, and particularly in pause duration and the subsequent change in direction, in

response to different visual stimuli, can thus offer valuable insights into the locust decision-

making process.

We have previously shown that a specific motion-sensitive descending interneuron (one of

many behaviorally-relevant descending interneurons (DINs, e.g. [35]), the descending

contralateral movement detector (DCMD), conveys information relevant to the locust

response to small, slow moving objects (such as other marching locusts [13] and see also

[36], [37]). Furthermore, this pathway was shown to demonstrate density-dependent phase-

related differences [13], [38], manifested in gregarious locusts being better suited than

solitarious ones to the repeated decision-making, and thereby facilitating and coordinating

the marching behavior of the swarm. Monitoring the DCMD response to various swarming-

related visual stimuli may offer some insights into the neural mechanisms behind the

decision-making process under focus in this study.

Here we explored swarming-related decision-making at the behavioral level in *S. gregaria*

nymphs, by analyzing different aspects of the individual locust's walking behavior. These

served in our investigation of the role of visual feature recognition and discrimination as

possible underlying mechanisms in decision-making. A complementary preliminary

electrophysiological study of the processing of visual-motion inputs, relevant to the

dynamic interactions between the individuals in a marching swarm, has lent further support

to our hypotheses.

Methods:

Animals: All experiments were carried out using V^{th}-instar larvae of *S. gregaria,* taken

from our high-density, gregarious phase locust lab-colony at the School of Zoology, Tel

Aviv University (rearing conditions were as recently described in [12].

The experimental setup: Individual locusts were tethered in a fixed (forward) head

direction, via a 1 cm long clear vinyl tube attached to their pronotum with epoxy resin, in

a natural-like typical walking posture, above an airflow-suspended Styrofoam trackball,

illuminated from above with LED lights. The ball was decorated with an irregular black

over white pattern in order to facilitate the tracking of its movement. Two parallel LCD

screens, 30 cm apart, were positioned one on either side of the locust, allowing the

presentation of controlled visual stimuli, while carefully monitoring the locust's behavioral

responses and movements of the ball by a high-speed video camera (figure 1B).

Experiments started after one hour of acclimation of the locust to the tether. In two sets of

behavioral experiments the locusts' responses were monitored using FicTrac [39], a

computer-vision tracking software that determines the angular position of the ball for each

frame. In an additional set of experiments, an optical mouse sensor was further utilized to

record the movement of the ball. The behavioral setup was complemented by a

corresponding electrophysiological setup, enabling the recording of the neural responses

of the locust DCMD interneurons to (similar) controlled visual stimuli (see

Electrophysiology section below).

The visual stimuli: Visual stimuli, designed using the Python programming language and

PsychoPy (an open source software package; [40]), were presented in the form of random

dot kinematograms (RDK) of black dots on a white background, at a maximum contrast of

100%. We chose RDK following previous reports of utilizing such stimuli for testing

multiple target processing, and specifically motion perception [41]. Unless stated

otherwise, the RDK comprised 40, 1.2 cm diameter dots, corresponding to a subtended

visual angle of 6.86° on the insect's eye (within the known size of the locusts). Each visual

stimulus was presented for 60 seconds.

We first presented the control stimuli: (1) blank (white screen), and (2) still dots on a white

screen. Next, we conducted a set of different behavioral experiments to investigate the

tethered locust's response to the following different tentative features of swarming-related

visual stimuli:

Direction of motion – The RDK comprised fully coherent, 5 cm/s moving dots, simulating a coherently moving locust swarm. Three types of stimuli were used, each with a different direction of motion: (1) both screens showing dots aligned with the direction of the tethered locust's heading; (2) both screens showing dots in a direction 180° to the tethered locust's heading; and (3) one screen showing aligned dots and the other with dots moving in the opposite direction.

Motion speed - Tethered locusts were presented with dots moving with 100% coherence on both screens, aligned with the tethered locust's heading, and at graded speeds. The tested motion speeds were 1, 3, 5, 10 and 15 cm/s, which cover a marching locust's speed range, as measured previously [12].

Coherence level - Tethered locusts were presented with dots moving on both screens at a motion speed of 5 cm/s and graded coherence levels: a fraction of the dots moved in alignment with the locust's heading while the remaining dots each moved in a random direction. Coherence levels tested were 0 (all dots moving in different random directions), 0.1, 0.25, 0.5, and 1 (all dots aligned with the locust's heading direction).

Competing stimuli - A fourth experiment was conducted to investigate situations of competing stimuli, i.e., decision-making in the presence of conflict. First, a more quantitative type of conflict was presented to the locusts: 2/3 of the dots on each screen moved in one direction, either aligned with or opposite to the tethered locust's heading, while the remaining 1/3 moved in the other direction. Next, a size difference was added (size mimicking proximity differences): the 1/3 dots moving in the opposite direction to the 2/3 were also double the size of the latter (2.4 cm diameter).

Behavioral analysis: The rotation angle, the difference between two angular positions of

the trackball in subsequent frames, was used to analyze the locust motion parameters. A

motion threshold was determined based on the extent of the rotation angle. A locust was

considered to be moving if the threshold was crossed for at least 10 consecutive frames.

Pausing was determined if the same threshold was not crossed for at least 20 consecutive

frames. Based on these indices, we calculated the fraction of time spent walking (walking

fraction) and the average pause duration. A sideways motion (positive and negative)

threshold was determined based on the direction of the trackball rotation. A locust was

considered to be moving sideways if this threshold was crossed for at least 10 consecutive

frames. The coordinate positions from the optical mouse sensor were used to determine the

walking parameters by calculating the walking distance, using the Cartesian formula.

Electrophysiology: Dissection and electrophysiological procedures followed Ariel et al.,

(2014). Briefly, following CO_2 induced anesthesia, the legs of the nymphs were removed

and a silver hook electrode was positioned around the ventral neck connectives for

extracellular recording of DCMD activity. The locusts were positioned above a plastic

platform in the same position and posture as on the airflow-trackball. The experiments were

performed using RDKs with graded speeds and coherence levels similar to the behavioral

experiments, and also using graded sizes of 0.4 ,0.8, 1.2, 1.4, 2.2, 5.5 and 6.8 cm diameter.

Each stimulus was presented for 20 seconds, with an inter-trial period of 1 min. The DCMD

action potential times, number, and frequency were analyzed.

Results:

187 ***The locusts respond to swarming-related visual stimuli with a preference to maintain***

188 ***their heading***

189 Locusts tethered in our setup exhibited the typical pause-and-go motion pattern even in the

190 absence of moving visual stimuli (see also (10,34)). Similar walking kinematics

191 (manifested in pause duration and walking fraction) were measured in response to the white

192 background only and to the background with motionless dots (Fig. S1). When testing the

193 effect of moving stimuli, the response of the locusts comprised several clear time-

194 dependent features (Fig. S2). Specifically, when the direction of the stimulus motion on

195 one or both screens was opposite to the tethered locust's head direction, the demonstrated

196 behavioral response was not consistent throughout the stimuli, but became exhausted prior

197 to termination of the stimuli. This time-limited response was probably due to the open-loop

198 nature of our experiments, i.e., to the fact that the locust's response did not induce any

199 (expected) directional change in the incoming visual inputs. Consequently, we limited our

200 comparative analysis of the different visual-stimuli-induced behavioral responses to the

201 first 40 seconds of each trial only. During this consistent and robust "responsive time-

202 interval", the locusts attempted to align themselves with the direction of motion of the dots,

203 and/or to join the motion (figure 2).

204 When characterizing the behavioral response to stimuli moving on both screens in the

205 locust's heading, compared to both screens showing stimuli moving in the opposite

206 direction, the latter generated a significantly decreased average pause duration (figure 2b,

207 n=26, Friedman test, $p < 0.05$, Dunn's multiple comparisons test, $p < 0.05$) and significantly

208 increased overall side motion (figure 2c, n=26, Friedman test, $p < 0.001$, Dunn's multiple

209 comparisons test, $p < 0.01$). When each screen displayed a different direction of motion, one

aligned with the locust's head direction and the other opposite to it, no significant difference

was observed in the above-noted parameters between this condition and the two others.

However, the locust's side motion towards the monitor displaying moving dots in a

direction aligned with its heading was significantly higher compared to its side motion

towards the other monitor (figure 2c, n=26, Wilcoxon matched-pairs signed rank test,

$p<0.05$). No such preference for motion towards a specific side was noted when both

screens displayed stimuli with the same direction of motion. Overall, these findings

confirm the swarming-related nature of our controlled stimuli, i.e., the locusts clearly

attempted to swarm alongside or to join the controlled visual stimuli presented in our

experimental setup, demonstrating a preference towards stimuli that were aligned with their

initial heading.

***Clear thresholds are demonstrated in the response to swarming-related visual stimuli at
both the individual conspecific and the group level.***

The next feature investigated for a possible effect on the locust's behavior was motion

speed. As can be seen in figure 3a, a clear dependence and a clear speed threshold were

demonstrated: in response to stimuli with motion speed greater than 3 cm/sec, significantly

higher walking fractions (figure 3a(i), n=15, Kruskal-Wallis test, $p<0.0001$) and shorter

pause durations figure 3a(ii), n=15, Kruskal-Wallis test, $p<0.0001$) were observed,

compared to the response to dots moving at speeds below this threshold.

Maintaining the speed of all the moving dots above the demonstrated threshold, and

changing the coherence level among the presented dots, revealed a second decision rule

based on yet another threshold (figure 3b): in response to stimuli with coherence level

above 25%, the locusts exhibited significantly larger walking fractions (figure 3b(i), n=16,

Kruskal-Wallis test, p<0.0001) and significantly shorter pause durations (figure 3b(ii),

n=16, Kruskal-Wallis test, p<0.0001) compared to their response to stimuli with coherence

levels below this threshold. It is important to note that while both the speed and the

coherence level are characteristic features of the visual inputs in a marching swarm, the

former is a feature of each individual group member, contributing to the collective motion

of the swarm; while the latter is a characteristic of the collective, or a group-level trait,

reflecting the common direction of motion.

The locust response to complex moving visual stimuli

As noted, the visual environment within a locust swarm is an intricate and noisy one,

intriguing us to investigate locust response to complex and conflicting stimuli (figure 4).

First, locusts were presented with visual stimuli comprising two groups: the 2/3 group of

dots moved in one direction and the 1/3 group of dots moved in the opposite direction All

dots moved at similar speeds, above the speed threshold demonstrated previously. Next,

a size difference was introduced, such as the dots in the minority group being twice the

size of those in the majority group. When 2/3 of the dots were moving in a direction

aligned with the locust's heading and the smaller group of dots were moving in the

opposite direction, no significant effect was observed in the locust's response following

an increase in size of the dots in the smaller group. However, when the majority of dots

were moving in the opposite direction to that of the locust's heading, changes in locust

kinematics were noted. When all the dots were equal in size, the locust's walking fraction

significantly decreased (figure 4a, n=19, Friedman test, p<0.05, Dunn's multiple

comparisons test, p<0.05). This was possibly due to the conflict between relative

abundance (2/3 of dots moving in the opposite direction) and the preferred motion

direction still present in the remaining 1/3. Doubling the size of the dots in the 1/3 group,

moving in the direction of the locust's heading partially restored walking fraction and

significantly increased pause duration (figure 4b, n=19, Friedman test, p<0.01, Dunn's

multiple comparisons test, p<0.05). This specific complex visual stimulus required more

intensive information processing by the locust, demonstrated by the larger pauses.

Overall, these findings reveal intricate interactions between stimulus number, size, and

direction, which together affect the locust decision-making process.

Neurophysiological correlates to the responses to swarming-related visual cues

Further exploration of the sensory-motor processing of swarming-related visual cues was

conducted through a series of neurophysiological experiments.

Based on the behavioral observations, we expected our different visual stimuli to induce

variable neuronal responses, depending on the stimuli motion speed, coherence level, and

dot size. This hypothesis was tested by studying the response of the DCMD interneuron,

a key participant in a well-described motion sensitive visual pathway [42,43], to similar

types of stimuli as above. The DCMD has been mostly studied in the context of looming

stimuli. Hence, it should be noted that the responses observed and monitored in our

experiments differ from those of the typical looming response (figure 5). The DCMD firing

rate in response to control stimuli (white background and still dots) was similar to its spontaneous firing rate reported in previous studies [13]. Manipulating the characteristics of swarming-related visual stimuli thus induced different responses:

Motion speed - We found a clear dependence of the DCMD firing rate on the speed of the moving dots. A significant difference was seen between the slowest tested motion speed – 1 cm/s, and the fastest one – 15 cm/s (figure 6a , n=7, Friedman test, $p<0.001$, Dunn's multiple comparisons test, $p<0.001$), comprising two extremes within the speed range of marching locusts [12]. The DCMD's responses to visual stimuli moving at intermediate speeds did not significantly differ from each other.

Coherence level - Low coherence levels elicited high DCMD firing rates, with the DCMD response declining with the increase in motion coherence level from 0/1 to 1 (figure 6b). DCMD firing rate with coherence levels of 0% or 10% was significantly higher compared to that in response to 100% coherent stimuli (figure 6b, n=6, Friedman test, $p<0.01$, Dunn's multiple comparisons test, $p<0.01$).

Size effect - Testing the DCMD response to swarming-related moving stimuli comprising dots of different sizes, revealed a size-dependent firing rate: a significant difference was noted between dots with a diameter of 0.4 cm and those with a diameter of 2.2, 5.5 or 6.8 cm (figure 6c, n=10, Kruskal-Wallis test, $p<0.0001$, Dunn's multiple comparisons test, $p<0.051$). A graded increase in firing rate was seen with the increase in size.

Overall, our neurophysiologic investigation revealed that the DCMD was sensitive to different speeds, coherence levels, and sizes of swarming-related visual stimuli, in a similar though not identical manner to that revealed in our behavioral experiments.

Discussion

Sensory information has a crucial role in ecological decision-making [15]. In order to enable sensory processing to be swift and context-appropriate, organisms are required to identify and extract highly specific, behaviorally relevant, signals from their surroundings [44]. Different strategies for rapidly coping with a visually cluttered environment have been suggested in previous studies of different organisms engaged in vision-based collective motion [13,45–47]. Flocking birds were reported to consider visual information from a fixed number of influential neighbours (i.e., a topological range; [48]). Zebrafish rely on visual occupancy for direction choice [45], and use bout-like movements for conspecific recognition [49]. Collectively moving *Drosophila* larvae depend on the number of conspecifics and cues related to their unique visual kinematic for decision-making [50]. Beyond the principal role of motion sensitivity in maintaining synchrony during collective marching [11,12], only very limited knowledge is available regarding how locusts utilize visual-sensory cues for swarming-related decision-making amidst their highly challenging visual surroundings.

Our findings have identified specific characteristics of the behaviorally-relevant visual inputs affecting decision-making in desert locust nymphs. Moreover, we show, to the best of our knowledge for the first time, that locusts can extract collective-motion relevant information at both the individual conspecific level (i.e. speed) and the group level (coherence or common direction), possibly by means of filtering and discrimination. While filtering can reduce the information processing load at the very first stage by differentiating relevant from non-relevant stimuli and ignoring the latter, discrimination can aid the

extraction of information from the relevant stimuli and subsequently facilitate critical

decision-making.

Desert locust nymphs walk at an average speed of ~5 cm/sec [12]. In response to coherent

stimuli moving at non-zero speeds below 3 cm/sec, the tethered individuals exhibited

longer pause durations and shorter walking fractions, possibly reflecting longer decision-

making time due to a mismatch between dot speed and the expected behaviorally relevant

conspecifics' speed. Hence, the locusts employed filtering at the level of the characteristic

of the individual. In natural settings, such a clear speed threshold may be exploited to

recognize marching conspecifics, such that anything moving at a speed below the threshold

is ignored. This behavioral threshold was discovered in the V[th]-larval instar nymphs

(corresponding to walking speed at this stage). Since locust collective marching appears

early on and is maintained throughout the different developmental (larval) stages [51], and

as development is accompanied by a marked change in size as well as in walking speed, an

interesting point for future research would be that of developmental plasticity within the

observed speed threshold.

The relatively limited walking behavior demonstrated in response to dots moving in a non-

coherent fashion (a non-decisive state manifested by unusually long pause durations), could

have resulted from a lack of appropriate relevant information at the level of the group

movement pattern. Importantly, our findings indicate that the locusts perceive a complete

absence of motion cues (i.e., still dots) differently to that of an absence of conclusive

information in the motion cues (i.e., non-coherently moving dots). This could be reflected

in gregarious locusts continuing to walk even when alone (albeit with altered kinematics

[14], but tending to pause and wait when surrounded by conspecifics moving randomly,

e.g., early morning at the initial organizing stages of a swarm in a natural setting[51,52].

This ability to extract the trajectory of the surrounding locusts seems to be a fundamental

characteristic of gregarious-phase locusts, instrumental for the decision-making in enabling

the collective motion of locust swarms.

In his review, Warrant [53] discusses different visual matched filters and their important

role in the ecology of vision in insects. These include peripheral matched filters for sex

(mate), for prey detection and pursuit, and for the physical environment (aspects of the

physical terrain). Central visual matched filters include filters for the insect's own

locomotion speed ("fast" or "slow" eyes) and for navigation (the celestial pattern of

polarized light). Our above findings suggest yet another possible matched filter that is

crucial for locust swarming behavior: this can be referred to as a "social environment"

matched filter, or maybe even filters, as we have demonstrated filtering at both the level of

the motion of individual neighboring conspecifics as well as of the surrounding group.

Relevant conflicting stimuli increase the difficulty imposed on information processing and

decision-making. A conflict can derive from a contradiction in one feature (e.g. opposing

motion directions), requiring a single-attribute decision, or from the complex interactions

between several features (e.g. direction, abundance, and size), becoming a multi-attribute

choice problem [54]. Locusts presented with directionally contradicting but otherwise

identical swarming-related visual stimuli, demonstrated a preference for stimuli with

motion direction aligned with their own heading. This preference to join in the marching

aligned to the locust's current heading is consistent with the observation that marching

locusts in experimental ring-shaped arenas only rarely change direction [11,13]. When

different parameters provide conflicting or inconsistent information, it becomes beneficial

to assign a higher weight to one over the other. Ants, for example, rank several attributes

when faced with a multi-feature problem [55], and the ranking is dynamic in relation to

their current situation [56]. When presented with mixed-type stimuli, size affected the

locust's behavior only when in a specific relation to the direction of motion and relative

abundance. The interplay between these three parameters significantly affected pause

duration, which is the kinematic phase assumed to be dedicated to information processing

and decision-making [13,34]. Size may act as a proxy for visual target distance, with

nearest neighbors being larger. More attention dedicated to larger dots, although less

abundant, indicates that immediate neighbors may influence the decision of an individual

more strongly than more distant members of the swarm. This is also consistent with

previous reports suggesting a limited functional radius of attention around an individual in

a group [13];[14].

The DCMD is only one out of several currently known descending interneurons that take

part in motion-sensitive pathways [57,58]. It has been widely researched for its

characteristic response to looming visual stimuli, and its function in predator and

collision avoidance maneuvers [59–62]. Nevertheless, the motion-sensitive pathway in

which it takes part is capable of responding to different, complex types of visual-motion

stimuli [37,62]. The DCMD was also shown to demonstrate activity changes with

developmental stages [36]. Ariel et al. [13] demonstrated the its ability to convey

information regarding motion types similar to those of marching locusts, and a specific

"tuning" of the response habituation rate in swarming, gregarious-phase locusts compared

to the non-swarming solitarious-phase ones. Hence, the DCMD response to swarming-

related visual stimuli, to which locusts demonstrated behavioral responses, can provide

information regarding the neurophysiological mechanisms underlying these behavioral

responses, and the related decision-making process. The DCMD in our experimental

setting demonstrated consistent responses to non-looming visual stimuli, with a clear

effect of changes in motion speed, coherence level, and stimulus size. The sensitivity to

changes in these specific visual stimulus features emphasizes their importance and

supports the involvement of these during visual information processing in swarming-

related decision-making.

In its simplest form, decision-making can be understood as the process of selecting between

two alternatives. The flexibility of decisions is accepted as a trademark of higher cognition

in organisms [63]. Given that cognitive performances and integral abilities are often

assumed to be positively correlated with brain size across species, it is no surprise that the

miniature brains of insects are believed to limit their computational power and cognitive

abilities [64–67]. However, with mounting evidence in support of highly sophisticated

behaviors in insects [68], this assumption currently holds little ground. Here, we have

provided additional insights into insect cognition by investigating decision-making in a

collectively marching insect species. We demonstrate that desert locusts use

discrimination, mediated by selective attention, to extract relevant information from a

complex, noisy, visual environment. The rules of decision-making (decision rules) in

gregarious desert locusts seem to be a function of multiple interacting factors, with the

ultimate goal of staying in sync with conspecifics. A differential weightage to parameters

such as direction, number, and size of the stimuli was also observed, with conflicting

information increasing the difficulty imposed on the decision-making process.

To conclude, locusts utilize different mechanisms that enable them to meet the challenges

presented by the overloaded and cluttered visual environment, and that support the sensory

perception and integration required for collective motion-related decision-making. These

mechanisms constitute an instrumental aspect of their ability to synchronize with

conspecifics and maintain the cohesion of the swarm, and thus probably also exist in all

animals demonstrating visual-based collective motion. Much further work is required,

however, in order to uncover and describe the details of the sensory-motor integration (e.g.

the role of feedback from the motor system), to fully elucidate the underlying

neurophysiological mechanisms (e.g. additional key neuronal pathways), and to provide

insights into related brain-level phenomena, such as the representation of conspecifics and

their behavior (speed, direction, etc.), the actual depiction or abstraction of the swarm as a

whole, and more.

FUNDING
This research was funded by The Israel Science Foundation (ISF), research grant
2306/18.

AUTHOR CONTRIBUTION
AA conceived the study; IB and PY carried out experiments and data analysis. AA and
IB and PY wrote the manuscript
The authors have no competing interests related to this study.

Supplementary material is available online on DRYAD:
https://doi.org/10.5061/dryad.jdfn2z3dw

Bibliography

1.   Gueron S, Levin SA, Rubenstein DI. 1996 The Dynamics of Herds: From
      Individuals to Aggregations. *J. Theor. Biol.* **182**, 85–98.
      (doi:10.1006/jtbi.1996.0144)

2. Petit O, Bon R. 2010 Decision-making processes: the case of collective
movements. *Behav. Processes* **84**, 635–647. (doi:10.1016/j.beproc.2010.04.009)

3. Herbert-Read JE. 2016 Understanding how animal groups achieve coordinated
movement. *J. Exp. Biol.* **219**, 2971–2983. (doi:10.1242/jeb.129411)

4. Heras FJH, Romero-Ferrero F, Hinz RC, de Polavieja GG. 2019 Deep attention
networks reveal the rules of collective motion in zebrafish. *PLoS Comput. Biol.*
15, e1007354. (doi:10.1371/journal.pcbi.1007354)

5. Gil MA, Hein AM, Spiegel O, Baskett ML, Sih A. 2018 Social information links
individual behavior to population and community dynamics. *Trends Ecol. Evol.*
33, 535–548. (doi:10.1016/j.tree.2018.04.010)

6. Yang W-C, Schmickl T. 2019 Collective motion as an ultimate effect in crowded
selfish herds. *Sci. Rep.* **9**, 6618. (doi:10.1038/s41598-019-43179-6)

7. Be'er A, Ariel G. 2019 A statistical physics view of swarming bacteria. *Mov.*
Ecol. **7**, 9. (doi:10.1186/s40462-019-0153-9)

8. Ayali A. 2019 The puzzle of locust density-dependent phase polyphenism. *Curr.*
Opin. Insect Sci. **35**, 41–47. (doi:10.1016/j.cois.2019.06.008)

9. Cullen DA *et al.* 2017 From molecules to management: mechanisms and
consequences of locust phase polyphenism. In *Insect Epigenetics*, pp. 167–285.
Elsevier. (doi:10.1016/bs.aiip.2017.06.002)

10. Ariel G, Ayali A. 2015 Locust collective motion and its modeling. *PLoS Comput.*
Biol. **11**, e1004522. (doi:10.1371/journal.pcbi.1004522)

11. Buhl J, Sumpter DJT, Couzin ID, Hale JJ, Despland E, Miller ER, Simpson SJ.
2006 From disorder to order in marching locusts. *Science* **312**, 1402–1406.
(doi:10.1126/science.1125142)

12. Knebel D, Ayali A, Guershon M, Ariel G. 2019 Intra- versus intergroup variance
in collective behavior. *Sci. Adv.* **5**, eaav0695. (doi:10.1126/sciadv.aav0695)

13. Ariel G, Ophir Y, Levi S, Ben-Jacob E, Ayali A. 2014 Individual pause-and-go
motion is instrumental to the formation and maintenance of swarms of marching
locust nymphs. *PLoS ONE* **9**, e101636. (doi:10.1371/journal.pone.0101636)

14. Knebel D, Sha-Ked C, Agmon N, Ariel G, Ayali A. 2021 Collective motion as a
distinct behavioral state of the individual. *iScience* **24**, 102299.
(doi:10.1016/j.isci.2021.102299)

15. Hein AM. 2022 Ecological decision-making: From circuit elements to emerging

principles. *Curr. Opin. Neurobiol.* **74**, 102551. (doi:10.1016/j.conb.2022.102551)

16.   Bernays EA, Wcislo WT. 1994 Sensory Capabilities, Information Processing, and
Resource Specialization. *Q. Rev. Biol.* **69**, 187. (doi:10.1086/418539)

17.   Kohn JR, Heath SL, Behnia R. 2018 Eyes matched to the prize: the state of
matched filters in insect visual circuits. *Front. Neural Circuits* **12**, 26.
(doi:10.3389/fncir.2018.00026)

18.   Benard J, Stach S, Giurfa M. 2006 Categorization of visual stimuli in the
honeybee Apis mellifera. *Anim. Cogn.* **9**, 257–270. (doi:10.1007/s10071-006-
0032-9)

19.   Stach S, Benard J, Giurfa M. 2004 Local-feature assembling in visual pattern
recognition and generalization in honeybees. *Nature* **429**, 758–761.
(doi:10.1038/nature02594)

20.   Wehner R. 1987 ?Matched filters? ? neural models of the external world. *J. Comp.*
Physiol. **161**, 511–531. (doi:10.1007/BF00603659)

21.   Nordström K. 2012 Neural specializations for small target detection in insects.
Curr. Opin. Neurobiol. **22**, 272–278. (doi:10.1016/j.conb.2011.12.013)

22.   O'Carroll D. 1993 Feature-detecting neurons in dragonflies. *Nature* **362**, 541–
543. (doi:10.1038/362541a0)

23.   Warrant EJ. 2016 Sensory matched filters. *Curr. Biol.* **26**, R976–R980.
(doi:10.1016/j.cub.2016.05.042)

24.   Wang M-Y, Ings TC, Proulx MJ, Chittka L. 2013 Can bees simultaneously
engage in adaptive foraging behaviour and attend to cryptic predators? *Animal*
Behaviour **86**, 859–866. (doi:10.1016/j.anbehav.2013.07.029)

25.   van Swinderen B. 2012 Competing visual flicker reveals attention-like rivalry in
the fly brain. *Front. Integr. Neurosci.* **6**, 96. (doi:10.3389/fnint.2012.00096)

26.   Römer H, Krusch M. 2000 A gain-control mechanism for processing of chorus
sounds in the afferent auditory pathway of the bushcricket Tettigonia viridissima
(Orthoptera; Tettigoniidae). *J. Comp. Physiol. A* **186**, 181–191.
(doi:10.1007/s003590050018)

27.   Nityananda V. 2016 Attention-like processes in insects. *Proc. Biol. Sci.* **283**.
(doi:10.1098/rspb.2016.1986)

28.   Giurfa M. 2004 Conditioning procedure and color discrimination in the honeybee
Apis mellifera. *Naturwissenschaften* **91**, 228–231. (doi:10.1007/s00114-004-

0530-z)

29.   Dyer AG, Chittka L. 2004 Fine colour discrimination requires differential conditioning in bumblebees. *Naturwissenschaften* **91**, 224–227. (doi:10.1007/s00114-004-0508-x)

30.   1995 Shape vision in bees: innate preference for flower-like patterns. *Phil. Trans. R. Soc. Lond. B* **347**, 123–137. (doi:10.1098/rstb.1995.0017)

31.   1994 Visual discrimination of pattern orientation by honeybees: performance and implications for "cortical" processing. *Phil. Trans. R. Soc. Lond. B* **343**, 199–210. (doi:10.1098/rstb.1994.0021)

32.   Tang S, Juusola M. 2010 Intrinsic Activity in the Fly Brain Gates Visual Information during Behavioral Choices. *Nature Precedings* (doi:10.1038/npre.2010.4325.1)

33.   Giurfa M. 2013 Cognition with few neurons: higher-order learning in insects. *Trends Neurosci.* **36**, 285–294. (doi:10.1016/j.tins.2012.12.011)

34.   Bazazi S, Bartumeus F, Hale JJ, Couzin ID. 2012 Intermittent motion in desert locusts: behavioural complexity in simple environments. *PLoS Comput. Biol.* **8**, e1002498. (doi:10.1371/journal.pcbi.1002498)

35.   Kien J, Altman JS. 1984 Descending interneurones from the brain and suboesophageal ganglia and their role in the control of locust behaviour. *J. Insect Physiol.* **30**, 59–72. (doi:10.1016/0022-1910(84)90108-2)

36.   Simmons PJ, Sztarker J, Rind FC. 2013 Looming detection by identified visual interneurons during larval development of the locust Locusta migratoria. *J. Exp. Biol.* **216**, 2266–2275. (doi:10.1242/jeb.083360)

37.   Dick PC, Gray JR. 2014 Spatiotemporal stimulus properties modulate responses to trajectory changes in a locust looming-sensitive pathway. *J. Neurophysiol.* **111**, 1736–1745. (doi:10.1152/jn.00499.2013)

38.   Matheson T, Rogers SM, Krapp HG. 2004 Plasticity in the visual system is correlated with a change in lifestyle of solitarious and gregarious locusts. *J. Neurophysiol.* **91**, 1–12. (doi:10.1152/jn.00795.2003)

39.   Moore RJD, Taylor GJ, Paulk AC, Pearson T, van Swinderen B, Srinivasan MV. 2014 FicTrac: a visual method for tracking spherical motion and generating fictive animal paths. *J. Neurosci. Methods* **225**, 106–119. (doi:10.1016/j.jneumeth.2014.01.010)

40.   Peirce J, Gray JR, Simpson S, MacAskill M, Höchenberger R, Sogo H, Kastman

E, Lindeløv JK. 2019 PsychoPy2: Experiments in behavior made easy. *Behav. Res. Methods* **51**, 195–203. (doi:10.3758/s13428-018-01193-y)

41. Gold JI, Shadlen MN. 2007 The neural basis of decision making. *Annu. Rev. Neurosci.* **30**, 535–574. (doi:10.1146/annurev.neuro.29.051605.113038)

42. Gabbiani F, Krapp HG, Laurent G. 1999 Computation of object approach by a wide-field, motion-sensitive neuron. *J. Neurosci.* **19**, 1122–1141. (doi:10.1523/JNEUROSCI.19-03-01122.1999)

43. Fotowat H, Gabbiani F. 2011 Collision detection as a model for sensory-motor integration. *Annu. Rev. Neurosci.* **34**, 1–19. (doi:10.1146/annurev-neuro-061010-113632)

44. Warren WH. 2021 Information Is Where You Find It: Perception as an Ecologically Well-Posed Problem. *Iperception* **12**, 20416695211000370. (doi:10.1177/20416695211000366)

45. Harpaz R, Nguyen MN, Bahl A, Engert F. 2021 Precise visuomotor transformations underlying collective behavior in larval zebrafish. *Nat. Commun.* **12**, 6578. (doi:10.1038/s41467-021-26748-0)

46. Fernández-Juricic E, Erichsen JT, Kacelnik A. 2004 Visual perception and social foraging in birds. *Trends Ecol. Evol.* **19**, 25–31. (doi:10.1016/j.tree.2003.10.003)

47. Dombrovski M, Kim A, Poussard L, Vaccari A, Acton S, Spillman E, Condron B, Yuan Q. 2019 A plastic visual pathway regulates cooperative behavior in drosophila larvae. *Curr. Biol.* **29**, 1866-1876.e5. (doi:10.1016/j.cub.2019.04.060)

48. Ballerini M *et al.* 2008 Interaction ruling animal collective behavior depends on topological rather than metric distance: evidence from a field study. *Proc Natl Acad Sci USA* **105**, 1232–1237. (doi:10.1073/pnas.0711437105)

49. Kappel JM *et al.* 2022 Visual recognition of social signals by a tectothalamic neural circuit. *Nature* (doi:10.1038/s41586-022-04925-5)

50. Slepian Z, Sundby K, Glier S, McDaniels J, Nystrom T, Mukherjee S, Acton ST, Condron B. 2015 Visual attraction in Drosophila larvae develops during a critical period and is modulated by crowding conditions. *J. Comp. Physiol. A Neuroethol. Sens. Neural Behav. Physiol.* **201**, 1019–1027. (doi:10.1007/s00359-015-1034-3)

51. Uvarov B. 1977 Grasshoppers and locusts. A handbook of general acridology. Volume 2. Behaviour, ecology, biogeography, population dynamics. *undefined*

52. Ellis PE, Ashall C. 1957 Field Studies on diurnal Behaviour, Movement and Aggregation in the Desert Locust (Schistocerca gregaria Forskål). *Field Studies on*

diurnal Behaviour, Movement and Aggregation in the Desert Locust (Schistocerca
gregaria Forskål).

53.   Warrant EJ. 2016 Matched filtering and the ecology of vision in insects. In *The*
ecology of animal senses (eds G von der Emde, E Warrant), pp. 143–167. Cham:
Springer International Publishing. (doi:10.1007/978-3-319-25492-0_6)

54.   Latty T, Trueblood JS. 2020 How do insects choose flowers? A review of multi-
attribute flower choice and decoy effects in flower-visiting insects. *J. Anim. Ecol.*
89, 2750–2762. (doi:10.1111/1365-2656.13347)

55.   Franks NR, Mallon EB, Bray HE, Hamilton MJ, Mischler TC. 2003 Strategies for
choosing between alternatives with different attributes: exemplified by house-
hunting ants. *Animal Behaviour* **65**, 215–223. (doi:10.1006/anbe.2002.2032)

56.   Sasaki T, Pratt SC. 2013 Ants learn to rely on more informative attributes during
decision-making. *Biol. Lett.* **9**, 20130667. (doi:10.1098/rsbl.2013.0667)

57.   Williamson R, Burns MD. 1982 Large neurones in locust neck connectives. *J.*
Comp. Physiol. **147**, 379–388. (doi:10.1007/BF00609672)

58.   Card GM. 2012 Escape behaviors in insects. *Curr. Opin. Neurobiol.* **22**, 180–186.
(doi:10.1016/j.conb.2011.12.009)

59.   Judge S, Rind F. 1997 The locust DCMD, a movement-detecting neurone tightly
tuned to collision trajectories. *J. Exp. Biol.* **200**, 2209–2216.
(doi:10.1242/jeb.200.16.2209)

60.   Schlotterer GR. 1977 Response of the locust descending movement detector
neuron to rapidly approaching and withdrawing visual stimuli. *Can. J. Zool.* **55**,
1372–1376. (doi:10.1139/z77-179)

61.   Pinter RB, Olberg RM, Abrams TW. 1982 Is the locust DCMD A looming
detector? *J. Exp. Biol.* **101**, 327–331. (doi:10.1242/jeb.101.1.327)

62.   Rind FC, Simmons PJ. 1992 Orthopteran DCMD neuron: a reevaluation of
responses to moving objects. I. Selective responses to approaching objects. *J.*
Neurophysiol. **68**, 1654–1666. (doi:10.1152/jn.1992.68.5.1654)

63.   Haberkern H, Jayaraman V. 2016 Studying small brains to understand the
building blocks of cognition. *Curr. Opin. Neurobiol.* **37**, 59–65.
(doi:10.1016/j.conb.2016.01.007)

64.   Mosqueiro TS, Huerta R. 2014 Computational models to understand decision
making and pattern recognition in the insect brain. *Curr. Opin. Insect Sci.* **6**, 80–
85. (doi:10.1016/j.cois.2014.10.005)

65.   Ratcliffe JM, Fenton MB, Shettleworth SJ. 2006 Behavioral flexibility positively
613   correlated with relative brain volume in predatory bats. *Brain Behav. Evol.* **67**,
614   165–176. (doi:10.1159/000090980)

615   66.   Shultz S, Dunbar RIM. 2010 Species differences in executive function correlate
616   with hippocampus volume and neocortex ratio across nonhuman primates. *J.*
617   *Comp. Psychol.* **124**, 252–260. (doi:10.1037/a0018894)

618   67.   Sol D, Timmermans S, Lefebvre L. 2002 Behavioural flexibility and invasion
619   success in birds. *Animal Behaviour* **63**, 495–502. (doi:10.1006/anbe.2001.1953)

620   68.   Chittka L, Giurfa M, Riffell JA. 2019 Editorial: the mechanisms of insect
621   cognition. *Front. Psychol.* **10**, 2751. (doi:10.3389/fpsyg.2019.02751)

(a)(i)

(a)(ii)

(b)(i)

Python work station

(b)(ii)

High-speed video

Locust

Track-ball

Mouse sensor

Screen 1

Screen 2

(a)

(b)

(c)