Water wisteria genome reveals environmental adaptation and heterophylly

regulation in amphibious plants

Gaojie Li[1,2], Xuyao Zhao[1,2], Jingjing Yang[1,2], Shiqi Hu[3], Jathish Ponnu[4],

Seisuke Kimura[5,6], Inhwan Hwang[7], Keiko U Torii[8,9,10], Hongwei Hou[1,2,]*

[1] The State Key Laboratory of Freshwater Ecology and Biotechnology, Institute of

Hydrobiology, Chinese Academy of Sciences, Wuhan, Hubei, 430072, China

[2] University of Chinese Academy of Sciences, Beijing 100049, China

[3] Zhejiang Marine Development Research Institute, Zhoushan, 316021, China

[4] Joseph Gottlieb Kölreuter Institute for Plant Sciences, Karlsruhe Institute of

Technology, 76131 Karlsruhe, Germany

[5] Faculty of Life Sciences, Kyoto Sangyo University, Kamigamo-motoyama, Kita-ku,

Kyoto-shi, Kyoto 603-8555, Japan

[6] Center for Plant Sciences, Kyoto Sangyo University, Kamigamo-motoyama, Kita-ku,

Kyoto-shi, Kyoto 603-8555, Japan

[7] Department of Life Science, Pohang University of Science and Technology, Pohang

37673, Republic of Korea

[8] Institute of Transformative Biomolecules (WPI-ITbM), Nagoya University, Chikusa,

Nagoya, Aichi, 464-8601, Japan

[9] Howard Hughes Medical Institute, The University of Texas at Austin, Austin, TX

78712, USA

[10] Department of Molecular Biosciences, The University of Texas at Austin, Austin,

TX 78712, USA

*Corresponding Author: Tel: (+86-27) 68788691; Email: houhw@ihb.ac.cn

Abstract

Heterophylly is a phenomenon in which an individual plant dramatically changes

its leaf shape in response to the surrounding environment. *Hygrophila difformis*

(Acanthaceae), also known as water wisteria, has recently emerged as a model plant to

study heterophylly because of its striking leaf shape variation in response to various

ecological factors. Under submerged conditions, *H. difformis* develops complex

leaves and in terrestrial conditions it develops simple leaves. Here, we sequenced and

assembled the chromosome-level genome of triploid *H. difformis* (scaffold N50:

60.43 Mb, genome size: 871.92 Mb), which reveals 36,099 predicted protein-coding

genes distributed over 15 pseudochromosomes. *H. difformis* diverged from its

relatives during the Oligocene climate-change period and expanded the gene families

related to its amphibious lifestyle. Genes involved in environmental stimuli, leaf

development, and other pathways are differentially expressed in submerged and

terrestrial conditions, possibly modulating morphological and physiological

acclimation to changing environments. We confirmed that auxin plays a role in the

heterophylly of *H. difformis*. Finally, we discovered candidate genes that respond to

different environmental conditions and elucidated the role of *LATE MERISTEM*

IDENTITY 1 (*LMI1*) in heterophylly. Our study establishes *H. difformis* as a model for

studying the interconnections between ecological adaptation and plant morphological

features.

Keywords: Heterophylly, Environmental adaptation, Genome, *Hygrophila difformis*,

Phenotypic plasticity

Introduction

Amphibious plants can grow both in water and on land, switching their

morphology and physiology according to various environmental cues (Agarie et al.,

1997). While genomic data are available for several aquatic plants (Lan et al., 2017;

Xue et al., 2020; Lu et al., 2021; Abramson et al., 2022; Shi et al., 2022), amphibious

plants remain largely unexplored at the genomic level. Climate change poses new

challenges for plant growth and survival. One of the strategies that plants use to cope

with these challenges is phenotypic plasticity (Alpert and Simms, 2002), the ability to

change their phenotypes in response to environmental cues. Phenotypic plasticity

connects plant evolution, ecology and molecular biology and helps to predict plant

responses to novel environments under climate change (Nicotra et al., 2010; Arnold et

al., 2019; Stotz et al., 2021). A striking example of phenotypic plasticity is

heterophylly, the ability of a plant to alter its leaf shape depending on the environment

(van Veen and Sasidharan, 2021). Heterophylly is common in amphibious plants,

which can grow both in land and water (Nakayama et al., 2014; Li et al., 2017; Kim et

al., 2018; Koga et al., 2021). Heterophyllous plants can adjust their leaf morphology

to various environmental factors such as submergence (Li et(Han et al., 2021). al.,

2017), light (Momokawa et al., 2011; Nakayama et al., 2014), temperature

(Nakayama and Kimura, 2015), and CO_2 (Titus and Gary Sullivan, 2001), bridging

plant ecology, evolution, and molecular biology, might enable to predict plant

responses to novel environments posed by climate change. Whereas studies on model

plants, like *Arabidopsis thaliana*, have greatly contributed to identifying genes and

processes regulating plant phenotypes, we need a suitable model system on

phenotypic plasticity to understand the relationships of environmental factors on plant

phenotypes. Interestingly, heterophyllous plants across diverse taxa often show similar

leaf variations to adapt to the amphibious environment, making them potential plant

models for convergent evolution and environmental adaptation.

Recently, several mechanisms regulating heterophylly have been suggested in a

few amphibious species. For example, ethylene and abscisic acid (ABA) control genes

that regulate adaxial–abaxial leaf polarity and induce heterophylly in *Ranunculus*

trichophyllus (Ranunculaceae) (Kim et al., 2018). Similarly, genes that affect cell

elongation and division contribute to leaf differentiation in *Callitriche palustris*

(Plantaginaceae) (Koga et al., 2021). In *Rorippa aquatica* (Brassicaceae),

KNOTTED1-LIKE HOMEOBOX (*KNOX1*) genes modulate the levels of endogenous

phytohormones gibberellic acid (GA) and cytokinin to influence

temperature-dependent heterophylly (Nakayama et al., 2014). However, despite the

availability of transcriptomic data for several heterophyllous plants (van Veen et al.,

2013; He et al., 2018; Han et al., 2021; Koga et al., 2021; Horiguchi et al., 2023), the

molecular mechanisms underlying heterophylly are still largely unknown.

Hygrophila difformis (Acanthaceae), also known as water wisteria, is an

amphibious plant that exhibits heterophylly by developing deeply lobed, complex

leaves under submergence and simple leaves in terrestrial conditions (Fig. 1A). *H.*

difformis has recently gained the attention of researchers as a promising new model

for heterophylly studies, due to its beneficial traits such as easy gene transformation,

fast propagation, and conspicuous leaf shape change in response to various

environmental stimuli (Li et al., 2017; Li et al., 2020). It also allows us to explicitly

address the interaction between environmental responses, morphological features, and

variation in gene expression in an ecological context (Li et al., 2021). Previously, we

showed that *SHOOT MERISTEMLESS* (*STM*) participates in the heterophylly

regulation of *H. difformis* (Li et al., 2022), but key genes and molecular mechanisms

underlying this phenomenon are still unclear. The genus *Hygrophila* comprises nearly

90 species, many of which are amphibious and display phenotypic plasticity (Fig. S1).

H. difformis, however, displays the most conspicuous form of heterophylly.

Hygrophila includes species that are grown as ornamental plants (Karatas et al., 2013),

used as medicinal herbs (Ingale et al., 2013), and even serve as models for

photosynthetic studies (Botts et al., 1990; Horiguchi et al., 2021). However, at present,

there are no genomic resources available for this genus.

Here, we report a chromosome-scale genome of *H. difformis*, assembled by

integrating PacBio, Illumina, and High-throughput Chromosome Conformation

Capture (Hi-C) sequencing technologies. We performed genome and transcriptome

analyses to reveal the molecular mechanisms underlying heterophylly in response to

the amphibious environment. We also identified and validated *LATE MERISTEM*

IDENTITY 1 (*LMI1*) as a candidate gene for leaf shape regulation. Our study provides

the first genome and transcriptome for the genus *Hygrophila*, establishing *H. difformis*

as a novel ecological model for investigating environmental impacts on plant

phenotypes. This study offers a comprehensive framework and functional data for

future genetic, genomic, and molecular studies of plant phenotypic plasticity. We

believe that our work will attract more attentions to the plant phenotypic plasticity and

environmental adaptation, and use *H. difformis* as an emerging model for plant

Eco/Evo/Devo studies.

Results

Genome sequencing, assembly, and annotation

We previously reported that *H. difformis* reproduces through vegetative ways (Li

et al., 2017), may be infertile and likely a triploid. Here we generated, 142.27 Gb

sequencing data (genome coverage: 150 ×), with 52,199,178,926 *k*-mers and a *k*-mer

depth of 54 (supplementary table S1, Fig. S2), which estimated a genome size of

913.38 Mb. The *K*-mer analysis confirmed that *H. difformis* is triploid (Fig. S2), as its

two peaks have triple values (*K*-mer depth =18 and 54) (Ranallo-Benavidez et al.,

2020). We also performed karyotype analysis and cytogenetic studies, which showed

the presence of 45 chromosomes (2n = 3X = 45) (Fig. S3). We further evaluated the

genome size by flow cytometry, and the detected genome size was 0.89 Gb (Fig. S4),

slightly smaller than the *k*-mer estimated genome size (supplementary table S1). The

genome of *H. difformis* was sequenced, separated, and assembled to the haploid

genome using the PacBio Sequel platform (Zhou et al., 2021). The initial genome

assembly consisted of 2,752 contigs covering 914.25 Mb. Using the Hi-C seq

platform (Burton et al., 2013; Durand et al., 2016), we obtained 15

pseudochromosome sequences covering 871.92 Mb (95.37%), with a Scaffold N50 of

60.43 Mb (supplementary table S2-4). BUSCO (Benchmarking Universal

Single-Copy Orthologs) (Simao et al., 2015) analysis suggested that 95.50% of the

total genes were recovered (supplementary table S5). Based on the Hi-C contig

contact matrix heatmap and genome features (Fig. 1B, C), we verified that the

clustering, ordering, and orientation of the contigs were successful.

We identified 36,099 protein-coding gene models (supplementary table S6)

according to homology-based and *de novo* predictions and 90.62% (32,714) of them

could be mapped with functional annotations using public databases (supplementary

table S7). We also performed BUSCO analysis using the predicted protein sequences

to assess the completeness of our annotation, which indicates that 94.40% of the total

genes were recovered (supplementary table S5). These results suggest a successful

assembly and annotation of major genic regions in the *H. difformis* genome.

Repetitive DNA sequences are a major component of eukaryotic genomes,

reflecting their organization and evolution (Mehrotra and Goyal, 2014). Using a

combination of homology-based searches and *de novo* prediction, we identified

repetitive sequences accounting for 74.92% of the genome (supplementary table S8),

which is higher compared to other reported related terrestrial species within

Acanthaceae family members, *Andrographis paniculata* (53.3%) (Sun et al., 2019),

Strobilanthes cusia (74.4%) (Xu et al., 2020), *Avicennia marina* (48.8%) (Ma et al.,

2022), and the reported related aquatic species *Utricularia gibba* (42.18%) (Lan et al.,

2017) within the order Lamiales. Among the repeat sequences in *H. difformis* genome,

long terminal repeats (LTRs) were dominant (62.39%) (supplementary table S8),

which is also higher compared to *U. gibba* (12.44%), *A. marina* (24.92%), *A.*

paniculata (38.54%), and *S. cusia* (51.32%). These results suggest that the genome of

H. difformis is relatively more repetitive than the genomes of other reported related

species.

Comparative genomics and genome evolution

To investigate the evolution of *H. difformis*, we compared its genome to fourteen

representative species of diverse evolutionary status, including one monocot, *Oryza*

sativa (Poaceae), and thirteen eudicots. We identified a total of 14,656 ortholog

groups, of which 1,362 were unique to *H. difformis*. (Supplementary table S9). We

identified 210 single-copy orthologous gene sets among the 15 genomes and

subsequently constructed a phylogenetic tree based on the molecular clock (Kumar et

al., 2017). The phylogenetic placement of these species is consistent with references

therein (Kordyum and Mosyakin, 2020). Among the genomes of other Lamiales, we

estimated the divergence times of around 69.8 Mya (Million years ago) between *O.*

europaea and the other six species, 57.8 Mya between *B. hygrometrica* and the

remaining five species, and 30.6 Mya between *A. paniculata* and *H. difformis* (Fig.

2A). The differentiation time of *H. difformis* was approximately 25.4–37.9 Mya,

coinciding with the evolution and diversification of East Asian Flora (EAF) during the

Oligocene (33.9 to 23.0 Mya) (Ma et al., 2019; O'Brien et al., 2020).

We then examined gene expansion and contraction in the phylogenetic analysis.

We found that *H. difformis* had more gene expansion than its related species (Fig. 2B).

A total of 5,089 ortholog groups were expanded in the *H. difformis* lineage, while

1,390 ortholog groups were contracted. GO pathway enrichment analysis showed that

the significantly expanded (Q value < 0.01) genes were enriched in traits related to

environmental adaptation, such as "tropism" (GO:0009606), "response to freezing"

(GO: 0050826), and "response to blue light" (GO:0009637), and also in energy

metabolisms and physiological processes, such as "carbohydrate metabolic process"

(GO:0005975), "oxidation-reduction process" (GO:0055114) and "regulation of

photosynthesis" (GO:0010109). We also detected expanded genes in the "hormone

metabolic process" (GO:0042445) and "regulation of hormone levels" (GO:0010817),

which may regulate the hormone levels and homeostasis in *H. difformis*

(supplementary table S10). The significantly contracted (Q value < 0.01) genes were

enriched in "carbon-oxygen lyase activity" (GO:0016838), "terpene synthase activity"

(GO:0010333), and some other pathways (supplementary table S11).

Whole genome duplication (WGD) events are major evolutionary forces and

have been associated with macroevolutionary changes and evolutionary innovations

that happened relatively frequently in plants (Ren et al., 2022). Many genera of

Lamiales underwent additional WGD events after the gamma event (Godden et al.,

2019). To study the evolution of *H. difformis* within Acanthaceae, we characterized

the synonymous substitution divergence (*Ks*) of each collinear gene pair within a

genome or between genomes for WGD analysis between *H. difformis* and other

reported species (Fig. 2C, Fig. S5). The *Ks* distribution of paralogs showed that *H.*

difformis, grape (*Vitis vinifera*, the outgroup of Acanthaceae) and *A. paniculata* have

only one peak at *Ks* = ~1.19, ~1.28 and ~1.41, respectively (Fig. 2C). It should be

mentioned that *H. difformis* likely experienced a recent WGD event that led to a

triploid karyotype, which was not captured in this analysis. The orthologous

distribution peaks for *H. difformis–A. paniculata*, *H. difformis–V. vinifera* also

correspond to *Ks* values of ~0.72 and ~1.53, respectively (Fig. 2C). These results

showed that the *H. difformis* genome may only experience the ancient whole-genome

triplication (γ), which is similar to its close-related species *A. paniculata* and *S. cusia*

within Acanthaceae (Sun et al., 2019; Xu et al., 2020). We revealed that the *Ks*

distribution peak within *V. vinifera* was less than the *Ks* peak of *H. difformis-V.*

vinifera, which indicated that the γ event occurred after their differentiation (Fig. 2C).

We also analyzed the *Ks* distribution of recent reported Acanthaceae species *A.*

marina and *S. cusia* (Fig. S5). In contrast, *Ks* distribution peaks of *H. difformis–S.*

cusia and *H. difformis–A. paniculata* were less than the *Ks* peak within *H. difformis*, *S.*

cusia and *A. paniculata* (Fig. 2C, Fig. S5), which indicated that these species

underwent a common γ event. In addition, *A. marina* (2n= 64) experienced a special

recent WGD (*Ks* = ~0.49) (Fig. S5), which is similar to the recently published

research (Ma et al., 2022).

Morphological, physiological, and molecular adaptation of *H. difformis* to the

terrestrial or submerged environment

Heterophyllous plants grown in submerged conditions often develop filiform,

dissected or linear leaves that lack cuticle and stomata, while the terrestrial leaves are

thicker, with differentiated mesophylls, and are cutinized with stomata (Koga et al.,

2021; Li et al., 2021). We found that leaf area, dissection index (DI), and epidermal

cell complexity of *H. difformis* were significantly increased in submerged conditions

(Student's *t*-test, $P < 0.01$), while stomatal density, vein density, and chlorophyll

content were significantly increased in terrestrial conditions (Student's *t*-test, $P < 0.01$)

(Fig. 3A, S6), which is similar to other heterophyllous plants (Kim et al., 2018; Koga

et al., 2021).

To elucidate the molecular mechanisms underlying the phenotypic plasticity in *H.*

difformis, we performed RNA-seq analysis on both submerged and terrestrial shoots

using annotations based on our genomic data. A total of 6,226 differentially expressed

genes (DEGs) (false discovery rate [FDR] < 0.05 and expression-level $\log_2$

[fold-change] > 1) were identified, with 4,100 upregulated genes and 2,126

downregulated genes in terrestrial shoots. We then analyzed the DEGs that are

involved in environmental sensing or response (Fig. 3B). The thermosensor gene

EARLY FLOWERING 3 (*ELF3*) (Jung et al., 2020), a negative regulator of

thermomorphogenesis, is downregulated in terrestrial conditions (Fig. 3B). *ELF3* has

been shown to function antagonistically with *PHYTOCHROME-INTERACTING*

FACTOR4 (*PIF4*) in thermomorphogenesis (Raschke et al., 2015), and we also

observed an inverse expression pattern of *PIF4* in *H. difformis* grown aerially. The

proton sensor gene *SLAC1 HOMOLOGUE 3* (*SLAH3*) (Lehmann et al., 2021), the

mechanical stimuli sensor gene *MSCS-LIKE 10* (*MSL10*) (Tran et al., 2021), and the

PLASMA MEMBRANE INTRINSIC PROTEIN (*PIP*) aquaporin genes (Yaneff et al.,

2015) were also upregulated in terrestrial conditions.

Morphological changes in leaf shape are the most obvious characteristics of

heterophyllous plants. The antagonistic interaction of *KNOX1* and *BELL1-LIKE*

HOMEOBOX (*BLH*) genes modulate leaf development, with *KNOX1* overexpression

enhancing leaf lobing (Hareven et al., 1996) and the *BLH* overexpression reducing

leaf serration (Kumar et al., 2007; Kimura et al., 2008). We observed increased

expression of *BLH* genes in terrestrial shoots, which may underlie the leaf

simplification observed in terrestrial conditions. *TfTCP13* (from *Torenia fournieri*)

overexpression in *Arabidopsis thaliana* leads to narrower leaves (Zhang et al., 2021),

and we detected increased expression of *TCP13* in submerged shoots of *H. difformis*.

TCP14 expression was reduced in the terrestrial leaves of *H. difformis*, mimicking the

broad, short leaf phenotype induced by the dominant negative construct

TCP14::SRDX in *A. thaliana* (Kieffer et al., 2011). *GROWTH-REGULATING*

FACTORs (*GRFs*) control leaf size in *A. thaliana* (Debernardi et al., 2014) and *GRF5*

was elevated in the submerged shoots of *H. difformis*, which may account for the

increased leaf size in submerged conditions (Fig. 3A, B). Genes involved in

chloroplast development and chlorophyll biosynthesis are also elevated in terrestrial

shoots (Fig. 3B, S6).

MUTE, *FAMA*, and *SLOW ANION CHANNEL-ASSOCIATED 1* (*SLAC1*), which

regulate stomatal development and movement, are upregulated in terrestrial shoots,

possibly facilitating stomatal formation in terrestrial leaves. Although *EPIDERMAL*

PATTERNING FACTOR-LIKE9 (*EPFL9/STOMAGEN*) is a positive regulator of the

stomatal development (Hunt et al., 2010), it is often absent in aquatic plants, including

Z. marina, *S. polyrhiza*, *Nymphaea colorata*, and *Wolffia australiana* (Park et al.,

2021). However, we identified its putative homologous gene (Hdi026079) in *H.*

difformis. Interestingly, its expression was not detected in either terrestrial or

submerged shoots, suggesting this gene might be a pseudogene or it acquired a

different function in *H. difformis*.

Genes involved in vascular development, lignin biosynthesis, and wax

biosynthesis are upregulated in terrestrial shoots (Fig. 3B), which may enhance

terrestrial acclimation. We also observed increased expression of genes responsive to

oxidative stress in the terrestrial environment, which is in agreement with our finding

that reactive oxygen species (ROS) accumulate in terrestrial leaves of *H. difformis*

(Fig. 3B, S7).

Roles of auxin and ABA in terrestrial and submerged acclimation

The role of phytohormones in the environmental control of morphogenesis has

been reviewed recently (Nakayama et al., 2017; Li et al., 2019). We detected a large

number of DEGs involved in the "response to hormone", especially related to the

"response to auxin" and "response to abscisic acid" (Fig. 3E). Our previous study

showed that exogenous ABA can induce leaf simplification in *H. difformis* (Li et al.,

2017). Here, we observed increased expression of genes related to ABA biosynthesis

(e.g., *NINE-CIS-EPOXYCAROTENOID DIOXYGENASE* [*NCED*], *ABA*

DEFICIENT1 [*ABA1*], and *ALDEHYDE OXIDASE* [*AAO*]) and ABA signaling-related

genes in terrestrial shoots, consistent with increased ABA content observed in

terrestrial shoots (Fig. 3C, F). *CYP707A*, which encodes an ABA-degradation enzyme,

was reduced in terrestrial conditions, as were the *PYRABACTIN RESISTANCE*

1-LIKE (*PYL*) genes. The expression pattern of these genes was consistent with their

orthologs of *C. palustris* (Koga et al., 2021), suggesting similar ABA pathways

regulate heterophylly in different amphibious species.

We also observed increased expression of auxin biosynthesis gene *YUCCA* (*YUC*)

and the *YUC*-related *NAKED PINS IN YUC MUTANTS 1* (*NPY1*) in submerged shoots,

along with increased auxin contents (Fig. 3D, F). *INDOLE-3-ACETIC ACID* (*IAA*)-

INDUCIBLE and *AUXIN RESPONSE FACTOR 10* (*ARF10*) were also elevated in

submerged conditions. Interestingly, the auxin influx transporter *AUXIN RESISTANT*

1 (*AUX1*) was upregulated in submerged conditions, while the auxin efflux carriers

PIN-FORMED (*PIN*) and *PIN-LIKES* (*PILS*) were downregulated.

To further examine the role of auxin transport in *H. difformis*, we treated

submerged plants with the polar auxin transport inhibitor naphthylphthalamic acid

(NPA), which caused a gradual reduction in leaf dissection and eventually the

formation of oval leaves with smooth margins (Fig. 3G). Previous studies have

reported the roles of ABA, ethylene (Kuwabara et al., 2003), GA, and cytokinin

(Nakayama et al., 2014) in heterophylly. Here, we demonstrate for the first time that

auxin metabolism and transport are also implicated in the regulation of heterophylly

in *H. difformis*.

Expression patterns of DEGs under diverse ecological factors

A previous study revealed expression patterns of candidate genes in the

heterophyllous plant *C. palustris* under different phytohormone treatments (Koga et

al., 2021). However, the response of heterophyllous plants to more natural and diverse

environmental signals remain unclear. To address this, *H. difformis*, plants were

grown in four sets of ecological factors (submerged/terrestrial, normal/low light

density, normal/low temperature, and high/normal humidity) and sampled for

RNA-seq (Fig. 4A). We also measured morphological and physiological

characteristics between these groups (Fig. S8), showing similarities and differences

affected by diverse ecological factors. Our transcriptomic analysis identified a total of

6,226 DEGs in the submerged/terrestrial group (Fig. 3), 9,213 DEGs in the

normal/low light density group, 2,851 DEGs in the normal/low-temperature group;

and 4,065 DEGs in the high/normal humidity group (Fig. 4B, 4C). A total of 180

genes were identified as upregulated DEGs and 119 genes were identified as

downregulated DEGs in all four comparisons (supplementary table S12-13). *NCED3*,

ETHYLENE-INSENSITIVE 3 (*EIN3*), *CYTOKININ OXIDASE 5* (*CKX5*), and *GA*

INSENSITIVE DWARF1A (*GID1A*) genes, which are related to phytohormone

pathways, were among those genes elevated in all groups (Fig. 4D). Genes related to

antioxidants were also upregulated, while *NPY1* was downregulated in all groups (Fig.

4E). A previous study showed that expression and functional divergence of *SUGARS*

WILL EVENTUALLY BE EXPORTED TRANSPORTERS (*SWEET*) genes contribute to

the specific developmental strategy of *Euryale ferox* leaves (Wu et al., 2022). Here,

we observed that three putative orthologs of *SWEET* were upregulated in these groups,

implying their potential roles in heterophylly. We also identified nineteen upregulated

TFs and eight downregulated TFs (Fig. 4F) from these common DEGs, which may

regulate the acclimation of *H. difformis* to different ecological conditions.

To further pinpoint key genes involved in heterophylly, co-expression analysis

was conducted on a total of 15,295 DEGs (all DEGs that were changed in at least one

group) and genes from 15 samples were filtered and clustered into ten modules using

weighted gene co-expression network analysis (WGCNA) (Fig. S9). We focused on

the yellow module which was associated with gene expression patterns and leaf

complexity (Fig. S9, S10C). We identified three leaf development-related genes,

putative orthologs of *LATE MERISTEM IDENTITY1* (*LMI1A*, *LMI1B*, and *LMI1C*),

which were positively co-expressed with three auxin-related genes (*NPY1*, *AUX1*, and

ARF16), two *GA-STIMULATED IN ARABIDOPSIS* (*GASA7* and *GASA14*) genes, the

ethylene synthesis gene *ACC SYNTHASE1* (*ACS1*), and the photosynthesis-related

gene *LIGHT-HARVESTING COMPLEX B7* (*LHCB7*) (Fig. 4G). It is known that the

phenomenon of heterophylly involves morphological, anatomical and physiological

adaptation (Horiguchi et al., 2019; Li et al., 2021). The candidate genes identified in

our work regulating leaf formation, along with phytohormone pathways and

physiological adaptation to environmental changes, might influence the heterophylly

of *H. difformis*.

HdLMI1 genes regulate the leaf shape of *H. difformis* in submerged conditions

LMI1 and its putative orthologs have been studied in several terrestrial species

and shown to have conserved roles in leaf shape diversity (Vlad et al., 2014; Andres et

al., 2017). Here, we identified three putative paralogous *LMI1* orthologs in *H.*

difformis, and examined their expression in different samples (Fig. 5A). These genes

were highly expressed in humid and aquatic samples (A1-A3, and C1-C3), which had

complex leaves (Fig. S10). In contrast, their expression was downregulated in

simple-leafed groups (Fig. 5A).

To further examine the roles of *HdLMI1s* in regulating leaf development in *H.*

difformis, we knocked down these genes using virus-induced gene silencing (VIGS)

(Zhang et al., 2020) with the Tobacco rattle virus (TRV) constructs *TRV2-HdLMI1A*,

TRV2-HdLMI1B and *TRV2-HdLMI1C*. Compared to mock plants, the

*TRV2-HdLM*I1A, *TRV2-HdLMI1B*, and *TRV2-HdLMI1C* transgenic plants grown in

submerged conditions were similar to each other, developing crinkled, complete, or

shallowly serrated leaf margin (Fig. 5B). DI indicated that the mock plants had

significantly higher leaf complexity than *TRV2-HdLMI1A*, *TRV2-HdLMI1B*, and

TRV2-HdLMI1C transgenic plants (Dunnett's test, $P < 0.01$) (Fig. 5C). In addition to

their morphological features, these plants showed a significantly reduced expression

of all three *HdLMI1s* compared to mock (Dunnett's test, $P < 0.01$) (Fig. 5D). A

previous study showed that orthologs of *LMI1* in *Medicago truncatula* cooperatively

regulate leaf margin development (Wang et al., 2021). These results confirmed our

hypothesis that *HdLMI1s* are important regulators of heterophylly in *H. difformis*.

Discussion

Our knowledge of genetic regulatory networks that contribute to the variation of

plant phenotypes has been advanced by the use of the latest technologies, but the

mechanisms of phenotypic plasticity still remain elusive mostly due to the scarcity of

suitable models (Hoffmann and Sgro, 2011; van Veen and Sasidharan, 2021). Our

published genome and transcriptome of *H. difformis* provide the resources for further

studies to expand the knowledge on adaptations that enable this plant to thrive in a

highly fluctuating environment. The genome of *H. difformis* also offers a chance to

investigate convergent evolution among heterophyllous species.

Many aquatic species became polyploid during speciation (Albert and Renner,

2020; Lu et al., 2021), and polyploidy was verified as a factor in plants expanding

their ranges or invading new environments as polyploid progenies often possess novel

traits (Cheng et al., 2021). Polyploidy and hybridization in aquatic or amphibious

plants are also associated with complex evolutionary histories (Les and Philbrick,

1993; Prancl et al., 2014). For example, water caltrop (*Trapa natans*) (Lu et al., 2021)

and *C. palustris* are tetraploid (Koga et al., 2021), while *Cabomba caroliniana* has

polyploid ($3\times$, $6\times$, $8\times$) and aneuploid variation between individuals (McCracken et al.,

2013). The number of chromosomes in *Hygrophila* species is varied, such as in *H.*

spinosa ($2n = 32$), *H. augustifolia* ($2n = 44$) and *H. balsamica* ($2n = 34$)

(Govindarajan and Subramanian, 1983). We previously reported that *H. difformis*

always reproduces through vegetative propagation (Li et al., 2017) and does not seem

to produce viable seeds. Our genome survey and cytogenetic studies revealed 45

chromosomes, and further analyses suggest that the sequenced *H. difformis* is triploid

(2n = 3x = 45). The family Lamiaceae exhibit widespread but asymmetrical gene

duplication patterns and ancient polyploidy (Godden et al., 2019). For example,

Paulownia tomentosa (Paulowniaceae) had 3 inferred WGDs, while there were 7-18

WGDs in Nepetoideae (Godden et al., 2019). Previous studies reported that *A.*

paniculata (2n = 48) and *S. cusia* (2n = 32) only experienced one γ event (Sun et al.,

2019; Xu et al., 2020). However, *A. marina* (2n = 64) was reported to have two

WGDs, and the recent WGD was attributed to a novel specific adaptation to global

warming that occurred during the palaeocene–Eocene maximum (Ma et al., 2022).

Our analysis confirmed the recent WGD in *A. marina*, but did not detect the common

γ event, which could be due to data or methodological differences. We also observed

that *H. difformis* possessed the same γ event as *A. paniculata* and *S. cusia*, and lacked

any additional WGD during its evolutionary history (Fig. 2E). Polyploidy is widely

regarded as a common speciation strategy that has pronounced implications for plant

evolution and ecology (Van De Peer et al., 2017). *H. difformis* appears to be a natural

triploid. There are only four Acanthaceae species with genome information reported

so far (Ma et al., 2022), and thus more data about species within the *Hygrophila* genus

is essential for evolutionary studies. Future studies on comparing subgenomes by the

haplotype-resolved genome assembly will also be helpful in this regard.

Our comparative genome data showed that the differentiation time of *H.*

difformis is about 30.6 Mya (Fig. 2A), coinciding with the evolution and

diversification of East Asian flora (EAF) during the Oligocene (33.9 to 23.0 Mya). It

was recorded that the decline of atmospheric CO_2 and global cooling culminated in

the Early Oligocene Aridification event at 31 Mya (Wasiljeff et al., 2022), which is

close to the differentiation time of *H. difformis*. We found significantly expanded (Q

value < 0.01) genes of *H. difformis* were enriched in "response to freezing",

"regulation of photosynthesis", and "carbohydrate metabolic process", which may

reflect the environmental response, physiological plasticity and metabolic adaptations

to the changing environments (supplementary table S10). In addition, the formation of

the Asian monsoon system in the Oligocene influenced the plant evolution and

diversification (Ma et al., 2019). The seasonality of rainfall increased progressively,

achieving modern monsoon-like wet summers and dry winters, by the early Oligocene

(Chen and Su, 2020). Therefore, plant species must have adapted to the seasonality of

rainfall and drought. For example, the habitat diversification (terrestrial and

aquatic/marshland) within *Pogostemon* (Lamiaceae) is related to adaptations to

seasonally changing climate that was triggered by the Asian monsoon in southern and

southeast Asia (Yao et al., 2016). It is known that monsoon systems involved in

seasonal rain and fluctuated water levels, and the water level change was regarded as

the driver of heterophylly in many amphibious species (Wanke, 2011; Nakayama et al.,

2014). *Hygrophila* species are widely spread in East Asia, South America, and

Southern Africa (https://www.gbif.org/species/9795574), and many of which are

amphibious plants (Fig. S1). However, due to lack of enough genome and fossil data

of *Hygrophila*, it would be premature to conclude that the amphibious habitat and

phenotypic plasticity of *H. difformis* are the innovation for climate change that

happened in Oligocene.

The leaflet development in *C. hirsuta* depends on *REDUCED COMPLEXITY*

(*RCO*), a homolog of *LMI1* that represses leaf flanks growth and promotes leaflets

formation (Vlad et al., 2014). *RCO* originated from gene duplication in the

Brassicaceae family and was lost in *A. thaliana*, leading to leaf simplification (Vlad et

al., 2014). However, *LMI1* has diverse functions in different species. For instance,

LMI1 regulates the formation of simple serrated leaves in *A. thaliana*. The *lmi1*

mutation resulted in decreased leaf serration but formed leaflets in *A. thaliana* (Saddic

et al., 2006), leaf serrations in *M. truncatula* (Wang et al., 2021), and rapeseed

(*Brassica napus*) (Hu et al., 2018), and leaf lobes in cotton (*Gossypium hirsutum*)

(Andres et al., 2017). To assess the function of *LMI1* in heterophylly, we silenced

HdLMI1 genes by VIGS and observed reduced leaf complexity in *H. difformis*

transgenic plants (Fig. 5). However, the three *HdLMI1* genes may have

cross-suppressed together by the TRV constructs due to their sequence similarities,

which points to functional redundancies among the *HdLMI1* genes. *In M. truncatula*,

two *LMI1* paralogous are separately expressed in leaf margin tooth and sinus, and

cooperatively regulate the leaf serrations (Wang et al., 2021). Thus, it is necessary to

distinguish the effects of silencing each *HdLMI1* gene via CRISPR/Cas9-mediated

genetic modifications. Moreover, heterophylly may involve epigenetic regulation such

as non-coding RNAs, DNA methylation, nucleosome assembly (Nakayama et al.,

2014; Li et al., 2021) etc., and more advanced analyses such as target mimicry, single-

cell sequencing, and CRISPR/Cas9 genetic modification are warranted in future

studies.

Auxin is a key phytohormone that orchestrates development and thus participates

in plant evolution (Moon and Hake, 2011; Finet and Jaillais, 2012). Auxin transport is

essential for the initiation of leaf primordia and the formation of leaf margins (Shani

et al., 2006). In Arabidopsis, auxin transport disruption leads to the absence of

serration in leaves, whereas *PIN1* localization determines the local auxin maxima at

the tips of emerging serrations (Hay et al., 2006). In compound-leaved species auxin

transport perturbation causes a dramatic simplification of leaf morphology (DeMason

and Chawla, 2004; Wang et al., 2005). Although auxin accumulation has been

detected in some waterlogged plants (Qi et al., 2019), its role in

heterophylly-induction of amphibious plants remains elusive. Previous studies

measured various phytohormones in *C. palustris* under submerged/terrestrial

conditions (Koga et al., 2021) , but they focused on the importance of ABA, GA and

did not analyze the role of auxin in detail. Here, we show that auxin levels are

significantly increased under submergence, and the genes involved in auxin

metabolism, transport and response (e.g., *NPY1*, *AUX1*, *PIN1*) are differentially

expressed (Fig. 3, 4). Moreover, we demonstrate that the auxin transport inhibition

abolishes heterophylly in *H. difformis*. The interplay between of auxin maxima and

KNOX1 genes underlies the evolution of serrations in *A. thaliana* and leaflets in *C.*

hirsuta (Canales et al., 2010).The *LMI1*-like and *KNOX1* genes also jointly control

leaf development in both cotton and *C. hirsuta* (Chang et al., 2019; Wang et al., 2022).

A recent study revealed that *LMI1* of *M. truncatula* regulates leaf margin development

by activating auxin transporter genes (Wang et al., 2021). Thus, putative orthologs of

LMI1, *KNOX1,* and the auxin transport genes may cooperate in *H. difformis*

heterophylly regulation. In addition, *PIF*-mediated auxin metabolism regulates plant

adaptation to various environmental factors (Sun et al., 2013; de Wit et al., 2014;

Burko et al., 2022). In this study, we identified *PIFs* among the DEGs involved in

heterophylly. Future research may establish *PIFs* as network hubs integrating various

environmental factors and developmental signals in the heterophylly of *H. difformis*.

We identified numerous DEGs associated with environmental stimuli (Fig. 3B),

which form complex networks with plant growth genes. For instance, ELF3 is a

thermosensor that regulates rhythmicity and temperature responsiveness (Jung et al.,

2020). Similarly, PIF4 directly activates the auxin biosynthetic gene *YUCCA8* (*YUC8*)

to control hypocotyl elongation (Sun et al., 2012), while the petiole phenotypes in *elf3*

mutants depend on PIF4-PIF5 mediated GA abundance (Filo et al., 2015). Heat shock

proteins also modulate light and temperature signaling via the ELF3-PIF4 module to

affect the hypocotyl growth (Zeng et al., 2023). *MSL10* participates in the

wound-triggered early signal transduction pathway (Zou et al., 2016), while *SLAH3*

respond to flooding stress in *A. thaliana* (Lehmann et al., 2021). In summary, genes

related to environmental stimuli can regulate signal sensing and phytohormone

metabolism, which may act as key links between external changes and internal

responses in *H. difformis*.

Flooding stress (submergence or waterlogging) is a major challenge to many

terrestrial plants, especially crops (Bailey-Serres et al., 2012) in the present era of

rapid climate change. Plants have evolved two main strategies to cope with the lethal

effects of submergence (Bailey-Serres and Voesenek, 2008; Akman et al., 2012;

Muller et al., 2021). The "escape" strategy involves shoot elongation to reach the

surface and restore air contact. In contrast, the "wait" strategy involves growth

reduction and energy conservation to survive until the floods recede. Recent studies

have mainly focused on the "escape" strategy in crops, especially in deep-water rice

(Kuroha et al., 2018; Nagai et al., 2020). Interestingly, the aquatic plant *N. nucifera*

also adopts the "escape" strategy during early submergence (Deng et al., 2022).

Heterophyllous plants appears to have evolved a third strategy, "variation", to adapt to

constant changes in submergence and dehydration (Fig. 6). Understanding the

mechanisms of heterophylly could provide novel avenues to engineer resilient crops

and develop agricultural practices to mitigate the effects of climate change.

Materials and Methods

Plant materials and phenotypic analysis

The sequenced individual of *H. difformis* was collected from South China

Botanical Garden, Chinese Academy of Sciences, Guangdong Province, China

(N113.373781º, E23.18794º). For the analysis of submergence/emergence-treated

samples, the shoots (at the same stage as our previous study (Li et al., 2022)) were

harvested from plants grown under different experimental conditions (A: terrestrial

plants grown at 23°C, 60% RH, 60 µmol m^{-2} s^{-1} illumination; B: terrestrial plants

grown at 18°C, 60% RH, 60 µmol m^{-2} s^{-1} illumination; C: submerged plants grown at

23°C, 60 µmol m^{-2} s^{-1} illumination, D: submerged plants grown at 23°C, 10 µmol m^{-2}

s^{-1} illumination, E: terrestrial plants grown at 23°C, 30% RH, 60 µmol m^{-2} s^{-1}

illumination). For morphological analysis, mature leaves were photographed with a

Canon EOS80D camera, and all light microscopy observations were performed under

a Sunny EX20 light microscope and photographed with a ToupCam TP605100A

digital camera. Images were integrated using MvImage media software (ToupCam).

Leaf complexity was estimated based on the dissection index (DI), calculated as

$DI = leaf\ perimeter/\sqrt{leaf\ area}$, and epidermal cell complexity was also quantified a

similar method of $Cell\ complexity = cell\ perimeter/\sqrt{cell\ area}$. Stomatal and vein

density are calculated as described (Li et al., 2017). All calculations were performed

using ImageJ 1.47v (http://rsb.info.nih.gov/ij/). Statistical differences were

determined using Student's *t*-test or Tukey's test, and data represent the results from at

least twenty independent individuals.

Estimation of genome size and ploidy

Short reads from the MGIseq2000 platform were filtered and used for genome size estimation by SOAPnuke (Chen et al., 2018). Initially, the adaptors were removed from the sequencing reads. Read pairs were then excluded if any single end has an average quality lower than 20, and subsequently, the ends of reads were trimmed if the average quality was lower than 20 in a sliding window size of 5 bp. Finally, read pairs with any end shorter than 75 bp were removed. These quality-filtered reads were used for genome size estimation. We generated the 17-mer occurrence distribution of sequencing reads from short libraries, and the genome size was estimated by 17-K-mer analysis based on the K-mer method (Manekar and Sathe, 2018) of $\text{Genome size} = \text{Kmer number}/\text{Kmer depth}$. The chromosome number of sequenced *H. difformis* was determined by a cytological method (Li et al., 2017). Root tips were fixed in an acetic-alcohol mixture (1:3) overnight. Cells were dissociated by 1 M hydrochloric acid at 60 °C for 1 h. After thorough washing with ultrapure water, they were stained by Modified Carbol fuchsin solution and squashed for chromosome number count. Genome size was estimated by flow cytometry as previously reported (Wang et al., 2014). We collected the young leaves from *H. difformis* and immediately carried out the flow cytometry (Dolezel and Bartos, 2005) to determine genome size by using *Sorghum bicolor* (BTX623, genome size = 0.71Gb) and *Zea mays* (Mo17, genome size = 2.18 Gb) as external standards.

***De novo* sequencing and genome assembly**

Tissue culture-raised plantlets from the *H. difformis* individual were utilized for the isolation of high molecular weight genomic DNA for genome sequencing. The quality and quantity of the extracted DNA were examined using a NanoDrop spectrophotometer (NanoDrop Technologies, Wilmington, DE, USA). Genomic DNA was isolated using the Qiagen DNeasy Plant Mini Kit (Qiagen, USA). We also collected a wide range of tissue samples (stem, shoot, root, flower, and leaves) from the same plant for RNA sequencing (RNA-seq) to support the genome assembly.

Two long-insert continuous long-read libraries (40 kb) were prepared following

PacBio Sequel platform (Pacific Biosciences, Frasergen, Wuhan, China). The SMRT

(Single Molecule Real-Time) library was constructed using the SMRTbell Express

Template Prep kit 2.0 (Pacific Biosciences). Briefly, the DNA was carried into the first

enzymatic reaction to remove single-stranded overhangs followed by treatment with

repair enzymes. Subsequently, the ends of the double-stranded fragments were

polished and tailed with T-overhang SMRTbell adapters and the SMRTbell library

was purified using AMPure PB beads. The size distribution and concentration of the

library were assessed using the FEMTO Pulse automated pulsed-field capillary

electrophoresis instrument (Agilent Technologies, Wilmington, DE) and the Qubit 3.0

Fluorometer (Life Technologies, Carlsbad, CA, USA). Following library

characterization, BluePippin system (Sage Science, Beverly, MA) was used to remove

SMRTbells of ≤25 kb. After size selection, the library was purified with AMPure PB

beads. Library size and quantity were assessed using the FEMTO Pulse and the Qubit

dsDNA HS reagents Assay kit. Sequencing primer and Sequel II DNA Polymerase

were annealed and bound, respectively, to the final SMRTbell library. SMRT

sequencing was performed using a single SMRT Cell on the Sequel II System with

Sequel II Sequencing Kit by Frasergen Bioinformatics Co., Ltd. (Wuhan, China). The

draft assembly of the sequenced genome was assembled using MECAT2 (v20190226)

with the default parameters (Xiao et al., 2017). The SMRT link toolkit was then

performed to correct errors after the initial assembly of the genome.

To anchor hybrid scaffolds onto chromosomes, genomic DNA was also used to

construct a Hi-C library. Briefly, samples were cross-linked under vacuum infiltration

and the cross-linked samples were subsequently lysed. Endogenous nuclease was

inactivated with 0.3% SDS, then chromatin DNA was digested by MboI (NEB), and

marked with biotin-14-dCTP (Invitrogen) and then ligated by T4 DNA ligase (NEB).

After reversing cross-links, the ligated DNA was extracted through QIAamp DNA

Mini Kit (Qiagen) according to manufacturers' instructions. Purified DNA was

sheared to 300- to 500-bp fragments and were further blunt-end repaired, A-tailed and

adaptor added, followed by purification through biotin-streptavidin–mediated

pull-down and PCR amplification. Finally, the Hi-C libraries were quantified and

sequenced on the Illumina Nova-seq platform (San Diego, CA, USA) or MGI-seq

platform (BGI, China). We obtained sequencing data using a BGI MGISEQ-2000

platform on 150 PE mode. For anchored contigs, 381,400,686 clean read pairs were

generated from the Hi-C library and mapped to the polished *H. difformis* genome

using BWA (bwa-0.7.16) (Li and Durbin, 2010). Lachesis (Burton et al., 2013) was

applied to order and orient the clustered contigs, and Juicer and Juicebox tool (v1.8.8)

with default parameters (Durand et al., 2016) were used to correct the assembly error

in the Hi-C assembled genome. Paired reads mapped to different contigs were used

for the Hi-C-associated scaffolding. Self-ligated, non-ligated, and other invalid reads

were filtered out. We applied 3D-DNA to order and orient the clustered contigs and

performed Juicer to filter the sequences and cluster them. Juicebox with default

parameters was applied to adjust chromosome construction manually. We also use

purge haplotigs to remove heterozygosity. We finally anchored the scaffolds on fifteen

chromosomes. In addition, the BUSCO pipeline was used to assess the completeness

and accuracy of our assembled *H. difformis* genome as previous method (Simao et al.,

2015).

Prediction and annotation of protein-coding genes

We predicted the protein-coding genes of *H. difformis* using three methods: *ab*

initio, homology-based, and transcriptomic prediction. AUGUSTUS (v3.3.1) (Stanke

et al., 2006) and GlimmerHMM (v3.0.4) (Majoros et al., 2004) were used to perform

ab initio gene prediction. We used EXONERATE (v2.2.0) (Slater and Birney, 2005)

for the homology-based prediction. First, the protein sequences were aligned to the

genome assembly and predicted coding genes using EXONERATE with the default

parameters. For the transcriptomic prediction, we used RNA-seq analysis. We first

assembled clean RNA-Seq reads into transcripts using trinity (Haas et al., 2013), and

the gene structures were formed using PASA (Trapnell et al., 2010). Finally, MAKER

(v3.00) (Cantarel et al., 2008) was used to integrate the prediction results. Gene

functions were inferred according to the best match of alignments to the

non-redundant database (NR) of the National Center for Biotechnology Information

(NCBI) (https://www.ncbi.nlm.nih.gov/), TrEMBL (Boeckmann et al., 2003),

SWISS-PROT (Boeckmann et al., 2003), and InterPro (Mitchell et al., 2015). Motifs

and protein domains were annotated using PfamScan (Mistry et al., 2007) and

InterProScan (http://www.ebi.ac.uk/InterProScan). We also performed BUSCO

analysis (dataset: embryophyta_odb10 models) using the predicted protein to assess

our annotation completeness of *H. difformis* genome.

Annotation of repetitive sequences

The homology-based and *de novo* prediction methods are combined to identify

the repeats in the genome. For homology-based analysis, we identified the known TEs

within the *H. difformis* genome using RepeatMasker (open-4.0.9) (Tarailo-Graovac

and Chen, 2009) with the Repbase TE library (Jurka et al., 2005). RepeatProteinMask

searches were also conducted using the TE protein database as a query library. For *de*

novo prediction, we constructed a *de novo* repeat library of the *H. difformis* genome

using RepeatModeler (http://www.repeatmasker.org/RepeatModeler/), which can

automatically execute two core *de novo* repeat-finding programs RECON (v1.08)

(Bao and Eddy, 2002) and RepeatScout (v1.0.5) (Price et al., 2005) to

comprehensively conduct, refine and classify consensus models of putative

interspersed repeats for the *H. difformis* genome. Furthermore, we performed a *de*

novo search for long terminal repeat (LTR) retrotransposons against the *H. difformis*

genome sequences using LTR_FINDER (v1.0.7) (Xu and Wang, 2007). Finally, we

merged the library files of the two methods and used Repeatmaker to identify the

repeat contents.

Gene family identification and phylogenetic analysis

We selected 14 species to construct putative gene families: *Oryza sativa*

(*Poaceae*), *Boea hygrometrica* (*Gesneriaceae*), *Mimulus guttatus* (*Phrymaceae*), *Olea*

europaea (*Oleaceae*), *Sesamum indicum* (*Pedaliaceae*), *Andrographis paniculata*

(*Acanthaceae*), *Genlisea aurea* (*Lentibulariaceae*), *Actinidia chinensis*

(*Actinidiaceae*), *Arabidopsis thaliana* (*Brassicaceae*), *Cucumis sativus*

(*Cucurbitaceae*), *Glycine max* (*Leguminosae*), *Ipomoea nil* (*Convolvulaceae*), *Vitis*

vinifera (*Vitaceae*) and *Solanum lycopersicum* (*Solanaceae*), and their genome data

were downloaded from the NCBI database (https://www.ncbi.nlm.nih.gov).

To cluster protein-coding genes into families, protein sequences from the longest

transcript of each gene from *H. difformis* and other species were extracted and aligned

to each other using the BLAST program with a maximal e-value of $1e^{-5}$. The genome

sequences and corresponding protein sequences were downloaded from NCBI

(https://www.ncbi.nlm.nih.gov/). Genes with identity $< 30\%$ or coverage $< 50\%$ were

filtered out to exclude putative fragmented genes. The OrthoMCL (v14-137) (Li et al.,

2003) method was used to cluster genes from these different species into gene

families. To reveal phylogenetic relationships among *H. difformis* and other closely

related species, protein sequences from 196 single-copy orthologous genes were used

for phylogenetic tree reconstruction. The protein sequences of the single-copy

orthologous genes were aligned with the MUSCLE (v3.8.31) program (Edgar, 2004),

and the corresponding coding sequence alignments were generated and concatenated

with the guidance of protein alignment. RAxML (v8.2.11) (Stamatakis, 2014) and

TimeTree (Hedges et al., 2006) were used to construct the phylogenetic tree using the

maximum likelihood method.

Gene family expansion and contraction analysis

Based on the identified gene families and constructed phylogenetic tree with

predicted divergence times of species, we used CAFÉ to analyze gene family

expansion and contraction (Mendes et al., 2020). A random birth or death model in

CAFÉ was used to study gene gain or loss in gene families across the phylogenetic

tree, and an adjusted P value (Q value) was calculated for the genes of *H. difformis* to

test the significant expansion or contraction (Fisher test, Q value < 0.01). These

expansion and contraction genes were mapped to GO pathways for functional

enrichment analysis, which was conducted using the enrichment methods. This

method implemented hypergeometric test algorithms and the Q-value (FDR, False

Discovery Rate) was calculated to adjust the p-value using the R package

(https://github.com/StoreyLab/qvalue).

Collinearity and WGD analysis

BLASTp (e-value $<1e^{-5}$) was used to detect orthologous genes in *H. difformis*, *S.*

cusia and *A. paniculata*, to assess the level of synteny between the species. Then,

syntenic paralogous blocks were identified with MCScanX (Wang et al., 2012)

integrated in TBtools software with default parameters, and visualized by TBtools

(Chen et al., 2020). BLASTp (e-value $<1e^{-5}$) was also used to detect orthologous

pairwise, and MCScanX analysis was then performed for *Ks* evaluation of *V. vinifera*,

A. marina, *H. difformis*, *S. cusia* and *A. paniculata*. Genome data and annotation of *V.*

vinifera and *A. paniculata* were downloaded from the NCBI database

(https://www.ncbi.nlm.nih.gov). Genome data and annotation of *S. cusia* were

downloaded from its database (http://indigoid-plant.iflora.cn) (Xu et al., 2020).

Genome data and annotation of *A. marina* were downloaded from its database

(https://datadryad.org/stash/dataset/doi:10.5061/dryad.3j9kd51f5) (Friis et al., 2021).

These analyses were performed using a haploid genome. We extracted all paralogous

and orthologous gene pairs from syntenic blocks in those species to further estimate

the *Ks* distribution and infer WGD dates by assuming a molecular clock (Schmutz et

al., 2010).

RNA sequencing, gene expression analysis, and co-expression analyses

Total RNA was extracted from shoots including leaf primordia of plants grown

for a month under each condition were collected for RNA-seq, with each sample

performing three biological replicates (Li et al., 2022). We used SOAPnuke (Chen et

al., 2017) to filter the raw reads, and HISAT2 (Kim et al., 2015) to align filtered clean

reads to the *H. difformis* genome. Read counts of these samples were calculated, and

differential gene expression analysis was conducted using the R package DESeq2

(Love et al., 2014) (false discovery rate [FDR] < 0.05 and expression-level $\log_2$

[fold-change] > 1). For DEGs identification and naming, putative orthologs were

identified by tBLASTn and BLASTp (e-value $<1e-5$) analysis to published

Arabidopsis databases (https://www.arabidopsis.org/). To identify relationships

between DEGs, co-expression networks were constructed using the WGCNA package

(Langfelder and Horvath, 2008) integrated in TBtools software with default

parameters, and DEGs were filtered according to their gene expression profiles

(discarded genes that FPKM < 0.9). Cytoscape (Shannon et al., 2003) was used to

visualize the network of genes related to phytohormones' responses. Expression

heatmaps were also generated using TBtools software (Chen et al., 2020).

Quantification of phytohormones and NPA treatments

To measure phytohormone levels, plants were grown in growth chambers at

23°C under a white light flux density of 60 µmol m^{-2} s^{-1} for 1 month. Shoots

including leaf primordia were dissected from the plants and frozen in liquid nitrogen

immediately after sampling. Extraction and quantification of endogenous hormones

(IAA and ABA) were performed as described previously (Nguyen et al., 2018). The

experiments were conducted in triplicate from three independent plants. For

naphthylphthalamic acid (NPA) treatments, plants grown in submerged conditions

were shifted to an aquarium that contained a 10 µM NPA solution. After 1-month of

treatment, emerging leaves were harvested for morphological analysis.

ROS detection

Terrestrial and submerged mature leaves (at same stage, P6) were harvested and

stained with DAB and NBT solution for H_2O_2 and O_2^- analysis, respectively, as

previously described (Chen et al., 2021). Leaves were harvested from twenty

independent plants and immediately stained by H_2O_2 distribution (1%

3,3-diaminobenzidine, DAB), and O_2^- distribution (0.1% nitro blue tetrazolium,

NBT) overnight. Leaves were then rinsed in an ethanol series and recorded for

analysis.

Gene functional studies and expression analysis

To amplify the cDNA sequences of *HdLMI1A*, *HdLMI1B*, and *HdLMI1C*, total

RNA was extracted from shoots including leaf primordia of plants grown for a month

and then used to synthesize cDNA as described previously (Li et al., 2020). Amplified

fragments of expected length were purified and cloned into a pEASY-T1 cloning

vector (TransGen, China) for sequencing. The functions of *HdLMI1A*, *HdLMI1B*, and

HdLMI1C were investigated using the VIGS method as previously described (Zhang

et al., 2020). Specific fragments (171 bp for *HdLMI1A*, 268 bp for *HdLMI1B*, and 252

bp for *HdLMI1C*) were amplified, purified, and independently introduced into the

pTRV2 vector (http://www.bt-lab.cn/) using a Trelief SoSoo Cloning Kit (Tsingke

Biological Technology, China) to generate the constructs TRV2-HdLMI1A,

TRV2-HdLMI1B, and TRV2-HdLMI1C. Recombinant vectors and empty vectors

were then introduced into *Agrobacterium tumefaciens* strain LBA4404 by the liquid

nitrogen freeze-thaw method. Empty vectors were introduced as mock controls. At

least twenty wild-type individuals were used for each treatment. After two rounds of

infiltration (7 and 14 days), plants were grown for 3 additional days in terrestrial

conditions and then shifted to aquaria. After 20 days of growth in submerged

conditions, representative leaves from ten individuals showing phenotypic changes

were photo-documented for analysis. The silencing efficiency of VIGS treatments was

determined by reverse transcription quantitative PCR (RT-qPCR) as previously

described (Li et al., 2022). The cDNAs from individual developing leaves of the

VIGS-treated plants were used as templates. The primers used in this study are

provided in supplementary table S14. Significant differences in relative expression

levels between mock and transgenic lines at $P < 0.01$ were evaluated using Dunnett's

test with the two-sided function in IBM SPSS version 24.

Data availability

The genome dataset is available on GenBank of NCBI BioProject with the

accession number PRJNA872356. Genome sequencing data (BGI seq reads, PacBio

Sequel reads, and Hi-C interaction reads) and environmental treatment RNA-Seq

reads are available in the NCBI Sequence Read Archive under accessions

PRJNA855313 and PRJNA859677. Sequence data from this article can be found in

the GenBank data library under the following accession number: *HdLMI1A*

(OR296694), *HdLMI1B* (OR296695) and *HdLMI1C* (OR296696).

Figure legends

Figure 1. Genome assembly of *Hygrophila difformis*.

(A) Morphology of the submerged plant (top, left), submerged leaf (top, right), terrestrial plant (bottom, left), and terrestrial leaf (bottom, right) of *H. difformis*. Bars = 1 cm.

(B) High-throughput chromosome conformation capture map of the *H. difformis* genome showing a strong intra-chromosomal interaction signal.

(C) Overview of *H. difformis* genome assembly. Chromosome (i), repeats (ii), gene density (iii), GC ratio (iv), and collinearity blocks (v) are displayed. The colour density from light to dark indicates the density of indicated features from low to high.

Figure 2. Evolutionary analysis of the *H. difformis* genome.

(A) Phylogenetic relationships and times of divergence between *H. difformis* and other species. Different color backgrounds indicate species of diverse evolutionary status. Shallow green, Lamiales; Dark green, Acanthaceae.

(B) Expansion and contraction of gene families. Red, contraction; Blue, expansion.

(C) Distribution of average synonymous substitution levels (*Ks*) between syntenic blocks. These paralogs and orthologous gene pairs are among *H. difformis*, *A. paniculata* and *Vitis vinfera*. Left, paralogs; right, orthologs.

Figure 3. Phenotype and transcriptomic analyses of *H. difformis* under terrestrial or submerged conditions.

(A) Morphological parameters (leaf area, dissection index, cell complexity, stomatal density, vein density, and chlorophyll content) of *H. difformis* in terrestrial or submerged conditions. Data from twenty biological replicates for each condition. ***, significant differences detected by Student's *t*-test ($P < 0.01$).

(B) DEGs (false discovery rate [FDR] < 0.05 and expression-level $\log_2$ [fold-change] > 1) involved in environmental response, leaf development, stomatal density, vein density, wax and antioxidants. Relative gene expressions were scaled to see their differences.

(C) DEGs that are involved in ABA metabolism and signalling. Relative gene

expressions were scaled to see their differences.

(D) DEGs that are involved in auxin metabolism and signalling. Genes' relative

expressions were scaled to see their differences.

(E) Diagram for phytohormone-related categories showing differential expression in

terrestrial or submerged conditions. The number of genes is in red and is represented

by the relative size of the circle corresponding to each gene category.

(F) ABA and IAA content in shoots of *H. difformis* under terrestrial or submerged

conditions. Error bars represent ± SD. Asterisks indicate significant differences

(Student's *t*-test: ***, $P < 0.001$). Data presented here are the means of three

independent experiments.

(G) Effect of NPA on *H. difformis* heterophylly. Photos show the emerging leaves

after 10 μM NPA treatment for 1 month. The oldest is on the left, and the youngest

leaf is on the right. The photo is of one representative from ten independent plants

treated with NPA. Bar = 1 cm.

Figure 4. Phenotypic and transcriptomic analyses of *H. difformis* grown under diverse

environmental treatments.

(A) Phenotypes of *H. difformis* grown in different conditions. Group 1: submerged *vs*

terrestrial plants. Group 2: Normal light density *vs* low light density. Group 3: Normal

temperature *vs* low temperature. Group 4: high humidity *vs* normal humidity. Bars =

1cm.

(B-C) Comparative transcriptome analyses of *H. difformis*. UpSet plots present the

overlapping DEGs in 4 comparison groups. UpSet plot of DEGs upregulated

compared to the control (B). UpSet plot of DEGs downregulated compared to the

control (C).

(D) Upregulated DEG expression profiles.

(E) Downregulated DEG expression profiles.

(F) Expression patterns of 27 putative TF genes.

(G) Co-expression network reconstruction identified *LMI1* genes as being clustered

with genes involved in auxin and GA signaling as well as ethylene synthesis and

photosynthesis.

Figure 5. Expression of *LMI1* orthologs and function of *HdLMI1s*.

(A) Relative expression of *LMI1* genes. Up, FPKM of *LMI1* genes in different

environmental conditions. Down, Relative expression of *LMI1* genes in different

groups. * Stars indicate DEGs (false discovery rate [FDR] < 0.05 and expression-level

$\log_2$ [fold-change] > 1).

(B) Phenotypes of whole plants and representative leaves in the mock,

TRV2-HdLMI1A, TRV2-HdLMI1B, and TRV2-HdLMI1C transgenic plants grown in

submerged conditions. Bars = 1 cm.

(C) Dissection index in the mock, TRV2-HdLMI1A, TRV2-HdLMI1B, and

TRV2-HdLMI1C transgenic plants grown in submerged conditions. Data from ten

biological replicates for each, with significant differences between mock and

transgenic plants detected by Dunnett's test (**, $P < 0.01$).

(D) Relative expression of *HdLMI1A*, *HdLMI1B*, and *HdLMI1C* in the mock,

TRV2-HdLMI1A, TRV2-HdLMI1B, and TRV2-HdLMI1C transgenic plants. Error

bars represent ± SD. Data presented here are the means from three independent

experiments. Asterisks indicate significant differences (Dunnett's test: **, $P < 0.01$).

Figure 6. Schematic model of the regulation of heterophylly in *H. difformis*.

The downregulated genes are indicated by blue arrows, whereas the upregulated genes

are indicated by red arrows.

Supplemental Data

Supplemental Figure S1. Phenotypic plasticity in the *Hygrophila* genus.

Supplemental Figure S2. The 17-mer distribution of Illumina short-read data.

Supplemental Figure S3. Mitotic metaphase chromosomes of the sequenced

individual of *H. difformis*.

Supplemental Figure S4. Flow cytometric analysis of the genome size of *H.*

difformis.

Supplemental Figure S5. Distribution of average synonymous substitution levels (Ks)

between syntenic blocks.

Supplemental Figure S6. Photosynthetic physiology analysis of *H. difformis* in

terrestrial and submerged conditions.

Supplemental Figure S7. ROS detection in the terrestrial or submerged leaves.

Supplemental Figure S8. Morphological parameters of *H. difformis* in different

groups.

Supplemental Figure S9. Gene dendrogram and corresponding module colors.

Supplemental Figure S10. Co-expression network and phenotypic analysis in the

yellow module.

Supplemental Table S1. Genome survey by K-mer method.

Supplemental Table S2. General statistics by Hi-C assisted assembly.

Supplemental Table S3. Genome features by Hi-C assisted assembly.

Supplemental Table S4. Information statistics for genome assembly.

Supplemental Table S5. BUSCO analysis for the genome.

Supplemental Table S6. General statistics of gene prediction.

Supplemental Table S7. Summary of gene function annotation.

Supplemental Table S8. Statistics of repeat sequence classification.

Supplemental Table S9. Statistics of gene family clustering.

Supplemental Table S10. Gene expansion analyzed by GO enrichment.

Supplemental Table S11. Gene contraction analyzed by GO enrichment

Supplemental Table S12. Annotation of 180 upregulated DEGs.

Supplemental Table S13. Annotation of 119 downregulated DEGs.

Supplemental Table S14. Primers used in this study.

Author contributions

897   G.J.L. and H.W.H. designed the experiments and wrote the manuscript; G.J.L.,
898   X.Y.Z., and J.J.Y. performed genome and transcriptome analyses; G.J.L. conducted
899   environmental treatment experiments and collected the materials; S.Q.H., J.P., S.K.,
900   K.U.T., and I.H. improved the manuscript. All authors read and approved the final
901   version of the manuscript.

Funding information

904   This study was supported by the International Partnership Program of the
905   Chinese Academy of Sciences (152342KYSB20200021), the National Natural
906   Science Foundation of China (32101254), and the National Key Research and
907   Development Program of China (2017YFE0128800). This work was also supported
908   by the China Scholarship Council.

Acknowledgements

911   We thank professor Shilin Chen of the Institute of Chinese Materia Medica,
912   China Academy of Chinese Medical Sciences for kindly providing the genome
913   annotation of *A. paniculata* and Dr. Markus Stetter, University of Cologne for critical
914   reading of the manuscript. We thank Li Xie and Huan Wang of Wuhan Frasergen
915   Bioinformatics Co., Ltd. for kindly providing the after service of our genome project.

References

Abramson BW, Novotny M, Hartwick NT, Colt K, Aevermann BD, Scheuermann RH, Michael TP (2022) The genome and preliminary single-nuclei transcriptome of Lemna minuta reveals mechanisms of invasiveness. Plant Physiology **188:** 879-897

Agarie S, Kai M, Takatsuji H, Ueno O (1997) Expression of C3 and C4 photosynthetic characteristics in the amphibious plant *Eleocharis vivipara*: structure and analysis of the expression of isogenes for pyruvate, orthophosphate dikinase. Plant Mol Biol **34:** 363-369

Akman M, Bhikharie AV, McLean EH, Boonman A, Visser EJ, Schranz ME, van Tienderen PH (2012) Wait or escape? Contrasting submergence tolerance strategies of *Rorippa amphibia*, *Rorippa sylvestris* and their hybrid. Ann Bot **109:** 1263-1276

Albert VA, Renner T (2020) Aquatic angiosperm ambiguities answered. Nature Plants **6:** 181-183

Alpert P, Simms EL (2002) The relative advantages of plasticity and fixity in different environments: when is it good for a plant to adjust? Evolutionary Ecology **16:** 285-297

Andres RJ, Coneva V, Frank MH, Tuttle JR, Samayoa LF, Han SW, Kaur B, Zhu LL, Fang H, Bowman DT, Rojas-Pierce M, Haigler CH, Jones DC, Holland JB, Chitwood DH, Kuraparthy V (2017) Modifications to a LATE MERISTEM IDENTITY1 gene are responsible for the major leaf shapes of Upland cotton (*Gossypium hirsutum* L.). Proceedings of the National Academy of Sciences of the United States of America **114:** E57-E66

Arnold PA, Kruuk LEB, Nicotra AB (2019) How to analyse plant phenotypic plasticity in response to a changing climate. New Phytologist **222:** 1235-1241

Bailey-Serres J, Lee SC, Brinton E (2012) Waterproofing crops: effective flooding survival strategies. Plant Physiol **160:** 1698-1709

Bailey-Serres J, Voesenek LA (2008) Flooding stress: acclimations and genetic

diversity. Annu Rev Plant Biol **59:** 313-339

Bao Z, Eddy SR (2002) Automated de novo identification of repeat sequence families
in sequenced genomes. Genome Res **12:** 1269-1276

Boeckmann B, Bairoch A, Apweiler R, Blatter MC, Estreicher A, Gasteiger E,
Martin MJ, Michoud K, O'Donovan C, Phan I, Pilbout S, Schneider M
(2003) The SWISS-PROT protein knowledgebase and its supplement TrEMBL
in 2003. Nucleic Acids Research **31:** 365-370

Botts PS, Lawrence JM, Witz BW, Kovach CW (1990) Plasticity in morphology,
proximate composition, and energy content of *Hygrophila polysperma* (Roxb)
Anders. Aquatic Botany **36:** 207-214

Burko Y, Willige BC, Seluzicki A, Novak O, Ljung K, Chory J (2022) PIF7 is a
master regulator of thermomorphogenesis in shade. Nature Communications **13**

Burton JN, Adey A, Patwardhan RP, Qiu R, Kitzman JO, Shendure J (2013)
Chromosome-scale scaffolding of de novo genome assemblies based on
chromatin interactions. Nat Biotechnol **31:** 1119-1125

Canales C, Barkoulas M, Galinha C, Tsiantis M (2010) Weeds of change:
Cardamine hirsuta as a new model system for studying dissected leaf
development. Journal of Plant Research **123:** 25-33

Cantarel BL, Korf I, Robb SMC, Parra G, Ross E, Moore B, Holt C, Alvarado
AS, Yandell M (2008) MAKER: An easy-to-use annotation pipeline designed for
emerging model organism genomes. Genome Research **18:** 188-196

Chang LJ, Mei GF, Hu Y, Deng JQ, Zhang TZ (2019) *LMI1*-like and KNOX1
genes coordinately regulate plant leaf development in dicotyledons. Plant
Molecular Biology **99:** 449-460

Chen CJ, Chen H, Zhang Y, Thomas HR, Frank MH, He YH, Xia R (2020)
TBtools: An integrative toolkit developed for interactive analyses of big
biological data. Molecular Plant **13:** 1194-1202

Chen WY, Su T (2020) Asian monsoon shaped the pattern of woody dicotyledon
richness in humid regions of China. Plant Diversity **42:** 148-154

Chen Y, Chen Y, Shi C, Huang Z, Zhang Y, Li S, Li Y, Ye J, Yu C, Li Z, Zhang X,
Wang J, Yang H, Fang L, Chen Q (2018) SOAPnuke: a MapReduce
acceleration-supported software for integrated quality control and preprocessing
of high-throughput sequencing data. Gigascience **7:** 1-6

Chen Y, Li GJ, Yang JJ, Zhao XY, Sun ZL, Hou HW (2021) Role of Nramp
transporter genes of Spirodela polyrhiza in cadmium accumulation.
Ecotoxicology and Environmental Safety **227:** 112907

Chen YX, Chen YS, Shi CM, Huang ZB, Zhang Y, Li SK, Li Y, Ye J, Yu C, Li Z,
Zhang XQ, Wang J, Yang HM, Fang L, Chen Q (2017) SOAPnuke: a
MapReduce acceleration-supported software for integrated quality control and
preprocessing of high-throughput sequencing data. Gigascience **7:** 1-6

Cheng JL, Li J, Zhang Z, Lu H, Chen GQ, Yao BB, Dong YX, Ma L, Yuan XX,
Xu JX, Zhang Y, Dai WM, Yang XH, Xue LF, Zhang Y, Zhang CB, Mauricio
R, Peng G, Hu SJ, Valverde BE, Song XL, Li Y, Stift M, Qiang S (2021)
Autopolyploidy-driven range expansion of a temperate-originated plant to
pan-tropic under global change. Ecological Monographs **91:** e01445

de Wit M, Lorrain S, Fankhauser C (2014) Auxin-mediated plant architectural
changes in response to shade and high temperature. Physiologia Plantarum **151:**
13-24

Debernardi JM, Mecchia MA, Vercruyssen L, Smaczniak C, Kaufmann K, Inze
D, Rodriguez RE, Palatnik JF (2014) Post-transcriptional control of GRF
transcription factors by microRNA miR396 and GIF co-activator affects leaf size
and longevity. Plant J **79:** 413-426

DeMason DA, Chawla R (2004) Roles for auxin during morphogenesis of the
compound leaves of pea (*Pisum sativum*). Planta **218:** 435-448

Deng X, Yang D, Sun H, Liu J, Song H, Xiong Y, Wang Y, Ma J, Zhang M, Li J,
Liu Y, Yang M (2022) Time-course analysis and transcriptomic identification of
key response strategies to complete submergence in *Nelumbo nucifera*. Hortic
Res **9:** uhac001

Dolezel J, Bartos J (2005) Plant DNA flow cytometry and estimation of nuclear

genome size. Annals of Botany **95:** 99-110

Durand NC, Shamim MS, Machol I, Rao SS, Huntley MH, Lander ES, Aiden EL

(2016) Juicer provides a one-click system for analyzing loop-resolution Hi-C

experiments. Cell Syst **3:** 95-98

Edgar RC (2004) MUSCLE: multiple sequence alignment with high accuracy and

high throughput. Nucleic Acids Research **32:** 1792-1797

Filo J, Wu A, Eliason E, Richardson T, Thines BC, Harmon FG (2015)

Gibberellin driven growth in elf3 mutants requires PIF4 and PIF5. Plant

Signaling & Behavior **10**

Finet C, Jaillais Y (2012) AUXOLOGY: When auxin meets plant evo-devo.

Developmental Biology **369:** 19-31

Friis G, Vizueta J, Smith EG, Nelson DR, Khraiwesh B, Qudeimat E,

Salehi-Ashtiani K, Ortega A, Marshell A, Duarte CM, Burt JA (2021) A

high-quality genome assembly and annotation of the gray mangrove, *Avicennia*

marina. G3-Genes Genomes Genetics **11**

Godden GT, Kinser TJ, Soltis PS, Soltis DE (2019) Phylotranscriptomic Analyses

Reveal Asymmetrical Gene Duplication Dynamics and Signatures of Ancient

Polyploidy in Mints. Genome Biol Evol **11:** 3393-3408

Govindarajan T, Subramanian D (1983) Karyomorphological Studies in South

Indian Acanthaceae. Cytologia **48:** 491-504

Haas BJ, Papanicolaou A, Yassour M, Grabherr M, Blood PD, Bowden J, Couger

MB, Eccles D, Li B, Lieber M, MacManes MD, Ott M, Orvis J, Pochet N,

Strozzi F, Weeks N, Westerman R, William T, Dewey CN, Henschel R,

LeDuc RD, Friedman N, Regev A (2013) De novo transcript sequence

reconstruction from RNA-seq using the Trinity platform for reference generation

and analysis. Nat Protoc **8:** 1494-1512

Han SJ, Xing ZF, Jiang HS, Li W, Huang WM (2021) Biological adaptive

mechanisms displayed by a freshwater plant to live in aquatic and terrestrial

environments. Environmental and Experimental Botany **191**

Hareven D, Gutfinger T, Parnis A, Eshed Y, Lifschitz E (1996) The making of a
compound leaf: genetic manipulation of leaf architecture in tomato. Cell **84:**
735-744

Hay A, Barkoulas M, Tsiantis M (2006) ASYMMETRIC LEAVES1 and auxin
activities converge to repress *BREVIPEDICELLUS* expression and promote leaf
development in Arabidopsis. Development **133:** 3955-3961

He D, Guo P, Gugger PF, Guo Y, Liu X, Chen J (2018) Investigating the molecular
basis for heterophylly in the aquatic plant *Potamogeton octandrus*
(Potamogetonaceae) with comparative transcriptomics. PeerJ **6:** e4448

Hedges SB, Dudley J, Kumar S (2006) TimeTree: a public knowledge-base of
divergence times among organisms. Bioinformatics **22:** 2971-2972

Hoffmann AA, Sgro CM (2011) Climate change and evolutionary adaptation. Nature
470: 479-485

Horiguchi G, Matsumoto K, Nemoto K, Inokuchi M, Hirotsu N (2021) Transition
from proto-kranz-type photosynthesis to HCO_3^- use photosynthesis in the
amphibious plant *Hygrophila polysperma*. Frontiers in Plant Science **12:** 675507

Horiguchi G, Nemoto K, Yokoyama T, Hirotsu N (2019) Photosynthetic
acclimation of terrestrial and submerged leaves in the amphibious plant
Hygrophila difformis. AoB Plants **11:** plz009

Horiguchi G, Oyama R, Akabane T, Suzuki N, Katoh E, Mizokami Y, Noguchi K,
Hirotsu N (2023) Cooperation of external carbonic anhydrase and HCO3-
transporter supports underwater photosynthesis in submerged leaves of the
amphibious plant *Hygrophila difformis*. Ann Bot

Hu L, Zhang H, Yang Q, Meng Q, Han S, Nwafor CC, Khan MHU, Fan C, Zhou
Y (2018) Promoter variations in a homeobox gene, BnA10.LMI1, determine
lobed leaves in rapeseed (*Brassica napus* L.). Theor Appl Genet **131:** 2699-2708

Hunt L, Bailey KJ, Gray JE (2010) The signalling peptide EPFL9 is a positive
regulator of stomatal development. New Phytologist **186:** 609-614

Ingale KG, Thakurdesai PA, Vyawahare NS (2013) Protective effect of *Hygrophila spinosa* against cisplatin induced nephrotoxicity in rats. Indian Journal of Pharmacology **45:** 232-236

Jung JH, Barbosa AD, Hutin S, Kumita JR, Gao M, Derwort D, Silva CS, Lai X, Pierre E, Geng F, Kim SB, Baek S, Zubieta C, Jaeger KE, Wigge PA (2020) A prion-like domain in ELF3 functions as a thermosensor in Arabidopsis. Nature **585:** 256-260

Jurka J, Kapitonov VV, Pavlicek A, Klonowski P, Kohany O, Walichiewicz J (2005) Repbase Update, a database of eukaryotic repetitive elements. Cytogenet Genome Res **110:** 462-467

Karatas M, Aasim M, Cinar A, Dogan M (2013) Adventitious Shoot Regeneration from Leaf Explant of Dwarf Hygro (*Hygrophila polysperma* (Roxb.) T. Anderson). Scientific World Journal

Kieffer M, Master V, Waites R, Davies B (2011) TCP14 and TCP15 affect internode length and leaf shape in Arabidopsis. Plant J **68:** 147-158

Kim D, Langmead B, Salzberg SL (2015) HISAT: a fast spliced aligner with low memory requirements. Nat Methods **12:** 357-360

Kim J, Joo Y, Kyung J, Jeon M, Park JY, Lee HG, Chung DS, Lee E, Lee I (2018) A molecular basis behind heterophylly in an amphibious plant, Ranunculus trichophyllus. Plos Genetics **14:** e1007208

Kimura S, Koenig D, Kang J, Yoong FY, Sinha N (2008) Natural variation in leaf morphology results from mutation of a novel *KNOX* gene. Curr Biol **18:** 672-677

Koga H, Kojima M, Takebayashi Y, Sakakibara H, Tsukaya H (2021) Identification of the unique molecular framework of heterophylly in the amphibious plant Callitriche palustris L. Plant Cell **33:** 3272-3292

Kordyum EL, Mosyakin SL (2020) Endosperm of angiosperms and genomic imprinting. Life (Basel) **10:** 104

Kumar R, Kushalappa K, Godt D, Pidkowich MS, Pastorelli S, Hepworth SR, Haughn GW (2007) The Arabidopsis BEL1-LIKE HOMEODOMAIN proteins

SAW1 and SAW2 act redundantly to regulate *KNOX* expression spatially in leaf margins. Plant Cell **19:** 2719-2735

Kumar S, Stecher G, Suleski M, Hedges SB (2017) TimeTree: A resource for timelines, timetrees, and divergence times. Mol Biol Evol **34:** 1812-1819

Kuroha T, Nagai K, Gamuyao R, Wang DR, Furuta T, Nakamori M, Kitaoka T, Adachi K, Minami A, Mori Y, Mashiguchi K, Seto Y, Yamaguchi S, Kojima M, Sakakibara H, Wu J, Ebana K, Mitsuda N, Ohme-Takagi M, Yanagisawa S, Yamasaki M, Yokoyama R, Nishitani K, Mochizuki T, Tamiya G, McCouch SR, Ashikari M (2018) Ethylene-gibberellin signaling underlies adaptation of rice to periodic flooding. Science **361:** 181-186

Kuwabara A, Ikegami K, Koshiba T, Nagata T (2003) Effects of ethylene and abscisic acid upon heterophylly in *Ludwigia arcuata* (Onagraceae). Planta **217:** 880-887

Lan T, Renner T, Ibarra-Laclette E, Farr KM, Chang TH, Cervantes-Perez SA, Zheng C, Sankoff D, Tang H, Purbojati RW, Putra A, Drautz-Moses DI, Schuster SC, Herrera-Estrella L, Albert VA (2017) Long-read sequencing uncovers the adaptive topography of a carnivorous plant genome. Proc Natl Acad Sci U S A **114:** E4435-E4441

Langfelder P, Horvath S (2008) WGCNA: an R package for weighted correlation network analysis. BMC Bioinformatics **9:** 559

Lehmann J, Jorgensen ME, Fratz S, Muller HM, Kusch J, Scherzer S, Navarro-Retamal C, Mayer D, Bohm J, Konrad KR, Terpitz U, Dreyer I, Mueller TD, Sauer M, Hedrich R, Geiger D, Maierhofer T (2021) Acidosis-induced activation of anion channel SLAH3 in the flooding-related stress response of Arabidopsis. Curr Biol **31:** 3575-3585 e3579

Les DH, Philbrick CT (1993) Studies of Hybridization and Chromosome-Number Variation in Aquatic Angiosperms - Evolutionary Implications. Aquatic Botany **44:** 181-228

Li G, Hu S, Hou H, Kimura S (2019) Heterophylly: Phenotypic plasticity of leaf

shape in aquatic and amphibious plants. Plants (Basel) **8:** 420

Li G, Hu S, Yang J, Schultz EA, Clarke K, Hou H (2017) Water-Wisteria as an

ideal plant to study heterophylly in higher aquatic plants. Plant Cell Reports **36:**

1225-1236

Li G, Hu S, Yang J, Zhao X, Kimura S, Schultz EA, Hou H (2020) Establishment

of an Agrobacterium mediated transformation protocol for the detection of

cytokinin in the heterophyllous plant *Hygrophila difformis* (Acanthaceae). Plant

Cell Rep **39:** 737-750

Li G, Hu S, Zhao X, Kumar S, Li Y, Yang J, Hou H (2021) Mechanisms of the

morphological plasticity induced by phytohormones and the environment in

plants. Int J Mol Sci **22:** 765

Li G, Yang J, Chen Y, Zhao X, Chen Y, Kimura S, Hu S, Hou H (2022) SHOOT

MERISTEMLESS participates in the heterophylly of *Hygrophila difformis*

(Acanthaceae). Plant Physiol**:** kiac382

Li H, Durbin R (2010) Fast and accurate long-read alignment with Burrows-Wheeler

transform. Bioinformatics **26:** 589-595

Li L, Stoeckert CJ, Roos DS (2003) OrthoMCL: Identification of ortholog groups

for eukaryotic genomes. Genome Research **13:** 2178-2189

Love MI, Huber W, Anders S (2014) Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. Genome Biol **15:** 550

Lu RS, Chen Y, Zhang XY, Feng Y, Comes HP, Li Z, Zheng ZS, Yuan Y, Wang

LY, Huang ZJ, Guo Y, Sun GP, Olsen KM, Chen J, Qiu YX (2021) Genome

sequencing and transcriptome analyses provide insights into the origin and

domestication of water caltrop (Trapa spp., Lythraceae). Plant Biotechnol J **20:**

761-776

Ma DN, Ding QS, Guo ZJ, Xu CQ, Liang PP, Zhao ZZ, Song SW, Zheng HL

(2022) The genome of a mangrove plant, Avicennia marina, provides insights

into adaptation to coastal intertidal habitats. Planta **256**

Ma XG, Wang ZW, Tian B, Sun H (2019) Phylogeographic Analyses of the East

Asian Endemic Genus Prinsepia and the Role of the East Asian Monsoon System

in Shaping a North-South Divergence Pattern in China. Frontiers in Genetics **10**

Majoros WH, Pertea M, Salzberg SL (2004) TigrScan and GlimmerHMM: two

open source ab initio eukaryotic gene-finders. Bioinformatics **20:** 2878-2879

Manekar SC, Sathe SR (2018) A benchmark study of k-mer counting methods for

high-throughput sequencing. Gigascience **7**

McCracken A, Bainard JD, Miller MC, Husband BC (2013) Pathways of

introduction of the invasive aquatic plant Cabomba caroliniana. Ecology and

Evolution **3:** 1427-1439

Mehrotra S, Goyal V (2014) Repetitive sequences in plant nuclear DNA: types,

distribution, evolution and function. Genomics Proteomics Bioinformatics **12:**

164-171

Mendes FK, Vanderpool D, Fulton B, Hahn MW (2020) CAFE 5 models variation

in evolutionary rates among gene families. Bioinformatics **36:** 5516-5518

Mistry J, Bateman A, Finn RD (2007) Predicting active site residue annotations in

the Pfam database. BMC Bioinformatics **8:** 298

Mitchell A, Chang HY, Daugherty L, Fraser M, Hunter S, Lopez R, McAnulla C,

McMenamin C, Nuka G, Pesseat S, Sangrador-Vegas A, Scheremetjew M,

Rato C, Yong SY, Bateman A, Punta M, Attwood TK, Sigrist CJA, Redaschi

N, Rivoire C, Xenarios I, Kahn D, Guyot D, Bork P, Letunic I, Gough J,

Oates M, Haft D, Huang HZ, Natale DA, Wu CH, Orengo C, Sillitoe I, Mi

HY, Thomas PD, Finn RD (2015) The InterPro protein families database: the

classification resource after 15 years. Nucleic Acids Research **43:** D213-D221

Momokawa N, Kadono Y, Kudoh H (2011) Effects of light quality on leaf

morphogenesis of a heterophyllous amphibious plant, *Rotala hippuris*. Ann Bot

108: 1299-1306

Moon J, Hake S (2011) How a leaf gets its shape. Current Opinion in Plant Biology

14: 24-30

Muller JT, van Veen H, Bartylla MM, Akman M, Pedersen O, Sun P, Schuurink

RC, Takeuchi J, Todoroki Y, Weig AR, Sasidharan R, Mustroph A (2021)
Keeping the shoot above water - submergence triggers antithetical growth
responses in stems and petioles of watercress (*Nasturtium officinale*). New
Phytol **229:** 140-155

Nagai K, Mori Y, Ishikawa S, Furuta T, Gamuyao R, Niimi Y, Hobo T, Fukuda M,
Kojima M, Takebayashi Y, Fukushima A, Himuro Y, Kobayashi M, Ackley
W, Hisano H, Sato K, Yoshida A, Wu J, Sakakibara H, Sato Y, Tsuji H,
Akagi T, Ashikari M (2020) Antagonistic regulation of the gibberellic acid
response during stem growth in rice. Nature **584:** 109-114

Nakayama H, Kimura S (2015) Leaves may function as temperature sensors in the
heterophylly of *Rorippa aquatica* (Brassicaceae). Plant Signaling & Behavior **10:**
e1091909

Nakayama H, Nakayama N, Seiki S, Kojima M, Sakakibara H, Sinha N, Kimura
S (2014) Regulation of the KNOX-GA gene module induces heterophyllic
alteration in North American lake cress. Plant Cell **26:** 4733-4748

Nakayama H, Sinha NR, Kimura S (2017) How do plants and phytohormones
accomplish heterophylly, leaf phenotypic plasticity, in response to environmental
cues. Frontiers in Plant Science **8:** 1717

Nguyen TN, Tuan PA, Mukherjee S, Son S, Ayele BT (2018) Hormonal regulation
in adventitious roots and during their emergence under waterlogged conditions in
wheat. Journal of Experimental Botany **69:** 4065-4082

Nicotra AB, Atkin OK, Bonser SP, Davidson AM, Finnegan EJ, Mathesius U,
Poot P, Purugganan MD, Richards CL, Valladares F, van Kleunen M (2010)
Plant phenotypic plasticity in a changing climate. Trends in Plant Science **15:**
684-692

O'Brien CL, Huber M, Thomas E, Pagani M, Super JR, Elder LE, Hull PM
(2020) The enigma of Oligocene climate and global surface temperature
evolution. Proc Natl Acad Sci U S A **117:** 25302-25309

Park H, Park JH, Lee Y, Woo DU, Jeon HH, Sung YW, Shim S, Kim SH, Lee KO,

Kim JY, Kim CK, Bhattacharya D, Yoon HS, Kang YJ (2021) Genome of the
world's smallest flowering plant, *Wolffia australiana*, helps explain its
specialized physiology and unique morphology. Communications Biology **4:** 900

Prancl J, Kaplan Z, Travnicek P, Jarolimova V (2014) Genome Size as a Key to
Evolutionary Complex Aquatic Plants: Polyploidy and Hybridization in
Callitriche (Plantaginaceae). Plos One **9**

Price AL, Jones NC, Pevzner PA (2005) De novo identification of repeat families in
large genomes. Bioinformatics **21 Suppl 1:** i351-358

Qi XH, Li QQ, Ma XT, Qian CL, Wang HH, Ren NN, Shen CX, Huang SM, Xu
XW, Xu Q, Chen XH (2019) Waterlogging-induced adventitious root formation
in cucumber is regulated by ethylene and auxin through reactive oxygen species
signalling. Plant Cell and Environment **42:** 1458-1470

Ranallo-Benavidez TR, Jaron KS, Schatz MC (2020) GenomeScope 2.0 and
Smudgeplot for reference-free profiling of polyploid genomes. Nat Commun **11:**
1432

Raschke A, Ibanez C, Ullrich KK, Anwer MU, Becker S, Glockner A, Trenner J,
Denk K, Saal B, Sun X, Ni M, Davis SJ, Delker C, Quint M (2015) Natural
variants of ELF3 affect thermomorphogenesis by transcriptionally modulating
PIF4-dependent auxin response genes. BMC Plant Biol **15:** 197

Ren G, Jiang Y, Li A, Yin M, Li M, Mu W, Wu Y, Liu J (2022) The genome
sequence provides insights into salt tolerance of *Achnatherum splendens*
(Gramineae), a constructive species of alkaline grassland. Plant Biotechnol J **20:**
116-128

Saddic LA, Huvermann BR, Bezhani S, Su YH, Winter CM, Kwon CS, Collum
RP, Wagner D (2006) The LEAFY target LMI1 is a meristem identity regulator
and acts together with LEAFY to regulate expression of CAULIFLOWER.
Development **133:** 1673-1682

Schmutz J, Cannon SB, Schlueter J, Ma JX, Mitros T, Nelson W, Hyten DL, Song
QJ, Thelen JJ, Cheng JL, Xu D, Hellsten U, May GD, Yu Y, Sakurai T,

Umezawa T, Bhattacharyya MK, Sandhu D, Valliyodan B, Lindquist E, Peto

M, Grant D, Shu SQ, Goodstein D, Barry K, Futrell-Griggs M, Abernathy B,

Du JC, Tian ZX, Zhu LC, Gill N, Joshi T, Libault M, Sethuraman A, Zhang

XC, Shinozaki K, Nguyen HT, Wing RA, Cregan P, Specht J, Grimwood J,

Rokhsar D, Stacey G, Shoemaker RC, Jackson SA (2010) Genome sequence

of the palaeopolyploid soybean. Nature **463:** 178-183

Shani E, Yanai O, Ori N (2006) The role of hormones in shoot apical meristem

function. Current Opinion in Plant Biology **9:** 484-489

Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, Amin N,

Schwikowski B, Ideker T (2003) Cytoscape: a software environment for

integrated models of biomolecular interaction networks. Genome Res **13:**

2498-2504

Shi T, Huneau C, Zhang Y, Li Y, Chen J, Salse J, Wang Q (2022) The

slow-evolving Acorus tatarinowii genome sheds light on ancestral monocot

evolution. Nat Plants **8:** 764-777

Simao FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM (2015)

BUSCO: assessing genome assembly and annotation completeness with

single-copy orthologs. Bioinformatics **31:** 3210-3212

Slater GS, Birney E (2005) Automated generation of heuristics for biological

sequence comparison. Bmc Bioinformatics **6:** 31

Stamatakis A (2014) RAxML version 8: a tool for phylogenetic analysis and

post-analysis of large phylogenies. Bioinformatics **30:** 1312-1313

Stanke M, Keller O, Gunduz I, Hayes A, Waack S, Morgenstern B (2006)

AUGUSTUS: ab initio prediction of alternative transcripts. Nucleic Acids

Research **34:** W435-W439

Stotz GC, Salgado-Luarte C, Escobedo VM, Valladares F, Gianoli E (2021)

Global trends in phenotypic plasticity of plants. Ecology Letters **24:** 2267-2281

Sun J, Qi L, Li Y, Zhai Q, Li C (2013) PIF4 and PIF5 transcription factors link blue

light and auxin to regulate the phototropic response in Arabidopsis. Plant Cell **25:**

2102-2114

Sun JQ, Qi LL, Li YN, Chu JF, Li CY (2012) PIF4-Mediated Activation of *YUCCA8* Expression Integrates Temperature into the Auxin Pathway in Regulating Arabidopsis Hypocotyl Growth. Plos Genetics **8**

Sun W, Leng L, Yin Q (2019) The genome of the medicinal plant Andrographis paniculata provides insight into the biosynthesis of the bioactive diterpenoid neoandrographolide (vol 97, pg 841, 2018). Plant Journal **97:** 996-996

Tarailo-Graovac M, Chen N (2009) Using RepeatMasker to identify repetitive elements in genomic sequences. Curr Protoc Bioinformatics **Chapter 4:** 4 10 11-14 10 14

Titus JE, Gary Sullivan P (2001) Heterophylly in the yellow waterlily, *Nuphar variegata* (Nymphaeaceae): effects of [CO_2], natural sediment type, and water depth. Am J Bot **88:** 1469-1478

Tran D, Girault T, Guichard M, Thomine S, Leblanc-Fournier N, Moulia B, de Langre E, Allain JM, Frachisse JM (2021) Cellular transduction of mechanical oscillations in plants by the plasma-membrane mechanosensitive channel MSL10. Proc Natl Acad Sci U S A **118:** e1919402118

Trapnell C, Williams BA, Pertea G, Mortazavi A, Kwan G, van Baren MJ, Salzberg SL, Wold BJ, Pachter L (2010) Transcript assembly and quantification by RNA-Seq reveals unannotated transcripts and isoform switching during cell differentiation. Nature Biotechnology **28:** 511-U174

Van De Peer Y, Mizrachi E, Marchal K (2017) The evolutionary significance of polyploidy. Nature Reviews Genetics **18:** 411-424

van Veen H, Mustroph A, Barding GA, Vergeer-van Eijk M, Welschen-Evertman RA, Pedersen O, Visser EJ, Larive CK, Pierik R, Bailey-Serres J, Voesenek LA, Sasidharan R (2013) Two Rumex species from contrasting hydrological niches regulate flooding tolerance through distinct mechanisms. Plant Cell **25:** 4691-4707

van Veen H, Sasidharan R (2021) Shape shifting by amphibious plants in dynamic

hydrological niches. New Phytol **229:** 79-84

Vlad D, Kierzkowski D, Rast MI, Vuolo F, Dello Ioio R, Galinha C, Gan XC, Hajheidari M, Hay A, Smith RS, Huijser P, Bailey CD, Tsiantis M (2014) Leaf shape evolution through duplication, regulatory diversification, and loss of a homeobox gene. Science **343:** 780-783

Wang H, Jones B, Li ZG, Frasse P, Delalande C, Regad F, Chaabouni S, Latche A, Pech JC, Bouzayen M (2005) The tomato Aux/IAA transcription factor IAA9 is involved in fruit development and leaf morphogenesis. Plant Cell **17:** 2676-2692

Wang LH, Yu S, Tong CB, Zhao YZ, Liu Y, Song C, Zhang YX, Zhang XD, Wang Y, Hua W, Li DH, Li D, Li F, Yu JY, Xu CY, Han XL, Huang S, Tai SS, Wang JY, Xu X, Li YR, Liu SY, Varshney RK, Wang J, Zhang XR (2014) Genome sequencing of the high oil crop sesame provides insight into oil biosynthesis. Genome Biology **15**

Wang X, Zhang JJ, Xie YY, Liu X, Wen LZ, Wang HF, Zhang J, Li J, Han L, Yu XL, Mysore KS, Wen JQ, Zhou C (2021) LATE MERISTEM IDENTITY1 regulates leaf margin development via the auxin transporter gene SMOOTH LEAF MARGIN1. Plant Physiology **187:** 218-235

Wang Y, Strauss S, Liu SD, Pieper B, Lymbouridou R, Runions A, Tsiantis M (2022) The cellular basis for synergy between RCO and KNOX1 homeobox genes in leaf shape diversity. Current Biology **32:** 3773-+

Wang YP, Tang HB, DeBarry JD, Tan X, Li JP, Wang XY, Lee TH, Jin HZ, Marler B, Guo H, Kissinger JC, Paterson AH (2012) MCScanX: a toolkit for detection and evolutionary analysis of gene synteny and collinearity. Nucleic Acids Research **40:** e49

Wanke D (2011) The ABA-mediated switch between submersed and emersed life-styles in aquatic macrophytes. Journal of Plant Research **124:** 467-475

Wasiljeff J, Salminen JM, Stenman J, Zhang ZQ, Kaakinen A (2022) Oligocene moisture variations as evidenced by an aeolian dust sequence in Inner Mongolia,

China. Scientific Reports **12**

Wu P, Zhang LK, Zhang K, Yin YL, Liu AL, Zhu Y, Fu Y, Sun FF, Zhao SP, Feng K, Xu XW, Chen XH, Cheng F, Li LJ (2022) The adaptive evolution of *Euryale ferox* to the aquatic environment through paleo-hexaploidization. Plant Journal **110:** 627-645

Xiao CL, Chen Y, Xie SQ, Chen KN, Wang Y, Han Y, Luo F, Xie Z (2017) MECAT : fast mapping, error correction, and de novo assembly for single-molecule sequencing reads. Nature Methods **14:** 1072-1074

Xu W, Zhang LB, Cunningham AB, Li S, Zhuang HF, Wang YH, Liu AZ (2020) Blue genome: chromosome-scale genome reveals the evolutionary and molecular basis of indigo biosynthesis in *Strobilanthes cusia*. Plant Journal **104:** 864-879

Xu Z, Wang H (2007) LTR_FINDER: an efficient tool for the prediction of full-length LTR retrotransposons. Nucleic Acids Res **35:** W265-268

Xue T, Zheng X, Chen D, Liang L, Chen N, Huang Z, Fan W, Chen J, Cen W, Chen S, Zhu J, Chen B, Zhang X, Chen Y (2020) A high-quality genome provides insights into the new taxonomic status and genomic characteristics of *Cladopus chinensis* (Podostemaceae). Hortic Res **7:** 46

Yaneff A, Vitali V, Amodeo G (2015) PIP1 aquaporins: Intrinsic water channels or PIP2 aquaporin modulators? Febs Letters **589:** 3508-3515

Yao G, Drew BT, Yi TS, Yan HF, Yuan YM, Ge XJ (2016) Phylogenetic relationships, character evolution and biogeographic diversification of Pogostemon s.l. (Lamiaceae). Mol Phylogenet Evol **98:** 184-200

Zeng Y, Wang JY, Huang S, Xie Y, Zhu TD, Liu LY, Li L (2023) HSP90s are required for hypocotyl elongation during skotomorphogenesis and thermomorphogenesis via the COP1-ELF3-PIF4 pathway in Arabidopsis. New Phytologist

Zhang L, Zhou L, Yung WS, Su W, Huang M (2021) Ectopic expression of Torenia fournieri TCP8 and TCP13 alters the leaf and petal phenotypes in *Arabidopsis thaliana*. Physiol Plant **173:** 856-866

Zhang R, Fu XH, Zhao CY, Cheng J, Liao H, Wang PP, Yao X, Duan XS, Yuan Y,
Xu GX, Kramer EM, Shan HY, Kong HZ (2020) Identification of the key
regulatory genes involved in elaborate petal development and specialized
character formation in *Nigella damascena* (Ranunculaceae). Plant Cell **32:**
3095-3112

Zhou XJ, Li JT, Wang HL, Han JW, Zhang K, Dong SW, Zhang YZ, Ya HY,
Cheng YW, Sun SS (2021) The chromosome-scale genome assembly,
annotation and evolution of *Rhododendron henanense* subsp. lingbaoense. Mol
Ecol Resour **22:** 988-1001

Zou Y, Chintamanani S, He P, Fukushige H, Yu L, Shao M, Zhu L, Hildebrand
DF, Tang X, Zhou JM (2016) A gain-of-function mutation in Msl10 triggers cell
death and wound-induced hyperaccumulation of jasmonic acid in Arabidopsis. J
Integr Plant Biol **58:** 600-609

1365