1 **Multi-site phosphorylation of the TOPBP1 ATR Activation Domain propels**
2 **ATR signaling during the response to DNA breaks**
3

Holly Senebandith[1], Kenna Ruis[1], Katrina Montales[1], Yasaman Jami-Alahmadi[2],

James A. Wohlschlegel[2], and W. Matthew Michael[1,3]

[1]Molecular and Computational Biology Section

Department of Biological Sciences

University of Southern California

Los Angeles, CA 90089

[2]Department of Biological Chemistry

Geffen School of Medicine at UCLA

Los Angeles, CA 90095

[3]Author for correspondence at: mattm@usc.edu

Abstract

TOPBP1 is a BRCT domain-containing scaffold protein that plays important roles in

a diverse array of cellular processes, including DNA damage signaling, DNA repair,

DNA replication, and mRNA transcription. For DNA damage signaling, TOPBP1

activates the crucial damage response kinase ATR and this occurs during replication

stress as well as during a DNA double-strand break response. ATR signaling allows

cells to survive genotoxic issues and represents a formidable barrier to

transformation to a tumorigenic state. Despite its importance to genome stability,

the biochemical mechanism for how TOPBP1 activates ATR is not fully understood.

TOPBP1 uses a discrete domain, termed the ATR activation domain, to stimulate

ATR kinase. Recent work has shown that the AAD must be in a multimeric state to

activate ATR. Other work has shown that phosphorylation of the AAD on serine

1131 (in Xenopus) is important for its function, and some have suggested that this is

linked to the formation of TOPBP1 condensates during a DNA damage response. In

this study we examine AAD phosphorylation in detail and we report three important

new findings. One, S1131 phosphorylation promotes ATR activation in a manner

independent of condensate formation and is instead linked to promoting

multimerization of the AAD. Two, we identify a novel sight of AAD phosphorylation,

on T1098, and show that is required for ATR activation. Three, we identify

additional, candidate phosphorylation sites, some of which fit the consensus for

casein kinase 2, and we show that casein kinase 2 activity is required for the AAD to

perform its function. These studies show that multi-site phosphorylation of the AAD

is an important component of the mechanism by which TOPBP1 activates ATR.

Key words

TOPBP1, ATR, AAD, CHK1, multi-site phosphorylation, casein kinase 2

1. Introduction

Ataxia telangiectasia and Rad3-related (ATR) kinase is a vital component of the cellular response to replication stress and DNA damage (Salvidar et al, 2017; Waterman et al, 2020). ATR is also important for the proper timing of cell cycle progression under unperturbed conditions (Saldivar et al, 2018). This large phosphatidylinositol 3-kinase-like kinase (PIKK) is hyper-activated at sites of DNA damage and it goes on to phosphorylate hundreds of substrates to allow cells to survive genotoxic stress. ATR signaling is also a barrier to cancer as it will drive damaged cells into senescence, thereby preventing their continued proliferation (Bartkova et al. 2006; Di Micco et al. 2006). Despite the importance of ATR signaling to human health and disease, it remains unclear how ATR is activated by damaged DNA or stalled replication forks.

So far, two proteins have been identified that help activate ATR at sites of damage, TOPBP1 and ETAA1 (reviewed in Niedzwiedz, 2016; Salvidar et al, 2017). The work presented here focuses on TOPBP1, which is composed of nine BRCT domains and a discrete and stand-alone ATR activation domain (AAD; Figure 1A). To initiate DNA damage signaling, for example at sites of DNA double-strand breaks (DSBs), ATR is recruited to the DSB via its binding partner ATRIP while TOPBP1 is recruited separately, via an unknown mechanism. At the DSB, TOPBP1 undergoes a conformational change that allows its AAD to contact ATR and ATRIP, in a manner that greatly stimulates ATR kinase activity. Recent work from our group, and others, has revealed that the conformational change is linked to multimerization of TOPBP1

at sites of damage (Kim et al, 2020; Thada et al, 2021; Ruis et al, 2021). Notably,

while the AAD by itself can activate ATR when present at high concentrations,

forced multimerization of the AAD renders it a far more potent activator (Kim et al,

2020; Thada et al, 2021). Indeed, forced multimerization of the full-length TOPBP1

protein also produces more robust ATR activation (Zhou et al, 2013). These studies

make it clear that the way TOPBP1 is regulated is through control of its oligomeric

state; TOPBP1 only assumes a multimeric state when it is bound to damaged DNA,

and thus ATR is only activated at sites of damage and is not constitutively activated

in solution.

These findings have now focused attention on how TOPBP1 multimers form

at sites of damage. Recent work from our group has shed light on this with the

derivation of a synthetic form of TOPBP1 that is a stripped-down version lacking all

sequences irrelevant to ATR signaling (Ruis et al, 2021). This protein, termed

TOPBP1 Jr, contains (1) the first three BRCT domains, which are needed for

recruitment to DSBs, (2) the AAD, and (3) the final two BRCT domains, which are

required for multimerization (Figure 1A). Other works have shown that sequences

within the AAD itself are important for regulation of TOPBP1. For example, it has

been shown that serine 1131 in Xenopus TOPBP1 must be phosphorylated by the

ataxia telangiectasia mutated (ATM) kinase in order for TOPBP1 to activate ATR

(Yoo et al, 2007). Other, more recent work has suggested that phosphorylation of

this same serine residue in human TOPBP1 causes the protein to undergo liquid-

liquid phase separation to create TOPBP1 condensates at sites of damage (Frattini et

al, 2021). These condensates are micron-sized, contain millions of TOPBP1

molecules, and are proposed to amplify ATR signaling.

In this work we use our <u>D</u>SB-<u>m</u>ediated <u>A</u>TR activation in *Xenopus* egg extract

system (DMAX; Montales et al, 2021; Ruis et al, 2022; Montales et al, 2022) to study

how TOPBP1's AAD is regulated. DMAX offers several powerful advantages for

mechanistic studies of ATR signaling. For example, ATR signaling is readily assessed

by scoring for phosphorylation of the key ATR substrate CHK1. Recruitment to DSBs

is also directly assayed in DMAX using DSBs that are immobilized on magnetic beads

in a pull-down assay. Structure-function analysis is another key feature of DMAX as

target proteins can be removed from *Xenopus* egg extract (XEE) via

immunodepletion and mutant forms, rapidly and easily produced by *in vitro*

transcription and translation (IVTT), are added back for phenotypic assessment.

Using DMAX, we show here that S1131 in the AAD plays an important role in

TOPBP1 multimerization independently of condensate formation. We also identify a

novel site of phosphorylation within the AAD, T1098, and show that this residue is

required for ATR activation. Lastly, we provide evidence that casein kinase 2 is

required for ATR activation.

2. Material and Methods

2.1 Materials

2.1.1 Plasmids

All plasmids used in this study are described below, in table format, and all are

Xenopus proteins. Specific details about plasmid construction are available upon

request.

Plasmid	Name	Vector	AA coordinates
5Xmyc-TOPBP1	Cut5	pCS2+MT	1-1513
5Xmyc-TOPBP1 S1131A	pSY29	pCS2+MT	1-1513
5Xmyc-TOPBP1 T1098A	pHS60	pCS2+MT	1-1513
5Xmyc-AAD-BRCT7&8	pHG133	pCS2+MT	971-1513
5Xmyc-TOPBP1 Jr	pKM116	pCS2+MT	1-333 & 992-1513
5Xmyc-AAD	pKR4	pCS2+MT	971-1279
5Xmyc-BRCT7&8	pKM1	pCS2+MT	1235-1470
GST-AAD	pHG117	pGEX-4T3	971-1279
GST-AAD S1131A	pKR10	pGEX-4T3	971-1279
GST-AAD S1131D	pKR11	pGEX-4T3	971-1279
GST-AAD-BRCT7&8	pKR32	pGEX-4T3	971-1513
GST-BRCT7&8	pHG08	pCS2+MT	1197-1513

2.1.2 Recombinant proteins

The recombinant proteins used in this study were GST, GST-AAD (wild type,

S1131A, S1131D), GST-BRCT7&8, GST-AAD-0BRCT7&8). All proteins were

expressed in *E. coli* BL21(DE3) cells, at 37°C for 4 hours, and all proteins were

purified from the soluble fraction according to standard procedures (detailed

below).

2.1.3 Antibodies

We used the following commercially sourced antibodies in this work: Myc

(Millipore Sigma #M4439) and GST (Millipore Sigma #05-782). We also used our

own antibody against *Xenopus* TOPBP1, HU142, which has been described (Van

Hatten et al., 2002).

2.2 Methods

2.2.1 Xenopus egg extracts

The high-speed supernatant (HSS) of XEE was used exclusively in this study. HSS

was prepared exactly as described (Smythe and Newport, 1991). For

immunodepletion of TOPBP1, the HU142 antibody was used and the procedure was

performed exactly as described (Van Hatten et al, 2002). Depleted XEEs were then

supplemented with IVTT- produced proteins (2.5 µl IVTT lysate in 20 ul of XEE) as

described (Montales et al, 2021).

2.2.2 IVTT production of proteins and determination of protein concentrations

IVTT reactions were performed using the SP6 TnT® Quick Coupled

Transcription/Translation System (Promega #L2080) according to the

manufacturer's instructions.

2.2.3 Recombinant Protein Expression and Purification

Plasmids were transformed in BL21 (DE3) Competent *E. Coli* cells and grown in LB

media (GST fusion proteins) or LB media with 0.2% glucose (MBP fusion proteins)

with ampicillin. At an OD600 of 0.6-0.8, cells were induced with 0.3 mM IPTG for 4

hours at 37°C. Cells were then harvested by centrifugation (4000 rpm for 20 min)

and washed with phosphate-buffered saline (PBS). Cells were lysed by sonication in

GST column buffer (40% PBS, 60% TEN buffer: 10 mM Tris-Cl, pH 8.0, 1 mM EDTA,

pH 8.0, 100 mM NaCl) supplemented with cOmplete protease inhibitor cocktail

tablets (Roche #04693159001) and 10 mg lysozyme. Lysates were cleared by

centrifugation (15,000 rcf for 20 min) and incubated with glutathione sepharose

(BioVision #6566) for 1 hour at 4°C. Beads were poured over a column and washed

with column buffer. GST fusion proteins were eluted with 20 mM glutathione in 75

mM HEPES, pH 7.5, 150 mM NaCl, 5 mM DTT, and 0.1% Triton-X. Proteins were

dialyzed overnight into PBS.

2.2.4 ATR activation (DMAX) assays

For DMAX assays, okadaic acid was first mixed with 20 µl of HSS to a final

concentration of 1 µM, as described (Montales et al, 2021). Linear dsDNA derived

from EcoRI-digested lambda DNA (Montales et al, 2021) was then added to the

mixture and reactions were incubated at room temperature for 30 min. Samples

were analyzed via Western blotting using standard conditions. In some cases, when

GST-AAD proteins were used, then no DNA was included in the assay. For the

experiments shown in Figure 4, the casein kinase 2 inhibitor CX-4945 was used, and

it was purchased from Selleckchem.com (catalog # S2248).

2.2.5 DSB binding assay.

These were performed exactly as described (Montales et al, 2021). Briefly, a 5 kb

PCR fragment was biotinylated via inclusion of a biotin moiety on the forward PCR

primer. After PCR and cleanup, the 5 kb PCR fragment was coupled to magnetic

streptavidin beads (Dyna- beads M-270 Streptavidin, ThermoFisher) according to

the manufacturer's instructions. These "DSB beads", containing 600 fmol of dsDNA

per assay, were then incubated in 20 μl of HSS supplemented with 5 μl of an IVTT

reaction programmed to produce the protein of interest. After incubation the beads

were collected on a magnetic stand and washed three times in PBS + 0.1% TritionX-

100. Bound proteins were then eluted with 2X SDS-PAGE sample buffer and

examined by Western blotting.

2.2.6 Protein binding assays.

For the GST pull-down assays, 10– 20 μL of IVTT proteins were mixed with 5–10 u g

of GST fusion protein and incubated for 60 min at 30 ∘C. The mixtures were diluted

into 400 μL of Binding Buffer (PBS + 0.1 % NP-40) and then incubated with

glutathione sepharose (BioVision #6566) for one hour at 4 ∘C. The beads were

washed four times with 500 μL of Binding Buffer and then eluted with 2X Sample

Buffer (Millipore Sigma #S3401-10VL). The input and bound fractions were then

analyzed via Western blotting using standard conditions.

Mass spectrometry analysis of GST-AAD

GST-AAD was incubated in XEE for 30 minutes at room temperature and purified

back out of the extract via glutathione-sepharose affinity chromatography. The

sample was run on SDS-PAGE and a Coomassie-stained band was excised from the

gel. Proteins in coomassie-stained gel slices were reduced, alkylated and digested by

trypsin as previously described (Kaiser and Wohlschlegel, 2005). After digestion,

peptides were extracted from the gel slices using 100 µl of 60% acetonitrile/0.1%

TFA, dried in a SpeedVac vacuum concentrator, desalted using C18 tips, and

reconstituted in 10 µl of 5% formic acid for LC-MS/MS analysis.

Desalted peptide mixtures were loaded onto a 25 cm long, 75 µm inner diameter

fused-silica capillary, packed in-house with bulk 1.9 µM ReproSil-Pur beads with

120 Å pores as described previously (Jami-Alahmadi et al, 2021). Peptides were

analyzed using a 140 min water-acetonitrile gradient delivered by a Dionex Ultimate

3000 UHPLC (Thermo Fisher Scientific) operated initially at 400 nL/min flow rate

with 1% buffer B (acetonitrile solution with 3% DMSO and 0.1% formic acid) and

99% buffer A (water solution with 3% DMSO and 0.1% formic acid). Buffer B was

increased to 6% over 5 min at which time the flow rate was reduced to 200 nl/min.

A linear gradient from 6-28% B was applied to the column over the course of 123

min. The linear gradient of buffer B was then further increased to 28-35% for 8 min

followed by a rapid ramp-up to 85% for column washing. Eluted peptides were

ionized via a Nimbus electrospray ionization source (Phoenix S&T) by application of

a distal voltage of 2.2 kV.

All mass spectrometry data were collected using data dependent acquisition on

Orbitrap Fusion Lumos Tribrid mass spectrometer (Thermo Fisher Scientific) with

an MS1 resolution of 120,000 followed by sequential MS2 scans at a resolution of

15,000. The Integrated Proteomics pipeline 2 was utilized to generate peptide and

protein identifications. MS2 spectra were searched using the ProLuCID algorithm

against the Xenopus reference proteome (xenbase.org) followed by filtering by

DTASelect using a decoy database estimated false discovery rate of < 1% (Xu et al,

2015; Cociorva et al, 2007).

3. Results and Discussion

3.1 Serine 1131 promotes TOPBP1's ability to activate ATR in a manner independent of condensate formation.

Previous work from the Dunphy group had identified S1131 in TOPBP1's AAD as an ATM phospho-acceptor site that is required for ATR activation in XEE under some conditions (Yoo et al, 2007). S1131 is required for ATR activation by small homopolymeric oligonucleotides like A_{70}-T_{70} (AT70), and by sperm chromatin digested with EcoRI to produce DSBs. Interestingly, S1131 is dispensable for ATR activation by stalled replication forks. We thus initiated our studies by checking the requirement for S1131 in our DMAX system. To do so, we performed immunodepletion, where either non-specific or anti-TOPBP1 antibodies are coupled to Protein A beads and then incubated with XEE. After incubation, the XEE is recovered away from the antibody beads containing the captured TOPBP1 and used for ATR activation assays. EcoRI-digested λ DNA was used as a source of DSBs. We also prepared recombinant forms of TOPBP1, using IVTT in rabbit reticulocyte lysates, so that they could be added back to the depleted extracts. As shown in Figure 1B, a mock-depleted extract (treated with non-specific antibody beads, lane 1, labeled "TOPBP1⁺ XEE") contained TOPBP1 and was able to activate ATR. By contrast, for the TOPBP1-depleted extract, TOPBP1 levels were substantially reduced, and this extract was deficient for ATR activation (Figure 1B, lane 2). When wild type TOPBP1 was added back to the TOPBP1-depleted extract, then ATR activation was restored (lane 3), and this did not occur when an S1131A mutant form of TOPBP1 was added back (lane 4). This experiment shows that S1131 is

required for ATR activation by TOPBP1 in our DMAX system. Previous work had not

addressed if S1131 is important for TOPBP1 recruitment to DSBs, and thus we

employed our DSB binding assay to examine this important issue. DSB beads were

prepared as previously described (Montales et al, 2021), where linear, 5kb dsDNAs

with biotin moieties attached to end are immobilized on magnetic streptavidin

beads. The beads are then incubated in XEE containing myc-tagged forms of either

wild type or S1131A TOPBP1. Since our previous work has repeatedly shown that

TOPBP1 does not bind to streptavidin beads lacking DNA under these assay

conditions (Montales et al, 2021; Ruis et al, 2022; Montales et al, 2022), we did not

include empty beads in this experiment. After incubation, the beads were isolated,

washed, and probed for TOPBP1 occupancy by Western blotting. As shown in Figure

1C, both wild type and S1131A TOPBP1 could bind to DSBs. We conclude S1131 is

dispensable for recruitment of TOPBP1 to DSBs, and likely functions after

recruitment to stimulate DNA-bound ATR kinase.

What is the role of S1131 in ATR activation? Previous work has shown that

phosphorylation of S1131 (to produce P-S1131) makes binding of TOPBP1 to ATR-

ATRIP more efficient (Yoo et al, 2007). More recently, it has been shown for human

TOPBP1 that P-S1138, the analogous residue, is required for TOPBP1's ability to

phase separate (Frattini et al, 2021). This, in turn, promotes robust ATR signaling.

Importantly, although the human AAD contains S1138, the AAD by itself cannot

undergo phase transition as additional sequences are required (Frattini et al, 2021).

The AAD by itself can, however, activate ATR when added as a GST fusion protein to

XEE (Kumagai et al, 2006). Given these circumstances, we reasoned that if S1131

functions exclusively in the context of condensate formation then it should be

irrelevant to the ability of the isolated AAD, which cannot form a condensate, to

activate ATR. To put this notion to the test, we prepared GST-AAD fusion proteins of

three types: wild type, S1131A, and S1131D. Previous work has shown that the

S1131D substitution, which mimics P-S1131, produces a dominant, gain-of-function

TOPBP1 that binds ATR-ATRIP more efficiently and activates ATR more readily (Yoo

et al, 2007). This previous work did not, however, test the S1131D mutation in the

context of the isolated AAD. The three different GST-AAD proteins were added to

XEE at the indicated concentrations and ATR signaling was assessed by anti P-CHK1

Western blotting (Figure 1D). The S1131A mutant produced less P-CHK1 than did

the wild type version when both were present at 4 uM (compare lanes 2 and 4 in

Figure 1D). This shows that S1131 is important for the ability of the isolated AAD to

activate ATR, and thus that S1131 can function independently of condensate

formation to promote ATR signaling. Interestingly, we observed the opposite effect

for the S1131D mutation which hyper-activated ATR (compare lanes 2 and 5 in

Figure 1D). This is consistent with previous data showing that S1131D in the

context of full-length TOPBP1 promotes ectopic activation of ATR (Yoo et al, 2007),

and our data extend these previous data by showing that S1131D can exert this

effect in the context of the AAD alone. Based on the data in Figure 1, we conclude

that the ability of S1131 to promote ATR activation is an intrinsic property of the

AAD, and thus can be disconnected from condensate formation. This is not to say

that it is not involved in condensate formation, but it is clearly doing something

separate from phase transition.

3.2 S1131 promotes TOPBP1 multimerization.

In recent work we used a synthetic form of TOPBP1, TOPBP1 Jr, to show that

just three functional domains are needed for regulated activation of ATR by TOPBP1

(Ruis et al, 2021). The BRCT0-2 domains mediate recruitment and are dispensable

thereafter while the AAD interfaces with ATR-ATRIP and the BRCT7&8 domains

promote multimerization in an unknown manner. To determine if S1131 is possibly

connected to TOPBP1 multimerization we first asked if the BRCT7&8 domains of

TOPBP1 could interact with the AAD. For this, a standard GST pull-down assay and

the GST fusion proteins we used are shown in Figure 2A. These proteins were used

in binding assays with myc-tagged BRCT7&8, produced by IVTT. As shown in Figure

2B, myc-BRCT7&8 did not bind to GST alone, but did bind to GST-AAD, GST-

BRCT7&8, and a fragment containing both the AAD and BRCT7&8. We next

examined how the same set of GST proteins would bind to myc-tagged AAD alone,

and again detectable binding was observed for all of the GST fusions but not GST

alone (Figure 2C). In this experiment, however, the binding efficiencies differed,

with the more efficient binding observed for GST-BRCT7&8, less efficient binding to

GST-AAD-BRCT7&8, and relatively weak binding to GST-AAD. These data show that

the AAD domain interacts with itself weakly, and that it binds BRCT7&8 more

robustly (Figure 2D). In addition, the data show that BRCT7&8 can self-associate.

We note that others have previously shown that the AAD can interact with itself

(Thada et al, 2019).

To pursue these observations, we next asked how S1131 would impact

interaction of the ADD with the TOPBP1 Jr. We chose TOPBP1 Jr for this analysis

because, unlike the other constructs, TOPBP1 Jr has been shown to activate ATR in

the context of DSBs (Ruis et al, 2021). We used the same GST proteins for this

experiment as were used in Figure 1D. As shown in Figure 2E, myc-tagged TOPBP1

Jr could bind to the wild type AAD, and binding was somewhat reduced with the

S1131A mutant. Interestingly, binding was increased, relative to wild type, with the

S1131D mutant. Thus the same phospho-mimic mutation that increases ATR

activation by the AAD also increases multimerization. We conclude that promoting

multimerization is a previously undisclosed function of S1131.

3.3 T1098 within the AAD is phosphorylated in XEE and required for ATR activation.

During the course of our studies of TOPBP1's AAD, we noticed that, after

incubation in XEE, that GST-AAD protein migrated with reduced mobility on SDS-

PAGE. This can be seen in Figure 3A, where GST-AAD that had been incubated in

XEE and then recovered back out of the extract using glutathione sepharose beads

was run side-by-side with the input material. This reduced mobility suggests that

post-translational modifications (PTMs) had been placed on GST-AAD during

incubation in XEE, and thus to gain a sense of what these PTMs are we excised the

band in Figure 3A, lane 3, and submitted it to analysis by mass spectrometry. This

analysis revealed several peptides containing phosphorylated serine or threonine

residues (Figure 3B). Figure 3C shows the sequence of the AAD from *Xenopus*

TOPBP1 and these candidate phospho-acceptor sites are highlighted in red. The

underlined portion shows the region in the frog protein that corresponds to the

minimally defined AAD from human TOPBP1 (Thada et al, 2019), and thus two of

the candidate phospho-acceptor sites (S1095 and T1098) fall within this region. We

note that S1131 is only phosphorylated during a DNA damage response (Yoo et al,

2007), and thus we did not expect to recover P-S1131 signals in this experiment.

To pursue these observations, we next asked two questions – are the

candidate phospho-acceptor sites identified by the mass spectrometry analysis

conserved in human TOPBP1 and, if so, have any of those sites been curated on the

website PhosphoSitePlus® (www.phosphosite.org), which catalogs phospho-sites

for all human proteins? The answers are displayed in Figure 3D, where seven of the

nine sites identified by mass spectrometry are conserved in human TOPBP1 and

two of these, S1008 and T1098, are listed on PhosphoSitePlus®. Because T1098 is

conserved in humans, listed on PhosphoSitePlus®, and falls within the minimally

defined AAD, we focused on it first. The MS/MS spectrum for the peptide harboring

pT1098 is displayed in Figure 4A. To determine if this residue is important for ATR

activation, we produced a T1098A point mutant in full-length TOPBP1 and we

tested this mutant for ATR activation, exactly as we did for S1131 in Figure 1B. As

shown in Figure 4B, the T1098A mutant fails to activate ATR. This result shows that

S1131 is not the only critical phospho-acceptor site within the AAD, as T1098 is also

important.

3.4 A casein kinase 2 inhibitor blocks both DSB- and ADD-mediated ATR activation.

One feature of the AAD region of TOPBP1 is that it contains an unusually high

number of consensus sequences for phosphorylation by casein kinase 2. The

consensus motif for casein kinase 2 is S/T-X-X-D/E (St-Denis et al, 2015), and this

motif is found nine times in the AAD region (Figure 5A), and one of these (S1266)

was identified in our mass spectrometry analysis and is conserved in human

TOPBP1. This prompted us to ask if casein kinase 2 activity is required for the AAD

to function. For this, we asked if DSBs could still activate ATR in XEEs treated with

the casein kinase 2 inhibitor (CK2i) CX-4945. We found that CX-4945 could block

ATR activation (Figure 5B). We have previously shown that CX-4945 does not

impair the recruitment of TOPBP1 to DSBs (Montales et al, 2022) and thus we

considered the possibility that the AAD requires casein kinase 2 activity in order to

function. To test this idea we added GST-AAD to XEE and observed that CX-4945

could still inhibit ATR activation. Based on this experiment, we conclude that the

AAD requires, either directly or indirectly, casein kinase 2 activity in order to

activate ATR.

In this paper we present data showing that phosphorylation of TOPBP1's

AAD is crucial for its ability to activate ATR. Previous work had identified S1131 as a

regulated phospho-acceptor site necessary for ATR activation by DSBs, but not

stalled replication forks. This previous work showed that S1131 is phosphorylated

by ATM, and that the manifestation of this phosphorylation was increased affinity

for ATR-ATRIP (Yoo et al, 2007). Here, we extend these observations by showing

that S1131 is important for multimerization of the AAD (Figure 3E). We also report

that an additional residue within the AAD, T1098, is also phosphorylated, but this is

constitutive phosphorylation as it occurs in the absence of a DDR response. Like

S1131, phosphorylation of T1098 is required for ATR activation. Summarizing these

findings, we see that under resting conditions, TOPBP1 is primed for action via

constitutive phosphorylation on T1098; however in its monomeric form, it is unable

to activate ATR (Figure 7). ATM-directed phosphorylation of S1131 occurs in

response to DSBs, and now TOPBP1 can assume the multimeric form that is

required for ATR activation (Figure 7). It is likely that additional sites within the

AAD are also phosphorylated, as this region contains multiple casein kinase 2

consensus motifs and we have shown here that casein kinase 2 activity is required

for AAD function (Figure 6). In conclusion, our data show that the AAD undergoes

multi-site phosphorylation, under both constitutive and regulated conditions, and

that at least some of these phosphorylations convert TOPBP1 from an inactive to an

active form via multimerization.

Author contribution

W.M.M. conceptualized the project, acquired the funding, and wrote the manuscript.

H.S., K.R., and K.M. performed the experiments. J.A.W. and Y.J. performed the mass

spectrometry experiments.

Conflict of interest statement

The authors declare no conflict of interest.

Acknowledgments

This work was supported by NIH grant R01GM12287 (W.M.M.) and R01GM089778

(J.A.W.). We thank Shan Yan and Hovik Gasparyan for the production of plasmid

constructs used in this study.

References
408
Bartkova J, Rezaei N, Liontos M, Karakaidos P, Kletsas D, Issaeva N, Vassiliou LV,
Kolettas E, Niforou K, Zoumpourlis VC, Takaoka M, Nakagawa H, Tort F, Fugger K,
Johansson F, Sehested M, Andersen CL, Dyrskjot L, Ørntoft T, Lukas J, Kittas C,
Helleday T, Halazonetis TD, Bartek J, Gorgoulis VG. Oncogene-induced senescence is
part of the tumorigenesis barrier imposed by DNA damage checkpoints. Nature.
2006 Nov 30;444(7119):633-7. doi: 10.1038/nature05268. PMID: 17136093.
415
416   Cociorva D, L Tabb D, Yates JR. Validation of tandem mass spectrometry database
417   search results using DTASelect. Curr Protoc Bioinformatics. 2007 Jan;Chapter
418   13:Unit 13.4. doi: 10.1002/0471250953.bi1304s16. PMID: 18428785.
419
420   Di Micco R, Fumagalli M, Cicalese A, Piccinin S, Gasparini P, Luise C, Schurra C, Garre'
421   M, Nuciforo PG, Bensimon A, Maestro R, Pelicci PG, d'Adda di Fagagna F. Oncogene-
422   induced senescence is a DNA damage response triggered by DNA hyper-replication.
423   Nature. 2006 Nov 30;444(7119):638-42. doi: 10.1038/nature05327. PMID:
424   17136094.
425
426   Frattini C, Promonet A, Alghoul E, Vidal-Eychenie S, Lamarque M, Blanchard MP,
427   Urbach S, Basbous J, Constantinou A. TopBP1 assembles nuclear condensates to
428   switch on ATR signaling. Mol Cell. 2021 Mar 18;81(6):1231-1245.e8. doi:
429   10.1016/j.molcel.2020.12.049. Epub 2021 Jan 26. PMID: 33503405.
430
431   Jami-Alahmadi Y, Pandey V, Mayank AK, Wohlschlegel JA. A Robust Method for
432   Packing High Resolution C18 RP-nano-HPLC Columns. J Vis Exp. 2021 May 14;(171).
433   doi: 10.3791/62380. PMID: 34057454.
434
435   Kaiser P, Wohlschlegel J. Identification of ubiquitination sites and determination of
436   ubiquitin-chain architectures by mass spectrometry. Methods Enzymol.
437   2005;399:266-77. doi: 10.1016/S0076-6879(05)99018-6. PMID: 16338362.
438
439   Kim A, Montales K, Ruis K, Senebandith H, Gasparyan H, Cowan Q, Michael WM.
440   Biochemical analysis of TOPBP1 oligomerization. DNA Repair (Amst). 2020
441   Dec;96:102973. doi: 10.1016/j.dnarep.2020.102973. Epub 2020 Sep 21. PMID:
442   32987353; PMCID: PMC7670859.
443
444   Kumagai A, Lee J, Yoo HY, Dunphy WG. TopBP1 activates the ATR-ATRIP complex.
445   Cell. 2006 Mar 10;124(5):943-55. doi: 10.1016/j.cell.2005.12.041. PMID: 16530042.
446
447   Montales K, Kim A, Ruis K, Michael WM. Structure-function analysis of TOPBP1's
448   role in ATR signaling using the DSB-mediated ATR activation in Xenopus egg
449   extracts (DMAX) system. Sci Rep. 2021 Jan 11;11(1):467. doi: 10.1038/s41598-020-
450   80626-1. PMID: 33432091; PMCID: PMC7801695.
451

452   Montales K, Ruis K, Lindsay H, Michael WM. MRN-dependent and independent
453   pathways for recruitment of TOPBP1 to DNA double-strand breaks. PLoS One. 2022
454   Aug 2;17(8):e0271905. doi: 10.1371/journal.pone.0271905. PMID: 35917319;
455   PMCID: PMC9345342.
456
457   Niedzwiedz W. Activating ATR, the devil's in the dETAA1l. Nat Cell Biol. 2016 Oct
458   27;18(11):1120-1122. doi: 10.1038/ncb3431. Erratum in: Nat Cell Biol. 2016 Nov
459   29;18(12):1367. PMID: 27784903.
460
461   Ruis K, Huynh O, Montales K, Barr NA, Michael WM. Delineation of a minimal
462   topoisomerase II binding protein 1 for regulated activation of ATR at DNA double-
463   strand breaks. J Biol Chem. 2022 Jul;298(7):101992. doi:
464   10.1016/j.jbc.2022.101992. Epub 2022 Apr 28. PMID: 35490781; PMCID:
465   PMC9257406.
466
467   Saldivar JC, Cortez D, Cimprich KA. The essential kinase ATR: ensuring faithful
468   duplication of a challenging genome. Nat Rev Mol Cell Biol. 2017 Oct;18(10):622-
469   636. doi: 10.1038/nrm.2017.67. Epub 2017 Aug 16. Erratum in: Nat Rev Mol Cell
470   Biol. 2017 Dec;18(12):783. PMID: 28811666; PMCID: PMC5796526.
471
472   Saldivar JC, Hamperl S, Bocek MJ, Chung M, Bass TE, Cisneros-Soberanis F, Samejima
473   K, Xie L, Paulson JR, Earnshaw WC, Cortez D, Meyer T, Cimprich KA. An intrinsic
474   S/G2 checkpoint enforced by ATR. Science. 2018 Aug 24;361(6404):806-810. doi:
475   10.1126/science.aap9346. PMID: 30139873; PMCID: PMC6365305.
476
477   Smythe C, Newport JW. Systems for the study of nuclear assembly, DNA replication,
478   and nuclear breakdown in Xenopus laevis egg extracts. Methods Cell Biol.
479   1991;35:449-68. doi: 10.1016/s0091-679x(08)60583-x. PMID: 1664032.
480
481   St-Denis N, Gabriel M, Turowec JP, Gloor GB, Li SS, Gingras AC, Litchfield DW.
482   Systematic investigation of hierarchical phosphorylation by protein kinase CK2. J
483   Proteomics. 2015 Apr 6;118:49-62. doi: 10.1016/j.jprot.2014.10.020. Epub 2014
484   Nov 5. PMID: 25449829.
485
486   Thada V, Cortez D. Common motifs in ETAA1 and TOPBP1 required for ATR kinase
487   activation. J Biol Chem. 2019 May 24;294(21):8395-8402. doi:
488   10.1074/jbc.RA119.008154. Epub 2019 Apr 2. PMID: 30940728; PMCID:
489   PMC6544864.
490
491   Thada V, Cortez D. ATR activation is regulated by dimerization of ATR activating
492   proteins. J Biol Chem. 2021 Jan-Jun;296:100455. doi: 10.1016/j.jbc.2021.100455.
493   Epub 2021 Feb 24. PMID: 33636182; PMCID: PMC7994790.
494
495   Van Hatten RA, Tutter AV, Holway AH, Khederian AM, Walter JC, Michael WM. The
496   Xenopus Xmus101 protein is required for the recruitment of Cdc45 to origins of
497   DNA replication. J Cell Biol. 2002 Nov 25;159(4):541-7. doi:

498   10.1083/jcb.200207090. Epub 2002 Nov 18. PMID: 12438414; PMCID:
499   PMC2173091.
500
501   Waterman DP, Haber JE, Smolka MB. Checkpoint Responses to DNA Double-Strand
502   Breaks. Annu Rev Biochem. 2020 Jun 20;89:103-133. doi: 10.1146/annurev-
503   biochem-011520-104722. Epub 2020 Mar 16. PMID: 32176524; PMCID:
504   PMC7311309.
505
506   Xu T, Park SK, Venable JD, Wohlschlegel JA, Diedrich JK, Cociorva D, Lu B, Liao L,
507   Hewel J, Han X, Wong CCL, Fonslow B, Delahunty C, Gao Y, Shah H, Yates JR 3rd.
508   ProLuCID: An improved SEQUEST-like algorithm with enhanced sensitivity and
509   specificity. J Proteomics. 2015 Nov 3;129:16-24. doi: 10.1016/j.jprot.2015.07.001.
510   Epub 2015 Jul 11. PMID: 26171723; PMCID: PMC4630125.
511
512   Yoo HY, Kumagai A, Shevchenko A, Shevchenko A, Dunphy WG. Ataxia-telangiectasia
513   mutated (ATM)-dependent activation of ATR occurs through phosphorylation of
514   TopBP1 by ATM. J Biol Chem. 2007 Jun 15;282(24):17501-6. doi:
515   10.1074/jbc.M701770200. Epub 2007 Apr 19. PMID: 17446169.
516
Zhou ZW, Liu C, Li TL, Bruhn C, Krueger A, Min W, Wang ZQ, Carr AM. An essential
function for the ATR-activation-domain (AAD) of TopBP1 in mouse development
and cellular senescence. PLoS Genet. 2013;9(8):e1003702. doi:
10.1371/journal.pgen.1003702. Epub 2013 Aug 8. PMID: 23950734; PMCID:
PMC3738440.

Figures and Legends

Figure 1. S1131 in TOPBP1's AAD is dispensable for recruitment of TOPBP1 to

DSBs but is required for ATR activation in the DMAX system.

A. Schematic representation of full length TOPBP1 and TOPBP1 Junior. **B.** A DMAX

assay where XEE was immunodepleted of endogenous TOPBP1 and then

supplemented with the indicated IVTT-produced TOPBP1 proteins prior to the

addition of DSBs. TOPBP1⁺ XEE (lane 1) refers to extract that was mock-depleted

using non-specific IgG. Blank (lane 2) refers to a TOPBP1-depleted extract that

received an unprogrammed IVTT reaction. After incubation, the samples were

probed by Western blotting for P-CHK1, CHK1, and TOPBP1 (using anti-TOPBP1

antibody HU142). The experiment shown is representative of two independently

performed biological replicates. **C.** HSS was combined with IVTT-produced and myc-

tagged TOPBP1 proteins. These proteins were the wild type form ("WT") and the

S1131A mutant. DSB beads were then added and the samples were incubated for 30

min. After incubation, the beads were washed and eluted and the samples were

probed by Western blot for the indicated protein. The myc antibody was used to

probe for TOPBP1. The bound samples were also stained by silver and a 12.5 kDa

low molecular weight (l.m.w) protein is shown, to demonstrate equal isolation of the

beads. The experiment shown is representative of two independently performed

biological replicates. **D.** The indicated GST proteins were added to XEE at the

indicated concentrations and incubation was carried out for 30 minutes at room

temperature. The samples were then probed for P-CHK1, CHK1, and GST. The

experiment shown is representative of two independently performed biological

replicates.

Figure 2. S1131 is important for multimerization of the AAD.

A. Schematic representation of the proteins used for protein binding assays. **B.** The

indicated GST proteins, expressed and purified from *E. coli*, were incubated in IVTT

reactions expressing myc-tagged BRCT7&8. After incubation the samples were

diluted in Binding Buffer and incubated with glutathione-sepharose beads. The

beads were then washed, and the samples were eluted and probed by Western

blotting for the indicated proteins. A sample of the total binding reaction, taken

prior to dilution, was also probed for myc (lane "input"). The experiment shown is

representative of two independently performed biological replicates. **C.** Same as

(B) except myc-AAD was used. The experiment shown is representative of two

independently performed biological replicates. **D.** Schematic showing how the AAD-

BRCT7&8 region interacts with itself. **E.** Same as (B) except myc-TOPBP1 Jr was

used with the indicated GST fusion proteins. The experiment shown is

representative of two independently performed biological replicates.

Figure 3. Mass spectrometry reveals multiple phospho-acceptor sies within

the AAD region.

A. A Coomassie blue stained SDS-PAGE gel showing altered mobility of GST-AAD

after incubation in XEE.   **B.** The amino acid sequence of the AAD region used for

mass spectrometry analysis. See main text for details. **C.** Summary of conservation

and PhosphoSitePlus® hits for the phospho-acceptor sites identified by mass

spectrometry.

Figure 4. T1098 is required for ATR activation by TOPBP1.

A. MS/MS spectra for the peptide harboring T1098. **B.** Same as Figure 1B except the

T1098A mutant was used. The experiment shown is representative of two

independently performed biological replicates.

Figure 5. Delineation of the NBS1 binding region within TOPBP1's BRCT1

domain.

A. Same as Figure 3C except the position of consensus casein kinase 2

phosphorylation sites is highlighted with grey shading. **B-C.** XEEs were treated as

indicated with either DMSO or CK2i and either DSBs (B.) or the indicated GST

proteins (C.) were added. After incubation the samples were probed by Western

blotting for P-CHK1 and CHK1. The experiments shown are representative of two

independently performed biological replicates.

Figure 6. Multi-site phosphorylation of the TOPBP1 AAD drives ATR activation

through a multimerization mechanism.

Please see text for details.

 Figure 1

591   **Figure 2**

594 **Figure 3**

A

B

^{972}EQQKRVPEALYPHTYNPKMSLDISAVQDGSYTASKF**S**AD**T**SLQQDENSELQLQQ
NNKFGE**T**SDDQVKKAAGDGNPQNPSKDVKGALTQTLEMRENFQRQLQEFMSATS
VVKPRGSVGRAGFDNS**S**PC**T**PEGARSTRNGRSRVLEALRQSRQAMTDLNTEPSQNE
QII**W**DDPTAREERAKLVSNLQWPDSPSQYSEQLQHNMNDAGGNYTPAKESLTDSEI
AELEACEFEPKSAMRTPVIENNLQSPTKPDHLTPTPQAPSIAFPLANPPVAPQPREK
PVQPFSKEETLKERRFQL**SS**LDPQERIDYSQL^{1167}

C

frog site	human equivalent	PhosphoSitePlus® hits
S1008	S1016	2
T1011	S1019	0
T1031	not conserved	na
S1095	not conserved	na
T1098	T1105	1
S1266	S1273	0
S1267	S1274	0

597   **Figure 4**

Figure 5

A

^{972}EQQKRVPEALYPHTYNPKMSLDISAVQDGSYTASKFSADTSLQQDENSELQLQQ
NNKFGETSDDQVKKAAGDGNPQNPSKDVKGALTQTLEMRENFQRQLQEFMSATS
VVKPRGSVGRAGFDNSPCTPEGARSTRNGRSRVLEALRQSRQAMTDLNTEPSQNE
QIIWDDPTAREERAKLVSNLQWPDSPSQYSEQLQHNMNDAGGNYTPAKESLTDSEI
AELEACEFEPKSAMRTPVIENNLQSPTKPDHLTPTPQAPSIAFPLANPPVAPQPREK
PVQPFSKEETLKERRFQLSSLDPQERIDYSQL^{1167}

603   **Figure 6**