Expression of thioredoxin-1 in the ASJ neuron corresponds with and enhances intrinsic regenerative capacity under lesion conditioning in *C. elegans*

Noa W.F. Grooms, Michael Q. Fitzgerald, Binyamin Zuckerman, Samuel E. Ureña, Leor S. Weinberger, Samuel H. Chung

Abstract

A conditioning lesion of the peripheral sensory axon triggers robust central axon regeneration in mammals. We trigger conditioned regeneration in the *C. elegans* ASJ neuron by laser surgery or genetic disruption of sensory pathways. Conditioning upregulates *trx-1* expression, as indicated by *trx-1* promoter-driven green fluorescent protein and fluorescence in situ hybridization, suggesting *trx-1* levels and associated fluorescence indicate regenerative capacity. Redox activity of *trx-1* functionally enhances conditioned regeneration, but both redox-dependent and -independent activity inhibit non-conditioned regeneration. Six strains isolated in a forward genetic screen for reduced fluorescence, which suggests diminished regenerative potential, also show reduced axon outgrowth. We demonstrate an association between *trx-1* expression and the conditioned state that we leverage to rapidly assess regenerative capacity.

Keywords

neuron regeneration; lesion conditioning; thioredoxin; genetic screen; *C. elegans*; fluorescence microscopy; redox biology

Abbreviations

CNS, central nervous system; EMS, ethyl methanesulfonate; DLK, dual-leucine kinase; TXN, thioredoxin; cAMP, cyclic adenosine monophosphate; CRE, cAMP responsive element; CRE-binding protein (CREB),

27   RAG, regeneration associated gene; RNAi, ribonucleic acid interference; a+d, concomitant axon + dendrite

28   surgery; FISH, fluorescence in situ hybridization

Introduction

The exquisitely complex and precise shape of neurons makes them highly susceptible to injury and disease. Compounding this weakness is the extremely limited regenerative capacity of the mammalian central nervous system (CNS), which severely limits functional recovery. Millions of people worldwide are affected by traumatic brain or spinal cord injury, amyotrophic lateral sclerosis (ALS), Alzheimer's, Parkinson's, or other neurodegenerative disorders (Feigin*, et al.*, 2021; James*, et al.*, 2019). The lack of effective treatments for these afflictions represents one of the areas of greatest need in modern medicine.

In a remarkable phenomenon known as lesion conditioning, damage to a neuron's peripheral sensory fiber triggers cellular mechanisms to drive regeneration in the peripheral and central nervous systems (McQuarrie and Grafstein, 1973; Richardson and Issa, 1984). Lesion conditioning fosters a pro-regenerative environment (Dubový*, et al.*, 2019; Xiong and Collins, 2012), enhances neuroregeneration (Hoffman, 2010) by overcoming CNS inhibitory growth cues (Chong*, et al.*, 1996; Oudega*, et al.*, 1994), and even dramatically reduces neurodegenerative markers in disease models (Franz*, et al.*, 2009). Therefore, lesion conditioning drives and enables regeneration that could treat CNS afflictions.

Our laboratory utilizes the roundworm *Caenorhabditis elegans* for its strongly conserved genetics with mammals, facile genetics, optical transparency, and rapid development. In a prior study, we established a lesion conditioning model in the ASJ sensory neuron (Chung*, et al.*, 2016). We discovered that mutation or pharmacological inhibition of genes in sensory transduction pathways can trigger conditioned regeneration, similar to the effect of a conditioning lesion. As shown in Tab. 1, these interventions

produce several observable phenotypes, including conditioned regeneration and a form

of ectopic axon outgrowth that has the same underlying genetic pathway.

In the prior study, we cell-specifically labelled the ASJ using a transgenic green

fluorescent protein (GFP) reporter driven by a promoter of *trx-1*, an ortholog of human

thioredoxin TXN. We noticed by eye that interventions that condition the ASJ also brighten

the neuron, indicating an upregulation of the gene driving our fluorescent label. Our visual

observations are consistent with prior studies. Following peripheral nerve axotomy, TXN

accumulates in the nucleus and dramatically increases throughout the cytoplasm

(Mansur*, et al.*, 1998), indicating that it is a regeneration-associated gene (RAG) (Ma and

Willis, 2015).

In this study, we quantify ASJ fluorescence and utilize it as an indicator of regenerative

capacity. We quantitatively show that mutations that condition the ASJ by disrupting

sensory signaling significantly brighten the neuron. We confirm this upregulation of *trx-1*

by labeling mRNA transcripts with fluorescence *in situ* hybridization (FISH). Similarly,

conditioning by lesion via laser surgery results in significantly increased fluorescence.

The intensity of fluorescence following these interventions roughly corresponds with the

neuron's regenerative capacity. Additionally, we show that the *trx-1* gene restrains non-

conditioned regeneration, under both redox-dependent and presumably redox-

independent functions and enhances conditioned regeneration via redox-dependent

functions. We leverage our fluorescent proxy to screen for mutations that underlie

conditioning in the ASJ neuron. We isolate twelve mutant lines, spanning six putative

mutations, with significantly reduced ASJ fluorescence. Six of these lines also display

reduced ectopic outgrowth, indicating a disruption in the conditioning pathway. Our proxy

enables efficient isolation of genes underlying lesion conditioning by enabling a visually

based screen that does not require time-consuming surgery.

Materials and methods

C. elegans cultivation, strains, and mutagenesis: We followed established procedure for

cultivation on agarose plates with OP50 bacteria at 20°C (Brenner, 1974). For fluorescent

bead measurements, we cultivated animals on Bacto agarose plates at 20°C for multiple

generations. We cultivated animals on Difco agarose plates at 20°C and 25°C for laser

surgeries and ectopic outgrowth experiments, respectively. We confirmed the genotypes

for all mutant strains by polymerase chain reaction followed by Sanger sequencing

through GENEWIZ (for single-nucleotide polymorphisms) or gel electrophoresis (for large

deletions). Tab. 2 lists the genetics of the strains utilized. We followed established

protocol for ethylmethanesulfonate (EMS) mutagenesis (Brenner, 1974).

FISH labeling and imaging: We used the designer tool from Stellaris (LGC Biosearch

Technologies) to create probes (Table S1) that detect *trx-1* transcripts. We designed the

probes under the following settings: a masking level of 5 (to avoid cross-hybridization to

commonly expressed RNAs at high levels) and at least 2 base pair spacing between

single probes. We generated 16 probes (CAL Fluor® Red 590 conjugated) for mature *trx-*

1, targeting all exons and both UTRs to allow maximal probe number. Of note, we were

unable to reach the optimal amount of probes recommended by the manufacturer (>24)

due to the short mature length of *trx-1*. We synchronized animals by bleaching, hatched

overnight in M9 buffer, and grown until young adult stage in standard conditions on NGM

plates. We used fixation and hybridization protocol as described (Bolkova and Lanctot,

2016). We mounted animals mounted with ProLong™ Glass Antifade Mountant with

NucBlue™ Stain (P36983, Invitrogen™). We acquired images using an Evident Scientific

(formerly Olympus) FLUOVIEW FV3000 laser scanning confocal microscope, equipped

with a high-sensitivity spectral detector unit (with Peltier-cooled GaAsP photomultiplier

modules) and X Line UPLXAPO60XO oil objective. For each animal, we acquired 17 z-

planes with a step size of 0.4 μm.

<u>Laser surgery, imaging of axon regeneration and ectopic outgrowth, and postprocessing:</u>

We followed established procedures for animal immobilization by sodium azide (Chung*,*

et al., 2006) and femtosecond laser surgery (Harreguy*, et al.*, 2020; Wang*, et al.*, 2022).

We severed only the axon (within 2 μm from the cell body), only the dendrite (2/3rd the

length distal from the cell body), or both the axon and dendrite (axon+dendrite), and we

reimaged 48 hours afterwards. We followed established procedures for immobilizing

animals on agarose-azide pads, imaging axon regeneration and outgrowth, and

postprocessing images to measure regenerated length (Chung*, et al.*, 2016). Note that

neurons in certain genetic backgrounds may appear dim in some images due to the

uniform normalization needed for comparison between backgrounds. These neurons are

sufficiently bright to unambiguously measure the regenerated length, particularly after

laser surgery, which upregulates *trx-1*-driven GFP.

Imaging with fluorescent beads: Fluorescence measurements are affected by numerous

factors, including the intensity of the light source and transmission through microscope

optics. These factors may change over time as the components age and their alignment

degrades. To account for these effects, we calibrated fluorescence measurements by

imaging animals alongside fluorescent beads and normalizing measurements across the

images taken. We thoroughly vortexed 0.3% and 3% 2.5 μm stock bead solution

(Invitrogen I7219) and mixed it with nematode growth medium (NGM) buffer (Chung, *et*

al., 2016) in a 5:1 NGM to bead volumetric ratio. We dropped 5.0 μL of the 5:1 solution to

the center of the agarose-azide pad, and then picked adults to the bead solution droplet.

To approximately match neuron to bead fluorescence, we used 3% beads for *tax-2* and

tax-4 strains and 0.3% beads for other strains and surgical conditions. We imaged worms

through a Nikon 0.30 NA, 10x microscope objective and an Andor Zyla sCMOS Plus

camera. Z-stack images were captured with 1-µm spacing.

Postprocessing of fluorescence measurements: We create maximum intensity

projections of our raw z-stack images in ImageJ and identify the brightest pixel within the

ASJ cell body and dendrite for each animal. The pixel values are the raw fluorescent

intensities. Additionally, we average the brightest pixel value within five representative

beads in each raw image. We normalize and scale all raw neuron intensities to our

reference (*tax-2*) intensities according to equation (1):

$(1)\ F_{normalized} = F_{raw} * \dfrac{\bar{F}_{ref\ bead}}{\bar{F}_{raw\ bead}} * \dfrac{F_{raw\ bead\ type}}{F_{ref\ bead\ type}}$, where,

F_{raw} is raw fluorescent intensity of ASJ cell body or dendrite.

$\bar{F}_{raw\,bead}$ is the average of the fluorescence intensity of five representative beads in

the raw image.

$\bar{F}_{ref\,bead}$ is average of the fluorescence intensity of all representative beads in the

set of reference images.

$\frac{F_{raw\,bead\,type}}{F_{ref\,bead\,type}}$ is the true ratio of bead fluorescent intensities according to

manufacturer data. This ratio is 0.08 if the raw image is calibrated with 0.3% beads

or 1.00 if calibrated with 3% beads like the reference image.

The first ratio of Eq. (1) normalizes for day-to-day variability in imaging parameters

between an image and the reference image set. If beads appear unusually bright in an

image due to stronger illumination or increased transmission, then this ratio will

compensate. If the beads are abnormally dim, the opposite will occur. The second ratio

compares the expected brightness of the bead type used in the image against the

reference bead type. Under ideal imaging conditions, this second ratio comparing the

bead intensities from manufacturer data should be the inverse of the first ratio that

compares the measured bead intensities. Therefore, the product of the two ratios should

center around 1.0 if the bead intensities match their manufacturer specifications.

To quantify FISH signal, we adapted an established procedure (Iourov, *et al.*, 2005).

We measured the intensity of the hybridization fluorescence in ImageJ by creating

maximum intensity projections of our z-stacks and acquiring the brightest pixel value for

each ASJ cell body. Since *tax-2* strongly upregulates *trx-1*, we could not quantify the

signal at a single molecule (smFISH) level.

<u>Statistics and interpretation of results:</u> For axon length and fluorescence measurements,

we calculated *P* values by the unpaired, unequal variance, two-tailed *t* test. The

conditioning effect is the difference between regenerated length after axon cut and after

axon+dendrite cut. Its standard deviation is the square root of the sum of the squares of

the regenerated length standard deviations. To compare conditioning effects in different

backgrounds, we calculated *P* values by the unpaired, unequal variance, two-tailed *t* test.

To compare frequencies of ectopic outgrowth, we used Fisher's exact test.

Results

Defects in sensory transduction upregulate *trx-1* expression in the ASJ neuron.

The ASJ is an amphid sensory neuron located in the nose of *C. elegans* (Fig. 1a, left).

The amphids are bilateral, bipolar neurons following a stereotyped morphology consisting

of a cell body, a sensory dendrite that extends to the nose, and an axon that mediates

synaptic connections in the nerve ring (White*, et al.*, 1986). The ASJ axon regenerates

when it is severed. Mutation of dual-leucine kinase, *dlk-1*, completely abolishes what we

call "conventional" (*i.e.*, single-axotomy) axon regeneration (Chung*, et al.*, 2016). The

regeneration of other *C. elegans* (Ghosh-Roy*, et al.*, 2010; Hammarlund*, et al.*, 2009) and

mammalian neurons (Itoh*, et al.*, 2009; Shin*, et al.*, 2012) also show a strong dependence

on *dlk-1*. We trigger DLK-independent regeneration by concomitantly severing the ASJ

axon and dendrite in our lesion conditioning model. This DLK-independent regeneration

exhibits several hallmarks of lesion-conditioned regeneration, including mediation by

cyclic adenosine monophosphate (cAMP) (Neumann*, et al.*, 2002; Qiu*, et al.*, 2002) and

inhibition by L-type voltage gated calcium channels (Enes*, et al.*, 2010). We also

uncovered several sensory mutations that significantly reduce ASJ neuronal activity

levels and thus condition the ASJ to regenerate in *dlk-1* mutants without a physical lesion

(Chung*, et al.*, 2016).

We characterized the effect of a subset of these mutations on the expression of *ptrx-*

1::trx-1::gfp translational fusion reporters *ofIs1* and *ofIs4*. The loci of crucial genes *egl-19*

and *unc-43* are on chromosome IV close to the integration site of *ofIs1*, which

necessitated the use of *ofIs4*. While our previous study utilized an extrachromosomal

array reporter, *ofIs1* and *ofIs4* are integrated into the genome, allowing more accurate

comparison of expression levels across animals. We generated fluorescent animals with

the mutations by crossing in *ofIs1* and *ofIs4* (Fig. 1a) and measured brightness of the ASJ

cell bodies and dendrites to roughly assess TRX-1 levels and localization at a subcellular

level (Fig. 1b). Scattered and out-of-focus light from the larger, brighter cell body often

hides fine or dim structures nearby. Thus, we were unable to consistently measure the

intensity of the axon. In wild-type (wt) animals, the ASJ cell body and dendrite are

relatively dim. In a *dlk-1* mutant background, the ASJ fluorescence remains dim with a

slight decrease in cell body fluorescence (*p < 0.001*), consistent with a non-conditioned

regenerative state. The remainder of the mutations significantly increase the brightness

of the ASJ fibers and especially cell bodies (*p < 0.001* for all mutations). The most

dramatic alteration in fluorescent intensity occurs under *tax-2* and *tax-4* (Fig. 1b, *p <*

0.001)*,* which also produce the strongest conditioned regeneration of the sensory

mutations we studied. The fluorescence in the dendrites of each conditioned strain

significantly increase except for *unc-36*, which may be consistent with the weaker

conditioning effect of *unc-36* (Chung*, et al.*, 2016). We compare *trx-1* expression via

fluorescence with regenerative capacity in several conditioned backgrounds (Fig. S1). For

ASJ cell bodies and dendrites, there is a general positive correlation between

regeneration and *trx-1* expression. However, the correlation is not strong enough to

reliably predict each mutation's impact on regeneration. Thus, the expression of *ptrx-*

1::trx-1::gfp in the ASJ generally corresponds with regenerative potential but is not

predictive at a single gene level.

We confirmed the upregulation of *trx-1* by targeting *Trx-1* mRNA with FISH probes. In

C. elegans, *trx-1* is significantly expressed in only the ASJ neuron and the intestines. We

labeled transcripts in wild-type and *tax-2*, our strongest conditioning mutation, animals.

As shown in Fig. 1c, to verify our FISH probes accurately targeted *trx-1*, we imaged the

ASJ GFP in green (top row) and FISH in red (middle row). The overlays demonstrate

clear colocalization of signals in 3D, indicating the probes accurately target *trx-1* (bottom

row). We quantified fluorescence of the FISH hybridization following the procedure used

for GFP images in Fig. 1a. We measured an 8x enhancement in ASJ fluorescence in *tax-*

2 animals compared to wild-type (Fig. 1d). This increase is similar to what we see in *ofIs1*

and *ofIs4* fluorescence, further confirming the upregulation of *trx-1* in response to

conditioning.

Dendrite cuts upregulate *trx-1* expression and stimulate *dlk*-independent

regeneration in the ASJ neuron.

We also examined changes in ASJ fluorescence in *dlk-1* mutant two days after cutting

the axon, the dendrite, or the axon and dendrite concomitantly (denoted as

"axon+dendrite" or "a+d"). Figure 2a shows images of the ASJ in *dlk-1* animals 48 hours

after surgery as well as in a chronically conditioned background, *tax-2*, to represent a

strongly conditioned fluorescence level. Figure 2b shows the measured cell body and

dendrite fluorescent intensities after each surgery. Our imaging procedure without surgery

(mock condition) does not significantly alter ASJ fluorescence. Surgical interventions on

the axon, dendrite, or axon+dendrite significantly increase fluorescence. Notably,

severing the dendrite or axon+dendrite significantly increases ASJ fluorescence ($p <$

0.001) of nearly the entire population to a brightness that begins to approach the level in

tax-2 animals (right side, Fig. 2b). The fluorescence after axon+dendrite surgery is not

significantly different from after dendrite surgery ($p = 0.163$), suggesting that cutting the

dendrite is sufficient for activating the conditioning response. Figure 2c scatter plots the

dendrite fluorescence against the cell body fluorescence post-surgery, clustering neurons

by fluorescent intensity for comparing to conditioned state. Data points tightly clustered

on the left side of the plot (dimmer ASJ fluorescence, mock and axon cut) represent a

non-conditioned state. Interestingly, changes in fluorescence after axon cut exhibit

bimodal distribution where half remain as dim as the ASJ in mock animals and the other

half noticeably brighten. The cell body and the dendrite fluorescent intensities are

correlated, leading to a clear demarcation between non-conditioned and conditioned

states. Figure 2c confirms that neurons are strongly brightened following dendrite and

axon+dendrite surgeries. The heightened levels of fluorescent intensity following dendrite

or axon+dendrite cut are consistent with these interventions' abilities to condition the

neuron to regenerate. These results are consistent with published findings that TXN

upregulates following peripheral axon injury, the hallmark of a RAG (Bai*, et al.*, 2003;

Mansur*, et al.*, 1998).

Postsurgery regenerated length in ASJ expressing the translational reporter *ofIs1*

match regeneration under a different translational reporter (Chung*, et al.*, 2016).

Conventional regeneration in the ASJ is largely *dlk*-dependent. Figure 2d displays length

regenerated by the ASJ in wild-type and *dlk-1* following axotomy or axon+dendrite cut.

The ASJ regenerates 43 μm on average following axotomy in wild-type but does not

regenerate in *dlk-1* (*p < 0.001*). The conditioning effect is fundamentally the increase in

regeneration arising from a dendrite cut, indicated by the arrows in the figure. Under our

model, axon+dendrite cut triggers conventional and conditioned regeneration, which both

contribute to the total length. In both wild-type and *dlk-1* background, the addition of the

dendrite significantly increases regenerated length (*p < 0.001*) indicating a significant

conditioning effect. Because *dlk-1* eliminates conventional ASJ regeneration, it exposes

regeneration due to conditioning. Thus, we associate regeneration in *dlk-1* with

conditioned regeneration. Note also that while axotomy alone does not produce

conditioned regeneration in *dlk-1*, it significantly increases fluorescence of some neurons.

This divergence indicates that the fluorescent reporter does not fully indicate the

conditioned regenerative state and that *trx-1* expression alone is not sufficient for

activating conditioned regeneration. We speculate that the significant increase in

fluorescence after axotomy alone could be an upregulation of *trx-1* in response to

neuroinjury without activating conditioning mechanisms.

Opposite effects of *trx-1* on two forms of ASJ regeneration

Given that TXN is a RAG (upregulated following neuron injury), we wanted to determine

if *trx-1* is involved in either form of regeneration. Our prior work utilized a translational

—13—

ptrx-1::trx-1::gfp fusion marker (*ofIs1 or ofIs4*), potentially resulting in an overexpression

of functional TRX-1 protein in the ASJ. Therefore, we visualized the ASJ with a

transcriptional marker *ofIs5[ptrx-1::gfp]* and probed the effects of *trx-1(ok1449)*, which is

a loss-of-function allele deleting a 860 bp region spanning part of the proximal promoter

region up to the last exon (Miranda-Vizuete*, et al.*, 2006). As shown in Fig. 3, the trends

in regeneration in wild-type and *dlk-1* are consistent with regeneration under translational

fusion reporters in Fig. 2 and in our prior study (Chung*, et al.*, 2016). In wild-type animals,

the axon regenerates at a basal rate when it is severed, and mutation of *dlk-1* eliminates

conventional regeneration. The addition of a dendrite cut increases regeneration in wild-

type (*p < 0.05*) and restores regeneration in *dlk-1* due to the conditioning effect (*p <*

0.001).

ASJ conventional regeneration in *trx-1(ok1449)* is significantly stronger than in wild-

type (*p < 0.001*), suggesting that *trx-1* restrains conventional regeneration. Introducing

the *dlk-1* mutation to *trx-1* mutant background nearly eliminates conventional

regeneration (Fig. 3), similar to results in translational fusions and *trx-1(+)*. The

conditioning effect in *dlk-1; trx-1(ok1449)* is significantly less than the conditioning effect

in *dlk-1*, as indicated by the arrows in Fig. 3 (*p < 0.05*). These data suggest that *trx-1*

enhances conditioned regeneration.

Redox functions of *trx-1* modulate both pathways of ASJ regeneration

To further uncover the role of *trx-1* in governing regeneration in the ASJ, we examined its

redox-dependent functions. Two cysteine residues at conserved active sites on

thioredoxin enable reversible oxidation (Lee*, et al.*, 2013). Substituting one or both

cysteine residues with serines, a redox-inactive amino acid, abolishes thioredoxin's redox

activity. In *C. elegans*, *trx-1(syb206)* mutants with both residues substituted (wild-type

CGPC → redox-inactive SGPS) were utilized to demonstrate that *trx-1* regulates

development through separate redox-dependent and -independent pathways (Sanzo-

Machuca*, et al.*, 2019). We assessed regeneration in the redox-inactive mutant to isolate

thioredoxin's redox-dependent effects (Fig. 3). Following axon cut only, regenerated

length in *trx-1(syb206)* is significantly longer than in wildtype (*p < 0.05*) but significantly

shorter than in the full *trx-1(ok1449)* null (*p < 0.05*). These results suggest that

conventional regeneration is inhibited by *trx-1* under both redox-dependent and putatively

redox-independent functions.   In contrast, there is no significant change in the

conditioning effect (indicated by arrows) between *dlk-1; trx(ok1449)* and *dlk-1; trx-*

1(syb206) ($p = 0.141$). This result suggest that conditioned regeneration is likely mediated

solely by redox-dependent functions of *trx-1*.

Screening of *trx-1* reporter isolates mutants with reduced regenerative potential.

One powerful technique for exposing genes underlying a phenotype is the unbiased

forward screen (Brenner, 1974). In brief, a screen stochastically introduces mutations into

the genomes of many animals. The correlation between *trx-1* expression and

regenerative capacity suggests that we can rapidly screen for genes involved in

conditioned regeneration utilizing a *trx-1* fluorescent reporter. We exposed our chronically

conditioned *tax-2* strain, labeled with the translational reporter *ofIs1*, to ethyl

methanesulfonate (EMS). Starting in the F_2 generation, we isolated mutant lines by

visually screening for decreased fluorescence in the ASJ neuron by eye (Fig. 4a) which

potentially indicates a defect in a conditioning-related gene. Often, it was difficult to detect

a significant reduction in cell body fluorescence, so we screened for significantly dimmer

dendrites that we could not observe at a set magnification. We generated twelve mutant

lines, spanning six putative mutations, with significantly reduced ASJ fluorescence in the

cell body (Fig. 4b) or dendrite (Fig. 4c). In a typical screen, there is a frequency of roughly

one null mutation in a given gene for every 2000 F_1s screened (Jorgensen and Mango,

2002). Given that we screened 100 F_1s, it is highly unlikely (though possible) that our

screen produced the same mutations or mutations in the same gene. Thus, by the

statistics, we strongly expect the six F_1 lines isolated to carry distinct mutations in distinct

genes. Ten of these lines display significantly reduced fluorescence in both the dendrite

and cell body (*p < 0.001*).

Mutations to genes involved in the sensory pathways (*i.e.*, conditioning mutations)

decrease ASJ neuronal activity and trigger regeneration. These sensory mutations,

including the mutations in Fig. 1, also alter axon morphology in *C. elegans* sensory

neurons and produce ectopic outgrowths (Coburn and Bargmann, 1996; Peckol*, et al.*,

1999). We found that ectopic outgrowth extensively shares genetic and molecular

mechanisms of axon growth with conditioned regeneration (Chung*, et al.*, 2016). Thus,

ectopic outgrowth indicates conditioned regenerative potential. As shown in Fig. 4d, six

of the strains we isolated exhibit significantly reduced frequencies of ectopic outgrowth

compared to the original pre-mutagenized strain (*p < 0.05* or *0.001*), suggesting that these

strains possess a mutation that restrains conditioned regeneration in the ASJ. As we

previously found, fluorescence is not always an accurate predictor of conditioned

outgrowth. We identified some strains with significantly reduced cell body and dendrite

fluorescence but unchanged ectopic outgrowth frequency. We expect these strains to

carry a mutation that decreases *trx-1* expression without impact to the conditioning

pathway. In summary, our results demonstrate the utility and limitations of our

fluorescence proxy as an indicator of ASJ intrinsic regenerative capacity.

Discussion

The change in expression of TXN following neuronal injury has been widely characterized

in mammalian models. TXN induction occurs after a variety of insults, including ischemia

(Tomimoto*, et al.*, 1993) and oxidative stress (Sugino*, et al.*, 1999). Importantly, motor

and sensory nerve axotomy in rats leads to significantly increased levels of TXN (Mansur*,*

et al., 1998; Stemme*, et al.*, 1985). Likewise, we note that laser surgery of the ASJ

neurites upregulates *trx-1* expression, as indicated by *trx-1::gfp* reporter. Our study offers

three lines of evidence for associating this enhanced *trx-1* expression with the conditioned

state. First, changes due to conditioning, including upregulation of RAGs, preferentially

occurs after lesioning the sensory neurite rather than the central, or synaptic, neurite.

Severing the sensory dendrite strongly upregulates *trx-1* in nearly all ASJ neurons while

severing the axon upregulates expression in only 50% of neurons to a lesser degree (Fig.

2b). Second, each mutation we tested that improves regeneration (Chung*, et al.*, 2016)

by conditioning the ASJ neuron also upregulates expression of *trx-1* as measured by GFP

(Fig. 1b) and FISH (Figs. 1cd). Third, *trx-1* expression generally corresponds with

conditioned regenerative potential (Fig. 1b). Out of the mutations tested, the *tax-2* and

tax-4 mutations brighten the ASJ the most and produce the highest levels of ectopic

outgrowth and regeneration (Chung*, et al.*, 2016). However, this relationship is not

comprehensive (Fig. S1). The *tax-2* and *tax-4* mutations produce marginally better

conditioned regeneration compared to other mutations, and the *unc-36* mutation

increases *trx-1* expression but minimally triggers regeneration (Chung*, et al.*, 2016). This

differential alteration of *trx-1* expression and regeneration suggests a divergence of

pathways underlying *trx-1* expression and conditioning.

To our knowledge, the role of *trx-1* in neuronal regeneration has not directly been

studied, although prior evidence suggests its role as a mediator. Overexpressing TXN in

mice can promote neuroprotective effects following stroke (Takagi*, et al.*, 1999), and TXN

is required for nerve growth factor enhancement of nerve outgrowth in a tumor cell line

(Bai*, et al.*, 2003). We directly test the role of *trx-1* in modulating conventional and

conditioned regeneration in the ASJ (Fig. 3). We show that *trx-1* inhibits conventional

regeneration, consistent with findings in the PLM sensory neuron in *C. elegans*. Animals

defective in thioredoxin reductase *trxr-1*, which enables *trx-1* activity (Stenvall*, et al.*,

2011), display increased PLM regeneration (Kim*, et al.*, 2018). We also show that *trx-1*

mediates *dlk*-independent conditioned regeneration, consistent with the upregulation of

trx-1 expression following conditioning interventions. Lesion conditioning is enabled by

changes in transcription that lead to an enhancement of regeneration (Hoffman, 2010).

The mammalian TXN gene contains a regulatory region with cAMP responsive element

(CRE) through which nerve growth factor acts to induce nerve outgrowth (Bai*, et al.*,

2003). The *trx-1* gene has a similar consensus CRE sequence, CGTCA (Montminy*, et*

al., 1986; Park*, et al.*, 2021). We speculate CRE-binding protein (CREB), an integral

transcription factor for many biological processes, could regulate *trx-1* transcription

through this region to enact the conditioning response. The cAMP pathway is one of the

best-described mechanisms for enhancing conditioned regeneration (Neumann*, et al.*,

2002; Qiu*, et al.*, 2002), and in *C. elegans*, cAMP signaling triggers conditioned

regeneration (Chung*, et al.*, 2016). Likewise, we recently found that CREB enhances

conditioned regeneration (Wang*, et al.*, 2023). Thus, cAMP and CREB may contribute to

conditioned regeneration in part via *trx-1* activation through its CRE region.

The mechanisms by which *trx-1* differentially affects conventional and conditioned

regeneration requires additional study. Dendrite injury preferentially upregulates *trx-1*,

which would presumably enhance conditioned regeneration over conventional

regeneration. This matches the perturbation and phenotype of a conditioning lesion. The

conventional and conditioned regeneration pathways are fundamentally neurite growth

mechanisms, and they likely converge or regulate each other to produce this growth.

Their interaction may favor one pathway over the other depending on context, such as

neuron development (regeneration recapitulates some aspects of development),

regeneration, or even cell death and stress response. Both *trx-1* and *dlk-1* are known to

interact with JNK/MAPK signaling cascades that initiate cell death in response to stress

(Fujino*, et al.*, 2007; Nix*, et al.*, 2011). The specific interaction of *dlk-1* and *trx-1* to drive

their biological functions remains undefined. However, our results in Fig. 3 suggest that

the inhibition of conventional regeneration by *trx-1* acts upstream or converges to *dlk-1*

given that loss of *trx-1* only significantly increases conventional regeneration with

functional DLK.

Redox activities of thioredoxin in *C. elegans* are important for modulating biological

processes (Sanzo-Machuca*, et al.*, 2019). In mammalian models, the redox-related

pathways of TXN have been heavily studied, particularly in regulating apoptosis and

protecting against neurodegenerative disorders. It is not surprising that *trx-1* plays an

integral biological role in these processes since a hallmark of cell death and

neurodegeneration is an accumulation of damaged proteins, reactive oxygen species,

and oxidative stress (Lin and Beal, 2006; Mattson and Magnus, 2006). Thioredoxin

inhibits apoptosis signal-regulating kinase 1 (ASK1) to mediate endoplasmic reticulum

stress-induced apoptosis (Nishitoh*, et al.*, 2002; Saitoh*, et al.*, 1998) and can delay retinal

neurodegeneration in the Tubby mouse model (Kong*, et al.*, 2007; Kong*, et al.*, 2010). It

is still unknown how these redox functions dictate regenerative potential, but stress

response and cell death pathways are important modulators of regeneration (Pinan-

Lucarre*, et al.*, 2012). We speculate that *trx-1* may impinge on or regulate these pathways

to affect axon regeneration.

Using our fluorescence proxy, we isolated twelve mutant lines with significantly

decreased ASJ fluorescence. Six of those lines display significantly reduced ectopic

outgrowth, which is related to conditioned regeneration, indicating that they contain a

mutation in the conditioning pathways. Additionally, we developed a fluorescent bead

normalization protocol to facilitate quantitative comparison of fluorescence in genetic

backgrounds or under different interventions. These tools enable us to pursue a rapid,

yet quantitative approach to the question of neuronal regeneration. Our approach is

complimentary to established techniques for identifying RAGs, such as microarray

analysis (Costigan*, et al.*, 2002). In contrast to these RAGs, genes identified by our

approach are highly likely to functionally underlie regeneration, by virtue of their

involvement in ectopic outgrowth.

Our proxy demonstrates significant potential as we isolated several strains with

reduced outgrowth potential by selecting for reduced ASJ fluorescence. However, we did

identify equally as many dim strains without reduced outgrowth potential (Fig. 4d). These

findings are not unexpected since we introduce stochastic mutations throughout the

genome and screen with a fluorescent indicator that only roughly corresponds with

regenerative capacity. False positives can be a common drawback to forward genetic

screens under mutagenesis due to their stochastic nature. In these isolated lines, we

expect that EMS disrupted pathways directly related to *trx-1* expression, genes that

interact upstream of *trx-1*, or the transgenic label itself, thus causing a significant

reduction in fluorescence. These effects can be mitigated by pursuing a reverse genetic

screen under RNAi to test genes that have an expected role in conditioned regeneration.

A similar screen was successfully carried out on transcription factors regulating non-

conditioned *trx-1* expression (Gonzalez-Barrios*, et al.*, 2015).

Additionally, we screen for decreased fluorescence by eye under the current workflow.

This reduction in fluorescence is often difficult to distinguish by eye. Also, the range of

fluorescence of the starting strain and mutant often overlap, which prevented outcrossing

and identification of the causative mutations. We could improve the throughput and

probability of success by implementing real-time quantitative measurements to more

precisely screen for changes in fluorescence. For instance, we could utilize microfluidic

devices to rapidly screen through animals for a reduction in fluorescence or implement

whole plate imaging (Wang*, et al.*, 2023) to quickly image entire populations of animals

to select for decreased fluorescence.

Outlook

Studying regeneration *in vivo* remains difficult for experimental reasons and due to

interacting pathways yet to be explored and fully defined. We have developed a single-

cell model to study the conventional and conditioned regeneration pathways in the *C.*

elegans ASJ neuron and a proxy to identify genes and potential therapeutic targets

involved. Our study could be extended in multiple directions. First, the mechanisms

through which *trx-1* (and its redox activities) modulates regeneration is of particular

interest because *trx-1* specifically enhances conditioned, but inhibits conventional,

regeneration. Second, the mechanisms that underlie *trx-1* expression and their relation

to conditioning pathways should be clarified. Third, while *trx-1* expression is specific to

the ASJ, our approach could be extended to other neurons by labeling with fluorescent

reporters driven by their RAGs. Neuron types and even subtypes have distinct

regeneration capacities (Duan*, et al.*, 2015), but broad regeneration studies in multiple

neurons are lacking. Examining regeneration in other neurons may illuminate new

regeneration pathways. Our work thus represents an important first step in identifying

undiscovered modulators of neuron regeneration and in establishing new approaches for

rapidly studying regeneration.

Acknowledgements

We thank members of the Chung laboratory for taking part in the revision process. We

thank Leilani Schulting for assistance with data postprocessing. We acknowledge Peter

Swoboda (Karolinska Institute, Huddinge, Sweden) for providing OE3417 and OE3265.

We would like to also thank Antonio Miranda-Vizuete (Biomedicine Institute of Sevilla,

Seville, Spain) for providing VZ763, VZ748, and invaluable feedback on the manuscript.

References

Bai, J., Nakamura, H., Kwon, Y.W., Hattori, I., Yamaguchi, Y., Kim, Y.C., Kondo, N., Oka, S., Ueda, S., Masutani, H.*, et al.* (2003). Critical roles of thioredoxin in nerve growth factor-mediated signal transduction and neurite outgrowth in PC12 cells. J Neurosci *23*, 503-509.

Bolkova, J., and Lanctot, C. (2016). Quantitative gene expression analysis in Caenorhabditis elegans using single molecule RNA FISH. Methods *98*, 42-49.

Brenner, S. (1974). The genetics of Caenorhabditis elegans. Genetics *77*, 71-94.

Chong, M.S., Woolf, C.J., Turmaine, M., Emson, P.C., and Anderson, P.N. (1996). Intrinsic versus extrinsic factors in determining the regeneration of the central processes of rat dorsal root ganglion neurons: The influence of a peripheral nerve graft. J Comp Neurol *370*, 97-104.

Chung, S.H., Awal, M.R., Shay, J., McLoed, M.M., Mazur, E., and Gabel, C.V. (2016). Novel DLK-independent neuronal regeneration in Caenorhabditis elegans shares links with activity-dependent ectopic outgrowth. Proc Natl Acad Sci USA *113*, E2852-E2860.

Chung, S.H., Clark, D.A., Gabel, C.V., Mazur, E., and Samuel, A.D. (2006). The role of the AFD neuron in C. elegans thermotaxis analyzed using femtosecond laser ablation. BMC Neurosci *7*, 30.

Coburn, C.M., and Bargmann, C.I. (1996). A putative cyclic nucleotide-gated channel is required for sensory development and function in C. elegans. Neuron *17*, 695-706.

Costigan, M., Befort, K., Karchewski, L., Griffin, R.S., D'Urso, D., Allchorne, A., Sitarski, J., Mannion, J.W., Pratt, R.E., and Woolf, C.J. (2002). Replicate high-density rat genome oligonucleotide microarrays reveal hundreds of regulated genes in the dorsal root ganglion after peripheral nerve injury. BMC Neuroscience *3*.

Duan, X., Qiao, M., Bei, F., Kim, I.-J., He, Z., and Sanes, Joshua R. (2015). Subtype-Specific Regeneration of Retinal Ganglion Cells following Axotomy: Effects of Osteopontin and mTOR Signaling. Neuron *85*, 1244-1256.

Dubový, P., Klusáková, I., Hradilová-Svíženská, I., Brázda, V., Kohoutková, M., and Joukal, M. (2019). A Conditioning Sciatic Nerve Lesion Triggers a Pro-regenerative State in Primary Sensory Neurons Also of Dorsal Root Ganglia Non-associated With the Damaged Nerve. Front Cell Neurosci *13*.

Enes, J., Langwieser, N., Ruschel, J., Carballosa-Gonzalez, M.M., Klug, A., Traut, M.H., Ylera, B., Tahirovic, S., Hofmann, F., Stein, V.*, et al.* (2010). Electrical Activity Suppresses Axon Growth through Ca(v)1.2 Channels in Adult Primary Sensory Neurons. Curr Biol *20*, 1154-1164.

Feigin, V.L., Vos, T., Alahdab, F., Amit, A.M.L., Bärnighausen, T.W., Beghi, E., Beheshti, M., Chavan, P.P., Criqui, M.H., Desai, R.*, et al.* (2021). Burden of Neurological Disorders Across the US From 1990-2017. JAMA Neurology *78*, 165.

Franz, C.K., Quach, E.T., Krudy, C.A., Federici, T., Kliem, M.A., Snyder, B.R., Raore, B., and Boulis, N.M. (2009). A Conditioning Lesion Provides Selective Protection in a Rat Model of Amyotrophic Lateral Sclerosis. PLOS ONE *4*, 8.

Fujino, G., Noguchi, T., Matsuzawa, A., Yamauchi, S., Saitoh, M., Takeda, K., and Ichijo, H. (2007). Thioredoxin and TRAF family proteins regulate reactive oxygen species-dependent activation of ASK1 through reciprocal modulation of the N-terminal homophilic interaction of ASK1. Mol Cell Biol *27*, 8152-8163.

Ghosh-Roy, A., Wu, Z.L., Goncharov, A., Jin, Y.S., and Chisholm, A.D. (2010). Calcium
and Cyclic AMP Promote Axonal Regeneration in Caenorhabditis elegans and Require
DLK-1 Kinase. J Neurosci *30*, 3175-3183.
Gonzalez-Barrios, M., Fierro-Gonzalez, J.C., Krpelanova, E., Mora-Lorca, J.A., Pedrajas,
J.R., Penate, X., Chavez, S., Swoboda, P., Jansen, G., and Miranda-Vizuete, A. (2015).
Cis- and Trans-Regulatory Mechanisms of Gene Expression in the ASJ Sensory Neuron
of Caenorhabditis elegans. Genetics *200*, 123-U269.
Hammarlund, M., Nix, P., Hauth, L., Jorgensen, E.M., and Bastiani, M. (2009). Axon
Regeneration Requires a Conserved MAP Kinase Pathway. Science *323*, 802-806.
Harreguy, M.B., Marfil, V., Grooms, N.W.F., Gabel, C.V., Chung, S.H., and Haspel, G.
(2020). Ytterbium-doped fibre femtosecond laser offers robust operation with deep and
precise microsurgery of C. elegans neurons. Sci Rep *10*, 4545.
Hoffman, P.N. (2010). A conditioning lesion induces changes in gene expression and
axonal transport that enhance regeneration by increasing the intrinsic growth state of
axons. Exp Neurol *223*, 11-18.
Iourov, I.Y., Soloviev, I.V., Vorsanova, S.G., Monakhov, V.V., and Yurov, Y.B. (2005). An
approach for quantitative assessment of fluorescence in situ hybridization (FISH) signals
for applied human molecular cytogenetics. J Histochem Cytochem *53*, 401-408.
Itoh, A., Horiuchi, M., Bannerman, P., Pleasure, D., and Itoh, T. (2009). Impaired
regenerative response of primary sensory neurons in ZPK/DLK gene-trap mice.
Biochemical and Biophysical Research Communications *383*, 258-262.
James, S.L., Theadom, A., Ellenbogen, R.G., Bannick, M.S., Montjoy-Venning, W.,
Lucchesi, L.R., Abbasi, N., Abdulkader, R., Abraha, H.N., Adsuar, J.C.*, et al.* (2019).
Global, regional, and national burden of traumatic brain injury and spinal cord injury,
1990–2016: a systematic analysis for the Global Burden of Disease Study 2016. The
Lancet Neurology *18*, 56-87.
Jorgensen, E.M., and Mango, S.E. (2002). The art and design of genetic screens:
caenorhabditis elegans. Nat Rev Genet *3*, 356-369.
Kim, K.W., Tang, N.H., Piggott, C.A., Andrusiak, M.G., Park, S., Zhu, M., Kurup, N.,
Cherra, S.J., Wu, Z., Chisholm, A.D.*, et al.* (2018). Expanded genetic screening in
Caenorhabditis elegans identifies new regulators and an inhibitory role for NAD+ in axon
regeneration. eLife *7*.
Kong, L., Tanito, M., Huang, Z., Li, F., Zhou, X., Zaharia, A., Yodoi, J., McGinnis, J.F.,
and Cao, W. (2007). Delay of photoreceptor degeneration in tubby mouse by
sulforaphane. J Neurochem *101*, 1041-1052.
Kong, L., Zhou, X., Li, F., Yodoi, J., McGinnis, J., and Cao, W. (2010). Neuroprotective
effect of overexpression of thioredoxin on photoreceptor degeneration in Tubby mice.
Neurobiol Dis *38*, 446-455.
Lee, S., Kim, S.M., and Lee, R.T. (2013). Thioredoxin and thioredoxin target proteins:
from molecular mechanisms to functional significance. Antioxid Redox Signal *18*, 1165-
1207.
Lin, M.T., and Beal, M.F. (2006). Mitochondrial dysfunction and oxidative stress in
neurodegenerative diseases. Nature *443*, 787-795.
Ma, T.C., and Willis, D.E. (2015). What makes a RAG regeneration associated? Front
Mol Neurosci *8*, 13.

Mansur, K., Iwahashi, Y., Kiryu-Seo, S., Su, Q., Namikawa, K., Yodoi, J., and Kiyama, H.
(1998). Up-regulation of thioredoxin expression in motor neurons after nerve injury. Mol
Brain Res *62*, 86-91.
Mattson, M.P., and Magnus, T. (2006). Ageing and neuronal vulnerability. Nat Rev
Neurosci *7*, 278-294.
McQuarrie, I.G., and Grafstein, B. (1973). Axon Outgrowth Enhanced by a Previous
Nerve Injury. Arch Neurol *29*, 53-55.
Miranda-Vizuete, A., Gonzalez, J.C., Gahmon, G., Burghoorn, J., Navas, P., and
Swoboda, P. (2006). Lifespan decrease in a Caenorhabditis elegans mutant lacking TRX-
1, a thioredoxin expressed in ASJ sensory neurons. FEBS Lett *580*, 484-490.
Montminy, M.R., Sevarino, K.A., Wagner, J.A., Mandel, G., and Goodman, R.H. (1986).
Identification of a cyclic-AMP-responsive element within the rat somatostatin gene. Proc
Natl Acad Sci U S A *83*, 6682-6686.
Neumann, S., Bradke, F., Tessier-Lavigne, M., and Basbaum, A.I. (2002). Regeneration
of sensory axons within the injured spinal cord induced by intraganglionic cAMP elevation.
Neuron *34*, 885-893.
Nishitoh, H., Matsuzawa, A., Tobiume, K., Saegusa, K., Takeda, K., Inoue, K., Hori, S.,
Kakizuka, A., and Ichijo, H. (2002). ASK1 is essential for endoplasmic reticulum stress-
induced neuronal cell death triggered by expanded polyglutamine repeats. Genes Dev
16, 1345-1355.
Nix, P., Hisamoto, N., Matsumoto, K., and Bastiani, M. (2011). Axon regeneration requires
coordinate activation of p38 and JNK MAPK pathways. Proceedings of the National
Academy of Sciences of the United States of America *108*, 10738-10743.
Oudega, M., Varon, S., and Hagg, T. (1994). Regeneration of Adult Rat Sensory Axons
into Intraspinal Nerve Grafts: Promoting Effects of Conditioning Lesion and Graft
Predegeneration. Exp Neurol *129*, 194-206.
Park, J., Oh, H., Kim, D., Cheon, Y., Park, Y.J., Hwang, H., Neal, S.J., Dar, A.R., Butcher,
R.A., Sengupta, P.*, et al.* (2021). CREB mediates the C. elegans dauer polyphenism
through direct and cell-autonomous regulation of TGF-beta expression. PLoS Genet *17*,
22.
Peckol, E.L., Zallen, J.A., Yarrow, J.C., and Bargmann, C.I. (1999). Sensory activity
affects sensory axon development in C. elegans. Development *126*, 1891-1902.
Pinan-Lucarre, B., Gabel, C.V., Reina, C.P., Hulme, S.E., Shevkoplyas, S.S., Slone, R.D.,
Xue, J., Qiao, Y.J., Weisberg, S., Roodhouse, K.*, et al.* (2012). The Core Apoptotic
Executioner Proteins CED-3 and CED-4 Promote Initiation of Neuronal Regeneration in
Caenorhabditis elegans. PLoS Biol *10*.
Qiu, J., Cai, C.M., Dai, H.N., McAtee, M., Hoffman, P.N., Bregman, B.S., and Filbin, M.T.
(2002). Spinal axon regeneration induced by elevation of cyclic AMP. Neuron *34*, 895-
903.
Richardson, P.M., and Issa, V.M.K. (1984). Peripheral Injury Enhances Central
Regeneration of Primary Sensory Neurons. Nature *309*, 791-793.
Saitoh, M., Nishitoh, H., Fujii, M., Takeda, K., Tobiume, K., Sawada, Y., Kawabata, M.,
Miyazono, K., and Ichijo, H. (1998). Mammalian thioredoxin is a direct inhibitor of
apoptosis signal-regulating kinase (ASK) 1. EMBO J *17*, 2596-2606.
Sanzo-Machuca, A., Monje Moreno, J.M., Casado-Navarro, R., Karakuzu, O., Guerrero-
Gomez, D., Fierro-Gonzalez, J.C., Swoboda, P., Munoz, M.J., Garsin, D.A., Pedrajas,

J.R*., et al.* (2019). Redox-dependent and redox-independent functions of Caenorhabditis
elegans thioredoxin 1. Redox Biol *24*, 101178.
Shin, J.E., Cho, Y.C., Beirowski, B., Milbrandt, J., Cavalli, V., and DiAntonio, A. (2012).
Dual Leucine Zipper Kinase Is Required for Retrograde Injury Signaling and Axonal
Regeneration. Neuron *74*, 1015-1022.
Stemme, S., Hansson, H.A., Holmgren, A., and Rozell, B. (1985). Axoplasmic transport
of thioredoxin and thioredoxin reductase in rat sciatic nerve. Brain Res *359*, 140-146.
Stenvall, J., Fierro-González, J.C., Swoboda, P., Saamarthy, K., Cheng, Q., Cacho-
Valadez, B., Arnér, E.S.J., Persson, O.P., Miranda-Vizuete, A., and Tuck, S. (2011).
Selenoprotein TRXR-1 and GSR-1 are essential for removal of old cuticle during molting
in <i>Caenorhabditis elegans</i>. Proceedings of the National Academy of Sciences *108*,
1064-1069.
Sugino, T., Nozaki, K., Takagi, Y., Hattori, I., Hashimoto, N., and Yodoi, J. (1999).
Expression and distribution of redox regulatory protein, thioredoxin after metabolic
impairment by 3-nitropropionic acid in rat brain. Neurosci Lett *275*, 145-148.
Takagi, Y., Mitsui, A., Nishiyama, A., Nozaki, K., Sono, H., Gon, Y., Hashimoto, N., and
Yodoi, J. (1999). Overexpression of thioredoxin in transgenic mice attenuates focal
ischemic brain damage. Proceedings of the National Academy of Sciences *96*, 4131-
4136.
Tomimoto, H., Akiguchi, I., Wakita, H., Kimura, J., Hori, K., and Yodoi, J. (1993). Astroglial
expression of ATL-derived factor, a human thioredoxin homologue, in the gerbil brain after
transient global ischemia. Brain Res *625*, 1-8.
Wang, Y.L., Grooms, N.W.F., and Chung, S.H. (2022). Transverse and axial resolution
of femtosecond laser ablation. J Biophotonics, e202200042.
Wang, Y.L., Grooms, N.W.F., Jaklitsch, E.L., Schulting, L.G., and Chung, S.H. (2023).
High-throughput submicron-resolution microscopy of Caenorhabditis elegans populations
under strong immobilization by cooling cultivation plates. iScience *26*, 105999.
White, J.G., Southgate, E., Thomson, J.N., and Brenner, S. (1986). The structure of the
nervous system of the nematode Caenorhabditis elegans. Philos Transact B-Biol Sci *314*,
1-340.
Xiong, X., and Collins, C.A. (2012). A conditioning lesion protects axons from
degeneration via the Wallenda/DLK MAP kinase signaling cascade. J Neurosci *32*, 610-
615.
660

interventions →	condition neuron	→ phenotype
surgery conditioning mutations pharmacology		regeneration ectopic outgrowth **fluorescence**

662 **Table 1**: Interventions condition neurons by disrupting sensory signaling. Conditioning generates visible
663 phenotypes. We examine *trx-1::gfp* fluorescence in this study.
664

mutation	translational marker		transcriptional marker
	ofIs1[ptrx-1::trx-1::gfp]	*ofIs4[ptrx-1::trx-1::gfp]*	*ofIs5[ptrx-1::gfp]*
none	BOS2	OE3417	BOS41
dlk-1(ju476)I	BOS71		BOS44
dlk-1(ju476)I; trx-1(ok1449)II			BOS42
dlk-1(ju476)I; trx-1(syb206)II			BOS82
egl-19(n582)IV		BOS70	
sax-1(ky211)X	BOS74		
tax-2(p691)I	BOS72	BOS76	
tax-4(p678)III		BOS73	
trx-1(ok1449)II			VZ763
trx-1(syb206)II			BOS80
unc-36(e251)III	BOS69		
unc-43(n1186n498)IV		BOS75	

Table 2: Strains and genotypes utilized.

Figure captions

Figure 1: Sensory mutations upregulate ASJ neuron expression of *trx-1*. (a)

Leftmost panels: Line drawing and typical ASJ neuron. Remainder of panels:

representative images of *ptrx-1::trx-1::gfp* in wild-type (wt), mutant backgrounds. Sensory

mutations, particularly *tax-2* and *tax-4*, upregulate expression. Identical brightness,

contrast for all images except leftmost. Fluorescent beads indicated by #. (b)

Quantification of fluorescence in part A. Data represented as average ± standard

deviation (SD). $n = 50$ for each data point. ** $p < 0.001$ (c) FISH confirms upregulation of

trx-1 in sensory mutant*.* Representative images of *trx-1*-targeted labels in GFP (top), FISH

(middle), and merged (bottom) in wild-type and *tax-2* animals. (d) Quantification of

fluorescence in part C. Each point represents measurement in one neuron. Data

represented as average $\pm$ SD.

Figure 2: Dendrite cuts upregulate *trx-1* expression and condition ASJ to

regenerate. (a) GFP fluorescence images of ASJ neuron. Fluorescent bead indicated by

#. Identical brightness, contrast for all images. (b) Quantification of fluorescence in part

A. Data represented as average $\pm$ SD. Each dot represents one animal. (c) ASJ cell

body and dendrite fluorescence distributions from part B. Dendrite cut brightens

fluorescence in entire population. (d) Total length of ASJ regeneration following axon or

axon+dendrite (a+d) cut in wt, *dlk-1*. Dendrite cut enhances ASJ regeneration under DLK-

independent mechanism. Arrows represent conditioning effect, or contribution of

conditioning to additional regenerated length from dendrite cut. Data represented as

average $\pm$ standard error of the mean (SEM). $n \geq 20$ for all conditions. ** $p < 0.001$.

Control and *tax-2* images and data replicated from Fig. 1.

Figure 3: *trx-1* modulates ASJ regeneration. Mutation of *dlk-1* and *trx-1* differentially

modulate regeneration after axon or a+d surgery. ok1449 is *trx-1* null allele; syb206 is *trx-*

1 redox-dead allele. Arrows represent conditioning effect. Redox-dependent and -

independent functions of *trx-1* modulate conventional and conditioned regeneration. Data

represented as average $\pm$ SEM. $n \geq 20$ for all conditions. * $p < 0.05$, ** $p < 0.001$.

Figure 4: Fluorescence proxy enables isolation of mutants with reduced

conditioned regenerative potential. (a) ASJ images in control and post-mutagenesis

animals. Fluorescent bead indicated by #. Identical brightness, contrast for all images.

(b) Quantification of cell body fluorescence in part A. Reduced fluorescence in 11 mutant

lines. (c) Quantification of dendrite fluorescence in part A. Reduced fluorescence in 11

mutant lines. (d) Ectopic outgrowth frequency for control and post-mutagenesis strains.

Reduced ectopic outgrowth in 6 lines. Fluorescence measurements: $n = 50$. Ectopic

outgrowth: $60 \leq n \leq 94$ for strains 3b, 6a, 6c; $n \geq 150$ for other strains. Strain 4a not viable

at cultivation temperature. Data represented as average $\pm$ SD. * $p < 0.05$, ** $p < 0.001$.

Control (*tax-2*) images and fluorescence data from Fig. 1.

Figure 1
Grooms, et al

Figure 2
Grooms, et al

Figure 3
Grooms, et al

Figure 4
Grooms, et al