Host genetic background is a barrier to vaccine-induced protection against infection

Rocky Lai[1], Diana Gong[2], Travis Williams[1], Abiola F Ogunsola[1], Kelly Cavallo[1], Cecilia Lindestam Arlehamn[3], Gillian L. Beamer[4], Martin Ferris[5], Christopher M. Sassetti[1], Doug Lauffenburger[2], Samuel M. Behar[1]

1. Department of Microbiology and Physiological Systems, University of Massachusetts Medical School, Worcester, Massachusetts, USA.
2. Department of Biological Engineering, Massachusetts Institute of Technology, Cambridge, MA, USA
3. Center for infectious disease and vaccine research, La Jolla Institute for Immunology., La Jolla, CA, USA
4. Texas Biomedical Research Institute, San Antonio, TX, USA
5. Department of Genetics, University of North Carolina at Chapel Hill, Chapel Hill, United States

*Correspondence:

Samuel M. Behar

E-mail address: samuel.behar@umassmed.edu

Summary (150)

The heterogeneity of immune responses observed in humans is difficult to model in

standard inbred laboratory mice. We used Collaborative Cross (CC) recombinant inbred

strains to capture the diversity of outbred populations and better understand how host

variation affects BCG-induced immunity against *Mycobacterium tuberculosis*. Using 24

CC strains, we show host genetics has a major role in determining whether BCG elicits

protective immunity to *M. tuberculosis*. Only 54% of BCG-vaccinated CC strains had

significantly reduced lung bacillary burdens after infection. Furthermore, BCG efficacy is

dissociable from inherent susceptibility to TB. Like people, CC mice have diverse

immunological responses after BCG vaccination and *M. tuberculosis* challenge. We

define immunological signatures associated with BCG-induced protection other than

canonical Th1 immunity. Like interferon-γ, lung homogenate levels of IL-17 correlated

with disease. However, the presence of Th1/17 CD4 T cells were an immune feature that

correlated with protection. CC mice can be used to define correlates of protection and to

identify vaccine strategies that protect a larger fraction of genetically diverse individuals

instead of optimizing protection for a single genotype.

Introduction

The introduction of Bacillus Calmette Guerin (BCG) as a vaccine in the last century has contributed significantly to our ability to control and contain tuberculosis (TB), which has been a threat to global health for millennia. BCG is the only vaccine approved for preventing tuberculosis and over 4 billion doses have been administered in over 150 countries (*1*). Although BCG prevents extrapulmonary TB in infants, it is less effective in averting pulmonary disease in adolescents and adults (*2-5*). Complicating the use of BCG is its variable efficacy in different regions of the world (*6*). While the development of more effective TB vaccines is recognized by the WHO as an important yet unmet medical need, the lack of identifiable correlates of protective immunity to TB is an impediment towards this goal.

Why is identifying immune correlates of protection following BCG difficult? TB is primarily a human disease that is transmitted person-to-person by aerosolized *Mycobacterium tuberculosis* (Mtb). No animal model recapitulates all the features of human TB, although non-human primates come close (*7, 8*). Although imperfect, the mouse model has facilitated mechanistic studies that defined many of the critical immunological pathways required for host resistance, permitted in vivo antibiotic testing, and accelerated preclinical vaccine development. As inbred mouse strains are genetically identical, a strength of the mouse model lies in its "infinite" reproducibility. For experimental immunology, C57BL/6 mice are the preferred inbred strain as most genetically modified mice are produced using its genetic background. It is also the most frequently used mouse strain for TB research. However, several features of C57BL/6 mice suggest it is an outlier strain for TB research. Immunologically, it only expresses a

single class II MHC molecule (i.e., I-A[b]). Pathologically, it develops atypical pathology

(i.e., pneumonitis) following Mtb infection. Finally, and perhaps most importantly, it is

among the most resistant of all inbred strains to Mtb infection, which makes it difficult to

identify conditions that make C57BL/6 mice more resistant to disease. Investigators have

addressed these limitations by using F1 hybrids and susceptible strains to increase MHC

and genetic diversity and identify strategies to increase host resistance (*9-12*). However,

even among different inbred strains, the genetic diversity is limited (*13*). This is a problem

not only for TB research, but for many areas of biomedical research.

The mouse genetics community responded to these limitations of inbred mice by

developing a mouse resource that captures the considerable genetic diversity of the *Mus*

species. Starting with eight founder mouse strains, including representatives from all

three *Mus musculus* subspecies and three wild-derived strains, an eight-way funnel

breeding scheme led to random assortment of the founder genomes among the progeny

(*14*). The progenies were used to derive recombinant inbred mouse strains, which

became known as Collaborative Cross (CC) strains. Each CC strain is a unique genetic

representation (i.e., genotype) of the eight founder strains. Importantly, as they are inbred,

they are a renewable resource, and an unlimited number of experiments can be

performed with the same genotype, which is important for identifying experimental

correlates and defining mechanisms.

We previously reported that the CC strains have a large variation in primary

susceptibility to TB (*15, 16*). Here we leverage the CC model to understand the how host

genetics affects vaccine-induced protection against Mtb. We find that the genetic

background of the host has a major effect on whether BCG induces protection. There was

a poor correlation between inherent host susceptibility and BCG-mediated protection,

indicating that vaccines can mediate protection even in individuals with a genetic

propensity to TB. Interestingly, the T cell response following BCG vaccination and Mtb

challenge was diverse, with some immune features universally correlating with outcome

of vaccination while others only were important in subsets of CC strains. We propose that

the use of CC mice as a preclinical model can facilitate the development of vaccines that

protect a diverse range of genotypes against TB.

Results.

BCG vaccination protects only a subset of CC strains against TB.

To establish how the host genetic background affects vaccine mediated protection against Mtb, 24 CC lines (Table 1) were vaccinated subcutaneously with BCG and rested for 12 weeks. Then, the mice were infected with lose dose aerosolized H37Rv.YFP (*17*). Four weeks after infection, the reduction in lung and spleen CFU were determined as a measurement of protection. Mice were initially tested in four batches, with each batch including six different CC strains and C57BL/6 mice as a reference strain.

The aggregated lung CFU from the unvaccinated and BCG-vaccinated CC strains (Fig.1A, top) and C57BL/6 mice (Fig.1A, bottom) shows that as a population, CC mice are protected by BCG similarly as C57BL/6 mice. However, the variance in the lung CFU of CC mice is greater. When the data is segregated by individual CC strains, we can see that the variance reflects different levels of protection conferred by BCG vaccination in each CC strain (Fig.1B, top). The CC strains were classified as being protected by BCG if there was a statistically significant difference between unvaccinated and vaccinated mice. The $\Delta\log_{10}$ lung CFU was calculated for each CC strain, and they were ranked from most protected (CC037, $\Delta\log_{10}$ CFU = 1.6) to least protected (CC040, $\Delta\log_{10}$ CFU = –0.6) (Fig.1B, bottom). BCG vaccination exacerbated subsequent Mtb infection in the lung for two CC strains (CC003 and CC040), similar to what had previously been observed in the NZO founder strain (*15*). Importantly, BCG only led to significant reductions in the lung Mtb burden in 13 of the 24 CC strains.

We next asked whether the protective effect of BCG vaccination correlated with the inherent ability (i.e., in the unvaccinated state) of each CC strain to control Mtb

replication in their lungs. We categorized the 24 CC strains relative to C57BL/6 mice,

which we used as a reference strain. The lung CFU of unvaccinated mice defined the

inherent susceptibility or resistance to Mtb (susceptibility, y-axis). Next, we considered

whether each CC strain was protected or not based on the $\Delta\log_{10}$ lung CFU (relative

BCG-induced protection, x-axis). The plot was centered on C57BL/6 mice, which

represented a median strain in terms of lung susceptibility and protection and divided the

24 CC strains into four different groups, represented by four quadrants (Fig.1C). Based

on lung CFU, there is a poor correlation between intrinsic susceptibility of the mouse

strain and protection conferred by BCG at this time point (r=0.104, p=0.62).

The same analysis was performed using the spleen bacterial burden from infected

mice. As was observed in the lung, the CC population was similarly protected as C57BL/6

(Fig.1D). The range of spleen protection extended from most protected (CC026, $\Delta\log_{10}$

CFU = 2.56) to least protected (CC045, $\Delta\log_{10}$ CFU = 0.13) (Fig.1E). BCG significantly

protected against systemic infection in 17 of the 24 CC strains (Fig.1E). There was also

no correlation between susceptibility and protection in the spleen (r=0.089, p=0.67).

Interestingly, while C57BL/6 mice were a "median" strain in the lung analysis, C57BL/6

mice had among the highest spleen CFU, suggesting they disseminated more, but were

among the most protected mouse strain following BCG vaccination (Fig.1F). Among the

CC strains, protection in the lung and the spleen were moderately correlated (r=0.513,

p=0.0087).

In follow-up studies, 13 of the 24 CC strains were retested to confirm the initial

phenotype, with emphasis placed on strains that represented varying levels of protection

and susceptibility (Table 1). Thus, using 24 CC strains, with diverse genotypes, we find

that less than 50% of the strains were protected by BCG, and that protection did not

correlate with the intrinsic susceptibility of the mouse to Mtb infection. Finally, the lack of

correlation between the reduction in lung versus spleen highlights the importance of

tissue-specific immunity and the problems inherent in protecting the lung against

infection. We next assessed how BCG vaccination altered the histological appearance of

TB lesions in the lung.

BCG modifies the lung lesions caused by Mtb in a strain-specific manner.

We assessed how BCG vaccination modified lung lesion development after Mtb

infection in 19 different CC strains plus C57BL/6 mice. A board-certified veterinarian

pathologist (G.B.), blinded to vaccination status of each strain, examined the lung for

necrosis, neutrophilic and lymphocytic infiltrates, lesion size and number. Within each

pair, the vaccinated group was correctly identified in 15 of the 20 strains analyzed based

on reduced severity of the histopathology. In some CC strains that were protected by

BCG, increased lymphocytic infiltration, and reduced necrosis were frequently observed

after vaccination (Table 2). CC025 (Fig. 2, top left) and CC037 (Fig. 2, middle left) were

both susceptible to primary Mtb infection, and the infected lung showed innate immune

cells and tissue necrosis. BCG vaccination protects these mice by promoting lymphocyte

recruitment and in CC025 BCG vaccination reduced necrosis. However, a reduction in

lung CFU associated with BCG vaccination was not always associated with an improved

histological appearance of the lung. Despite a significant reduction in lung CFU after

BCG, no discernable histopathological differences were observed between unvaccinated

and BCG vaccinated CC072 mice, and both groups had non-necrotizing lymphocyte-rich

granulomas (Fig. 2, bottom left).

BCG did not significantly reduce lung CFU in eight of 20 CC strains. Two of the

non-protected CC strains, CC003 (Fig. 2, top right) and CC024 (Fig. 2, middle right) had

no discernable histological differences in granuloma structure, content, or severity

between control and BCG vaccination when evaluated blindly. In these 2 strains, lesions

were generally small to medium-sized, non-necrotizing, with abundant perivascular and

peribronchiolar lymphocytes. Given the intermediate susceptibility of the CC003 strain,

and BCG's inability to alter lung CFU or change the lesions, the data suggests that BCG

does not change CC003 innate or adaptive immune responses to Mtb challenge.

Likewise, CC024 is unprotected by BCG vaccination, nor are lung lesions altered,

suggesting that neither the innate nor adaptive responses of this resistant strain to

primary Mtb infection is further enhanced by BCG vaccination. The natural resistance of

CC024 strain was not further improved. In contrast, CC004 (Fig. 2, bottom right) did have

discernable histopathological changes when controls and BCG vaccinees were evaluated

blinded. The differences attributable to BCG vaccination were the greater lymphocytic

infiltration with denser perivascular and peribronchiolar lymphocytes. The altered lesions

detected after vaccination suggests that the immunity induced by BCG in CC004 mice

fails to control bacilli. Together, these data show that the host genetic background impacts

the granuloma content, structure, and quality of the immune response to BCG and Mtb

challenge in a manner that is not wholly predictable.

BCG-protected CC strains have an altered lung microenvironment after Mtb infection

The Mtb-infected lung is a complex environment of resident and recruited cells that produce inflammatory mediators and counter-regulatory (e.g., anti-inflammatory) signals. We sought to determine whether BCG vaccination primes the immune response to generate a different lung microenvironment upon Mtb challenge, perhaps one that is less conducive to bacillary replication and transmission. Cytokine levels in lung homogenates from PBS control and BCG-vaccinated C57BL/6 or CC mice were measured 4 weeks post infection using the MD31 Discovery Assay Array.

Given the critical role of IFNγ in Mtb control, its levels in control and BCG-vaccinated CC mice were compared. In C57BL/6 mice and many of the protected CC strains, including CC037, CC025, CC059, CC031, and CC011, IFNγ concentrations in BCG-vaccinated mice were reduced compared to unvaccinated mice (Fig.3A). However, this pattern did not hold for all protected strains. No changes in IFNγ levels were detected the CC032, CC042, CC072, CC026 and CC001 strains, despite being protected by BCG. Thus, when considered alone, neither the absolute level of IFNγ nor its changes after vaccination could be used to predict protection.

We next asked if there was a correlation between the different cytokines in the lung homogenate and lung CFU, in either unvaccinated or BCG-vaccinated mice. We asked if this correlation differed among protected and unprotected CC strains, and C57BL/6 mice. (Fig.3B, top left). There was a strong correlation between CFU and lung IFNγ levels (R=0.71, p=0.0004), in unvaccinated C57BL/6 mice but not in BCG-vaccinated C57BL/6 mice. Thus, IFNγ correlated with bacterial burden but not with protection, as observed in

prior studies (*18, 19*). The CC strains protected by BCG had a similar pattern. However,

for unprotected CC strains, a similar correlation between IFNγ and CFU was identified in

unvaccinated, and BCG vaccinated mice, which we suggest is evidence that these strains

were unprotected by BCG vaccination. This pattern held for T cell cytokines (IFNγ and IL-

17), mediators produced by myeloid cells (IL-1β, IL-6, KC, and MIP-1α), and TNF (Fig.3B

and not shown). These data suggest that protected and unprotected CC strains respond

differently to BCG, and that the protected lung environment differs immunologically from

an unprotected environment.

Other cytokines were of interest. LIX (CXCL2) was negatively associated with lung

CFU in unvaccinated but not in BCG-vaccinated C57BL/6 mice (–0.67, p=0.0013). This

correlation was only observed in C57BL/6 mice, and LIX levels poorly correlated with lung

CFU in CC strains (Fig.3C). RANTES strongly correlated with lung CFU in C57BL/6 mice,

independent of vaccination status, but was only weakly correlated in CC strains (Fig.3D).

Like RANTES, MIP-1β was also correlated with lung CFU, independently of vaccination

status (Fig.3E). This was most convincing in unprotected CC mice (R=0.71-0.78), with

moderate correlations in vaccinated C57BL/6 and protected CC strains (R=0.46-0.49).

Four cytokines (G-CSF, M-CSF, MIP-2 and LIF) were strongly correlated with lung

CFU in unvaccinated CC strains, regardless of protection status (R=0.57-0.8, Fig.3F).

Again, these correlations were lost in the BCG-vaccinated protected CC strains. None of

these cytokines were made in significant amounts by C57BL/6 mice or were poorly

correlated with CFU. These data show how differences in the host genetic background,

result in varied lung histology and cytokines, which dramatically alter the lung

inflammatory environment during Mtb infection.

Host genetic background and not BCG vaccination determines T cell subset

distribution in the lung after Mtb infection.

We hypothesized that changes in the frequency of T cell subsets would correlate

with protection induced by BCG vaccination. An 18-parameter flow cytometry panel was

used to identify and enumerate different immune subsets (Figure S1). As T cells are

important in protection against TB, we determined the frequency and relative number of

different T cell subsets modulated by BCG vaccination. Not all parameters were obtained

for all 24 CC strains (see methods). We quantified CD4 and CD8 T cells, MAIT cells and

NKT cells (Fig.4A, and Figure S2). Memory and effector T cell populations were defined

as central memory (T_{CM}; CD44$^+$CD62L$^+$CD127$^+$), effector memory (T_{EM}; CD44$^+$CD62L$^-$

CD127$^+$), effector (T_{Eff}; CD44$^+$CD62L$^-$CD127$^-$), and resident memory (T_{RM};

CD44$^+$CD103$^+$CD69$^+$) (Fig.4B, 4C). CD4 T cells subsets were defined in all CC strains

based on their expression of the transcription factors Foxp3, Eomes, Tbet and RORγt to

identify Treg, Th1, Th2, and Th17 cells, respectively (Fig.4D). For these T cell subsets,

we observed tremendous strain-to-strain variation. With one or two exceptions, BCG

vaccination did not modulate the distribution of T cells subsets nor was there any

correlation between the relative frequency of these T cell subsets and the outcome of

BCG vaccination. Thus, the frequency of different T cells subsets in the lungs of CC mice

after Mtb infection was largely determined by each strain's genetic background.

Expression of CXCR3 and CX$_3$CR1, which have been used to discriminate

between T cells in the lung parenchymal vs. the vasculature, was analyzed in a limited

number of CC strains (*20*). CXCR3 also marks Th1 cells, and in conjunction with CCR6,

can differentiate between Th1 and Th17 cells, respectively (*21*). Like other markers, we

observed variation in CXCR3 expression between different CC strains, but little or no

variation that correlated with vaccination status or susceptibility. In addition to allelic

variation, CC strains also carry private mutations. For example, we previously reported

that CC042 mice lacks cell surface CD11a expression due to a mutation in the *Itgal* gene

on Chromosome 7 (*22*). Here, we report that CC003 mice don't express CXCR3 (Fig.4E).

This phenotype is associated with the CXCR3 allele of the PWK founder. Finally, a

reduction in CX3CR1$^+$ CD4 and CD8 T cells was observed in several of the protected CC

strains but not in unprotected strains (Fig.4E). This pattern mirrored the reduction in IFNγ

levels observed in some protected CC mice (Fig.3), suggesting that these changes reflect

Th1 modulation.

Finally, we used principal components analysis to holistically visualize the variation

in the T cell phenotyping data. What is striking is the small effect of BCG vaccination in

altering T cell phenotypes, in comparison to large effect of genetic strain differences.

These data all indicate that genetics differences between the CC strains determine the

differences among T cell subsets recruited to the lung, and not BCG vaccination or Mtb

infection. Therefore, we next wished to determine whether there were functional

differences among the T cell subsets in protected vs. nonprotected CC strains.

The patterns of cytokine production differ in CC strains after BCG vaccination and

Mtb infection.

As the cytokines produced by T cells, including IFNγ, TNF, and IL-21, among

others, are important in control of Mtb infection, we hypothesized that changes in T cell

function would be associated with BCG-induced protection. To measure function, we

stimulated lung mononuclear cells (MNC) from each mouse with anti-CD3 or the P300

megapool, which contains 300 peptides representing epitopes from 70 Mtb proteins that

are frequently recognized by human CD4 T cells and are also recognized by murine CD4

and CD8 T cells (23, 24). Twenty-six cytokines and chemokines in the culture

supernatants were measured using the Legendplex platform 24 hours after stimulation.

There was little difference in the ability of lung MNC from mock or BCG vaccinated Mtb-

infected C57BL/6 mice to secrete different cytokines or chemokines following P300

stimulation. In contrast, we detected differential production of IFNγ, TNF, IL-2 or IL-17, all

cytokines that have been associated with immunity to Mtb infection, among mock vs. BCG

vaccinated CC strains (Fig.5A). Interestingly, the MNC from BCG vaccinated CC037,

CC031, CC72 and CC001 strains all produced more IL-17 than their respective controls.

In contrast, none of the unprotected strains produced significant amounts of IL-17, except

for CC023, and its lung MNC produced less IL-17 after BCG vaccination.

In parallel, a 14-parameter flow cytometry panel measured intracellular levels of

IFNγ, IL-2, IL-17, TNF, IL10, and IL-4, and surface expression of CD107 in both CD4 and

CD8 T cells (Figure S3). BCG vaccination had a minimal effect on CD8 T cell cytokine

responses to P300 stimulation in both C57BL/6 mice as well as most CC lines examined

(Fig.5B). However, BCG clearly modified the CD4 T cell cytokine responses in some CC

strains (Fig.5C). Patterns of cytokine production were observed in BCG vaccinated CC

mice that were not observed in C57BL/6 mice. For example, while BCG vaccination did

not alter the frequency of IFNγ[+] or TNF[+] CD4 T cells in CC037 mice, it did increase IL-2

and IL-17 responses. BCG vaccination of CC031 mice increased the frequency of CD4 T

cells producing IFNγ, IL-2 and TNF but not IL-17. Finally, CD4 T cells from BCG

vaccinated had CC072 had increased IFNγ and IL-2 but not TNF or IL-17. Interestingly,

no discernable differences in cytokine responses were observed in most of the non-

protected CC mice, except for a decrease in IFNγ responses in CC023. Thus, the cytokine

responses in both control and BCG vaccinated CC mice differ greatly from classic

C57BL/6 model. Increased cytokine responses after T cell stimulation were more likely to

be detected in protected CC strains. Moreover, BCG vaccination led to quantitative and

qualitative changes in the type of T cell responses that persisted after Mtb challenge.

Correlates of Protection.

We then turned to multivariate approaches to discern how BCG-induced immune

changes in the lung mechanistically lead to reductions in lung CFU in CC strains. To do

so, the data from the lung homogenate cytokines, intracellular cytokine staining, T cell

phenotyping, and LEGENDplex were combined, and for each vaccinated mouse the

transformed feature data was calculated as: [(vaccinated CC subject–

mean(unvaccinated CC strain]. Then, a partial least squares regression (PLSR) model

with elastic net feature selection was created to identify features that changed with BCG

vaccination in a way that maximally covaried with BCG-induced changes in lung CFU.

The model performed significantly better than null models, and the first two latent

variables in PLSR model account for 24% and 8%, respectively, of the variability in the

lung features (Figure 6A, Figure S4A). The elastic net-selected features in the model

included features from all four datasets (Fig.6B, 6C). By plotting the scaled values of the

elastic net-selected features for all CC strains, ordered from most protected to least

protected, what is most apparent is that the protected strains all had BCG-induced

reductions in several inflammatory cytokines such as MIG, CCL3 and CCL4 (i.e., MIP-1α

and -1β), IL-6, RANTES, and IL-17. The decrease in these inflammatory mediators could

be a sign of resolved infection, in contrast to the unprotected strains which had more

inflammatory cytokines which could signify unimpeded infection. Protected strains also

had a greater vaccination-induced increase in activated CD4 T cells. Contrarily, protected

strains had decreased activated CD4⁻CD8⁻ T cells producing T-bet. Importantly, IL-2,

unlike several other cytokines, was increased in protected strains. Thus, BCG vaccination

leads to qualitative and quantitative changes in the T cell response and in the immune

environment of the lung after Mtb infection in CC genotypes that are associated with

protection.

Immune responses in protected and unprotected CC strains differ following Mtb

infection

While the PLSR model highlights potential correlates of lung protection across all

genotypes, it may miss correlates that only apply to certain subsets of CC strains. We

hypothesized that naturally resistant and susceptible strains may be protected in different

ways. To test this, we created separate partial least squares discriminant analysis

(PLSDA) models for resistant and susceptible CC strains and examined the features that

best separated the BCG-protected and non-protected strains in each of these categories.

(Figure 1C). The data used for the models was processed the same way as in the PLSR

models, although we also looked at the vaccinated and unvaccinated mice separately in

addition to the vaccine-induced differences. Our first analysis compared resistant CC

strains that were protected by BCG (n=4) or not (n=8), based on lung CFU. As described

earlier (Figure 4), the unvaccinated and vaccinated mice of each CC strain were very

similar and clustered together when the elastic net-selected features of the model were

hierarchically clustered, indicating that the genetic background was an important

determinant of outcome (Figure 7A). The protected and unprotected mice were both split

into multiple clusters, showing that there were multiple distinct types of immune

responses that were associated with protective immunity in resistant mice. The model

could predict protected vs non-protected CC strains accurately and thus there is a

multivariate signature of these categories (Figure S4B). However, there were no clear

individual features distinguishing the categories. In contrast, the vaccine-induced model

comparing protected and non-protected resistant strains did yield more univariate trends

in addition to the strong multivariate signature. Namely, both CD4 and CD8 T cells making

IL-2 are increased in protected strains, while several lung homogenate cytokines as well

as double negative T cells expressing CD107a and producing IL-17 decreased in

protected strains (Figure 7B).

We next analyzed CC strains that were inherently more susceptible than B6 mice,

and were protected by BCG (n=4) or not (n=8) (Figure 7C). This yielded two clear

phenotypes of susceptible, non-protected CC strains. One subcluster comprising CC004

and CC044 had high levels of T_{CM}, CD4$^+$Foxp3$^+$, CD8$^+$PD-1$^+$ and polyfunctional CD4$^-$

CD8$^-$ T cells. The other cluster had high levels of lung MIP-1a and CD3$^-$Foxp3$^+$ T cells,

and fewer polyfunctional CD4$^-$CD8$^-$ T cells. Interestingly, the unprotected strains with

high CD4$^+$Foxp3$^+$T cells had low CD8$^+$Foxp3$^+$ T cells, and the unprotected strains with

high CD8$^+$Foxp3$^+$ T cells had low CD4$^+$Foxp3$^+$ T cells. When looking at the protection

model with the susceptible strains, several features were similar to the model with all

mice. In this model it is evident that while IL-17 in the lung homogenate decreases with

BCG vaccination in protected strains, the amount of IL-17 when stimulated with the P300

peptide pool increases. (Figure 7D). Overall, the features that distinguish protected and

non-protected strains are similar for naturally resistant and susceptible mice.

Finally, we compared susceptible CC strains that were unprotected by BCG (i.e.,

poor immunity) to inherently resistant CC strains that were protected by BCG (i.e., good

immunity). Unsurprisingly, there is clear separation of categories in the model with all

mice. "Poor immunity" CC strains tended to have high levels of inflammatory monokines

(Figure 7E, 7F). "Good immunity" CC strains high levels of RANTES after stimulation with

anti-CD3. A subset of good immunity strains including CC031 and CC011 also have high

frequencies of IFNγ-producing CD8 T cells and various subsets of CD4$^-$CD8$^-$ T cells

including those producing IFNγ or expressing CD11a. In contrast, a subset of poor

immunity strains had high levels of polyfunctional CD4$^-$CD8$^-$ T cells and ILCregs. The

protection model revealed similar features with the interesting addition of activated CD4

T_{RM} cells increasing with BCG vaccination in poor immunity but not good immunity CC

strains.

We find that immune features can be identified using PLSDA that can categorize

the inherent susceptibility and response to BCG of CC strains following Mtb infection. The

expression of different immune features was variable between CC strains even in the

same protection category, indicating no single path to protection. Looking at more refined

subgroups of CC strains allowed us to find features that define subclasses of protected

and non-protected strains and highlight features that did not emerge in the PLSR model

such as non-T_{RM} effector and central memory T cell populations.

Th1/17 cells detected in a subset of CC mice that are protected by BCG vaccination

An aim of this study was to identify factors that were associated with protection in

genetically diverse CC mice after vaccination and Mtb challenge. Vaccination enhanced

IL-17 responses in the CC037, CC001, CC072 and CC031 strains, which were all

protected by BCG (Figure 5A, 5B). As the CC strains that produced IL-17 after antigen

stimulation also had strong IFNγ response, we assessed these T cell responses for dual

production of IFNγ and IL-17. BCG vaccinated CC037 mice had the highest IL-17 levels

in lung homogenate (Figure 5A, 5B), and after stimulation with P300 or anti-CD3 mAb,

CD4 T cells were identified that produced both IFNγ and IL-17 (Figure 8A). Dual IFNγ$^+$IL-

17$^+$ CD4 T cells were not detected in C57BL/6 mice after P300 stimulation, regardless of

vaccination status. BCG vaccination significantly increased the frequency of IFNγ$^+$IL-17$^+$-

producing CD4 T cells in the CC037, CC072 and CC001 (Figure 8A, 8B). Further analysis

identified BCG primed CD4 T cells from CC037 and CC001 to coexpress both Tbet and

RORγt after Mtb infection (Figure 8C, 8D), which are the master transcription factors for

Th1 and Th17 cells, respectively.

We next measured the polyfunctionality of the T cell response and expanded our

cytokine analysis to include IL-2 and TNF. In addition to IFNγ and IL-17, an increase in

the proportion of CD4 T cells producing IL-2 or TNF was detected (Figure 8E), indicating

that CD4 T cell polyfunctionality was increased in BCG vaccinated mice. A detailed

analysis of the different polyfunctional populations revealed different distributions of

Th1/17 cells within each CC strain. IFNγ⁺IL-2⁺TNF⁺IL-17⁺ cells were significantly

increased in CC037 following BCG vaccination (Figure S5, red box). Similarly,

IFNγ⁺TNF⁺IL-17⁺ cells were increased following BCG vaccination in CC072 and CC001

(Figure S5, blue box). BCG led to a decrease in the frequency of IFNγ "single positive'

CD4 T cells in all four CC lines (Fig. S5, purple box). However, among these CC strains

only CC037 had a significant decrease in IFNγ⁺TNF⁺ cells. This contrasts with C57BL/6

mice where BCG vaccination had a minimal effect on CD4 T cell polyfunctionality.

Significant decreases were observed for IFNγ⁺TNF⁺ dual producers and IFNγ single

positive cells in Mtb-infected BCG-vaccinated C57BL/6 mice, consistent with the

observation that Th1 dominant response is associated with decrease bacterial burden in

this background.

Finally, we performed a correlation analysis between all 60 immune features and

the normalized lung CFU from unvaccinated control and BCG vaccinated mice. For

C57BL/6 mice, IFNγ-producing and polyfunctional CD4 T cells correlated with lung CFU.

In CC037 mice, IFNγ-producing CD8 T cells correlated with lung CFU. Importantly, many

features were negatively correlated with lung bacterial burden in CC037 mice (Figure 8F).

The frequency of CD4⁺Roryt⁺, and antigen-stimulated IL-17 secretion and IFNγ/IL-17

dual-producing CD4 T cells were significant even after correction for multiple testing,

suggesting that these features were associated with protection following BCG. There was

a strong correlation between the frequency of IFNγ/IL-17 dual-producing CD4 T cells and

lung CFU in CC037 mice (Spearman *r* = -0.8947) (Figure 8G).

Discussion

Using CC mice to model an outbred population, our study of 24 CC strains sought to determine how pre-existing immunity to BCG modified the immune response to Mtb infection in a genetically diverse population. An important outcome of this study is our ranking of how BCG induced protection varies across the 24 CC strains were studied, compared to control unvaccinated mice and relative to C57BL/6 mice, which were used as a reference strain. We categorized the different CC strains as being either protected or unprotected by BCG against Mtb infection, based on whether vaccination led to a statistically significant reduction in lung bacillary burden measured four weeks after low dose aerosol infection. Based on whether the different mice were susceptible or resistant, and protected or not, relative to C57BL/6 mice, we divided the 24 CC strains into four phenotypes (i.e., quadrants). We found that different immune parameters were associated with protection in the four different groups. The surprising result was that BCG does not protect all mouse genotypes against Mtb, and no protection was detected in nearly half of the CC strains tested. Thus, these results extend our analysis of the eight CC founder strains, which also varied in their ability to be protected by BCG vaccination (*15*). Not only do environmental mycobacteria and genetic variations in BCG strains used around the world interfere with the ability of BCG-induced protection, but we report now that host genetics should be considered a third important barrier that prevents vaccine-induced protection to TB.

An important question is why BCG doesn't induce protection in all CC strains? Protection was largely defined based on reductions in lung CFU. Four weeks after low dose Mtb infection was chosen as it is a standard time point used in many murine vaccine

studies. Although the complexity of this experiment precluded performing other time

points, it is possible that our rankings may have changed if other time points were used,

just as the rankings differ for lung vs. splenic CFU reduction. Interestingly, the CC strains

that were unprotected by BCG can be subdivided based on their lung pathology and

immune responses. None of the unprotected CC strains had immune responses that were

modified by BCG vaccination, but some (e.g., CC004, CC040) had changes in lung

histology (e.g., more lymphocytic infiltrates) that suggested that a memory recall

response. Thus, we suggest that BCG induced an immune response in some CC strains

(e.g., CC004, CC040), but it was insufficient to control lung CFU. Other strains may have

generated a suboptimal response to vaccination (i.e., low responders; CC003, CC024,

CC041).

We believe that the variation in host responses makes this an ideal platform for

testing the efficacy of preclinical vaccines in the context of a genetically diverse

background. Only a fraction of individuals infected with Mtb will develop disease. While

there exist well-defined risk factors for the development of TB, the factors that drive the

emergence of disease is largely unknown in most people. Similarly, the factors that

influence BCG efficacy remains controversial. In addition to risk factors that disrupt cell

mediated immunity, there are also genetic factors (i.e., MSMD, polygenic), which

predispose certain people to TB. Similar genetic factors could also contribute to variation

in BCG efficacy against pulmonary Mtb. Our analysis shows that while certain mouse

genotypes cannot be protected by BCG, the inherent susceptibility or resistance of the

mouse strain is not a predictor of vaccine efficacy. One caveat is that we have only tested

the efficacy of one vaccine in the CC strains. It is possible that vaccines that work through

alternate mechanisms may be able to protect CC mice that are refractory to BCG

vaccination.

Genetics affects host resistance to TB (*16*). Our studies show how dramatically

host genetics affects vaccine-induced immunity. Different inbred recombinant mouse

strains derived from the same eight founders can be protected or not by BCG. Importantly,

the ability of these mouse strains to be protected by BCG was largely independent of their

inherent susceptibility to Mtb infection. This suggests that a priori, vaccines are

appropriate interventions even in individuals that might have a genetic susceptibility to

TB. Importantly, we believe that diverse human populations include genotypes that

generate suboptimal, or even detrimental, responses to BCG. We expect that our data

will help to establish CC strains as a novel animal resource that can facilitate the discovery

of correlates of vaccine-induced protection and accelerate preclinical vaccine testing.

Instead of optimizing vaccine concepts in a single mouse genotype (e.g., C57BL/6 mice),

identifying vaccines that are effective in numerous different genotypes (i.e., genetically

distinct individuals), as modeled by CC or DO mice, represents an alternative paradigm

for preclinical vaccine development.

Levels of IL-17 in lung homogenate is strongly correlated with lung CFU in CC

strains, co-varies with IL-1β and IL-6, two cytokines that are involved in skewing of Th17

cells, and with TNF, KC, MIP2, and G-CSF, all which are induced by IL-17. In BCG

vaccinated CC strains that were protected, the correlation with IL-17 was lost. We

suggest that like IFNγ, IL-17 levels are a correlate of disease. In contrast, in three of the

12 protected strains (i.e., CC037, CC072 and CC001), we detected CD4 T cells that

produced IFNγ and IL-17 after Mtb infection. These cells are phenotypically similar to

Th1/17 cells, or Th1*, that have been identified following TB infection (*25, 26*), and have

recently been associated with late-forming, low burden granulomas in a NHP TB infection

model (*27*). None of the 13 unprotected strains had this phenotype, and these dual

producers were detected only in BCG vaccinated mice and not in sham controls, nor were

they detected in the C57BL/6 reference strain. Thus, this is an example of an immune

response that BCG primes but is not manifested until the CD4 T cell response is recalled

by Mtb infection.

Finally, we wished to identify the components of the immune response stimulated

by BCG, which were subsequently recalled after Mtb infection and associated with

protection. We hypothesized that CC strains protected by BCG would share common

immunological features. We postulated that by comparing protected to unprotected CC

strains would allow us to filter out components of the immune response that were elicited

by vaccination or infection but not associated with protection. As BCG is known to elicit T

cell responses that can mediate protection, we extensively characterized the T cell

immunity. Analysis of the different CC strains showed that there was considerable

diversity in the composition of the T cells recruited and maintained in the lung after

infection. However, most of the variability was determined by the host genetic background

and BCG vaccination had little impact on the T cell subsets recruited to the lung. In

contrast, several of the CC strains protected by BCG vaccination had evidence of

functional skewing of the T cell responses. Despite there being several different flavors

of immunity, we were able to identify an immune signature that was associated with BCG-

elicited protection against TB.

Acknowledgements

The following reagents were obtained through the NIH Tetramer Core Facility (contract number

75N93020D00005) for providing MR-1 and CD1d tetramers. We thank the Vanderbilt Medical

Center Digital Shared Histology Resource (DHSR) for scanning and hosting our histology data.

Author contribution

Declaration of interests

The authors declare no competing interests.

Funding

Supported by P01 AI123286 (C.M.S., M.F., S.M.B,), Digitization of glass slides was funded by

R01 HL14541 (G.L.B).

Materials and Methods

Mice

Female C57BL/6/J (#0664) were purchased from The Jackson Laboratory. Female mice from 24 CC strains were purchased from the UNC Systems Genetics Core Facility (SGCF) between July 2018 and August 2019. The 24 CC strains used in this study include: CC001/Unc, CC002/Unc, CC003/Unc, CC004/TauUnc, CC010/GeniUnc, CC011/Unc, CC012/GeniUnc, CC023/GeniUnc, CC024/GeniUnc, CC025/GeniUnc, CC026/GeniUnc, CC031/GeniUnc, CC032/GeniUnc, CC035/Unc, CC036/GemUnc, CC037/TauUnc, CC040/TauUnc, CC041/TauUnc, CC042/GeniUnc, CC044/Unc, CC045/GeniUnc, CC059/TauUnc, CC061/GeniUnc,, CC072/TauUnc. More information regarding the CC strains can be found at http://csbio.unc.edu/CCstatus/index.py?run=AvailableLines.information.

All animal studies were conducted using the relevant guidelines and regulations, and approved by the Institutional Animal Care and Use Committee at the University of Massachusetts Medical School (UMMS) (Animal Welfare A3306-01), using the recommendations from the Guide for the Care and Use of Laboratory Animals of the National Institutes of Health and the Office of Laboratory Animal Welfare.

Bacterial strains

Bacteria expressing yellow fluorescent protein (sfYFP) were generated as described (17). Mice were vaccinated with 105 CFU BCG (BCG-SSI strain, resuspended in 0.04% Tween/PBS) subcutaneously (100 uL/injection).

<u>Aerosolized Mtb infection of mice</u>

Mice were infected by the aerosol route as previously described (*17*). Frozen bacterial

stocks were thawed and sonicated for 1 minute and then diluted into 5 ml of 0.01%

Tween-80 in PBS. The diluted bacterial suspension was aerosolized to infect mice using

a Glas-Col chamber (Terre Haute). The average number of bacteria delivered into the

lung was determined for each experiment by plating lung homogenate from 4-5 mice 24

hours after infection and ranged between 40-150 CFU/mouse.

<u>Bacterial burden in lung and spleen</u>

Infected mice were euthanized at pre-determined timepoints, and the left lung or whole

spleen were homogenized using 2 mm zirconium oxide beads (Next Advance) in a

FastPrep homogenizer (MP Biomedicals). Tissue homogenates were serially diluted and

plated on 7H11 agar plates (Hardy Diagnosis). CFU was enumerated after 19-21 days

of incubation at 37°C and 5% CO_2.

<u>Lung cell preparation</u>

Single cell suspensions were prepared by homogenizing lungs using a GentleMACS

tissue dissociator (Miltenyi), digesting with 300 U/ml collagenase (Sigma) in complete

RPMI (10% FBS, 2 mM L-Glutamine, 100 units/ml Penicillin/Streptomycin, 1 mM Na-

Pyruvate, 1X Non-essential amino acids, 0.5X Minimal essential amino acids, 25 mM of

HEPES, and 7.5 mM of NaOH) at 37°C for 30 minutes, and followed by a second run of

dissociation using the GentleMACS. Suspensions were then sequentially filtered through

70 μm, treated with ACK lysis buffer (Gibco), then filtered through 40 μm strainers.

<u>Flow cytometric analysis</u>

Cells were first stained with Zombie Fixable Viability dye (Biolegend) for 10 minutes at

room temperature (RT), after which cells were stained with antibodies in autoMACS

running buffer (Miltenyi) containing 5 ug/ml of anti-mouse CD16/32 (BioXcell) for 10

minutes at 4°C. In some experiments, MAIT cells were identified using the MR-1-PE

tetramer (NIH tetramer core facility) at 37°C for 1hr. Cell were then stained with surface

antibody cocktail for 20 minutes at 4°C. In some experiments, CD1d-APC tetramer (NIH

tetramer core facility) were also added at this time to identify NKT cells. To measure

transcription factor (TF) expression, cells were fixed and permeabilized for 30 minutes at

RT, followed by staining for 30 minutes with antibodies to the different TF at RT using the

Foxp3/TF staining buffer set (ThermoFisher). Samples were acquired on Aurora (Cytek)

or MACSquant (Miltenyi).

For intracellular staining, cells were first stimulated for 5 hours with either anti-CD3

antibodies (1 ug/mL, Biolegend) or P300 peptide megapool (*23*) which contains 300

peptides representing 90 antigens, and contains epitopes recognized by both murine CD4

and CD8 T cells. GolgiPlug (BD Biosciences) was also given at this time (0.5ug/well).

Viability and surface staining were performed as described, after which cells were fixed

instead with Fixation/Permeabilization kit for 20 minutes at 4°C. Cells were then stained

with an intracellular cytokine antibody cocktail for 20 minutes at 4°C.

To inactivate the bacteria, samples were fixed with 1% paraformaldehyde/PBS for 1 hour

at room temperature and then washed with MACS buffer. Flow data were analyzed using

FlowJo v10.7.1. Polyfunctionality was examined using the SPICE platform (*28*).

LEGENDplex cytokine analysis

Lung cells were stimulated with P300 peptide megapool (1ug/mL) or anti-CD3 (1ug/mL)

at a concentration of 10^6 cells/mL for 24hrs. Culture supernatants were filtered through

0.2um filter, after which cytokine levels were measured with LEGENDplex kits from

Biolegend. Specifically, we used their mouse T helper cytokine panel (IFNγ, TNF, IL-2,

IL-4, IL-5, IL-6, IL-9, IL-10, IL-13, IL-17a, IL-17f, IL-21, IL-22) as well as their mouse

proinflammatory chemokine panel (RANTES, MIP-3α, Eotaxin, TARC, KC, MCP-1, MIG,

IP-10, MIP-1α, MIP-1β, BLC, LIX, MDC) for a total of 26 cytokines and chemokines.

Lung homogenate

Cytokines and chemokiens were measured from lung homogenates using the 31-plex

discovery Assay (MD31) from Eve technologies (Calgary, Canada).

Histology analysis

The right middle lung lobe from each mouse was immersion fixed in 10% formalin

immediately following euthanasia. Lung tissue was processed, embedded in paraffin,

sectioned at 5um, and stained with hematoxylin and eosin at the Morphology Core Facility

at UMass Chan Medical School for histological examination by a board-certified

veterinary pathologist (GB). Blind classification (BCG vs PBS) within each CC strain was

determined based on the assumption that BCG vaccination would have three possible

effects on lungs: (i) reduce necrosis/neutrophilic infiltrates; (ii) increase lymphocytes

and/or plasma cell abundance; and (iii) reduce granuloma size. Accuracy of the blind

classification was assessed by an independent investigator (AO). All glass slides were

subsequently re-evaluated to more closely examine and describe histopathology features

attributable to BCG vaccination and attempt to find unique responses. Glass slides were

then digitized by Vanderbilt University Medical Center (Nashville, TN) and whole images

generated using Aperio ScanScope (Buffalo Grove, IL, USA) at 400 times normal

magnification with jpeg compression. Representative images were captured using Aperio

ImageScope magnified 40 and 400 times and imported into GraphPad Prism 9 to

generate the panel Figure. Digital images were not adjusted. Whole slide imaging and

quantification of immunostaining were performed in the Digital Histology Shared

Resource (DHSR) at Vanderbilt University Medical Center.

<u>Univariate correlations</u>

To evaluate how C57BL/6 mice, protected CC strains, and non-protected CC strains

compared in their relationships between lung cytokine production and Mtb burden, a

series of univariate Pearson correlations were conducted. Lung homogenate cytokines

and chemokines measurements in the form of concentrations were natural logged and

each of their Pearson correlations with the log10 lung CFU or spleen CFU were

evaluated, with both the cytokine and CFU measurements separated into groups of

C57BL/6, protected CC strains, and non-protected CC strains. A Bonferroni multiple

hypothesis correction was applied to account for the number of correlations calculated.

<u>Multivariate analyses</u>

Multivariate analyses involved combining all four immunological datasets in this study:

the lung homogenate cytokine/chemokine measurements, the ICS, the LEGENDplex

cytokine analysis, and the T cell phenotypic flow cytometry. The analyses also included

all CC mice used in the study in addition to the control C57/BL6 mice.

For the ICS, the data were analyzed as combinatorics of the seven markers measured in

order to best capture polyfunctionality, and further divided by CD4 T cell, CD8 T cell, CD4$^-$

CD8$^-$ T cell, and CD3$^-$ categories. The features were filtered by a minimum mean and

variance percentage of 0.000001. Only P300 stimulated samples were used from the ICS

dataset. The flow cytometry data was in the form of percentages of CD3$^+$ or CD3$^-$ cells,

while the immunoassay data was in the form of concentrations. The combined dataset

was natural log transformed and subsequently z-scored by measurement across all

samples. For models that involved BCG-induced protection, these z-scored values were

further transformed by subtracting the features from each vaccinated mouse and the

mean of the features of the unvaccinated mice from the same mouse strain.

For the PLSR and PLSDA models, elastic net feature selection was performed in 100

trials of 5-fold cross validation, where features that were present in at least 80% of trials

were included in the final model. Each model was compared to two null models: one with

permuted lung CFUs and one with randomly chosen features. The mean squared error

of the prediction of lung CFU (for the PLSR) or the classification accuracy (for the

PLSDAs) was compared to that of the null models in 50 trials of 5-fold cross validation.

<u>Quantification and Statistical Analysis</u>.

Statistical analysis was performed using Graphpad Prism 9. P-values were calculated

using unpaired t test, one-way ANOVA, or two-way ANOVA as indicated in the figure

legends. Multivariate analyses were performed in R (v3.6.0 and 4.0.4) using the ropls

Bioconductor package.

CC strain[a]	Batch[b]	MHC Haplotype[c]	ΔLung CFU$_1$[d] reduction (logCFU)	ΔLung CFU$_2$[d] reduction (logCFU)	ΔSpleen CFU$_1$[e] reduction (logCFU)	ΔSpleen CFU$_2$[e] reduction (logCFU)
CC001/Unc	4	b	0.94	1.45	2.79	1.45
CC002/Unc	3	b	0.83	n/a	0.46	n/a
CC003/Unc	3	g	0.07	-0.61	0.77	0.15
CC004/TauUnc	4	b	0.05	0.08	-0.16	-0.12
CC010/GeniUnc	1	b	0.53	n/a	0.92	n/a
CC011/Unc	3	g7	1.04	n/a	2.08	n/a
CC012/GeniUnc	4	wsb	0.79	n/a	0.89	n/a
CC023/GeniUnc	1	b	1.17	-0.09	1.47	0.32
CC024/GeniUnc	2	cast	0.15	0.24	0.38	0.62
CC025/GeniUnc	3	pwk	1.40	n/a	0.43	n/a
CC026/GeniUnc	2	cast	0.93	1.05	3.75	1.87
CC031/GeniUnc	3	b	1.22	1.35	1.26	1.60
CC032/GeniUnc	1	b	1.16	n/a	1.06	n/a
CC035/Unc	4	wsb	0.91	n/a	1.13	n/a
CC036/Unc	3	g7	0.62	n/a	1.84	n/a
CC037/TauUnc	1	b	1.37	1.84	1.07	1.05
CC040/TauUnc	2	b	-0.64	-0.28	-0.35	1.14
CC041/TauUnc	2	b	0.14	0.94	0.71	0.94
CC042/GeniUnc	1	b	1.65	0.96	1.00	1.32
CC044/Unc	4	b/g7	0.08	0.13	0.12	0.39
CC045/GeniUnc	4	z	0.24	n/a	0.13	n/a
CC059/TauUnc	2	b/a	1.39	n/a	1.17	n/a
CC061/GeniUnc	1	b	0.93	n/a	1.58	n/a
CC072/TauUnc	2	b	0.97	1.37	1.06	1.41

Table 1. A description of the CC analysis batches.

[a] A group of C57BL/6 mice were included in all batches as a reference strain. Data was normalized to internal C57BL/6 control as indicated in the methods.
[b] The first cohort of CC mice were analyzed in 4 separate batches as indicated.
[c] MHC haplotype of each CC mouse as inherited from CC founders.
[d, e] Thirteen of the 24 CC strains were analyzed a second time to validate the original results and perform more in-depth immune analysis. ΔLungCFU[d] and ΔSpleenCFU[e] for the first experiment and the repeat experiment are listed. Mice that were exacerbated by BCG vaccination are highlighted in red.

[a] Strain	[b] BCG-induced protection	[c] Change in pathology	[d] How did the vaccinated mice differ?
CC003	N	N	No differences. Both groups had mild lesions with perivascular lymphocytic cuffs and a few macrophage-rich lesions
CC041	N	N	No differences. Both groups had mild lesions with perivascular lymphocytic cuffs and a few macrophage-rich lesions
CC024	N	N	No differences. Both groups had small hard-to-find lymphocytic lesions with perivascular and peribroncial lymphocytic cuffs
CC002	N	Y	Ctrl had mild lesions with perivascular lymphocytic cuffs. BCG mice had larger moderate sized macrophage-rich lesions. Lymphocytes were similarly abundant.
CC004	N	Y	Ctrl mice had mild to moderate lesions with lymphohistiocytic infiltrates. BCG mice have more lymphocytes with regions of dense perivascular/peribronchiolar lymphocytic cuffs.
CC036	N	Y	Ctrl mice have mild disease with lymphohistiocytic infiltrates. BCG mice also have lymphohistiocytic infiltrates with few very dense lymphocytic areas
CC040	N	Y	Ctrl mice have moderate necrotizing lesions with few lymphocytes. BCG mice have mild disease - small & hard to find lesions that are lymphocyte rich without necrosis.
CC044	N	Y	Unusual. Ctrl mice have lymphohistiocytic infiltrates with dense lymphocytic cuffs. BCG mice have lymphohistiocytic infiltrates with dense lymphocytic cuffs and lymphoid aggregates/follicules in parenchyma.
CC061	P	N	No differences. Unusual. Mild lesions. Both groups have dense lymphoid aggregates/follicles in parenchyma. Also, perivascular lymphocytic cuffs.
CC072	P	N	No differences. Mild disease. perivascular lymphocytic cuffs with a few macrophage-rich lesions. One mouse has a foreign-body granulomatous bronchitis/pneumonia with intralesional food/bedding/plant.
C57BL/6	P	Y	Ctrl mice had mild to moderate lymphohistiocytic lesions. BCG mice had with more, dense lymphocytic infiltrates
CC001	P	Y	Ctrl mice had moderate lesions with lymphohistiocytic infiltrates and equivocal necrosis. BCG mice had mild lesions with perivascular lymphocytic cuffs and a few macrophage-rich lesions without necrosis
CC023	P	Y	Ctrl mice had moderate to severe lesions with necrosis and minimal lymphocytes. BCG mice have moderate lesions with more lymphocytes and less necrosis.
CC025	P	Y	Ctrl mice have moderate disease with mild lymphocytic infiltrates and necrosis. BCG mice have abundant dense perivascular/peribronchiolar lymphocytic ifiltrates and lymphohistiocytic granulomas and less necrosis.
CC026	P	Y	Ctrl mice had moderate lesions with lymphohistiocytic infiltrates. BCG mice had mild lesions with lymphohistiocytic infiltrates and a few very dense lymphocytic areas
CC031	P	Y	Ctrl mice have moderate disease with lymphohistiocytic infiltrates. BCG mice have abundant dense perivascular/peribronchiolar lymphocytes and lymphohistiocytic granulomas
CC032	Protected	Y	Ctrl mice have mild disease with lymphohistiocytic infiltrates. BCG mice have more lymphocyte rich lesions that are small & hard to find
CC037	Protected	Y	Ctrl mice have moderate disease with lymphohistiocytic infiltrates. BCG mice have abundant dense perivascular/peribronchiolar lymphocytes and lymphohistiocytic granulomas
CC042	Protected	Y	Ctrl mice have mild to moderate disease with few lymphocytes and necrosis. BCG mice have mild disease with more lymphocytes, less necrosis
CC059	Protected	Y	Ctrl mice had mild disease with lymphohistiocytic infiltrates with small regions of necrosis. BCG mice have abundant dense perivascular/peribronchiolar lymphocytes and lymphohistiocytic granulomas without necrosis

Table 2. Summary of histological changes of CC mice.
[a] Nineteen of the 24 CC strains were available for analysis.
[b] The effect of BCG on the lung CFU is from Figure 1.
[c] Assessment by a pathologist of whether differences existed between sham and BCG vaccinated
mice after Mtb infection (n=5/strain/condition).
[d] Summary of observed differences between the groups.

Figure legends

Figure 1. BCG-induced protection in 24 CC strains and C57BL/6 mice.

(A) Lung and (D) spleen CFU of all 24 CC mice (n=120/group; top) or all C57BL/6 mice

(n=20/group; bottom) at 4 weeks post infection. Mice were vaccinated with BCG or

administered PBS and rested for 12 weeks prior to infection. Each point represents an

individual mouse. Line, median. Student's t-test; ****, p < 0.0001.

(B) Protection in the lung and (E) was assessed by comparing the CFU in BCG vs. sham

vaccinated mice for each individual mouse strain (n=5/group) using the Benjamini-

Hochberg procedure to determine the false discovery rate (FDR) (top). The ΔCFU of BCG

vs. sham vaccinated was calculated (bottom). The open bar is the reference strain

C57BL/6 strain (n=20/group). *, p < 0.05; **, p < 0.01; ***, p < 0.001; ****, p < 0.0001.

(C) Correlation in the lung and (F) spleen between susceptibility and protection. The

relative susceptibility (y-axis, defined by CFU in unvaccinated group, relative to the

C57BL/6 reference strain) and the protection conferred by BCG (x-axis, defined by ΔCFU

of vaccinated mice compared to unvaccinated controls, relative to the C57BL/6 reference

strain). The graph is centered on C57BL/6 mice, leading to the different CC strains being

placed in four quadrants based on their relative susceptibility and protection.

Figure 2. Lung histology of CC mice after BCG vaccination and Mtb challenge

Representative lung histology of three CC strains (CC025, CC037, CC072) that were

protected by BCG, after sham or BCG vaccination and Mtb challenge (Mtb and BCG+Mtb,

respectively) (left). Representative lung histology of three CC strains (CC003, CC004,

CC024) that were unprotected by BCG (right). Original magnification, 40X (inset, 400X).

Scale bar, 500 μm (inset, 50 μm).

Figure 3. BCG-protected CC mice have an altered lung microenvironment of CC mice

after Mtb infection.

(A) IFNγ levels were measured in the lungs of either PBS or BCG vaccinated CC mice at 4 weeks

post infection. A two-way ANOVA was performed using Sidak's correction for multiple

comparisons. ***, $p < 0.001$; ****, $p < 0.0001$".

(B) – (F) Pearson correlation between lung CFU and (B) IFNγ, IL-17, TNF, IL-1β, KC and IL-6 (C)

LIX (D) RANTES (E) MIP-1β and (F) M-CSF, MIP-2, G-CSF, LIF, comparing PBS and BCG

vaccinated mice at 4 weeks post infection in C57BL/6 mice (B6), non-protected (NP) or protected

(P) CC mice.

Figure 4. Frequency of various T cell populations in protected and unprotected CC mice

following Mtb infection.

(A) Proportion of CD3$^+$ cell, CD3$^+$CD4$^+$ and CD3$^+$CD8$^+$ T cells in the lungs of PBS and

BCG vaccinated CC mice at 4 weeks post infection.

(B) Proportion of memory CD4 and CD8 T cells in the lungs of either PBS or BCG

vaccinated CC mice at 4 weeks post infection, as defined by CD44$^+$ and various

combinations of CD62L and CD127 expression; CD62L$^+$ CD127$^+$ (central memory),

CD62L$^-$ CD127$^-$ (effector), CD62L$^-$ CD127$^+$ (effector memory).

(C) Proportion of resident memory CD4 and CD8 T cells in the lungs of either PBS or

BCG vaccinated CC mice at 4 weeks post infection, as defined by CD44$^+$ CD103$^+$ CD69$^+$.

(D) Proportion of Th1, Treg and Th17 cells in the lungs of either PBS or BCG vaccinated

CC mice at 4 weeks post infection, as defined by $CD44^+$ and $Tbet^+$ (Th1), $Foxp3^+$ (Treg)

or RoRγt (Th17) expression.

(E) Proportion of CD4 and CD8 T cells expressing either CXCR3 or CX_3CR1 in the lungs

of either PBS or BCG vaccinated CC mice at 4 weeks post infection.

(F) PCA of the T cell phenotyping data. Each point represents the average of usually 5

mice within a given mouse strain that are either BCG vaccinated or unvaccinated. Open

symbols, unvaccinated; closed symbols, BCG vaccinated. Each color represents a

different mouse strain.

(A–E) The data represent the mean ± SEM from one experiment. Two-way analysis of

variance with the Benjamini and Hochberg multiple comparsion method. The FDR was

set to 0.05 and the numbers in the figures are the q value.

Figure 5. Cytokine response in protected and unprotected CC strains following Mtb

infection.

(A) Cells were isolated from the lungs of CC mice at 4 weeks post infection and stimulated

for 24 hours with the P300 peptide pool. IFNγ, IL-2, TNF and IL-17a were measured in

the supernatants.

(B) and (C) Cells from the lungs of CC mice at 4 weeks post infection were stimulated for

5 hours with the P300 peptide pool. Frequencies of CD4 or CD8 T cells that expressed

IFNγ, IL-2, TNF or IL-17 was determined using intracellular cytokine staining.

(A–C) The data represent the mean ± SEM from one experiment. Two-way analysis of

variance with Sidak's multiple comparison test was used to determine significance. *, p <

0.05; **, p < 0.01; ***, p < 0.001; ****, p < 0.0001.

Figure 6. Multivariate correlates of protection across all 24 CC strains

(A) A partial least squares regression (PLSR) model was performed to associate BCG-

induced changes in lung immune features with BCG-induced changes in lung CFU. Each

data point represents a vaccinated mouse that has the mean of the unvaccinated mice

from the same strain subtracted from it. The scores of the top two latent variables of the

model are shown, with the color of each data point representing the changed in lung CFU

induced by BCG vaccination. The percents of the variance in the lung features captured

by each latent variable are shown as percentages in the axes.

(B) The bar graph depicts the loadings of the first latent variable of the PLSR model in

(A). Positive loadings indicate immune features that increase from BCG vaccination in

samples that are on the positive x-axis in the scores plot, which generally indicates they

have worse protection. Negative loadings conversely indicate immune features that

generally increase from BCG vaccination in samples that are protected by vaccination.

(C) The heatmap illustrates all features selected to be included in the model, indicating

they were important for associating strains with the lung protection continuum. Colors are

representations of z-scored features, with a different color representing each 10%

quantile of the data. Columns are averages of the mice in the CC strains (n=5 for most

cases). Feature names are colored based on the dataset from which they come. In (B)

and (C), features from the ICS dataset indicate production of the cytokines listed and no

production of any measured cytokines not listed.

Figure 7. Identifying signatures of protection in subclasses of CC strains

(A), (C), (E) Scores plot (top) and heatmaps (bottom) of selected features for PLSDA

models. Each dot represents a single mouse. The two colors in the scores plots signify

the two classes being compared. An ellipse is drawn around the 95% confidence interval

for each class. Percent of variance of the immune feature data is shown on the axes.

Heatmaps show features that were most important for distinguishing the two classes.

Colors are representations of z-scored features. Columns are averages of the mice in the

CC strains, with vaccinated and unvaccinated mice being separated (n=5 for most cases).

Feature names are colored based on the dataset from which they come (blue=T cell

phenotyping, brown=lung homogenate cytokines, green=ICS, magenta=LEGENDplex).

NP = non-protected, P = protected, R = resistant, S = susceptible.

(B), (D), (F) PLSDA models performed as in (A), (C), and (E), except that features are the

changes induced by BCG vaccination as in Figure 6.

Figure 8. Th1/17 cells correlate with BCG-mediated protection in a subset of protected

CC mice.

(A) Cells were isolated from the lungs of CC mice at 4 weeks post infection and stimulated

for 5 hours with either aCD3, P300 peptide pool or media control. Representative flow

plots of IFNγ and IL-17 producing CD4 T cells from CC037 and C57BL/6 are shown. Data

is representative of two independent experiments.

(B) Frequencies of CD4 T cells that expressed both IFNγ and IL-17 was determined using

intracellular cytokine staining. Data is representative of two independent experiments.

One-way analysis of variance with Sidak's multiple comparison test was used to

determine significance. **, $p < 0.01$; ***, $p < 0.001$; ****, $p < 0.0001$.

(C) Representative flow plots of CD4 T cells expressing either T-bet or RORγt in CC037

or C57BL/6 mice. Data is representative of two independent experiments.

(D) Frequencies of CD4 T cells that expressed both T-bet or RORγt in CC037 or C57BL/6

mice. Data is representative of two independent experiments. One-way analysis of

variance with Sidak's multiple comparison test was used to determine significance. **, p

< 0.01; ***, $p < 0.001$.

(E) Proportion of CD4 T cells that express various combinations of IFNγ, TNF, IL-2 or IL-

17 in either PBS or BCG vaccinated mice at 4 weeks post Mtb infection. Plots are

generated using the SPICE platform (*28*).

(F) Volcano plots. Correlation between 60 immune features and the lung CFU determined

for CC037 and C57BL/6 mice (from Batch 1). Dotted line, $p = 0.05$. *, $q < 0.05$, corrected

with the Benjamini and Hochberg multiple comparison method (FDR = 0.05)

(G) Correlation of IFNγ$^+$IL-17$^+$ cells and bacterial burden in the lungs of either PBS or

BCG vaccinated CC037 mice at 4 weeks post Mtb infection. Colors represent BCG

vaccinated mice (red) or unvaccinated mice (blue). Spearman *r* and p value are shown.

SUPPLEMENTAL INFORMATION

Supplementary tables

Table S1. –Pathological descriptions of changes lung histology after BCG vaccination in

CC mice.

Table S2. A description of select CC strains that were repeated for in-depth immune

analysis.

Supplementary figure legends

Figure S1. Gating strategy for the analysis of the cell surface T cell phenotype.

(A) Lymphocytes were identified using FSC and SSC properties, after which viability dyes

were used to identify live cells. Singlet gates were used to exclude doublets, after which

T cells and non-T cells were identified using the lineage marker CD3.

(B) NKT and MAIT cells were identified from the CD3$^+$ gate using their antigen-loaded

CD1d and MR1 tetramers, respectively. CD4$^+$, CD8$^+$ and CD4$^-$CD8$^-$ T cells were also

identified from this population for further analysis.

(C) Expression of various markers were also examined within the CD3$^-$ population. As an

example, flow plot represents expression of T-bet within the CD3$^-$ population.

(D) Identification of memory T cells. CD44 was used to identify antigen-experienced T

cells. Resident memory T cells (T$_{RM}$) were identified based on CD103 expression. From

the CD103$^-$ fraction, CD127 and CD62L expression identified central memory (T$_{CM}$),

effector (T$_E$) and effector memory (T$_{EM}$). Subset identification of various T cell populations

(ex: Tbet⁺CD4⁺ T cells) as well as activation markers (ex: PD1, CD11a) were also

examined in CD4⁺, CD8⁺ and CD4⁻CD8⁻ T cell subsets.

(E) Naïve T cells identified within the CD44⁻ compartment using CD62L expression.

Figure S2. Frequencies of non-conventional T cells in CC mice. Frequencies of MR-1-

restricted MAIT cells and CD1d-restricted NKT cells were measured in the lungs of PBS

or BCG vaccinated CC mice at 4 weeks post Mtb infection. The data in the graphs

represent the mean ± SD of one experiment. Two-way analysis of variance with the

Benjamini and Hochberg multiple comparsion method. The FDR was set to 0.05 and the

numbers in the figures are the q value.

Figure S3. Gating strategy for the analysis of T cell function (intracellular cytokine

staining).

(A) Lymphocytes were identified as using FSC and SSC properties, after which viability

dyes were used to identify live cells. Singlet gates were used to exclude doublets, after

which T cells were identified using the lineage marker CD3.

(B) CD4, CD8 and CD4⁻CD8⁻ T cells were also identified from the CD3⁺ population for

further analysis. Cytokine expression (ex: IFNγ) were examined in all 3 T cell sub-

populations.

(C) Cytokine expression was also examined within the CD3⁻ population.

Figure S4. Multivariate model accuracies

(A) The accuracy of our protection model (Figure 6) was compared to that of two different

types of null model, where our measure of accuracy was mean squared error of prediction

of Δlog (lung CFU). Our PLSR model performed significantly better than both a null model

consisting of random features instead of our elastic-net chosen features and a null model

where the outcomes of Δlog (lung CFU) were scrambled. Our PLSR model and each of

the null models were run 50 times, and each dot represents one of these trials. Labeled

p-values are from one-way ANOVAs.

(B) Our models highlighting the immune features that distinguish pairs of

protection/susceptibility categories (Figure 7) were compared to two null models as in (A).

Each of our models, both the ones involving all mice and the protection models were

significantly better than both null models. Here, the accuracy metric is classification

accuracy. Labeled p-values are from one-way ANOVAs.

Figure S5. Examining the Polyfunctionality of CD4 T cells in select CC strains. Boolean

gating was used to identify various sub-populations amongst the CD4 T cells following

P300 stimulation in certain CC mice. An increase in various IFNγ$^+$IL-17$^+$ CD4 T cells (that

also produced TNF or IL-2) were detected in CC037, CC072, CC031 and CC001 but not

in C57BL/6. In contrast, a noticeable decrease in the frequency of IFNγ "single positive"

CD4 T cells was observed in all strains including C57BL/6. The data in the graphs

represent the mean ± SD of two experiment. One-way analysis of variance with Sidak's

multiple comparison test was used to determine significance. *, $p < 0.05$; **, $p < 0.01$; ***,

$p < 0.001$; ****, $p < 0.0001$.

910   References

912 1.   WHO, Global Tuberculosis Report 2021. (2021).

913 2.   G. A. Colditz *et al.*, Efficacy of BCG vaccine in the prevention of tuberculosis. Meta-
914   analysis of the published literature. *JAMA* **271**, 698-702 (1994).

915 3.   A. Roy *et al.*, Effect of BCG vaccination against Mycobacterium tuberculosis infection in
916   children: systematic review and meta-analysis. *BMJ* **349**, g4643 (2014).

917 4.   P. Mangtani *et al.*, Protection by BCG vaccine against tuberculosis: a systematic review
918   of randomized controlled trials. *Clin Infect Dis* **58**, 470-480 (2014).

919 5.   I. Abubakar *et al.*, Systematic review and meta-analysis of the current evidence on the
920   duration of protection by bacillus Calmette-Guerin vaccination against tuberculosis. *Health*
921   *Technol Assess* **17**, 1-372, v-vi (2013).

922 6.   P. E. Fine, Variation in protection by BCG: implications of and for heterologous immunity.
923   *Lancet* **346**, 1339-1345 (1995).

924 7.   J. L. Flynn, H. P. Gideon, J. T. Mattila, P. L. Lin, Immunology studies in non-human primate
925   models of tuberculosis. *Immunol. Rev.* **264**, 60-73 (2015).

926 8.   M. Ahmed *et al.*, Immune correlates of tuberculosis disease and risk translate across
927   species. *Sci Transl Med* **12**, (2020).

928 9.   C. Aagaard *et al.*, Immunization with Mycobacterium tuberculosis-Specific Antigens
929   Bypasses T Cell Differentiation from Prior Bacillus Calmette-Guerin Vaccination and
930   Improves Protection in Mice. *J Immunol* **205**, 2146-2155 (2020).

931 10.   J. S. Woodworth *et al.*, A Mycobacterium tuberculosis-specific subunit vaccine that
932   provides synergistic immunity upon co-administration with Bacillus Calmette-Guerin. *Nat*
933   *Commun* **12**, 6658 (2021).

934 11.   I. Kramnik, G. Beamer, Mouse models of human TB pathology: roles in the analysis of
935   necrosis and the development of host-directed therapies. *Semin Immunopathol* **38**, 221-
936   237 (2016).

937 12.   J. Nemeth *et al.*, Contained Mycobacterium tuberculosis infection induces concomitant
938   and heterologous protection. *PLoS Pathog* **16**, e1008655 (2020).

939 13.   K. E. Noll, M. T. Ferris, M. T. Heise, The Collaborative Cross: A Systems Genetics
940   Resource for Studying Host-Pathogen Interactions. *Cell Host Microbe* **25**, 484-498 (2019).

941 14.   G. A. Churchill *et al.*, The Collaborative Cross, a community resource for the genetic
942   analysis of complex traits. *Nat Genet* **36**, 1133-1137 (2004).

943 15.   C. M. Smith *et al.*, Tuberculosis Susceptibility and Vaccine Protection Are Independently
944   Controlled by Host Genotype. *MBio* **7**, (2016).

16.   C. M. Smith *et al.*, Host-pathogen genetic interactions underlie tuberculosis susceptibility
in genetically diverse mice. *Elife* **11**, (2022).

17.   J. Lee *et al.*, CD11cHi monocyte-derived macrophages are a major cellular compartment
infected by Mycobacterium tuberculosis. *PLoS Pathog* **16**, e1008621 (2020).

18.   B. Nandi, S. M. Behar, Regulation of neutrophils by interferon-gamma limits lung
inflammation during tuberculosis infection. *J Exp Med* **208**, 2251-2262 (2011).

19.   E. Sahiratmadja *et al.*, Plasma granulysin levels and cellular interferon-gamma production
correlate with curative host responses in tuberculosis, while plasma interferon-gamma
levels correlate with tuberculosis disease activity in adults. *Tuberculosis (Edinb)* **87**, 312-
321 (2007).

20.   S. Sakai *et al.*, Cutting edge: control of Mycobacterium tuberculosis infection by a subset
of lung parenchyma-homing CD4 T cells. *J Immunol* **192**, 2965-2969 (2014).

21.   S. Becattini *et al.*, Functional heterogeneity of human memory CD4(+) T cell clones primed
by pathogens or vaccines. *Science* **347**, 400-406 (2015).

22.   C. M. Smith *et al.*, Functionally Overlapping Variants Control Tuberculosis Susceptibility
in Collaborative Cross Mice. *mBio* **10**, (2019).

23.   C. S. Lindestam Arlehamn *et al.*, A Quantitative Analysis of Complexity of Human
Pathogen-Specific CD4 T Cell Responses in Healthy M. tuberculosis Infected South
Africans. *PLoS Pathog* **12**, e1005760 (2016).

24.   Y. R. Patankar *et al.*, Limited recognition of Mycobacterium tuberculosis-infected
macrophages by polyclonal CD4 and CD8 T cells from the lungs of infected mice. *Mucosal*
Immunol **13**, 140-148 (2020).

25.   C. S. Lindestam Arlehamn *et al.*, Memory T cells in latent Mycobacterium tuberculosis
infection are directed against three antigenic islands and largely contained in a
CXCR3+CCR6+ Th1 subset. *PLoS Pathog* **9**, e1003130 (2013).

26.   E. V. Acosta-Rodriguez *et al.*, Surface phenotype and antigenic specificity of human
interleukin 17-producing T helper memory cells. *Nat Immunol* **8**, 639-646 (2007).

27.   H. P. Gideon *et al.*, Multimodal profiling of lung granulomas in macaques reveals cellular
correlates of tuberculosis control. *Immunity* **55**, 827-846 e810 (2022).

28.   M. Roederer, J. L. Nozzi, M. C. Nason, SPICE: exploration and analysis of post-cytometric
complex multivariate datasets. *Cytometry A* **79**, 167-174 (2011).

Figure 1. BCG-induced protection in 24 CC strains and C57BL/6 mice.
(A) Lung and (D) spleen CFU of all 24 CC mice (n=120/group; top) or all C57BL/6 mice (n=20/group; bottom) at 4 weeks post infection. Mice were vaccinated with BCG or administered PBS and rested for 12 weeks prior to infection. Each point represents an individual mouse. Line, median. Student's t-test; ****, $p < 0.0001$. (B) Protection in the lung and (E) was assessed by comparing the CFU in BCG vs. sham vaccinated mice for each individual mouse strain (n=5/group) using the Benjamini-Hochberg procedure to determine the false discovery rate (FDR) (top). The ΔCFU of BCG vs. sham vaccinated was calculated (bottom). The open bar is the reference strain C57BL/6 strain (n=20/group). *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$; ****, $p < 0.0001$. (C) Correlation in the lung and (F) spleen between susceptibility and protection. The relative susceptibility (y-axis, defined by CFU in unvaccinated group, relative to the C57BL/6 reference strain) and the protection conferred by BCG (x-axis, defined by ΔCFU of vaccinated mice compared to unvaccinated controls, relative to the C57BL/6 reference strain). The graph is centered on C57BL/6 mice, leading to the different CC strains being placed in four quadrants based on their relative susceptibility and protection.

CC strain[a]	Batch[b]	MHC Haplotype[c]	ΔLung CFU$_1$ [d] reduction (logCFU)	ΔLung CFU$_2$ [d] reduction (logCFU)	ΔSpleen CFU$_1$ [e] reduction (logCFU)	ΔSpleen CFU$_2$ [e] reduction (logCFU)
CC001/Unc	4	b	0.94	1.45	2.79	1.45
CC002/Unc	3	b	0.83	n/a	0.46	n/a
CC003/Unc	3	g	0.07	-0.61	0.77	0.15
CC004/TauUnc	4	b	0.05	0.08	-0.16	-0.12
CC010/GeniUnc	1	b	0.53	n/a	0.92	n/a
CC011/Unc	3	g7	1.04	n/a	2.08	n/a
CC012/GeniUnc	4	wsb	0.79	n/a	0.89	n/a
CC023/GeniUnc	1	b	1.17	-0.09	1.47	0.32
CC024/GeniUnc	2	cast	0.15	0.24	0.38	0.62
CC025/GeniUnc	3	pwk	1.40	n/a	0.43	n/a
CC026/GeniUnc	2	cast	0.93	1.05	3.75	1.87
CC031/GeniUnc	3	b	1.22	1.35	1.26	1.60
CC032/GeniUnc	1	b	1.16	n/a	1.06	n/a
CC035/Unc	4	wsb	0.91	n/a	1.13	n/a
CC036/Unc	3	g7	0.62	n/a	1.84	n/a
CC037/TauUnc	1	b	1.37	1.84	1.07	1.05
CC040/TauUnc	2	b	-0.64	-0.28	-0.35	1.14
CC041/TauUnc	2	b	0.14	0.94	0.71	0.94
CC042/GeniUnc	1	b	1.65	0.96	1.00	1.32
CC044/Unc	4	b/g7	0.08	0.13	0.12	0.39
CC045/GeniUnc	4	z	0.24	n/a	0.13	n/a
CC059/TauUnc	2	b/a	1.39	n/a	1.17	n/a
CC061/GeniUnc	1	b	0.93	n/a	1.58	n/a
CC072/TauUnc	2	b	0.97	1.37	1.06	1.41

Table S1. A description of select CC strains that were repeated for in-depth immune analysis.
a A group of C57BL/6 mice were included in all batches as a reference strain. Data was normalized to internal C57BL/6 control as indicated in the methods.
b The first cohort of CC mice were analyzed in 4 separate batches as indicated.
c MHC haplotype of each CC mouse as inherited from CC founders.
d, e Thirteen of the 24 CC strains were analyzed a second time to validate the original results and perform more in-depth immune analysis. ΔLungCFU[d] and ΔSpleenCFU[e] for the first experiment and the repeat experiment are listed. Mice that were exacerbated by BCG vaccination are highlighted in red.

Strain[a]	BCG effect[b] (lung CFU)	Were differences observed?[c]	How did the vaccinated mice differ?[d]
CC003/TauUNC	Not protected	N	No differences. Both groups had mild lesions with perivascular lymphocytic cuffs and a few macrophage-rich lesions
CC041/TauUNC	Not protected	N	No differences. Both groups had mild lesions with perivascular lymphocytic cuffs and a few macrophage-rich lesions
CC024/GeniUNC	Not protected	N	No differences. Both groups had small hard-to-find lymphocytic lesions with perivascular and peribroncial lymphocytic cuffs
CC002/TauUNC	Not protected	Y	Ctrl had mild lesions with perivascular lymphocytic cuffs. BCG mice had larger moderate sized macrophage-rich lesions. Lymphocytes were similarly abundant.
CC004/TauUnc	Not protected	Y	Ctrl mice had mild to moderate lesions with lymphohistiocytic infiltrates. BCG mice have more lymphocytes with regions of dense perivascular/peribronchiolar lymphocytic cuffs.
CC036/GeniUnc	Not protected	Y	Ctrl mice have mild disease with lymphohistiocytic infiltrates. BCG mice also have lymphohistiocytic infiltrates with few very dense lymphocytic areas
CC040	Not protected	Y	Ctrl mice have moderate necrotizing lesions with few lymphocytes. BCG mice have mild disease - small & hard to find lesions that are lymphocyte rich without necrosis.
CC044/GeniUnc	Not protected	Y	Unusual. Ctrl mice have lymphohistiocytic infiltrates with dense lymphocytic cuffs. BCG mice have lymphohistiocytic infiltrates with dense lymphocytic cuffs and lymphoid aggregates/follicles in parenchyma.
CC061/GeniUNC	Protected	N	No differences. Unusual. Mild lesions. Both groups have dense lymphoid aggregates/follicles in parenchyma. Also, perivascular lymphocytic cuffs.
CC072/TauUNC	Protected	N	No differences. Mild disease. perivascular lymphocytic cuffs with a few macrophage-rich lesions. One mouse has a foreign-body granulomatous bronchitis/pneumonia with intralesional food/bedding/plant.
C57BL/6	Protected	Y	Ctrl mice had mild to moderate lymphohistiocytic lesions. BCG mice had with more, dense lymphocytic infiltrates
CC001/TauUnc	Protected	Y	Ctrl mice had moderate lesions with lymphohistiocytic infiltrates and equivocal necrosis. BCG mice had mild lesions with perivascular lymphocytic cuffs and a few macrophage-rich lesions without necrosis
CC023	Protected	Y	Ctrl mice had moderate to severe lesions with necrosis and minimal lymphocytes. BCG mice have moderate lesions with more lymphocytes and less necrosis.
CC025/TauUNC	Protected	Y	Ctrl mice have moderate disease with mild lymphocytic infiltrates and necrosis. BCG mice have abundant dense perivascular/peribronchiolar lymphocytic ifiltrates and lymphohistiocytic granulomas and less necrosis.
CC026	Protected	Y	Ctrl mice had moderate lesions with lymphohistiocytic infiltrates. BCG mice had mild lesions with lymphohistiocytic infiltrates and a few very dense lymphocytic areas
CC031/GeniUnc	Protected	Y	Ctrl mice have moderate disease with lymphohistiocytic infiltrates. BCG mice have abundant dense perivascular/peribronchiolar lymphocytes and lymphohistiocytic granulomas
CC032/GeniUNC	Protected	Y	Ctrl mice have mild disease with lymphohistiocytic infiltrates. BCG mice have more lymphocyte rich lesions that are small & hard to find
CC037/TauUNC	Protected	Y	Ctrl mice have moderate disease with lymphohistiocytic infiltrates. BCG mice have abundant dense perivascular/peribronchiolar lymphocytes and lymphohistiocytic granulomas
CC042	Protected	Y	Ctrl mice have mild to moderate disease with few lymphocytes and necrosis. BCG mice have mild disease with more lymphocytes, less necrosis
CC059/TauUNC	Protected	Y	Ctrl mice had mild disease with lymphohistiocytic infiltrates with small regions of necrosis. BCG mice have abundant dense perivascular/peribronchiolar lymphocytes and lymphohistiocytic granulomas without necrosis

Table S2. Pathological descriptions of changes lung histology after BCG vaccination in CC mice
a Nineteen of the 24 CC strains were available for analysis.
b The effect of BCG on the lung CFU is from Figure 1.
c Assessment by a pathologist of whether differences existed between sham and BCG vaccinated mice after Mtb infection (n=5/strain/condition).
d Summary of observed differences between the groups.

.

Figure 2. Lung histology of CC mice after BCG vaccination and Mtb challenge
Representative lung histology of three CC strains (CC025, CC037, CC072) that were protected
by BCG, after sham or BCG vaccination and Mtb challenge (Mtb and BCG+Mtb, respectively)
(left). Representative lung histology of three CC strains (CC003, CC004, CC024) that were
unprotected by BCG (right). Original magnification, 40X (inset, 400X). Scale bar, 500 μm
(inset, 50 μm).

Figure 3. BCG-protected CC mice have an altered lung microenvironment of CC mice after Mtb infection.

(A) IFNγ levels were measured in the lungs of either PBS or BCG vaccinated CC mice at 4 weeks post infection. A two-way ANOVA was performed using Sidak's correction for multiple comparisons. ***, p < 0.001; ****, p < 0.0001". (B) – (F) Pearson correlation between lung CFU and (B) IFNγ, IL-17, TNF, IL-1β, KC and IL-6 (C) LIX (D) RANTES (E) MIP-1β and (F) M-CSF, MIP-2, G-CSF, LIF, comparing PBS and BCG vaccinated mice at 4 weeks post infection in C57BL/6 mice (B6), non-protected (NP) or protected (P) CC mice.

Figure S1. Gating strategy for the analysis of the cell surface T cell phenotype.
(A) Lymphocytes were identified using FSC and SSC properties, after which viability dyes were used to identify live cells. Singlet gates were used to exclude doublets, after which T cells and non-T cells were identified using the lineage marker CD3.
(B) NKT and MAIT cells were identified from the CD3+ gate using their antigen-loaded CD1d and MR1 tetramers, respectively. CD4+, CD8+ and CD4−CD8− T cells were also identified from this population for further analysis.
(C) Expression of various markers were also examined within the CD3− population. As an example, flow plot represents expression of T-bet within the CD3− population.
(D) Identification of memory T cells. CD44 was used to identify antigen-experienced T cells. Resident memory T cells (T$_{RM}$) were identified based on CD103 expression. From the CD103− fraction, CD127 and CD62L expression identified central memory (T$_{CM}$), effector (T$_E$) and effector memory (T$_{EM}$). Subset identification of various T cell populations (ex: Tbet+CD4+ T cells) as well as activation markers (ex: PD1, CD11a) were also examined in CD4+, CD8+ and CD4−CD8− T cell subsets.
(E) Naïve T cells identified within the CD44− compartment using CD62L expression.

Figure 4. Frequency of various T cell populations in protected and unprotected CC mice following Mtb infection.

(A) Proportion of CD3+ cell, CD3+CD4+ and CD3+CD8+ T cells in the lungs of PBS and BCG vaccinated CC mice at 4 weeks post infection.

(B) Proportion of memory CD4 and CD8 T cells in the lungs of either PBS or BCG vaccinated CC mice at 4 weeks post infection, as defined by CD44+ and various combinations of CD62L and CD127 expression; CD62L+ CD127+ (central memory), CD62L− CD127− (effector), CD62L− CD127+ (effector memory).

(C) Proportion of resident memory CD4 and CD8 T cells in the lungs of either PBS or BCG vaccinated CC mice at 4 weeks post infection, as defined by CD44+ CD103+ CD69+.

(D) Proportion of Th1, Treg and Th17 cells in the lungs of either PBS or BCG vaccinated CC mice at 4 weeks post infection, as defined by CD44+ and Tbet+ (Th1), Foxp3+ (Treg) or RoRγt (Th17) expression.

(E) Proportion of CD4 and CD8 T cells expressing either CXCR3 or CX$_3$CR1 in the lungs of either PBS or BCG vaccinated CC mice at 4 weeks post infection.

(F) PCA of the T cell phenotyping data. Each point represents the average of usually 5 mice within a given mouse strain that are either BCG vaccinated or unvaccinated. Open symbols, unvaccinated; closed symbols, BCG vaccinated. Each color represents a different mouse strain.

(A–E) The data represent the mean ± SEM from one experiment. Two-way analysis of variance with the Benjamini and Hochberg multiple comparison method. The FDR was set to 0.05 and the numbers in the figures are the q value.

Figure S2. Frequencies of non-conventional T cells in CC mice. Frequencies of MR-1-restricted MAIT cells were measured in the lungs of PBS or BCG vaccinated CC mice at 4 weeks post Mtb infection. The data in the graphs represent the mean ± SD of one experiment. Two-way analysis of variance with the Benjamini and Hochberg multiple comparsion method. The FDR was set to 0.05 and the numbers in the figures are the q value.

Figure S3. Gating strategy for the analysis of T cell function (intracellular cytokine staining).

(A) Lymphocytes were identified as using FSC and SSC properties, after which viability dyes were used to identify live cells. Singlet gates were used to exclude doublets, after which T cells were identified using the lineage marker CD3.

(B) CD4, CD8 and CD4-CD8- T cells were also identified from the CD3+ population for further analysis. Cytokine expression (ex: IFNγ) were examined in all 3 T cell sub-populations.

(C) Cytokine expression was also examined within the CD3- population.

Figure 5. Cytokine response in protected and unprotected CC strains following Mtb infection.
(A) Cells were isolated from the lungs of CC mice at 4 weeks post infection and stimulated for 24 hours with the P300 peptide pool. IFNγ, IL-2, TNF and IL-17a were measured in the supernatants. (B) and (C) Cells from the lungs of CC mice at 4 weeks post infection were stimulated for 5 hours with the P300 peptide pool. Frequencies of CD4 or CD8 T cells that expressed IFNγ, IL-2, TNF or IL-17 was determined using intracellular cytokine staining. (A–C) The data represent the mean ± SEM from one experiment. Two-way analysis of variance with Sidak's multiple comparison test was used to determine significance. *, p < 0.05; **, p < 0.01; ***, p < 0.001; ****, p < 0.0001.

Figure 6. Multivariate correlates of protection across all 24 CC strains

(A) A partial least squares regression (PLSR) model was performed to associate BCG-induced changes in lung immune features with BCG-induced changes in lung CFU. Each data point represents a vaccinated mouse that has the mean of the unvaccinated mice from the same strain subtracted from it. The scores of the top two latent variables of the model are shown, with the color of each data point representing the changed in lung CFU induced by BCG vaccination. The percents of the variance in the lung features captured by each latent variable are shown as percentages in the axes. (B) The bar graph depicts the loadings of the first latent variable of the PLSR model in (A). Positive loadings indicate immune features that increase from BCG vaccination in samples that are on the positive x-axis in the scores plot, which generally indicates they have worse protection. Negative loadings conversely indicate immune features that generally increase from BCG vaccination in samples that are protected by vaccination. (C) The heatmap illustrates all features selected to be included in the model, indicating they were important for associating strains with the lung protection continuum. Colors are representations of z-scored features, with a different color representing each 10% quantile of the data. Columns are averages of the mice in the CC strains (n=5 for most cases). Feature names are colored based on the dataset from which they come. In (B) and (C), features from the ICS dataset indicate production of the cytokines listed and no production of any measured cytokines not listed.

Resistant CC strains: Protected vs. not protected

Susceptible CC strains: Protected vs. not protected

Good immunity (protecte resistant) vs. poor immunity (not protected susceptible)

All mice models

Protection models (vacc-unvacc)

Figure 7. Identifying signatures of protection in subclasses of CC strains

(A), (C), (E) Scores plot (top) and heatmaps (bottom) of selected features for PLSDA models. Each dot represents a single mouse. The two colors in the scores plots signify the two classes being compared. An ellipse is drawn around the 95% confidence interval for each class. Percent of variance of the immune feature data is shown on the axes. Heatmaps show features that were most important for distinguishing the two classes. Colors are representations of z-scored features. Columns are averages of the mice in the CC strains, with vaccinated and unvaccinated mice being separated (n=5 for most cases). Feature names are colored based on the dataset from which they come (blue=T cell phenotyping, brown=lung homogenate cytokines, green=ICS, magenta=LEGENDplex). NP = non-protected, P = protected, R = resistant, S = susceptible. (B), (D), (F) PLSDA models performed as in (A), (C), and (E), except that features are the changes induced by BCG vaccination as in Figure 6.

Figure S4. Multivariate model accuracies

(A) The accuracy of our protection model (Figure 6) was compared to that of two different types of null model, where our measure of accuracy was mean squared error of prediction of Δlog (lung CFU). Our PLSR model performed significantly better than both a null model consisting of random features instead of our elastic-net chosen features and a null model where the outcomes of Δlog (lung CFU) were scrambled. Our PLSR model and each of the null models were run 50 times, and each dot represents one of these trials. Labeled p-values are from one-way ANOVAs.

(B) Our models highlighting the immune features that distinguish pairs of protection/susceptibility categories (Figure 7) were compared to two null models as in (A). Each of our models, both the ones involving all mice and the protection models were significantly better than both null models. Here, the accuracy metric is classification accuracy. Labeled p-values are from one-way ANOVAs.

Figure 8

Figure 8. Th1/17 cells correlate with BCG-mediated protection in a subset of protected CC mice.

(A) Cells were isolated from the lungs of CC mice at 4 weeks post infection and stimulated for 5 hours with either aCD3, P300 peptide pool or media control. Representative flow plots of IFNγ and IL-17 producing CD4 T cells from CC037 and C57BL/6 are shown. Data is representative of two independent experiments.

(B) Frequencies of CD4 T cells that expressed both IFNγ and IL-17 was determined using intracellular cytokine staining. Data is representative of two independent experiments. One-way analysis of variance with Sidak's multiple comparison test was used to determine significance. **, $p < 0.01$; ***, $p < 0.001$; ****, $p < 0.0001$.

(C) Representative flow plots of CD4 T cells expressing either T-bet or RORγt in CC037 or C57BL/6 mice. Data is representative of two independent experiments.

(D) Frequencies of CD4 T cells that expressed both T-bet or RORγt in CC037 or C57BL/6 mice. Data is representative of two independent experiments. One-way analysis of variance with Sidak's multiple comparison test was used to determine significance. **, $p < 0.01$; ***, $p < 0.001$.

(E) Proportion of CD4 T cells that express various combinations of IFNγ, TNF, IL-2 or IL-17 in either PBS or BCG vaccinated mice at 4 weeks post Mtb infection. Plots are generated using the SPICE platform (*28*).

(F) Volcano plots. Correlation between 60 immune features and the lung CFU determined for CC037 and C57BL/6 mice (from Batch 1). Dotted line, $p = 0.05$. *, $q < 0.05$, corrected with the Benjamini and Hochberg multiple comparison method (FDR = 0.05)

(G) Correlation of IFNγ+IL-17+ cells and bacterial burden in the lungs of either PBS or BCG vaccinated CC037 mice at 4 weeks post Mtb infection. Colors represent BCG vaccinated mice (red) or unvaccinated mice (blue). Spearman *r* and p value are shown.

Figure S5. Examining the Polyfunctionality of CD4 T cells in select CC strains. Boolean gating was used to identify various sub-populations amongst the CD4 T cells following P300 stimulation in certain CC mice. An increase in various IFNγ+IL-17+ CD4 T cells (that also produced TNF or IL-2) were detected in CC037, CC072, CC031 and CC001 but not in C57BL/6. In contrast, a noticeable decrease in the frequency of IFNγ "single positive" CD4 T cells was observed in all strains including C57BL/6. The data in the graphs represent the mean ± SD of two experiment. One-way analysis of variance with Sidak's multiple comparison test was used to determine significance. *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$; ****, $p < 0.0001$.