Host genetic background is a barrier to vaccine-induced protection against infection

Rocky Lai[1], Diana Gong[2], Travis Williams[1], Abiola F Ogunsola[1], Kelly Cavallo[1], Cecilia Lindestam

Arlehamn[3], Sarah Acolatse[2], Gillian L. Beamer[4], Martin Ferris[5], Christopher M. Sassetti[1], Doug

Lauffenburger[2], Samuel M. Behar[1]

1. Department of Microbiology and Physiological Systems, University of Massachusetts Medical

School, Worcester, Massachusetts, USA.

2. Department of Biological Engineering, Massachusetts Institute of Technology, Cambridge,

MA, USA

3. Center for infectious disease and vaccine research, La Jolla Institute for Immunology., La Jolla,

CA, USA

4. Texas Biomedical Research Institute, San Antonio, TX, USA

5. Department of Genetics, University of North Carolina at Chapel Hill, Chapel Hill, United States

21   *Correspondence:

22   Samuel M. Behar

23   E-mail address: samuel.behar@umassmed.edu

Summary

The heterogeneity of immune responses observed in humans is difficult to model in standard inbred laboratory mice. We used Collaborative Cross (CC) recombinant inbred strains to capture the diversity of outbred populations and better understand how host variation affects BCG-induced immunity against *Mycobacterium tuberculosis*. Using 24 CC strains, we show host genetics has a major role in determining whether BCG elicits protective immunity to *M. tuberculosis*. Only 54% of BCG-vaccinated CC strains had significantly reduced lung bacillary burdens after infection. Furthermore, BCG efficacy is dissociable from inherent susceptibility to TB. Like people, CC mice have diverse immunological responses after BCG vaccination and *M. tuberculosis* challenge. We define immunological signatures associated with BCG-induced protection other than canonical Th1 immunity. Like interferon-γ, lung homogenate levels of IL-17 correlated with disease. However, the presence of Th1/17 CD4 T cells were an immune feature that correlated with protection. CC mice can be used to define correlates of protection and to identify vaccine strategies that protect a larger fraction of genetically diverse individuals instead of optimizing protection for a single genotype.

Introduction

The introduction of Bacillus Calmette Guerin (BCG) as a vaccine in the last century has contributed significantly to our ability to control and contain tuberculosis (TB), which has been a threat to global health for millennia. BCG is the only vaccine approved for preventing tuberculosis and over 4 billion doses have been administered in over 150 countries (1). Although BCG prevents extrapulmonary TB in infants, it is less effective in averting pulmonary disease in adolescents and adults (2-5). Complicating the use of BCG is its variable efficacy in different regions of the world (6). While the development of more effective TB vaccines is recognized by the WHO as an important yet unmet medical need, the lack of identifiable correlates of protective immunity to TB is an impediment towards this goal.

Why is identifying immune correlates of protection following BCG difficult? TB is primarily a human disease that is transmitted person-to-person by aerosolized *Mycobacterium tuberculosis* (Mtb). No animal model recapitulates all the features of human TB, although non-human primates come close (7, 8). Although imperfect, the mouse model has facilitated mechanistic studies that defined many of the critical immunological pathways required for host resistance, permitted in vivo antibiotic testing, and accelerated preclinical vaccine development. As inbred mouse strains are genetically identical, a strength of the mouse model lies in its "infinite" reproducibility. For experimental immunology, C57BL/6 mice are the preferred inbred strain as most genetically modified mice are produced using its genetic background. It is also the most frequently used mouse strain for TB research. However, several features of C57BL/6 mice suggest it is an outlier strain for TB research. Immunologically, it only expresses a single class II MHC molecule (i.e., I-A^b). Pathologically, it develops atypical pathology (i.e., pneumonitis) following Mtb infection. Finally, and perhaps most importantly, it is among the most resistant of all inbred strains to Mtb infection, which makes it difficult to identify conditions that make C57BL/6 mice more resistant to disease. Investigators have addressed these limitations by using F1 hybrids and susceptible strains to increase MHC and genetic diversity and identify strategies to increase host resistance

(9-12). However, even among different inbred strains, the genetic diversity is limited (13). This is

a problem not only for TB research, but for many areas of biomedical research.

The mouse genetics community responded to these limitations of inbred mice by

developing a mouse resource that captures the considerable genetic diversity of the *Mus* species.

Starting with eight founder mouse strains, including representatives from all three *Mus musculus*

subspecies and three wild-derived strains, an eight-way funnel breeding scheme led to random

assortment of the founder genomes among the progeny (14). The progenies were used to derive

recombinant inbred mouse strains, which became known as Collaborative Cross (CC) strains.

Each CC strain is a unique genetic representation (i.e., genotype) of the eight founder strains.

Importantly, as they are inbred, they are a renewable resource, and an unlimited number of

experiments can be performed with the same genotype, which is important for identifying

experimental correlates and defining mechanisms.

We previously reported that the CC strains have a large variation in primary susceptibility

to TB (15, 16). Here we leverage the CC model to understand the how host genetics affects

vaccine-induced protection against Mtb. We find that the genetic background of the host has a

major effect on whether BCG induces protection. There was a poor correlation between inherent

host susceptibility and BCG-mediated protection, indicating that vaccines can mediate protection

even in individuals with a genetic propensity to TB. Interestingly, the T cell response following

BCG vaccination and Mtb challenge was diverse, with some immune features universally

correlating with outcome of vaccination while others only were important in subsets of CC strains.

We propose that the use of CC mice as a preclinical model can facilitate the development of

vaccines that protect a diverse range of genotypes against TB.

Results.

BCG vaccination protects only a subset of CC strains against TB.

To establish how the host genetic background affects vaccine mediated protection against Mtb, 24 CC lines (Table 1) were vaccinated subcutaneously with BCG and rested for 12 weeks. Then, the mice were infected with lose dose aerosolized H37Rv.YFP (17). Four weeks after infection, the reduction in lung and spleen CFU were determined as a measurement of protection. Mice were initially tested in four batches, with each batch including six different CC strains and C57BL/6 mice as a reference strain.

The aggregated lung CFU from the unvaccinated and BCG-vaccinated CC strains (Fig.1A, top) and C57BL/6 mice (Fig.1A, bottom) shows that as a population, CC mice are protected by BCG similarly as C57BL/6 mice. However, the variance in the lung CFU of CC mice is greater. When the data is segregated by individual CC strains, we can see that the variance reflects different levels of protection conferred by BCG vaccination in each CC strain (Fig.1B, top). The CC strains were classified as being protected by BCG if there was a statistically significant difference between unvaccinated and vaccinated mice. The $\Delta\log_{10}$ lung CFU was calculated for each CC strain, and they were ranked from most protected (CC037, $\Delta\log_{10}$ CFU = 1.6) to least protected (CC040, $\Delta\log_{10}$ CFU = –0.6) (Fig.1B, bottom). BCG vaccination exacerbated subsequent Mtb infection in the lung for two CC strains (CC003 and CC040), similar to what had previously been observed in the NZO founder strain (15). Importantly, BCG only led to significant reductions in the lung Mtb burden in 13 of the 24 CC strains.

We next asked whether the protective effect of BCG vaccination correlated with the inherent ability (i.e., in the unvaccinated state) of each CC strain to control Mtb replication in their lungs. We categorized the 24 CC strains relative to C57BL/6 mice, which we used as a reference strain. The lung CFU of unvaccinated mice defined the inherent susceptibility or resistance to Mtb (susceptibility, y-axis). Next, we considered whether each CC strain was protected or not based on the $\Delta\log_{10}$ lung CFU (relative BCG-induced protection, x-axis). The plot was centered on

C57BL/6 mice, which represented a median strain in terms of lung susceptibility and protection

and divided the 24 CC strains into four different groups, represented by four quadrants (Fig.1C).

Based on lung CFU, there is a poor correlation between intrinsic susceptibility of the mouse strain

and protection conferred by BCG at this time point (r=0.104, p=0.62).

The same analysis was performed using the spleen bacterial burden from infected mice.

As was observed in the lung, the CC population was similarly protected as C57BL/6 (Fig.1D). The

range of spleen protection extended from most protected (CC026, Δlog_{10} CFU = 2.56) to least

protected (CC045, Δlog_{10} CFU = 0.13) (Fig.1E). BCG significantly protected against systemic

infection in 17 of the 24 CC strains (Fig.1E). There was also no correlation between susceptibility

and protection in the spleen (r=0.089, p=0.67). Interestingly, while C57BL/6 mice were a "median"

strain in the lung analysis, C57BL/6 mice had among the highest spleen CFU, suggesting they

disseminated more, but were among the most protected mouse strain following BCG vaccination

(Fig.1F). Among the CC strains, protection in the lung and the spleen were moderately correlated

(r=0.513, p=0.0087).

In follow-up studies, 13 of the 24 CC strains were retested to confirm the initial phenotype,

with emphasis placed on strains that represented varying levels of protection and susceptibility

(Table 1). Thus, using 24 CC strains, with diverse genotypes, we find that less than 50% of the

strains were protected by BCG, and that protection did not correlate with the intrinsic susceptibility

of the mouse to Mtb infection. Finally, the lack of correlation between the reduction in lung versus

spleen highlights the importance of tissue-specific immunity and the problems inherent in

protecting the lung against infection. We next assessed how BCG vaccination altered the

histological appearance of TB lesions in the lung.

BCG modifies the lung lesions caused by Mtb in a strain-specific manner.

We assessed how BCG vaccination modified lung lesion development after Mtb infection

in 19 different CC strains plus C57BL/6 mice. A board-certified veterinarian pathologist (G.B.),

blinded to vaccination status of each strain, examined the lung for necrosis, neutrophilic and

lymphocytic infiltrates, lesion size and number. Within each pair, the vaccinated group was

correctly identified in 15 of the 20 strains analyzed based on reduced severity of the

histopathology. In some CC strains that were protected by BCG, increased lymphocytic infiltration,

and reduced necrosis were frequently observed after vaccination (Table 2). CC025 (Fig. 2, top

left) and CC037 (Fig. 2, middle left) were both susceptible to primary Mtb infection, and the

infected lung showed innate immune cells and tissue necrosis. BCG vaccination protects these

mice by promoting lymphocyte recruitment and in CC025 BCG vaccination reduced necrosis.

However, a reduction in lung CFU associated with BCG vaccination was not always associated

with an improved histological appearance of the lung. Despite a significant reduction in lung CFU

after BCG, no discernable histopathological differences were observed between unvaccinated

and BCG vaccinated CC072 mice, and both groups had non-necrotizing lymphocyte-rich

granulomas (Fig. 2, bottom left).

BCG did not significantly reduce lung CFU in eight of 20 CC strains. Two of the non-

protected CC strains, CC003 (Fig. 2, top right) and CC024 (Fig. 2, middle right) had no discernable

histological differences in granuloma structure, content, or severity between control and BCG

vaccination when evaluated blindly. In these 2 strains, lesions were generally small to medium-

sized, non-necrotizing, with abundant perivascular and peribronchiolar lymphocytes. Given the

intermediate susceptibility of the CC003 strain, and BCG's inability to alter lung CFU or change

the lesions, the data suggests that BCG does not change CC003 innate or adaptive immune

responses to Mtb challenge. Likewise, CC024 is unprotected by BCG vaccination, nor are lung

lesions altered, suggesting that neither the innate nor adaptive responses of this resistant strain

to primary Mtb infection is further enhanced by BCG vaccination. The natural resistance of CC024

strain was not further improved. In contrast, CC004 (Fig. 2, bottom right) did have discernable

histopathological changes when controls and BCG vaccinees were evaluated blinded. The

differences attributable to BCG vaccination were the greater lymphocytic infiltration with denser

perivascular and peribronchiolar lymphocytes. The altered lesions detected after vaccination

suggests that the immunity induced by BCG in CC004 mice fails to control bacilli. Together, these

data show that the host genetic background impacts the granuloma content, structure, and quality

of the immune response to BCG and Mtb challenge in a manner that is not wholly predictable.

BCG-protected CC strains have an altered lung microenvironment after Mtb infection

The Mtb-infected lung is a complex environment of resident and recruited cells that

produce inflammatory mediators and counter-regulatory (e.g., anti-inflammatory) signals. We

sought to determine whether BCG vaccination primes the immune response to generate a

different lung microenvironment upon Mtb challenge, perhaps one that is less conducive to

bacillary replication and transmission. Cytokine levels in lung homogenates from PBS control and

BCG-vaccinated C57BL/6 or CC mice were measured 4 weeks post infection using the MD31

Discovery Assay Array.

Given the critical role of IFNγ in Mtb control, its levels in control and BCG-vaccinated CC

mice were compared. In C57BL/6 mice and many of the protected CC strains, including CC037,

CC025, CC059, CC031, and CC011, IFNγ concentrations in BCG-vaccinated mice were reduced

compared to unvaccinated mice (Fig.3A). However, this pattern did not hold for all protected

strains. No changes in IFNγ levels were detected the CC032, CC042, CC072, CC026 and CC001

strains, despite being protected by BCG. Thus, when considered alone, neither the absolute level

of IFNγ nor its changes after vaccination could be used to predict protection.

We next asked if there was a correlation between the different cytokines in the lung

homogenate and lung CFU, in either unvaccinated or BCG-vaccinated mice. We asked if this

correlation differed among protected and unprotected CC strains, and C57BL/6 mice. (Fig.3B, top

left). There was a strong correlation between CFU and lung IFNγ levels (R=0.71, p=0.0004), in

unvaccinated C57BL/6 mice but not in BCG-vaccinated C57BL/6 mice. Thus, IFNγ correlated with

bacterial burden but not with protection, as observed in prior studies (18, 19). The CC strains

protected by BCG had a similar pattern. However, for unprotected CC strains, a similar correlation

between IFNγ and CFU was identified in unvaccinated, and BCG vaccinated mice, which we

suggest is evidence that these strains were unprotected by BCG vaccination. This pattern held

for T cell cytokines (IFNγ and IL-17), mediators produced by myeloid cells (IL-1β, IL-6, KC, and

MIP-1α), and TNF (Fig.3B and not shown). These data suggest that protected and unprotected

CC strains respond differently to BCG, and that the protected lung environment differs

immunologically from an unprotected environment.

Other cytokines were of interest. LIX (CXCL2) was negatively associated with lung CFU

in unvaccinated but not in BCG-vaccinated C57BL/6 mice (–0.67, p=0.0013). This correlation was

only observed in C57BL/6 mice, and LIX levels poorly correlated with lung CFU in CC strains

(Fig.3C). RANTES strongly correlated with lung CFU in C57BL/6 mice, independent of vaccination

status, but was only weakly correlated in CC strains (Fig.3D). Like RANTES, MIP-1β was also

correlated with lung CFU, independently of vaccination status (Fig.3E). This was most convincing

in unprotected CC mice (R=0.71-0.78), with moderate correlations in vaccinated C57BL/6 and

protected CC strains (R=0.46-0.49).

Four cytokines (G-CSF, M-CSF, MIP-2 and LIF) were strongly correlated with lung CFU

in unvaccinated CC strains, regardless of protection status (R=0.57-0.8, Fig.3F). Again, these

correlations were lost in the BCG-vaccinated protected CC strains. None of these cytokines were

made in significant amounts by C57BL/6 mice or were poorly correlated with CFU. These data

show how differences in the host genetic background, result in varied lung histology and

cytokines, which dramatically alter the lung inflammatory environment during Mtb infection.

Host genetic background and not BCG vaccination determines T cell subset distribution

in the lung after Mtb infection.

We hypothesized that changes in the frequency of T cell subsets would correlate with

protection induced by BCG vaccination. An 18-parameter flow cytometry panel was used to

identify and enumerate different immune subsets (Figure S1). As T cells are important in

protection against TB, we determined the frequency and relative number of different T cell subsets

modulated by BCG vaccination. Not all parameters were obtained for all 24 CC strains (see

methods). We quantified CD4 and CD8 T cells, MAIT cells and NKT cells (Fig.4A, and Figure S2).

Memory and effector T cell populations were defined as central memory (T_{CM};

CD44$^+$CD62L$^+$CD127$^+$), effector memory (T_{EM}; CD44$^+$CD62L$^-$CD127$^+$), effector (T_{Eff};

CD44$^+$CD62L$^-$CD127$^-$), and resident memory (T_{RM}; CD44$^+$CD103$^+$CD69$^+$) (Fig.4B, 4C). CD4 T

cells subsets were defined in all CC strains based on their expression of the transcription factors

Foxp3, Eomes, Tbet and RORγt to identify Treg, Th1, Th2, and Th17 cells, respectively (Fig.4D).

For these T cell subsets, we observed tremendous strain-to-strain variation. With one or two

exceptions, BCG vaccination did not modulate the distribution of T cells subsets nor was there

any correlation between the relative frequency of these T cell subsets and the outcome of BCG

vaccination. Thus, the frequency of different T cells subsets in the lungs of CC mice after Mtb

infection was largely determined by each strain's genetic background.

Expression of CXCR3 and CX$_3$CR1, which have been used to discriminate between T

cells in the lung parenchymal vs. the vasculature, was analyzed in a limited number of CC strains

(20). CXCR3 also marks Th1 cells, and in conjunction with CCR6, can differentiate between Th1

and Th17 cells, respectively (21). Like other markers, we observed variation in CXCR3 expression

between different CC strains, but little or no variation that correlated with vaccination status or

susceptibility. In addition to allelic variation, CC strains also carry private mutations. For example,

we previously reported that CC042 mice lacks cell surface CD11a expression due to a mutation

in the *Itgal* gene on Chromosome 7 (22). Here, we report that CC003 mice don't express CXCR3

(Fig.4E). This phenotype is associated with the CXCR3 allele of the PWK founder. Finally, a

reduction in CX3CR1$^+$ CD4 and CD8 T cells was observed in several of the protected CC strains

but not in unprotected strains (Fig.4E). This pattern mirrored the reduction in IFNγ levels observed

in some protected CC mice (Fig.3), suggesting that these changes reflect Th1 modulation.

Finally, we used principal components analysis to holistically visualize the variation in the

T cell phenotyping data. What is striking is the small effect of BCG vaccination in altering T cell

phenotypes, in comparison to large effect of genetic strain differences. These data all indicate

that genetics differences between the CC strains determine the differences among T cell subsets

recruited to the lung, and not BCG vaccination or Mtb infection. Therefore, we next wished to

determine whether there were functional differences among the T cell subsets in protected vs.

nonprotected CC strains.

The patterns of cytokine production differ in CC strains after BCG vaccination and Mtb

infection.

As the cytokines produced by T cells, including IFNγ, TNF, and IL-21, among others, are

important in control of Mtb infection, we hypothesized that changes in T cell function would be

associated with BCG-induced protection. To measure function, we stimulated lung mononuclear

cells (MNC) from each mouse with anti-CD3 or the P300 megapool, which contains 300 peptides

representing epitopes from 70 Mtb proteins that are frequently recognized by human CD4 T cells

and are also recognized by murine CD4 and CD8 T cells (23, 24). Twenty-six cytokines and

chemokines in the culture supernatants were measured using the Legendplex platform 24 hours

after stimulation. There was little difference in the ability of lung MNC from mock or BCG

vaccinated Mtb-infected C57BL/6 mice to secrete different cytokines or chemokines following

P300 stimulation. In contrast, we detected differential production of IFNγ, TNF, IL-2 or IL-17, all

cytokines that have been associated with immunity to Mtb infection, among mock vs. BCG

vaccinated CC strains (Fig.5A). Interestingly, the MNC from BCG vaccinated CC037, CC031,

CC72 and CC001 strains all produced more IL-17 than their respective controls. In contrast, none

of the unprotected strains produced significant amounts of IL-17, except for CC023, and its lung

MNC produced less IL-17 after BCG vaccination.

In parallel, a 14-parameter flow cytometry panel measured intracellular levels of IFNγ, IL-

2, IL-17, TNF, IL10, and IL-4, and surface expression of CD107 in both CD4 and CD8 T cells

(Figure S3). BCG vaccination had a minimal effect on CD8 T cell cytokine responses to P300

stimulation in both C57BL/6 mice as well as most CC lines examined (Fig.5B). However, BCG

clearly modified the CD4 T cell cytokine responses in some CC strains (Fig.5C). Patterns of

cytokine production were observed in BCG vaccinated CC mice that were not observed in

C57BL/6 mice. For example, while BCG vaccination did not alter the frequency of IFNγ$^+$ or TNF$^+$

CD4 T cells in CC037 mice, it did increase IL-2 and IL-17 responses. BCG vaccination of CC031

mice increased the frequency of CD4 T cells producing IFNγ, IL-2 and TNF but not IL-17. Finally,

CD4 T cells from BCG vaccinated had CC072 had increased IFNγ and IL-2 but not TNF or IL-17.

Interestingly, no discernable differences in cytokine responses were observed in most of the non-

protected CC mice, except for a decrease in IFNγ responses in CC023. Thus, the cytokine

responses in both control and BCG vaccinated CC mice differ greatly from classic C57BL/6

model. Increased cytokine responses after T cell stimulation were more likely to be detected in

protected CC strains. Moreover, BCG vaccination led to quantitative and qualitative changes in

the type of T cell responses that persisted after Mtb challenge.

Correlates of Protection.

We then turned to multivariate approaches to discern how BCG-induced immune changes

in the lung mechanistically lead to reductions in lung CFU in CC strains. To do so, the data from

the lung homogenate cytokines, intracellular cytokine staining, T cell phenotyping, and

LEGENDplex were combined, and for each vaccinated mouse the transformed feature data was

calculated as: [(vaccinated CC subject– mean(unvaccinated CC strain]. Then, a partial least

squares regression (PLSR) model with elastic net feature selection was created to identify

features that changed with BCG vaccination in a way that maximally covaried with BCG-induced

changes in lung CFU. The model performed significantly better than null models, and the first two

latent variables in PLSR model account for 24% and 8%, respectively, of the variability in the lung

features (Figure 6A, Figure S4A). The elastic net-selected features in the model included features

from all four datasets (Fig.6B, 6C). By plotting the scaled values of the elastic net-selected

features for all CC strains, ordered from most protected to least protected, what is most apparent

is that the protected strains all had BCG-induced reductions in several inflammatory cytokines

such as MIG, CCL3 and CCL4 (i.e., MIP-1α and -1β), IL-6, RANTES, and IL-17. The decrease in

these inflammatory mediators could be a sign of resolved infection, in contrast to the unprotected

strains which had more inflammatory cytokines which could signify unimpeded infection.

Protected strains also had a greater vaccination-induced increase in activated CD4 T cells.

Contrarily, protected strains had decreased activated $CD4^-CD8^-$ T cells producing T-bet.

Importantly, IL-2, unlike several other cytokines, was increased in protected strains. Thus, BCG

vaccination leads to qualitative and quantitative changes in the T cell response and in the immune

environment of the lung after Mtb infection in CC genotypes that are associated with protection.

Immune responses in protected and unprotected CC strains differ following Mtb infection

While the PLSR model highlights potential correlates of lung protection across all

genotypes, it may miss correlates that only apply to certain subsets of CC strains. We

hypothesized that naturally resistant and susceptible strains may be protected in different ways.

To test this, we created separate partial least squares discriminant analysis (PLSDA) models for

resistant and susceptible CC strains and examined the features that best separated the BCG-

protected and non-protected strains in each of these categories. (Figure 1C). The data used for

the models was processed the same way as in the PLSR models, although we also looked at the

vaccinated and unvaccinated mice separately in addition to the vaccine-induced differences. Our

first analysis compared resistant CC strains that were protected by BCG (n=4) or not (n=8), based

on lung CFU. As described earlier (Figure 4), the unvaccinated and vaccinated mice of each CC

strain were very similar and clustered together when the elastic net-selected features of the model

were hierarchically clustered, indicating that the genetic background was an important

determinant of outcome (Figure 7A). The protected and unprotected mice were both split into

multiple clusters, showing that there were multiple distinct types of immune responses that were

associated with protective immunity in resistant mice. The model could predict protected vs non-

protected CC strains accurately and thus there is a multivariate signature of these categories

(Figure S4B). However, there were no clear individual features distinguishing the categories. In

contrast, the vaccine-induced model comparing protected and non-protected resistant strains did

yield more univariate trends in addition to the strong multivariate signature. Namely, both CD4

and CD8 T cells making IL-2 are increased in protected strains, while several lung homogenate

cytokines as well as double negative T cells expressing CD107a and producing IL-17 decreased

in protected strains (Figure 7B).

We next analyzed CC strains that were inherently more susceptible than B6 mice, and

were protected by BCG (n=4) or not (n=8) (Figure 7C). This yielded two clear phenotypes of

susceptible, non-protected CC strains. One subcluster comprising CC004 and CC044 had high

levels of T_{CM}, CD4$^+$Foxp3$^+$, CD8$^+$PD-1$^+$ and polyfunctional CD4$^-$CD8$^-$ T cells. The other cluster

had high levels of lung MIP-1a and CD3$^-$Foxp3$^+$ T cells, and fewer polyfunctional CD4$^-$CD8$^-$ T

cells. Interestingly, the unprotected strains with high CD4$^+$Foxp3$^+$T cells had low CD8$^+$Foxp3$^+$ T

cells, and the unprotected strains with high CD8$^+$Foxp3$^+$ T cells had low CD4$^+$Foxp3$^+$ T cells.

When looking at the protection model with the susceptible strains, several features were similar

to the model with all mice. In this model it is evident that while IL-17 in the lung homogenate

decreases with BCG vaccination in protected strains, the amount of IL-17 when stimulated with

the P300 peptide pool increases. (Figure 7D). Overall, the features that distinguish protected and

non-protected strains are similar for naturally resistant and susceptible mice.

Finally, we compared susceptible CC strains that were unprotected by BCG (i.e., poor

immunity) to inherently resistant CC strains that were protected by BCG (i.e., good immunity).

Unsurprisingly, there is clear separation of categories in the model with all mice. "Poor immunity"

CC strains tended to have high levels of inflammatory monokines (Figure 7E, 7F). "Good

immunity" CC strains high levels of RANTES after stimulation with anti-CD3. A subset of good

immunity strains including CC031 and CC011 also have high frequencies of IFNγ-producing CD8

T cells and various subsets of CD4⁻CD8⁻ T cells including those producing IFNγ or expressing

CD11a. In contrast, a subset of poor immunity strains had high levels of polyfunctional CD4⁻CD8⁻

T cells and ILCregs. The protection model revealed similar features with the interesting addition

of activated CD4 T_{RM} cells increasing with BCG vaccination in poor immunity but not good

immunity CC strains.

We find that immune features can be identified using PLSDA that can categorize the

inherent susceptibility and response to BCG of CC strains following Mtb infection. The expression

of different immune features was variable between CC strains even in the same protection

category, indicating no single path to protection. Looking at more refined subgroups of CC strains

allowed us to find features that define subclasses of protected and non-protected strains and

highlight features that did not emerge in the PLSR model such as non-T_{RM} effector and central

memory T cell populations.

Th1/17 cells detected in a subset of CC mice that are protected by BCG vaccination

An aim of this study was to identify factors that were associated with protection in

genetically diverse CC mice after vaccination and Mtb challenge. Vaccination enhanced IL-17

responses in the CC037, CC001, CC072 and CC031 strains, which were all protected by BCG

(Figure 5A, 5B). As the CC strains that produced IL-17 after antigen stimulation also had strong

IFNγ response, we assessed these T cell responses for dual production of IFNγ and IL-17. BCG

vaccinated CC037 mice had the highest IL-17 levels in lung homogenate (Figure 5A, 5B), and

after stimulation with P300 or anti-CD3 mAb, CD4 T cells were identified that produced both IFNγ

and IL-17 (Figure 8A). Dual IFNγ⁺IL-17⁺ CD4 T cells were not detected in C57BL/6 mice after

P300 stimulation, regardless of vaccination status. BCG vaccination significantly increased the

frequency of IFNγ$^+$IL-17$^+$-producing CD4 T cells in the CC037, CC072 and CC001 (Figure 8A,

8B). Further analysis identified BCG primed CD4 T cells from CC037 and CC001 to coexpress

both Tbet and RORγt after Mtb infection (Figure 8C, 8D), which are the master transcription

factors for Th1 and Th17 cells, respectively.

We next measured the polyfunctionality of the T cell response and expanded our cytokine

analysis to include IL-2 and TNF. In addition to IFNγ and IL-17, an increase in the proportion of

CD4 T cells producing IL-2 or TNF was detected (Figure 8E), indicating that CD4 T cell

polyfunctionality was increased in BCG vaccinated mice. A detailed analysis of the different

polyfunctional populations revealed different distributions of Th1/17 cells within each CC strain.

IFNγ$^+$IL-2$^+$TNF$^+$IL-17$^+$ cells were significantly increased in CC037 following BCG vaccination

(Figure S5, red box). Similarly, IFNγ$^+$TNF$^+$IL-17$^+$ cells were increased following BCG vaccination

in CC072 and CC001 (Figure S5, blue box). BCG led to a decrease in the frequency of IFNγ

"single positive' CD4 T cells in all four CC lines (Fig. S5, purple box). However, among these CC

strains only CC037 had a significant decrease in IFNγ$^+$TNF$^+$ cells. This contrasts with C57BL/6

mice where BCG vaccination had a minimal effect on CD4 T cell polyfunctionality. Significant

decreases were observed for IFNγ$^+$TNF$^+$ dual producers and IFNγ single positive cells in Mtb-

infected BCG-vaccinated C57BL/6 mice, consistent with the observation that Th1 dominant

response is associated with decrease bacterial burden in this background.

Finally, we performed a correlation analysis between all 60 immune features and the

normalized lung CFU from unvaccinated control and BCG vaccinated mice. For C57BL/6 mice,

IFNγ-producing and polyfunctional CD4 T cells correlated with lung CFU. In CC037 mice, IFNγ-

producing CD8 T cells correlated with lung CFU. Importantly, many features were negatively

correlated with lung bacterial burden in CC037 mice (Figure 8F). The frequency of CD4$^+$Roryt$^+$,

and antigen-stimulated IL-17 secretion and IFNγ/IL-17 dual-producing CD4 T cells were

significant even after correction for multiple testing, suggesting that these features were

associated with protection following BCG. There was a strong correlation between the frequency

of IFNγ/IL-17 dual-producing CD4 T cells and lung CFU in CC037 mice (Spearman r = -0.8947)

(Figure 8G).

Discussion

We sought to determine how pre-existing immunity to BCG modifies the immune response

to Mtb infection in a genetically diverse population using CC mice to model an outbred population.

An important outcome is our ranking of how BCG induced protection varies across 24 CC strains,

compared to control unvaccinated mice and relative to C57BL/6 mice, which were used as a

reference strain. We kept constant the BCG strain, growth conditions, dose, and vaccination

route. All mice were rested for 12 weeks after vaccination, challenged with the same Mtb strain,

and analyzed four weeks after infection. The different CC strains were categorized as being either

protected or unprotected by BCG against Mtb infection, based on whether vaccination led to a

statistically significant reduction in lung bacillary burden measured four weeks after low dose

aerosol infection. After controlling for these variables, the surprising result was that BCG protected

only half of the CC strains tested. As these strains differed primarily in the genes and alleles they

inherited from the CC founder strains, we conclude that the host genetic background has a major

influence on whether BCG confers protection against Mtb infection. These results extend our

analysis of the eight CC founder strains, which also vary in their ability to be protected by BCG

vaccination (15). Not only do environmental mycobacteria and genetic variations in BCG strains

used around the world interfere with the ability of BCG-induced protection, but we suggest that

host genetics should be considered a third important barrier to vaccine-mediated protection to TB.

Why BCG doesn't induce protection in all CC strains? Protection was largely defined

based on reductions in lung CFU four weeks after aerosol Mtb infection, which is a standard time

point in murine vaccine studies. Interestingly, the CC strains that were not protected by BCG can

be categorized based on their lung pathology. In half of the unprotected CC strains (e.g., CC004),

the BCG-vaccinated mice had changes in lung histology (e.g., more lymphocytic infiltrates) that

differed from the unvaccinated controls. We conclude that immunization generated a memory

recall response but it was insufficient to control lung CFU. Histological changes couldn't be

identified in other strains (e.g., CC024), and we suggest that BCG generated a suboptimal

immune response.

Importantly, we wished to identify the components of the immune response stimulated by

BCG, which were subsequently recalled after Mtb infection and associated with protection. We

hypothesized that CC strains protected by BCG would share common immunological features.

We postulated that by comparing protected to unprotected CC strains would allow us to filter out

components of the immune response that were elicited by vaccination or infection but not

associated with protection. The T cell immune response following BCG vaccination and Mtb

challenge was extensively characterized as T cells are essential for protection against Mtb.

Although considerable diversity was observed, BCG vaccination had little impact on the

composition of T cells recruited and maintained in the lung after infection. Instead, the variability

was largely shaped by the genetic background. In contrast, functional skewing of the T cell

responses by BCG was evident in several of the protected CC strains. While many of the

protected CC strains had unique flavors of immunity, we identified immune signatures associated

with BCG-elicited protection against TB. We found that different immune parameters were

associated with protection in the four different groups

Like lung homogenate IFNγ levels, lung IL-17 levels strongly correlated with lung CFU.

However, the correlation between IL-17 and lung CFU was lost in BCG protected CC strains after

vaccination and challenge. These data suggest that like IFNγ, IL-17 levels are a correlate of

disease, and not of protection. Interestingly, IL-17 concentrations co-varies with IL-1β and IL-6,

two cytokines that are involved in skewing of Th17 cells, and with TNF, KC, MIP2, and G-CSF,

all which are induced by IL-17. IL-17 could be more generally important in driving lung

inflammation than predicted by the C57BL/6 model. In contrast, the CD4 T cells from four of the

13 protected strains by BCG co-produced IFNγ and IL-17 after Mtb infection. None of the 11

unprotected strains had this phenotype, and these dual producers were detected only in BCG

vaccinated mice and not in sham controls, nor were they detected in the C57BL/6 reference strain.

Thus, this is an example of an immune response that BCG primes but is not manifested until the

CD4 T cell response is recalled by Mtb infection. These cells are similar to Th1/17 cells, or Th1*,

that are identified following TB infection, and are associated with late-forming, low burden

granulomas in a NHP TB infection model (25-27). However, an important difference is that IL-17

is not produced by Th1* cells. Importantly, these dual IFNγ/IL-17 producing CD4 T cells were

among the strongest correlates of protection. These data present a paradox. On the one hand,

IL-17 is a correlate of disease; conversely, IL-17-producing or RoRγt-expressing CD4 T cells are

a correlate of protection. We hypothesize that a cellular feature other than IL-17 production, such

the capacity for self-renewal, migration, or plasticity, is associated with Mtb control. The CC strains

will be invaluable in further unraveling the role of IL-17 and IL-17-producing T cells in immunity to

TB.

While extrinsic factors, many that disrupt cell mediated immunity, increase the risk of

developing TB, what drives the emergence of disease in most people is largely unknown. Host

genetics affects susceptibility to TB both in mice and people, and genetic loci have been identified

that increase TB risk (i.e., MSMD, polygenic) (16)[ref]. The factors that influence BCG efficacy

remains controversial. We hypothesize that genetic factors contribute to variation in the

effectiveness of BCG against pulmonary Mtb. Our analysis shows that while certain mouse

genotypes cannot be protected by BCG, the inherent susceptibility or resistance of the mouse

strain is not a predictor of vaccine efficacy. This suggests that a priori, vaccines are appropriate

interventions even in individuals that might have a genetic susceptibility to TB.

One caveat is that we have tested the efficacy of single vaccine in CC mice. The variation

in host responses of CC mice could be leveraged as an ideal platform for testing the efficacy of

preclinical vaccines in the context of a genetically diverse background. It is possible that vaccines

that work through alternate mechanisms may be able to protect CC mice that are refractory to

BCG vaccination. Instead of optimizing vaccine concepts in a single mouse genotype (e.g.,

C57BL/6 mice), identifying vaccines that are effective in numerous different genotypes (i.e.,

genetically distinct individuals), as modeled by CC or DO mice, represents an alternative

paradigm for preclinical vaccine development. Our data establish CC strains as a novel animal

resource that can facilitate the discovery of correlates of vaccine-induced protection and

accelerate preclinical vaccine testing.

Acknowledgements

We thank Clare Smith and Megan Proulx for helpful discussions, as well as other members of the

Behar lab. The MR-1 and CD1d tetramers were obtained through the NIH Tetramer Core Facility

(contract 75N93020D00005). The Vanderbilt Medical Center Digital Shared Histology Resource

(DHSR) digitized and hosted our histology data.

Author contribution

Declaration of interests

The authors declare no competing interests.

Funding

Supported by P01 AI123286 (C.M.S., M.F., S.M.B.), NSF Graduate Research Fellowship Grant

No.1745302 (D.G.), and R01 HL14541 (G.L.B). This work is also supported by NIAID IMPAcTB

and Army Institute for Collaborative Biotechnologies (D.L., D.G.).

Materials and Methods

Mice

Female C57BL/6/J (#0664) were purchased from The Jackson Laboratory. Female mice from 24

CC strains were purchased from the UNC Systems Genetics Core Facility (SGCF) between July

2018 and August 2019. The 24 CC strains used in this study include: CC001/Unc, CC002/Unc,

CC003/Unc, CC004/TauUnc, CC010/GeniUnc, CC011/Unc, CC012/GeniUnc, CC023/GeniUnc,

CC024/GeniUnc, CC025/GeniUnc, CC026/GeniUnc, CC031/GeniUnc, CC032/GeniUnc,

CC035/Unc, CC036/GemUnc, CC037/TauUnc, CC040/TauUnc, CC041/TauUnc,

CC042/GeniUnc, CC044/Unc, CC045/GeniUnc, CC059/TauUnc, CC061/GeniUnc,,

CC072/TauUnc. More information regarding the CC strains can be found at

http://csbio.unc.edu/CCstatus/index.py?run=AvailableLines.information.

All animal studies were conducted using the relevant guidelines and regulations, and approved

by the Institutional Animal Care and Use Committee at the University of Massachusetts Medical

School (UMMS) (Animal Welfare A3306-01), using the recommendations from the Guide for the

Care and Use of Laboratory Animals of the National Institutes of Health and the Office of

Laboratory Animal Welfare.

Bacterial strains

Bacteria expressing yellow fluorescent protein (sfYFP) were generated as described (17). Mice

were vaccinated with 105 CFU BCG (BCG-SSI strain, resuspended in 0.04% Tween/PBS)

subcutaneously (100 uL/injection).

Aerosolized Mtb infection of mice

Mice were infected by the aerosol route as previously described (17). Frozen bacterial stocks

were thawed and sonicated for 1 minute and then diluted into 5 ml of 0.01% Tween-80 in PBS.

The diluted bacterial suspension was aerosolized to infect mice using a Glas-Col chamber (Terre

Haute). The average number of bacteria delivered into the lung was determined for each

experiment by plating lung homogenate from 4-5 mice 24 hours after infection and ranged

between 40-150 CFU/mouse.

<u>Bacterial burden in lung and spleen</u>

Infected mice were euthanized at pre-determined timepoints, and the left lung or whole spleen

were homogenized using 2 mm zirconium oxide beads (Next Advance) in a FastPrep

homogenizer (MP Biomedicals). Tissue homogenates were serially diluted and plated on 7H11

agar plates (Hardy Diagnosis). CFU was enumerated after 19-21 days of incubation at 37°C and

5% CO_2.

<u>Lung cell preparation</u>

Single cell suspensions were prepared by homogenizing lungs using a GentleMACS tissue

dissociator (Miltenyi), digesting with 300 U/ml collagenase (Sigma) in complete RPMI (10% FBS,

2 mM L-Glutamine, 100 units/ml Penicillin/Streptomycin, 1 mM Na-Pyruvate, 1X Non-essential

amino acids, 0.5X Minimal essential amino acids, 25 mM of HEPES, and 7.5 mM of NaOH) at

37°C for 30 minutes, and followed by a second run of dissociation using the GentleMACS.

Suspensions were then sequentially filtered through 70 μm, treated with ACK lysis buffer (Gibco),

then filtered through 40 μm strainers.

<u>Flow cytometric analysis</u>

Cells were first stained with Zombie Fixable Viability dye (Biolegend) for 10 minutes at room

temperature (RT), after which cells were stained with antibodies in autoMACS running buffer

(Miltenyi) containing 5 ug/ml of anti-mouse CD16/32 (BioXcell) for 10 minutes at 4°C. In some

experiments, MAIT cells were identified using the MR-1-PE tetramer (NIH tetramer core facility)

at 37°C for 1hr. Cell were then stained with surface antibody cocktail for 20 minutes at 4°C. In

some experiments, CD1d-APC tetramer (NIH tetramer core facility) were also added at this time

to identify NKT cells. To measure transcription factor (TF) expression, cells were fixed and

permeabilized for 30 minutes at RT, followed by staining for 30 minutes with antibodies to the

different TF at RT using the Foxp3/TF staining buffer set (ThermoFisher). Samples were acquired

on Aurora (Cytek) or MACSquant (Miltenyi).

For intracellular staining, cells were first stimulated for 5 hours with either anti-CD3 antibodies (1

ug/mL, Biolegend) or P300 peptide megapool (23) which contains 300 peptides representing 90

antigens, and contains epitopes recognized by both murine CD4 and CD8 T cells. GolgiPlug (BD

Biosciences) was also given at this time (0.5ug/well). Viability and surface staining were

performed as described, after which cells were fixed instead with Fixation/Permeabilization kit for

20 minutes at 4°C. Cells were then stained with an intracellular cytokine antibody cocktail for 20

minutes at 4°C.

To inactivate the bacteria, samples were fixed with 1% paraformaldehyde/PBS for 1 hour at room

temperature and then washed with MACS buffer. Flow data were analyzed using FlowJo v10.7.1.

Polyfunctionality was examined using the SPICE platform (28).

LEGENDplex cytokine analysis

Lung cells were stimulated with P300 peptide megapool (1ug/mL) or anti-CD3 (1ug/mL) at a

concentration of 10^6 cells/mL for 24hrs. Culture supernatants were filtered through 0.2um filter,

after which cytokine levels were measured with LEGENDplex kits from Biolegend. Specifically,

we used their mouse T helper cytokine panel (IFNγ, TNF, IL-2, IL-4, IL-5, IL-6, IL-9, IL-10, IL-13,

IL-17a, IL-17f, IL-21, IL-22) as well as their mouse proinflammatory chemokine panel (RANTES,

MIP-3α, Eotaxin, TARC, KC, MCP-1, MIG, IP-10, MIP-1α, MIP-1β, BLC, LIX, MDC) for a total of

26 cytokines and chemokines.

Lung homogenate

Cytokines and chemokiens were measured from lung homogenates using the 31-plex discovery

Assay (MD31) from Eve technologies (Calgary, Canada).

Histology analysis

The right middle lung lobe from each mouse was immersion fixed in 10% formalin immediately

following euthanasia. Lung tissue was processed, embedded in paraffin, sectioned at 5um, and

stained with hematoxylin and eosin at the Morphology Core Facility at UMass Chan Medical

School for histological examination by a board-certified veterinary pathologist (GB). Blind

classification (BCG vs PBS) within each CC strain was determined based on the assumption that

BCG vaccination would have three possible effects on lungs: (i) reduce necrosis/neutrophilic

infiltrates; (ii) increase lymphocytes and/or plasma cell abundance; and (iii) reduce granuloma

size. Accuracy of the blind classification was assessed by an independent investigator (AO). All

glass slides were subsequently re-evaluated to more closely examine and describe

histopathology features attributable to BCG vaccination and attempt to find unique responses.

Glass slides were then digitized by Vanderbilt University Medical Center (Nashville, TN) and

whole images generated using Aperio ScanScope (Buffalo Grove, IL, USA) at 400 times normal

magnification with jpeg compression. Representative images were captured using Aperio

ImageScope magnified 40 and 400 times and imported into GraphPad Prism 9 to generate the

panel Figure. Digital images were not adjusted. Whole slide imaging and quantification of

immunostaining were performed in the Digital Histology Shared Resource (DHSR) at Vanderbilt

University Medical Center.

<u>Univariate correlations</u>

To evaluate how C57BL/6 mice, protected CC strains, and non-protected CC strains compared in their relationships between lung cytokine production and Mtb burden, a series of univariate Pearson correlations were conducted. Lung homogenate cytokines and chemokines measurements in the form of concentrations were natural logged and each of their Pearson correlations with the log10 lung CFU or spleen CFU were evaluated, with both the cytokine and CFU measurements separated into groups of C57BL/6, protected CC strains, and non-protected CC strains. A Bonferroni multiple hypothesis correction was applied to account for the number of correlations calculated.

<u>Multivariate analyses</u>

Multivariate analyses involved combining all four immunological datasets in this study: the lung homogenate cytokine/chemokine measurements, the ICS, the LEGENDplex cytokine analysis, and the T cell phenotypic flow cytometry. The analyses also included all CC mice used in the study in addition to the control C57/BL6 mice.

For the ICS, the data were analyzed as combinatorics of the seven markers measured in order to best capture polyfunctionality, and further divided by CD4 T cell, CD8 T cell, CD4$^-$CD8$^-$ T cell, and CD3$^-$ categories. The features were filtered by a minimum mean and variance percentage of 0.000001. Only P300 stimulated samples were used from the ICS dataset. The flow cytometry data was in the form of percentages of CD3$^+$ or CD3$^-$ cells, while the immunoassay data was in the form of concentrations. The combined dataset was natural log transformed and subsequently z-scored by measurement across all samples. For models that involved BCG-induced protection, these z-scored values were further transformed by subtracting the features from each vaccinated mouse and the mean of the features of the unvaccinated mice from the same mouse strain.

For the PLSR and PLSDA models, elastic net feature selection was performed in 100 trials of 5-

fold cross validation, where features that were present in at least 80% of trials were included in

the final model. Each model was compared to two null models: one with permuted lung CFUs and

one with randomly chosen features. The mean squared error of the prediction of lung CFU (for

the PLSR) or the classification accuracy (for the PLSDAs) was compared to that of the null models

in 50 trials of 5-fold cross validation.

Quantification and Statistical Analysis

Statistical analysis was performed using Graphpad Prism 9. P-values were calculated using

unpaired t test, one-way ANOVA, or two-way ANOVA as indicated in the figure legends.

Multivariate analyses were performed in R (v3.6.0 and 4.0.4) using the ropls Bioconductor

package.

^a CC strain	^b Batch	^c MHC Haplotype	^d log(ΔCFU) Exp1 (lung)	^d log(ΔCFU) Exp2 (lung)	^e log(ΔCFU) Exp1 (spleen)	^e log(ΔCFU) Exp2 (spleen)
CC001	4	b	0.94	1.45	2.79	1.45
CC002	3	b	0.83	n/a	0.46	n/a
CC003	3	g	0.07	-0.61	0.77	0.15
CC004	4	b	0.05	0.08	-0.16	-0.12
CC010	1	b	0.53	n/a	0.92	n/a
CC011	3	g7	1.04	n/a	2.08	n/a
CC012	4	wsb	0.79	n/a	0.89	n/a
CC023	1	b	1.17	-0.09	1.47	0.32
CC024	2	cast	0.15	0.24	0.38	0.62
CC025	3	pwk	1.40	n/a	0.43	n/a
CC026	2	cast	0.93	1.05	3.75	1.87
CC031	3	b	1.22	1.35	1.26	1.60
CC032	1	b	1.16	n/a	1.06	n/a
CC035	4	wsb	0.91	n/a	1.13	n/a
CC036	3	g7	0.62	n/a	1.84	n/a
CC037	1	b	1.37	1.84	1.07	1.05
CC040	2	b	-0.64	-0.28	-0.35	1.14
CC041	2	b	0.14	0.94	0.71	0.94
CC042	1	b	1.65	0.96	1.00	1.32
CC044	4	b/g7	0.08	0.13	0.12	0.39
CC045	4	z	0.24	n/a	0.13	n/a
CC059	2	b/a	1.39	n/a	1.17	n/a
CC061	1	b	0.93	n/a	1.58	n/a
CC072	2	b	0.97	1.37	1.06	1.41

Table 1. A description of the CC analysis batches.

^a A group of C57BL/6 mice were included in all batches as a reference strain. Data was normalized
to internal C57BL/6 control as indicated in the methods.
^b The first cohort of CC mice were analyzed in 4 separate batches as indicated.
^c MHC haplotype of each CC mouse as inherited from CC founders.
^{d, e} Thirteen of the 24 CC strains were analyzed a second time to validate the original results and
perform more in-depth immune analysis. ΔLungCFU^d and ΔSpleenCFU^e for the first experiment
and the repeat experiment are listed. Mice that were exacerbated by BCG vaccination are
highlighted in red.
655

[a] Strain	[b] BCG-induced protection	[c] Change in pathology	[d] How did the vaccinated mice differ?
CC003	N	N	No differences. Both groups had mild lesions with perivascular lymphocytic cuffs and a few macrophage-rich lesions
CC041	N	N	No differences. Both groups had mild lesions with perivascular lymphocytic cuffs and a few macrophage-rich lesions
CC024	N	N	No differences. Both groups had small hard-to-find lymphocytic lesions with perivascular and peribroncial lymphocytic cuffs
CC002	N	Y	Ctrl had mild lesions with perivascular lymphocytic cuffs. BCG mice had larger moderate sized macrophage-rich lesions. Lymphocytes were similarly abundant.
CC004	N	Y	Ctrl mice had mild to moderate lesions with lymphohistiocytic infiltrates. BCG mice have more lymphocytes with regions of dense perivascular/peribronchiolar lymphocytic cuffs.
CC036	N	Y	Ctrl mice have mild disease with lymphohistiocytic infiltrates. BCG mice also have lymphohistiocytic infiltrates with few very dense lymphocytic areas
CC040	N	Y	Ctrl mice have moderate necrotizing lesions with few lymphocytes. BCG mice have mild disease - small & hard to find lesions that are lymphocyte rich without necrosis.
CC044	N	Y	Unusual. Ctrl mice have lymphohistiocytic infiltrates with dense lymphocytic cuffs. BCG mice have lymphohistiocytic infiltrates with dense lymphocytic cuffs and lymphoid aggregates/follicules in parenchyma.
CC061	P	N	No differences. Unusual. Mild lesions. Both groups have dense lymphoid aggregates/follicles in parenchyma. Also, perivascular lymphocytic cuffs.
CC072	P	N	No differences. Mild disease. perivascular lymphocytic cuffs with a few macrophage-rich lesions. One mouse has a foreign-body granulomatous bronchitis/pneumonia with intralesional food/bedding/plant.
C57BL/6	P	Y	Ctrl mice had mild to moderate lymphohistiocytic lesions. BCG mice had with more, dense lymphocytic infiltrates
CC001	P	Y	Ctrl mice had moderate lesions with lymphohistiocytic infiltrates and equivocal necrosis. BCG mice had mild lesions with perivascular lymphocytic cuffs and a few macrophage-rich lesions without necrosis
CC023	P	Y	Ctrl mice had moderate to severe lesions with necrosis and minimal lymphocytes. BCG mice have moderate lesions with more lymphocytes and less necrosis.
CC025	P	Y	Ctrl mice have moderate disease with mild lymphocytic infiltrates and necrosis. BCG mice have abundant dense perivascular/peribronchiolar lymphocytic ifiltrates and lymphohistiocytic granulomas and less necrosis.
CC026	P	Y	Ctrl mice had moderate lesions with lymphohistiocytic infiltrates. BCG mice had mild lesions with lymphohistiocytic infiltrates and a few very dense lymphocytic areas
CC031	P	Y	Ctrl mice have moderate disease with lymphohistiocytic infiltrates. BCG mice have abundant dense perivascular/peribronchiolar lymphocytes and lymphohistiocytic granulomas
CC032	Protected	Y	Ctrl mice have mild disease with lymphohistiocytic infiltrates. BCG mice have more lymphocyte rich lesions that are small & hard to find
CC037	Protected	Y	Ctrl mice have moderate disease with lymphohistiocytic infiltrates. BCG mice have abundant dense perivascular/peribronchiolar lymphocytes and lymphohistiocytic granulomas
CC042	Protected	Y	Ctrl mice have mild to moderate disease with few lymphocytes and necrosis. BCG mice have mild disease with more lymphocytes, less necrosis
CC059	Protected	Y	Ctrl mice had mild disease with lymphohistiocytic infiltrates with small regions of necrosis. BCG mice have abundant dense perivascular/peribronchiolar lymphocytes and lymphohistiocytic granulomas without necrosis

Table 2. Summary of histological changes of CC mice.
[a] Nineteen of the 24 CC strains were available for analysis.
[b] The effect of BCG on the lung CFU is from Figure 1.
[c] Assessment by a pathologist of whether differences existed between sham and BCG vaccinated
mice after Mtb infection (n=5/strain/condition).
[d] Summary of observed differences between the groups.

Figure legends

Figure 1. BCG-induced protection in 24 CC strains and C57BL/6 mice.

(A) Lung and (D) spleen CFU of all 24 CC mice (n=120/group; top) or all C57BL/6 mice

(n=20/group; bottom) at 4 weeks post infection. Mice were vaccinated with BCG or administered

PBS and rested for 12 weeks prior to infection. Each point represents an individual mouse. Line,

median. Student's t-test; ****, $p < 0.0001$.

(B) Protection in the lung and (E) was assessed by comparing the CFU in BCG vs. sham

vaccinated mice for each individual mouse strain (n=5/group) using the Benjamini-Hochberg

procedure to determine the false discovery rate (FDR) (top). The ΔCFU of BCG vs. sham

vaccinated was calculated (bottom). The open bar is the reference strain C57BL/6 strain

(n=20/group). *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$; ****, $p < 0.0001$.

(C) Correlation in the lung and (F) spleen between susceptibility and protection. The relative

susceptibility (y-axis, defined by CFU in unvaccinated group, relative to the C57BL/6 reference

strain) and the protection conferred by BCG (x-axis, defined by ΔCFU of vaccinated mice

compared to unvaccinated controls, relative to the C57BL/6 reference strain). The graph is

centered on C57BL/6 mice, leading to the different CC strains being placed in four quadrants

based on their relative susceptibility and protection.

Figure 2. Lung histology of CC mice after BCG vaccination and Mtb challenge

Representative lung histology of three CC strains (CC025, CC037, CC072) that were protected

by BCG, after sham or BCG vaccination and Mtb challenge (Mtb and BCG+Mtb, respectively)

(left). Representative lung histology of three CC strains (CC003, CC004, CC024) that were

unprotected by BCG (right). Original magnification, 40X (inset, 400X). Scale bar, 500 μm (inset,

50 μm).

Figure 3. BCG-protected CC mice have an altered lung microenvironment of CC mice after Mtb

infection.

(A) IFNγ levels were measured in the lungs of either PBS or BCG vaccinated CC mice at 4 weeks

post infection. A two-way ANOVA was performed using Sidak's correction for multiple

comparisons. ***, $p < 0.001$; ****, $p < 0.0001$".

(B) – (F) Pearson correlation between lung CFU and (B) IFNγ, IL-17, TNF, IL-1β, KC and IL-6 (C)

LIX (D) RANTES (E) MIP-1β and (F) M-CSF, MIP-2, G-CSF, LIF, comparing PBS and BCG

vaccinated mice at 4 weeks post infection in C57BL/6 mice (B6), non-protected (NP) or protected

(P) CC mice.

Figure 4. Frequency of various T cell populations in protected and unprotected CC mice following

Mtb infection.

(A) Proportion of CD3$^+$ cell, CD3$^+$CD4$^+$ and CD3$^+$CD8$^+$ T cells in the lungs of PBS and BCG

vaccinated CC mice at 4 weeks post infection.

(B) Proportion of memory CD4 and CD8 T cells in the lungs of either PBS or BCG vaccinated CC

mice at 4 weeks post infection, as defined by CD44$^+$ and various combinations of CD62L and

CD127 expression; CD62L$^+$ CD127$^+$ (central memory), CD62L$^-$ CD127$^-$ (effector), CD62L$^-$

CD127$^+$ (effector memory).

(C) Proportion of resident memory CD4 and CD8 T cells in the lungs of either PBS or BCG

vaccinated CC mice at 4 weeks post infection, as defined by CD44$^+$ CD103$^+$ CD69$^+$.

(D) Proportion of Th1, Treg and Th17 cells in the lungs of either PBS or BCG vaccinated CC mice

at 4 weeks post infection, as defined by CD44$^+$ and Tbet$^+$ (Th1), Foxp3$^+$ (Treg) or RoRγt (Th17)

expression.

(E) Proportion of CD4 and CD8 T cells expressing either CXCR3 or CX$_3$CR1 in the lungs of either

PBS or BCG vaccinated CC mice at 4 weeks post infection.

(F) PCA of the T cell phenotyping data. Each point represents the average of usually 5 mice within

a given mouse strain that are either BCG vaccinated or unvaccinated. Open symbols,

unvaccinated; closed symbols, BCG vaccinated. Each color represents a different mouse strain.

(A–E) The data represent the mean ± SEM from one experiment. Two-way analysis of variance

with the Benjamini and Hochberg multiple comparsion method. The FDR was set to 0.05 and the

numbers in the figures are the q value.

Figure 5. Cytokine response in protected and unprotected CC strains following Mtb infection.

(A) Cells were isolated from the lungs of CC mice at 4 weeks post infection and stimulated for 24

hours with the P300 peptide pool. IFNγ, IL-2, TNF and IL-17a were measured in the supernatants.

(B) and (C) Cells from the lungs of CC mice at 4 weeks post infection were stimulated for 5 hours

with the P300 peptide pool. Frequencies of CD4 or CD8 T cells that expressed IFNγ, IL-2, TNF

or IL-17 was determined using intracellular cytokine staining.

(A–C) The data represent the mean ± SEM from one experiment. Two-way analysis of variance

with Sidak's multiple comparison test was used to determine significance. *, $p < 0.05$; **, $p < 0.01$;

***, $p < 0.001$; ****, $p < 0.0001$.

Figure 6. Multivariate correlates of protection across all 24 CC strains

(A) A partial least squares regression (PLSR) model was performed to associate BCG-induced

changes in lung immune features with BCG-induced changes in lung CFU. Each data point

represents a vaccinated mouse that has the mean of the unvaccinated mice from the same strain

subtracted from it. The scores of the top two latent variables of the model are shown, with the

color of each data point representing the changed in lung CFU induced by BCG vaccination. The

percents of the variance in the lung features captured by each latent variable are shown as

percentages in the axes.

(B) The bar graph depicts the loadings of the first latent variable of the PLSR model in (A). Positive

loadings indicate immune features that increase from BCG vaccination in samples that are on the

positive x-axis in the scores plot, which generally indicates they have worse protection. Negative

loadings conversely indicate immune features that generally increase from BCG vaccination in

samples that are protected by vaccination.

(C) The heatmap illustrates all features selected to be included in the model, indicating they were

important for associating strains with the lung protection continuum. Colors are representations

of z-scored features, with a different color representing each 10% quantile of the data. Columns

are averages of the mice in the CC strains (n=5 for most cases). Feature names are colored

based on the dataset from which they come. In (B) and (C), features from the ICS dataset indicate

production of the cytokines listed and no production of any measured cytokines not listed.

Figure 7. Identifying signatures of protection in subclasses of CC strains

(A), (C), (E) Scores plot (top) and heatmaps (bottom) of selected features for PLSDA models.

Each dot represents a single mouse. The two colors in the scores plots signify the two classes

being compared. An ellipse is drawn around the 95% confidence interval for each class. Percent

of variance of the immune feature data is shown on the axes. Heatmaps show features that were

most important for distinguishing the two classes. Colors are representations of z-scored features.

Columns are averages of the mice in the CC strains, with vaccinated and unvaccinated mice

being separated (n=5 for most cases). Feature names are colored based on the dataset from

which they come (blue=T cell phenotyping, brown=lung homogenate cytokines, green=ICS,

magenta=LEGENDplex). NP = non-protected, P = protected, R = resistant, S = susceptible.

(B), (D), (F) PLSDA models performed as in (A), (C), and (E), except that features are the changes

induced by BCG vaccination as in Figure 6.

Figure 8. Th1/17 cells correlate with BCG-mediated protection in a subset of protected CC mice.

(A) Cells were isolated from the lungs of CC mice at 4 weeks post infection and stimulated for 5 hours with either aCD3, P300 peptide pool or media control. Representative flow plots of IFNγ and IL-17 producing CD4 T cells from CC037 and C57BL/6 are shown. Data is representative of two independent experiments.

(B) Frequencies of CD4 T cells that expressed both IFNγ and IL-17 was determined using intracellular cytokine staining. Data is representative of two independent experiments. One-way analysis of variance with Sidak's multiple comparison test was used to determine significance. **, $p < 0.01$; ***, $p < 0.001$; ****, $p < 0.0001$.

(C) Representative flow plots of CD4 T cells expressing either T-bet or RORγt in CC037 or C57BL/6 mice. Data is representative of two independent experiments.

(D) Frequencies of CD4 T cells that expressed both T-bet or RORγt in CC037 or C57BL/6 mice. Data is representative of two independent experiments. One-way analysis of variance with Sidak's multiple comparison test was used to determine significance. **, $p < 0.01$; ***, $p < 0.001$.

(E) Proportion of CD4 T cells that express various combinations of IFNγ, TNF, IL-2 or IL-17 in either PBS or BCG vaccinated mice at 4 weeks post Mtb infection. Plots are generated using the SPICE platform (28).

(F) Volcano plots. Correlation between 60 immune features and the lung CFU determined for CC037 and C57BL/6 mice (from Batch 1). Dotted line, $p = 0.05$. *, $q < 0.05$, corrected with the Benjamini and Hochberg multiple comparison method (FDR = 0.05)

(G) Correlation of IFNγ$^+$IL-17$^+$ cells and bacterial burden in the lungs of either PBS or BCG vaccinated CC037 mice at 4 weeks post Mtb infection. Colors represent BCG vaccinated mice (red) or unvaccinated mice (blue). Spearman *r* and p value are shown.

 SUPPLEMENTAL INFORMATION

Supplementary tables

Table S1. –Pathological descriptions of changes lung histology after BCG vaccination in CC mice.

Table S2. A description of select CC strains that were repeated for in-depth immune analysis.

Supplementary figure legends

Figure S1. Gating strategy for the analysis of the cell surface T cell phenotype.

(A) Lymphocytes were identified using FSC and SSC properties, after which viability dyes were

used to identify live cells. Singlet gates were used to exclude doublets, after which T cells and

non-T cells were identified using the lineage marker CD3.

(B) NKT and MAIT cells were identified from the CD3$^+$ gate using their antigen-loaded CD1d and

MR1 tetramers, respectively. CD4$^+$, CD8$^+$ and CD4$^-$CD8$^-$ T cells were also identified from this

population for further analysis.

(C) Expression of various markers were also examined within the CD3$^-$ population. As an

example, flow plot represents expression of T-bet within the CD3$^-$ population.

(D) Identification of memory T cells. CD44 was used to identify antigen-experienced T cells.

Resident memory T cells (T$_{RM}$) were identified based on CD103 expression. From the CD103$^-$

fraction, CD127 and CD62L expression identified central memory (T$_{CM}$), effector (T$_E$) and effector

memory (T$_{EM}$). Subset identification of various T cell populations (ex: Tbet$^+$CD4$^+$ T cells) as well

as activation markers (ex: PD1, CD11a) were also examined in CD4$^+$, CD8$^+$ and CD4$^-$CD8$^-$ T cell

subsets.

(E) Naïve T cells identified within the CD44$^-$ compartment using CD62L expression.

Figure S2. Frequencies of non-conventional T cells in CC mice. Frequencies of MR-1-restricted

MAIT cells and CD1d-restricted NKT cells were measured in the lungs of PBS or BCG vaccinated

CC mice at 4 weeks post Mtb infection. The data in the graphs represent the mean ± SD of one

experiment. Two-way analysis of variance with the Benjamini and Hochberg multiple comparsion

method. The FDR was set to 0.05 and the numbers in the figures are the q value.

Figure S3. Gating strategy for the analysis of T cell function (intracellular cytokine staining).

(A) Lymphocytes were identified as using FSC and SSC properties, after which viability dyes were

used to identify live cells. Singlet gates were used to exclude doublets, after which T cells were

identified using the lineage marker CD3.

(B) CD4, CD8 and CD4$^-$CD8$^-$ T cells were also identified from the CD3$^+$ population for further

analysis. Cytokine expression (ex: IFNγ) were examined in all 3 T cell sub-populations.

(C) Cytokine expression was also examined within the CD3$^-$ population.

Figure S4. Multivariate model accuracies

(A) The accuracy of our protection model (Figure 6) was compared to that of two different types

of null model, where our measure of accuracy was mean squared error of prediction of Δlog (lung

CFU). Our PLSR model performed significantly better than both a null model consisting of random

features instead of our elastic-net chosen features and a null model where the outcomes of Δlog

(lung CFU) were scrambled. Our PLSR model and each of the null models were run 50 times,

and each dot represents one of these trials. Labeled p-values are from one-way ANOVAs.

(B) Our models highlighting the immune features that distinguish pairs of protection/susceptibility

categories (Figure 7) were compared to two null models as in (A). Each of our models, both the

ones involving all mice and the protection models were significantly better than both null models.

Here, the accuracy metric is classification accuracy. Labeled p-values are from one-way

ANOVAs.

Figure S5. Examining the Polyfunctionality of CD4 T cells in select CC strains. Boolean gating

was used to identify various sub-populations amongst the CD4 T cells following P300 stimulation

in certain CC mice. An increase in various IFNγ$^+$IL-17$^+$ CD4 T cells (that also produced TNF or

IL-2) were detected in CC037, CC072, CC031 and CC001 but not in C57BL/6. In contrast, a

noticeable decrease in the frequency of IFNγ "single positive" CD4 T cells was observed in all

strains including C57BL/6. The data in the graphs represent the mean ± SD of two experiment.

One-way analysis of variance with Sidak's multiple comparison test was used to determine

significance. *, p < 0.05; **, p < 0.01; ***, p < 0.001; ****, p < 0.0001.

849   References
850

1.   WHO. Global Tuberculosis Report 2021. 2021.

2.   Colditz GA, Brewer TF, Berkey CS, Wilson ME, Burdick E, Fineberg HV, et al. Efficacy of
BCG vaccine in the prevention of tuberculosis. Meta-analysis of the published literature.
JAMA. 1994;271(9):698-702.

3.   Roy A, Eisenhut M, Harris RJ, Rodrigues LC, Sridhar S, Habermann S, et al. Effect of
BCG vaccination against Mycobacterium tuberculosis infection in children: systematic
review and meta-analysis. *BMJ.* 2014;349:g4643.

4.   Mangtani P, Abubakar I, Ariti C, Beynon R, Pimpin L, Fine PE, et al. Protection by BCG
vaccine against tuberculosis: a systematic review of randomized controlled trials. *Clin*
Infect Dis. 2014;58(4):470-80.

5.   Abubakar I, Pimpin L, Ariti C, Beynon R, Mangtani P, Sterne JA, et al. Systematic review
and meta-analysis of the current evidence on the duration of protection by bacillus
Calmette-Guerin vaccination against tuberculosis. *Health Technol Assess.* 2013;17(37):1-
372, v-vi.

6.   Fine PE. Variation in protection by BCG: implications of and for heterologous immunity.
Lancet. 1995;346(8986):1339-45.

7.   Flynn JL, Gideon HP, Mattila JT, and Lin PL. Immunology studies in non-human primate
models of tuberculosis. *Immunol Rev.* 2015;264(1):60-73.

8.   Ahmed M, Thirunavukkarasu S, Rosa BA, Thomas KA, Das S, Rangel-Moreno J, et al.
Immune correlates of tuberculosis disease and risk translate across species. *Sci Transl*
Med. 2020;12(528).

9.   Aagaard C, Knudsen NPH, Sohn I, Izzo AA, Kim H, Kristiansen EH, et al. Immunization
with Mycobacterium tuberculosis-Specific Antigens Bypasses T Cell Differentiation from
Prior Bacillus Calmette-Guerin Vaccination and Improves Protection in Mice. *J Immunol.*
2020;205(8):2146-55.

10.   Woodworth JS, Clemmensen HS, Battey H, Dijkman K, Lindenstrom T, Laureano RS, et
al. A Mycobacterium tuberculosis-specific subunit vaccine that provides synergistic
immunity upon co-administration with Bacillus Calmette-Guerin. *Nat Commun.*
2021;12(1):6658.

11.   Kramnik I, and Beamer G. Mouse models of human TB pathology: roles in the analysis of
necrosis and the development of host-directed therapies. *Semin Immunopathol.*
2016;38(2):221-37.

12.   Nemeth J, Olson GS, Rothchild AC, Jahn AN, Mai D, Duffy FJ, et al. Contained
Mycobacterium tuberculosis infection induces concomitant and heterologous protection.
PLoS Pathog. 2020;16(7):e1008655.

13.   Noll KE, Ferris MT, and Heise MT. The Collaborative Cross: A Systems Genetics
Resource for Studying Host-Pathogen Interactions. *Cell Host Microbe.* 2019;25(4):484-98.

14.   Churchill GA, Airey DC, Allayee H, Angel JM, Attie AD, Beatty J, et al. The Collaborative
889   Cross, a community resource for the genetic analysis of complex traits. *Nat Genet.*
890   2004;36(11):1133-7.

891   15.   Smith CM, Proulx MK, Olive AJ, Laddy D, Mishra BB, Moss C, et al. Tuberculosis
892   Susceptibility and Vaccine Protection Are Independently Controlled by Host Genotype.
893   *MBio.* 2016;7(5).

894   16.   Smith CM, Baker RE, Proulx MK, Mishra BB, Long JE, Park SW, et al. Host-pathogen
895   genetic interactions underlie tuberculosis susceptibility in genetically diverse mice. *Elife.*
896   2022;11.

897   17.   Lee J, Boyce S, Powers J, Baer C, Sassetti CM, and Behar SM. CD11cHi monocyte-
898   derived macrophages are a major cellular compartment infected by Mycobacterium
899   tuberculosis. *PLoS Pathog.* 2020;16(6):e1008621.

900   18.   Nandi B, and Behar SM. Regulation of neutrophils by interferon-gamma limits lung
901   inflammation during tuberculosis infection. *J Exp Med.* 2011;208(11):2251-62.

902   19.   Sahiratmadja E, Alisjahbana B, Buccheri S, Di Liberto D, de Boer T, Adnan I, et al. Plasma
903   granulysin levels and cellular interferon-gamma production correlate with curative host
904   responses in tuberculosis, while plasma interferon-gamma levels correlate with
905   tuberculosis disease activity in adults. *Tuberculosis (Edinb).* 2007;87(4):312-21.

906   20.   Sakai S, Kauffman KD, Schenkel JM, McBerry CC, Mayer-Barber KD, Masopust D, et al.
907   Cutting edge: control of Mycobacterium tuberculosis infection by a subset of lung
908   parenchyma-homing CD4 T cells. *J Immunol.* 2014;192(7):2965-9.

909   21.   Becattini S, Latorre D, Mele F, Foglierini M, De Gregorio C, Cassotta A, et al. Functional
910   heterogeneity of human memory CD4(+) T cell clones primed by pathogens or vaccines.
911   *Science.* 2015;347(6220):400-6.

912   22.   Smith CM, Proulx MK, Lai R, Kiritsy MC, Bell TA, Hock P, et al. Functionally Overlapping
913   Variants Control Tuberculosis Susceptibility in Collaborative Cross Mice. *mBio.*
914   2019;10(6).

915   23.   Lindestam Arlehamn CS, McKinney DM, Carpenter C, Paul S, Rozot V, Makgotlho E, et
916   al. A Quantitative Analysis of Complexity of Human Pathogen-Specific CD4 T Cell
917   Responses in Healthy M. tuberculosis Infected South Africans. *PLoS Pathog.*
918   2016;12(7):e1005760.

919   24.   Patankar YR, Sutiwisesak R, Boyce S, Lai R, Lindestam Arlehamn CS, Sette A, et al.
920   Limited recognition of Mycobacterium tuberculosis-infected macrophages by polyclonal
921   CD4 and CD8 T cells from the lungs of infected mice. *Mucosal Immunol.* 2020;13(1):140-
922   8.

923   25.   Lindestam Arlehamn CS, Gerasimova A, Mele F, Henderson R, Swann J, Greenbaum JA,
924   et al. Memory T cells in latent Mycobacterium tuberculosis infection are directed against
925   three antigenic islands and largely contained in a CXCR3+CCR6+ Th1 subset. *PLoS
926   Pathog.* 2013;9(1):e1003130.

927   26.   Acosta-Rodriguez EV, Rivino L, Geginat J, Jarrossay D, Gattorno M, Lanzavecchia A, et
928   al. Surface phenotype and antigenic specificity of human interleukin 17-producing T helper
929   memory cells. *Nat Immunol.* 2007;8(6):639-46.

930   27.   Gideon HP, Hughes TK, Tzouanas CN, Wadsworth MH, 2nd, Tu AA, Gierahn TM, et al.
931   Multimodal profiling of lung granulomas in macaques reveals cellular correlates of
932   tuberculosis control. *Immunity.* 2022;55(5):827-46 e10.

933   28.   Roederer M, Nozzi JL, and Nason MC. SPICE: exploration and analysis of post-cytometric
934   complex multivariate datasets. *Cytometry A.* 2011;79(2):167-74.

Figure 1. BCG-induced protection in 24 CC strains and C57BL/6 mice.
(A) Lung and (D) spleen CFU of all 24 CC mice (n=120/group; top) or all C57BL/6 mice (n=20/group; bottom) at 4 weeks post infection. Mice were vaccinated with BCG or administered PBS and rested for 12 weeks prior to infection. Each point represents an individual mouse. Line, median. Student's t-test; ****, p < 0.0001.(B) Protection in the lung and (E) was assessed by comparing the CFU in BCG vs. sham vaccinated mice for each individual mouse strain (n=5/group) using the Benjamini-Hochberg procedure to determine the false discovery rate (FDR) (top). The ΔCFU of BCG vs. sham vaccinated was calculated (bottom). The open bar is the reference strain C57BL/6 strain (n=20/group). *, p < 0.05; **, p < 0.01; ***, p < 0.001; ****, p < 0.0001. (C) Correlation in the lung and (F) spleen between susceptibility and protection. The relative susceptibility (y-axis, defined by CFU in unvaccinated group, relative to the C57BL/6 reference strain) and the protection conferred by BCG (x-axis, defined by ΔCFU of vaccinated mice compared to unvaccinated controls, relative to the C57BL/6 reference strain). The graph is centered on C57BL/6 mice, leading to the different CC strains being placed in four quadrants based on their relative susceptibility and protection.

CC strain[a]	Batch[b]	MHC Haplotype[c]	ΔLung CFU$_1$ reduction[d] (logCFU)	ΔLung CFU$_2$ reduction[d] (logCFU)	ΔSpleen CFU$_1$ reduction[e] (logCFU)	ΔSpleen CFU$_2$ reduction[e] (logCFU)
CC001/Unc	4	b	0.94	1.45	2.79	1.45
CC002/Unc	3	b	0.83	n/a	0.46	n/a
CC003/Unc	3	g	0.07	-0.61	0.77	0.15
CC004/TauUnc	4	b	0.05	0.08	-0.16	-0.12
CC010/GeniUnc	1	b	0.53	n/a	0.92	n/a
CC011/Unc	3	g7	1.04	n/a	2.08	n/a
CC012/GeniUnc	4	wsb	0.79	n/a	0.89	n/a
CC023/GeniUnc	1	b	1.17	-0.09	1.47	0.32
CC024/GeniUnc	2	cast	0.15	0.24	0.38	0.62
CC025/GeniUnc	3	pwk	1.40	n/a	0.43	n/a
CC026/GeniUnc	2	cast	0.93	1.05	3.75	1.87
CC031/GeniUnc	3	b	1.22	1.35	1.26	1.60
CC032/GeniUnc	1	b	1.16	n/a	1.06	n/a
CC035/Unc	4	wsb	0.91	n/a	1.13	n/a
CC036/Unc	3	g7	0.62	n/a	1.84	n/a
CC037/TauUnc	1	b	1.37	1.84	1.07	1.05
CC040/TauUnc	2	b	-0.64	-0.28	-0.35	1.14
CC041/TauUnc	2	b	0.14	0.94	0.71	0.94
CC042/GeniUnc	1	b	1.65	0.96	1.00	1.32
CC044/Unc	4	b/g7	0.08	0.13	0.12	0.39
CC045/GeniUnc	4	z	0.24	n/a	0.13	n/a
CC059/TauUnc	2	b/a	1.39	n/a	1.17	n/a
CC061/GeniUnc	1	b	0.93	n/a	1.58	n/a
CC072/TauUnc	2	b	0.97	1.37	1.06	1.41

Table S1. A description of select CC strains that were repeated for in-depth immune analysis.
a A group of C57BL/6 mice were included in all batches as a reference strain. Data was normalized to internal C57BL/6 control as indicated in the methods.
b The first cohort of CC mice were analyzed in 4 separate batches as indicated.
c MHC haplotype of each CC mouse as inherited from CC founders.
d, e Thirteen of the 24 CC strains were analyzed a second time to validate the original results and perform more in-depth immune analysis. ΔLungCFU[d] and ΔSpleenCFU[e] for the first experiment and the repeat experiment are listed. Mice that were exacerbated by BCG vaccination are highlighted in red.

Strain[a]	BCG effect[b] (lung CFU)	Were differences observed?[c]	How did the vaccinated mice differ?[d]
CC003/TauUNC	Not protected	N	No differences. Both groups had mild lesions with perivascular lymphocytic cuffs and a few macrophage-rich lesions
CC041/TauUNC	Not protected	N	No differences. Both groups had mild lesions with perivascular lymphocytic cuffs and a few macrophage-rich lesions
CC024/GeniUNC	Not protected	N	No differences. Both groups had small hard-to-find lymphocytic lesions with perivascular and peribroncial lymphocytic cuffs
CC002/TauUNC	Not protected	Y	Ctrl had mild lesions with perivascular lymphocytic cuffs. BCG mice had larger moderate sized macrophage-rich lesions. Lymphocytes were similarly abundant.
CC004/TauUnc	Not protected	Y	Ctrl mice had mild to moderate lesions with lymphohistiocytic infiltrates. BCG mice have more lymphocytes with regions of dense perivascular/peribronchiolar lymphocytic cuffs.
CC036/GeniUnc	Not protected	Y	Ctrl mice have mild disease with lymphohistiocytic infiltrates. BCG mice also have lymphohistiocytic infiltrates with few very dense lymphocytic areas
CC040	Not protected	Y	Ctrl mice have moderate necrotizing lesions with few lymphocytes. BCG mice have mild disease - small & hard to find lesions that are lymphocyte rich without necrosis.
CC044/GeniUnc	Not protected	Y	Unusual. Ctrl mice have lymphohistiocytic infiltrates with dense lymphocytic cuffs. BCG mice have lymphohistiocytic infiltrates with dense lymphocytic cuffs and lymphoid aggregates/follicles in parenchyma.
CC061/GeniUNC	Protected	N	No differences. Unusual. Mild lesions. Both groups have dense lymphoid aggregates/follicles in parenchyma. Also, perivascular lymphocytic cuffs.
CC072/TauUNC	Protected	N	No differences. Mild disease. perivascular lymphocytic cuffs with a few macrophage-rich lesions. One mouse has a foreign-body granulomatous bronchitis/pneumonia with intralesional food/bedding/plant.
C57BL/6	Protected	Y	Ctrl mice had mild to moderate lymphohistiocytic lesions. BCG mice had with more, dense lymphocytic infiltrates
CC001/TauUnc	Protected	Y	Ctrl mice had moderate lesions with lymphohistiocytic infiltrates and equivocal necrosis. BCG mice had mild lesions with perivascular lymphocytic cuffs and a few macrophage-rich lesions without necrosis
CC023	Protected	Y	Ctrl mice had moderate to severe lesions with necrosis and minimal lymphocytes. BCG mice have moderate lesions with more lymphocytes and less necrosis.
CC025/TauUNC	Protected	Y	Ctrl mice have moderate disease with mild lymphocytic infiltrates and necrosis. BCG mice have abundant dense perivascular/peribronchiolar lymphocytic ifiltrates and lymphohistiocytic granulomas and less necrosis.
CC026	Protected	Y	Ctrl mice had moderate lesions with lymphohistiocytic infiltrates. BCG mice had mild lesions with lymphohistiocytic infiltrates and a few very dense lymphocytic areas
CC031/GeniUnc	Protected	Y	Ctrl mice have moderate disease with lymphohistiocytic infiltrates. BCG mice have abundant dense perivascular/peribronchiolar lymphocytes and lymphohistiocytic granulomas
CC032/GeniUNC	Protected	Y	Ctrl mice have mild disease with lymphohistiocytic infiltrates. BCG mice have more lymphocyte rich lesions that are small & hard to find
CC037/TauUNC	Protected	Y	Ctrl mice have moderate disease with lymphohistiocytic infiltrates. BCG mice have abundant dense perivascular/peribronchiolar lymphocytes and lymphohistiocytic granulomas
CC042	Protected	Y	Ctrl mice have mild to moderate disease with few lymphocytes and necrosis. BCG mice have mild disease with more lymphocytes, less necrosis
CC059/TauUNC	Protected	Y	Ctrl mice had mild disease with lymphohistiocytic infiltrates with small regions of necrosis. BCG mice have abundant dense perivascular/peribronchiolar lymphocytes and lymphohistiocytic granulomas without necrosis

Table S2. Pathological descriptions of changes lung histology after BCG vaccination in CC mice
a Nineteen of the 24 CC strains were available for analysis.
b The effect of BCG on the lung CFU is from Figure 1.
c Assessment by a pathologist of whether differences existed between sham and BCG vaccinated mice after Mtb infection (n=5/strain/condition).
d Summary of observed differences between the groups.

.

Figure 2. Lung histology of CC mice after BCG vaccination and Mtb challenge
Representative lung histology of three CC strains (CC025, CC037, CC072) that were protected by BCG, after sham or BCG vaccination and Mtb challenge (Mtb and BCG+Mtb, respectively) (left). Representative lung histology of three CC strains (CC003, CC004, CC024) that were unprotected by BCG (right). Original magnification, 40X (inset, 400X). Scale bar, 500 μm (inset, 50 μm).

Figure 3. BCG-protected CC mice have an altered lung microenvironment of CC mice after Mtb infection.

(A) IFNγ levels were measured in the lungs of either PBS or BCG vaccinated CC mice at 4 weeks post infection. A two-way ANOVA was performed using Sidak's correction for multiple comparisons. ***, p < 0.001; ****, p < 0.0001". (B) – (F) Pearson correlation between lung CFU and (B) IFNγ, IL-17, TNF, IL-1β, KC and IL-6 (C) LIX (D) RANTES (E) MIP-1β and (F) M-CSF, MIP-2, G-CSF, LIF, comparing PBS and BCG vaccinated mice at 4 weeks post infection in C57BL/6 mice (B6), non-protected (NP) or protected (P) CC mice.

Figure S1. Gating strategy for the analysis of the cell surface T cell phenotype.
(A) Lymphocytes were identified using FSC and SSC properties, after which viability dyes were used to identify live cells. Singlet gates were used to exclude doublets, after which T cells and non-T cells were identified using the lineage marker CD3.
(B) NKT and MAIT cells were identified from the CD3+ gate using their antigen-loaded CD1d and MR1 tetramers, respectively. CD4+, CD8+ and CD4−CD8− T cells were also identified from this population for further analysis.
(C) Expression of various markers were also examined within the CD3− population. As an example, flow plot represents expression of T-bet within the CD3− population.
(D) Identification of memory T cells. CD44 was used to identify antigen-experienced T cells. Resident memory T cells (T_{RM}) were identified based on CD103 expression. From the CD103− fraction, CD127 and CD62L expression identified central memory (T_{CM}), effector (T_E) and effector memory (T_{EM}). Subset identification of various T cell populations (ex: Tbet+CD4+ T cells) as well as activation markers (ex: PD1, CD11a) were also examined in CD4+, CD8+ and CD4−CD8− T cell subsets.
(E) Naïve T cells identified within the CD44− compartment using CD62L expression.

Figure 4. Frequency of various T cell populations in protected and unprotected CC mice following Mtb infection.

(A) Proportion of CD3+ cell, CD3+CD4+ and CD3+CD8+ T cells in the lungs of PBS and BCG vaccinated CC mice at 4 weeks post infection.

(B) Proportion of memory CD4 and CD8 T cells in the lungs of either PBS or BCG vaccinated CC mice at 4 weeks post infection, as defined by CD44+ and various combinations of CD62L and CD127 expression; CD62L+ CD127+ (central memory), CD62L− CD127− (effector), CD62L− CD127+ (effector memory).

(C) Proportion of resident memory CD4 and CD8 T cells in the lungs of either PBS or BCG vaccinated CC mice at 4 weeks post infection, as defined by CD44+ CD103+ CD69+.

(D) Proportion of Th1, Treg and Th17 cells in the lungs of either PBS or BCG vaccinated CC mice at 4 weeks post infection, as defined by CD44+ and Tbet+ (Th1), Foxp3+ (Treg) or RoRγt (Th17) expression.

(E) Proportion of CD4 and CD8 T cells expressing either CXCR3 or CX3CR1 in the lungs of either PBS or BCG vaccinated CC mice at 4 weeks post infection.

(F) PCA of the T cell phenotyping data. Each point represents the average of usually 5 mice within a given mouse strain that are either BCG vaccinated or unvaccinated. Open symbols, unvaccinated; closed symbols, BCG vaccinated. Each color represents a different mouse strain.

(A–E) The data represent the mean ± SEM from one experiment. Two-way analysis of variance with the Benjamini and Hochberg multiple comparison method. The FDR was set to 0.05 and the numbers in the figures are the q value.

Figure S2. Frequencies of non-conventional T cells in CC mice. Frequencies of MR-1-restricted MAIT cells were measured in the lungs of PBS or BCG vaccinated CC mice at 4 weeks post Mtb infection. The data in the graphs represent the mean ± SD of one experiment. Two-way analysis of variance with the Benjamini and Hochberg multiple comparsion method. The FDR was set to 0.05 and the numbers in the figures are the q value.

Figure S3. Gating strategy for the analysis of T cell function (intracellular cytokine staining).

(A) Lymphocytes were identified as using FSC and SSC properties, after which viability dyes were used to identify live cells. Singlet gates were used to exclude doublets, after which T cells were identified using the lineage marker CD3.

(B) CD4, CD8 and CD4-CD8- T cells were also identified from the CD3+ population for further analysis. Cytokine expression (ex: IFNγ) were examined in all 3 T cell sub-populations.

(C) Cytokine expression was also examined within the CD3− population.

Figure 5. Cytokine response in protected and unprotected CC strains following Mtb infection.
(A) Cells were isolated from the lungs of CC mice at 4 weeks post infection and stimulated for 24 hours with the P300 peptide pool. IFNγ, IL-2, TNF and IL-17a were measured in the supernatants. (B) and (C) Cells from the lungs of CC mice at 4 weeks post infection were stimulated for 5 hours with the P300 peptide pool. Frequencies of CD4 or CD8 T cells that expressed IFNγ, IL-2, TNF or IL-17 was determined using intracellular cytokine staining. (A–C) The data represent the mean ± SEM from one experiment. Two-way analysis of variance with Sidak's multiple comparison test was used to determine significance. *, p < 0.05; **, p < 0.01; ***, p < 0.001; ****, p < 0.0001.

Figure 6. Multivariate correlates of protection across all 24 CC strains

(A) A partial least squares regression (PLSR) model was performed to associate BCG-induced changes in lung immune features with BCG-induced changes in lung CFU. Each data point represents a vaccinated mouse that has the mean of the unvaccinated mice from the same strain subtracted from it. The scores of the top two latent variables of the model are shown, with the color of each data point representing the changed in lung CFU induced by BCG vaccination. The percents of the variance in the lung features captured by each latent variable are shown as percentages in the axes. (B) The bar graph depicts the loadings of the first latent variable of the PLSR model in (A). Positive loadings indicate immune features that increase from BCG vaccination in samples that are on the positive x-axis in the scores plot, which generally indicates they have worse protection. Negative loadings conversely indicate immune features that generally increase from BCG vaccination in samples that are protected by vaccination. (C) The heatmap illustrates all features selected to be included in the model, indicating they were important for associating strains with the lung protection continuum. Colors are representations of z-scored features, with a different color representing each 10% quantile of the data. Columns are averages of the mice in the CC strains (n=5 for most cases). Feature names are colored based on the dataset from which they come. In (B) and (C), features from the ICS dataset indicate production of the cytokines listed and no production of any measured cytokines not listed.

Figure 7. Identifying signatures of protection in subclasses of CC strains

(A), (C), (E) Scores plot (top) and heatmaps (bottom) of selected features for PLSDA models. Each dot represents a single mouse. The two colors in the scores plots signify the two classes being compared. An ellipse is drawn around the 95% confidence interval for each class. Percent of variance of the immune feature data is shown on the axes. Heatmaps show features that were most important for distinguishing the two classes. Colors are representations of z-scored features. Columns are averages of the mice in the CC strains, with vaccinated and unvaccinated mice being separated (n=5 for most cases). Feature names are colored based on the dataset from which they come (blue=T cell phenotyping, brown=lung homogenate cytokines, green=ICS, magenta= LEGENDplex). NP = non-protected, P = protected, R = resistant, S = susceptible. (B), (D), (F) PLSDA models performed as in (A), (C), and (E), except that features are the changes induced by BCG vaccination as in Figure 6.

Figure S4. Multivariate model accuracies

(A) The accuracy of our protection model (Figure 6) was compared to that of two different types of null model, where our measure of accuracy was mean squared error of prediction of Δlog (lung CFU). Our PLSR model performed significantly better than both a null model consisting of random features instead of our elastic-net chosen features and a null model where the outcomes of Δlog (lung CFU) were scrambled. Our PLSR model and each of the null models were run 50 times, and each dot represents one of these trials. Labeled p-values are from one-way ANOVAs.

(B) Our models highlighting the immune features that distinguish pairs of protection/susceptibility categories (Figure 7) were compared to two null models as in (A). Each of our models, both the ones involving all mice and the protection models were significantly better than both null models. Here, the accuracy metric is classification accuracy. Labeled p-values are from one-way ANOVAs.

Figure 8

Figure 8. Th1/17 cells correlate with BCG-mediated protection in a subset of protected CC mice.

(A) Cells were isolated from the lungs of CC mice at 4 weeks post infection and stimulated for 5 hours with either aCD3, P300 peptide pool or media control. Representative flow plots of IFNγ and IL-17 producing CD4 T cells from CC037 and C57BL/6 are shown. Data is representative of two independent experiments.

(B) Frequencies of CD4 T cells that expressed both IFNγ and IL-17 was determined using intracellular cytokine staining. Data is representative of two independent experiments. One-way analysis of variance with Sidak's multiple comparison test was used to determine significance. **, $p < 0.01$; ***, $p < 0.001$; ****, $p < 0.0001$.

(C) Representative flow plots of CD4 T cells expressing either T-bet or RORγt in CC037 or C57BL/6 mice. Data is representative of two independent experiments.

(D) Frequencies of CD4 T cells that expressed both T-bet or RORγt in CC037 or C57BL/6 mice. Data is representative of two independent experiments. One-way analysis of variance with Sidak's multiple comparison test was used to determine significance. **, $p < 0.01$; ***, $p < 0.001$.

(E) Proportion of CD4 T cells that express various combinations of IFNγ, TNF, IL-2 or IL-17 in either PBS or BCG vaccinated mice at 4 weeks post Mtb infection. Plots are generated using the SPICE platform (28).

(F) Volcano plots. Correlation between 60 immune features and the lung CFU determined for CC037 and C57BL/6 mice (from Batch 1). Dotted line, $p = 0.05$. *, $q < 0.05$, corrected with the Benjamini and Hochberg multiple comparison method (FDR = 0.05)

(G) Correlation of IFNγ⁺IL-17⁺ cells and bacterial burden in the lungs of either PBS or BCG vaccinated CC037 mice at 4 weeks post Mtb infection. Colors represent BCG vaccinated mice (red) or unvaccinated mice (blue). Spearman *r* and p value are shown.

Figure S5. Examining the Polyfunctionality of CD4 T cells in select CC strains. Boolean gating was used to identify various sub-populations amongst the CD4 T cells following P300 stimulation in certain CC mice. An increase in various IFNγ+IL-17+ CD4 T cells (that also produced TNF or IL-2) were detected in CC037, CC072, CC031 and CC001 but not in C57BL/6. In contrast, a noticeable decrease in the frequency of IFNγ "single positive" CD4 T cells was observed in all strains including C57BL/6. The data in the graphs represent the mean ± SD of two experiment. One-way analysis of variance with Sidak's multiple comparison test was used to determine significance. *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$; ****, $p < 0.0001$.