Title: Genomic effects of the glucocorticoid receptor guide the acute stress-induced delayed

anxiety and basolateral amygdala spine plasticity in rats

Running Title: GCs, NOR, and the delayed effects of acute stress

Leonardo Santana Novaes[1], Leticia Morais Bueno-de-Camargo[1], Amadeu Shigeo de

Almeida[1], Vitor Augusto Laurino Juliano[1], Nilton Barreto dos Santos[1†], Ki Ann Goosens[2],

Carolina Demarchi Munhoz[1*].

Affiliation: [1]Department of Pharmacology, Universidade de Sao Paulo Instituto de Ciencias

Biomedicas, São Paulo, Brazil, 05508-000.

†*In memorian.*

[2]Department of Psychiatry, Icahn School of Medicine at Mount Sinai, New York, NY 10029

*****Corresponding author**: Carolina Demarchi Munhoz at Department of Pharmacology, Av.

Professor Lineu Prestes, 1524, room 323. Universidade de Sao Paulo Instituto de Ciencias

Biomedicas, São Paulo - SP, 05508-000, Brazil. Phone Number: + 55 11 3091 7990.

Email: cdmunhoz@usp.br

Abstract

Anxiety, a state related to anticipatory fear, can be adaptive in the face of environmental threats or stressors. However, anxiety can also become persistent and manifest as anxiety- and stress-related disorders, such as generalized anxiety or post-traumatic stress disorder (PTSD). In rodents, systemic administration of glucocorticoids (GCs) or short-term restraint stress induces anxiety-like behaviors and dendritic branching within the basolateral complex of the amygdala (BLA) ten days later. Additionally, increased arousal-related memory retention mediated by elevated GCs requires concomitant noradrenaline (NE) signaling, both acting in the BLA. It is unknown whether GCs and NE play a role in the delayed acute stress effects on BLA dendritic plasticity and behavior. Here, we show that inhibiting corticosterone (CORT) elevation during the two-hour restraint stress prevents stress-induced increases in delayed anxiety-like behaviors and BLA dendritic spine density in rats. Also, we show that acute stress's behavioral and morphological delayed effects are critically dependent on glucocorticoid receptor (GR) genomic actions in the BLA. Unlike CORT, the pharmacological enhancement of NE signaling in the BLA was insufficient to drive delayed anxiety-related behavior. Nonetheless, the delayed anxiety-like behavior 10 days after acute stress requires NE signaling in the BLA at the time it is expressed. Therefore, we define essential roles for two stress-related hormones acting at two different times, for the late stress consequences: CORT, via GR, immediately at stress; NE, via beta-adrenoceptors, during the expression of delayed behavior.

Significance Statement

The dysregulation in orchestrating and finetuning major stress-related neural circuitries leads to enhanced reactivity and other altered ways of coping with threatening situations. These predispose humans to multiple psychiatric disorders, including anxiety and PTSD. Given the tremendous burden of affective disorders, we must advance our understanding of stress neurobiology and translate this into improved treatments. Here we showed that the absence of neuronal GR genomic signaling in the BLA prevented delayed effects on dendritic spine density and anxiety-like behavior ten days after stressor exposure. We also define essential roles for two stress-related hormones acting at two different times, for the late stress consequences: CORT, via GR, immediately at stress; NE, via beta-adrenoceptors, during the expression of delayed behavior.

Keywords: stress; neuroplasticity; amygdala; glucocorticoid signaling; anxiety; behavior.

Introduction

The body's physiological and behavioral responses to stressful stimuli have critical adaptive value as it prepares to deal with threatening conditions. The almost immediate release of norepinephrine (NE) in the brain during emotional arousal binds to alpha and beta-adrenergic receptors (1). Tens of minutes later, the release of glucocorticoid hormones (GCs) reaches a peak; GCs drive many acute adaptations to stress through actions at glucocorticoid receptors (GRs) [for a review, (2)]. A seminal work showed that elevation of corticosterone (CORT; the principal GC in rodents) enhances memory retention in tasks with high arousal content (3). Interestingly, the authors showed that arousal is set by the NE levels that reach the amygdala's basolateral complex (BLA) (3).

However, maladaptation can sometimes follow stressor exposure (4). Exacerbated fear responses to stimuli that would not usually elicit such a response, avoidance behavior, and persistent anxiety symptoms are characteristic of anxiety- and stress-related disorders, such as generalized anxiety disorder and post-traumatic stress disorder (PTSD) (5). The relationship between an acute stressor and the delayed manifestation of anxiety symptoms has been widely studied, mainly because it resembles stress-related psychiatric disorders, including PTSD.

Several studies have shown that a single two-hour stress session is sufficient to generate anxiety-like behavior ten days later (6-9). Also, both two-hour restraint stress and the single prolonged stress protocols, in addition to promoting delayed anxiety-related behavior in rats, are associated with deficits in fear extinction memory (10, 11).

Some studies have shown that either acute two-hour immobilization stress or a systemic elevation of CORT promotes, 10 to 12 days later, anxiety-like behavior in rats accompanied by increased spine density and dendritic arborization in rats BLA (7, 12, 13). This delayed effect of stress on behavior is associated with morphological changes in the BLA, suggesting that increased synaptic contacts in this brain area are the basis for the increased anxiogenic drive. Hyperexcitability of the amygdala has also been linked to anxiety (14). Thus, it seems logical that hypertrophy of BLA dendritic trees plays a causal role in anxiety.

Although there is a clear relationship between a transient increase in CORT levels and emergent anxiety-related behavior, it is unclear whether CORT acts directly upon BLA neurons to cause behavioral and morphological dendritic plasticity effects after an acute

episode of stress. It also remains to be clarified whether these morphological changes in the
BLA are sufficient to promote anxiety or if there is a dependence on another mediator, in
addition to CORT, for the delayed behavioral manifestation associated with acute stress.
Thus, in this work, we sought to better understand the role of GR, particularly the genomic
actions of this receptor, in the BLA at the time of stress in stress-induced behavioral and
morphological outcomes. Also, considering the arousal-inducing effect of catecholamines
and their modulation on neuronal excitability (15), we explored the role of the adrenergic
receptors in the BLA in expressing stress-induced anxiety-related behavior.

Materials and Methods

Subjects

Male Wistar rats (60-days old) were obtained from either the Facility for SPF Rat
Production at the Institute of Biomedical Sciences - Animal Facility Network at the
University of São Paulo (USP) or Harlan Sprague-Dawley (United States) and maintained in
the vivarium of the USP Pharmacology Department (Unit 1) or Massachusetts Institute of
Technology (MIT). Neonatal Wistars rats (1 to 4 days old) were used for the cortical
primary culture. Upon arrival, the animals were pair-housed in standard polypropylene
cages (30 X 40 X 18 cm) with ad libitum access to food and water, kept in a 12-hour light-
dark cycle (light on at 6:00 am), and under controlled temperature (23 ± 2 ºC). All
experiments were conducted under the standards of the Ethics Committee for Animal Use of
the Institute of Biomedical Sciences/University of São Paulo (CEUA - ICB 85/2016) and by
the Committee for Animal Care at MIT. All methods were in accordance with the Guidelines
of the Brazilian National Council for the Control of Animal Experimentation (CONCEA) and
the US National Institutes of Health (NIH) Guide for the Care and Use of Laboratory Animals.

Animal experimental procedures

Acute restraint stress

The animals were placed in a ventilated PVC restraint tube for 2 hours in a lighted and
soundproof room. The behavioral experiments were conducted 10 days after the end of the
stress session.

Behavioral tests

The behavioral tests were conducted in a room of soundproof and low light (10-14 lux). All
tests were videotaped using a digital video camera (Logitech C920, Lausanne, Switzerland)
from the top of the apparatuses. After each session, the apparatuses were cleaned with 5%
alcohol to remove olfactory cues.

Elevated plus maze

Animals were placed in the center of the maze (elevation of 50 cm, 50 cm × 10 cm each
arm), facing one of the open arms, and allowed to freely explore the apparatus for 5 min. We
scored entries, time spent in each arm, and the total displacement in the maze using
Ethovision software (Noldus, the Netherlands). Anxiety index (16, 17) was calculated as
follows: 1 - [(time spent in open arms/total time on the maze) + (number of entries to the
open arms/Total exploration on the maze)/2].

Light/dark box

128 Animals were placed in the middle of the light chamber (45 x 41 x 21 cm light chamber, 65
129 lx; 35 x 41 x 21 cm dark chamber, 0 lx) facing the opposite side of the connecting door. Each
130 trial lasted 5 min, and the duration of time spent in the light chamber was used as a
131 parameter to address anxiety-like behavior.

Stereotactic surgery

Drugs and viral vectors were injected into the BLA through stereotaxic surgery as described
by (18), with some modifications. Rats were anesthetized with Isoflurane (4% oxygen
saturation for induction and 2-3% for maintenance), received a dose of anti-inflammatory
for preventive analgesia (ketoprofen, 5 mg/kg, subcutaneous), and were set up in the
stereotaxic instrument (Kopf Instruments). After disinfection of the skin (70% ethanol, 1%
iodine, and 2% glycerin), the skin covering the skull was incised, and small holes were
drilled in the skull surface for the passage of cannulae or injection needles. The stainless-
steel guide cannulae (23-gauge, World Precision Instruments) were positioned bilaterally
1.0 mm dorsal to the BLA, as the following coordinates: A/P, -2.4; M/L, ± 5.1; D/V, -6.0 mm
relative to bregma and the surface of the brain (19). The cannulae were fixed with
polyacrylic cement and anchored to the skull with stainless steel screws. Drug infusions
were made through injectors (30-gauge stainless steel that extended 1.0mm beyond the tip
of each guide cannula) after the animal's complete recovery from cannula implantation

surgery (4 to 5 days). The injectors were connected to a Hamilton syringe (5 μl, Hamilton
Co., Reno, NV) through a polypropylene tube, and the infusion rate was 0.1μl/min with 1
min for diffusion before the injector was removed from the guide cannula. The viral
particles were injected with the aid of stainless-steel needles coupled to a Hamilton syringe
(5 μl, Hamilton Co, Reno, NV) fixed in an infusion pump (Pump 11 Elite, Harvard Apparatus)
coupled to the stereotaxic apparatus. The injections were placed bilaterally in the BLA,
following the coordinates: A/P, -2.4; M/L, ± 5.1; D/V, -7.0 mm. The viruses were infused at a
rate of 0.1μl/min for 20 min (total of 2μl per injection site, totaling 2 X 106 viral particles).
The needles remained in place for 10 minutes before being removed. The injections were
performed 4-5 days before stress, a critical time to reach the viral expression peak 5. Only
animals with bilateral viral infection in the BLA were considered in the behavioral analysis.
After surgery, the animals were under observation for 3 days, and if they showed any sign of
pain or discomfort, they received a daily dose of anti-inflammatory (ketoprofen, 5 mg/kg,
subcutaneous).

Cannula placement verification, viral infection visualization, and immunofluorescence

Cannula placement verification was done by Nissl staining. The expression of GFP indicated
viral infection into BLA. Animals in which the cannulae were not correctly positioned
bilaterally on the BLA, i.e., GFP expression was not restricted to the BLA, were excluded
from the study. The GFP detection and immunofluorescence assay are detailed below.

Drug administration

Metyrapone (i.p.; 150 mg in 1 ml of saline; Sigma-Aldrich) was administered 1.5 hours
before restraint stress. Bilateral intra-BLA infusion of RU-486 (10ng in 0.5 μl of 5% DMSO;
Sigma-Aldrich), dexamethasone (30 ng in 1 μl of 0.5% ethanol), propranolol (2.5 μg in 0.5 μl
of saline), or yohimbine (2.5 μg in 0.5 μl of saline), or their corresponding vehicle controls,
were also performed.

Viral vector construct

Viral vectors containing the dnGR receptor gene sequence were constructed as previously
described (20-22). The dnGR gene construct is a chimera composed of the rat GR alpha
subunit (Genbank M14053), containing the ligand-binding domain, fused to the human GR
beta subunit (Genbank X03348), which contains the DNA-binding site domain. The viral

construct was obtained from the MIT viral core facility (McGovern Institute, MIT). dnGR was

packaged in the HSV p1005 vector (short-term expression), a Herpes Simplex Virus 1 (HSV-

1) derived amplicon plasmid with an added transcription cassette expressing GFP (separate

promoter and poly(A)) (23, 24). The empty vector (lacking the dnGR sequence) was used as

a control. The target gene is driven by the IE 4/5 promoter, while a CMV promoter drives

the GFP. The two transcription cassettes are in a nose-to-tail orientation. Adherent 2.2 cells

were transfected with the plasmid of interest (containing dnGR). The transfection was done

using the Lipofectamine LTX reagent (Invitrogen). Twenty-four hours after transfection, the

cells were super-infected with Helper Virus 5dl1.2, with a multiplicity of infection = 0.03.

The cells and media were harvested 48 hours later when the virus's cytopathic effect

reached 100%. The solutions were frozen and thawed three times, sonicated to release

infectious viral particles, and then centrifuged to clear the media of cell debris. The resulting

supernatant passed onto 2-2 cells. This amplification procedure was repeated twice. After

the final sonication and centrifugation, the supernatant was purified on a sucrose gradient,

viral bands were centrifuged, and viral pellets were resuspended in 10% sucrose in D-PBST.

The vector titration was quantified in Vero cells by counting the cells expressing the GFP

protein. An average titration of approximately 1 x 106 viral particles per µl was achieved.

Dendritic spine density analysis

For some experiments, the brains were prepared using the Rapid Golgi Kit (FD

NeuroTechnologies, Inc.) according to the manufacturer's instructions. An

immunohistochemistry assay was performed for other experiments to amplify the GFP

signal to analyze the dendritic spine density in the BLA neurons previously infected with

the HSV vector. Brains were incubated in impregnation solution for 14 days followed by

incubation in cryoprotectant solution for 48 h, coronally sliced (120 µm) using a Leica

CM3050-S cryostat (Leica Biosystems) and mounted on gelatin-coated slides. Brain sections

were dehydrated in ethanol, diaphonized in xylene, and coverslipped. The BLA images were

captured in a bright field with the slide scanner Zeiss Axio Scan.Z1 (Zeiss), using a 40X

objective. A z-stack was created with a step size of 410 nm, and the images were analyzed in

ZEN 2 (Zeiss) software with a magnification of 600×. At least 5 neurons per animal (4

animals per group) were analyzed according to the following criteria: only neurons with the

soma wholly localized in the BLA, heavily impregnated, without truncated dendrites. After

neuronal soma identification, we classified each dendritic segment as primary (originating

from the soma), secondary (originating from the primary branch), tertiary (originating from

the secondary branch), or quaternary (originating from the tertiary branch). Regardless of

the morphologic classification, all protrusions originating from a dendrite were considered

spines. After we determined the origin and the end of each branch, the dendritic branch

length was calculated by the software.

Brain tissue processing for immunofluorescence and immunohistochemistry

Brains were sectioned in a cryostat (Leica CM3050-S, Leica Biosystems, Germany) in a

coronal plane (40-μm-thick) at -22 °C and stored in an anti-freeze solution (0.05 M sodium

phosphate buffer, 150 g of sucrose, and 300 ml of ethylene glycol) at -20 °C.

We performed the immunofluorescence assay to visualize GR expression in the BLA. Brain

slices were incubated in blocking solution (0.02 M potassium phosphate solution containing

0.3% X-100 triton and 1% bovine albumin) for 1 hour, followed by incubation in blocking

solution containing the primary antibody (rabbit anti-GR, dilution 1: 1000, Santa Cruz;

mouse anti-MAP-2, dilution 1: 5000, Sigma-Aldrich) for 18h (4 °C). The slices were then

incubated for 2h (room temperature) in a 0.02M potassium phosphate solution containing

the secondary donkey antibody conjugated to a fluorochrome (AlexaFluor 594 anti-rabbit,

dilution 1: 500, Thermo Fisher Scientific; AlexaFluor 488 anti-mouse, dilution 1: 500,

Thermo Fisher Scientific). Brain slices were mounted on positively charged slides

(Fisherbrand Superfroast Plus, Thermo Fisher Scientific) and covered with a coverslip using

Fluoromount-G as a mounting medium (SouthernBiotech). The images were acquired in a

confocal microscope (Nikon A1R +, Nikon Instruments Inc., Japan), sectioned in the Z plane

at 0.6 μm.

Although the HSV vector carries the gene that encodes GFP, an immunohistochemistry

procedure was performed to intensify the signal and improve GFP particles' detection

throughout the BLA dendritic tree. Brain slices were incubated in blocking solution (0.02 M

potassium phosphate solution containing 0.3% X-100 triton and 3% normal donkey serum)

followed by incubation in blocking solution containing the primary antibody anti-GFP

raised in rabbit (dilution 1: 1000; Thermo Fisher Scientific) for 18 hours (4 °C). The slices

were then incubated in a 0.02 M potassium phosphate solution containing a biotinylated

donkey antibody (dilution 1: 200; Vector) and 0.3% X-100 triton for 2 h at room

temperature. The antigen-antibody complex was visualized with the biotin-avidin technique

(Elite kit; Vector) and 3,3'-diaminobenzidine (DAB) as a chromogen through a peroxidase

reaction composed of a 2.5% solution of nickel sulfate and 0.003% hydrogen peroxide in

0.02M potassium phosphate buffer. The sections were mounted on gelatin-coated slides,

air-dried, and dipped in a 0.05% aqueous solution of osmium tetroxide for 20 s to enhance

the labeling's visibility, dehydrated, transferred into xylene, and coverslipped with DPX.

Spine density was quantified under a bright field, as detailed for the Golgi staining assay.

Western Blot

The GR expression and its phosphorylated form were analyzed by western blot as

previously (17).

Protein extraction

Cytosolic and nuclear protein extracts were obtained using the Cel-LyticTM NuCLEARTM

Extraction Kit (Sigma–Aldrich Co.). BLA tissues stored at -80 ºC were thawed and

mechanically homogenized with a conical plastic pestle (Thermo Fisher Scientific) in cold

lysis buffer (10 mM HEPES, pH 7.9, 10 mM KCl, 1.5 mM MgCl2, 0.5 mm PMSF, 1 mm DTT)

supplemented with protease and phosphatase inhibitors (Halt Protease Inhibitor Cocktail,

ThermoFisher Scientific), and centrifuged for 20 min at 11,000xg at 4 °C. The supernatant

(cytosolic fraction) was collected in a tube and stored at -80 °C. The remaining pellets were

homogenized in the extraction buffer (20 mM HEPES, pH 7.9, 300 mM NaCl, 1.5 mM MgCl2,

0.25 mM EDTA, 0.5 mM PMSF, 0.5 mM DTT 25% glycerol) supplemented with protease and

phosphatase inhibitors, kept on ice for 30 min, and centrifuged for 5 min at 20,000xg (4 °C).

The resulting supernatants (nuclear proteins) were stored at -80 °C. Protein concentration

was determined using the Bradford method (Bio-Rad Laboratories, Inc).

Electrophoresis

Protein samples (20 µg/lane) were size-separated in 10% SDS-PAGE gel (90 V) and then

blotted onto Immobilon® PVDF membrane (EMD Millipore Corporation). The Ponceau

method for immunoblot was used to ensure equal protein loading (25). Blots were blocked

with 5% bovine serum albumin (BSA) diluted in TBS-T buffer (50 mM Tris-HCl, 150 mM

NaCl, 0.1% Tween 20, pH 7.5) for 1 h at room temperature and subsequently incubated

overnight at 4 °C with specific primary antibodies: monoclonal rabbit anti-GR (dilution

1:1000, Santa Cruz) and polyclonal rabbit anti-Ser211phospho-GR (dilution 1:1000, Cell

Signaling). After incubation with the primary antibodies, the membrane was then probed

with a secondary antibody conjugated to horseradish peroxidase (dilution of 1:2000,

Kirkegaard & Perry Laboratories) for 2 h at room temperature, and the signal was obtained

by the Luminata Forte Western HRP substrate (Merck Millipore, Darmstadt, Germany)

using the ChemiDoc system (Bio-Rad Laboratories). Several exposure times were

performed to ensure the linearity of the band intensities. Each band's relative density was

normalized to the value of α-Tubulin (sc-5286, dilution: 1:20,000, Santa Cruz

Biotechnology).

Primary cortical cell culture

Neonatal rats, between 1 and 4 days post-natal, were sacrificed by decapitation, and then

the frontal cortex was dissected in a sterile medium (Hanks buffer: 160 mM NaCl, 5.3 mM

KCl, 0.44 Kh2PO4, 0.33 mM Na2HPO4, 5.5 mM glucose, 4.0 mM NaHCO3) supplemented

with antibiotics (penicillin/streptomycin, Invitrogen), at pH 7.4. The isolated frontal cortex

was maintained in Hanks buffer containing 0.05% trypsin solution (Vitrocell) and 0.001%

DNAse (Sigma-Aldrich) at 37 °C for 10 minutes and then transferred to Dulbecco's Modified

Eagle Medium culture medium (DMEN; Vitrocel) supplemented with 10% fetal bovine

serum (FBS, Vitrocel) and 2 mM glutamine (Sigma-Aldrich). The suspension was gently

mechanically grounded with the aid of a Pasteur pipette. The solution was filtered through a

70 μM membrane (Becton-Dickinson), and the cells were further cultured in a 24-well plate

and maintained at 37 °C with 5% CO2 in a humidified atmosphere (80% relative humidity)

in an incubator. For immunofluorescence assay, cells were cultured on borosilicate

coverslips covered with poly-L-lysine (0.01%, Sigma-Aldrich) in a 24-well plate. To

calculate the number of viral particles necessary for efficient infection, the number of cells

was counted before plating using the Trypan Blue exclusion method (Sigma-Aldrich, MO,

USA). The cells were maintained in DMEN supplemented with antibiotics and 10% SFB. The

medium was changed on the second and fifth days after culturing, and the cells were

maintained until the tenth day.

Infection of cells by viral particles and analysis of sub-localization and genomic activity of GR

To check the sublocalization of the dnGR receptor in the cell compartments, in response to

CORT, culture cells were infected with the viral particles (HSV-dnGR and HSV-GFP, MOI of

1) diluted in culture medium (DMEN + 2% SFB) on the eighth culture day. Forty-eight hours

later (tenth day of culture), the cells were treated with 10 µM CORT in the culture medium (DMEN + 2% SFB) for 1h. Then, the medium was removed, and the coverslips were washed with PBS and then fixed with 4% formaldehyde (Amresco) for 20 minutes at room temperature. After 5 washes with PBS, the cells adhered to the coverslips were incubated in the blocking solution (5% donkey serum, 0.05% Triton X-100 in PBS) for 2h at room temperature. Subsequently, the cells were incubated in a blocking solution containing the primary anti-GR antibody (1: 1000, Santa Cruz) for 24 hours and then, after washing with PBS, incubated in PBS containing the secondary antibody (AlexaFluor 594, 1: 2000) and 0.05% Triton X-100 for 2h. Finally, the coverslips were incubated in DAPI solution (1: 100000, Sigma-Aldrich; PBS with 0.05% Triton X-100) for 20 minutes, washed with PBS, and mounted on slides in Fluoromount-G mounting medium (SouthernBiotech, USA). The slides were analyzed under a fluorescence microscope (Nikon Eclipse 80i; Nikon Instruments Inc., Japan).

For the analysis of GR genomic activity in cells infected by tdGR, the polymerase chain reaction (PCR) assay was performed to amplify the *gilz* and *fkbp5* genes described below. The cells were obtained using the same protocol for image analysis, but the cells were cultured on 6-well plates. At the end of the treatment with CORT, the cells were washed 5 times with PBS and collected in TRIzol® (0.75 µl per sample; Thermo Fisher Scientific). RNA extraction followed the manufacturer's instructions.

Polymerase chain reaction (PCR)

The cDNA was obtained using the SuperScript III Reverse Transcriptase Kit (Thermo Fisher Scientific) according to the manufacturer's instructions. For the reaction, 12 µl of final solution were used, containing 5 µg of RNA, 1 µl of dNTPs (0.5nM each), 1 µl of Random Primer, and 5 µg of total RNA in sterile water.

For the amplification, 2 µl of the cDNA (20 ng / µl), 5 µl of Amp Buffer, 0.75 ul MgCl2, 0.5 µl of dNTPs, 1.5 µl of the forward primer, 1.5 µl of the reverse primer, 0.5 µl of Taq Polymerase (Promega Co), and Milli-Q water (Merck Millipore) sufficient for 25 µl were added in a microtube. The reaction conditions in the thermal cycler (LifeECO, Bulldog Bio, Inc) were: 95 ° C for 2 initial minutes, followed by 30 cycles at 95 ° C for 1 minute, 55 ° C for 30 seconds, and 72 ° C for 1 minute. After the last cycle, the samples remained for 5 minutes at 72 ° C. The product was separated using agarose gel electrophoresis (2%), stained with

BlueGreen (Thermo Fisher Scientific), and visualized with a UV transilluminator (ChemiDoc Imager; Bio-Rad Laboratories, Inc). The sequences of the primers used are as follows:

gapdh: forward, 5'-GGTGTGAACGGATTTGGC-3'; reverse, 5'-CTGG-AAGATGGTGATGGGTT-3';

gilz: forward, 5'-TGGGGCCTAGTAACACCAAG-3'; reverse, 5'-GAGCACACTGGCATCACATC-3'

fkbp5: forward, 5'-GAACCCAATGCTGAGCTTATG-3'; reverse, 5'-ATGTACTTGCCTCCCTTGAAG-3'.

Euthanasia of the animals and obtaining the analyzed material

Guillotine beheading, BLA dissection, and blood collection

The animals were deeply anesthetized with Isoflurane (maximum exposure of the 30s) and euthanatized by decapitation. The brain was quickly removed in a cold solution of saline-phosphate buffer (137 mM NaCl, 2.68 mM KCl, 1.27 mM KH_2PO_4, 8.06 mM Na_2HPO_4). The BLA was bilaterally dissected on a cold glass plate and frozen in a conical tube (1.5 ml) on dry ice. The brain structures were stored at –80 ºC for later use. Brains were placed in a 1 mm thick coronal cut acrylic brain matrix (Zivic Laboratories Inc.), and the dissection was performed under a stereomicroscope (Tecnival - Prolab). The animals' blood was obtained from the animal's trunk at the time of decapitation.

Transcardiac perfusion

The animals were deeply anesthetized with a solution of ketamine hydrochloride (9 mg / 100 g, i.p.) and xylazine (1 mg / 100g, i.p.) and infused transcardiacally using a peristaltic pump (Cole Parmer), with 200 ml of 0.9% saline followed by about 500 ml of 4% formaldehyde solution in sodium phosphate buffer (pH 7.4) at 4 ° C for 20 minutes. After 3 hours, the brain was removed from the skull and kept in a 30% sucrose solution in potassium phosphate buffer (0.02M) at 4 ° C for 48 hours. After this cryoprotection period, the brain was removed from the sucrose solution and frozen at -80 ° C until histological processing.

Blood collection

Blood was sampled from the body trunk after decapitation, immediately after stress termination. Plasma was obtained by adding EDTA (4.5 mM) to the collected blood and centrifugation at 4,000 g for 10 minutes. Samples were stored at -80 ºC.

 Analysis of the concentration of corticosterone in the blood

We used the Corticosterone EIA kit (Enzo Life Sciences, Inc) to analyze blood CORT levels, following the manufacturer's guidelines. Plasma samples were diluted 30 times in the assay buffer provided by the kit to fit the standard curve's optimal range. The values depicted in the figures were corrected for the dilution factor.

Statistical analysis

Statistical analysis was performed using two-way ANOVA. Post hoc Tukey's multiple comparison test examined differences between individual groups. Unpaired Student's t-test was used to compare two experimental groups. Sample size and animal numbers were estimated based on previous studies, and outlier samples were checked by the ROUT method. Quantification of data was blind for all experiments. GraphPad Prism software 9.0 was used for the statistical analysis, and the level of statistical significance was p < 0.05. Data are presented as mean ± SEM. Detailed statistical information for each test performed is reported in the figures.

Results

Systemic CORT release is necessary for the delayed stress-induced anxiety-like behavior and increased BLA dendritic spine density.

To confirm whether the acute elevation of CORT plasma levels is essential to these stress-related behavioral and neuroplastic effects, we inhibited the stress-induced rise in this hormone's blood levels by administering metyrapone (Met). This compound inhibits the synthesis of GCs, before stress exposure. We assessed the behavioral outcomes ten days later (Fig. 1*A*). As shown in Fig. 1*B*, pre-stress Met administration effectively prevented the acute stress-induced increase in CORT plasma levels measured immediately after stress exposure, as the stressed animals administered with saline presented higher plasma CORT concentrations than non-stressed animals ($P < 0.001$) or stressed animals treated with Met ($P < 0.001$).

Ten days after restraint stress, we assessed the anxiety-like behavior on an elevated plus-maze test (EPM). Likewise, pre-stress Met administration effectively blunted the occurrence of stress-induced anxiety-like behavior. Met-administered stressed animals presented higher open arms exploration than saline-treated stressed animals ($P < 0.01$, Fig. 1*C*; $P <$

0.05, Fig. 1*D*) and the same exploration profile as the saline non-stressed animals (Fig. 1*C*

and *D*). The anxiety index confirmed these data, being higher in saline stressed animals than

in non-stressed ($P < 0.01$) and Met-treated stressed animals ($P < 0.01$) (Fig. 1*E*). Finally,

neither stress nor Met treatment influenced the animals' motor behavior, indicated by the

maze's total displacement (Fig. 1*F* and *G*).

Next, we analyzed the effects of pre-stress Met administration on the delayed stress-

induced increase in dendritic spine density in the BLA using Golgi staining in a subset of the

rats that underwent behavioral testing (n = 4/group). We confirmed that a single 2-hr

restraint stress exposure augmented the spine density in the primary ($P < 0.001$, Fig. 2*A*),

secondary ($P < 0.001$, Fig. 2*B*), tertiary ($P < 0.001$, Fig. 2*C*), and quaternary ($P < 0.01$, Fig.

2*D*) segments of the BLA dendritic neurons compared to the non-stressed animals. Met

administration blunted the stress-induced increase in dendritic spine density in the BLA

with no effects in non-stressed animals in all dendritic segments (Fig. 2), suggesting a

coordinated role of stress-induced systemic CORT elevation at the time of stress exposure in

both BLA neuroplasticity and the late development of anxiety-like behavior.

The delayed effects of acute stress on anxiety-like behavior and BLA dendritic spine density increase depend on GR's neuronal genomic activity.

Next, we examined whether neuronal GR activation in BLA was implicated in the late stress

outcomes associated with acute stress-induced plasma CORT elevation. Two hours of

restraint stress increased GR nuclear translocation into BLA neurons immediately after

stress (Fig. S1A and *B*), and Western blot analysis revealed not only an increase in GR

nuclear translocation in the BLA ($P = 0.0346$; Fig. S1*C*) but also in the phosphorylation at the

Ser211 site ($P = 0.0166$; Fig. S1*D*), known to be associated with a gain in GR transcriptional

function (26, 27). Notably, ten days after stress and immediately after the EPM test, there

was no difference in the GR nuclear translocation in the BLA of the stressed animals (Fig.

S2). Likewise, in a previous study, we found no changes in serum CORT levels of animals

submitted to EPM previously stressed 10 days before (8).

These findings collectively suggest that CORT signaling through GR at the time of stress

ultimately triggers delayed changes in the BLA. In line with this, animals infused with the

GR antagonist RU486 directly into the BLA immediately prior to stress (Fig. S3*A*) showed,

ten days later, a higher percentage of entries ($P = 0.0161$, Fig. S3*B*) and time spent ($P =$

0.0215; Fig. S3*C*) in the EPM open arms, with no changes in the total distance traveled in the

maze (Fig. S3*D*), when compared to stressed animals that received vehicle infusions.

A rise in CORT serum levels leads, via GR, to rapid (non-genomic) and long-lasting

(genomic) changes in the BLA neuronal excitability and synaptic plasticity (28, 29). Because

of the delayed effect of stress-associated GR signaling on anxiety-like behavior and

remodeling of dendritic spines in the BLA, we expressed a transcriptionally inactive,

truncated form of GR in the BLA, the so-called dominant negative GR (dnGR). Cultured cells

infected with dnGR responded to CORT treatment with the receptor's translocation to the

cell nucleus as control GFP-infected cells (Fig. S4*A* and *B*). However, even with a high GR

expression in the nucleus, dnGR-infected cells expressed fewer GR-related mRNA

transcription products when compared to GFP-infected cells (Fig. S4*C*).

The HSV-based plasmid has a peak of transgene expression three to four days after

infection, with a subsequent rapid decline in transgene expression (20). Thus, we applied

restraint stress four to five days after intra-cranial delivery of the virus (Fig. 3*A*) to

maximize any inhibition of GR genomic activity during the stress session. Notably, the

behavioral test was administered ten days later, a time point at which viral expression of

the transgene should be minimal. Stressed GFP-infected animals exhibited anxiety-like

behavior, as shown by reduced entries ($P < 0.01$; Fig. 3*B*) and time spent ($P < 0.01$; Fig. 3*C*)

in open arms compared to the GFP non-stressed animals. On the other hand, the expression

of dnGR in the BLA prevented the anxiogenic effect of acute stress. These animals, even

when stressed, displayed similar levels of open arm exploration compared to the GFP non-

stressed and significantly elevated open arm exploration relative to the GFP stressed

animals. This situation was observed for open arm entries ($P < 0.05$) and the anxiety index

($P < 0.05$) (Fig. 3*B* and *D*). There were no effects of stress, and dnGR expression on distance

traveled in the maze, suggesting differences in motor activity did not contribute to the

behavioral difference (Fig. 3*E* and *F*).

Additionally, dnGR expression did not impact the release of CORT during acute stress:

similar stress-induced levels of CORT were observed in dnGR-Stress and GFP-stress groups

(Fig 3*G)* ($P < 0.001$ and $P < 0.01$, respectively) and exhibited similar hormone levels relative

to each other. Thus, the blunting of the delayed behavioral consequences of stress by dnGR

is not due to altered CORT release.

To determine whether dnGR expression also prevented stress-induced changes in BLA
spine density, we explored the GFP protein signal throughout all subcellular compartments
in brain sections taken from the animals after the EPM and performed the analysis of spine
density. Fig. 4 shows a stress-induced increase in the dendritic spine density in animals
previously infected with the GFP control construct, compared to non-stressed animals, in
the primary (P < 0.05; Fig. 4A), secondary (P < 0.05; Fig. 4B), and tertiary (P < 0.001; Fig. 4C)
dendritic segments. The overexpression of dnGR in the BLA prevented the stress-induced
increase in the density of spines in the secondary (P < 0.05; Fig. 4B) and tertiary (P < 0.001;
Fig. 4C) but not in the primary (Fig. 4A), dendritic segments.

An increase in GR activity in the BLA is sufficient to induce a delayed anxiogenic
effect

To test whether acute elevation of CORT in the BLA was sufficient to induce delayed anxiety,
we stereotaxically injected dexamethasone (Dex) into the BLA and, ten days later, tested the
anxiety-related behavior (Fig. S5A). The EPM test showed that Dex infusion lowered the
percentage of open-arms entries (P = 0.0482; Fig. S5B) and time spent in the open arms (P =
0.0285; Fig. S5C), and elevated anxiety index (P = 0.0127; Fig. S5D), compared to saline-
infused stressed animals. We performed a second anxiety-related behavioral test since the
Dex-treated animals moved less in the maze than the saline ones (P = 0.0048; Fig. S5E and
F). Using the light/dark box test, we confirmed that Dex-treated animals displayed more
remarkable anxiety-like behavior than saline-treated animals since Dex-treated animals
spent less time in the light compartment than saline-administered animals (P = 0.0369; Fig.
S5G).

BLA beta-adrenergic receptor activation is necessary for the expression of delayed
stress-related anxiety-like behavior

The acute anxiogenic effect of NE is widely known and involves the noradrenergic
projections from locus coeruleus (LC) to BLA (30). However, it is not known whether NE
has delayed anxiogenic effects. We next addressed this question by infusing the alpha-2
adrenergic receptor antagonist yohimbine into the BLA and examining the acute and
delayed effects of this manipulation on anxiety-like behavior (Fig. S6A). Antagonizing alpha-
2 autoreceptors in the BLA results in increased firing from noradrenergic nerve terminals
(12, 13). Consistent with previous reports, we observed that increased activity of NE in BLA

had an acute anxiogenic effect in the open field test (Fig S6*B, C, D, E*). In contrast with CORT,

intra-BLA administration of yohimbine was insufficient to induce anxiety-like behavior in

the EPM test ten days later (Fig. S6*F, G, H, I*).

Because increasing NE activity in the BLA has an acute, but not delayed, anxiogenic effect,

we next wanted to examine whether beta-adrenergic receptor activity is critical for the

delayed manifestation of anxiety-like behavior in previously stressed animals. Thus, we

administered saline or propranolol, a beta-adrenoceptor antagonist, into the BLA of the

animals before the behavioral test, 10 days after the stress (Fig. 5*A*). As shown in Fig. 5,

stressed propranolol-treated animals presented a higher percentage of entries ($P < 0.05$;

Fig. 5*B*) and time spent ($P < 0.05$; Fig. 5C) in the EPM than stressed saline-administered

animals. These effects were not accompanied by changes in the animals' locomotor activity

(Fig. 5*D*). Therefore, these results suggest that even at a delayed time point at which stress-

dependent increases in spine density are present (Fig. 2), stress-related changes in anxiety

still depend on the activation of beta-receptors in the BLA during the behavioral task.

Discussion

Here, we report that inhibition of CORT synthesis during acute stress prevented stress-

related behavioral and morphological effects that emerged after ten days. This finding

complements a previous study reporting that single systemic administration of CORT,

simulating an acute stress exposure, induced a delayed anxiogenic effect, along with parallel

increases in dendritic length, branch points, and spine density in BLA neurons (12, 13). It is

important to note that we showed that inhibiting CORT alone (Fig. 1-2), but not

catecholamines (Fig. 5, S6), was sufficient to hinder these effects of stress. Together with

previous reports in the literature (12, 31), in this study, we have shown that CORT release is

essential for the late effects of stress on anxiety-like behavior and dendritic remodeling of

the BLA.

Amygdala hyperactivity is considered a biological substrate of stress- and anxiety-related

disorders in human and animal models (32). Stimulation of BLA projection neurons triggers

anxiety-like behavior in rodents (33), while acute and chronic stress increases the intrinsic

excitability of BLA pyramidal neurons (34, 35). Likewise, symptoms of anxiety manifested

by patients suffering from stress-related disorders, such as PTSD, are correlated with

amygdala hyperactivity (36, 37). The greater concentration of synaptic contacts in dendritic

spines, compared to the rest of the neuronal surface, confers heightened ionic conductance
in this area, leading to a higher probability of evoked excitatory postsynaptic potentials
[EPSPs, see (38)]. Likely, delayed morphological changes in the BLA dendritic tree following
acute restraint stress produce a more excitable neuronal phenotype, although this remains
untested. Accordingly, the delayed anxiogenic effect of single prolonged stress seen in rats
co-occurs with hypertrophy in the BLA dendritic tree and increased miniature excitatory
and inhibitory postsynaptic frequency currents in this brain area (39). Curiously, in the
present study, the lack of GR genomic signaling in the BLA neurons prevented the stress-
related increase in the spine density in the most distal dendritic branches (Fig. 4B, C)
without affecting the most proximal ones (Fig. 4A). This effect is consistent with other
reports that distal synaptic inputs on BLA neurons are more likely to be excitatory. In
contrast, proximal synapses are more likely to be inhibitory (40).

Acute elevations in blood CORT levels have a modulatory action on the turnover of dendritic
spines in the cortex, producing a short-term effect on the elimination and formation of new
dendritic spines (41). Interestingly, acute stress and a single systematic injection of CORT
promote delayed (at least 10 days later) but not short-term (1 day later) changes in the BLA
dendritic tree (7, 31).

Protein synthesis is a critical step for the formation of new spines (42), and our findings not
only showed that the stress-related anxiety-like behavior and the increase in the BLA
dendritic spine density are both governed by the direct action of CORT in this brain region,
but that they are phenomena dependent on GR genomic activity. The GR mobilization by
dexamethasone injection into BLA generated a delayed anxiety-like behavior even without
stress. The mechanism by which GR influences the origin and stabilization of new spines is
not yet fully understood. However, growing evidence has shown a synergistic and
interdependent action between GR and the TRKB-BDNF receptor in the motor cortex during
motor-related learning (43-45).

The role of NE in modulating behavior is well known and is closely related to the immediate
effects on changing/increasing the perception of threats under a stress exposure (46). Acute
stress leads to a rapid release of NE in the limbic system, including BLA and hippocampus,
primarily via projections of noradrenergic neurons from the LC (30, 47). The activation of
beta-adrenergic receptors in the BLA promotes increased excitatory synaptic transmission

and increased neuronal spike firing rates [see (15)]. In the present study, we confirmed an
acute anxiogenic effect of increased availability of NE in the BLA (via yohimbine
administration) (Fig. S6), but this increase in NE release was insufficient to promote
delayed behavioral changes. In contrast, Dex administration in the BLA was sufficient to
promote delayed behavioral changes (Fig. S5). Furthermore, blocking NE signaling directly
into BLA with propranolol at the time of behavioral testing prevented the anxious-like
manifestation in previously stressed animals (Fig. 7).

The synergism of GR and noradrenergic transmission in the BLA has already been critical in
potentiating the effects of arousal on memory retention (48). Here, we reported a non-
simultaneous but coordinated and (possibly) non-direct dependence of the delayed
manifestation of stress-induced anxiety on both GR and beta-adrenoceptor signaling in the
BLA. It is important to note that previously we did not see any increase in CORT release ten
days after the stress and immediately after the EPM test (8), suggesting no role for elevated
CORT in the expression of anxiety-like behavior. Thus, we propose that the stress-related
delayed plastic changes in the BLA's dendritic arbor depend on GR signaling, and these
changes may potentiate the stress response to NE. Therefore, given the morphological
effects of stress, the delayed behavioral outcome depends on NE signaling via beta-
adrenoceptors.

Acknowledgments

We gratefully thank Guiomar Wiesel for her technical assistance.

Funding

This work was supported by the Fundação de Amparo à Pesquisa do Estado de São Paulo
(FAPESP 2016/03572-3 and 2019/00908-9) and Conselho Nacional de Desenvolvimento
Científico e Tecnológico (CNPq 422523/2016-0) to CDM and a grant from the US Army
Research Office and the Defense Advanced Research Projects Agency (grant W911NF-10-1-
0059) to KAG. LSN was supported by FAPESP (2012/24002-0 and 2018/19599-3). LMBC
was supported by CAPES, CNPq (142260/2019-3), and FAPESP (2018/15982-7). NBS was
supported by FAPESP (2018/17991-3). ASA was supported by FAPESP (2019/00400-5 and
2021/12488-4) and CAPES. VALJ was supported by FAPESP (2020/06914-8 and
2021/13524-4) and CNPq (PIBIC 2019-1429). C.D.M. is a research fellow from CNPq. This
study was financed in part by the Coordenação de Aperfeiçoamento de Pessoal de Nível
Superior - Brasil (CAPES) - Finance Code 001.

Competing Interest Statement

The authors declared no conflicts of interest concerning their authorship or the publication
of this article.

Author Contributions:

LSN: designed research, performed research, analyzed data, wrote the paper; LMBC:
performed research, analyzed data; ASA: performed research, analyzed data; NBS:
performed research, analyzed data; KAG: designed research, contributed unpublished
reagents/analytic tools, wrote the paper; CDM: designed research, analyzed data,
contributed unpublished reagents/analytic tools, wrote the paper.

References

1.   H. J. Krugers, H. Karst, M. Joels, Interactions between noradrenaline and
corticosteroids in the brain: from electrical activity to cognitive performance. *Front
Cell Neurosci* **6**, 15 (2012).
2.   E. R. de Kloet, M. Joels, F. Holsboer, Stress and the brain: from adaptation to disease.
Nature reviews. Neuroscience **6**, 463-475 (2005).

3.   B. Roozendaal, S. Okuda, E. A. Van der Zee, J. L. McGaugh, Glucocorticoid
enhancement of memory requires arousal-induced noradrenergic activation in the
basolateral amygdala. *Proceedings of the National Academy of Sciences of the United*
States of America **103**, 6741-6746 (2006).

4.   B. S. McEwen, H. Akil, Revisiting the Stress Concept: Implications for Affective
Disorders. *The Journal of neuroscience : the official journal of the Society for*
Neuroscience **40**, 12-21 (2020).

5.   D. A. Regier, E. A. Kuhl, D. J. Kupfer, The DSM-5: Classification and criteria changes.
World Psychiatry **12**, 92-98 (2013).

6.   P. Chakraborty, S. Datta, B. S. McEwen, S. Chattarji, Corticosterone after acute stress
prevents the delayed effects on the amygdala. *Neuropsychopharmacology : official*
publication of the American College of Neuropsychopharmacology **45**, 2139-2146
(2020).

7.   R. Mitra, S. Jadhav, B. S. McEwen, A. Vyas, S. Chattarji, Stress duration modulates the
spatiotemporal patterns of spine formation in the basolateral amygdala. *Proceedings*
of the National Academy of Sciences of the United States of America **102**, 9371-9376
(2005).

8.   L. S. Novaes, N. B. Dos Santos, J. G. Perfetto, K. A. Goosens, C. D. Munhoz,
Environmental enrichment prevents acute restraint stress-induced anxiety-related
behavior but not changes in basolateral amygdala spine density.
Psychoneuroendocrinology **98**, 6-10 (2018).

9.   F. Yasmin *et al.*, Stress-induced modulation of endocannabinoid signaling leads to
delayed strengthening of synaptic connectivity in the amygdala. *Proceedings of the*
National Academy of Sciences of the United States of America **117**, 650-655 (2020).

10.   P. R. Abdullahi *et al.*, Protective effects of morphine in a rat model of post-traumatic
stress disorder: Role of hypothalamic-pituitary-adrenal axis and beta- adrenergic
system. *Behavioural brain research* **395**, 112867 (2020).

11.   L. S. Novaes, L. M. Bueno-de-Camargo, C. D. Munhoz, Environmental enrichment
prevents the late effect of acute stress-induced fear extinction deficit: the role of
hippocampal AMPA-GluA1 phosphorylation. *Transl Psychiatry* **11**, 18 (2021).

12.   R. Mitra, R. M. Sapolsky, Acute corticosterone treatment is sufficient to induce
anxiety and amygdaloid dendritic hypertrophy. *Proceedings of the National Academy*
of Sciences of the United States of America **105**, 5573-5578 (2008).

13.   R. P. Rao, S. Anilkumar, B. S. McEwen, S. Chattarji, Glucocorticoids protect against the
delayed behavioral and cellular effects of acute stress on the amygdala. *Biological*
psychiatry **72**, 466-475 (2012).

14.   N. Daviu, M. R. Bruchas, B. Moghaddam, C. Sandi, A. Beyeler, Neurobiological links
between stress and anxiety. *Neurobiol Stress* **11**, 100191 (2019).

15.   A. E. Fink, J. E. LeDoux, beta-Adrenergic enhancement of neuronal excitability in the
lateral amygdala is developmentally gated. *J Neurophysiol* **119**, 1658-1664 (2018).

16.   H. Cohen, M. A. Matar, D. Buskila, Z. Kaplan, J. Zohar, Early post-stressor intervention
with high-dose corticosterone attenuates posttraumatic stress response in an
animal model of posttraumatic stress disorder. *Biological psychiatry* **64**, 708-717
(2008).

17.   L. S. Novaes *et al.*, Environmental enrichment protects against stress-induced
anxiety: Role of glucocorticoid receptor, ERK, and CREB signaling in the basolateral
amygdala. *Neuropharmacology* **113**, 457-466 (2017).

18.   L. S. Novaes, S. J. Shammah-Lagnado, Projections from the anteroventral part of the
641   medial amygdaloid nucleus in the rat. *Brain research* **1421**, 30-43 (2011).

642   19.   G. Paxinos, C. Watson, *The Rat Brain in Stereotaxic Coordinates.* (Elsevier, London,
643   ed. 6th Edition, 2007), pp. 476.

644   20.   D. Kaufer *et al.*, Restructuring the neuronal stress response with anti-glucocorticoid
645   gene delivery. *Nature neuroscience* **7**, 947-953 (2004).

646   21.   E. D. Kirby *et al.*, Basolateral amygdala regulation of adult hippocampal
647   neurogenesis and fear-related activation of newborn neurons. *Molecular psychiatry*
648   **17**, 527-536 (2012).

649   22.   R. M. Meyer, A. Burgos-Robles, E. Liu, S. S. Correia, K. A. Goosens, A ghrelin-growth
650   hormone axis drives stress-induced vulnerability to enhanced fear. *Molecular*
651   *psychiatry* **19**, 1284-1294 (2014).

652   23.   R. L. Neve, F. Lim, Generation of high-titer defective HSV-1 vectors. *Current protocols*
653   *in neuroscience / editorial board, Jacqueline N. Crawley ... [et al.]* **Chapter 4**, Unit 4 13
654   (2013).

655   24.   S. J. Russo *et al.*, Nuclear factor kappa B signaling regulates neuronal morphology
656   and cocaine reward. *The Journal of neuroscience : the official journal of the Society for*
657   *Neuroscience* **29**, 3529-3537 (2009).

658   25.   O. Salinovich, R. C. Montelaro, Reversible staining and peptide mapping of proteins
659   transferred to nitrocellulose after separation by sodium dodecylsulfate-
660   polyacrylamide gel electrophoresis. *Analytical biochemistry* **156**, 341-347 (1986).

661   26.   W. Chen *et al.*, Glucocorticoid receptor phosphorylation differentially affects target
662   gene expression. *Mol Endocrinol* **22**, 1754-1766 (2008).

663   27.   M. D. Krstic, I. Rogatsky, K. R. Yamamoto, M. J. Garabedian, Mitogen-activated and
664   cyclin-dependent protein kinases selectively and differentially modulate
665   transcriptional enhancement by the glucocorticoid receptor. *Mol Cell Biol* **17**, 3947-
666   3954 (1997).

667   28.   S. Duvarci, D. Pare, Glucocorticoids enhance the excitability of principal basolateral
668   amygdala neurons. *The Journal of neuroscience : the official journal of the Society for*
669   *Neuroscience* **27**, 4482-4491 (2007).

670   29.   H. Karst, S. Berger, G. Erdmann, G. Schutz, M. Joels, Metaplasticity of amygdalar
671   responses to the stress hormone corticosterone. *Proceedings of the National*
672   *Academy of Sciences of the United States of America* **107**, 14449-14454 (2010).

673   30.   J. G. McCall *et al.*, Locus coeruleus to basolateral amygdala noradrenergic projections
674   promote anxiety-like behavior. *Elife* **6** (2017).

675   31.   H. Kim *et al.*, Regional differences in acute corticosterone-induced dendritic
676   remodeling in the rat brain and their behavioral consequences. *BMC Neurosci* **15**, 65
677   (2014).

678   32.   B. M. Sharp, Basolateral amygdala and stress-induced hyperexcitability affect
679   motivated behaviors and addiction. *Transl Psychiatry* **7**, e1194 (2017).

680   33.   G. G. Calhoon, K. M. Tye, Resolving the neural circuits of anxiety. *Nature neuroscience*
681   **18**, 1394-1404 (2015).

682   34.   A. Hetzel, J. A. Rosenkranz, Distinct effects of repeated restraint stress on basolateral
683   amygdala neuronal membrane properties in resilient adolescent and adult rats.
684   *Neuropsychopharmacology : official publication of the American College of*
685   *Neuropsychopharmacology* **39**, 2114-2130 (2014).

686   35.   J. A. Rosenkranz, E. R. Venheim, M. Padival, Chronic stress causes amygdala
687   hyperexcitability in rodents. *Biological psychiatry* **67**, 1128-1136 (2010).

36. A. M. Hull, Neuroimaging findings in post-traumatic stress disorder. Systematic
review. *Br J Psychiatry* **181**, 102-110 (2002).
37. R. K. Pitman *et al.*, Biological studies of post-traumatic stress disorder. *Nature*
reviews. Neuroscience **13**, 769-787 (2012).
38. A. T. Gulledge, N. T. Carnevale, G. J. Stuart, Electrical advantages of dendritic spines.
PloS one **7**, e36007 (2012).
39. H. H. Zhang *et al.*, Traumatic Stress Produces Delayed Alterations of Synaptic
Plasticity in Basolateral Amygdala. *Front Psychol* **10**, 2394 (2019).
40. P. M. Klenowski *et al.*, Structural and functional characterization of dendritic arbors
and GABAergic synaptic inputs on interneurons and principal cells in the rat
basolateral amygdala. *J Neurophysiol* **114**, 942-957 (2015).
41. C. Liston, W. B. Gan, Glucocorticoids are critical regulators of dendritic spine
development and plasticity in vivo. *Proceedings of the National Academy of Sciences*
of the United States of America **108**, 16074-16079 (2011).
42. J. Tanaka *et al.*, Protein synthesis and neurotrophin-dependent structural plasticity
of single dendritic spines. *Science* **319**, 1683-1687 (2008).
43. M. Arango-Lievano *et al.*, Persistence of learning-induced synapses depends on
neurotrophic priming of glucocorticoid receptors. *Proceedings of the National*
Academy of Sciences of the United States of America **116**, 13097-13106 (2019).
44. M. Arango-Lievano *et al.*, Neurotrophic-priming of glucocorticoid receptor signaling
is essential for neuronal plasticity to stress and antidepressant treatment.
Proceedings of the National Academy of Sciences of the United States of America **112**,
15737-15742 (2015).
45. F. Jeanneteau, M. J. Garabedian, M. V. Chao, Activation of Trk neurotrophin receptors
by glucocorticoids provides a neuroprotective effect. *Proceedings of the National*
Academy of Sciences of the United States of America **105**, 4862-4867 (2008).
46. G. Aston-Jones, J. Rajkowski, J. Cohen, Role of locus coeruleus in attention and
behavioral flexibility. *Biological psychiatry* **46**, 1309-1320 (1999).
47. R. J. Valentino, E. Van Bockstaele, Convergent regulation of locus coeruleus activity
as an adaptive response to stress. *European journal of pharmacology* **583**, 194-203
(2008).
48. D. de Quervain, L. Schwabe, B. Roozendaal, Stress, glucocorticoids and memory:
implications for treating fear-related disorders. *Nature reviews. Neuroscience* **18**, 7-
19 (2017).

Figure 1. CORT synthesis inhibition during stress prevented the delayed emergence
of anxiety-like behavior. (A) Schematic representation of the experimental design
showing the timeline of vehicle (saline) or metyrapone (Met) injections, the stress
submission, and the EPM behavioral test. One cohort of animals was euthanized
immediately after the end of the stress session. (B) Previous administration of Met

prevented the stress-induced increase in CORT plasma levels measured immediately after
stress (n = 6–7 animals for each group, two-way ANOVA: stress $F_{1, 21}$ = 40.34, P < 0.0001;
Met $F_{1, 21}$ = 22.81, P = 0.0001; stress-Met interaction $F_{1, 21}$ = 44.35, P < 0.0001). (C and D)
Stressed animals treated with saline exhibited reduced open arms entries (C; n = 10–12
animals for each group, two-way ANOVA: stress $F_{1, 40}$ = 4.692, P = 0.0363; Met $F_{1, 40}$ =
3.23, P = 0.0799; stress-Met interaction $F_{1, 40}$ = 8.9, P = 0.0048) and time spent (D; n = 10–
12 animals for each group, two-way ANOVA: stress $F_{1, 39}$ = 8.712, P = 0.0053; Met $F_{1, 39}$ =
5.783, P = 0.0210; stress-Met interaction $F_{1, 39}$ = 2.352, P = 0.1332) in the EPM test
compared to saline-treated non-stressed animals and to both stressed and non-stressed
Met-treated animals. (E) Stress increased the anxiety index in saline-treated but not in Met-
treated rats (n = 10–12 animals for each group, two-way ANOVA: stress $F_{1, 39}$ = 6.85, P =
0.0126; Met $F_{1, 39}$ = 3.786, P = 0.0589; stress-Met interaction $F_{1, 39}$ = 9.282, P = 0.0041).
(F) Neither stress nor Met administration influenced the animal's locomotor activity in the
EPM (n = 9–12 animals for each group, two-way ANOVA: stress $F_{1, 34}$ = 2.388, P = 0.1315;
Met $F_{1, 34}$ = 3.78, P = 0.0602; stress-Met interaction $F_{1, 34}$ = 0.1839, P = 0.6707). (G)
Representative tracking plots for the average motion of each experimental group in the
EPM. Results are represented as mean ± SEM. Tukey's multiple comparisons post-hoc test.
Significance differences between groups are indicated as * P < 0.05; ** P < 0.01; and *** P <
0.001.

Figure 2. Pre-stress Met administration prevented the stress-induced increase in BLA dendritic spine density. (A–D) Stressed saline-treated animals showed an increase in spine density, effect prevented by pre-stress Met administration, in the primary (A; n = 15–20 neurons for each group, two-way ANOVA: stress $F_{1, 69}$ = 9.351, P = 0.0032; Met $F_{1, 69}$ = 7.786, P = 0.0079; stress-Met interaction $F_{1, 69}$ = 7.72, P = 0.0070), secondary (B; n = 15–20 neurons for each group, two-way ANOVA: stress $F_{1, 69}$ = 9.871, P = 0.0025; Met $F_{1, 69}$ = 20.65, P < 0.0001; stress-Met interaction $F_{1, 69}$ = 8.895, P = 0.0038), tertiary (C; n = 15–20 neurons for each group, two-way ANOVA: stress $F_{1, 69}$ = 42.17, P < 0.0001; Met $F_{1, 69}$ = 39.75, P < 0.0001; stress-Met interaction $F_{1, 69}$ = 9.809, P = 0.0025), and quaternary (D; n = 14–20 neurons for each group, two-way ANOVA: stress $F_{1, 65}$ = 18.62, P < 0.0001; Met $F_{1, 65}$ = 15.92, P = 0.0002; stress-Met interaction $F_{1, 65}$ = 6.267, P = 0.0148) dendrite segments compared to non-stressed animals. (Top, left) Representative photomicrography of a BLA's secondary dendrite segment. (Scale bar, 10 μm). Results are represented as mean ± SEM.

763   Tukey's multiple comparisons post-hoc test). Significance differences between groups are

764   indicated   as   **   P   <   0.01   and   ***   P   <   0.001.

Figure 3. dnGR overexpression in the BLA prevented the delayed emergence of stress-induced anxiety-like behavior. (A, left) Schematic representation of the experimental design shows the virus injection timeline into BLA (GFP or dnGR), the stress submission, and the EPM behavioral test. A cohort of animals was euthanized immediately after the end of the stress session. (Right) Representative figure showing the viral injection into BLA and photomicrography of an infected BLA (GFP, green). (Scale bar, 100 μm). CeA: central

nucleus of the amygdala. (B) GFP stressed animals exhibited reduced open arm entries in

the EPM test compared to GFP non-stressed animals and to both stressed and non-stressed

animals overexpressing dnGR in the BLA (n = 10–12 animals for each group, two-way

ANOVA: stress F1, 39 = 7.527, P = 0.0091; dnGR F1, 39 = 2.612, P = 0.1114; stress-dnGR

interaction F1, 39 = 6.724, P = 0.0133). (C) GFP stressed animals spent less time in the EPM

open arms compared to GFP non-stressed animals. dnGR animals (stressed and non-

stressed) exhibited no differences in time spent in open arms compared to GFP non-

stressed animals (n = 10–12 animals for each group, two-way ANOVA: stress F1, 39 = 6.595,

P = 0.0142; dnGR F1, 39 = 0.6368, P = 0.4297; stress-dnGR interaction F1, 39 = 8.864, P =

0.0050). (D) Stress increased the anxiety index in GFP-expressing but not in dnGR-

expressing rats (n = 10–12 animals for each group, two-way ANOVA: stress F1, 39 = 7.943, P

= 0.0075; dnGR F1, 39 = 2.06, P = 0.1592; stress-dnGR interaction F1, 39 = 8.047, P =

0.0072). (E) Neither stress nor BLA virus infection influenced the animal's locomotor

activity in the EPM (n = 10–12 animals for each group, two-way ANOVA: stress F1, 42 =

0.212, P = 0.6476; dnGR F1, 42 = 2.257, P = 0.1405; stress-dnGR interaction F1, 42 = 2.283,

P = 0.1383). (F) Representative tracking plots for the average motion of each experimental

group in the EPM. (G) Acute restraint stress promotes, immediately after the stressor

stimulus, an increase in CORT plasma levels in both GFP- and dnGR-overexpressing animals

(n = 6–9 animals for each group, two-way ANOVA: stress F1, 25 = 36.25, P < 0.0001; dnGR

F1, 25 = 0.007222, P = 0.9330; stress-dnGR interaction F1, 25 = 0,07801, P = 0.7823).

Results are represented as mean ± SEM. Tukey's multiple comparisons post-hoc test.

Significance differences between groups are indicated as * P < 0.05, ** P < 0.01, and *** P <

0.001.

Figure 4. dnGR expression in the BLA prevented the stress-related increase in BLA's dendritic spine density. (A–C) GFP stressed animals showed an increase in spine density, effect prevented by the dnGR overexpressing in the BLA, in the primary (A; n = 19–27 neurons for each group, two-way ANOVA: stress $F_{1, 83}$ = 17.32, P < 0.0001; dnGR $F_{1, 83}$ = 0.188, P = 0.6657; stress-dnGR interaction $F_{1, 83}$ = 0.02408, P = 0.8771), secondary (B; n = 19–27 neurons for each group, two-way ANOVA: stress $F_{1, 85}$ = 10.49, P = 0.0017; dnGR $F_{1, 85}$ = 10.34, P = 0.0018; stress-dnGR interaction $F_{1, 85}$ = 0.6021, P = 0.4399), and tertiary (C; n = 15–25 neurons for each group, two-way ANOVA: stress $F_{1, 69}$ = 16.8, P = 0.0001; dnGR

$F_{1, 69} = 8.421$, $P = 0.0050$; stress-dnGR interaction $F_{1, 69} = 7.85$, $P = 0.0066$) dendrite

segments compared to non-stressed animals. (Top, left) Representative photomicrography

of a BLA's secondary dendrite segment. (Scale bar, 10 µm). Results are represented as mean

± SEM. Tukey's multiple comparisons post-hoc test. Significance differences between groups

are   indicated   as   * P <   0.05,   ** P <   0.01,   and   *** P <   0.001.

Figure 5. Pre-EPM test propranolol administration in the BLA prevented the delayed
manifestation of the stress-induced anxiety-like behavior. (A, left) Schematic
representation of the experimental design shows the propranolol or saline injection
timeline into BLA, the stress submission, and the EPM test. (Right) Representative figure
and photomicrography showing a representative location of cannulae and injection needle
tips into BLA. (B–E) Stressed, intra-BLA saline administered animals exhibited reduced
open arms entries (B; n = 8 animals per group, two-tailed Student's t-test: P = 0.0441) and
time spent (C; n = 8 animals per group, two-tailed Student's t-test: P = 0.0218) in the EPM
test comparing to stressed, intra-BLA propranolol administered animals. (D) There was no
influence of intra-BLA propranolol administration on the animal's locomotor activity in the
EPM (n = 8-9 animals per group, two-tailed Student's t-test: P = 0.0285). (E) Representative
tracking plots for the average motion of each experimental group in the EPM. Results are
represented as mean ± SEM. Significance differences between groups are indicated as * P <
0.05.

 Supplementary Figures and Tables

Figure S1. Acute restraint stress affects GR nuclear translocation and GRser211
phosphorylation in the BLA. (A and B) BLA immunofluorescence showing increased
accumulation of GR (red) in the nucleus of neurons (green, MAP-2) from stressed animals
(B) compared to non-stressed animals immediately after the cessation of stress (A). DAPI
(blue) was used as a nuclear marker. Arrowheads indicate the nuclear region of neurons.
(Scale bars on smaller and larger panels correspond to 25 μm and 8 μm, respectively).
Images were acquired in a confocal microscope sectioned in the Z plane at 0.6 μm. (C and D)
Western blot analysis showed an increase in GR nuclear translocation (C; n = 4-5 for each

group, two-tailed student's t-test: P = 0.0346), as well as in GRser211 phosphorylation (D; n

= 3 for each group, two-tailed student's t-test: P = 0.0166) in stressed animals compared to

unstressed ones. (Bottom, right) Representative autoradiography of Western blot. Results

are represented as mean ± SEM. Significance differences between groups are indicated as *

P　　　　　　　　　　　　　　　　<　　　　　　　　　　　　　　　　0.05.

Figure S2. GR expression in the BLA 10 days after the stress. Acute restraint stress has no
delayed effect on GR nuclear translocation in the BLA. (Left) Representative
autoradiography of Western blot. (A–C) Western blot analysis showed no effect of stress on
nuclear GR (A; n = 5 per group, two-tailed student's t-test: P = 0.7837), cytosolic GR (B; n = 5
per group, two-tailed student's t-test: P = 0.9408), or the nuclear-cytosolic ratio (C; n = 5
per group, two-tailed student's t-test: P = 0.7571). Results are represented as mean ± SEM.

Figure S3. Pre-stress intra-BLA GR antagonist RU-486 administration prevented the
delayed emergence of stress-induced anxiety-like behavior. (A) Schematic representation of
the experimental design showing the timeline of RU-486 or saline injection into BLA, the
stress submission, and the EPM behavioral test. (B–D) Pre-stress intra-BLA saline
administered animals exhibited reduced open arms entries (B; n = 12 animals per group,
two-tailed student's t-test: P = 0.0161) and time spent (C; n = 12 animals per group, two-
tailed student's t-test: P = 0.0215) in the EPM test comparing to RU-486 administered
animals. (D) There was no influence of intra-BLA RU-486 administration on the animal's
locomotor activity in the EPM (n = 12 animals per group, two-tailed student's t-test: P =
0.6168). Results are represented as mean ± SEM. Significance differences between groups
are   indicated   as   *   P   <   0.05.

Figure S4. dnGR- and GFP-infected cells response to the challenge with CORT. (A and B)
Primary culture immunofluorescence of rat cortex showing GR (red) distribution in the
cytoplasmic and nuclear compartments of neurons. Neurons were infected with GFP (A) or
dnGR (B) and stimulated with CORT (10 μM, 1h). DAPI was used as a nuclear marker (Blue),
and GFP is represented in green. (Scale bars on smaller and larger panels correspond to 30
μm and 10 μm, respectively). (C) RT-PCR cDNA amplicons for GILZ, FKBP5, and GAPDH in
GFP-   and   dnGR-infected   cells   treated   with   CORT   (10   μM,   1h).

Figure S5. Intra-BLA dexamethasone administration promotes an anxiety-like behavior ten

days after infusion. (A, left) Schematic representation of the experimental design showing

the timeline of dexamethasone (Dex) or saline injection into BLA, the stress submission, and

the behavioral tests. (Right) Representative figure and photomicrography showing a

representative location of cannulae and injection needle tips into BLA. (Scale bar, 100 μm);

(B–D) Intra-BLA dexamethasone administered animals exhibited reduced open arms

entries (B; n = 8 animals per group, two-tailed student′s t-test: P = 0.0482) and time spent

(C; n = 7-8 animals per group, two-tailed student′s t-test: P = 0.0285), as well as increased

anxiety index (D; n = 7-8 animals per group, two-tailed student′s t-test: P = 0.0127) in the

EPM test comparing to saline administered animals. (E) Intra-BLA dexamethasone

promoted a delayed reduced motor activity in the EPM test (n = 8 animals per group, two-

tailed student's t-test: P = 0.0048). (F) Intra-BLA dexamethasone administration promoted

a delayed decrease in time spent in the lit compartment compared to saline administered

animals in the light/dark (L/D) test (n = 7 animals per group, two-tailed student's t-test: P =

0.0369). (G) Representative tracking plots for the average motion of each experimental

group in the EPM. Results are represented as mean ± SEM. Significance differences between

groups   are   indicated   as   *   P   <   0.05.

Figure S6. Intra-BLA yohimbine administration promotes an acute, but not delayed,

anxiety-like behavior. (A, left) Schematic representation of the experimental design showing

the timeline of yohimbine or saline injection into BLA and the behavioral tests. (Right)

Representative figure and photomicrography showing a representative location of cannulae

and injection needle tips in the BLA. (B-C) Animals that received intra-BLA yohimbine

exhibited reduced time spent (B; n = 7 animals per group, two-tailed student's t-test: P =

0.0225) and entries (C; n = 7 animals per group, two-tailed student's t-test: P = 0.0234) in

the center area of the OF. (D) Intra-BLA yohimbine promoted an acute decrease in motor

activity in the OF (n = 7 animals per group, two-tailed student's t-test: P = 0.00462). (E)

Representative tracking plots for the average motion of each experimental group in the OF.

(F-G) There were no differences in the open arms entries (F; n = 7-8 animals per group,

two-tailed student's t-test: P = 0.9058) and time spent (G; n = 7-8 animals per group, two-

tailed student's t-test: P = 0.8491) in the EPM test of BLA yohimbine administered animals

comparing to saline administered animals. (H) Intra-BLA yohimbine did not influence the

animal's locomotor activity in the EPM (n = 7-8 animals per group, two-tailed student's t-

test: P = 0.1617). (I) Representative tracking plots for the average motion of each

experimental group in the EPM. Results are represented as mean ± SEM. Significance

differences between groups are indicated as * P < 0.05.