Monkey dorsolateral prefrontal cortex represents abstract visual sequences during a no-report

task

Authors: Nadira Yusif Rodriguez[1,2], Theresa H. McKim [1^], Debaleena Basu[1], Aarit Ahuja[1,2], and

Theresa M. Desrochers[1,2,3]*

Affiliations:

1. Department of Neuroscience, Brown University

2. Robert J. and Nancy D. Carney Institute for Brain Science

^ current address: Department of Child and Adolescent Psychiatry, Psychosomatics and

Psychotherapy, University Hospital Würzburg, Germany

3. Department of Psychiatry and Human Behavior

* Corresponding Author: Theresa M. Desrochers, theresa_desrochers@brown.edu

Abbreviated title: Monkey DLPFC represents abstract visual sequences

Pages: 57

Figures: 8 main text, 2 extended data

Tables: 8

Words:

Abstract: 231

Introduction: 823

Discussion: 1,554

Conflict of Interest Statement: The authors declare no competing financial interests.

Acknowledgements: We thank Matthew Maestri for his assistance with animal training and data

collection. We would also like to thank Dr. Michael Worden, Lynn Fanella, Fabienne McEleney

and Brown University's MRI Facilities staff for their support and guidance throughout this

project. We also thank Dr. David Sheinberg, Dr. Amitai Shenhav, Dr. Katherine Conen, and Dr.

Diana Burk for their continued support throughout this project and preparation of this

publication, as well as members of both the Sheinberg and the Desrochers Labs for many helpful

discussions and contributions. This study was supported by the NSF-EPSCoR Neural Basis of

Attention 1632738 (N.Y.R, T.M.D.), the NIGMS-NIH Initiative to Maximize Student

Development IMSD R25GM083270 (N.Y.R.), the NIH-NIGMS COBRE P20GM103645-07, and

the Carney Institute for Brain Science Innovation Award (T.M.D.). Part of this research was

conducted using computational resources and services at the Center for Computation and

Visualization, Brown University, S10 OD016366.

Abstract

Monitoring sequential information is an essential component of our daily lives. Many of these

sequences are abstract, in that they do not depend on the individual stimuli, but do depend on an

ordered set of rules (e.g., chop then stir when cooking). Despite the ubiquity and utility of

abstract sequential monitoring, little is known about its neural mechanisms. Human rostrolateral

prefrontal cortex (RLPFC) exhibits specific increases in neural activity (i.e., "ramping") during

abstract sequences. Monkey dorsolateral prefrontal cortex (DLPFC) has been shown to represent

sequential information in motor (not abstract) sequence tasks, and contains a sub-region, area 46,

with homologous functional connectivity to human RLPFC. To test the prediction that area 46

may represent abstract sequence information, and do so with parallel dynamics to those found in

humans, we conducted functional magnetic resonance imaging (fMRI) in monkeys. When

monkeys performed no-report abstract sequence viewing, we found that left and right area 46

responded to abstract sequential changes. Interestingly, responses to rule and number changes

overlapped in right area 46 and left area 46 exhibited responses to abstract sequence rules with

changes in ramping activation, similar to that observed in humans. Together, these results

indicate that monkey DLPFC monitors abstract visual sequential information, potentially with a

preference for different dynamics in the two hemispheres. More generally, these results show

that abstract sequences are represented in functionally homologous regions across monkeys and

humans.

Significance Statement

Daily, we complete sequences that are "abstract" because they depend on an ordered set of rules (e.g., chop then stir when cooking) rather than the identity of individual items. Little is known about how the brain tracks, or monitors, this abstract sequential information. Based on previous human work showing abstract sequence related dynamics in an analogous area, we tested if monkey dorsolateral prefrontal cortex (DLPFC), specifically area 46, represents abstract sequential information using awake monkey fMRI. We found that area 46 responded to abstract sequence changes, with a preference for more general responses on the right and dynamics similar to humans on the left. These results suggest that abstract sequences are represented in functionally homologous regions across monkeys and humans.

Introduction

Sequential tasks that require monitoring are prevalent in daily life. For example, taking a bus requires tracking familiar sequences of buildings (e.g., three houses then a library), enabling the detection of deviations from this sequence (e.g., if there is a detour). Similarly, many cognitive processes occur serially, and often demand that we maintain an internal representation of the previous steps to complete the next. Even in tasks that are not explicitly sequential, a system for tracking transitions between steps, such as when completing a mathematical operation, may be essential.

Sequence monitoring is this active process of tracking the order of subsequent "states" or steps. Monitoring is distinct from other well-studied sequence processes, such as explicit memorization, or potentially more automatic behaviors, such as a series of motor outputs (e.g., playing the piano) or statistical sequence learning (Desrochers et al., 2019) . Such sequential processes likely contain monitoring operations within them but are also comprised of other cognitive computations. *Abstract sequences* are sequences that are not dependent on the individual stimuli but can instead be described by the rule they follow (e.g., three same, one different or AAAB) (Desrochers et al., 2022). Therefore, *abstract sequence monitoring* entails sequences of sensory stimuli that possess abstract structure and active monitoring of this structure. While it may be assumed that abstract sequence monitoring underlies many aforementioned sequence types (including motor sequences), it is rarely studied in isolation and the neural underpinnings of abstract sequence monitoring remain largely unknown.

Multiple modes of evidence in humans indicate that activity in rostrolateral prefrontal cortex (RLPFC) is crucial to sequence monitoring (Desrochers et al., 2015, 2019; McKim and

Desrochers, 2022). Functional magnetic resonance imaging (fMRI) revealed systematic

increasing activity ("ramping") from the beginning to the end of each sequence in human

RLPFC. Across studies, this activity occurs either bilaterally, or in the left RLPFC. Further,

noninvasive transcranial magnetic stimulation (TMS) showed that the left RLPFC was necessary

for sequential tasks in humans. Other studies have also demonstrated the involvement of RLPFC

as part of a frontoparietal network active during complex sequential tasks (Farooqui et al., 2012;

Wang et al., 2019; Wen et al., 2020). While consistent with the findings discussed above, these

studies frequently involve other cognitive phenomena, like decision-making, leaving open their

specific role of sequence monitoring.

Studies in nonhuman primates also suggest a role of lateral prefrontal cortex in abstract visual

sequence monitoring. Motor sequence studies show that neurons in the dorsolateral prefrontal

cortex (DLPFC) are selective for serial position (Barone and Joseph, 1989; Averbeck et al.,

2006; Shima et al., 2007; Berdyyeva and Olson, 2010) and sequence boundaries (Fujii and

Graybiel, 2003), and include neural dynamics that could underlie the ramping observed in human

BOLD activation(Desrochers et al., 2015, 2019; McKim and Desrochers, 2022). Neurons in the

DLPFC also show ordinal selectivity during visual object sequences (Ninokura et al., 2004;

Warden and Miller, 2010; Naya et al., 2017). A rich literature also supports the involvement of

DLPFC in representing non-sequential abstract rules (Hoshi et al., 1998; White and Wise, 1999;

Wallis et al., 2001; Eiselt and Nieder, 2013). Responses in the DLPFC can also selectively

represent sequential regularities (Vergnieux and Vogels, 2020). Together, these physiological

studies suggest that the monkey DLPFC is well-positioned to monitor abstract visual sequences.

We hypothesized that a specific sub-region of monkey DLPFC (area 46) monitors visual abstract

sequential information. In humans, abstract sequence monitoring has been localized to the

RLPFC, which is distinct from the rostromedial prefrontal cortex (Koechlin et al., 2000; Burgess

et al., 2003; Gilbert et al., 2010; Moayedi et al., 2015; Henssen et al., 2016; Du et al., 2020).

While rostromedial prefrontal cortex has similar connectivity in monkeys and humans,

anatomical evidence suggests that monkey area 46 contains the most similar connectivity

patterns to human RLPFC (Sallet et al., 2013; Neubert et al., 2014), and overlapping high-level

visual representations (Xu et al., 2022). Therefore, we predicted that abstract visual sequence

monitoring would be supported by monkey area 46, and that similar ramping dynamics, as

observed in humans, would localize to this same area.

To directly test these predictions, we conducted event-related fMRI in awake nonhuman

primates, and used deviations from established abstract visual sequences during a no-report task

to index abstract sequence monitoring. We found that nonhuman primate DLPFC distinctly

represents abstract visual sequence information, independent from other task constraints.

Additionally, we find that these abstract sequences elicit ramping dynamics similar to those

observed in humans during abstract sequence performance. Intriguingly, deviant responses with

differing primary dynamics were observed in the two hemispheres: an onset-based signal on the

right, and ramping on the left. These findings indicate that a specific sub-region of DLPFC

preferentially supports abstract sequence monitoring in monkeys and may be functionally

homologous to humans. Further, these results establish an important connection between human

and monkey complex cognition and the neural substrates that mediate it, providing a foundation

for understanding more complex behaviors across species in the future.

Materials and Methods

Subjects

We tested three adult male rhesus macaques (ages spanning 6-12 years during data collection, 9-14 kg). All procedures followed the NIH Guide for Care and Use of Laboratory Animals and were approved by Institutional Animal Care and Use Committee (IACUC) at Brown University.

No-Report Abstract Visual Sequence Task

All visual stimuli used in this study were displayed using an OpenGL-based software system developed by Dr. David Sheinberg at Brown University. The experimental task was controlled by a QNX real-time operating system using a state machine. Eye position was monitored using video eye tracking (Eyelink 1000, SR Research). Stimuli were displayed at the scanner on a 24-inch BOLDscreen flat-panel display (Cambridge Systems). The general design of the visual sequence paradigm was based on a similar auditory sequence task (Wang et al., 2015).

Stimuli

Each image presentation consisted of fractal stimulus (approximately 8° visual angle) with varying colors and features. Fractals were generated using MATLAB for each scanning session using custom scripts based on stimuli from (Kim and Hikosaka, 2013) following the instructions outlined in (Miyashita et al., 1991). For each scan session, new, luminance matched, fractal sets were generated. All stimuli were presented on a gray background, with a fixation spot that was always present on the screen superimposed on the images. To provide behavioral feedback, the fixation spot was yellow when the monkey was successfully maintaining fixation and red if the monkey was not fixating. Stimuli were displayed for 0.1 to 0.3 s each, depending on the sequence type and timing template, detailed as follows.

Sequence Types

There are five sequence types in this task (**Figure 1**): habituation sequences and four deviant

sequence types. Across these sequence types, there were a total of nine different timing templates

used. These templates were included to counterbalance for stimulus and sequence duration

across the sequence types and to provide a greater variety of sequential timings during

habituation. The inter-sequence interval was jittered to decorrelate across timing templates (mean

2 s, 0.25-8 s).

Habituation Sequences

Habituation sequences were composed of images drawn from a pool of four possible fractals. We

will refer to the habituation image pool as [A, B, C, D]. Sequences were composed from these

images in one of two possible rules: three the same, one different (e.g., AAAB, DDDC) and four

the same (e.g., AAAA, CCCC). All sequences contained four images and followed one of three

possible general timings based on the total duration of the sequence: short (1.1 s), medium (1.7

s), and long (2.3 s). Each total sequence duration, in turn, had two possible timing templates

within it, one with longer stimulus durations and one with shorter stimulus durations: short 0.1 s

and 0.2 s, medium 0.1 s and 0.3 s, long 0.2 s and 0.3 s. Inter-stimulus intervals were arranged to

evenly space the four stimulus presentations within the total sequence duration.

Deviant Sequences

Deviant sequences were composed of images drawn from a different pool of three possible

fractals. We will refer to the deviant image pool as [E, F, G]. All deviant images were displayed

for 0.2 s, regardless of deviant type. Across deviant types, the total sequence durations were

matched to the short, medium, and long habituation timing templates. There were four deviant

types, detailed as follows:

New Items, Same Rule (NISR): These deviants use images that come from the deviant pool of

images, but do not differ from the habituation rule. For example, if the habituation rule was three

the same, one different then NISR sequences would follow the same rule with new images (e.g.,

GGGF and FFFE) Alternatively, if the habituation rule was four same, an example NISR would

be EEEE. All sequences were four items and had a total duration of 1.7 s.

Rule deviants: These deviants do not follow the same rule as habituation, but instead follow the

alternate rule. If the habituation rule was three the same, one different, example deviants would

follow the four the same rule, e.g., EEEE and GGGG. All deviants contained four images and

had a total sequence duration of 1.7 s, the same as medium habituation sequences.

Number Deviants: These deviants follow the same rule as habituation but contain a different

number of images (either two or six). If the habituation rule was three the same, one different,

example deviants would be EG and FFFFFE. Two-item deviants had a total sequence duration of

1.1 s, the same as short habituation sequences, and six-item deviants had a total sequence

duration of 2.3 s, the same as long habituation sequences.

Double Deviants: These deviants combine Rule and Number deviant types. If the habituation

rule was three the same, one different, example deviants would be EE and GGGGGG. The

timing was the same as number deviants.

Block Structure

Each block contained 30 sequences and lasted approximately 112 s on average. Habituation blocks contained equal numbers of the six possible timing templates (two of each: short, medium, and long). Habituation sequences were presented in pseudo-random order such that a sequence could not begin with the same fractal as the final fractal of the previous sequence. Deviant blocks were composed of 24 habituation sequences and six deviant sequences. All deviant sequences within a block were of the same sequence type. The six deviant sequences were pseudo-randomly interspersed throughout the block such that deviant sequences did not occur in the first 6 sequences of the block (to avoid block initiation confounds), and deviant sequences were not presented consecutively to each other. If deviant sequences contained a variable number of items (i.e., number deviants and double deviants), then an equal number of two- and six-item sequences were included within a single block. The 24 habituation sequences within deviant blocks were presented in the same manner as in habituation blocks (i.e., evenly distributed timing templates and avoiding between-sequence fractal image repeats).

Run Structure

Each run was composed of five blocks, interleaved with 14 s fixation blocks (**Figure 1**). The first block of each run contained only habituation sequences. The four subsequent blocks were one of each of the four possible deviant types, with their order counterbalanced across runs. The same habituation rule was used for the entirety of a single run. Runs lasted approximately 10.5 min. The sequence rule (*three same, one different* or *four same*) used for each run was counterbalanced across each scanning session so as to have an equal number of runs for each rule. Monkeys typically completed 4-8 runs of this task (among other tasks not reported on here) in a single scanning session (one day).

Runs were initiated according to the monkey's fixation behavior to ensure that the monkey was

not moving and engaged in the task before acquiring functional images. During this pre-scan

period, a fixation spot was presented. Once the monkey successfully acquired this fixation spot

and received approximately four liquid rewards (12 – 16 s), functional image acquisition and the

first habituation block were initiated.

Reward

The timing of liquid rewards was not contingent upon sequential events, only on the monkey

maintaining fixation. Rewards were delivered on a graduated schedule such that the longer the

monkey maintained fixation, the more frequent rewards were administered (Leite et al., 2002).

The first reward was given after 4 s of continuous fixation. After two consecutive rewards of the

same fixation duration, the fixation duration required to obtain reward was decreased by 0.5 s.

The minimum duration between rewards that the monkey could obtain was 0.5 s. Fixation had to

be maintained within a small window (typically 3° of visual angle) around the fixation spot to

not break fixation. The only exception was a brief time window (0.32 s) provided for blinks. If

the monkey's eyes left the fixation window and returned within that time window, it would not

trigger a fixation break. If fixation was broken, the reward schedule would restart at the

maximum 4 s duration required to obtain reward.

FMRI Data Acquisition

Monkeys were trained to sit in the "sphynx" position in a custom MR-safe primate chair

(Applied Prototype, Franklin, MA or custom-made by Brown University). The monkey's head

was restrained from moving via a plastic "post" (PEEK, Applied Prototype, Franklin, MA)

affixed to the monkeys' head and the primate chair. Monkeys were habituated to contrast agent

injection procedures, recorded MRI sounds, wearing earplugs (Mack's Soft Moldable Silicone

Putty Ear Plugs, Kid's size), and transportation to the scanner prior to MRI scanning sessions.

Monkeys were trained on the behavioral task with different images that were never used during

scanning.

Prior to each scanning session, monkeys were intravenously injected with a contrast agent:

monocrystalline iron oxide nanoparticle (MION, Feraheme (ferumoxytol), AMAG

Pharmaceuticals, Inc., Waltham, MA, 30 mg per mL or BioPal Molday ION, Biophysics Assay

Lab Inc., Worcester, MA, 30 mg per mL). MION to improves the contrast-to-noise ratio ~3-fold

(Vanduffel et al., 2001; Leite et al., 2002) and enhances spatial selectivity of MR signal changes

(Zhao et al., 2006). MION was injected, approximately 30-60 min before scanning, into the

saphenous vein below the knee (7 mg/kg), then flushed with a volume of sterile saline

approximately double the volume of the MION injected. No additional MION was added during

scanning, as MION has a long blood half-life (15.3 +/- 3.5 hr) (Leite et al., 2002).

A Siemens 3T PRISMA MRI system with a custom six-channel surface coil (ScanMed, Omaha,

NE) at the Brown University MRI Research Facility was used for whole-brain imaging.

Anatomical scans consisted of a T1-MPRAGE (repetition time, TR, 2700 ms; echo time, TE,

3.16 ms; flip angle, 9°; 208 sagittal slices; 0.5 x 0.5 x 0.5 mm), a T2 anatomical (TR, 3200 ms;

TE 410 ms; variable flip angle; 192 interleaved transversal slices; 0.4 x 0.4 x 0.4 mm), and an

additional high resolution T2 anatomical (TR, 8020 ms; TE 44 ms; flip angle, 122°; 30

interleaved transversal slices; 0.4 x 0.4 x 1.2 mm). Functional images were acquired using a fat-

saturated gradient-echoplanar sequence (TR, 1.8 s; TE, 15 ms; flip angle, 80°; 40 interleaved

axial slices; 1.1 x 1.1 x 1.1 mm).

The target sample size (number of runs per monkey) was calculated using pilot data from a

previous version of this task not included in the current data set. A region of interest was

constructed from a cluster of deviant > NISR activation and the number of runs calculated for a

significant effect in this region (using the beta values of the onset GLM, see below) at 80%

power and alpha = 0.05 (G-Power). Guided by this power analysis and similar studies (Wang et

al., 2015), we estimated a total of 200 runs across the three animals would be necessary.

FMRI Data Analysis

The majority of the following analyses were performed in Matlab using SPM 12 (http://www.fil.

Ion.ucl.ac.uk/spm). Prior to analysis, data were preprocessed using the following steps:

reorienting (to ensure proper assignment of the x,y,z planes), motion correction (realignment),

normalization, and spatial smoothing (2 mm isotropic Gaussian kernel separately for gray matter

and white matter). All steps were performed on individual runs separately. The T1-MPRAGE

anatomical image was skull stripped using FSL BET brain extraction tool

(http://www.fmrib.ox.ac.uk/fsl/) to facilitate normalization. All images were normalized to the

112-RM SL macaque atlas (McLaren et al., 2009).

Runs were included for analysis only if they met the following criteria: the monkey had to be

performing well and a sufficient number of acquisition volumes within the run had to pass data

quality checks. The monkey's performance was evaluated by calculating the percentage of time

within a run that fixation was maintained. Runs were excluded if the monkey was fixating < 80%

of the time (similar criteria as in (Vanduffel et al., 2001; Leite et al., 2002; Wang et al., 2015).

Approximately 20% of runs were excluded due to poor fixation: 10% from monkey J, 3% from

monkey W and 7% from monkey B. To evaluate data quality, we used the ART toolbox (Artifact

Detection Tools, https://www.nitrc.org/projects/artifact_detect) to detect outlier volumes. Any

volumes that had motion greater than one voxel (1.1 mm) in any direction were excluded. Any

run with greater than 12% of volumes excluded was excluded from analysis (0% runs excluded

for monkey J, 0.5% of runs excluded for monkey W, and 15% of runs excluded for monkey B).

Runs with poor image quality due to artifact or banding to pre-process or analyze were also

excluded. These accounted for 2% of the data for monkey J, 5% for monkey W, and 0.5% for

monkey B. After applying these criteria, a total of 232 runs (average of 340 volumes per run for

all animals, 93 sessions in total across animals) were included for analysis from 3 monkeys:

monkey W: 97 runs (32 sessions); monkey J: 65 runs (32 sessions), and monkey B: 70 runs (32

sessions).

Models

Within-subject statistical models were constructed under the assumptions of the general linear

model (GLM) in SPM 12 for each pseudo-subject bin. For all models, data were binned into

approximately 10-run pseudo-subject bins. Each bin contained data from only one monkey. Runs

were pseudo-randomly assigned to bins to balance the number of runs which followed each of

the two sequential rules (three same one different or four same) and the distribution of runs from

earlier and later scanning sessions. Condition regressors were all convolved with a gamma

function (shape parameter = 1.55, scale parameter = 0.022727) to model the MION

hemodynamic response function (Vanduffel and Farivar, 2014). The first six sequences in a run

and reward times were included as nuisance conditions. Additional nuisance regressors were

included for the six motion estimate parameters (translation and rotation), outlier volumes, and

image variability (standard deviation of within run image movement variability, calculated using

the ART toolbox). Outlier volumes were determined using the ART toolbox (standard global

mean; global signal detection outlier detection threshold = 4.5; motion threshold = 1.1mm; scan

to scan motion and global signal change for outlier detection) and one additional regressor with a

"1" at only that volume was included for each volume to be "scrubbed".

Regressors were estimated using a bin-specific fixed-effects model. Whole-brain estimates of

bin-specific effects were entered into second-level analyses that treated bin as a random effect.

One-sample t-tests (contrast value vs zero, p < 0.005) were used to assess significance. These

effects were corrected for multiple comparisons when examining whole-brain group voxelwise

effects using extent thresholds at the cluster level to yield false discovery rate (FDR) error

correction (p < 0.05). Group contrasts were rendered on an inflated MNI canonical brain using

Caret (Van Essen et al., 2001). Prior to selecting GLM's we used the model assessment,

comparison, and selection toolbox (MACS, https://github.com/JoramSoch/MACS, (Soch and

Allefeld, 2018) to determine models that would be the best fit. Three GLMs were applied to the

data as follows:

Onsets Model: To assess the univariate effects of deviant sequences, we constructed a model

using instantaneous stimulus onset regressors for the first item in each sequence with the

following nine condition regressors for different sequence types: short, medium, and long

habituation sequence timing templates; NISR; rule deviants; two- and six-item number deviants;

and two- and six-item rule and number deviants.

Parametric Last Item versus Unique Ramp Model: To directly test whether variance could be

better accounted for by a phasic response at the last item in the sequence or ramping activation,

we constructed a pair of models to allow last item and ramp regressors to compete for variance

within the same model. Onset regressors were constructed with an instantaneous stimulus onset

regressor at each position in the sequence with the same nine condition regressors for the

different sequence types as in the Onsets Model: short, medium, and long habituation sequence

timing templates; NISR; rule deviants; two- and six-item number deviants; and two- and six-item

rule and number deviants. Including an onset at each position effectively modeled sustained

activation throughout the sequence and enabled the inclusion of the following parametric

regressors. The last item parametric was added as ones at the first sequence positions and an

arbitrarily larger value (6) at the last item. The ramp parametric was entered as the sequence

position (1-4, 1-2, or 1-6) for each sequence. Parametric regressors were implemented

hierarchically in the GLM. Therefore, variance explained by the last parametric regressor (in this

case, ramping), is above and beyond what could be explained by the onsets or last item

regressors.

Parametric Ramp versus Unique Last Item Model: This second model of the pair sought to

identify variance uniquely explained by the last item regressor, above and beyond variance

explained by the onsets or ramping regressors. All other aspects of the model were the same as

the unique ramp model above.

<u>ROI Analyses</u>

The primary bilateral regions of interest were constructed from the coordinates of a seed region

centered in macaque monkey area 46d. These coordinates were determined, using diffusion

weighted and functional MRI, to be most similar to the lateral portion of human area 10 (Gilbert

et al., 2010; Sallet et al., 2013). Human lateral area 10 overlaps with areas of ramping activation

observed in human RLPFC in previous studies (Desrochers et al., 2015, 2019; McKim and

Desrochers, 2022). A 5 mm sphere was created around the center coordinate for the seed region

in macaque Montreal Neurological Institute (MNI) space. The sphere was then transformed into

112RM-SL space using RheMap (Sirmpilatze & Klink, 2020, resulting in a sphere centered at

xyz = 12.7, 32.6, 22.5 in 112RM-SL space. For identification of brain areas we also utilized the

NIMH Macaque Template (NMT v02, Macaque Atlas, Jung et al., 2021; Seidlitz et al., 2018).

Additional ROIs were constructed with the explicit purpose of comparing nearby regions in

DLPFC that were significant clusters of activation last item versus ramping models. Specifically,

the significant left DLPFC cluster of activation for Unique Ramp, Rule Deviants > NISR in the

unique ramp model (center xyz = -12.2, 36, 23) and the significant left DLPFC cluster of

activation for Unique Last, Rule Deviants > NISR in the unique last item model (center xyz = -

12.3, 42.9, 21.8) were taken for comparison.

To compare activation within and across ROIs in a manner that controlled for variance, we

extracted t-values from the condition of interest over baseline using the Marsbar toolbox (Jean-

Baptiste Poline, 2002). T-values (one for each pseudo-subject bin, n = 22 bins) were entered into

RM-ANOVAs with the identity of the monkey entered as a covariate.

<u>Results</u>

Three monkeys (*macaca mulatta*) performed no-report abstract sequence viewing while

undergoing awake fMRI scanning. The monkeys were trained to fixate on a central spot while

viewing a stream of fractal images arranged into four-item visual sequences (based on Wang, et

al., **Figure 1**). This task did not require responses, only fixation, and thus was termed "no-

report". The task was performed in runs (~10 min each), that each contained five blocks. For

each run, the first block habituated animals to one of two possible sequential rules AAAB, or

AAAA (A and B represent different images drawn from a pool of four possible images; 30

sequences in total per block). Habituation sequences each had one of six possible timing

templates to balance stimulus and sequence durations across sequence types. Each subsequent

block contained rare deviants (6 of the 30 sequence repetitions per block) of one of the following

four possible types: new images following the same rule, number deviants (2 or 6 items), rule

deviants (e.g., AAAA), or double deviants. All deviant images were drawn from a separate three-

image pool. The five total blocks were interleaved with 16 s fixation blocks. To encourage

animals to maintain fixation throughout, reward was administered on a graduated schedule not

correlated with sequence presentation: the longer they maintained fixation, the shorter the

duration between rewards. Reward was thus decorrelated from the four-item visual sequences. A

total of 232 runs were analyzed (97 monkey W, 65 monkey J, 70 monkey B). Monkeys

performed the task well and fixated for 95% of the time in included runs (see Methods for those

excluded).

Figure 1. No-report abstract sequence viewing task. A. Schematic representation of human rostrolateral

prefrontal cortex (RLPFC; left) and monkey dorsolateral prefrontal cortex (DLPFC; right) depicting the main

questions that were the focus of this study: Does monkey DLPFC monitor abstract sequences, as shown by

responses to deviant sequences? and does monkey DLPFC exhibit ramping activation, as found in human RLPFC

during sequence monitoring? **B.** Monkeys only fixate throughout runs. Scanning is performed in the "sphynx"

position. **C.** Example partial habituation block for sequence rule *three same, one different* (AAAB). **D.** Example

stimulus pools (top) show a set of images that would be used in a single scanning session. New images are used each session. Six possible timing templates for habituation sequences (bottom, left) and deviant sequences (bottom, right) illustrated with gray rectangles indicating single images. Total sequence durations are listed for each template type. **E.** Examples of the five sequence types if the sequence rule in use is *three same, one different*. **F.** Example run, with each bar indicating one multi-image sequence: four images in habituation, new items same rule (NISR), and rule deviants; two or six images in number and double deviants. The first block contains only habituation sequences and subsequent blocks contain only one of the four deviant types. Sequence blocks alternate with fixation blocks. Blue water droplets schematize reward delivery, which is decoupled from sequence viewing and delivered on a graduated schedule based on the duration the monkey has maintained fixation.

Monkey DLPFC represents changes in abstract visual sequences

Our first goal was to test whether area 46 differentially responds when there is a change in the abstract visual sequence. Because this task is no-report, we examined this question using neural responses (BOLD) to deviant sequences. Previous work has shown that such deviant responses disappear with inattention and are robust in brain areas processing sequence related information (Bekinschtein et al., 2009; Dehaene et al., 2015). These results suggest that deviant neural responses indicate that individuals are attending to the sequences, even in the absence of a report. Therefore, we used neural responses to rare deviant sequences to indicate awareness of changes to an established abstract sequence, as is in similar auditory tasks (Uhrig et al., 2014; Wang et al., 2015). To specifically query the responses to abstract sequence changes, we could not simply compare habituated sequences to deviant sequences, as the deviant sequences were composed from a different pool of images than the habituation sequences, and any differences observed between habituation and deviant sequences could have resulted from differences in image identity. Therefore, to specifically examine changes in abstract sequence structure, we compared new items of the same rule (NISR) to number deviants and rule deviants. All images in this

comparison were drawn from the same pool of (deviant) images. Double deviants were not

included in analyses because of the inability to dissociate between changes due to rule and

number.

We first constructed an unbiased region of interest (ROI) for monkey area 46 in each hemisphere

to compare activity between rule and number deviants and NISR. Monkey area 46 has many

potential functional subdivisions (Gerbella et al., 2010, 2013; Borra et al., 2011, 2019; Saleem et

al., 2014); therefore, we created a 5 mm sphere centered on a seed region identified as having the

most similar connectivity with human RLPFC in monkey diffusion and functional MRI (Sallet et

al., 2013, center xyz = 12.7, 32.6, 22.5 in area 46d, see Methods). The resulting sphere spanned a

small region of area 46 that encompassed 46d, 46f, and 46v (NIMH Macaque Template, NMT

v2.0 Macaque Atlas, Jung et al., 2021; Seidlitz et al., 2018). Because sequence related activity in

human RLPFC was observed in both hemispheres, we used identical spheres (mirrored

coordinates) in the left and right hemispheres (referred to as L46 and R46, respectively)

throughout. To compare activity between rule and number deviants and NISR in these ROIs, we

created a model that included separate regressors for each habituation timing and deviant type,

modeled as zero-duration onsets. Statistical testing was performed on ~10 run bins (n = 22), each

consisting of data from a single monkey (see Methods). We compared t-values from the contrast

of each condition over baseline (e.g., Rule Deviants > Baseline vs. NISR > Baseline) to account

for potential differences in variance across conditions. This type of comparison was used to

examine ROI activity throughout, and we refer to comparisons by the conditions of interest

(without listing the contrast over baseline, e.g., Rule Deviants > NISR). All statistical tests on

ROIs were performed on binned data and included a covariate for monkey identity (n = 3). While

we report the effect of monkey in the following analyses, the main focus of the study was not on

individual differences, and our discussion focuses on condition effects.

We found that R46 represented abstract sequence changes, showing greater deviant activation

across both deviant types (**Figure 2**;**Table 1**). Responses were reliably greater for rule deviants

compared to NISR (sequence type: $F(1, 19) = 4.6$, $p = 0.046$, $\eta_p^2 = 0.19$) and marginally greater

for number deviants compared to NISR (sequence type: $F(1, 19) = 3.9$, $p = 0.062$, $\eta_p^2 = 0.17$).

Even though deviant responses compared to NISR in L46 did not reach statistical significance

(**Table 1**), there were no reliable differences between responses in R46 and L46 (**Table 2**). These

results suggest that a specific region of monkey DLPFC, area 46, monitors abstract visual

sequence structure.

Figure 2. Area 46 represents abstract visual sequences. T-values for the condition of interest > baseline are

shown. The locations of area 46 regions of interest (ROIs), L46 and R46, are outlined in black on coronal sections (y

= 33). **A.** Rule deviants compared to new items, same rule (NISR) in L46. **B.** Number deviants compared to NISR

in L46. **C.** Rule deviants compared to NISR in R46 showed a reliable difference. **D.** Number deviants compared to

NISR in R46 showed a marginal difference. Comparisons in L46 showed similar trends as in R46. Error bars are

95% confidence intervals (1.96 x standard error of the within-bin mean).

Table 1. Repeated measures ANOVAs comparing rule and number deviants to NISR in L46 and R46.

Factor	dfs	L46			R 46		
		F	p	η_p^2	F	p	η_p^2
Rule Deviants > NISR							
Sequence Type	1, 19	0.39	0.54	0.02	4.6	0.046	0.19
Monkey	2, 19	3.55	0.049	0.27	1.4	0.28	0.13
Monkey x Sequence Type	2, 19	1.15	0.34	0.11	2.9	0.078	0.24
Number Deviants > NISR							
Sequence Type	1, 19	0.51	0.48	0.03	3.9	0.062	0.17
Monkey	2, 19	3.8	0.041	0.29	0.76	0.48	0.074
Monkey x Sequence Type	2, 19	0.098	0.91	0.01	5.15	0.016	0.35
Habituation > NISR							
Sequence Type	1, 19	0.0053	0.94	0.0003	1.6233	0.33	0. 051
Monkey	2, 19	4.35	0.028	0.31	3.37	0.48	0.26
Monkey x Sequence Type	2, 19	0.45	0.64	0.05	1.6	0.22	0.15

Table 2. Repeated measures ANOVAs comparing deviant responses in L46 and R46.

Factor	dfs	Rule Deviants > NISR			Number Deviants > NISR		
		F	p	η_p^2	F	p	η_p^2
R46 > L46							
Sequence Type	1, 40	3.6	0.07	0.08	0.85	0.36	0.02
Monkey	2, 40	4.15	0.02	0.17	1.14	0.33	0.05
Brain Area	1, 40	0.005	0.94	0.0001	0.67	0.42	0.016
Monkey x Sequence Type	2, 40	0.54	0.6	0.026	2.88	0.07	0.13
Brain Area x Sequence Type	1, 40	0.51	0.48	0.012	2.7	0.11	0.06

As a control, we also examined conditions where the pool of images differed, but the abstract

sequential structure did not. If area 46 was responding specifically to a change in the abstract

sequential structure, then a change in the images should not change its activation level. We

examined the difference in contrast t-values between NISR and habituation trials with

comparable stimulus durations ("medium" timing, as in **Figure 1D**). We did not find any

significant differences between these conditions in either R46 or L46 (**Table 1**), indicating that

changes in activation in area 46 were specific to changes in abstract sequential structure.

Results from area 46 ROIs were supported by whole-brain contrasts examining responses to

number and rule deviants. Contrasts of Rule Deviants > NISR and Number Deviants > NISR

both showed significant clusters of activation in right area 46 (**Figure 3**, **Table 3**, **Extended**

Data Figure 3-1). Other significant clusters of activation were located in areas such as the

caudate nucleus, high-level auditory cortex (rostromedial belt region), and dorsal premotor

cortex, areas also observed in a similar auditory sequence task (Wang et al., 2015). Further,

deviant responses in earlier sensory areas (e.g., V2) may be analogous to responses in auditory

cortex previously observed. Though we could not address the question of sensory generality

within the current experiment, these results raise the intriguing possibility that these previously

indicated areas could be sensory-modality general in their responses to abstract sequential

structure.

Figure 3. Whole-brain deviant activity shows area 46 represents both rule and number deviants. A. Voxel

wise contrast of Rule Deviants > NISR false discovery rate (FDR) error cluster corrected for multiple comparisons

(FDRc < 0.05, height p < 0.005 unc., extent = 130) are shown. Individual monkey data shown in **Extended Data**

Figure 3-1A. **B.** Voxelwise contrast of Number Deviants > NISR (FDRc < 0.05, height p < 0.005 unc., extent =

132). Individual monkey data shown in **Extended Data Figure 3-1B**. **C.** Overlap of Rule Deviants > NISR and

Number Deviants > NISR contrasts showed significant, unique conjunction (violet outlined in black) only in the

DLPFC. **D.** Voxelwise contrast of Rule Deviants > Number Deviants showing no significant clusters of activation in

area 46 (FDRc < 0.05, height p < 0.005 unc., extent = 102). **E.** As in (C) for Number Deviants > Rule Deviants

(FDRc < 0.05, height p < 0.005 unc., extent = 111). Black outline on inflated brains indicates location of L46 or R46

(depending on the hemisphere shown) for reference. Lateral Sulcus (LS), Dorsal Lateral Pre-frontal Cortex

(DLPFC), Ventral Lateral Pre-frontal Cortex (VLPFC), Secondary Visual Cortex (V2), Orbital Prefrontal Area

(12o).

Table 3. Rule and number deviants compared to NISR contrast activation coordinates. Area labels as in the NIMH NMT v02 Macaque Atlas.

Contrast Location	Extent (voxels)	x	y	z	Peak t-val
Rule Deviant > NISR					
Dorsolateral Prefrontal Cortex (46d)	282	10.5	34.5	21.5	4.24
		14	36	26	3.88
Medial Prefrontal Cortex (10mr)	195	0	47.5	17	5.96
Dorsal Pre-motor Cortex(6DR)		13	21	25	
Motor Cortex (F1)	213	-11	16.5	32	5.58
Caudate Nucleus (cd)	130	14.5	13	8	3.88
	138	-14.5	11.5	7.5	4.09
Lateral Sulcus/Auditory Cortex (RM)	282	-21	13.5	9	5.59
Secondary Visual Cortex (V2)	207	8.5	-15	21	4.34
Cerebellum					
Number Deviants > NISR					
Dorsolateral Prefrontal Cortex (Area 46d)	168	9	35	21.5	6.44
Orbital Prefrontal Area (12o)	162	-16	32	9.5	5.13
Secondary Visual Cortex (V2)	132	4.5	-16.5	18.5	4.96

To determine if the observed responses to number and rule deviants were similar in area 46, we directly examined whether these responses generalized across deviant types. The t-values in R46 were not different between rule and number deviants (sequence type: $F(1,19) = 0.0011$, $p = 0.92$, $\eta_p^2 = 0.11$, monkey: $F(2, 19) = 2.5$, $p = 0.11$, $\eta_p^2 = 0.21$; monkey x sequence type: $F(2, 19) = 1.15$, $p = 0.34$, $\eta_p^2 = 0.11$). Next, we performed a conjunction analysis to determine the areas of activation that overlapped in the Rule Deviants > NISR and Number Deviants > NISR contrasts. We found that the only cluster of significant overlap between the deviant contrasts was in right area 46 (**Figure 3C**). In support of this finding, whole-brain direct contrasts of rule and number deviants showed significant activation clusters in VLPFC, insula, and pre-motor cortex, but no significant clusters in area 46 (**Figure 3D, E**). In summary, these results suggest that abstract

visual sequential structure is monitored in area 46, and that this monitoring is both unique to area

46 and general across different kinds of deviations.

Ramping activation reflects sequence monitoring in monkey DLPFC

We next tested the prediction that area 46 would display similar dynamics to those observed in

humans during abstract sequences. Specifically, previous experiments in humans showed that

BOLD activity increased ("ramped") from the beginning to the end of sequences in the RLPFC

(Desrochers et al., 2015, 2019; McKim and Desrochers, 2022). Given the similarity in

connectivity between monkey area 46 and human RLPFC, we hypothesized that changes in

abstract sequence structure would also produce changes in ramping activation in area 46 if

abstract sequence monitoring underlies this dynamic.

To test if ramping dynamics were present in area 46 during this task, we first designed a model

to isolate these dynamics. This model included regressors for the three dominant potential

dynamics (**Figure 4A**, **Extended Data Figure 4-1**, see also Materials and Methods). First,

instantaneous onsets were included for each image presentation, effectively modeling sustained

activation throughout each sequence. Then, two parametric regressors were included: last item

change and ramping. The ramping regressor parallels the analysis that revealed ramping in

human RLPFC: it increases linearly from the first to the last item in the sequence, and resets at

each new sequence. The last item change regressor is low at the first three positions of each

sequence, and high at the last item in the sequence. This regressor was designed to account for

the fact that differences in the rule that the sequence followed would occur at the last item (e.g.,

the difference between AAAA and AAAB occurs at the fourth item), and a dynamic associated

with this change could have variance mistakenly assigned to a ramping regressor.

These parametric regressors were orthogonalized in a stepwise fashion and only absorbed

variance above and beyond variance accounted for by the onset regressor. The last regressor,

therefore, contained "unique" variance. A pair of models, one with unique variance assigned to

ramp and one with it assigned to last item, were created to examine these dynamics. The

correlation between the resulting regressors was, as expected, low. For example, in one bin of

this parametric model with the unique variance assigned to ramp, the average correlation

coefficient between last item change and ramp regressors was 0.00005 ($\pm$ 0.0001 standard

deviation). While there are likely nearly infinite variations of dynamics possible that lie across a

spectrum between the last item change and ramping (e.g., exponential), our purpose in designing

these models was not to explore the space of all possible dynamics, but to test for ramping

dynamics in area 46.

We found ramping dynamics in monkey area 46 related to abstract sequence monitoring. When

comparing t-values of contrasts between deviants and baseline using the same spherical area 46

ROIs described above, we observed significant, unique variance ascribed to ramping activation

in L46 during rule deviants compared to NISR (sequence type: $F(1, 19) = 5.03$, $p = 0.037$, $\eta_p^2 =$

0.21; **Table 4**; **Figure 4B**). Unique ramping activation showed a numerical trend in the same

direction for number deviants compared to NISR in L46, but it did not reach statistical

significance (**Table 4**). Activity in L46 was not reliably different between the two deviant types

(sequence type: $F(1,19) = 2.06$, $p = 0.17$, $\eta_p^2 = 0.098$, monkey: $F(2, 19) = 0.57$, $p = 0.64$, $\eta_p^2 =$

0.06; monkey x sequence type: $F(2, 19) = 2.01$, $p = 0.16$, $\eta_p^2 = 0.17$). In R46, changes in unique

ramping activation during rule and number deviants were not significant (**Table 4**). Despite

apparent differences between L46 and R46, unique ramping was not reliably different between

these ROIs (**Table 5**). These results suggest that area 46 shows ramping dynamics for sequential

rule changes. Interestingly, ramping may be preferentially present in L46, suggesting that while

both hemispheres detect abstract sequence deviations, they may do so with different dynamics.

Further, these results suggest similar sequential monitoring processes may be present across

species in analogous areas.

Figure 4. Area 46 shows ramping activity for deviations to an established sequence rule. Parametric models and

T-values for the condition of interest > baseline shown. Coronal brain slices ($y = 33$) show locations of area 46

ROIs, L46 and R46, outlined in black. **A.** Example of regressors used to model parametric ramp and parametric last

item. Example regressors through the orthogonalization process shown in **Extended Data Figure 4-1**. **B.** Unique

ramping during rule deviants compared to NISR in L46 showed a reliable difference. **C.** Unique ramping during

number deviants compared to NISR in L46. **D.** Unique ramp number deviants compared to NISR in R46. **E.** Unique

ramping during number deviants compared to NISR in R46. Comparisons that were not reliably different showed

similar trends. Error bars are 95% confidence intervals (1.96 x standard error of the within-bin mean).

Table 4. Repeated measures ANOVAs comparing unique ramping activity during rule and number deviants to NISR

in L46 and R46.

Factor	dfs	L46			R 46		
		F	p	η_p^2	F	p	η_p^2

		Unique Ramp, Rule Deviants > NISR			Unique Ramp, Number Deviants > NISR		
Unique Ramp, Rule Deviants > NISR							
Sequence Type	1, 19	5.03	0.037	0.21	0.01	0.91	0.00075
Monkey	2, 19	0.82	0.45	0.08	0.64	0.54	0.63
Monkey x Sequence Type	2, 19	0.23	0.8	0.0235	0.81	0.46	0.08
Unique Ramp, Number Deviants > NISR							
Sequence Type	1, 19	0.42	0.52	0.02	0.03	0.86	0.0016
Monkey	2, 19	2.39	0.12	0.2	0.24	0.45	0.08
Monkey x Sequence Type	2, 19	5.15	0.63	0.047	5.15	0.84	0.0235

Table 5. Repeated measures ANOVAs comparing unique ramping deviant responses in L46 and R46.

Factor	*dfs*	Unique Ramp, Rule Deviants > NISR			Unique Ramp, Number Deviants > NISR		
		F	**p**	η_p^2	**F**	**p**	η_p^2
R46 > L46							
Sequence Type	1, 40	0.38	0.54	0.009	2.4	0.13	0.06
Monkey	2, 40	1.62	0.21	0.07	1.5	0.24	0.07
Brain Area	1, 40	0.02	0.89	0.0005	1.5	0.25	0.03
Monkey x Sequence Type	2, 40	0.64	0.53	0.03	0.12	0.89	0.006
Brain Area x Sequence Type	1, 40	0.14	0.71	0.003	2.7	0.11	0.06

Because variance due to changes at the last item of the sequence could be misattributed to

ramping regressors, we directly compared activity in area 46 that could be accounted for by

ramping and last item change regressors. In this control analysis, we found that activity was

significantly greater for unique ramping than unique last item change during rule deviants in L46

(sequence type: $F(1, 19) = 9.53$, $p = 0.006$, $\eta_p^2 = 0.33$; **Table 6**; **Figure 5A**). Number deviants in

L46 and both deviants in R46 showed similar numerical trends for unique ramping accounting

for greater variance than last item change but did not reach statistical significance (**Figure 5B-D**;

Table 6). These results suggest that area 46 dynamics during abstract sequence monitoring are

best accounted for by a ramping function, rather than a last item change.

Figure 5. Area 46 shows greater unique ramping than unique last item activity during abstract sequence deviants. T-values for the condition of interest > baseline shown. **A.** Unique ramp compared to unique last item during rule deviants in L46 showed a reliable difference. **B.** Unique ramp compared to unique last item during number deviants in L46. **C.** Unique ramp compared to unique last item during rule deviants in R46. **D.** Unique ramp compared to unique last item during number deviants in R46. Comparisons that were not reliably different showed similar trends. Error bars are 95% confidence intervals (1.96 x standard error of the within-bin mean).

Table 6. Repeated measures ANOVAs comparing unique ramping activity to unique last item activity during rule and number deviants to NISR in L46 and R46.

Factor	dfs	L46			R 46		
		F	p	η_p^2	F	p	η_p^2
Unique Ramp, Rule Deviants > Unique Last Item, Rule Deviants							
Sequence Type	1, 19	9.53	0.006	0.33	0.008	0.93	0.0004
Monkey	2, 19	1.02	0.38	0.97	0.72	0.5	0.07
Monkey x Sequence Type	2, 19	0.37	0.7	0.38	0.09	0.91	0.01
Unique Ramp, Number Deviants > Unique Last Item, Number Deviants							
Sequence Type	1, 19	0.196	0.66	0.01	0.07	0.8	0.0036
Monkey	2, 19	2.08	0.15	0.18	1.47	0.25	0.13
Monkey x Sequence Type	2, 19	1.26	0.31	0.12	0.15	0.86	0.015

ROI results were supported by whole-brain contrasts that examined ramping and last item

dynamics. Unique ramping was present in left area 46 during rule deviants compared to NISR

(**Figure 6A, Table 7, Extended Data Figure 3-1C**). Other clusters of activation were present in

the visual cortex and superior temporal gyrus that were similar to those observed for ramping

activation in humans (Desrochers et al., 2015, 2019). As expected from the ROI analyses, the

number deviant ramping contrast had significant whole brain clusters in areas such as visual

cortex and putamen but no significant clusters in area 46 (**Figure 6B, Table 7**). The localization

of significant unique last item clusters was different than ramping. Specifically, a more anterior

region of ventral area 46 (46v), in contrast to more posterior and dorsal area 46 (46d) observed

for unique ramping, showed significant activation for unique last item change in rule deviants

compared to NISR (**Figure 7A, Table 8**). Other significant clusters of activation for last item

change included the somatosensory cortex and central orbitofrontal cortex (area 13m). We also

contrasted unique last item variance in number deviants compared to NISR, and observed

activation in areas such as anterior cingulate gyrus, insula and visual cortices but no surviving

clusters in area 46 (**Figure 7B, Table 8**). These results suggest that subregions of area 46 within

the same hemisphere may reflect separable aspects of changes in sequential structure.

Figure 6. Area 46 shows unique ramping during abstract sequence deviants. A. Unique Ramp, Rule Deviants >

NISR (FDRc p < 0.05, height p < 0.005 unc., ext. 82). Individual monkey data shown in **Extended Data Figure 3-**

1C. B. Unique Ramp, Number Deviants > NISR (FDRc p < 0.05, height p < 0.005 unc., ext. 89). Black outline on

inflated brains indicates location of L46 or R46 (depending on the hemisphere shown) for reference. Medial

Prefrontal Cortex (A32) Dorsal Lateral Pre-frontal Cortex (DLPFC), Primary Visual Cortex (V1), Insular Cortex

(Ins), Anterior Lateral Belt Region of the Auditory Cortex (AL), Anterior Intraparietal Area (AIP).

616 **Table 7**. Unique ramp rule and number deviants compared to NISR contrast activation coordinates.

Contrast Location	Extent (voxels)	x	y	z	Peak t-val
Unique Ramp, Rule Deviants > NISR					
Primary Motor Cortex (F1)	153	11	13.5	29.5	6.48
Caudal Dorsal Pre-Motor Cortex (PMdc)	89	12	18	34.5	6.17
Dorsal Dorsal Lateral Prefrontal Cortex (46d)	130	-12.5	36	23	5.34
Intraparietal Area (IPa)	121	16	18.5	2	5.33
Rostral Area 12 (12r)	128	-15	29	14.5	5.19
Caudate Nucleus (Cd)	385	-10.5	19	23	5.15
Caudal Pre-Motor Cortex (PMdc)	113	-18.5	20	30	5
Secondary Visual Cortex (V2)	82	28	-7.5	14.5	4.67
Primary Visual Cortex (V1)	101	-20.5	-11	27	4.66
Body of the Fornix (bfx)	120	-1.5	12.5	22.5	4.49

Somatosensory Parietal Cortex (1,2)	120	-13	7.5	33.5	4.4
Primary Motor Cortex (M1)	97	-3.5	14.5	38	4.3
Medial Septum (ms)	82	-1	18.5	13	4.18
Secondary/Primary Visual Cortex (V2/V1)	129	9	-11	19	4.16
Intraparietal Area (IPa)	97	18.5	7.5	16	4.01
Globus Pallidus (MGPi)	250	-6.5	13	8	3.99
Pre-genual Cortex (24a)	101	-2	38.5	17	3.96
Unique Ramp, Number Deviants > NISR					
Visual Area 3 (V3)	163	7.5	-12.5	26.5	6.42
Anterior Lateral Belt Region (AL)	130	28	17	12.5	6.35
Agranular and Dysgranular Insula (Ia/Id)	103	21	19	10.5	5.62
Putamen (Pu)	127	7	24.5	10.5	5.5
Caudate Nucleus (Cd)	772	-5.5	20	18	5.29
Primary Visual Cortex (V1)	132	9	-11.5	19	5.16
Cerebellum	250	7	-15.5	12	4.85
Somatosensory Parietal Cortex (1,2)	89	-4.5	4	35	4.72
Cerebellum	138	4	-18	5.5	4.67
Anterior Intraparietal Area (AIP)	123	18.5	10.5	25.5	4.65
Somatosensory Parietal Cortex (1,2)	115	-21	13	32.5	4.5
Caudal Dorsal Pre-Motor Cortex (PMdc)	324	-10.5	20	28	4.45
Ventral Tegmental Area (VTA)	282	-4	4	3	4.37
Posterior Cingulate Gyrus (23c)	100	-7.5	5.5	30.5	4.07
Somatosensory Parietal Cortex (1,2)	152	-12.5	6.5	35	4.01

Figure 7. DLPFC does not show significant responses to unique last items during abstract sequence deviants.

A. Unique Last Item, Rule Deviants > NISR (FDRc < 0.05, height p < 0.005 unc., extent = 148). **B.** Unique Last

Item, Number Deviants > NISR (FDRc < 0.05, height p < 0.005 unc., extent = 82). Black outline on inflated brains

622   indicates location of L46 or R46 (depending on the hemisphere shown) for reference. Ventral Dorsal Lateral

623   Prefrontal Cortex (vDLPFC), Medial Superior Temporal Cortex (MST), Superior Temporal Sulcus Dorsal Bank

624   (TPO), Orbital Medial Prefrontal Cortex, Fundus of the Dorsal Lateral Prefrontal Cortex (46f).

626   **Table 8**. Unique last item rule and number deviants compared to NISR contrast activation coordinates.

Contrast Location	Extent (voxels)	x	y	z	Peak t-val
Unique Last Item, Rule Deviants > NISR					
Ventral Dorsal Lateral Prefrontal Cortex (46v)	148	-12	43	22	5.23
Granular Insula (Ig)	170	19	11	20	5.06
Medial Area 11 (11m)	156	3.5	41	17.5	4.93
Ventral Intraparietal Area/White Matter (VIP/WM)	360	8.5	-1	22.5	4.1
Unique Last Item, Number Deviants > NISR					
Area TFO (TFO)	121	-14	4	1.5	5.75
Caudate Nucleus	82	1	29.5	16.5	5.32
Primary Visual Cortex (V1)	247	-8.5	-18.5	12	5.13
Area 13 (13m)	184	4	38	18.5	4.92
Cerebellum	90	12.5	-12.5	5	4.89
Granular Insula (Ig)	117	17.5	16	15.5	4.36
Temporal Parietooccipital Associated Area (TPO)	88	21.5	14	7.5	4.32
Caudomedial Belt Region (CM)	116	-15	2.5	22	4.19
Anterior Cingulate Gyrus (24a)	85	-2	28.5	20.5	4.14
Posterior Cingulate Gyrus (29)	151	7	0.5	20.5	4.03

To determine if clusters of activation for unique ramping and unique last item were separable in

the frontal cortex, we directly compared the amount of variance assigned to each cluster for both

dynamics. One possibility is that, due to thresholding at the whole-brain level, there was similar

activation for each dynamic across areas 46d and 46v, but that the peak, and thus the location of

the thresholded cluster, differed slightly in location. To address this possibility, we created two

ROIs from the clusters of significant activation in Unique Ramp, Rule Deviants > NISR contrast

(area 46d, center *xyz* = -12.2, 36, 23.8 mm), and Unique Last Item, Rule Deviants > NISR

contrast (area 46v, center *xyz* = -12.3, 42.9, 21.8 mm). We found a significant interaction

between ROI and model in rule deviants compared to NISR (**Figure 8**; sequence type: F(1, 40) =

0.13, *p*= 0.71, η_p^2 = 0.003; monkey: F(2, 40) = 2.23, p = 0.12, η_p^2= 0.1; brain area: F(1, 40) =

0.09, p = 0.77, η_p^2 = 0.002; monkey x sequence type: F(2, 40) = 0.33, p = 0.72, η_p^2 =0.016; brain

area x sequence type: F(1, 40) = 29.3, p < 0.001, η_p^2 = 0.42), indicating that responses in the

ramping and last item clusters were reliably different. As expected from the whole-brain

contrasts, there was no significant interaction for number deviants compared to NISR (not

shown; F(1, 40) = 0.796, *p* = 0.38, η_p^2 = 0.02). These results show that these nearby clusters in

dorsal and ventral area 46 are separable in their dynamics.

Figure 8. Nearby regions of DLPFC show significantly different dynamics. ROIs constructed from significant

areas of activation in unique ramping (46d, dDLPFC) and unique last item (46v, vDLPFC) contrasts shown on

coronal sections. T-values for the condition of interest > baseline shown. Models show a double dissociation in left

area 46 during rule deviants. Error bars are 95% confidence intervals (1.96 x standard error of the within-bin mean).

<u>Discussion</u>

In this study, we examined if and how monkey DLPFC (area 46) monitors abstract sequential

information. We tested two main hypotheses: First, that evidence of abstract sequence

monitoring would occur in the area of monkey DLPFC analogous to human RLPFC previously

shown to be critical for abstract sequential tasks (Desrochers et al., 2015, 2019; McKim and

Desrochers, 2022); and, second, that ramping dynamics would be associated with abstract

sequence monitoring and show changes when abstract sequential structure changed. These

hypotheses were tested in a no-report sequence viewing task that allowed the isolation of

dynamics associated with sequence monitoring from other potential confounds such as motor

preparation. We found evidence to support both hypotheses. Right area 46 responded to abstract

sequence changes in both rule and number. Interestingly, left area 46 also responded to changes

in abstract sequential rules, but with ramping dynamics that were similar to those observed in

humans and separable from an increase only at the last item. These results suggest that a specific

subregion of monkey DLPFC is specialized for monitoring general visual abstract sequential

information and is a point of critical potential functional homology between human and monkey

PFC during higher-level cognitive function.

The activation patterns found, with ramping dynamics primarily on the left and onset-based on

the right, are consistent with prior findings. Though not explicitly tested, in humans ramping was

observed preferentially on the left during abstract sequential tasks (Desrochers et al., 2015,

2019), Experiment 1) in contrast to bilateral ramping activation observed in tasks where

sequences were based on stimulus identity (i.e., ordered visual items; (Desrochers et al., 2019),

Experiment 2; (McKim and Desrochers, 2022). These human results are also consistent with the

long-standing literature that emphasizes abstract cognitive functions in the left hemisphere of

humans (e.g., (Badre and D'Esposito, 2007; Bunge et al., 2009; Wendelken et al., 2012), with the

most famous such example being language (Broca, 1861; Milner, 1971; Petrides, 2013).

Ramping activity has been observed in human frontal cortex during sequential language

information processing (i.e., sentence comprehension) with electrocorticography (ECoG)

(Fedorenko et al., 2016). While generally consistent with prior work, future studies of the distinct

left and right hemisphere activation dynamics will be needed to determine their underlying

drivers and their potential cognitive import. In particular, the present findings suggest that neural

dynamics underlying DLPFC sequence monitoring may be distinct to each hemisphere. The

monkey fMRI findings described here can provide a guide for such recordings. In sum, our

results raise the possibility that the representation in and contribution of DLPFC to abstract

sequence monitoring is lateralized in monkeys, and provide a road map to future studies.

An advantage of fMRI is the whole brain view that is not afforded by typical

electrophysiological techniques in macaques. This view leads to potential insights about

functional organization of brain areas without the limitations of a recording chamber (Milham et

al., 2022). For example, in recent literature, fMRI has enabled the mapping of projections to and

from the PFC with a level of specificity and across distances not previously possible on this scale

(Xu et al., 2022). This work found an overlap in topographically organized high-level visual

maps from the dorsal and ventral streams in primate lateral prefrontal cortex. These results raise

the possibility that the localization of abstract visual sequence monitoring in DLPFC results from

its position near the apex of highly organized visuo-spatial maps. The overarching organization

693 of more cognitive processes in monkey frontal cortex has remained more elusive (Hutchison and

694 Everling, 2014; Neubert et al., 2014, 2014; Saleem et al., 2014), and the results presented here

695 represent a critical step forward in understanding their topography.

696 We observed similarities and differences to a previous study using a similar auditory task (Wang

697 et al., 2015) that may reflect the modality employed and the capacity for generalization. Areas of

698 the brain that responded to deviant sequences in both the current visual and prior auditory studies

699 may be modality general. These areas included premotor cortex, caudate nucleus, and the

700 auditory cortex rostromedial belt. In contrast, brain areas that uniquely responded to deviants in

701 the auditory or this visual task may be modality specific for abstract sequence changes. For

702 example, deviant responses in ventral LPFC and superior temporal sulcus were unique to the

703 auditory study. In this visual task, deviant responses in DLPFC, visual cortex, and mPFC were

704 observed that were not observed in the auditory task. Though we cannot draw strong conclusions

705 without direct comparison between the modalities, these results suggest that networks of brain

706 areas that partially overlap may constitute abstract sequence tracking across modalities. One

707 further important difference in the studies is that in the auditory study, there was a lack of

708 overlap between areas that respond to rule and number deviants in the monkey (in contrast to the

709 human). Here, we observed overlap in these responses in area 46, suggesting a higher level of

710 visual integration in the monkey. The question of sensory domain generality and integration

711 remains open to further investigation.

712 The results observed here in this no-report abstract sequence viewing task in monkeys are similar

713 to those observed in humans during sequential tasks in important ways. First, ramping activation

714 was observed in similar regions in the frontal cortex in the monkey (area 46) and human

(RLPFC). Other similar areas included visual cortex, putamen, and pre-motor cortex (Desrochers

et al., 2015, 2019; McKim and Desrochers, 2022). Though the current experiment was only

designed to detect the presence of ramping in relation to abstract sequences, by analogy, the

function may be similar in the lateral prefrontal cortex across species. Ramping activity has been

ascribed to many possible functions across species, including accumulating evidence (de Lange

et al., 2010; Krueger et al., 2017; Darriba and Waszak, 2018; Lin et al., 2020), keeping time

(Nobre et al., 2007; Berdyyeva and Olson, 2011; Cueva et al., 2020), reward anticipation

(Roesch and Olson, 2007; Horst and Laubach, 2013; Chiew et al., 2016; Falcone et al., 2019;

McKim and Desrochers, 2022) and monitoring sequence position (Desrochers et al., 2015,

2019). These possibilities are not mutually exclusive, as recent evidence in humans suggests that

reward anticipation and sequence information may be present simultaneously in this signal

(McKim and Desrochers, 2022). The current experiment identified ramping in the DLPFC as

being sequence related, but did not examine other potential influences and therefore remains an

open avenue of future inquiry. Similarly, other brain areas that display ramping activation remain

open for investigation.

Though this study bears resemblance to a field of literature using statistical learning paradigms,

the majority of those studies use tasks that rely on the identity of the stimuli themselves.

Importantly, this study is distinct from most statistical learning studies because the identity of the

stimulus alone cannot predict the following item (i.e., knowing the current fractal is the green

one does not determine the next stimulus without also having sequential rule information). A

subset of work in the infant learning literature examines statistical learning that is not dependent

on stimulus identity (i.e., "artificial grammar"), but mostly auditory tasks were used (e.g.,

(Saffran et al., 1996)). To our knowledge, only one behavioral study examined violations to a

visual (and auditory) artificial grammar where sequences of different lengths are constructed

according to set transition probabilities (Milne et al., 2018). While the findings were important

because violations were detected similarly in monkeys and humans, there were no neural data

presented. Further, while transition probabilities created the sequences, there was not a set "rule"

by which they were constructed. Therefore, the present study is unique in examining neural

responses to abstract visual sequences.

This study contained the following limitations. First, the timing of the stimuli in the current

design did not allow examination of dynamics of individual sequence items, only across the

sequence as a whole. In a single experiment, it was not feasible to separate each sequential item

by the time required to model each separately in an event-related design. Therefore, future work

will aim to examine the dynamics of individual sequential items in greater detail. Second, while

the no-report task allowed the elimination of motor preparatory confounds, it did not allow for

direct correlation with behavioral performance. Although the observed signals will potentially

also underlie tasks that require responses, this assertion remains to be tested and the present

study is an important foundation for further experiments. Third, we have focused here on the

DLPFC because although its importance in cognitive processes in monkeys has been established,

its response to visual abstract sequences and potential correspondence to dynamics in humans

remained unknown. The DLPFC is part of a network of areas active in this task, and although

they are outside the scope of the current experiment, they remain an important avenue of future

research.

In summary, we provide evidence that a specific subregion of monkey DLPFC monitors abstract

visual sequences and generalizes across different sequence violations (number and rule). Further,

sequence related ramping dynamics were also observed in DLPFC. Importantly, this region is

possibly analogous to human RLPFC, where necessary sequence-related ramping signals have

been identified in the past. These results suggest functional homology across the species as to

where and how more general abstract visual sequential information is represented in the brain.

These findings, in turn, inform future models of how abstract sequential information is

represented during more complex behaviors across species.

References

Averbeck BB, Sohn J-W, Lee D (2006) Activity in prefrontal cortex during dynamic selection of
action sequences. Nature neuroscience 9:276–282.

Badre D, D'Esposito M (2007) Functional magnetic resonance imaging evidence for a
hierarchical organization of the prefrontal cortex. Journal of Cognitive Neuroscience
19:2082–2099.

Barone P, Joseph J-P (1989) Prefrontal cortex and spatial sequencing in macaque monkey. Exp
Brain Res 78 Available at: http://link.springer.com/10.1007/BF00230234 [Accessed
February 9, 2022].

Bekinschtein TA, Dehaene S, Rohaut B, Tadel F, Cohen L, Naccache L (2009) Neural signature
of the conscious processing of auditory regularities. Proc Natl Acad Sci USA 106:1672–
1677.

Berdyyeva TK, Olson CR (2010) Rank Signals in Four Areas of Macaque Frontal Cortex During

Selection of Actions and Objects in Serial Order. J Neurophysiol 104:141–159.

Berdyyeva TK, Olson CR (2011) Relation of ordinal position signals to the expectation of

reward and passage of time in four areas of the macaque frontal cortex. Journal of

Neurophysiology 105:2547–2559.

Borra E, Ferroni CG, Gerbella M, Giorgetti V, Mangiaracina C, Rozzi S, Luppino G (2019)

Rostro-caudal Connectional Heterogeneity of the Dorsal Part of the Macaque Prefrontal

Area 46. Cerebral Cortex 29:485–504.

Borra E, Gerbella M, Rozzi S, Luppino G (2011) Anatomical Evidence for the Involvement of

the Macaque Ventrolateral Prefrontal Area 12r in Controlling Goal-Directed Actions. J

Neurosci 31:12351–12363.

Broca P (1861) Remarques sur le siège de la faculté du langage articulé, suivies d'une

observation d'aphémie (perte de la parole). Bulletin et Memoires de la Societe

anatomique de Paris 6:330–357.

Bunge SA, Helskog EH, Wendelken C (2009) Left, but not right, rostrolateral prefrontal cortex

meets a stringent test of the relational integration hypothesis. NeuroImage 46:338–342.

Burgess PW, Scott SK, Frith CD (2003) The role of the rostral frontal cortex (area 10) in

prospective memory: a lateral versus medial dissociation. Neuropsychologia 41:906–918.

Chiew KS, Stanek JK, Adcock RA (2016) Reward Anticipation Dynamics during Cognitive

Control and Episodic Encoding: Implications for Dopamine. Front Hum Neurosci 10

Available at: https://www.frontiersin.org/articles/10.3389/fnhum.2016.00555/full

[Accessed February 12, 2020].

Cueva CJ, Saez A, Marcos E, Genovesio A, Jazayeri M, Romo R, Salzman CD, Shadlen MN,

Fusi S (2020) Low-dimensional dynamics for working memory and time encoding.

PNAS 117:23021–23032.

Darriba Á, Waszak F (2018) Predictions through evidence accumulation over time. Scientific

Reports 8:1–15.

de Lange FP, Jensen O, Dehaene S (2010) Accumulation of evidence during sequential decision

making: the importance of top-down factors. J Neurosci 30:731–738.

Dehaene S, Meyniel F, Wacongne C, Wang L, Pallier C (2015) The Neural Representation of

Sequences: From Transition Probabilities to Algebraic Patterns and Linguistic Trees.

Neuron 88:2–19.

Desrochers TM, Ahuja A, Maechler MR, Shires J, Yusif Rodriguez N, Berryhill ME (2022)

Caught in the ACTS: Defining Abstract Cognitive Task Sequences as an Independent

Process. Journal of Cognitive Neuroscience 34:1103–1113.

Desrochers TM, Chatham CH, Badre D (2015) The necessity of rostrolateral prefrontal cortex

for higher-level sequential behavior. Neuron 87:1357–1368.

Desrochers TM, Collins AGE, Badre D (2019) Sequential Control Underlies Robust Ramping

Dynamics in the Rostrolateral Prefrontal Cortex. The Journal of Neuroscience 39:1471–

1483.

Du J, Rolls ET, Cheng W, Li Y, Gong W, Qiu J, Feng J (2020) Functional connectivity of the

orbitofrontal cortex, anterior cingulate cortex, and inferior frontal gyrus in humans.

Cortex 123:185–199.

Eiselt A-K, Nieder A (2013) Representation of Abstract Quantitative Rules Applied to Spatial

and Numerical Magnitudes in Primate Prefrontal Cortex. J Neurosci 33:7526–7534.

Falcone R, Weintraub DB, Setogawa T, Wittig JH, Chen G, Richmond BJ (2019) Temporal

Coding of Reward Value in Monkey Ventral Striatal Tonically Active Neurons. J

Neurosci 39:7539–7550.

Farooqui AA, Mitchell D, Thompson R, Duncan J (2012) Hierarchical Organization of

Cognition Reflected in Distributed Frontoparietal Activity. J Neurosci 32:17373–17381.

Fedorenko E, Scott TL, Brunner P, Coon WG, Pritchett B, Schalk G, Kanwisher N (2016)

Neural correlate of the construction of sentence meaning. Proc Natl Acad Sci USA

113:E6256–E6262.

Fujii N, Graybiel AM (2003) Representation of Action Sequence Boundaries by Macaque

Prefrontal Cortical Neurons. Science 301:1246–1249.

Gerbella M, Belmalih A, Borra E, Rozzi S, Luppino G (2010) Cortical Connections of the

Macaque Caudal Ventrolateral Prefrontal Areas 45A and 45B. Cerebral Cortex 20:141–

168.

Gerbella M, Borra E, Tonelli S, Rozzi S, Luppino G (2013) Connectional Heterogeneity of the

Ventral Part of the Macaque Area 46. Cerebral Cortex 23:967–987.

Gilbert SJ, Gonen-Yaacovi G, Benoit RG, Volle E, Burgess PW (2010) Distinct functional

connectivity associated with lateral versus medial rostral prefrontal cortex: A meta-

analysis. NeuroImage 53:1359–1367.

Henssen A, Zilles K, Palomero-Gallagher N, Schleicher A, Mohlberg H, Gerboga F, Eickhoff

SB, Bludau S, Amunts K (2016) Cytoarchitecture and probability maps of the human

medial orbitofrontal cortex. Cortex 75:87–112.

Horst NK, Laubach M (2013) Reward-related activity in the medial prefrontal cortex is driven by

consumption. Front Neurosci 7 Available at:

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3622888/ [Accessed February 10, 2020].

Hoshi E, Shima K, Tanji J (1998) Task-Dependent Selectivity of Movement-Related Neuronal

Activity in the Primate Prefrontal Cortex. Journal of Neurophysiology 80:3392–3397.

Hutchison RM, Everling S (2014) Broad intrinsic functional connectivity boundaries of the

macaque prefrontal cortex. NeuroImage 88:202–211.

Jean-Baptiste Poline MB (2002) Region of interest analysis using an SPM toolbox.

Jung B, Taylor PA, Seidlitz J, Sponheim C, Perkins P, Ungerleider LG, Glen D, Messinger A

(2021) A comprehensive macaque fMRI pipeline and hierarchical atlas. NeuroImage

235:117997.

Kim HF, Hikosaka O (2013) Distinct Basal Ganglia Circuits Controlling Behaviors Guided by

Flexible and Stable Values. Neuron 79:1001–1010.

Koechlin E, Corrado G, Pietrini P, Grafman J (2000) Dissociating the role of the medial and

lateral anterior prefrontal cortex in human planning. Proceedings of the National

Academy of Sciences 97:7651–7656.

Krueger PM, van Vugt MK, Simen P, Nystrom L, Holmes P, Cohen JD (2017) Evidence

accumulation detected in BOLD signal using slow perceptual decision making. J

Neurosci Methods 281:21–32.

Leite FP, Tsao D, Vanduffel W, Fize D, Sasaki Y, Wald LL, Dale AM, Kwong KK, Orban G a,

Rosen BR, Tootell RBH, Mandeville JB (2002) Repeated fMRI using iron oxide contrast

agent in awake, behaving macaques at 3 Tesla. NeuroImage 16:283–294.

Lin Z, Nie C, Zhang Y, Chen Y, Yang T (2020) Evidence accumulation for value computation in

the prefrontal cortex during decision making. Proceedings of the National Academy of

Sciences 117:30728–30737.

McKim TH, Desrochers TM (2022) Reward Value Enhances Sequence Monitoring Ramping

Dynamics as Ending Rewards Approach in the Rostrolateral Prefrontal Cortex. eNeuro

9:ENEURO.0003-22.2022.

McLaren DG, Kosmatka KJ, Oakes TR, Kroenke CD, Kohama SG, Matochik JA, Ingram DK,

Johnson SC (2009) A Population-Average MRI-Based Atlas Collection of the Rhesus

Macaque. NeuroImage 2:52–59.

Milham M et al. (2022) Toward next-generation primate neuroscience: A collaboration-based

strategic plan for integrative neuroimaging. Neuron 110:16–20.

Milne AE, Petkov CI, Wilson B (2018) Auditory and Visual Sequence Learning in Humans and

Monkeys using an Artificial Grammar Learning Paradigm. Neuroscience 389:104–117.

Milner B (1971) Interhemispheric differences in the localization of psychological processes in

man. British medical bulletin.

Miyashita Y, Higuchi S-I, Sakai K, Masui N (1991) Generation of fractal patterns for probing the

visual memory. Neuroscience Research 12:307–311.

Moayedi M, Salomons TV, Dunlop KAM, Downar J, Davis KD (2015) Connectivity-based

parcellation of the human frontal polar cortex. Brain Struct Funct 220:2603–2616.

Naya Y, Chen H, Yang C, Suzuki WA (2017) Contributions of primate prefrontal cortex and

medial temporal lobe to temporal-order memory. PNAS 114:13555–13560.

Neubert F-X, Mars RB, Thomas AG, Sallet J, Rushworth MFS (2014) Comparison of Human

Ventral Frontal Cortex Areas for Cognitive Control and Language with Areas in Monkey

Frontal Cortex. Neuron 81:700–713.

Ninokura Y, Mushiake H, Tanji J (2004) Integration of temporal order and object information in

the monkey lateral prefrontal cortex. J Neurophysiol 91:555–560.

Nobre A, Correa A, Coull J (2007) The hazards of time. Current Opinion in Neurobiology

17:465–470.

Petrides M (2013) Neuroanatomy of language regions of the human brain. Academic Press.

Roesch MR, Olson CR (2007) Neuronal activity related to anticipated reward in frontal cortex:

does it represent value or reflect motivation? Ann N Y Acad Sci 1121:431–446.

Saffran JR, Aslin RN, Newport EL (1996) Statistical Learning by 8-Month-Old Infants. Science

274:1926–1928.

Saleem KS, Miller B, Price JL (2014) Subdivisions and connectional networks of the lateral

prefrontal cortex in the macaque monkey. Journal of Comparative Neurology 522:1641–

1690.

Sallet J, Mars RB, Noonan MP, Neubert F-X, Jbabdi S, O'Reilly JX, Filippini N, Thomas AG,

Rushworth MF (2013) The Organization of Dorsal Frontal Cortex in Humans and

Macaques. J Neurosci 33:12255–12274.

Seidlitz J, Sponheim C, Glen D, Ye FQ, Saleem KS, Leopold DA, Ungerleider L, Messinger A

(2018) A population MRI brain template and analysis tools for the macaque. NeuroImage

170:121–131.

Shima K, Isoda M, Mushiake H, Tanji J (2007) Categorization of behavioural sequences in the

prefrontal cortex. Nature 445:315–318.

Sirmpilatze N, Klink PC (2020) RheMAP: Non-linear warps between common rhesus macaque

brain templates. Available at: https://zenodo.org/record/3668510 [Accessed June 28,

2022].

Soch J, Allefeld C (2018) MACS – a new SPM toolbox for model assessment, comparison and

selection. Journal of Neuroscience Methods 306:19–31.

Uhrig L, Dehaene S, Jarraya B (2014) A Hierarchy of Responses to Auditory Regularities in the

Macaque Brain. J Neurosci 34:1127–1132.

Van Essen DC, Drury HA, Dickson J, Harwell J, Hanlon D, Anderson CH (2001) An Integrated

Software Suite for Surface-based Analyses of Cerebral Cortex. Journal of the American

Medical Informatics Association 8:443–459.

Vanduffel W, Farivar R (2014) Functional MRI of Awake Behaving Macaques Using Standard

Equipment. IntechOpen. Available at:

https://www.intechopen.com/chapters/undefined/state.item.id [Accessed June 2, 2022].

Vanduffel W, Fize D, Mandeville JB, Nelissen K, Van Hecke P, Rosen BR, Tootell RBH, Orban

GA (2001) Visual motion processing investigated using contrast agent-enhanced fMRI in

awake behaving monkeys. Neuron 32:565–577.

Vergnieux V, Vogels R (2020) Statistical Learning Signals for Complex Visual Images in

Macaque Early Visual Cortex. Frontiers in Neuroscience 14 Available at:

https://www.frontiersin.org/article/10.3389/fnins.2020.00789 [Accessed June 15, 2022].

Wallis JD, Anderson KC, Miller EK (2001) Single neurons in prefrontal cortex encode abstract

rules. Nature 411:953–956.

Wang L, Amalric M, Fang W, Jiang X, Pallier C, Figueira S, Sigman M, Dehaene S (2019)

Representation of spatial sequences using nested rules in human prefrontal cortex.

NeuroImage 186:245–255.

Wang L, Uhrig L, Jarraya B, Dehaene S (2015) Representation of numerical and sequential

patterns in macaque and human brains. Curr Biol 25:1966–1974.

Warden MR, Miller EK (2010) Task-Dependent Changes in Short-Term Memory in the

Prefrontal Cortex. J Neurosci 30:15801–15810.

Wen T, Duncan J, Mitchell DJ (2020) Hierarchical Representation of Multistep Tasks in

Multiple-Demand and Default Mode Networks. J Neurosci 40:7724–7738.

Wendelken C, Chung D, Bunge SA (2012) Rostrolateral prefrontal cortex: Domain-general or

domain-sensitive? Human Brain Mapping 33:1952–1963.

White IM, Wise SP (1999) Rule-dependent neuronal activity in the prefrontal cortex. Exp Brain

Res 126:315–335.

Xu R, Bichot NP, Takahashi A, Desimone R (2022) The cortical connectome of primate lateral

prefrontal cortex. Neuron 110:312-327.e7.

Zhao F, Wang P, Hendrich K, Ugurbil K, Kim S-G (2006) Cortical layer-dependent BOLD and

CBV responses measured by spin-echo and gradient-echo fMRI: insights into

hemodynamic regulation. NeuroImage 30:1149–1160.

  Extended Data

Figure 3-1. Overlaid individual monkey contrasts for monkeys W, J, and B. Three second level contrasts, one

for each monkey, were created using only data bins that contained runs from each animal (W = 9, J = 6, B = 7 bins

each). As described in Materials and Methods, each bin contained approximately 10 runs. A liberal height threshold

of $p < 0.05$ was chosen for illustrative purposes before applying and extent (extents listed for each monkey by

contrast) and false discovery rate (FDR) error cluster corrected for multiple comparisons to $p < 0.05$. Slice number

in the y direction is listed under each coronal section. **A.** Voxel wise contrast of Rule Deviants > New Items, Same

Rule (NISR) (W extent = 560, J = 1296, B = 623 voxels). **B.** Voxel wise contrast of Number Deviants > NISR (W

962   extent = 772, J = 750, B = 667 voxels). **C.** Voxel wise contrast of Unique Ramp, Rule Deviants > NISR (W extent =

963   560, J = 630, B = 581 voxels).

966   **Figure 4-1. Example Ramp and Last Item regressors through the orthogonalization process.** Note that SPM

967   first creates regressors from onsets (in seconds, shown in **A.**) in samples using higher resolution to be convolved,

968   and then they are down-sampled before being orthogonalized and entered in the GLM.