Temporally resolved kinase regulatory networks control endothelial barrier

integrity

Ling Wei,[1] Selasi Dankwa,[1] Kamalakannan Vijayan,[1] Joseph D. Smith,[1,2,3,*] and Alexis

Kaushansky[1,2,3,4,*]

[1]Center for Global Infectious Disease Research, Seattle Children's Research Institute,

Seattle, WA 98109, USA

[2]Department of Pediatrics, University of Washington, Seattle, WA 98105, USA

[3]Department of Global Health, University of Washington, Seattle, WA 98105, USA

[4]Lead contact

*Correspondence: alexis.kaushansky@seattlechildrens.org (A.K.),

joseph.smith@seattlechildrens.org (J.D.S.)

SUMMARY

Breakdown of the blood-brain barrier is triggered by a range of physiological and pathological

stimuli and is detrimental for brain function. Yet, the underlying signaling networks regulating

barrier integrity are incompletely understood. Here, we present a novel and generalizable tool,

Temporally REsolved KInase Network Generation (TREKING), that combines machine learning

and network reconstruction to build time-resolved, functional phosphosignaling networks. We

investigated kinase-driven pathways that modulate barrier permeability in brain endothelial cells

in the presence of inflammatory stimuli. Our results reveal that >100 kinases are functional

during barrier insult and provide time-resolved molecular insights into the differential networks

that drive barrier disruption and recovery. The resulting models suggest a multi-layered rewiring

of thrombin-induced barrier disruption following TNFα priming, including the timing of common

signaling pathways, condition-specific phosphosignaling networks, and a skewing at major

signaling hubs towards barrier-weakening activities. TREKING provides a novel tool for

dissecting temporal phosphosignaling networks in biological systems.

Keywords: Endothelial barrier permeability, thrombin, TNFα, kinase regression, self-organizing

map, phosphosignaling network

INTRODUCTION

Homeostasis of the central nervous system (CNS) is tightly regulated by the blood-brain barrier (BBB). Dysregulation of the BBB can cause ions, neurotoxic plasma components, pathogens and immune cells to enter the CNS, which leads to massive disruption and degeneration (Daneman and Prat, 2015). BBB derangements are associated with the pathogenesis of a variety of diseases, including infectious causes such as cerebral malaria and non-infectious causes such as multifactorial neurodegenerative diseases (Miller et al., 2013; Zhao et al., 2015).

The permeability of endothelial barriers is tightly regulated via cell-cell junctions and focal adhesions. These points of cell-cell and cell-substrate contact are controlled by contractile mechanisms, whose functions are heavily regulated by phosphosignaling (Mehta and Malik, 2006). Kinases, via phosphorylation, can rapidly (on the scale of seconds to minutes) alter cellular states in response to barrier-disruptive stimuli and can promote long-term regulatory functions during barrier recovery (Kuppers et al., 2014). Whereas kinases that mediate barrier function have been intensively explored, work to date has largely focused on characterizing specific molecular mediators of barrier perturbation, without evaluating their complex network-level connectivity with other functional modules (Dankwa et al., 2021; Komarova et al., 2017; Kuppers et al., 2014; Mehta and Malik, 2006). Moreover, there is limited understanding of the phosphosignaling pathways involved in restoration of barrier integrity. Understanding the temporal dynamics of barrier regulation is critical for designing successful targeted therapeutic interventions.

During infection or disease, endothelial cells must integrate complex inflammatory signals. For instance, both thrombin, a multi-functional enzyme, and TNFα, an inflammatory cytokine, can promote morphological remodeling of the endothelium that leads to the loss of barrier integrity (Marcos-Ramiro et al., 2014; Oldenburg and de Rooij, 2014). TNFα exacerbates thrombin-induced barrier disruption (Anrather et al., 1997; Liu et al., 2004; Tiruppathi et al.,

2001), which strongly indicates that endothelial signaling pathways and/or their kinetics are

altered by combined inflammatory stimuli. High-resolution and time-resolved phosphoproteomic

analyses have cataloged phosphorylation events that occur in response to thrombin or TNFα-

induced endothelial activation (Beguin et al., 2019; van den Biggelaar et al., 2014). Separate

efforts have developed computational tools and modeling approaches that use -omics data for

network generation, including logic modeling such as Boolean logic models and logic ordinary

differential equations (ODEs), and enrichment-based approaches such as kinase-substrate

enrichment analysis (KSEA) (Casado et al., 2013; Schafer et al., 2019; Vaga et al., 2014).

However, these computational tools typically rely on large-scale phosphoproteomics data that

can typically not be collected with fine temporal resolution. Furthermore, proteomics only

indirectly assesses kinase functionality, as inferred by phosphorylation states. Given these

limitations, existing network-generation tools have not yet been applied to elucidating the time-

resolved molecular functional underpinnings of endothelial barrier integrity.

Kinase regression (KiR) combines a chemical screen with elastic net regression to

broadly interrogate the function of hundreds of kinases in a cellular phenotype (Arang et al.,

2017; Dankwa et al., 2021; Gujral et al., 2014). We have previously shown that kinase inhibitors

have dramatically different effects on endothelial barrier permeability and their effects exhibit

temporal features in barrier disruption and barrier recovery phases (Dankwa et al., 2021),

suggesting that different sets of kinase regulators are functional at different stages of barrier

perturbation. Here, we develop a novel approach, Temporally REsolved KInase Network

Generation (TREKING), to investigate phosphosignaling in disrupted endothelial monolayers

induced by thrombin in the presence or absence of TNFα pre-conditioning. TREKING allows 1)

assessment of time-resolved kinase functionality associated with barrier regulation, 2)

construction of phenotype-driving kinase-mediated phosphosignaling networks that recapitulate

signaling events at different stages of barrier perturbation, and 3) systematic comparison of

phosphosignaling between conditions. Our results reveal commonalities and divergences of

kinase signaling in response to classic pro-inflammatory stimuli, and insight into signaling

pathways with barrier-weakening or barrier-strengthening activity. This novel approach is easily

generalizable to other biological systems and will be particularly useful in deconvolving

condition-specific, temporally resolved signaling mechanisms important for many phenotypes of

interest.

RESULTS

TNFα pre-conditioning alters kinase determinants of thrombin-induced barrier

perturbations

In monolayers of human brain microvascular endothelial cells (HBMECs), TNFα pre-

conditioning exacerbated thrombin-induced barrier disruption, as demonstrated by the increased

maximum barrier permeability and prolonged barrier recovery (Fig. 1A). While this finding is

consistent with previous reports (Tiruppathi et al., 2001), the molecular networks that regulate

these differences remain unknown.

Here, we introduce TREKING, a generalizable methodology that leverages temporal KiR

to build time-resolved networks of kinase signaling from temporal datasets (Fig. S1). First, to

identify kinases involved in barrier integrity, a KiR screen was conducted with 28 kinase

inhibitors for their impact on barrier permeability in monolayers of HBMECs treated with

thrombin (-/+TNFα pre-conditioning) (Dankwa et al., 2021). Kinase inhibitors were added six

minutes after thrombin treatment and barrier permeability was assessed in real time using

xCELLigence. Different kinase inhibitors blunt, have minimal effect, or exacerbate thrombin-

induced barrier disruption and exhibit temporal features during the 6-hour time window (Fig. 1B)

(Dankwa et al., 2021).

We reasoned that if identical kinases control the barrier integrity in the two inflammatory

conditions, we expect perfect correlation between the activities of the 28 kinase inhibitors in

thrombin alone and TNFα pre-conditioned settings. If entirely different kinases control the barrier

integrity in the two conditions, we would expect little correlation between the activities of the

kinase inhibitors. Thus, the correlation between kinase inhibitor activities serves as an indirect

assessment of the similarity of the underlying barrier regulatory phosphosignaling networks

across conditions. Using area under the curve (AUC) of the normalized cell index as the metric,

we compared the barrier permeability between +thrombin and TNFα pre-conditioning +thrombin

conditions across the following time windows: maximum barrier disruption (0-10 minutes), early

barrier recovery (25-35 minutes), late barrier recovery (115-125 minutes for +thrombin

condition, 235-245 minutes for TNFα pre-conditioning +thrombin condition). The correlation was

the lowest within the first ten minutes of kinase inhibitor treatment (Pearson's r = 0.72) and

increased during early barrier recovery (Pearson's r = 0.90 to 0.91) (Fig. 1C-D). This suggests

that phosphosignaling networks are more divergent during barrier disruption, whereas more

similar phosphosignaling networks regulate barrier restoration. The correlation reduced in the

late barrier recovery phase (Pearson's r = 0.85 to 0.77) (Fig. 1C-D), perhaps suggesting that a

subset of different kinase networks regulate the transition to barrier homeostasis after injury.

This analysis is consistent with a model that TNFα pre-conditioning alters kinase regulatory

networks that dictate the extent and kinetics of thrombin-induced barrier disruption and recovery

in brain endothelial cells.

Time-resolved predictions of barrier-weakening and barrier-strengthening kinases

between the two inflammatory conditions

Previously, we predicted 29 and 25 kinases that regulate the HBMEC barrier in response to

thrombin treatment with or without TNFα pre-conditioning, respectively. Fifteen kinases were

shared between the two conditions (Dankwa et al., 2021). However, these KiR predictions were

based on the full AUC (6-hour time window, post-thrombin treatment), and did not take into

account when a kinase was functionally active (i.e., a "functional window"). To address this lack

of temporal resolution, we performed a temporal KiR (tKiR), using a sliding window analysis; a

5-minute sliding time window was applied to the normalized cell index data, which slides at 1-

minute steps for the first two hours after kinase inhibitor treatment and at 5-minute steps

afterwards (Fig. 2A). Among the 300 protein kinases used for tKiR (STAR Methods), the number

of functional kinases increased to 120 and 108 in HBMECs treated with thrombin -/+TNFα pre-

conditioning, respectively (Fig. 2A, Table S1). The higher resolution of tKiR is because the

sliding window analysis allows for the detection of kinases whose activity is brief or sporadic. In

total, 159 kinases (~30% of the human kinome) were predicted to regulate barrier function

within the 6-hour timeframe, with 69 kinases being common to the two conditions (Fig. 2A).

As a further refinement, the elastic net models were used to predict the direction of

barrier activity (weakening or strengthening) in a time-resolved manner (Fig. 2B-C, Fig. S1). For

HBMECs not pre-conditioned with TNFα, 53 and 60 kinases were predicted to play a barrier-

weakening and barrier-strengthening roles, respectively (Fig. 2B). In addition, seven kinases

were predicted to play both barrier-weakening and barrier-strengthening roles during different

time windows; we termed these "switch kinases" (Fig. 2B). For HBMECs pre-conditioned with

TNFα, 58 and 47 kinases were predicted respectively to be barrier-weakening and barrier-

strengthening, and 3 switch kinases were predicted (Fig. 2C). Overall, a greater number of

barrier-weakening kinases were predicted early during the time course, whereas barrier-

strengthening kinases were predominant during late barrier recovery and plateau phases (Fig.

2D). Nevertheless, multiple barrier-strengthening kinases were identified at the earliest time

windows following thrombin treatment, indicating that cells propagate both types of barrier

activities concurrently (Fig. 2D).

154 **The tKiR predictions of kinase activity recapitulate the canonical signaling pathways**

155 **involved in barrier regulation**

156 Assessing position of predicted kinases within canonical signaling pathways is one way to

157 evaluate the integrity of tKiR. For instance, if kinase A phosphorylates kinase B, we would

158 expect kinase A to be active at earlier time points than kinase B. Previous work has established

159 that mitogen-activated protein kinase (MAPK) cascades are involved in thrombin signaling,

160 including the extracellular signal-regulated kinases (ERKs), Jun N-terminal kinases (JNKs) and

161 p38 MAPKs (Mehta and Malik, 2006; Minami et al., 2004; Radeva and Waschke, 2018). The

162 topology of MAPKs is well-established; cell surface receptors such as receptor tyrosine kinases

163 (RTKs) provide a link between extracellular stimuli and the cascade (Morrison, 2012). After

164 activation, most MAPK pathways have a four-tier kinase architecture (MAP3K-MAP2K-MAPK-

165 MAPKAPK) (Pimienta and Pascual, 2007). Given the strict directionality of the network, we

166 reasoned that RTKs would be predicted to regulate the barrier during the earliest time points

167 after thrombin treatment, followed sequentially by MAP3Ks, MAP2Ks, MAPKs and then

168 MAPKAPKs.

169 The ERK, JNK, and p38 pathways were all implicated in thrombin signaling, with multiple

170 kinases in each signaling cascade predicted by tKiR (Fig. 3). The predicted kinases could be

171 assembled into pathways that were consistent with known information flow through the MAPK

172 cascade. Overall, the ERK and JNK pathways were associated with barrier-weakening activity

173 and the p38 pathway had a mixture of barrier-weakening and barrier-strengthening activities

174 mediated by different p38 family members (Fig. 3). For example, within the ERK pathway, all the

175 predicted kinases were barrier-weakening, and their activity profiles followed the expected order

176 of MAP2Ks preceding MAPKs, which preceded four downstream MAPKAPKs (Fig. 3). Likewise,

177 a mid-to-late acting barrier-strengthening p38 pathway could be assembled from known

178 components (Fig. 3).

The same MAPK pathways were predicted with TNFα pre-conditioning (Fig. 3).

However, the timing and duration of the ERK and JNK pathways were shifted earlier and

shortened for most kinases. We also observed a second wave of barrier-weakening activity by

MAPK1/ERK2, and multiple additional kinases in the JNK pathway were predicted, consistent

with previous work indicating that combined TNFα and thrombin treatment alters the magnitude

and duration of JNK phosphorylation activation marks in endothelial cells (Liu et al., 2004).

Furthermore, the p38 pathway became nearly entirely barrier-strengthening with TNFα pre-

conditioning and MAPKAPK2/MK2 was no longer predicted to be a switch kinase with dual

barrier-weakening and barrier-strengthening activities. Consequently, this analysis suggests that

core MAPK signaling pathways involved in thrombin-induced barrier disruption remain intact

after TNFα pre-conditioning, but the timing, duration, and importance of specific MAPKs are

altered.

To experimentally assess the model, we evaluated kinase phosphorylation of a subset of

tKiR predicted kinases in ERK, JNK and p38 pathways via western blot. HBMECs (-/+TNFα pre-

conditioning) were treated with thrombin for 0, 5, 15, 30, 60, 120, 180, 240 or 360 minutes. In

each experimental condition, the level of phosphorylated kinase after thrombin treatment was

compared to its phosphorylation at basal level (media only). By western blot, increased levels of

activated kinases were detected in each of the three MAPK pathways at different time frames

during barrier disruption and barrier recovery (Fig. 4A). In both +thrombin and TNFα pre-

conditioning +thrombin conditions the activity of MAP2K1/2 (MEK1/2) peaked at an earlier time

point compared to its downstream targets MAPK1/ERK2 and MAPK3/ERK1, which supports the

tKiR model predictions of early barrier-weakening functionality of these MAPKs (Fig. 4A).

Results from western blot also showed that the activity of MAPK9/JNK2 increased upon

thrombin treatment and remained higher than the basal level six hours after thrombin treatment,

at which point the barriers were mostly recovered (Fig. 4A). Similarly, the activity of p38 MAPKs

was increased shortly after thrombin treatment, and in both +thrombin and TNFα pre-

conditioning +thrombin conditions an early (barrier disruption phase) and a late (late barrier

recovery phase) activity peak were detected. Overall, in most cases there was increasing

phosphorylation immediately preceding or overlapping corresponding time windows of predicted

barrier activity for all three canonical MAPK pathways (Fig. 4B).

Construction of phosphosignaling networks that mediate barrier disruption

To investigate how endothelial cells propagate phosphosignals to and from key kinase

regulators beyond canonical linear cascades, we reconstructed kinase phosphosignaling

networks by first using self-organizing maps (SOMs), a dimensionality reduction method that

can capture the topographic relationships of time series data (Kohonen, 1982). Kinases of

similar temporal functionality were clustered into one neuron, and neurons having similar mean

temporal functionality are more closely located on the SOM (Fig. 5A, Fig. S2A-C). To generate

the SOMs, we used a grid size of 6×6 (resulting in a maximum of 36 neurons). In +thrombin

condition, we obtained 30 neurons and each neuron contained between 1 and 11 kinases (Fig.

5A, Fig. S2D, Table S2). After TNFα-preconditioning, we obtained 35 neurons and each neuron

contained between 1 and 9 kinases (Fig. S2A, D, Table S2).

Since kinases assigned to the same neuron exhibit similar temporal barrier functionality,

we reasoned that they are more likely to act within a connected signaling module. To build

phosphosignaling pathways and investigate systems-level interconnections, we built TREKING

models of barrier phosphosignaling networks (Fig. S1, Fig. S2E). For each neuron a local

phosphosignaling network was built to describe the paths through which phosphosignals may

propagate, using the kinase-substrate phosphorylation database PhosphoSitePlus® to predict

upstream and downstream kinases and to infer intermediate kinases in the pathways (Fig. S2E)

(Hornbeck et al., 2015). We were able to build 26 local networks from the 30 neurons generated

in +thrombin condition, and 27 local networks from the 35 neurons generated in TNFα pre-

conditioning +thrombin condition (Table S2). The remaining neurons included only a single

kinase that does not self-phosphorylate or multiple unconnected kinases. The size and topology

of the phosphosignaling networks varied across neurons. In +thrombin condition, most networks

have a maximum node-to-node shortest path length below 10 kinases, while the deepest

networks built from neurons (0,0) and (3,3) have a maximum shortest path length of 14 kinases

(Fig. S2E). In TNFα pre-conditioning +thrombin condition, most networks have a maximum

shortest path length ranging from 3 to 9 kinases, except for the network built from neuron (2,4)

that has a maximum shortest path length of 16 kinases (Fig. S2E).

The SOM analysis revealed substantial interconnections between MAPK and NF-κB

signaling pathways in both barrier-weakening and barrier-strengthening activities. For instance,

in +thrombin condition, the barrier-weakening neurons (0,0) and (1,0) include a mixture of ERK,

JNK, and canonical NF-κB signaling pathways with MAPK3/ERK1, CHUK/IKKα and

MAPKAPK3/MK3 in neuron (0,0) and MAPK1/ERK2 and MAPK8/JNK1 in neuron (1,0) (Fig 5A-

B, Table S2). Likewise, a mid-to-late barrier-strengthening neuron (5,5) contained the switch

kinase MAPKAPK2/MK2 in a p38 pathway (MAP3K20/ZAK-MAPK14/p38α-MAPKAPK2/MK2),

CSK (C-terminal Src kinase; a negative regulator of Src-family kinases (SFKs)), and two

members of non-canonical NF-κB signaling pathways (MAP3K14/NIK and IKBKB/IKKβ) (Fig.

5C-D). Therefore, MAPK and NF-κB interconnections were implicated in both barrier disruption

and barrier recovery phases. Notably, with TNFα pre-conditioning, this barrier-strengthening

pathway shrunk to MAP3K20/ZAK-MAPK14/p38α (p38 pathway) and MAP3K14/NIK (NFκB

signaling pathway). Overall, 77 of the 120 predicted kinases in +thrombin condition and 61 of

the 108 predicted kinases in TNFα pre-conditioning +thrombin condition could be assembled by

TREKING into signaling pathways of two or more kinases that acted with similar temporal

kinetics and identified potential network-level interconnections between individual barrier

signaling pathways within the broader signaling networks.

TNFα pre-conditioning alters the landscape of kinase regulators and associated

pathways in the barrier disruption and barrier recovery phases

To further investigate the phosphosignaling networks associated with thrombin-induced barrier

perturbation, and how these networks were altered in response to TNFα pre-conditioning, we

decided to investigate barrier disruption and barrier recovery phases separately. To investigate

networks that led to barrier disruption, we first built the composite phosphosignaling networks

important for barrier regulation during the barrier disruption phase (Fig. S6, Fig. S7).

Specifically, networks built from neurons that were functional within the first 30 minutes after

thrombin treatment (-/+TNFα) were combined to form a composite network that illustrates the

connections between functional kinases in the barrier disruption phase. Interestingly, the

composite networks reveal overlapping kinase nodes within barrier-weakening and barrier-

strengthening pathways, illustrating nexuses where specific kinases have the capacity to lead to

both phenotypes (Fig. S6, Fig. S7).

In the first 30 minutes after thrombin treatment, more kinases were predicted to be

barrier-weakening in TNFα pre-conditioning +thrombin condition (40 kinases) than in +thrombin

condition (33 kinases), and eighteen barrier-weakening kinases were shared, including multiple

members of the MAPK family (Fig. S3A). Among the twenty-two kinases uniquely predicted in

TNFα pre-conditioning +thrombin condition were all three members of the Janus kinase (JAK)

family (JAK1, JAK2 and JAK3) (Fig. S3A), which are known to be activated by TNFα (Guo et al.,

2016).

At the network level, pre-conditioning with TNFα led to a multi-layered rewiring of

phosphosignaling including a larger fraction of barrier-weakening signaling pathways, which is

manifested both as more barrier-weakening kinase regulators (Fig. S3A) and more barrier-

weakening kinase connections (Fig. S6, Fig. S7) in the barrier disruption phase. For instance,

the connectivity (in/out-degrees of freedom) of kinase nodes is skewed towards barrier-

weakening connections with TNFα pre-conditioning (Fig. S3B). Moreover, the ratio of barrier

signaling activities shifted at major signaling hubs. For example, AKT1 is the most highly

connected kinase in the composite networks. Whereas a mixture of signals flow through AKT1

under both inflammatory conditions, barrier-weakening connections predominate with TNFα pre-

conditioning (Fig. S3C). This suggests that connectivity of AKT1 is rewired to associate with

factors that promote barrier disruption when cells are pre-conditioned with TNFα. Similarly,

MAPK1 is balanced between barrier-weakening and barrier-strengthening connections when

thrombin is treated alone but skewed towards barrier-weakening connections in the context of

TNFα pretreatment (Fig. S6, Fig. S7). Thus, TREKING describes network-level mechanisms by

which barrier-weakening activities might be favored after TNFα pre-conditioning.

In addition to repurposing the phosphosignaling network towards barrier-weakening

activities in the barrier disruption phase, TNFα priming also altered the network during the

barrier recovery phase (Fig. 1). We reasoned that barrier-strengthening signaling activities may

begin prior to the repair phase, so we constructed composite networks for the full 6-hour time

course (Fig. S8, Fig. S9). At the network level, pre-conditioning with TNFα led to a reduction in

the fraction of barrier-strengthening pathways over the entire 6-hour time course. There were

several notable differences between the barrier-strengthening networks that might describe the

delayed recovery in the presence of TNFα. In the absence of TNFα, networks involving

megakaryocyte-associated tyrosine kinase (MATK/CTK) and CSK, both negative regulators of

SFKs, are engaged (Fig. S8). During early barrier recovery, the MATK pathway (EPHA1-

MATK/CTK-ABL1-FYN-PDGFRA-ARAF) was predicted to be functional, then later the CSK

pathway (CSK-MAPK14/p38α-MAPKAPK2/MK2) was predicted to drive barrier recovery (Fig.

5D, Fig. S8). Together, these pathways were predicted to be functional throughout barrier

recovery and drive a robust and sustained repair in the absence of TNFα pre-conditioning. In

contrast, when cells are pre-conditioned with TNFα, the activity of the CSK pathway is lost (Fig.

S9), providing one hypothesis of why the kinetics of late barrier recovery is distinct after TNFα

pre-conditioning. Taken together, this analysis suggests that TNFα pre-conditioning both

enhances the number of barrier-weakening networks and rewires barrier-strengthening

regulators to alter the magnitude and duration of barrier disruption.

TREKING predicts barrier-specific kinase functionality with temporal resolution

The TREKING model (Fig. S1) predicts activity windows for kinases and the networks through

which phosphosignals propagate with temporal resolution. To systematically evaluate the

accuracy of the networks predicted to underlie the barrier functionality in response to thrombin,

we first used the TREKING model to generate time-resolved predictions of the activation states

of kinases in response to thrombin (-/+TNFα pre-conditioning). By considering all the signaling

connections that a kinase is associated with, we can comprehensively explore its role in barrier

regulation over time. To assess the barrier function of a kinase at a particular time point, a 10-

minute time window centered at that time point was applied (Fig. S5A). If any of the signaling

pathways that the kinase is associated with are active within the time window, the model

predicts the kinase to be functionally active within that time window; otherwise, the model

predicts the kinase to be functionally inactive within the time window (Fig. S5A).

To evaluate the accuracy of the TREKING model for predicting kinase activity, we

treated HBMECs with thrombin in the presence or absence of TNFα pre-conditioning. We

collected cellular lysates at nine time points and evaluated activation of the 14 kinases using

western blot (Table S3). TREKING predicted 78 instances of active barrier function across all

time points in the absence of TNFα; 61 differential phosphorylation events were observed by

western blot (Fig. 6A, Fig. S5B). TREKING predicted 71 instances of active barrier function

across all time points after thrombin treatment with TNFα pre-conditioning; we detected 47

differential phosphorylation events by western blot (Fig. 6A, Fig. S5B). These data suggest that

TREKING outperforms random models (STAR Methods, Fig. 6B, Fig. S5C-D). TREKING

highlights many examples of altered kinase functionality after TNFα pre-conditioning (Fig. 6A,

Fig. S4). For instance, PTK2/FAK was predicted to have sustained activity in the absence of,

but not in the presence of TNFα pre-conditioning. PTK2/FAK also exhibits reduced connectivity

after TNFα pre-conditioning (Fig. 6A, Fig. S4). The western blot data were consistent with the

model predictions that without TNFα pre-conditioning the activity of PTK2/FAK was different

from non-treated HBMECs at all time points assessed, while in TNFα pre-conditioned cells a

significant reduction in phosphorylation was detected only in the barrier disruption phase (Fig.

6A, Fig. S4).

For highly connected kinases, such as SRC, p38 MAPK, PDPK1, and AKT1, TREKING

predicted that upon thrombin treatment (-/+TNFα pre-conditioning) they actively participate in

barrier regulation throughout the time course, although the kinases and/or signaling pathways

that they interact with are altered with TNFα pre-conditioning (Fig. S8, Fig. S9). Overall, the

accuracy of TREKING predictions vary by specific kinase. For example, active SRC, p38 MAPK

and PDPK1 were observed by western blot at most time points predicted by TREKING, but

TREKING failed to make predictions on AKT1 in the presence of TNFα, perhaps due to the

complexity of cell signaling through this extensively connected kinase (Fig. S5B).

Most previous and ongoing research has focused on certain stages of barrier

perturbation or sampled sparsely during a time course. We therefore asked how TREKING

predictions compared to the existing literature in their breadth and temporal resolution. To do

this, we visualized portions of the phosphosignaling network that had been described in the

literature to regulate the endothelial barrier in response to thrombin (STAR Methods, Table S4).

We then performed the same visualization with kinases that had been predicted by tKiR, and

finally, by TREKING. When comparing to the literature, TREKING substantially increased the

number of kinases associated with thrombin-induced barrier regulation and was able to

assemble most predicted kinases into a time-resolved phosphosignaling network (Fig. 7). This

highlights the power of TREKING to broadly dissect the functionally important phophosignaling,

with high temporal resolution, in barrier regulation.

DISCUSSION

Kinase signaling pathways are highly implicated in endothelial barrier regulation (Komarova et

al., 2017; Kuppers et al., 2014; Mehta and Malik, 2006) and consequently kinase inhibitors are

being explored for treatment of vascular injury (Aman et al., 2012; Botros et al., 2020; Rizzo et

al., 2015). However, a lack of resolution that describes the barrier activity of specific kinase

signaling networks and their importance in barrier disruption and repair has hindered therapeutic

applications. Here, we introduce TREKING, which uses kinase regression to broadly interrogate

kinase signaling pathways involved in barrier regulation with temporal resolution and to study

their interconnectedness in signaling networks.

In agreement with previous work (Mehta and Malik, 2006), our results using tKiR

implicated multiple MAPK signaling pathways in thrombin-induced barrier changes. Because of

the systematic nature of the tKiR, it was possible to assign barrier functions to specific MAPK

pathways with ERK and JNK pathways involved in early barrier disruption and the p38 pathway

in a mid-to-late acting barrier restoration. In each case, multiple kinases within each pathway

were predicted and the barrier phenotypes followed the expected flow of phosphosignaling in

the linear cascade, and kinase activation by western blots was consistent with these predictions.

Beyond MAPKs, our findings highlight the complexity of endothelial barrier regulation with over

100 kinases were predicted to regulate thrombin-induced change of barrier integrity. With tKiR,

"druggable windows" of the key kinase regulators mediating barrier permeability can be

identified, which provides valuable insight into designing therapeutic interventions to block

barrier leakage and restore barrier functionality, or in some circumstances, to temporarily disrupt

the barrier for controlled drug delivery into the brain (Hashimoto et al., 2020; Luo and Shusta,

2020; Yanagida et al., 2017).

Kinase activity is frequently inferred by assessing phosphorylation via western blot or

mass spectrometry. However, kinase activity is distinct from function and many kinases play

multiple distinct biological roles by targeting different sets of substrates with different kinetics.

While this behavior was coined "moonlighting", suggesting there was one "main" kinase function

with other ancillary activities, our data suggests that instead kinases can have multiple, equally

important, and sometimes opposing functions in regulating the barrier. Since TREKING predicts

functionality, rather than enzymatic activity, our work highlights these higher-order, complex

activities of kinases. By reconstructing networks with temporal resolution, TREKING enables us

to systematically dissect the "functional windows" of the kinases associated with barrier

regulation.

We termed a subset of kinases "switch kinases", as TREKING predicted their

functionality to be barrier-weakening and barrier-strengthening during different kinetic windows.

As one example of a switch kinase, MAPKAPK2/MK2 was predicted to be barrier-weakening

during barrier disruption and then to switch to barrier-strengthening during late barrier recovery.

MAPKAPK2/MK2 is phosphorylated and activated by MAPK14/p38α upon activation of the p38

pathway. Activation of MAPKAPK2/MK2 leads to phosphorylation of LIMK1 and heat shock

protein beta-1 (HSPB1), both of which regulate actin filament dynamics in endothelial cells

(Gorovoy et al., 2005; Rada et al., 2021). Together, this suggests that the p38 pathway may

regulate cytoskeleton remodeling during both barrier disruption and repair through different

signaling pathways. The temporal functionality of kinases in the MAP3K20/ZAK-MAPK14/p38α-

MAPKAPK2/MK2 axis of the p38 pathway is coherent in response to thrombin alone, as

highlighted by our SOM analysis, where all three kinases were predicted to be barrier-

strengthening during mid-to-late barrier recovery. In contrast, with TNFα pre-conditioning,

MAPKAPK2/MK2 no longer promotes barrier strengthening but retains its early barrier-

weakening function. In parallel, we no longer observe MAPKAPK2/MK2 signaling through p38

pathway in this setting. Interestingly, our identifying of switch kinases could explain reports of

kinases having contradictory activities. For instance, Src has been described to have both

barrier-weakening and barrier-strengthening properties (Birukova et al., 2013; Garcia et al.,

2001; Han et al., 2013; Klomp et al., 2019; Knezevic et al., 2009; McVerry and Garcia, 2004;

Vouret-Craviari et al., 2002). Our model suggests that these disparate reports might simply be

describing different barrier signaling activities that are kinetically resolved.

Several examples illustrate additional molecular and kinetic details that can be obtained

from TREKING in contrast to traditional, less systematic approaches, including shifts in the

timing of core MAPK signaling pathways (ERK/JNK pathways), condition-specific

phosphosignaling networks (e.g., MAPKAPK2/MK2 switch kinase), and a skewing of barrier

signaling activities at the major signaling hubs (e.g., AKT1) that alters the biological function of a

key regulatory kinase. TREKING also highlights functional kinetics of cascades that are altered

in TNFα pre-conditioned cells, suggesting how long-term inflammatory signals may alter the

phosphosignaling network topology towards different barrier tolerances. These differences are

often buried by conventional investigations that use tools such as genetic knockdown or

knockout, which respond on the time scale of days rather than minutes. This highlights the

importance of developing tools to systematically study the molecular details of kinase-driven

signaling and to capture the differences in cell signaling between conditions.

The global, time-resolved and mechanistic details revealed by TREKING has

ramifications beyond our fundamental understanding of phosphosignaling during inflammation.

Beyond controlling the vasculature, TREKING is broadly generalizable to other cellular systems,

and could be applied towards developing a molecular, kinetically resolved picture of any cellular

phenotype.

Limitations of the study

Despite its power to elucidate kinase signaling with temporal resolution, TREKING is not without

limitations. Firstly, the kinase-compound biochemical data used for KiR contains information

only on a subset of the human kinome (300 of 518 kinases) (Anastassiadis et al., 2011). To

partially overcome this limitation, TREKING reconstructs phosphosignaling networks and

reveals functional kinases not predicted by tKiR and therefore can fill in missing gaps by

leveraging kinase proteomic datasets. As additional biochemical data is available, these models

can be refined. Secondly, PhosphoSitePlus®, the knowledgebase used for building the

phosphosignaling networks, despite being the most comprehensive catalog available to us,

does not cover the complete kinase-substrate phosphorylation interactions in human cells, nor

is it specific to endothelial cells.

ACKNOWLEDGEMENTS

We would like to thank Mary-Margaret Dols for experimental guidance. This work was funded by

National Institutes of Health (NIH) grants R01 AI148802 and R61 HL154250 (to J.D.S. and

A.K.).

AUTHOR CONTRIBUTIONS

Conceptualization, L.W., S.D., J.D.S., and A.K.; Funding acquisition, J.D.S. and A.K.;

Investigation, L.W. and S.D.; Methodology, L.W., S.D., and K.V.; Supervision, J.D.S. and A.K.;

Writing – Original Draft, L.W. with help from S.D., J.D.S. and A.K.; Writing – Review & Editing,

L.W., S.D., J.D.S., and A.K.

DECLARATION OF INTERESTS

The authors declare no competing interests.

Figure 1. Pre-conditioning with TNFα exacerbated thrombin-induced barrier disruption.

(A) Left: representative xCELLigence data showing barrier response to thrombin with or without

TNFα pre-conditioning. Right: Quantification of the maximum change and time to 50% recovery

for data on left. P-values from student's t-test are shown. (B) Graphic of KiR screen conducted

with 28 kinase inhibitors for their impact on barrier permeability in thrombin treated HBMECs (-

/+TNFα pre-conditioning). Monolayers of HBMECs were treated with thrombin followed at 6

minutes by kinase inhibitor treatment and barrier permeability was measured in real time using

xCELLigence system. (C) Correlations of the 28 kinase inhibitors between +thrombin and TNFα

pre-conditioning +thrombin conditions at different stages of barrier perturbation. Kinase

inhibitors had similar but distinct impact on barrier permeability in response to thrombin with or

without TNFα pre-conditioning. Area under the corresponding permeability curve (AUC) within

representative time windows (10-minute) is displayed as scatter plots (blue circle: DMSO-

treated; black circles: kinase inhibitor-treated; black solid line: linear fitting line excluding DMSO

control; black dashed line: identity line). The time windows corresponding to the scatter plots are

indicated in (A) with lavender shading. (D) Pearson correlation coefficients between AUC in

+thrombin and TNFα pre-conditioning +thrombin conditions. See also Figure S1.

Figure 2. Using kinase regression to inform kinases important for barrier regulation with

temporal resolution. (A) Left: schematic describing analysis of using temporal KiR (tKiR) to

predict kinases important for barrier regulation. A 5-minute sliding time window was applied to

the normalized cell index data from the xCELLigence assay conducted with the 28 kinase

inhibitors. The elastic net regularization algorithm predicted barrier "active" kinases within the

corresponding time window, based on area under the curve (AUC). Right: Venn diagram

showing the total number of kinases predicted in each condition across all time windows. (B)-(C)

Left: time-resolved barrier activity profiles of predicted kinases in (B) +thrombin and (C) TNFα

pre-conditioning +thrombin conditions. Blue and red colors represent barrier-weakening and

barrier-strengthening functionality, respectively. Predicted kinases are hierarchically clustered

using Euclidean distance metric. Right: number of kinases predicted to be barrier-weakening,

barrier-strengthening, or of dual functionality across the time course ("switch kinases"). (D) Venn

diagrams showing the number of kinases predicted to be barrier-weakening or barrier-

strengthening at different stages of barrier perturbation. See also Table S1.

Figure 3. Temporal KiR recapitulates the phosphosignaling events in canonical MAPK

pathways. Members of three canonical MAPK pathways (ERK, JNK, and p38) were predicted

to regulate barrier function in thrombin treated HBMECs with or without TNFα pre-conditioning.

Kinases were mapped onto the corresponding signaling cascades along with their temporal

functionality profiles, where blue and red shadings represent barrier-weakening and barrier-

strengthening activity, respectively. Solid arrows represent direct phosphorylation interactions

reported previously and dashed arrows represent indirect or inferred interactions. See also

Table S1.

Figure 4. MAPK pathways are activated in response to thrombin treatment. (A) HBMECs

were stimulated with TNFα (10 ng/ml) for 21 hours. A separate set of cells was left unstimulated.

Cells were then treated with thrombin (5 nM) for the indicated times. Phosphorylation of

members in three canonical MAPK pathways (ERK, JNK, and p38) was probed by western blot.

Cells pre-conditioned with TNFα but not treated with thrombin are labeled "0-T". Data was

normalized to non-treated, media only condition ("0-M"). GAPDH was used as loading control.

Arrows in ERK pathway indicate the maximum phosphorylation level among all the time points

investigated in this study. (B) Comparison of western blot measurements and tKiR predictions.

For each kinase, the top row is the mean fold change of its phosphorylation with respect to

basal level ("0-M"), where the arrows represent the direction (↑: increase; ↓: decrease) and

magnitude of the rate of fold change from the previous time point (fold change/minute) in log

scale (the fold change is set to 1 if it is not considered different from the basal level); the bottom

row is the tKiR predictions on the temporal barrier activity of kinases, with blue and red

representing barrier-weakening and barrier-strengthening functionality, respectively. White

indicates the kinase was not predicted by tKiR. "NE" stands for "not evaluated" by tKiR, since

kinase inhibitors were added six minutes after thrombin treatment and no model predictions

were done within that time window. Purple represents instances where different p38 kinases

with barrier-weakening or barrier-strengthening functionality were predicted at that time point.

See also Figure S4 and Table S3.

Figure 5. Building local phosphosignaling networks that describe thrombin-mediated

barrier regulation. (A) Self-organizing map (SOM) of kinases predicted in response to thrombin

without TNFα pre-conditioning (30 neurons). In each neuron, the black line is the mean

functionality of all the kinases within that neuron, with "+1" indicating all kinases are barrier-

strengthening and "-1" indicating all kinases are barrier-weakening. Blue and red shadings

represent barrier-weakening and barrier-strengthening functionality, respectively. (B) A

representative barrier-weakening neuron containing barrier-weakening kinases of similar

temporal barrier kinetics. (C) A representative barrier-strengthening-dominant neuron containing

barrier-strengthening kinases of similar temporal barrier kinetics, as well as a switch kinase

(MAPKAPK2/MK2). (D) The local phosphosignaling network reconstructed from neuron (5,5).

Kinases predicted by tKiR are labeled in orange. Kinases and connections associated with

MAPK signaling, Src-family signaling and NF-κB signaling are highlighted in pink, yellow and

cyan, respectively. See also Figure S2 and Table S2.

Figure 6. TREKING predicts kinase regulators and kinase-mediated signaling pathways

functionally important for barrier regulation. (A) Left: comparison of western blot

measurements, tKiR predictions and TREKING predictions on a subset of kinases involved in

thrombin-induced barrier regulation. For each kinase, the top row is the mean fold change of its

phosphorylation with respect to basal level (non-treated, media only), where the arrows

represent the direction (↑: increase; ↓: decrease) and magnitude of the rate of fold change from

the previous time point (fold change/minute) in log scale (the fold change is set to 1 if it is not

considered different from the basal level); the middle and bottom rows are the tKiR and

TREKING predictions on the temporal barrier activity of kinases, with black representing being

predicted by the models to be functional (barrier-weakening, barrier-strengthening, or dual

functionality) and white representing not being predicted by the models. "NE" stands for "not

evaluated". Right: connectivity (in/out-degrees of freedom) of kinases in the composite

TREKING network. Light orange and orange represent in- and out-degrees of freedom in

+thrombin condition, respectively; light red and red represent in- and out-degrees of freedom in

TNFα pre-conditioning +thrombin condition, respectively. (B) Overall prediction accuracy of

TREKING compared to random models. Pink: correct predictions; gray: incorrect predictions.

See also Figures S4 and S5, Tables S1, S2 and S3.

Figure 7. tKiR and TREKING expand the current understanding of kinases and associated

phosphosignaling in barrier regulation. The reported time points in the literature were

expanded to 6-minute time windows (3 minutes before/after the reported time points) to account

for any experimental variations. The background network in gray includes all kinase-kinase

phosphorylation interactions from the kinase-substrate phosphorylation database

PhosphoSitePlus®. Blue and red nodes are the kinases reported by literature or predicted by

tKiR to have barrier-weakening and barrier-strengthening functionality, respectively; purple

nodes are the kinases having conflicting literature reports. Blue and red edges are the

interactions predicted by TREKING that are associated with only barrier-weakening and barrier-

strengthening activity, respectively; purple edges are the interactions associated with both

barrier-weakening and barrier-strengthening activities. The "combined" panel in the bottom is

the composite literature reports or model predictions across all time points. See also Table S4

and Supplemental file 1.

STAR★METHODS

RESOURCE AVAILABILITY

Lead contact

Further information and requests for resources and reagents should be directed to and will be

fulfilled by the lead contact, Alexis Kaushansky (alexis.kaushansky@seattlechildrens.org).

Materials available

This study did not generate new unique reagents.

Data and code availability

Codes and data generated in this study are included in supplemental information.

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Cell lines

Primary human brain microvascular endothelial cells (HBMECs; Cell Systems Cat# ACBRI 376)

were cultured on collagen (5 µg/cm^2; Corning Cat# 354236) in HBMEC culture media (Lonza

Cat# CC-3202) at 37°C and 5% CO_2. HBMECs were used until passage 9.

METHOD DETAILS

xCELLigence data acquisition

The kinetic data in this study are published and were acquired from xCELLigence assays

(Dankwa et al. 2021), which are described here in brief. HBMECs were grown to confluency in

xCELLigence e-plates. On the day of the assay, HBMECs were equilibrated in serum-free

culture media for 1-2 hours, and thrombin was added at 5 nM. To capture thrombin-induced

barrier disruption, the cell index was measured every minute for 6 minutes, after which kinase

inhibitors were added in triplicate at 0.5 µM. The cell index was then measured every minute for

2 hours and thereafter, every 5 minutes for 4 hours. The assays with TNFα pre-conditioning

were performed identically except that HBMECs were activated with 10 ng/ml TNFα for ~22

hours before media equilibration. Data analysis was performed as described previously

(Dankwa et al. 2021) with some modifications. The cell index was normalized to baseline (0) at

the time point prior to addition of thrombin and area under the curve (AUC) was determined for

each kinase inhibitor treatment. For this study, the AUC was determined within 5-minute sliding

windows over the 6-hour time course. AUC values were normalized by subtracting the AUC of

cells treated with thrombin+DMSO or TNFα pre-conditioning +thrombin+DMSO. These values

were then linearly transformed, with the most negative normalized AUC value within a 5-minute

window being set to 0 and the normalized value for the control sample being set to 100.

Lysate preparation

HBMECs were seeded in 6-well plates at 55,000 cells/well and grown for 3 days. Cells were

then activated with 10 ng/ml TNFα for 21 hours or kept in fresh media for the same period. On

the day of lysate preparation, cells were equilibrated in serum-free culture media for 1 hour and

then treated with thrombin at a final concentration of 5 nM for 5, 15, 30, 60, 120, 180, 240, 360

minutes. After the indicated incubation periods, cells were washed twice with ice-cold phosphate

buffered saline (PBS) and lysed in sodium dodecyl sulfate (SDS) lysis buffer (50 mM Tris-HCl,

2% SDS, 5% glycerol, 5 mM ethylenediaminetetraacetic acid, 1 mM sodium fluoride, 10 mM β-

glycerophosphate, 1 mM phenylmethylsulfonyl fluoride, 1 mM sodium orthovanadate, 1 mM

dithiothreitol, supplemented with a cocktail of protease inhibitors (Roche Cat# 4693159001) and

phosphatase inhibitors (Sigma-Aldrich Cat# P5726)). Cell lysates were clarified in filter plates at

3500 rpm for 30 minutes, after which they were stored at -80°C until use.

Western blot

All the gel electrophoresis was performed using Bolt™ 4-12% Bis-Tris mini protein gels.

Proteins were transferred to PVDF membranes using iBlot 2 dry blotting system (Thermo Fisher

Scientific Cat# IB21001). Primary antibodies were used at concentrations recommended by the

vendor (see Table S3 for antibody dilutions). Antibody to GAPDH (Cell Signaling Technology

Cat# 97166) was used as loading control at 1:2000 dilution. Blots were imaged using Bio-Rad

ChemiDoc imaging system and signals were quantified using ImageJ2 (version 2.3.0).

Background correction was done for each band by subtracting background signals nearby the

band. The signals from proteins of interest were first normalized to the signals from GAPDH,

and then the signals at each time point were normalized to the signals at time zero for the fold

change of phosphorylation from basal level.

Elastic net regularization

The elastic net regularization algorithm used for this study was published previously (Dankwa et

al. 2021). For each time window, the algorithm was applied on the normalized AUC values and

kinases with non-zero coefficients were predicted to be informative for barrier permeability. The

sign of the coefficients indicates the functionality of the kinases in regulating barrier integrity,

with kinases having positive and negative coefficients being informed to have barrier-

strengthening and barrier-weakening function, respectively.

Self-organizing map

The NumPy-based SOM implementation MiniSom (https://github.com/JustGlowing/minisom)

was used to cluster the tKiR predicted kinases by their temporal characteristics. Grid size of 6×6

was used to generate SOMs.

Path search

Search for the shortest paths between kinases was done using NetworkX

(https://github.com/networkx/networkx), a Python package for analyzing complex network

structures.

Generation of random models

The random models were generated following the TREKING pipeline. During each iteration, a

random background phosphosignaling network was generated using the configuration model

function in NetworkX package, which creates random directed graphs with given degree

sequences. The random network contains 518 nodes, corresponding to the total number of

human protein kinases, and the in- and out-degree sequences followed the same distribution as

the background phosphosignaling network used in TREKING. All the kinases were shuffled and

randomly assigned to each of the nodes in the random network. At each time point, kinases of

barrier-weakening or barrier-strengthening functionality were randomly selected from the 300

kinases used for KiR, with probabilities the same as in the TREKING model at the

corresponding time point. The random sampling of kinases was performed at all time points

corresponding to the TREKING model. Each of the kinases selected at any time point during the

time course was randomly assigned to one of the 36 clusters, corresponding to the total number

of neurons generated by SOMs. For each cluster the phosphosignaling network was

reconstructed by searching for the shortest paths between any pair of kinases within the cluster,

using the random phosphosignaling network generated during the iteration as the background

network. In total, 100 random models were generated and used to compare with TREKING

predictions.

Literature search

To compare the scope of previous research with the current study, a comprehensive literature

search on protein kinases reported to regulate barrier function was done using the free search

engine PubMed (https://pubmed.ncbi.nlm.nih.gov). The search terms were "endothelial barrier

thrombin" plus kinase gene names, and the search results were filtered so that only articles

published on or after year 2000 were shown. The search was done on each of the 518 human

protein kinases. The search was not restricted to studies on HBMECs; instead, studies on

thrombin-induced barrier disruption using endothelial cell lines, primary endothelial cells isolated

from different organs, or *in vivo* studies using mice or rats, were all included. To compare with

our work, studies or data beyond a 6-hour time window (after thrombin treatment) were

excluded.

Visualization

Schematics were created with BioRender.com. Phosphosignaling networks were visualized

using Cytoscape (https://cytoscape.org), with the hierarchic layout from yFiles layout algorithms

(https://www.yworks.com/products/yfiles-layout-algorithms-for-cytoscape).

QUANTIFICATION AND STATISTICAL ANALYSIS

Statistical analysis

Student's t-test was used to evaluate the difference of kinase activity between non-treated and

thrombin-treated conditions. SciPy statistical package was used to perform the t-tests. At each

time point, both p-value and fold change of phosphorylation with respect to non-treated cells

were reported to determine if the two sets of data are different from each other. A p-value below

0.05 or all biological replicates reporting a fold change increasing/decreasing by at least 20%

compared to non-treated cells indicates that the kinase activity is different between non-treated

and thrombin-treated conditions.

 SUPPLEMENTAL INFORMATION

Figure S1. Workflow for building TREKING models from tKiR predictions, related to STAR

Methods. Top: temporal kinase functionality was predicted using the elastic net regularization

algorithm. Kinase inhibitor screen data (Dankwa et al., 2021) and kinase-compound biochemical

data (Anastassiadis et al., 2011) were used for tKiR. Temporal AUC was calculated within a 5-

minute sliding time window that slides at 1-minute steps for the first 2 hours of kinase inhibitor

treatment and at 5-minute steps afterwards. The direction of barrier activity of kinases (barrier-

weakening or barrier-strengthening functionality) was predicted at each time window indicated

above based on the slope of the 28 kinase inhibitors AUC values versus the residual kinase

activity from the Anastassiadis et al dataset. Middle: workflow for building TREKING networks to

describe kinase phosphosignaling in thrombin-activated barriers. Self-organizing maps (SOMs)

were used to identify kinases with similar time-resolved barrier activity across the 6-hour

timeframe following thrombin treatment. Local phosphosignaling networks for each neuron were

built by searching for the shortest paths between any pair of kinases within that neuron using

the kinase-kinase phosphorylation database on PhosphoSitePlus® as the background network.

Bottom: for a subset of kinases in the tKiR and TREKING models, their phosphorylation levels

were measured by western blot at multiple time points in the barrier disruption and barrier

recovery phases. Western blot measurements were used to evaluate if tKiR and TREKING

correctly predicted the functional associations of these kinases in thrombin-induced barrier

perturbation.

Figure S2. Build local phosphosignaling networks based on kinase functional kinetics,

related to Figure 5. (A) Self-organizing map (SOM) of kinases predicted in TNFα pre-

conditioning +thrombin condition (35 neurons). In each neuron, the black line is the mean

functionality of all the kinases within that neuron, with "+1" indicating all kinases are barrier-

strengthening and "-1" indicating all kinases are barrier-weakening. Blue and red shadings

represent barrier-weakening and barrier-strengthening functionality, respectively. (B)-(C)

Distance maps of SOM clustering in (B) +thrombin and (C) TNFα pre-conditioning +thrombin

conditions. Each cell is the normalized sum of the Euclidean distances between a neuron and

its neighboring neurons. (D) Distribution of the number of kinases assigned to the SOM

neurons. (E) Left: example of how a local phosphosignaling network was built for each neuron

by searching for the shortest paths between any pair of kinases within that cluster using the

kinase-kinase phosphorylation database on PhosphoSitePlus® as the background network.

Right: the maximum path length in the local phosphosignaling networks.

Figure S3. TNFα pre-conditioning altered the key kinase regulators and skewed the

signaling networks towards barrier-weakening activities in the barrier disruption phase.

(A) Venn diagram showing the tKiR predicted barrier-weakening kinases during the barrier

disruption phase (6-30 minutes after thrombin treatment). (B) Connectivity (in/out-degrees of

freedom) of kinases functional in the barrier disruption phase in +thrombin (top) and TNFα pre-

conditioning +thrombin (bottom) conditions. Blue and red represent connections associated with

only barrier-weakening and barrier-strengthening activities, respectively; gray represents

connections associated with both barrier-weakening and barrier-strengthening activities. (C)

Left: network connections to and from AKT1 in the barrier disruption phase in +thrombin (top)

and TNFα pre-conditioning +thrombin (bottom) conditions. Blue and red represent connections

associated with only barrier-weakening and barrier-strengthening activities, respectively; gray

represents connections associated with both barrier-weakening and barrier-strengthening

activities. Right: connectivity of AKT1 in the TREKING network reconstructed in the barrier

disruption phase.

Figure S4. Summary of western blot measurements performed in this study, related to

Figures 4 and 6. The kinases are ordered according to their network connectivity (low to high).

Left: functionality profiles predicted by TREKING. For each kinase of interest, the mean

functionality profiles of all the associated neurons in the composite TREKING network were

pooled and displayed on the same graph (orange: +thrombin condition; red: TNFα pre-

conditioning +thrombin condition). Middle: fold change of phosphorylation in +thrombin

condition; right: fold change of phosphorylation in TNFα pre-conditioning +thrombin condition.

GAPDH was used as loading control.

Figure S5. Evaluate the accuracy of TREKING predictions using western blot, related to

Figure 6. (A) For each kinase, the mean functionality profiles of all the associated neurons in

the composite TREKING network were pooled. Within a 10-minute time window centered at the

designated time point, the kinase is functionally active if TREKING predicts its activity in any of

the neurons or associated networks within the time window. In each experimental condition, the

kinase phosphorylation after thrombin treatment was compared to its phosphorylation at basal

level (media only) to determine if the kinase is activated in response to thrombin. If TREKING

predicted the kinase to be functional at a given time, and the kinase activity as measured by

western blot is different from cells not treated with thrombin (i.e., higher or lower than basal

level), the model made correct predictions on its functionality in barrier regulation; otherwise, the

model failed to make predictions on its functionality at the corresponding time point. (B)

Summary of model validations in +thrombin (left) and TNFα pre-conditioning +thrombin (right)

conditions. Pink indicates that TREKING correctly predicted the kinase functionality in barrier

regulation and grey indicates incorrect predictions. Kinase connectivity (in/out-degrees of

freedom) in the composite TREKING network is shown to the right of the panel. Light orange

and orange represent in- and out-degrees of freedom in +thrombin condition, respectively; light

red and red represent in- and out-degrees of freedom in TNFα pre-conditioning +thrombin

condition, respectively. (C)-(D) Comparison of TREKING and random models in (C) +thrombin

and (D) TNFα pre-conditioning +thrombin conditions. Top: accuracy of model predictions for

each kinase across the time course; bottom: accuracy of model predictions at each time window

for kinases evaluated in this study.

Figure S6. Composite TREKING network for the barrier disruption phase in +thrombin

condition. Kinases predicted by tKiR are labeled in blue (barrier-weakening) and red (barrier-

strengthening). Blue and red lines represent connections associated with only barrier-

weakening and barrier-strengthening activities, respectively; gray lines represent connections

associated with both barrier-weakening and barrier-strengthening activities.

Figure S7. Composite TREKING network for the barrier disruption phase in TNFα pre-

conditioning +thrombin condition. Kinases predicted by tKiR are labeled in blue (barrier-

weakening) and red (barrier-strengthening). Blue and red lines represent connections

associated with only barrier-weakening and barrier-strengthening activities, respectively; gray

lines represent connections associated with both barrier-weakening and barrier-strengthening

activities.

Figure S8. Composite TREKING network for the entire 6-hour time course in +thrombin

condition. Kinases predicted by tKiR in the barrier disruption phase are labeled in blue (barrier-

weakening) and red (barrier-strengthening); kinases predicted by tKiR in the barrier recovery

and plateau phases are labeled in light blue (barrier-weakening) and light red (barrier-

strengthening); switch kinases are labeled in gray. Blue and red lines represent connections

associated with only barrier-weakening and barrier-strengthening activities, respectively; gray

lines represent connections associated with both barrier-weakening and barrier-strengthening

activities.

Figure S9. Composite TREKING network for the 6-hour time course in response to TNFα

pre-conditioning +thrombin. Kinases predicted by tKiR in the barrier disruption phase are

labeled in blue (barrier-weakening) and red (barrier-strengthening); kinases predicted by tKiR in

the barrier recovery and plateau phases are labeled in light blue (barrier-weakening) and light

red (barrier-strengthening); switch kinases are labeled in gray. Blue and red lines represent

connections associated with only barrier-weakening and barrier-strengthening activities,

respectively; gray lines represent connections associated with both barrier-weakening and

barrier-strengthening activities.

Table S1. Predicted temporal kinase functionality in +thrombin and TNFα pre-

conditioning +thrombin conditions, related to Figure 2. This is a Microsoft Excel workbook

containing two spreadsheets.

Table S2. SOM assignment of tKiR predicted kinases and edges of phosphosignaling

networks built in +thrombin and TNFα pre-conditioning +thrombin conditions, related to

Figure 5. This is a Microsoft Excel workbook containing four spreadsheets.

Table S3. Antibody information and western blot results on the kinases validated in this

study, related to Figures 4 and 6. This is a Microsoft Excel workbook containing 14

spreadsheets.

Table S4. Literature search on protein kinases reported to regulate endothelial barrier

integrity in response to thrombin, related to Figure 7. PubMed ID of the article is followed by

the barrier functionality reported in the corresponding article ("-1": barrier-weakening; "1":

barrier-strengthening). This is a Microsoft Excel workbook containing one spreadsheet.

Supplemental file 1. Movie comparing literature reported with tKiR and TREKING

predicted kinases and signaling pathways important for barrier regulation. This is an

MPEG-4 movie file.

Supplemental file 2. Jupyter notebook containing the scripts for tKiR, SOM generation

and network construction. Also including the scripts for making frames for the supplemental

movie (Supplemental file 1) and random model generation.

REFERENCES

Aman, J., van Bezu, J., Damanafshan, A., Huveneers, S., Eringa, E.C., Vogel, S.M.,
Groeneveld, A.B., Vonk Noordegraaf, A., van Hinsbergh, V.W., and van Nieuw Amerongen,
G.P. (2012). Effective treatment of edema and endothelial barrier dysfunction with imatinib.
Circulation *126*, 2728-2738.
Anastassiadis, T., Deacon, S.W., Devarajan, K., Ma, H., and Peterson, J.R. (2011).
Comprehensive assay of kinase catalytic activity reveals features of kinase inhibitor selectivity.
Nat Biotechnol *29*, 1039-1045.
Anrather, D., Millan, M.T., Palmetshofer, A., Robson, S.C., Geczy, C., Ritchie, A.J., Bach, F.H.,
and Ewenstein, B.M. (1997). Thrombin activates nuclear factor-kappaB and potentiates
endothelial cell activation by TNF. J Immunol *159*, 5620-5628.
Arang, N., Kain, H.S., Glennon, E.K., Bello, T., Dudgeon, D.R., Walter, E.N.F., Gujral, T.S., and
Kaushansky, A. (2017). Identifying host regulators and inhibitors of liver stage malaria infection
using kinase activity profiles. Nat Commun *8*, 1232.
Beguin, E.P., van den Eshof, B.L., Hoogendijk, A.J., Nota, B., Mertens, K., Meijer, A.B., and van
den Biggelaar, M. (2019). Integrated proteomic analysis of tumor necrosis factor alpha and
interleukin 1beta-induced endothelial inflammation. J Proteomics *192*, 89-101.
Birukova, A.A., Tian, X., Tian, Y., Higginbotham, K., and Birukov, K.G. (2013). Rap-afadin axis
in control of Rho signaling and endothelial barrier recovery. Mol Biol Cell *24*, 2678-2688.
Botros, L., Pronk, M.C.A., Juschten, J., Liddle, J., Morsing, S.K.H., van Buul, J.D., Bates, R.H.,
Tuinman, P.R., van Bezu, J.S.M., Huveneers, S.*, et al.* (2020). Bosutinib prevents vascular
leakage by reducing focal adhesion turnover and reinforcing junctional integrity. J Cell Sci *133*.
Casado, P., Rodriguez-Prados, J.C., Cosulich, S.C., Guichard, S., Vanhaesebroeck, B., Joel,
S., and Cutillas, P.R. (2013). Kinase-substrate enrichment analysis provides insights into the
heterogeneity of signaling pathway activation in leukemia cells. Sci Signal *6*, rs6.
Daneman, R., and Prat, A. (2015). The blood-brain barrier. Cold Spring Harb Perspect Biol *7*,
a020412.
Dankwa, S., Dols, M.M., Wei, L., Glennon, E.K.K., Kain, H.S., Kaushansky, A., and Smith, J.D.
(2021). Exploiting polypharmacology to dissect host kinases and kinase inhibitors that modulate
endothelial barrier integrity. Cell Chem Biol.
Garcia, J.G., Liu, F., Verin, A.D., Birukova, A., Dechert, M.A., Gerthoffer, W.T., Bamberg, J.R.,
and English, D. (2001). Sphingosine 1-phosphate promotes endothelial cell barrier integrity by
Edg-dependent cytoskeletal rearrangement. J Clin Invest *108*, 689-701.
Gorovoy, M., Niu, J., Bernard, O., Profirovic, J., Minshall, R., Neamu, R., and Voyno-
Yasenetskaya, T. (2005). LIM kinase 1 coordinates microtubule stability and actin
polymerization in human endothelial cells. J Biol Chem *280*, 26533-26542.
Gujral, T.S., Peshkin, L., and Kirschner, M.W. (2014). Exploiting polypharmacology for drug
target deconvolution. Proc Natl Acad Sci U S A *111*, 5048-5053.
Guo, H., Zhou, H., Lu, J., Qu, Y., Yu, D., and Tong, Y. (2016). Vascular endothelial growth
factor: an attractive target in the treatment of hypoxic/ischemic brain injury. Neural Regen Res
11, 174-179.
Han, J., Zhang, G., Welch, E.J., Liang, Y., Fu, J., Vogel, S.M., Lowell, C.A., Du, X., Cheresh,
D.A., Malik, A.B.*, et al.* (2013). A critical role for Lyn kinase in strengthening endothelial integrity
and barrier function. Blood *122*, 4140-4149.
Hashimoto, Y., Tachibana, K., and Kondoh, M. (2020). Tight junction modulators for drug
delivery to the central nervous system. Drug Discov Today *25*, 1477-1486.
Hornbeck, P.V., Zhang, B., Murray, B., Kornhauser, J.M., Latham, V., and Skrzypek, E. (2015).
PhosphoSitePlus, 2014: mutations, PTMs and recalibrations. Nucleic Acids Res *43*, D512-520.

Klomp, J.E., Shaaya, M., Matsche, J., Rebiai, R., Aaron, J.S., Collins, K.B., Huyot, V.,
Gonzalez, A.M., Muller, W.A., Chew, T.L.*, et al.* (2019). Time-Variant SRC Kinase Activation
Determines Endothelial Permeability Response. Cell Chem Biol *26*, 1081-1094 e1086.
Knezevic, N., Tauseef, M., Thennes, T., and Mehta, D. (2009). The G protein betagamma
subunit mediates reannealing of adherens junctions to reverse endothelial permeability increase
by thrombin. J Exp Med *206*, 2761-2777.
Kohonen, T. (1982). Self-organized formation of topologically correct feature maps. Biological
cybernetics *43*, 59-69.
Komarova, Y.A., Kruse, K., Mehta, D., and Malik, A.B. (2017). Protein Interactions at Endothelial
Junctions and Signaling Mechanisms Regulating Endothelial Permeability. Circ Res *120*, 179-
206.
Kuppers, V., Vockel, M., Nottebaum, A.F., and Vestweber, D. (2014). Phosphatases and
kinases as regulators of the endothelial barrier function. Cell Tissue Res *355*, 577-586.
Liu, Y., Pelekanakis, K., and Woolkalis, M.J. (2004). Thrombin and tumor necrosis factor alpha
synergistically stimulate tissue factor expression in human endothelial cells: regulation through
c-Fos and c-Jun. J Biol Chem *279*, 36142-36147.
Luo, H., and Shusta, E.V. (2020). Blood-Brain Barrier Modulation to Improve Glioma Drug
Delivery. Pharmaceutics *12*.
Marcos-Ramiro, B., Garcia-Weber, D., and Millan, J. (2014). TNF-induced endothelial barrier
disruption: beyond actin and Rho. Thromb Haemost *112*, 1088-1102.
McVerry, B.J., and Garcia, J.G. (2004). Endothelial cell barrier regulation by sphingosine 1-
phosphate. J Cell Biochem *92*, 1075-1085.
Mehta, D., and Malik, A.B. (2006). Signaling mechanisms regulating endothelial permeability.
Physiol Rev *86*, 279-367.
Miller, L.H., Ackerman, H.C., Su, X.Z., and Wellems, T.E. (2013). Malaria biology and disease
pathogenesis: insights for new treatments. Nat Med *19*, 156-167.
Minami, T., Sugiyama, A., Wu, S.Q., Abid, R., Kodama, T., and Aird, W.C. (2004). Thrombin
and phenotypic modulation of the endothelium. Arterioscler Thromb Vasc Biol *24*, 41-53.
Morrison, D.K. (2012). MAP kinase pathways. Cold Spring Harb Perspect Biol *4*.
Oldenburg, J., and de Rooij, J. (2014). Mechanical control of the endothelial barrier. Cell Tissue
Res *355*, 545-555.
Pimienta, G., and Pascual, J. (2007). Canonical and alternative MAPK signaling. Cell Cycle *6*,
2628-2632.
Rada, C.C., Mejia-Pena, H., Grimsey, N.J., Canto Cordova, I., Olson, J., Wozniak, J.M.,
Gonzalez, D.J., Nizet, V., and Trejo, J. (2021). Heat shock protein 27 activity is linked to
endothelial barrier recovery after proinflammatory GPCR-induced disruption. Sci Signal *14*,
eabc1044.
Radeva, M.Y., and Waschke, J. (2018). Mind the gap: mechanisms regulating the endothelial
barrier. Acta Physiol (Oxf) *222*.
Rizzo, A.N., Aman, J., van Nieuw Amerongen, G.P., and Dudek, S.M. (2015). Targeting Abl
kinases to regulate vascular leak during sepsis and acute respiratory distress syndrome.
Arterioscler Thromb Vasc Biol *35*, 1071-1079.
Schafer, A., Gjerga, E., Welford, R.W., Renz, I., Lehembre, F., Groenen, P.M., Saez-Rodriguez,
J., Aebersold, R., and Gstaiger, M. (2019). Elucidating essential kinases of endothelin signalling
by logic modelling of phosphoproteomics data. Mol Syst Biol *15*, e8828.
Tiruppathi, C., Naqvi, T., Sandoval, R., Mehta, D., and Malik, A.B. (2001). Synergistic effects of
tumor necrosis factor-alpha and thrombin in increasing endothelial permeability. Am J Physiol
Lung Cell Mol Physiol *281*, L958-968.
Vaga, S., Bernardo-Faura, M., Cokelaer, T., Maiolica, A., Barnes, C.A., Gillet, L.C., Hegemann,
B., van Drogen, F., Sharifian, H., Klipp, E.*, et al.* (2014). Phosphoproteomic analyses reveal

novel cross-modulation mechanisms between two signaling pathways in yeast. Mol Syst Biol *10*,
767.
van den Biggelaar, M., Hernandez-Fernaud, J.R., van den Eshof, B.L., Neilson, L.J., Meijer,
A.B., Mertens, K., and Zanivan, S. (2014). Quantitative phosphoproteomics unveils temporal
dynamics of thrombin signaling in human endothelial cells. Blood *123*, e22-36.
Vouret-Craviari, V., Bourcier, C., Boulter, E., and van Obberghen-Schilling, E. (2002). Distinct
signals via Rho GTPases and Src drive shape changes by thrombin and sphingosine-1-
phosphate in endothelial cells. J Cell Sci *115*, 2475-2484.
Yanagida, K., Liu, C.H., Faraco, G., Galvani, S., Smith, H.K., Burg, N., Anrather, J., Sanchez,
T., Iadecola, C., and Hla, T. (2017). Size-selective opening of the blood-brain barrier by
targeting endothelial sphingosine 1-phosphate receptor 1. Proc Natl Acad Sci U S A *114*, 4531-
4536.
Zhao, Z., Nelson, A.R., Betsholtz, C., and Zlokovic, B.V. (2015). Establishment and Dysfunction
of the Blood-Brain Barrier. Cell *163*, 1064-1078.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

KiR models of temporal kinase functionality and barrier phenotypes

TREKING models of barrier phosphosignaling networks

Comparison of western blot, tKiR functionality, and TREKING barrier phosphosignaling networks

Figure S1

Figure S2

Figure S3

Figure S4

Figure S4

Figure S4

Figure S4

Figure S5

Figure S6

Figure S7

Figure S8

Figure S9