Getting Smaller by Denaturation: Acid-Induced Compaction of Antibodies

Hiroshi Imamura[1,2,3], Ayako Ooishi[1], Shinya Honda[1*]

[1]Biomedical Research Institute, National Institute of Advanced Industrial Science and Technology,

1-1-1, Higashi, Tsukuba, Ibaraki 305-8566, Japan

[2]Department of Applied Chemistry, College of Life Sciences, Ritsumeikan University, 1-1-1 Noji-

Higashi, Kusatsu, Shiga 525-8577, Japan

[3]Department of Bio-Science, Nagahama Institute of Bio-Science and Technology, 1266 Tamura,

Nagahama 526-0829, Japan

ORCID: Hiroshi Imamura, 0000-0001-8924-747X; Shinya Honda, 0000-0002-7878-3944

*Corresponding Author:

Shinya Honda

Biomedical Research Institute

National Institute of Advanced Industrial Science and Technology

1-1-1, Higashi, Tsukuba, Ibaraki 305-8566, Japan

E-mail: s.honda@aist.go.jp

https://unit.aist.go.jp/bmd/biomed-mcd/t/

Abstract

Protein denaturation is a ubiquitous process that occurs both *in vitro* and *in vivo*. While the molecular understanding of the denatured structures of proteins is limited, it is commonly accepted that the loss of unique intramolecular contacts makes proteins larger. Herein, we report compaction of the immunoglobulin G1 (IgG1) protein upon acid denaturation. Small-angle X-ray scattering coupled with size-exclusion chromatography revealed that IgG1 radii of gyration at pH 2 were ~75% of those at neutral pH. Scattering profiles showed a globular shape with approximately triaxial ellipsoids, supported by analytical ultracentrifugation. A theoretical model indicated that the acid denaturation of proteins with size reduction was energetically costly, and thus acid-induced compaction required an attractive force for domain reorientation. Such intramolecular aggregation may be widespread in multidomain proteins as non-canonical structures. Herein, we discuss the potential biological significance of these non-canonical structures of immunoglobulins.

Introduction

Proteins are polymer chains composed of amino acid residues and optimised sequences that have the distinct thermodynamic states,[1] one of which is a native state and another is a denatured state. Folded structure in the native state has been determined by extensive biophysical investigations and predicted via bioinformatic methods. In contrast, the molecular pictures for the structures in the denatured state are limited. The denatured state comprises a variety of dynamic conformers, including fully and partially unfolded structures, which has technically hindered the elucidation of their fine structures. Despite this, recent studies have explored the difference in denatured structure causing different biological and immunological consequences; for example, expanded conformation of Tau protein is favorable for the biomolecular condensation, which promotes amyloid formation.[2] Whether unfolded structures of a protein (PNt) are expanded or collapsed determines the efficiency of the secretion to the cell surface.[3] Reversal effects of therapeutic proteins are potentially associated with the morphology of the denatured aggregates.[4,5] These studies indicate significance of accumulating the knowledge about the denatured structures of proteins.

Both amino acid side chains and main chains stabilise the native protein structure by forming non-covalent bonds among them, where the constituent atoms are uniquely organised. Upon denaturation, these native contacts are replaced by non-specific water-protein contacts that increase the spatial and temporal dynamics of the proteins.[6] Thus, the size of proteins is always increased by denaturation,[7-9] which is a major principle of protein folding.

Here, we report an exceptional case where denaturation makes proteins smaller, using an immunoglobulin G1 (IgG1) monoclonal antibody. Acid-solution induces the denaturation of IgG1, including the loss of the binding function[10] and native structures.[11,12] However, the structures of

the denatured IgG1 remain unclarified. The present study used a small-angle X-ray scattering

(SAXS) technique[13] to demonstrate the size reduction of IgG1 by probing the global solution

structure of the antibody under neutral and acidic pH conditions. While the biological significance

of the reduced size of denatured IgG1 was unclear, the current models include the molecular

requirements of non-canonical contacts between domains, suggesting the universality of the

phenomenon of denaturation reducing the size of multidomain proteins.

Figure 1. The mAb-1 pH-dependence of the small-angle X-ray scattering (SAXS) at 25 °C. The

data are vertically shifted for clarity. Black line shows the theoretical SAXS profile of the IgG1

model (PDB: 1HZH).

Results

Figure 1 depicts the acid-induced changes in an IgG1 monoclonal antibody, mAb-1, when

measured by SAXS. The SAXS profiles were similar at pH 7.1–3.1, with the characteristic

shoulders (Figure 1, arrows). These scattering patterns were consistent with the theoretical

scattering calculated from the crystal structure of IgG1 (black line in Figure 1; denoted as "Model")

and previous SAXS data reported in the literature.[14,15] Conversely, the scatterings at pH < 2.0

were distinct from those at pH 7.1–3.1; the characteristic shoulders in the log-log plots (Figure 1)

and peaks in the Kratky plots (Figure S1b) diminished at pH < 2.0, with more scattering in the

low-q range at pH 1.1 (Figure S1a).

The SAXS data indicated a drastic conformational alteration of mAb-1 around pH 2, but

the contaminating aggregates hampered further structural analysis (see Supporting Information

and Figure S1). Therefore, we performed SEC-SAXS analysis[16-18] *in situ* SAXS during size

exclusion chromatography (SEC), which allowed us to monitor the SAXS of monomeric mAb-1

at pH 2. Chromatographic analysis was conducted of the integral SAXS intensities and ultraviolet

(UV) absorbance at 280 nm using an elution buffer solution of 0.1 M glycine-HCl (pH 2) with 0

M or 0.2 M NaCl (Figure 2a). The peaks at ~169 min (0 M NaCl) and ~66 min (0.2 M NaCl)

represented monomeric mAb-1 that were well separated from the aggregates, while the peaks at

~124 min (0 M NaCl) and ~56 min (0.2 M NaCl) indicated dimeric mAb-1. The aggregates (>

trimer) manifested as peaks that presented early in the analysis. These assignments are supported

by the chromatogram of the integrated SAXS intensities, which have more pronounced aggregates

than the UV absorbance. The variation in the retention time between the data (0 M and 0.2 M

NaCl) was due to the difference in the flow rates used.

Based on the chromatograms, we determined the concentration-normalized SAXS profiles

($I(q)/c$) of monomeric mAb-1 at pH 2 (Figure 2b). Because the concentrations were < 0.5 mg/mL

in the sample cell during the chromatographic analysis, the intermolecular interference was

negligible. The radius of gyration (R_g) was 50.8 ± 0.3 Å for 0 M NaCl and 40.3 ± 0.1 Å for 0.2 M

NaCl and the concentration-normalized zero-angle scattering ($I(0)/c$) was 0.0894 ± 0.0003 cm^2

mg^{-1} (122 kg/mol) for 0 M NaCl and 0.0783 ± 0.0003 cm^2 mg^{-1} (108 kg/mol) for 0.2 M NaCl,

which were comparable with the native monomeric mAb-1 of 0.0880 ± 0.0007 cm^2 mg^{-1} (121

kg/mol). These results indicated that the SEC-SAXS was able to capture the monomeric acid-

denatured mAb-1 by removing the aggregates. The deviations from the exact molar mass (M) of

mAb-1 (148 kg/mol) were due to a systematic error. In the log-log and Kratky plots of the SAXS

profiles (Figure 2b and 2c), the characteristic shoulders/peaks observed for the native mAb-1 at

pH 7.1 (Figure 1 and Figure S1b) was not observed at pH 2. The distance distribution function

($P(r)$) at pH 2 indicated a lack of the characteristic native interdomain distance distribution (~80

Å) observed at pH 7.1 (Figure 2d), suggesting the distances and orientations of the Fab(antigen-

binding fragment)-Fc(crystallizable fragment) and Fab-Fab regions were not canonical at pH 2.

Furthermore, the maximum distance (D_{max}) at pH 2 were < ~200 Å (0 M NaCl) and < ~150 Å (0.2

M NaCl). The former was comparable with D_{max} at pH 7.1.

A fitting analysis of the SAXS profiles characterised the shape of mAb-1 at pH 2 (Figure

2b). Among the theoretical scattering functions of the sphere, cylinder, elliptical cylinder, ellipsoid,

and triaxial ellipsoid shapes,[19,20] the scattering function of the triaxial ellipsoid ($a \neq b$) gave the

best fit, with lengths, a, b, and c defining the shape of the triaxial ellipsoid (Figure 2b). The

scattering function of the ellipsoid ($a = b$) produced deviations from the experimental scatterings

in the region of $q > 0.1$ $Å^{-1}$ (Figure 2b and 2c); such a deviation was also found in $P(r)$ functions

(Figure 2d). These deviations for $I(q)/c$ and $P(r)$ at 0 M NaCl were larger than those at 0.2 M NaCl,

indicating that the shape of mAb-1 at 0 M NaCl was more flattened than that at 0.2 M NaCl.

Figure 2e shows the triaxial ellipsoids with the best parameters for this fitting. The

representation of mAb-1 at pH 2 with 0 M NaCl (grey) and 0.2 M NaCl (blue) was also flattened

compararable with that of the native IgG1 (red). We observed that 0.2 M NaCl made mAb-1 more

compact with a smaller maximum length and globular shape than 0 M NaCl. Figure 2f shows the reconstructed 3D-models of mAb-1 at pH 2 from the SAXS profiles and $P(r)$ by the *ab initio* electron density determination, DENSS (DENsity from Solution Scattering) algorithm.[21] The flattened shapes of mAb-1 at pH 2 were also consistent with those obtained via the fitting analysis that used the theoretical scattering functions. In addition, these 3D models illustrated some anisotropic electron distributions, such as a lower density region that looked like a "neck" in 0 M NaCl. Another remarkable characteristic of mAb-1 at pH 2 included a smaller R_g than that of the native mAb-1 (Figure 2g). The addition of NaCl led to a smaller R_g at pH 2, while R_g was not dependent on NaCl at pH 7. mAb-1 at pH 2 was very plastic and presented as globular due to the shielding effect of the salt, which reduced the intramolecular electrostatic repulsion of the acid-denatured mAb-1. We noted that the incubation time at pH 2 did not significantly change the R_g.

We further conducted the SEC-SAXS analysis on another IgG1, mAb-2, to test whether a variation of the sequence in the variable region alter the conclusion of the size-reduction. We verified an acid-induced compaction of mAb-2, which showed $R_g = 53.0 \pm 0.4$ Å at pH 7.2 and $R_g = 46.5 \pm 0.3$ Å at pH 2 and 0.2 M NaCl.

Figure 2. Size exclusion chromatography small-angle X-ray scattering (SEC-SAXS) analysis on

mAb-1 at pH 2. The results of mAb-1 at 0 M NaCl and 0.2 M NaCl are represented as grey and

cyan, respectively. (a) Overlaid elution SAXS and UV profiles where the SAXS-derived

chromatogram is an integrated SAXS intensity vs. retention time. The elution UV profiles were

acquired by simultaneous recording of mAb-1 absorbance at 280 nm. (b) SAXS profiles of

monomeric mAb-1. The scattering for mAb-1 in 0.2 M NaCl was multiplied by 10^2 for clarity.

Theoretical scattering of the triaxial ellipsoid (red solid line) and ellipsoid (dashed black line) with

best fit parameters for mAb-1 are shown. An elliptical object indicates where *a*, *b*, and *c* are the

length parameters. (c) Kratky plots of IgG1 SAXS profiles and theoretical scattering. (d) The

distance distribution function, $P(r)$, by the direct Fourier transformation of the SAXS profiles. $P(r)$

of the triaxial ellipsoid (red solid line) and ellipsoid (black dashed line) were calculated from the

scatterings in (b). $P(r)$ of the native mAb-1 at pH 7.1 (Figure S1e) is overlaid for comparison. This

$P(r)$ was scaled such that the area is equivalent to the $P(r)$ of the mAb-1 at pH 2 and 0 M NaCl.

The data from mAb-1 in 0.2 M NaCl presented in (c) and (d) are vertically shifted for clarity. (e)

A point cloud representation of the triaxial ellipsoids using best fit parameters for the scatterings

of mAb-1 at 0 M NaCl and 0.2 M NaCl. A canonical IgG1 structure,[22] where the C_α atoms (red

point clouds) are placed to fit with the triaxial ellipsoids with characteristic lengths. (f) 3D

reconstructed models of the mAb-1 at pH 2 using *ab initio* electron density determination

(DENSS).[21] The canonical mAb-1 models are also shown, indicating the *ab initio* model

reconstructed from the SAXS data (light pink), overlaid with the atomic coordinates of the IgG1

(red ribbons).[22] All the volumes inside the model surfaces are set at 258,000 $Å^3$, where the Porod

volume was determined for the scattering of mAb-1 at 0 M NaCl. (g) Dependence of mAb-1 R_g at

pH 2 on incubation time and NaCl concentration. The R_g of the native mAb-1 at pH 7 (with 0 M

NaCl and 0.1 M NaCl) is indicated for comparison.

An orthogonal analysis of the acid-denatured mAb-1 using an analytical ultracentrifugation (AUC)

strengthened the SAXS analysis. The sedimentation velocity data (Figure S2a) showed the

distribution of the sedimentation coefficients (s) of the mAb-1 samples at pH 7 and pH 2 (Figure

S2b), in which prominent peaks at $s = \sim6.5$–7.0 S were consistent with the expected s value for the

known molar mass of the monomeric mAb-1 (148 kg/mol; Table S1). We found a small but

significant variation in $s_{20,\text{w}}$ (s at 20 °C in water) among the samples, where the effects of the

viscosities and densities of the solvents were corrected. This variation indicates the differences in

the molecular envelope, represented as the friction factor in the AUC. The variation in $s_{20,\text{w}}$ due to

the association of the molecules was ruled out; for example, the dimer of mAb-1 had a $s = \sim9.1$ S

at pH 2 and 0 M NaCl (Figure S2b). According to equations (S8–S11), the friction factor (f/f_0) and

the corrected friction factor ($f/f_{0,\text{sp}}$) were determined. The f/f_0 of the native mAb-1 (1.43) was

comparable with a previously reported value (1.55);[23] the deviation should be due to the

difference in the specific volume (v) used. Empirically, approximately globular proteins,

moderately elongated proteins, and highly elongated proteins were reported to display f/f_0 of 1.2–

1.3, 1.5–1.9 and 2.0–3.0, respectively.[24] The current f/f_0 indicated that the native mAb-1 and the

acid-denatured mAb-1 were both globular rather than elongated. The f/f_0 at pH 2 decreased from

1.58 to 1.35 with the addition of 0.2 M NaCl, consistent with the SAXS result that mAb-1 at pH 2

was more globular in 0.2 M NaCl. The corrected friction factor ($f/f_{0,\text{sp}}$) in equations (S9–S11)

provided information regarding the protein shape and can be calculated for any triaxial

ellipsoid.[25,26] We calculated $f/f_{0,sp}$ using the triaxial ellipsoid lengths a, b, and c, as determined

using the SAXS analysis (Figure 2). The experimentally determined $f/f_{0,sp}$ at pH 2 decreased by

0.19 (1.32 at 0 M NaCl and 1.13 at 0.2 M NaCl), which was comparable with the decrease in the

calculated $f/f_{0,sp}$ by 0.14 (Table S1). The differences between the absolute values of the

experimental $f/f_{0,sp}$ and the calculated value were due to uncertainties in the β_1, v, and η used in

equations (S7 and S8). Together, the results of SAXS—the non-elongated shape of the acid-

denatured mAb-1 and its plastic globularisation by the addition of NaCl— are well supported by

the AUC.

Discussion

Protein denaturation increase their size independent of perturbations like temperature,

pressure, or chemical denaturation.[8] Indeed, the R_g for proteins reported in the literature indicated

that the underlying power laws of R_g vs. chain lengths (L) are distinct between native and unfolded

protein structures (Figure 3). The R_g of the native state of the multidomain protein mAb-1 followed

the power-law relationship for the native structures, although the representation of R_g in the

literature[9] is dominated by single-domain proteins. In contrast, the R_g of non-native mAb-1 was

not predicted by the power law relationship and anomaly smaller than that of the native state. The

R_g of the non-native mAb-1 dimer, apart from the scaling law for unfolded proteins, also indicates

a non-random coil structure.

Figure 3. R_g dependence on protein chain length. The data in literature includes native (green,

closed circles), native oligomeric (dark orange, closed circles), and unfolded (black, closed

triangles) proteins, curated by Po-Min Shih et al.[9] The dotted lines represent the power law

scaling of $R_g = R_0 L^{\xi}$, where R_0 is a prefactor (2.04, native; 4.47, native oligomer; and 2.05,

unfolded), L is the amino acid (a.a.) chain length, and ξ is the scaling exponent (0.45, native; 0.32,

native oligomer; and 0.58, unfolded). R_g of mAb-1 at pH 7 (red, open circle), R_g of mAb-1 at pH

2 with 0.2 M NaCl (red, open triangle), and R_g of mAb-1 dimer at pH 2 with 0.2 M NaCl (red,

open diamond) determined in this study are overlaid onto previously reported results.[9] The R_g

value of mAb-1 at pH 7 is given as an infinite dilution of the protein (Figure S1h).

11   Protein acid denaturation is caused by the repulsion of positive charges on the surface of the

12 protein that are described by simple models such as the Linderstrøm-Lang smeared charge

13 model.[27,28] This model indicates that the size expansion decreases the Gibbs energy by

14 denaturation (see Supporting Information) and describes how acid denaturation that reduces

15 protein size is energetically costly and anomalous. No previous reports have shown acid-

16 denaturation without size expansion; even the molten-globule state of cytochrome *c* at pH 2 and

0.1 M NaCl (17.4 Å), which is described as a "compact" denatured state, is larger than its native

state (13.5 Å).[29]

The charge distributions shown in Figure 4a identified a positive potential on the surface of

mAb-1 that widely emerged at pH 2. We hypothesise alternative, energetically favourable

interactions that hold the parts of the acid-denatured IgG1 molecule together by compensating for

the energetic cost of maintaining the protein's size. Additional van der Waals interactions would

assist in this process. The acid-denatured state at pH 2 was more compact with the addition of

NaCl, indicating that shielding the coulomb repulsion among the charged residues and

unfavourable salt-induced solvation of the exposed hydrophobic protein surface assisted the sub-

complete packing of the protein.

Figure 4. Acid effects on IgG structure. (a) Charge distribution in IgG1 at various pH. The

electrostatic properties were calculated using the pK_a predictor PROPKA[30,31] and the adaptive

Poisson–Boltzmann solver (APBS) software[32] using the atomic coordinates for IgG1[22] and the

mAb-1 sequence. (b) The conformational states of the IgG domains at pH 2.0–2.6 (with/without

salts) in the context of the isolated domain, Fc/Fab regions, the full-length IgG reported in the

literature; the references are presented in the table are [a]pH 2.0;[33] [b]aglycosylated C_H2, pH 2;[34]

[c]pH 2;[35] [d]aglycosylated Fc, pH 2.6;[36] [e]aglycosylated Fc, pH 2.5;[37] [f]glycosylated Fc, pH 2.6;[36]

[g]pH 2;[38] [h]pH 2.0;[39] [i]pH 2;[40] [j]pH 2.5;[41] [k]pH 2.1;[42] [l]glycosylated Fc, pH 2.0;[43] [m]pH 2.0;[11] and

[n]pH 2.0.[12] The canonical IgG1 structure,[22] its domains (C_H3, C_H2, C_H1, C_L, V_H, and V_L), and the

regions (Fc and Fab) are presented.

The gain of native contacts in folded proteins contributes to their size reduction.[7]

Therefore, the acid-denatured state for mAb-1 could be regarded as an alternatively folded state

(AFS) that had alternative contacts that prevented expansion. Buchner et $al.$[44] referred to AFS

hitherto as "The antibody conformation at low pH had been termed 'alternatively folded state'

because it exhibits characteristics of the folded state (e.g. remarkable stability against unfolding),

but the available spectroscopic information suggests that it is structurally significantly different

from the native state. It was first described for a complete IgG1 antibody,[11] but single domains

such as C_H3 also can adopt this state.[35]" The original paper published in 1991[11] characterised

MAK33, a murine κ/IgG1 subtype, at pH 2. The $s_{20,w}$ of 5.53 for the AFS (pH 2) of MAK33 was

smaller than $s_{20,w}$ of 6.60 for the native protein, and much larger than $s_{20,w}$ of 3.2 for the denatured

protein in 6 M guanidine hydrochloride at pH 2. These AUC results indicated that the molecules were neither significantly aggregated nor fully extended. Nevertheless, the small amount of the aggregates functioned as impurities hindered the discovery that the AFS state of IgG1 can be smaller than the native state. As the SEC-SAXS and the sedimentation coefficient distribution analysis for the AUC have not been established previously, the technical limitations also hindered the discovery. The circular dichroism (CD) spectra of the AFS state of MAK33 at pH 2 were similar to the current CD spectra (Figure S3a), further supporting the AFS of the present acid-denatured mAb-1 in which the β-sheets are dominant (Table S2) and slightly reorganised, as indicated using infrared (IR) analysis (Figure S3c). Subsequent studies on AFS focused on the isolated protein domains such as C_H3 or V_L domains, rather than full-length antibodies. Buchner *et al.*[45] reported that the MAK33 V_L domain could form stable high-molecular weight oligomers at pH 2 that had previously been reported as AFS. Although the term AFS is loosely defined, it can be concluded that the association between protein domains is a common mechanism for AFS irrespective of whether full-length antibodies or isolated domains are evaluated.

Studies have also reported that the other IgG molecules (murine monoclonal IgG2a antibody CB4-1,[42] monoclonal chimeric antibody IgG1 BR96[12]) can adopt the AFS state at pH 2. In addition, Sedlák et al[46] proposed an inter-domain interaction of heat-denatured IgG, referred to as "intramolecular aggregation". The authors compared differential scanning calorimetry (DSC) thermograms of full-length IgG and its Fab portion. The difference in melting temperatures of the Fab domain suggests a Fab-Fab interaction in the heat-denatured state of the protein, which was supported by the lack of protein concentration-dependence in the thermograms.[46] The concept of "intramolecular aggregation" would be universal for IgG irrespective of the denaturing perturbations (e.g., acid or heating).

We also investigated the domains that participated in interdomain interactions in acid-

denatured IgG1. The canonical IgG1 protein consists of 12 domains in the Fc or Fab regions

(Figure 3b). In literature,[11,12,33-42,47] the stabilities of the domains depend on the molecular

contexts (Figure 3b), e.g., isolated domains or domains in a full-length IgG. Isolated C_H3[34] at pH

2, and C_H3 in an aglycosylated Fc[37] at pH 2.5 unfold, hence the native dimerised C_H3-C_H3

interface are lost.[34,37] However, the ordered structure of C_H3 in the full-length IgG is possibly

retained[38,39] with local disorder that decreases the left-twisting of β-strands (Figure S3 and Table

S2). The C_H2 unfolds at pH 2 regardless of the presence of glycans in any molecular context.[34,36-

39] The isolated domains (C_H1, C_L, and V_L) of the Fab region unfold at pH 2,[33] whereas the Fab

region is structured when the Fab region protein[40-42] and full-length IgG[38,39,42] protein are

examined. Therefore, the possible mechanism of acid-induced intramolecular aggregation of IgG1

is as follows: at pH 2, C_H3 avoids complete unfolding by interacting with the Fab region, where

the flexibility of the unfolded C_H2 does not hamper the interdomain contacts. Salt weakens the

Fab-Fab repulsion, which brings the two Fab regions close to each other and IgG1 smaller. Self-

interactions between C_H3 are stabilised by salt, indicated by the AFS of the isolated C_H3

oligomer.[35] We speculate that the hydrophobic V_H-V_L and C_H1-C_L interfaces[48] would be exposed

upon breaking the native interfacial contacts and the inter-domain salt bridges.[49] All those effects

could make non-canonical contacts between the Fab and Fc regions.

It is intriguing whether this non-canonical structure of the acid-denatured IgG1 functions

similar to an α-lactalbumin in the molten globule state (human α-lactalbumin, HAMLET), which

is lethal to tumour cells.[50] A few reports have shown the biological significance of alternatively

folded state of immunoglobulins (Ig); for example, the oral administration of Ig (IgG and IgA) as

a passive immunity treatment. Orally administered Ig is exposed to the acidic pH of the

gastrointestinal (GI) tract, but is resistant to complete digestion.[51] According to the literature,[11] Jasion and Burnett suggested that transient exposure of IgG to the acidic conditions in the stomach may increase its stability as it passed into the GI tract and against proteolytic enzymes.[51] The non-canonical structure of acid-denatured IgG1 presented in this study supports the idea that reducing the flexibility of locally unfolded regions through intramolecular aggregation is beneficial for the protein's resistance to enzymatic digestion. The ions present in physiological conditions could also support the tight packing of the acid-denatured IgG1 (Figure 2). Another example is regarding the expression and secretion of natural or recombinant Ig proteins. Onitsuka *et al.*[52] found antibody-producing Chinese hamster ovary cells secreted aggregates of recombinant IgG from the intracellular environment into the extracellular milieu, further evidenced by a new live-cell imaging technique[53] that uses a fluorescently labelled AF.2A1 as a protein probe for non-native IgG structures.[47,54] Similar intracellular particles, known as Russell bodies, are found in plasma cells. Russell bodies are natural inclusion bodies usually found in cells that overexpress Ig. Mammalian cells generally secrete only correctly folded proteins; therefore, a naturally occurring AFS could pass through an intracellular quality control system, such as endoplasmic reticulum (ER)-associated degradation.[52,53] If the naturally occurring AFS was similar to the acid-denatured IgG1, the compact form may be favourable for Ig intracellular transport, secretion, internalisation, and transcytosis in various circumstances.

The acid-denatured state of IgG1 is relevant to downstream processes of biopharmaceutical manufacturing where antibodies experience temporal exposure to acid during affinity chromatographic purification and virus inactivation, which is followed by neutralisation. The non-native conformation of IgG1 induced by this pH shift is aggregation-prone at neutral pH, with structural origins that remain unclear.[10,55-60] Because the acid-denatured state is unstable at neutral

pH, the IgG1 molecules transition toward native or aggregated states. It is unclear whether the molecular characteristics of the acid-denatured IgG1, such as intramolecular aggregation, slow or accelerate these reactions. In addition, the precise prediction of antibody stability in an acidic environment requires the characterisation of the acid-denatured structure, which is important aspect of the engineering process of antibodies.[61]

Conclusion

In conclusion, the current knowledge-based principle of proteins getting larger by denaturation that has been established by many studies using single-domain proteins should be reconsidered. New view should now include the potential for proteins to get smaller by denaturation, as we demonstrated for acid-denatured IgG1 proteins (mAb-1 and mAb-2). The reported cases are limited; nevertheless, as they shared the characteristics of the previously reported AFS for IgGs, an acid-induced compaction can be common among various IgG proteins. Multidomain proteins possess additional higher-order structures, such as domain-domain spatial arrangements or orientations, which are lost upon denaturation. These degrees of freedom in multidomain proteins are prerequisites for a more compact structure by the rearrangement of the domains, in which there are minor secondary/tertiary structural changes. In contrast, single domain proteins are almost completely packed, with no degrees of freedom for further compaction. The compact denatured structure induced by intramolecular aggregation may be ubiquitous in multidomain proteins, including other classes of Ig, as a non-canonical structures.

Experimental Section

Two monoclonal antibodies (mAb) were used; one is mAb-1, a humanised IgG1

monoclonal antibody (148 kDa; calculated pI[62] = 8.5), and another is mAb-2, a human IgG1

monoclonal antibody (147 kDa; calculated pI[62] = 8.4). IgG1 was dissolved in a buffer solution

(0.01 M sodium phosphate, 0.15 M NaCl and 0.005% (v/v) polyoxyethylene (20) sorbitan

monolaurate, pH 7.4) to be ~1–100 mg/mL (stock solution of IgG1). The IgG1 stock solution was

diluted to given concentrations (based on the experiment) with 0.01 M sodium phosphate buffer

solution (pH 7.4) and dialyzed against 0.01 M sodium phosphate buffer solutions at pH 3, 4, 6,

and 7, or 0.1 M glycine-HCl buffer solutions at pH 1, 2, and 3 at 4 °C for 7 h using a Toru-kun

TOR-14K microdialyzer (NIPPON Genetics Co., Ltd., Japan) with a membrane with a cut-off of

14 kDa. During dialysis and incubation, the protein aggregated at pH 1–2 (Figure S1). These

samples were stored at 4 °C for ~14 h until the batch SAXS measurement (Figure 1). Both the

sodium phosphate and glycine-HCl buffer solutions at pH 3 produced comparable SAXS profiles

(the latter is presented in Figure 1). For the SEC-SAXS measurement (Figure 2), the ~10 or ~20

mg/mL IgG1 stock solutions were dialyzed against 0.1 M glycine-HCl buffer solution (pH 2.0)

with a membrane cut-off of 12−14 kDa (Scienova GmbH, Germany) and stored at 4 °C for 1 h, 1

d, 3 d, or 7–8 d before measurement.

SAXS The SAXS experiments were performed with a BL-10C beam line at the Photon Factory

(PF) of the High Energy Accelerator Research Organization (KEK) in Tsukuba, Japan.[63] The X-

ray wavelengths (λ) used were 0.10, 0.12, or 0.15 nm. The camera length was 1 or 2 m, and was

calibrated using the scattering pattern of silver behenate.[64] The scattering parameter q is defined

as $q = |\boldsymbol{q}| = 4\pi\sin\theta/\lambda$, where $\boldsymbol{q}$ is the scattering vector and 2θ is the scattering angle of the X-rays.

X-ray intensities were recorded with a PILATUS3 2M detector (DECTRIS Ltd., Switzerland). For

the batch SAXS measurement (Figure 1), the samples were measured at 25 ± 0.1ºC in a cell with

quartz windows; 30 images were collected, where one image was recorded during a 2-s exposure

time; the data from samples damaged by radiation were eliminated.

For the SEC-SAXS measurement (Figure 1), the IgG1 samples (~7–11 mg/mL, 100–150

μL) were passed through a stainless-steel sample cell via an in-line size exclusion column

(Superdex200^{TM} (10/300) GL; GE Healthcare UK Ltd., England; Lot. 10111292 and 10216439)

operated by an HPLC system (Acquity UPLC H-class and Empower3 FR2; Waters Co., Milford,

MA or Nexera-i; Shimadzu Co., Kyoto, Japan) installed with a beam line BL-10C[63,65] and using

flow rates of 0.2 mL min^{-1} for the initial 30 min and then 0.05 mL min^{-1} (Figure 2a, lower data)

or 0.2 mL min^{-1} (Figure 2a, upper data). The temperature of the sample cell and the column was

maintained at 25.0 °C. The exposure time for recording the image was 20 s. The sample solution

was eluted into the sample cell with 0.02 mm-thick quartz glass windows and 1 mm-path length,

where UV–vis absorption spectra (~2 mm-path length by the diagonal incident light on the cell)

were simultaneously recorded using a fibre-coupled UV–vis spectrometer, QEpro, with Opwave+

version 2.14–2.16 (Ocean Optics, Inc., Largo, FL). A fitting analysis using theoretical scattering

functions was conducted using the codes provided by the NIST Center for Neutron Research

(NCNR).[66] Unless otherwise indicated, all analyses were performed using IGOR Pro version

6.22A (WaveMetrics, Portland, OR, USA). Analyses of the scattering data are detailed in

Supporting Information.

AUC mAb-1 samples at 0.7 mg/mL were used. Sedimentation velocity measurements were

performed in a ProteomeLab XL-A (Beckman Coulter, Inc., Brea, CA) with rotor speeds of 42

krpm at 25 ± 0.2 ºC. An absorbance of 280 nm was used to monitor the protein concentration in a

double-sector cell. The sedimentation velocity data were processed with the SEDFIT program

using a sedimentation coefficient distribution analysis.[67] The parameters (s, f/f_0 and $f/f_{0,sp}$) were

determined according the equations in Supporting Information.

CD and IR mAb-1 samples at 1 mg/mL were measured. The procedures are detailed in

Supporting Information.

ACKNOWLEDGMENTS

This research is partially supported by the Japan Society for the Promotion of Science (grant

numbers JP19H03363 [to S.H.] and JP21K06503 [to H.I.]). The SAXS experiment in this work

was performed under the approval of the Photon Factory Program Advisory Committee (Proposal

No. 2015G061 and No. 2017G068). Computations were partially performed on the NIG

supercomputer at ROIS National Institute of Genetics. The authors thank Dr. Tomoshi Kameda

(AIST, Tokyo, Japan) and Dr. Seiki Yageta (AIST, Tsukuba, Japan) for stimulating discussion and

comments.

Author Contributions:

H.I. and S.H. designed research. H.I. and A.O. performed the experiments. H.I. analysed the data.

H.I and S.H. wrote the main manuscript text. All authors contributed to data interpretation,

critically reviewed the manuscript, and given approval to the final version of the manuscript.

Conflict of Interest:

The authors declare no competing financial interest.

REFERENCES

[1] A. M. Lesk, *Introduction to Protein Science: Architecture, Function, and Genomics*, Oxford University Press, Oxford, **2016**.

[2] J. Wen, L. Hong, G. Krainer, Q.-Q. Yao, T. P. J. Knowles, S. Wu, S. Perrett, *J. Am. Chem. Soc.* **2021**, *143*, 13056–13064.

[3] M. A. Bowman, J. A. Riback, A. Rodriguez, H. Guo, J. Li, T. R. Sosnick, P. L. Clark, *Proc. Natl. Acad. Sci. U. S. A.* **2020**, *117*, 23356–23364.

[4] M. K. Joubert, M. Hokom, C. Eakin, L. Zhou, M. Deshpande, M. P. Baker, T. J. Goletz, B. A. Kerwin, N. Chirmule, L. O. Narhi, et al., *J. Biol. Chem.* **2012**, *287*, 25266–25279.

[5] *Guidance for Industry: Immunogenicity Assessment for Therapeutic Protein Products*, Food and Drug Administration, Center for Drug Evaluation and Research (CDER), Center for Biologics Evaluation and Research (CBER), **2014**.

[6] P. L. Clark, K. W. Plaxco, T. R. Sosnick, *J. Mol. Biol.* **2020**, *432*, 2882–2889.

[7] V. N. Uversky, A. L. Fink, *FEBS Lett.* **2002**, *515*, 79–83.

[8] I. S. Millett, S. Doniach, K. W. Plaxco, *Adv. Protein. Chem.* **2002**, *62*, 241–262.

[9] P.-M. Shih, I. Wang, Y.-T. C. Lee, S.-J. Hsieh, S.-Y. Chen, L.-W. Wang, C.-T. Huang, C.-T. Chien, C.-Y. Chang, S.-T. D. Hsu, *J. Phys. Chem. B* **2015**, *119*, 5437–5443.

[10] L. Zhou, J. Zhang, E. DiGiammarino, A. Kavishwar, B. Yan, C. Chumsae, P. M. Ihnat, D. Powers, J. Harlan, W. B. Stine, *Anal. Chem.* **2018**, *90*, 12221–12229.

[11] J. Buchner, M. Renner, H. Lilie, H. J. Hinz, R. Jaenicke, T. Kiefhaber, R. Rudolph, *Biochemistry* **1991**, *30*, 6922–6929.

[12] M. Kats, P. C. Richberg, D. E. Hughes, *Anal. Chem.* **1997**, *69*, 338–343.

[13] D. I. Svergun, M. H. J. Koch, P. A. Timmins, R. P. May, *Small Angle X-Ray and Neutron Scattering From Solutions of Biological Macromolecules*, Oxford University Press, Oxford, **2013**.

[14] X. Tian, A. E. Langkilde, M. Thorolfsson, H. B. Rasmussen, B. Vestergaard, *J. Pharm. Sci.* **2014**, *103*, 1701–1710.

[15] X. Tian, B. Vestergaard, M. Thorolfsson, Z. Yang, H. B. Rasmussen, A. E. Langkilde, *IUCrJ* **2015**, *2*, 9–18.

[16] E. Mathew, A. Mirza, N. Menhart, *J. Synchrotron Rad.* **2004**, *11*, 314–318.

[17] E. Brookes, P. Vachette, M. Rocco, J. Pérez, *J. Appl. Crystallogr.* **2016**, *49*, 1827–1841.

[18] M. A. Graewert, S. Da Vela, T. W. Gräwert, D. S. Molodenskiy, C. E. Blanchet, D. I. Svergun, C. M. Jeffries, *Crystals* **2020**, *10*, 975.

[19] A. Guinier, G. Fournet, *Small-Angle Scattering of X-Rays*, John Wiley & Sons, Inc., New York, **1955**.

[20] L. A. Feigin, D. I. Svergun, *Structure Analysis by Small-Angle X-Ray and Neutron Scattering*, Plenum Press, New York, **1987**.

[21] T. D. Grant, *Nat. Methods* **2018**, *15*, 191–193.

[22] J. P. Brandt, T. W. Patapoff, S. R. Aragon, *Biophys. J.* **2010**, *99*, 905–913.

[23] R. Paul, A. Graff-Meyer, H. Stahlberg, M. E. Lauer, A. C. Rufer, H. Beck, A. Briguet, V. Schnaible, T. Buckel, S. Boeckle, *Pharm. Res.* **2012**, *29*, 2047–2059.

[24] H. P. Erickson, *Biol Proced Online* **2009**, *11*, 32–51.

[25] S. E. Harding, J. C. Horton, H. Cölfen, *Eur. Biophys. J.* **1997**, *25*, 347–359.

[26] J. G. de L. Torre, S. E. Harding, *Biophys. Rev.* **2013**, *5*, 195–206.

[27] K. Linderstrøm-Lang, *C. R. Trav. Lab. Carlsberg* **1924**, *15*, 70–95.

[28] M. S. Kay, R. L. Baldwin, *Biochemistry* **1998**, *37*, 7859–7868.

[29] M. Kataoka, Y. Hagihara, Y. Goto, *J. Mol. Biol.* **1993**, *229*, 591–596.

[30] C. R. Søndergaard, M. H. M. Olsson, M. Rostkowski, J. H. Jensen, *J. Chem. Theory Comput.* **2011**, *7*, 2284–2295.

[31] M. H. M. Olsson, C. R. Søndergaard, M. Rostkowski, J. H. Jensen, *J. Chem. Theory Comput.* **2011**, *7*, 525–537.

[32] E. Jurrus, D. Engel, K. Star, K. Monson, J. Brandi, L. E. Felberg, D. H. Brookes, L. Wilson, J. Chen, K. Liles, et al., *Protein Sci.* **2018**, *27*, 112–128.

[33] M. J. Feige, E. R. Simpson, E. M. Herold, A. Bepperling, K. Heger, J. Buchner, *J. Mol. Biol.* **2010**, *399*, 719–730.

[34] S. Yageta, T. M. Lauer, B. L. Trout, S. Honda, *Mol. Pharm.* **2015**, *12*, 1443–1455.

[35] M. J. W. Thies, R. Kammermeier, K. Richter, J. Buchner, *J. Mol. Biol.* **2001**, *309*, 1077–1085.

[36] S. Yageta, R. Shibuya, H. Imamura, S. Honda, *Mol. Pharm.* **2017**, *14*, 699–711.

[37] R. F. Latypov, S. Hogan, H. Lau, H. Gadgil, D. Liu, *J. Biol. Chem.* **2012**, *287*, 1381–1396.

[38] S. P. Martsev, Z. I. Kravchuk, A. P. Vlasov, G. V. Lyakhnovich, *FEBS Lett.* **1995**, *361*, 173–175.

[39] A. P. Vlasov, Z. I. Kravchuk, S. P. Martsev, *Biokhimiia* **1996**, *61*, 212–235.

[40] H. Lilie, J. Buchner, *FEBS Lett.* **1995**, *362*, 43–46.

[41] V. M. Tischenko, V. P. Zav'yalov, G. A. Medgyesi, S. A. Potekhin, P. L. Privalov, *Eur. J. Biochem.* **1982**, *126*, 517–521.

[42] K. Welfle, R. Misselwitz, G. Hausdorf, W. Höhne, H. Welfle, *Biochim. Biophys. Acta* **1999**, *1431*, 120–131.

[43] D. Kanmert, Structure and Interactions of Human IgG-Fc, Linköpings universitet, **2011**.

[44] M. J. Feige, L. M. Hendershot, J. Buchner, *Trends Biochem. Sci.* **2010**, *35*, 189–198.

[45] M. Hora, R. Sarkar, V. Morris, K. Xue, E. Prade, E. Harding, J. Buchner, B. Reif, *PLoS ONE* **2017**, *12*, e0181799.

[46] E. Sedlák, J. V. Schaefer, J. Marek, P. Gimeson, A. Plückthun, *Protein Sci.* **2015**, *24*, 1100–1113.

[47] T. Miyafusa, H. Watanabe, S. Honda, *Int. J. Biol. Macromol.* **2021**, *182*, 1697–1703.

[48] M. Hebditch, J. Warwicker, *Sci. Rep.* **2019**, *9*, 1969–9.

[49] C. Zhang, N. Codina, J. Tang, H. Yu, N. Chakroun, F. Kozielski, P. A. Dalby, *Comput. Struct. Biotechnol. J.* **2021**, *19*, 2726–2741.

[50] T. Nakamura, T. Aizawa, R. Kariya, S. Okada, M. Demura, K. Kawano, K. Makabe, K. Kuwajima, *J. Biol. Chem.* **2013**, *288*, 14408–14416.

[51] V. S. Jasion, B. P. Burnett, *Nutr. J.* **2015**, *14*, 1–8.

[52] M. Onitsuka, Y. Kadoya, T. Omasa, *J. Biosci. Bioeng.* **2019**, *127*, 752–757.

[53] Y. Senga, M. Doi, M. Onitsuka, S. Honda, *Cell Chem. Biol.* **2021**, *28*, 1–13.

[54] H. Watanabe, S. Yageta, H. Imamura, S. Honda, *Anal. Chem.* **2016**, *88*, 10095–10101.

[55] V. Filipe, B. KüKrer, A. Hawe, W. Jiskoot, *J. Pharm. Sci.* **2012**, *101*, 2327–2339.

[56] O.-O. Oyetayo, H. Kiefer, *Pharm. Res.* **2016**, *33*, 1359–1369.

[57] H. Imamura, S. Honda, *J. Phys. Chem. B* **2016**, *120*, 9581–9589.

[58] T. Skamris, X. Tian, M. Thorolfsson, H. S. Karkov, H. B. Rasmussen, A. E. Langkilde, B. Vestergaard, *Pharm. Res.* **2016**, *33*, 716–728.

1   [59]   H. Imamura, A. Sasaki, S. Honda, *J. Phys. Chem. B* **2017**, *121*, 8085–8093.
2   [60]   H. Imamura, S. Honda, *Methods Enzymol.* **2019**, *622*, 329–345.
3   [61]   M. Hebditch, R. Kean, J. Warwicker, *Comput. Struct. Biotechnol. J.* **2020**, *18*, 897–905.
4   [62]   E. Gasteiger, C. Hoogland, A. Gattiker, S. Duvaud, M. R. Wilkins, R. D. Appel, A.
5   Bairoch, Humana Press, Totowa, NJ, **2005**, pp. 571–607.
6   [63]   N. Shimizu, T. Mori, Y. Nagatani, H. Ohta, S. Saijo, H. Takagi, M. Takahashi, K.
7   Yatabe, T. Kosuge, N. Igarashi, *AIP Conf. Proc.* **2019**, *2054*, 060041.
8   [64]   T. C. Huang, H. Toraya, T. N. Blanton, Y. Wu, *J. Appl. Crystallogr.* **1993**, *26*, 180–184.
9   [65]   P. Bernadó, N. Shimizu, G. Zaccai, H. Kamikubo, M. Sugiyama, *Biochim. Biophys. Acta*
10   **2018**, *1862*, 253–274.
11   [66]   S. R. Kline, *J. Appl. Crystallogr.* **2006**, *39*, 895–900.
12   [67]   P. Schuck, *Biophys. J.* **2000**, *78*, 1606–1619.