Getting Smaller by Denaturation: Acid-Induced Compaction of Antibodies

Hiroshi Imamura[1,2,3], Ayako Ooishi[1], Shinya Honda[1*]

[1]Biomedical Research Institute, National Institute of Advanced Industrial Science and Technology, 1-1-1, Higashi, Tsukuba, Ibaraki 305-8566, Japan

[2]Department of Applied Chemistry, College of Life Sciences, Ritsumeikan University, 1-1-1 Noji-Higashi, Kusatsu, Shiga 525-8577, Japan

[3]Department of Bio-Science, Nagahama Institute of Bio-Science and Technology, 1266 Tamura, Nagahama 526-0829, Japan

ORCID: Hiroshi Imamura, 0000-0001-8924-747X; Shinya Honda, 0000-0002-7878-3944

*Corresponding Author:

Shinya Honda

Biomedical Research Institute

National Institute of Advanced Industrial Science and Technology

1-1-1, Higashi, Tsukuba, Ibaraki 305-8566, Japan

E-mail: s.honda@aist.go.jp

https://unit.aist.go.jp/bmd/biomed-mcd/t/

Abstract

Protein denaturation is a ubiquitous process that occurs both *in vitro* and *in vivo*. While the molecular understanding of the denatured structures of proteins is limited, it is commonly accepted that the loss of unique intramolecular contacts makes proteins larger. Herein, we report compaction of the immunoglobulin G1 (IgG1) protein upon acid denaturation. Small-angle X-ray scattering coupled with size exclusion chromatography revealed that IgG1 radii of gyration at pH 2 were ~75% of those at a neutral pH. Scattering profiles showed a compact globular shape, supported by analytical ultracentrifugation. The acid denaturation of proteins with size reduction is energetically costly, and acid-induced compaction requires an attractive force for domain reorientation. Such intramolecular aggregation may be widespread in immunoglobulin proteins as non-canonical structures. Herein, we discuss the potential biological significance of these non-canonical structures of antibodies.

1 Introduction

Proteins are polymer chains composed of amino acid residues and optimised sequences

that have the distinct thermodynamic states,[1] one of which is a native state and another a denatured

state. The folded structure in the native state has been determined by extensive biophysical

investigations and predicted via bioinformatic methods. In contrast, the molecular pictures for the

structures in the denatured state are limited. The denatured state comprises a variety of dynamic

conformers, including fully and partially unfolded structures, which has technically hindered the

elucidation of their fine structures. Despite this, recent studies have explored the difference in the

denatured structure causing different biological and immunological consequences; for example,

an expanded conformation of the Tau protein is favourable for biomolecular condensation, which

promotes amyloid formation.[2] Whether unfolded structures of a protein (PNt) are expanded or

collapsed determines the efficiency of the secretion to the cell surface.[3] Reversal effects of

therapeutic proteins are potentially associated with the morphology of the denatured aggregates.[4,5]

These studies indicate the significance of accumulating knowledge about the denatured structures

of proteins.

Both amino acid side chains and main chains stabilise the native protein structure by

forming non-covalent bonds among them, where the constituent atoms are uniquely organised.

Upon denaturation, these native contacts are replaced by non-specific water-protein contacts that

increase the spatial and temporal dynamics of the proteins.[6] Thus, the size of proteins is always

increased by denaturation,[7-9] which is a major principle of protein folding.

Here, we report an exceptional case where denaturation makes proteins smaller via an

immunoglobulin G1 (IgG1) monoclonal antibody. Acid-solution induces the denaturation of IgG1,

including the loss of the binding function[10] and native structures.[11,12] However, the structures of

the denatured IgG1 remain unclarified. This study used a small-angle X-ray scattering (SAXS)

technique[13] to demonstrate the size reduction of IgG1 by probing the global solution structure of

the antibody under neutral and acidic pH conditions. While the biological significance of the

reduced size of denatured IgG1 was unclear, the current models include the molecular

requirements of non-canonical contacts between domains, suggesting the universality of the

phenomenon of denaturation reducing the size of immunoglobulin proteins.

11 Figure 1. The mAb-A pH-dependence of the small-angle X-ray scattering (SAXS) at 25 °C.

12 Concentration-normalised SAXS profiles are presented as (a) log–log and (b) Kratky plots. The

13 protein concentration was 3.5 ± 0.1 mg/mL. The data are vertically shifted for clarity. The black

14 line shows the theoretical SAXS profile of the IgG1 model (PDB: 1HZH). The arrows in (a)

15 indicate the shoulders at $q = \sim 0.08$ and ~ 0.16 Å^{-1}.

Results

Figure 1 depicts the acid-induced changes in an IgG1 monoclonal antibody, mAb-A, when

measured by SAXS. The SAXS profiles were similar at pH 7.1–3.1, with the characteristic

shoulders in the log–log plots (Figure 1a, arrows) and peaks in the Kratky plots (Figure 1b) at $q =$

~0.08 and ~0.16 Å^{-1}. The former trait (q = ~0.08 Å^{-1}) corresponds to the Fc (crystallizable

fragment)–Fab (antigen-binding fragment) or Fab–Fab distance (~80 Å). The latter trait (~0.16 Å^{-}

^{1}) is sensitive to the inter-domain distances/orientations between the adjacent domains.[14] These

scattering patterns share these characteristics with the theoretical scattering calculated from the

crystal structure of IgG1 (black line in Figure 1; denoted as "Model") and previous SAXS data

reported in the literature.[15,16] Conversely, the scatterings at pH < 2.0 were distinct from those at

pH 7.1–3.1; the characteristic shoulders (Figure 1a) and peaks (Figure 1b) diminished at pH < 2.0,

with more scattering in the low-q range at pH 1.1 (Figure S1a).

The SAXS data indicated a drastic conformational alteration of mAb-A around pH 2, but

the contaminating aggregates hampered further structural analysis (see Supporting Information

and Figure S1). Therefore, we performed SEC-SAXS analysis[17-19] *in situ* SAXS during size

exclusion chromatography (SEC), which allowed us to monitor the SAXS of monomeric mAb-A

at pH 2. Chromatographic analysis was conducted of the integral SAXS intensities and ultraviolet

(UV) absorbance at 280 nm using an elution buffer solution of 0.1 M glycine-HCl (pH 2) with 0

M or 0.2 M NaCl (Figure 2a). The peaks at ~169 min (0 M NaCl) and ~66 min (0.2 M NaCl)

represented monomeric mAb-A that was well separated from the aggregates, while the peaks at

~124 min (0 M NaCl) and ~56 min (0.2 M NaCl) indicated dimeric mAb-A. The aggregates (>

trimer) manifested as peaks that presented early in the analysis. These assignments are supported

by the chromatogram of the integrated SAXS intensities, which have more pronounced aggregates

than the UV absorbance. The variation in the retention time between the data (0 M and 0.2 M

NaCl) was due to the difference in the flow rates used.

Based on the chromatograms and their frame-by-frame analyses (Figure S2), we

determined the concentration-normalised SAXS profiles ($I(q)/c$) of monomeric mAb-A at pH 2

(Figure 2b). Because the concentrations were < 0.5 mg/mL in the sample cell during the

chromatographic analysis, the intermolecular interference was negligible. The radius of gyration

(R_g) was 50.8 ± 0.3 Å for 0 M NaCl and 40.5 ± 0.4 Å for 0.2 M NaCl, and the concentration-

normalised zero-angle scattering ($I(0)/c$) was 0.0894 ± 0.0003 cm^2 mg^{-1} (122 kg/mol) for 0 M

NaCl and 0.0783 ± 0.0004 cm^2 mg^{-1} (108 kg/mol) for 0.2 M NaCl, which were comparable with

the native monomeric mAb-A of 0.0880 ± 0.0007 cm^2 mg^{-1} (121 kg/mol). The comparable molar

masses (Table S1) were found via Bayesian inference[20] These results indicated that the SEC-

SAXS was able to capture the monomeric acid-denatured mAb-A by removing the aggregates.

The deviations from the exact molar mass (M) of mAb-A (148 kg/mol) were due to a systematic

error. In the log–log and Kratky plots of the SAXS profiles (Figure 2b and S2h), the characteristic

shoulders/peaks observed for the native mAb-A at pH 7.1 (Figure 1) was not observed at pH 2.

The distance-distribution function ($P(r)$) at pH 2 indicated a lack of the characteristic distance

distribution (~80 Å) observed at pH 7.1 (Figure 2d), suggesting the distances and orientations of

the Fab-Fc and Fab–Fab regions were not canonical at pH 2. Furthermore, the maximum distance

(D_{max}) at pH 2 were < ~200 Å (0 M NaCl) and < ~150 Å (0.2 M NaCl). The former was comparable

with D_{max} at pH 7.1.

A fitting analysis of the SAXS profiles characterised the shape of mAb-A at pH 2 (Figure

2b). Among the theoretical scattering functions of the sphere, cylinder, elliptical cylinder, ellipsoid,

and triaxial ellipsoid shapes,[21,22] the scattering function of the triaxial ellipsoid ($a \neq b$) was the

best approximation, with lengths, a, b, and c defining the shape of the triaxial ellipsoid (Figure 2b).

The scattering function of the ellipsoid ($a = b$) produced deviations from the experimental

scatterings in the region of $q > 0.1$ Å^{-1} in Figure 2b and $qR_g > 4$ in Figure 2c; such a deviation was

also found in $P(r)$ functions (Figure 2d). These deviations for $I(q)/c$ and $P(r)$ at 0 M NaCl were

larger than those at 0.2 M NaCl, indicating that the shape of mAb-A at 0 M NaCl was more

flattened than that at 0.2 M NaCl.

Figure 2c shows the dimensionless Kratky plots. The SAXS of an ideal globular protein

obeys Gunier's law, $I(q) = I(0)\exp(-(R_g q)^2/3)$, and shows the maximum value where the black

dotted lines intersect in the dimensionless Kratky plots:[23,24] $qR_g = 3^{1/2}$ and $(qR_g)^2 I(q)/I(0) = 3\exp(-$

$1) = 1.104$. The maximum peak, 0.2 M NaCl, of the dimensionless Kratky plot indicates that the

shape of mAb-A closely resembles that of the ideal globular proteins. Slight deviation of the

maximum at 0 M NaCl from the "ideal" maximum would be due to the flattened ellipsoidal shape,

although the relevance thereof to conformational flexibility could not be rejected.

Figure 2e visualises the triaxial ellipsoidal approximation with the best fit parameters for

the scatterings of mAb-A. The representation of mAb-A at pH 2 with 0 M NaCl (grey) and 0.2 M

NaCl (blue) was flattened, in comparison with that of the native IgG1 (red). We observed that 0.2

M NaCl made mAb-A more compact with a smaller maximum length and globular shape than 0

M NaCl. Figure 2f shows the reconstructed 3D-models of mAb-A at pH 2 from the SAXS profiles

and $P(r)$ by the *ab initio* electron density determination, DENSS (DENsity from Solution

Scattering), algorithm.[25] The flattened shapes of mAb-A at pH 2 were also consistent with those

obtained via the fitting analysis that used the theoretical scattering functions. In addition, these 3D

models illustrated some anisotropic electron distributions, such as a lower density region that

looked like a "neck" in 0 M NaCl. Another remarkable characteristic of mAb-A at pH 2 included

a smaller R_g than that of the native mAb-A (Figure 2g). The addition of NaCl led to a smaller R_g

at pH 2, while R_g was not dependent on NaCl at pH 7. mAb-A at pH 2 was very plastic and

presented as globular due to the shielding effect of the salt, which reduced the intramolecular

electrostatic repulsion of the acid-denatured mAb-A. We noted that the incubation time at pH 2

did not significantly change the R_g.

We further conducted SEC-SAXS analysis on another IgG1, mAb-B, to test whether a

variation of the sequence in the variable region alter the conclusion of the size reduction. We

verified an acid-induced compaction of mAb-B, which showed $R_g = 53.0 \pm 0.4$ Å at pH 7.2 and R_g

$= 46.5 \pm 0.3$ Å at pH 2 and 0.2 M NaCl.

Figure 2. Size exclusion chromatography small-angle X-ray scattering (SEC-SAXS) analysis on

mAb-A at pH 2. The results of mAb-A at 0 M NaCl and 0.2 M NaCl are represented as grey and

cyan, respectively. (a) Overlaid elution SAXS and UV profiles where the SAXS-derived

chromatogram is an integrated SAXS intensity vs. retention time. The elution UV profiles were

acquired by simultaneous recording of mAb-A absorbance at 280 nm. (b) SAXS profiles of

monomeric mAb-A. The scattering for mAb-A in 0.2 M NaCl was multiplied by 10^2 for clarity.

Theoretical scattering of the triaxial ellipsoid (red solid line) and ellipsoid (dashed black line) with

best fit parameters for mAb-A are shown. An elliptical object indicates where a, b, and c are the

length parameters. (c) Dimensionless Kratky plots of IgG1 SAXS profiles and theoretical

scattering. (d) The distance-distribution function, $P(r)$, by direct Fourier transformation of the

SAXS profiles. $P(r)$ of the triaxial ellipsoid (red solid line) and ellipsoid (black dashed line) were

calculated from the scatterings in (b). $P(r)$ of the native mAb-A at pH 7.1 (Figure S1e) is overlaid

for comparison. This $P(r)$ was scaled such that the area is equivalent to the $P(r)$ of the mAb-A at

pH 2 and 0 M NaCl. (e) A point cloud representation of the triaxial ellipsoids using best fit

parameters for the scatterings of mAb-A at 0 M NaCl and 0.2 M NaCl. A canonical IgG1

structure,[26] where the C_α atoms (red point clouds) are placed to fit with the triaxial ellipsoids with

characteristic lengths. (f) 3D reconstructed models of the mAb-A at pH 2 using *ab initio* electron

density determination (DENSS).[25] The canonical mAb-A models are also shown, indicating the

ab initio model reconstructed from the SAXS data (light pink), overlaid with the atomic

coordinates of the IgG1 (red ribbons).[26] All the volumes inside the model surfaces are set at

258,000 $Å^3$, where the Porod volume was determined for the scattering of mAb-A at 0 M NaCl.

(g) Dependence of mAb-A R_g at pH 2 on incubation time and NaCl concentration. The R_g of the

native mAb-A at pH 7 (with 0 M NaCl and 0.1 M NaCl) is indicated for comparison.

An orthogonal analysis of the acid-denatured mAb-A using an analytical ultracentrifugation

(AUC) strengthened the SAXS analysis. The sedimentation velocity data (Figure S3a) showed the

distribution of the sedimentation coefficients (s) of the mAb-A samples at pH 7 and pH 2 (Figure

S3b), in which prominent peaks at $s = {\sim}6.5{-}7.0$ S were consistent with the expected s value for the

known molar mass of monomeric mAb-A (148 kg/mol; Table S2). We found a small but significant

variation in $s_{20,w}$ (s at 20 °C in water) among the samples, where the effects of the viscosities and

densities of the solvents were corrected. This variation indicates the differences in the molecular

envelope, represented as the friction factor in the AUC. The variation in $s_{20,w}$ due to the association

of the molecules was ruled out; for example, the dimer of mAb-A had a $s = {\sim}9.1$ S at pH 2 and 0

M NaCl (Figure S3b). According to equations (S8–S11), the friction factor (f/f_0) and the corrected

friction factor ($f/f_{0,sp}$) were determined. The f/f_0 of the native mAb-A (1.43) was comparable with

a previously reported value (1.55);[27] the deviation should be due to the difference in the specific

volume (v) used. Empirically, approximately globular proteins, moderately elongated proteins, and

highly elongated proteins were reported to display f/f_0 of 1.2–1.3, 1.5–1.9, and 2.0–3.0,

respectively.[28] The current f/f_0 indicated that the native mAb-A and the acid-denatured mAb-A

were both globular rather than elongated. The f/f_0 at pH 2 decreased from 1.58 to 1.35 with the

addition of 0.2 M NaCl, consistent with the SAXS result that mAb-A at pH 2 was more globular

in 0.2 M NaCl. The corrected friction factor ($f/f_{0,sp}$) in equations (S9–S11) provided information

regarding the protein shape and can be calculated for any triaxial ellipsoid.[29,30] We calculated $f/f_{0,sp}$

using the triaxial ellipsoid lengths a, b, and c, as determined using the SAXS analysis (Figure 2).

The experimentally determined $f/f_{0,sp}$ at pH 2 decreased by 0.19 (1.32 at 0 M NaCl and 1.13 at 0.2

M NaCl), which was comparable with the decrease in the calculated $f/f_{0,sp}$ by 0.14 (Table S2). The

differences between the absolute values of the experimental $f/f_{0,\mathrm{sp}}$ and the calculated value were

due to uncertainties in the β_1, v, and η used in equations (S7 and S8). Together, the results of

SAXS—the non-elongated shape of the acid-denatured mAb-A and its plastic globularisation by

the addition of NaCl— are well supported by the AUC.

Discussion

Protein denaturation increases the size of proteins independent of perturbations such as

temperature, pressure, or chemical denaturation.[8] Indeed, the R_g for proteins reported in the

literature indicated that the underlying power laws of R_g vs. chain lengths (L) are distinct between

native and unfolded protein structures (Figure 3). The R_g of the native state of the multidomain

protein mAb-A followed the power–law relationship for the native structures, although the

representation of R_g in the literature[9] is dominated by single-domain proteins. In contrast, the R_g

of non-native mAb-A was not predicted by the power–law relationship and anomaly smaller than

that of the native state. The R_g of the non-native mAb-A dimer, apart from the scaling law for

unfolded proteins, also indicates a non-random coil structure.

Figure 3. R_g dependence on protein chain length. The data in literature includes native (green,

closed circles), native oligomeric (dark orange, closed circles), and unfolded (black triangles,

where closed and open triangles indicate disulfide-free and disulfide-bonded) proteins, curated by

Po-Min Shih *et al.*[9] The dotted lines represent the power–law scaling of $R_g = R_0 L^\xi$, where R_0 is a

prefactor (2.04, native; 4.47, native oligomer; and 2.05, unfolded), L is the amino acid (a.a.) chain

length, and ξ is the scaling exponent (0.45, native; 0.32, native oligomer; and 0.58, unfolded). R_g

of mAb-A at pH 7 (red, open circle), R_g of mAb-A at pH 2 with 0.2 M NaCl (red, open triangle),

and R_g of mAb-A dimer at pH 2 with 0.2 M NaCl (red, open diamond) determined in this study

are overlaid onto previously reported results.[9] The R_g value of mAb-A at pH 7 is given as an

infinite dilution of the protein (Figure S1h).

Protein acid denaturation is caused by the repulsion of positive charges on the surface of the

protein that are approximated by simple models such as the Linderstrøm–Lang smeared charge

model.[31,32] This model indicates that the size expansion decreases the Gibbs energy by

denaturation (see Supporting Information) and describes how acid denaturation that reduces

protein size is energetically costly and anomalous. No previous reports have shown acid

denaturation without size expansion; even the molten-globule state of cytochrome c at pH 2 and

0.1 M NaCl (17.4 Å), which is described as a "compact" denatured state, is larger than that at its

native state (13.5 Å).[33]

The charge distributions shown in Figure 4a identified a positive potential on the surface of

mAb-A that widely emerged at pH 2. We hypothesise that alternative, energetically favourable

interactions hold the parts of the acid-denatured IgG1 molecule together by compensating for the

energetic cost of maintaining the size of the protein. Additional van der Waals interactions would

assist in this process. The acid-denatured state at pH 2 was more compact with the addition of

NaCl, indicating that shielding the coulomb repulsion among the charged residues and

unfavourable salt-induced solvation of the exposed hydrophobic protein surface assisted the sub-

complete packing of the protein. These effects could be common for inter- and intra-molecular

interactions. Lower pH increases intermolecular repulsive interactions, which is depicted as the

intermolecular interference (Figure S1f-i). Counteracting favourable interactions between the

monomers could generate the aggregates at pH 2.

Figure 4. Acid effects on IgG structure. (a) Charge distribution in IgG1 at various pHs. The electrostatic properties were calculated using the pK_a predictor PROPKA[34,35] and the adaptive Poisson–Boltzmann solver (APBS) software[36] using the atomic coordinates for IgG1[26] and the mAb-A sequence. (b) The conformational states of the IgG domains at pH 2.0–2.6 (with/without salts) in the context of the isolated domain, Fc/Fab regions, the full-length IgG reported in the literature; the references are presented in the table are [a]pH 2.0;[37] [b]aglycosylated C_H2, pH 2;[38] [c]pH 2;[39] [d]aglycosylated Fc, pH 2.6;[40] [e]aglycosylated Fc, pH 2.5;[41] [f]glycosylated Fc, pH 2.6;[40] [g]pH 2;[42] [h]pH 2.0;[43] [i]pH 2;[44] [j]pH 2.5;[45] [k]pH 2.1;[46] [l]glycosylated Fc, pH 2.0;[47] [m]pH 2.0;[11] and [n]pH 2.0.[12] The canonical IgG1 structure,[26] the domains thereof (C_H3, C_H2, C_H1, C_L, V_H, and V_L), and the regions (Fc and Fab) are presented.

The gain in native contacts in folded proteins contributes to their size reduction.[7]

Therefore, the acid-denatured state for mAb-A could be regarded as an alternatively folded state

(AFS) that had alternative contacts that prevented expansion. Buchner *et al.*[48] referred to AFS

hitherto as "The antibody conformation at low pH had been termed 'alternatively folded state'

because it exhibits characteristics of the folded state (e.g. remarkable stability against unfolding),

but the available spectroscopic information suggests that it is structurally significantly different

from the native state. It was first described for a complete IgG1 antibody,[11] but single domains

such as C_H3 also can adopt this state.[39]" The original paper published in 1991[11] characterised

MAK33, a murine κ/IgGl subtype, at pH 2. The $s_{20,w}$ of 5.53 for the AFS (pH 2) of MAK33 was

smaller than the $s_{20,w}$ of 6.60 for the native protein, and much larger than $s_{20,w}$ of 3.2 for the

denatured protein in 6 M guanidine hydrochloride at a pH 2. These AUC results indicated that the

molecules were neither significantly aggregated nor fully extended. Nevertheless, the small

amount of the aggregates functioned as impurities hindered the discovery that the AFS state of

IgG1 can be smaller than the native state. As the SEC-SAXS and the sedimentation coefficient

distribution analysis for the AUC have not been established previously, the technical limitations

also hindered the discovery. The circular dichroism (CD) spectra of the AFS state of MAK33 at

pH 2 were similar to the current CD spectra (Figure S4a), further supporting the AFS of the present

acid-denatured mAb-A in which the β-sheets are dominant (Table S3) and slightly reorganised, as

indicated using infrared (IR) analysis (Figure S4c). Subsequent studies on AFS focused on the

isolated protein domains such as C_H3 or V_L domains, rather than full-length antibodies. Buchner

et al.[49] reported that the MAK33 V_L domain could form stable high-molecular weight oligomers

at pH 2 that had previously been reported as AFS. Although the term AFS is loosely defined, it

can be concluded that the association between protein domains is a common mechanism for AFS

irrespective of whether full-length antibodies or isolated domains are evaluated.

Studies have also reported that other IgG molecules (murine monoclonal IgG2a antibody

CB4-1[46] and monoclonal chimeric antibody IgG1 BR96[12]) can adopt the AFS state at pH 2. In

addition, Sedlák *et al*[50] proposed an inter-region or inter-domain interaction of heat-denatured

IgG, referred to as "intramolecular aggregation". The authors of that study compared differential

scanning calorimetry (DSC) thermograms of full-length IgG and its Fab portion. The difference in

the melting temperatures of the Fab domain suggested a Fab–Fab interaction in the heat-denatured

state of the protein, which was supported by the lack of protein concentration-dependence in the

thermograms.[50] The concept of "intramolecular aggregation" would be universal for IgG

irrespective of the denaturing perturbations (e.g., acid or heating).

We also investigated the domains that participated in inter-domain interactions in acid-

denatured IgG1. The canonical IgG1 protein consists of 12 domains in the Fc or Fab regions

(Figure 4b). According to the literature,[11,12,37-46,51] the stabilities of the domains depend on the

molecular contexts (Figure 4b), e.g., isolated domains or domains in a full-length IgG. Isolated

C_H3[38] at pH 2 and C_H3 in an aglycosylated Fc[41] at pH 2.5 unfold; hence, the native dimerised

C_H3–C_H3 interface is lost.[38,41] However, the ordered structure of C_H3 in the full-length IgG is

possibly retained[42,43] with local disorder that decreases the left-twisting of β-strands (Figure S4

and Table S3). The C_H2 unfolds at pH 2 regardless of the presence of glycans in any molecular

context.[38,40-43] The isolated domains (C_H1, C_L, and V_L) of the Fab region unfold at pH 2,[37] whereas

the Fab region is structured when the Fab region protein[44-46] and full-length IgG[42,43,46] protein are

examined. Therefore, the possible mechanism of acid-induced intramolecular aggregation of IgG1

is as follows: at pH 2, C_H3 avoids complete unfolding by interacting with the Fab region, where

the flexibility of the unfolded C_H2 does not hamper the inter-domain contacts. Salt weakens the

Fab–Fab repulsion, which brings the two Fab regions close to each other and makes IgG1 smaller.

Self-interactions between C_H3 are stabilised by salt, indicated by the AFS of the isolated C_H3

oligomer.[39] We speculated that the hydrophobic V_H–V_L and C_H1–C_L interfaces[52] would be

exposed upon breaking the native interfacial contacts and the inter-domain salt bridges.[53] All those

effects could make non-canonical contacts between the Fab and Fc regions.

A lack of atomistic modelling or the ensemble representation thereof is a limitation of this

study. A flexible hinge in IgG allows for it to enable diverse positioning of the Fab and Fc regions.

Tian *et al*. [16] used this ensemble of models to improve the fitting of a native IgG1 SAXS profile

instead of using a single model. The possible distribution of R_g of the native IgG1 was ca. 45–58

Å. This indicates that the minimal requirement for getting smaller is the non-canonical positioning

of the Fab and Fc regions, which never populate in the native state. The acid-induced structural

changes, manifested as the decline of the Kratky peak at $q = 0.16$ Å^{-1} (quaternary structure), and

the CD/IR spectral change (secondary structure) make the irregular Fab/Fc arrangements

energetically favourable. The present shape approximation of the acid-denatured mAb-A using the

theoretical scatterings of the defined shapes and the DENSS method does not rule out the ensemble

of conformations that is compatible with the SAXS profile but represents the dominant structures.

It is intriguing to determine whether this non-canonical structure of the acid-denatured

IgG1 functions similarly to an α-lactalbumin in the molten globule state (human α-lactalbumin,

HAMLET), which is lethal to tumour cells.[54] A few reports have shown the biological significance

of the AFS of immunoglobulins; for example, the oral administration of Ig (IgG and IgA) has been

reported as a passive immunity treatment. Orally administered Ig is exposed to the acidic pH of

the gastrointestinal (GI) tract but is resistant to complete digestion.[55] Jasion and Burnett[11]

suggested that transient exposure of IgG to the acidic conditions in the stomach may increase the

stability of IgG as it passed into the GI tract and against proteolytic enzymes.[55] The non-canonical

structure of acid-denatured IgG1 presented herein supports the idea that reducing the flexibility of

locally unfolded regions via intramolecular aggregation is beneficial for the protein's resistance to

enzymatic digestion. The ions present in physiological conditions could also support the tight

packing of the acid-denatured IgG1 (Figure 2). Another example regards the expression and

secretion of natural or recombinant Ig proteins: Onitsuka *et al*.[56] found that antibody-producing

Chinese hamster ovary cells secreted aggregates of recombinant IgG from the intracellular

environment into the extracellular milieu, further evidenced by a new live–cell imaging

technique[57] that uses a fluorescently labelled AF.2A1 as a protein probe for non-native IgG

structures.[51,58] Similar intracellular particles, known as Russell bodies, are found in plasma cells.

Russell bodies are natural inclusion bodies usually found in cells that overexpress Ig. Mammalian

cells generally secrete only correctly folded proteins; therefore, a naturally occurring AFS could

pass through an intracellular quality control system, such as endoplasmic reticulum (ER)-

associated degradation.[56,57] If the naturally occurring AFS was similar to the acid-denatured IgG1,

the compact form may be favourable for Ig intracellular transport, secretion, internalisation, and

transcytosis in various circumstances.

The acid-denatured state of IgG1 is relevant to downstream processes of biopharmaceutical

manufacturing where antibodies experience temporal exposure to acid during affinity

chromatographic purification and virus inactivation, which is followed by neutralisation. The non-

native conformation of IgG1 induced by this pH shift is aggregation-prone at a neutral pH, with

structural origins that remain unclear.[10,59-64] Because the acid-denatured state is unstable at a

neutral pH, the IgG1 molecules transition toward native or aggregated states. It is unclear whether

the molecular characteristics of the acid-denatured IgG1, such as intramolecular aggregation, slow

or accelerate these reactions. In addition, the precise prediction of antibody stability in an acidic

environment requires the characterisation of acid-denatured structures, including in the dimeric

forms (Figure S5), which is important aspect of the engineering process of antibodies.[65]

Conclusion

The current knowledge-based principle of proteins getting larger by denaturation that has

been established by many studies using single-domain proteins should be reconsidered. New view

should now include the potential for proteins to get smaller by denaturation, as we demonstrated

for acid-denatured IgG1 proteins (mAb-A and mAb-B). The reported cases are limited;

nevertheless, as they shared the characteristics of the previously reported AFS for IgGs, an acid-

induced compaction can be common among various IgG proteins. Multidomain proteins possess

additional higher-order structures, such as domain–domain spatial arrangements or orientations,

which are lost upon denaturation. These degrees of freedom in multidomain proteins are

prerequisites for a more compact structure by the rearrangement of the domains, in which there

are minor secondary or tertiary structural changes. In contrast, single-domain proteins are almost

completely packed, with no degrees of freedom for further compaction. The compact denatured

structure induced by intramolecular aggregation may be ubiquitous in immunoglobulin proteins as

non-canonical structures.

1 Materials and Methods

Two monoclonal antibodies (mAb) commercially available as biotherapeutics were used;

one is mAb-A, a humanised IgG1 monoclonal antibody (148 kDa; calculated pI[66] = 8.5), and the

other is mAb-B, a human IgG1 monoclonal antibody (147 kDa; calculated pI[66] = 8.4). The molar

mass includes the glycan mass. IgG1 was dissolved in a buffer solution (0.01 M sodium phosphate,

0.15 M NaCl and 0.005% (v/v) polyoxyethylene (20) sorbitan monolaurate, pH 7.4) to be ~1–100

mg/mL (stock solution of IgG1). The IgG1 stock solution was diluted to the given concentrations

(based on the experiment) with 0.01 M sodium phosphate buffer solution (pH 7.4) and dialysed

against 0.01 M sodium phosphate buffer solutions at pH 3, 4, 6, and 7, or 0.1 M glycine-HCl buffer

solutions at pH 1, 2, and 3 at 4 °C for 7 h using a Toru-kun TOR-14K microdialyzer (NIPPON

Genetics Co., Ltd., Tokyo, Japan) with a membrane with a cut-off of 14 kDa. During dialysis and

incubation, the protein aggregated at pH 1–2 (Figure S1). These samples were stored at 4 °C for

~14 h until the batch SAXS measurement (Figure 1). Both the sodium phosphate and glycine-HCl

buffer solutions at pH 3 produced comparable SAXS profiles (the latter is presented in Figure 1).

For the SEC-SAXS measurement (Figure 2), the ~10 or ~20 mg/mL IgG1 stock solutions were

dialysed against 0.1 M glycine-HCl buffer solution (pH 2.0) with a membrane cut-off of 12–14

kDa (Scienova GmbH, Jena, Germany) and stored at 4 °C for 1 h, 1 d, 3 d, or 7–8 d before

measurement.

SAXS The SAXS experiments were performed with a BL-10C beam line at the Photon Factory of

the High Energy Accelerator Research Organization (KEK) in Tsukuba, Japan.[67] The X-ray

wavelengths (λ) used were 0.10, 0.12, or 0.15 nm. The camera length was 1 or 2 m, and was

calibrated using the scattering pattern of silver behenate.[68] The scattering parameter q is defined

as $q = |\boldsymbol{q}| = 4\pi\sin\theta/\lambda$, where $\boldsymbol{q}$ is the scattering vector and 2θ is the scattering angle of the X-rays.

X-ray intensities were recorded with a PILATUS3 2M detector (DECTRIS Ltd., Baden, Switzerland). For the batch SAXS measurement (Figure 1), the samples were measured at 25 ± 0.1 ℃ in a cell with quartz windows; 30 images were collected, where one image was recorded during a 2-s exposure time; the data from samples damaged by radiation were eliminated.

For the SEC-SAXS measurement (Figure 1), the IgG1 samples (~7–11 mg/mL, 100–150 µL) were passed through a stainless-steel sample cell via an in-line size exclusion column (Superdex200™ (10/300) GL; GE Healthcare UK Ltd., Amersham, England; Lot. 10111292 and 10216439) operated by an HPLC system (Acquity UPLC H-class and Empower3 FR2; Waters Co., Milford, MA, USA; or Nexera-i; Shimadzu Co., Kyoto, Japan) installed with a beam line BL-10C[67,69] and using flow rates of 0.2 mL min^{-1} for the initial 30 min and then 0.05 mL min^{-1} (Figure 2a, lower data) or 0.2 mL min^{-1} (Figure 2a, upper data). The temperature of the sample cell and the column was maintained at 25.0 °C. The exposure time for recording the image was 20 s. The sample solution was eluted into the sample cell with 0.02 mm-thick quartz glass windows and 1 mm-path length, where UV–vis absorption spectra (~2 mm-path length by the diagonal incident light on the cell) were simultaneously recorded using a fibre-coupled UV–vis spectrometer, QEpro, with Opwave+ version 2.14–2.16 (Ocean Optics, Inc., Largo, FL, USA). A fitting analysis using theoretical scattering functions was conducted using the codes provided by the NIST Center for Neutron Research (NCNR).[70] Unless otherwise indicated, all analyses were performed using IGOR Pro version 6.22A (WaveMetrics, Portland, OR, USA). SAXS measurements and the analyses of data are detailed in the Supporting Information and Table S4 therein.

AUC mAb-A samples at 0.7 mg/mL were used. Sedimentation velocity measurements were performed in a ProteomeLab XL-A (Beckman Coulter, Inc., Brea, CA, USA) with rotor speeds of 42 krpm at 25 ± 0.2 ℃. An absorbance of 280 nm was used to monitor the protein

concentration in a double-sector cell. The sedimentation velocity data were processed with the

SEDFIT program using a sedimentation coefficient distribution analysis.[71] The parameters (s, f/f_0

and $f/f_{0,\mathrm{sp}}$) were determined according the equations in the Supporting Information.

CD and IR mAb-A samples at 1 mg/mL were measured. The procedures are detailed in

the Supporting Information.

Acknowledgements

This research is partially supported by the Japan Society for the Promotion of Science (grant

numbers: JP19H03363 [to S.H.] and JP21K06503 [to H.I.]). The SAXS experiment in this work

was performed under the approval of the Photon Factory Program Advisory Committee (Proposal

No. 2015G061, No. 2017G068, and No. 2020G077). Computations were partially performed on

the NIG supercomputer at ROIS National Institute of Genetics. The authors thank Dr. Tomoshi

Kameda (AIST, Tokyo, Japan) and Dr. Seiki Yageta (AIST, Tsukuba, Japan) for stimulating

discussions and comments.

Author Contributions:

H.I. and S.H. designed the research. H.I. and A.O. performed the experiments. H.I. analysed the

data. H.I and S.H. wrote the main manuscript text. All authors contributed to data interpretation,

critically reviewed the manuscript, and approved the final version of the manuscript.

Conflict of Interest:

The authors declare no competing financial interests.

Data Availability. All study data are included in the article and/or *SI Appendix*.

REFERENCES

[1] Lesk, A. M. (2016). *Introduction to protein science: Architecture, function, and Genomics*. Oxford University Press.

[2] Wen, J., Hong, L., Krainer, G., Yao, Q.-Q., Knowles, T. P., Wu, S., & Perrett, S. (2021). Conformational expansion of tau in condensates promotes irreversible aggregation. *Journal of the American Chemical Society*, *143*(33), 13056–13064. https://doi.org/10.1021/jacs.1c03078

[3] Bowman, M. A., Riback, J. A., Rodriguez, A., Guo, H., Li, J., Sosnick, T. R., & Clark, P. L. (2020). Properties of protein unfolded states suggest broad selection for expanded conformational ensembles. *Proceedings of the National Academy of Sciences of the United States of America*, *117*(38), 23356–23364. https://doi.org/10.1073/pnas.2003773117

[4] Joubert, M. K., Hokom, M., Eakin, C., Zhou, L., Deshpande, M., Baker, M. P., Goletz, T. J., Kerwin, B. A., Chirmule, N., Narhi, L. O., & Jawa, V. (2012). Highly aggregated antibody therapeutics can enhance the in vitro innate and late-stage T-cell immune responses. *Journal of Biological Chemistry*, *287*(30), 25266–25279. https://doi.org/10.1074/jbc.m111.330902

[5] Center for Drug Evaluation and Research. (2014). *Immunogenicity assessment for therapeutic protein products*. U.S. Food and Drug Administration. Retrieved from https://www.fda.gov/regulatory-information/search-fda-guidance-documents/immunogenicity-assessment-therapeutic-protein-products

[6] Clark, P. L., Plaxco, K. W., & Sosnick, T. R. (2020). Water as a good solvent for unfolded proteins: Folding and collapse are fundamentally different. *Journal of Molecular Biology*, *432*(9), 2882–2889. https://doi.org/10.1016/j.jmb.2020.01.031

[7] Uversky, V. N., & Fink, A. L. (2002). The chicken-egg scenario of protein folding revisited. *FEBS Letters*, *515*(1-3), 79–83. https://doi.org/10.1016/s0014-5793(02)02441-9

[8] Millett, I. S., Doniach, S., & Plaxco, K. W. (2002). Toward a taxonomy of the denatured state: Small angle scattering studies of unfolded proteins. *Unfolded Proteins*, 241–262. https://doi.org/10.1016/s0065-3233(02)62009-1

[9] Shih, P.-M., Wang, I., Lee, Y.-T. C., Hsieh, S.-J., Chen, S.-Y., Wang, L.-W., Huang, C.-T., Chien, C.-T., Chang, C.-Y., & Hsu, S.-T. D. (2015). Random-coil behavior of chemically denatured topologically knotted proteins revealed by small-angle X-ray scattering. *The Journal of Physical Chemistry B*, *119*(17), 5437–5443. https://doi.org/10.1021/acs.jpcb.5b01984

[10] Zhou, L., Zhang, J., DiGiammarino, E., Kavishwar, A., Yan, B., Chumsae, C., Ihnat, P. M., Powers, D., Harlan, J., & Stine, W. B. (2018). Pulse SPR: A high throughput method to evaluate the domain stability of antibodies. *Analytical Chemistry*, *90*(20), 12221–12229. https://doi.org/10.1021/acs.analchem.8b03452

[11] Buchner, J., Renner, M., Lilie, H., Hinz, H. J., Jaenicke, R., Kiefhaber, T., & Rudolph, R. (1991). Alternatively folded states of an immunoglobulin. *Biochemistry*, *30*(28), 6922–6929. https://doi.org/10.1021/bi00242a016

[12]   Kats, M., Richberg, P., & Hughes, D. (1997). Ph-dependent isoform transitions of a monoclonal antibody monitored by micellar electrokinetic capillary chromatography. *Analytical Chemistry*, *69*(3), 338–343. https://doi.org/10.1021/ac9606517

[13]   Svergun, D. I., Koch, M. H., Timmins, P. A., & May, R. P. (2013). Basics of small angle scattering. *Small Angle X-Ray and Neutron Scattering from Solutions of Biological Macromolecules*, 13–26. https://doi.org/10.1093/acprof:oso/9780199639533.003.0002

[14]   Yageta, S., Imamura, H., Shibuya, R., & Honda, S. (2018). Ch2 domain orientation of human immunoglobulin G in solution: Structural comparison of glycosylated and aglycosylated FC regions using small-angle X-ray scattering. *MAbs*, *11*(3), 453–462. https://doi.org/10.1080/19420862.2018.1546086

[15]   Tian, X., Langkilde, A. E., Thorolfsson, M., Rasmussen, H. B., & Vestergaard, B. (2014). Small-angle X-ray scattering screening complements conventional biophysical analysis: Comparative structural and biophysical analysis of monoclonal antibodies igg1, IgG2, and igg4. *Journal of Pharmaceutical Sciences*, *103*(6), 1701–1710. https://doi.org/10.1002/jps.23964

[16]   Tian, X., Vestergaard, B., Thorolfsson, M., Yang, Z., Rasmussen, H. B., & Langkilde, A. E. (2015). In-depth analysis of subclass-specific conformational preferences of IGG antibodies. *IUCrJ*, *2*(1), 9–18. https://doi.org/10.1107/s205225251402209x

[17]   Mathew, E., Mirza, A., & Menhart, N. (2004). Liquid-chromatography-coupled saxs for accurate sizing of aggregating proteins. *Journal of Synchrotron Radiation*, *11*(4), 314–318. https://doi.org/10.1107/s0909049504014086

[18]   Brookes, E., Vachette, P., Rocco, M., & Pérez, J. (2016). *US-somo* HPLC-SAXS module: Dealing with capillary fouling and extraction of pure component patterns from poorly resolved SEC-Saxs Data. *Journal of Applied Crystallography*, *49*(5), 1827–1841. https://doi.org/10.1107/s1600576716011201

[19]   Graewert, M. A., Da Vela, S., Gräwert, T. W., Molodenskiy, D. S., Blanchet, C. E., Svergun, D. I., & Jeffries, C. M. (2020). Adding size exclusion chromatography (SEC) and light scattering (LS) devices to obtain high-quality small angle X-ray scattering (SAXS) data. *Crystals*, *10*(11), 975. https://doi.org/10.3390/cryst10110975

[20]   Hajizadeh, N. R., Franke, D., Jeffries, C. M., & Svergun, D. I. (2018). Consensus bayesian assessment of protein molecular mass from solution X-ray scattering data. *Scientific Reports*, *8*(1). https://doi.org/10.1038/s41598-018-25355-2

[21]   Guinier, A., & Fournet, G. (1955). *Small-angle scattering of X-rays (Oversat Fra Fransk)*. John Wiley.

[22]   Svergun, D. I., & Feĭgin, L. A. (1987). *Structure analysis by small-angle X-ray and neutron scattering*. Plenum Press.

[23]   Durand, D., Vivès, C., Cannella, D., Pérez, J., Pebay-Peyroula, E., Vachette, P., & Fieschi, F. (2010). NADPH oxidase activator p67phox behaves in solution as a multidomain protein with semi-flexible linkers. *Journal of Structural Biology*, *169*(1), 45–53. https://doi.org/10.1016/j.jsb.2009.08.009

[24]   Burger, V. M., Arenas, D. J., & Stultz, C. M. (2016). A structure-free method for quantifying conformational flexibility in proteins. *Scientific Reports*, *6*(1). https://doi.org/10.1038/srep29040

[25]   Grant, T. D. (2018). Ab initio electron density determination directly from solution scattering data. *Nature Methods*, *15*(3), 191–193. https://doi.org/10.1038/nmeth.4581

[26]   Brandt, J. P., Patapoff, T. W., & Aragon, S. R. (2010). Construction, MD simulation, and hydrodynamic validation of an all-atom model of a monoclonal IGG antibody. *Biophysical Journal*, *99*(3), 905–913. https://doi.org/10.1016/j.bpj.2010.05.003

[27]   Paul, R., Graff-Meyer, A., Stahlberg, H., Lauer, M. E., Rufer, A. C., Beck, H., Briguet, A., Schnaible, V., Buckel, T., & Boeckle, S. (2012). Structure and function of purified monoclonal antibody dimers induced by different stress conditions. *Pharmaceutical Research*, *29*(8), 2047–2059. https://doi.org/10.1007/s11095-012-0732-6

[28]   Erickson, H. P. (2009). Size and shape of protein molecules at the nanometer level determined by sedimentation, gel filtration, and electron microscopy. *Biological Procedures Online*, *11*(1), 32–51. https://doi.org/10.1007/s12575-009-9008-x

[29]   Harding, S. E., Horton, J. C., & Cölfen, H. (1997). The ellips suite of macromolecular conformation algorithms. *European Biophysics Journal*, *25*(5-6), 347–359. https://doi.org/10.1007/s002490050048

[30]   García de la Torre, J., & Harding, S. E. (2013). Hydrodynamic modelling of protein conformation in solution: ELLIPS and Hydro. *Biophysical Reviews*, *5*(2), 195–206. https://doi.org/10.1007/s12551-013-0102-6

[31]   Linderstrøm-Lang, K. (1924). *On the ionisation of proteins, by K. Linderström-Lang*. H. Hagerup.

[32]   Kay, M. S., & Baldwin, R. L. (1998). Alternative models for describing the acid unfolding of the apomyoglobin folding intermediate. *Biochemistry*, *37*(21), 7859–7868. https://doi.org/10.1021/bi9802061

[33]   Kataoka, M., Hagihara, Y., Mihara, K., & Goto, Y. (1993). Molten globule of cytochrome c studied by small angle X-ray scattering. *Journal of Molecular Biology*, *229*(3), 591–596. https://doi.org/10.1006/jmbi.1993.1064

[34]   Søndergaard, C. R., Olsson, M. H., Rostkowski, M., & Jensen, J. H. (2011). Improved treatment of ligands and coupling effects in empirical calculation and rationalization of Pka values. *Journal of Chemical Theory and Computation*, *7*(7), 2284–2295. https://doi.org/10.1021/ct200133y

[35]   Olsson, M. H., Søndergaard, C. R., Rostkowski, M., & Jensen, J. H. (2011). Propka3: Consistent treatment of internal and surface residues in empirical pka predictions. *Journal of Chemical Theory and Computation*, *7*(2), 525–537. https://doi.org/10.1021/ct100578z

[36]   Jurrus, E., Engel, D., Star, K., Monson, K., Brandi, J., Felberg, L. E., Brookes, D. H., Wilson, L., Chen, J., Liles, K., Chun, M., Li, P., Gohara, D. W., Dolinsky, T., Konecny, R., Koes, D. R., Nielsen, J. E., Head-Gordon, T., Geng, W., … Baker, N. A. (2017). Improvements to the apbs biomolecular solvation software suite. *Protein Science*, *27*(1), 112–128. https://doi.org/10.1002/pro.3280

[37]   Feige, M. J., Simpson, E. R., Herold, E. M., Bepperling, A., Heger, K., & Buchner, J. (2010). Dissecting the alternatively folded state of the Antibody Fab Fragment. *Journal of Molecular Biology*, *399*(5), 719–730. https://doi.org/10.1016/j.jmb.2010.04.032

[38]   Yageta, S., Lauer, T. M., Trout, B. L., & Honda, S. (2015). Conformational and colloidal stabilities of isolated constant domains of human immunoglobulin G and their

impact on antibody aggregation under acidic conditions. *Molecular Pharmaceutics*, *12*(5), 1443–1455. https://doi.org/10.1021/mp500759p

[39]   Thies, M. J. W., Kammermeier, R., Richter, K., & Buchner, J. (2001). The alternatively folded state of the antibody CH3 domain. *Journal of Molecular Biology*, *309*(5), 1077–1085. https://doi.org/10.1006/jmbi.2001.4707

[40]   Yageta, S., Shibuya, R., Imamura, H., & Honda, S. (2017). Conformational and colloidal stabilities of human immunoglobulin G FC and its cyclized variant: Independent and compensatory participation of domains in aggregation of multidomain proteins. *Molecular Pharmaceutics*, *14*(3), 699–711. https://doi.org/10.1021/acs.molpharmaceut.6b00983

[41]   Latypov, R. F., Hogan, S., Lau, H., Gadgil, H., & Liu, D. (2012). Elucidation of acid-induced unfolding and aggregation of human immunoglobulin igg1 and IGG2 FC. *Journal of Biological Chemistry*, *287*(2), 1381–1396. https://doi.org/10.1074/jbc.m111.297697

[42]   Martsev, S. P., Kravchuk, Z. I., Vlasov, A. P., & Lyakhnovich, G. V. (1995). Thermodynamic and functional characterization of a stable IGG conformer obtained by Renaturation from a partially structured low ph-induced state. *FEBS Letters*, *361*(2-3), 173–175. https://doi.org/10.1016/0014-5793(95)00145-y

[43]   Vlasov, A. P., Kravchuk, Z. I., Martsev, S. P. (1996) Non-native conformational states of immunoglobulins, thermodynamic and functional analysis of rabbit IgG. *Biokhimiia*, *61*, 212–235

[44]   Lilie, H., & Buchner, J. (1995). Domain interactions stabilize the *alternatively folded state* of an antibody fab fragment. *FEBS Letters*, *362*(1), 43–46. https://doi.org/10.1016/0014-5793(95)00203-l

[45]   Tischenko, V. M., Zav'yalov, V. P., Medgyesi, G. A., Potekhin, S. A., & Privalov, P. L. (1982). A thermodynamic study of cooperative structures in Rabbit Immunoglobulin G. *European Journal of Biochemistry*, *126*(3), 517–521. https://doi.org/10.1111/j.1432-1033.1982.tb06811.x

[46]   Welfle, K., Misselwitz, R., Hausdorf, G., Höhne, W., & Welfle, H. (1999). Conformation, ph-induced conformational changes, and thermal unfolding of anti-p24 (HIV-1) monoclonal antibody CB4-1 and its Fab and FC Fragments. *Biochimica Et Biophysica Acta (BBA) - Protein Structure and Molecular Enzymology*, *1431*(1), 120–131. https://doi.org/10.1016/s0167-4838(99)00046-1

[47]   Kanmert, D. (2011). *Structure and interactions of human Igg-Fc* (dissertation). Linköping University Electronic Press, Linköping.

[48]   Feige, M. J., Hendershot, L. M., & Buchner, J. (2010). How antibodies fold. *Trends in Biochemical Sciences*, *35*(4), 189–198. https://doi.org/10.1016/j.tibs.2009.11.005

[49]   Hora, M., Sarkar, R., Morris, V., Xue, K., Prade, E., Harding, E., Buchner, J., & Reif, B. (2017). Mak33 antibody light chain amyloid fibrils are similar to oligomeric precursors. *PLOS ONE*, *12*(7). https://doi.org/10.1371/journal.pone.0181799

[50]   Sedlák, E., Schaefer, J. V., Marek, J., Gimeson, P., & Plückthun, A. (2015). Advanced analyses of kinetic stabilities of iggs modified by mutations and glycosylation. *Protein Science*, *24*(7), 1100–1113. https://doi.org/10.1002/pro.2691

[51]   Miyafusa, T., Watanabe, H., & Honda, S. (2021). Local disorder of the C-terminal segment of the heavy chain as a common sign of stressed antibodies evidenced with a

peptide affinity probe specific to non-native IGG. *International Journal of Biological Macromolecules*, *182*, 1697–1703. https://doi.org/10.1016/j.ijbiomac.2021.05.137

[52]   Hebditch, M., & Warwicker, J. (2019). Web-based display of protein surface and ph-dependent properties for assessing the developability of Biotherapeutics. *Scientific Reports*, *9*(1). https://doi.org/10.1038/s41598-018-36950-8

[53]   Zhang, C., Codina, N., Tang, J., Yu, H., Chakroun, N., Kozielski, F., & Dalby, P. A. (2021). Comparison of the ph- and thermally-induced fluctuations of a therapeutic antibody fab fragment by Molecular Dynamics Simulation. *Computational and Structural Biotechnology Journal*, *19*, 2726–2741. https://doi.org/10.1016/j.csbj.2021.05.005

[54]   Nakamura, T., Aizawa, T., Kariya, R., Okada, S., Demura, M., Kawano, K., Makabe, K., & Kuwajima, K. (2013). Molecular mechanisms of the cytotoxicity of human α-lactalbumin made lethal to tumor cells (hamlet) and other protein-oleic acid complexes. *Journal of Biological Chemistry*, *288*(20), 14408–14416. https://doi.org/10.1074/jbc.m112.437889

[55]   Jasion, V. S., & Burnett, B. P. (2015). Survival and digestibility of orally-administered immunoglobulin preparations containing IGG through the gastrointestinal tract in humans. *Nutrition Journal*, *14*(1). https://doi.org/10.1186/s12937-015-0010-7

[56]   Onitsuka, M., Kadoya, Y., & Omasa, T. (2019). Secretory leakage of igg1 aggregates from recombinant Chinese hamster ovary cells. *Journal of Bioscience and Bioengineering*, *127*(6), 752–757. https://doi.org/10.1016/j.jbiosc.2018.11.015

[57]   Senga, Y., Doi, M., Onitsuka, M., & Honda, S. (2022). Live-cell imaging to analyze intracellular aggregation of recombinant IGG in Cho Cells. *Cell Chemical Biology*, *29*(1). https://doi.org/10.1016/j.chembiol.2021.08.010

[58]   Watanabe, H., Yageta, S., Imamura, H., & Honda, S. (2016). Biosensing probe for quality control monitoring of the structural integrity of therapeutic antibodies. *Analytical Chemistry*, *88*(20), 10095–10101. https://doi.org/10.1021/acs.analchem.6b02526

[59]   Filipe, V., Kükrer, B., Hawe, A., & Jiskoot, W. (2012). Transient molten globules and metastable aggregates induced by brief exposure of a monoclonal IGG to low ph. *Journal of Pharmaceutical Sciences*, *101*(7), 2327–2339. https://doi.org/10.1002/jps.23157

[60]   Oyetayo, O.-O., & Kiefer, H. (2016). Experimental model system to study ph shift-induced aggregation of monoclonal antibodies under controlled conditions. *Pharmaceutical Research*, *33*(6), 1359–1369. https://doi.org/10.1007/s11095-016-1878-4

[61]   Imamura, H., & Honda, S. (2016). Kinetics of antibody aggregation at neutral ph and ambient temperatures triggered by temporal exposure to acid. *The Journal of Physical Chemistry B*, *120*(36), 9581–9589. https://doi.org/10.1021/acs.jpcb.6b05473

[62]   Skamris, T., Tian, X., Thorolfsson, M., Karkov, H. S., Rasmussen, H. B., Langkilde, A. E., & Vestergaard, B. (2015). Monoclonal antibodies follow distinct aggregation pathways during production-relevant acidic incubation and neutralization. *Pharmaceutical Research*, *33*(3), 716–728. https://doi.org/10.1007/s11095-015-1821-0

[63]   Imamura, H., Sasaki, A., & Honda, S. (2017). Fate of a stressed therapeutic antibody tracked by fluorescence correlation spectroscopy: Folded monomers survive aggregation. *The Journal of Physical Chemistry B*, *121*(34), 8085–8093. https://doi.org/10.1021/acs.jpcb.7b05603

[64]   Imamura, H., & Honda, S. (2019). PH-shift stress on antibodies. *Methods in Enzymology*, 329–345. https://doi.org/10.1016/bs.mie.2019.02.021

[65]   Hebditch, M., Kean, R., & Warwicker, J. (2020). Modelling of ph-dependence to develop a strategy for stabilising mAbs at acidic steps in production. *Computational and Structural Biotechnology Journal*, *18*, 897–905. https://doi.org/10.1016/j.csbj.2020.03.002

[66]   Gasteiger, E., Hoogland, C., Gattiker, A., Duvaud, S., Wilkins, M. R., Appel, R. D., & Bairoch, A. (2005). Protein identification and analysis tools on the expasy server. *The Proteomics Protocols Handbook*, 571–607. https://doi.org/10.1385/1-59259-890-0:571

[67]   Shimizu, N., Mori, T., Nagatani, Y., Ohta, H., Saijo, S., Takagi, H., Takahashi, M., Yatabe, K., Kosuge, T., & Igarashi, N. (2019). BL-10C, the small-angle X-ray scattering beamline at the photon factory. *AIP Conference Proceedings*. https://doi.org/10.1063/1.5084672

[68]   Huang, T. C., Toraya, H., Blanton, T. N., & Wu, Y. (1993). X-ray powder diffraction analysis of Silver Behenate, a possible low-angle diffraction standard. *Journal of Applied Crystallography*, *26*(2), 180–184. https://doi.org/10.1107/s0021889892009762

[69]   Bernadó, P., Shimizu, N., Zaccai, G., Kamikubo, H., & Sugiyama, M. (2018). Solution scattering approaches to dynamical ordering in Biomolecular Systems. *Biochimica Et Biophysica Acta (BBA) - General Subjects*, *1862*(2), 253–274. https://doi.org/10.1016/j.bbagen.2017.10.015

[70]   Kline, S. R. (2006). Reduction and analysis of SANS and USANS data using Igor Pro. *Journal of Applied Crystallography*, *39*(6), 895–900. https://doi.org/10.1107/s0021889806035059

[71]   Schuck, P. (2000). Size-distribution analysis of macromolecules by sedimentation velocity ultracentrifugation and Lamm equation modeling. *Biophysical Journal*, *78*(3), 1606–1619. https://doi.org/10.1016/s0006-3495(00)76713-0