1   **Dynamic Kinetic Models Capture Cell-Free Metabolism for Improved**

2   **Butanol Production**

8   Jacob P. Martin[1,2,3], Blake J. Rasor[1,2,3], Jonathon DeBonis[1], Ashty S. Karim[1,2,3],

9   Michael C. Jewett[1,2,3], Keith E.J. Tyo[1,2,3], Linda J. Broadbelt[1,2]*

13   [1]Department of Chemical and Biological Engineering, Northwestern University,

14   Evanston, IL, USA 60208

15   [2]Center for Synthetic Biology, Northwestern University, Evanston, IL, USA 60208

16   [3]Chemistry of Life Processes Institute, Northwestern University, Evanston, IL, USA

17   60208

ABSTRACT

Cell-free systems are useful tools for prototyping metabolic pathways and

optimizing the production of various bioproducts. Mechanistically-based kinetic models

are uniquely suited to analyze dynamic experimental data collected from cell-free

systems and provide vital qualitative insight. However, to date, dynamic kinetic models

have not been applied with rigorous biological constraints or trained on adequate

experimental data to the degree that they would give high confidence in predictions and

broadly demonstrate the potential for widespread use of such kinetic models. In this

work, we construct a large-scale dynamic model of cell-free metabolism with the goal of

understanding and optimizing butanol production in a cell-free system. Using a novel

combination of parameterization methods, the resultant model captures experimental

metabolite measurements across two experimental conditions for nine metabolites at

timepoints between 0 and 24 hours. We present analysis of the model predictions,

provide recommendations for butanol optimization, and identify the aldehyde/alcohol

dehydrogenase as the primary bottleneck in butanol production. Sensitivity analysis

further reveals the extent to which various parameters are constrained, and our

approach for probing valid parameter ranges can be applied to other modeling efforts.

KEY WORDS

Metabolic Modeling, Kinetic Modeling, Ensemble Modeling, ODE Models, Dynamic

Models, Cell-Free Systems, Butanol Production, Parameter Optimization, Parameter

Estimation, Metabolic Control Analysis, Systems Biology, Computational Biology

1. INTRODUCTION

Widespread commercial success of metabolic engineering projects – from the production of biofuels (Atsumi et al., 2008; Fackler et al., 2021; Liew et al., 2022) to pharmaceuticals (Galanie et al., 2015; Ro et al., 2006) to polymer precursors (Arvay et al., 2021; Chung et al., 2015) – remains a challenge due to sub-optimal titers and yields (Biggs et al., 2021) and difficulty in optimizing biosynthetic pathways for production (Naseri & Koffas, 2020). This challenge is due to several factors, including the competition in living cells between non-native biosynthesis and cell viability (Dudley et al., 2015; Rasor et al., 2021; Wu et al., 2016), the highly connected nature of metabolic networks which obscures simple design choices for optimization (Tsiantis & Banga, 2020), and the difficulty of obtaining high-quality, kinetic data from living cells (Costa et al., 2016; Karim & Jewett, 2016). Cell-free systems addresses some of these issues, as cell viability no longer interferes with pathway flux, and time-course metabolomics are readily available without the need to extract intracellular species (Horvath et al., 2020; Karim et al., 2020; Miguez et al., 2019, 2021; Silverman et al., 2020; Vogeli et al., 2022). However, unlike purified systems, cell extract-based cell-free systems maintain much of the complexity of living metabolic networks (Bowie et al., 2020), which allows for native energy and cofactor regeneration to fuel biosynthesis (Jewett et al., 2008) but also preserves the complexity arising from network connectivity, making optimization nontrivial. While this complexity might seem problematic, it provides a means of collecting large data sets to understand and unravel metabolic complexity (Bowie et al., 2020; Bujara et al., 2011; Dudley et al., 2015, 2020).

Various computational models have been used to explain and analyze complex metabolic phenomena that would be difficult if not impossible to understand by manual inspection alone. For example, constraint-based models (CBMs), such as flux balance analysis (FBA)(Orth, Thiele, et al., 2010) and its many extensions (Jenior et al., 2020;

Lewis et al., 2010; Sánchez et al., 2017), have been used to simulate and explain

cellular behaviors. However, CBMs can only simulate steady-state solutions with limited

exceptions (Mahadevan et al., 2002), must assume cell-growth or other reasonable

cellular objectives for optimization, and do not explicitly represent metabolites or enzyme

abundances necessary to capture effects of enzyme saturation and allosteric regulation.

Kinetic models, which use a system of ordinary differential equations (ODEs)

where metabolites are tracked through time by explicitly formulated reaction rates, can

overcome these limitations (Strutz et al., 2019; Suthers et al., 2021). The primary

challenge in creating kinetic models is finding parameter values for each reaction rate,

as there are many more model parameters than experimental data points and

parameters are often unobservable, making the calculation a single "correct" or "best"

set of parameters impossible. To address this problem, an ensemble modeling (EM)

approach ca be applied to kinetic models, where several independently parameterized

models, or parameter sets, are used simultaneously as an ensemble of models (Tran et

al., 2008) to make predictions from the available data without overly relying on any

single set of parameters.

Many kinetic modeling formalisms exist which highly depend on the type of

system being modeled and the type and amount of data (Saa & Nielsen, 2017). For

example, when modeling living-cell metabolism at steady-state, kinetic models have

used "top-down" approaches for parameterization. First, known steady-state fluxes –

either experimentally measured (Tran et al., 2008) or predicted from tools like flux

balance analysis (J. Greene et al., 2019) – can be used as a "reference state" to lower

the degrees of freedom of parameterization, though this tends to produce solutions

which are more accurate nearer the reference state (J. L. Greene et al., 2017; St John et

al., 2019). Second, steady-state models can forgo simulations of absolute metabolite

concentrations in favor of tracking "relative" concentrations, again lowering the degrees

of freedom (Tran et al., 2008) (**Supplementary Methods S1**). Many methods are then

available to estimate the remaining parameters in steady-state models, including Monte

Carlo methods using random sampling (Tran et al., 2008), Bayesian estimation (Saa &

Nielsen, 2016; St John et al., 2019), local optimization (Gopalakrishnan et al., 2020), and

global parameter optimization (Khodayari et al., 2014).

In contrast, dynamic kinetic models, which can capture transient metabolism

found in non-steady state systems, typically require a "bottom-up" approach in which

model parameters are initially set at experimentally measured values if available (or at

reasonable estimates) and later optimized (Kim et al., 2018). These models have been

used for smaller-scale systems biology such as studying signaling or small isolated

biosynthetic pathways, where there are fewer model parameters and the primary hurdle

is network inference as opposed to parameter estimation (Min Lee et al., 2008). These

pathway-level models have frequently chosen to not explicitly model cofactors (Jia et al.,

2012; van Eunen et al., 2012), a choice which simplifies parameterization but limits the

scope of utility. Conversely, some dynamic metabolic models have included significant

portions metabolism but have "lumped" many reactions together into a single term or

reduced model structure (Buffing et al., 2018; Kurata & Sugimoto, 2018). While these

strategies make dynamic metabolic models more accessible, they limit the ability of the

model to fit more complex metabolic interactions, such as those found in unpurified cell-

free systems, and so more general methods were needed for this work. Recently, a

large-scale, high-resolution, and fully dynamic kinetic model of metabolism in an

Escherichia coli-derived cell-free system was developed using a literature-based

ensemble with Markov Chain Monte Carlo parameter optimization to study metabolic

impacts on cell-free protein synthesis (Horvath et al., 2020). While this work fit data from

a single experiment, this effort nonetheless demonstrated the feasibility of large-scale,

bottom-up parameterization of dynamic models in biological systems.

In this work, we aimed to develop a dynamic kinetic model of *E. coli*-based cell-

free metabolism for the study of biosynthetic capacity that can capture complex cell-free

metabolic behavior across multiple experimental conditions. To do this, we chose to

model the heterologous butanol production pathway, comprising five enzymatic steps to

catalyze the transformations from acetyl-CoA to butanol (Atsumi et al., 2008), which has

been successfully implemented in extract-based cell-free systems (Karim et al., 2019,

2020; Karim & Jewett, 2016). We chose this pathway in part because butanol is highly

reduced and requires several cofactors, and so was expected to be highly connected

with the larger cell-free metabolic network. Our model contains a large-scale

representation of core *E. coli* metabolism and the heterologous butanol pathway, and

model reactions are formulated with rate laws derived from mass-action kinetics of

elementary enzyme mechanisms. To parameterize these models, we combined a Monte

Carlo-based ensemble screening approach with a second step of local parameter

optimization, which were used in series to successfully fit measurements of butanol and

several species from various pathways of metabolism. Notably, both steps incorporated

thermodynamic and literature constraints that have not previously been used in the

bottom-up parameterization of large-scale dynamic models. The tools of metabolic

control analysis (MCA) were then used to give insight into the driving factors of cell-free

metabolism, as well as to give predictions regarding the optimization of butanol in this

cell-free system. We anticipate our model along with the cell-free platform will facilitate

both high-throughput prototyping of biosynthetic pathways for cellular design and cell-

free biomanufacturing.

2. MATERIALS AND METHODS

2.1. CELL-FREE METABOLIC EXPERIMENTS AND DATA

Metabolically active cell extracts were prepared according to (Grubbe et al.,

2020) from BL21-Star(DE3) *E. coli*. First, during cell-free protein synthesis (CFPS), each

enzyme in the butanol pathway was separately expressed in cell-free extracts and

quantified via [14]C-leucine incorporation (Rasor et al., 2022). Next, cell-free metabolic

engineering (CFME) experiments were conducted as described in (Karim et al., 2020;

Rasor et al., 2022), wherein each separately expressed butanol enzyme was added to

fresh cell-free extract, along with 120 mM glucose and several other cofactors, salts, and

other compounds. For the concentrations of each compound in the experiment present

from CFPS or added at the beginning of the CFME experiment, see **Supplementary**

Table S7.

Metabolite concentrations were measured by either high-performance liquid

chromatography (flowing 5 mM sulfuric acid at 0.6 mL/min on an Aminex Rezex™

ROA-Organic Acid H+ (8%) Column) or gas chromatography – mass spectrometry

(using N,O-Bis(trimethylsilyl)trifluoroacetamide derivatization and an Agilent HP-5MS

column with helium carrier gas). These measurements were taken every half hour from 0

hours (defined when glucose is first added) through 7.5 hours and triplicate

measurements were taken at 8 hours and 24 hours. Two experimental conditions were

tested: a butanol-positive (production) condition where all enzymes in the butanol

pathway were present, and a butanol-negative condition where the second enzyme in

the butanol pathway, 3-hydroxybutryryl-CoA dehydrogenase (HBD), was not added, thus

preventing butanol production. All metabolite measurements are given in

Supplementary Table S11.

2.2. NETWORK CONSTRUCTION

The metabolic network used to represent cell-free *E. coli* metabolism was

adapted from the model presented by Greene and coworkers (J. L. Greene et al., 2017),

which itself contained all metabolites and reactions in the BiGG *E. coli* core model

(Norsigian et al., 2020; Orth, Palsson, et al., 2010), as well as reactions described in a

previous model of *E. coli* metabolism (Khodayari et al., 2014). To simulate the non-

steady state cell-free environment, our model removed any reactions allowing the

transport of species between compartments as well as exchange reactions allowing

species to enter or leave the system. Duplicate reactions from isozymes were also

removed due to the lack of flux or literature data needed to distinguish each enzyme. All

enzyme-inhibitor pairs described in recent kinetic models of *E..coli* metabolism were

used (J. L. Greene et al., 2017; Horvath et al., 2020; Khodayari et al., 2014), as well as

E. coli enzyme inhibition described in BRENDA (Chang et al., 2021) or MetaCyc (Caspi

et al., 2014). All model regulation was either competitive or uncompetitive inhibition as

reported in the literature, and noncompetitive inhibition and allosteric inhibition and

activation were not considered due to lack of literature data. A full list of reactions and

inhibition is given in **Supplementary Table S6**. Lastly, the reactions for the engineered

pathway from acetyl-CoA to butanol (Atsumi et al., 2008) were included. The final model

contained 63 enzymatic reactions, 63 metabolites, and 100 inhibitor-enzyme pairs, and

is shown in **Figure 1**, as well as a more detailed representation in **Supplementary Fig.**

S1.

To construct a kinetic model, an approximate rate law was derived and applied to

each reaction. This equation to describe the rate of each enzymatic reaction expands

the random-order ternary complex enzyme mechanism and rate law described in

(Cornish-Bowden, 1979) with the additional assumption that the dissociation constant of

each substrate in multi-substrate reactions is unaffected by binding order. This

assumption halves the number of parameters per substrate compared to an order-

dependent rate form (Cornish-Bowden, 1979). The resulting rate law is mechanistically

grounded and is able to use experimental Michaelis constants as estimates for

dissociation constants, yet does not require manual curation to specific enzyme

mechanisms, is expandable to an arbitrary number of substrates, products, and

inhibitors, and reduces the hurdle of parameter estimation. In the case of two substrates

and two products:

$$A + B \overset{rxn\ j}{\rightleftharpoons} P + Q$$

205   The resultant net rate equation for reaction j ($v_{j,net}$), or the rates of the forward

206   and reverse elementary steps ($v_{j,f}$ and $v_{j,r}$), is given below:

$$v_{j,net} = v_{j,f} - v_{j,r} = \frac{V_{max_{j,f}}[A][B] - V_{max_{j,r}}[P][Q]}{1 + \frac{[A]}{K_{d_{j,A}}} + \frac{[B]}{K_{d_{j,B}}} + \frac{[A][B]}{K_{d_{j,A}}K_{d_{j,B}}} + \frac{[P]}{K_{d_{j,P}}} + \frac{[Q]}{K_{d_{j,Q}}} + \frac{[P][Q]}{K_{d_{j,P}}K_{d_{j,Q}}}}$$

The parameters $V_{max_{j,f}}$ and $V_{max_{j,f}}$ represent the forward and reverse maximal

rates, or rate constants, and $K_{d_{j,i}}$ represents the dissociation constant for species *i*. For

details, see **Supplementary Methods S2**.

A few unique metabolites and reactions were implemented for this model. First,

because the concentration of water is assumed to be constant, no dissociation constants

for water are used in any rate law, and instead is implicitly included in the rate constant.

Hydrogen ions are assumed to be buffered and are similarly not included in any rate

laws. Gas exchange for carbon dioxide and oxygen between the aqueous and gas

phases was included in the model, as CFME experiments were performed in a small

volume of liquid with a large headspace. Preliminary partial differential equation (PDEs)

modeling predicted spatial gradients of aqueous gas concentrations were negligible and

so were not further considered. Gas exchange at the surface of the liquid was assumed

to be quasi-equilibrated and was modeled kinetically using Henry's law constants as

constraints. See **Supplementary Methods S11** for more details.

2.3. MODEL PARAMETERIZATION

Due to the difficulty of parameter estimation for large-scale kinetic models and

the model equations being inherently underdetermined by the data, this work employed

an ensemble modeling approach in which many models, known as an "ensemble" of

models, each with the same structure but with independent parameter sets, are used

simultaneously. This approach not only alleviates the difficulty of finding a single set of

parameters that accurately describes the experimental data, but also acknowledges the

inability to determine a single optimal parameter value, known as parameter identifiability

or observability, that is a hallmark of large-scale kinetic models and provides a metric by

which uncertainty in both model parameter values and model behavior can be quantified.

Initial values, or parameter priors, for model dissociation constants were set to

experimentally measured Michaelis constants from BRENDA (Chang et al., 2021) and

MetaCyc (Caspi et al., 2014) when available. Of the 319 dissociation constants in the

model, 116 of these enzyme-substrate pairs had at least one Michaelis constant

available in these databases. When multiple experimental values were available, the

geometric mean was used. Where literature values were not available, the

corresponding dissociation constant was set to 0.1 mM, which was the median value of

all measured Michaelis constants (see **Supplementary Table S9 and S10** for all initially

available literature values). Of the 100 inhibition constants in the model, 13 values were

available on either BRENDA or MetaCyc, and the remainder were initially set to 0.5 mM,

which was the median value of the available constants.

Because turnover numbers are typically reported as specific quantities relative to

enzyme levels, the use of literature values for rate constants would require accurate and

broad proteomics for the cell-free experimental system, which were not available.

Instead, rate constants in a previously published model of cell-free protein synthesis

from Horvath, *et al.* (Horvath et al., 2020), which contained many of the same reactions

in central *E. coli* metabolism, were used as parameter priors. For reactions used in this

work that were not in the Horvath model, the initial parameter prior was set to the

geometric mean of all rate constants in the Horvath model. The direct use of the Horvath

rate constants in our model gave very poor fit to our data, as the data fit by the Horvath

model qualitatively differed from this study due to various experimental differences.

Nonetheless, the Horvath parameter values provided an initial point from which

parameters in this study could be further tuned.

Once priors for all parameters were obtained (**Figure 2a**), a single parameter set

was generated by sampling all model parameters within one order of magnitude in each

direction of each prior. Within each reaction, these parameters were further constrained

by the Haldane relationship, which provided physiological feasibility by ensuring

consistency with known thermodynamics. This equation related the equilibrium constant

in reaction j ($K_{eq,j}$) with the forward and reverse rate constants ($V_{max,f}$, $V_{max,r}$) and the

dissociation constants for each substrate or product i in reaction j ($K_{d_{j,i}}$) along with their

stoichiometry ($S_{i,j}$) and is given below in Eq. 1 (see **Supplementary Methods S3** for

details). Equilibrium constants were calculated using eQuilibrator 3.0 (Beber et al.,

2022).

$$K_{eq,j} = \frac{V_{max,f}}{V_{max,r}} \prod_{i \in \{substrates, products\}} K_{d_{j,i}}^{S_{i,j}}$$

264   **[1]**

This procedure was repeated to generate an ensemble of 10^7 independent

parameter sets (**Figure 2b**). Each parameter set was simulated by integrating the

system of ordinary differential equations (ODEs) which describe the change of each

metabolite concentration over time (**Figure 2c**) and were solved in MATLAB 2020b (The

MathWorks, Natick, MA) (**Supplementary Methods S7**). Initial ODE metabolite

concentrations were set, in order of priority, first by available measurements, then by

experimentally added concentrations, and finally by known *E. coli* intracellular

concentrations as measured by (Bennett et al., 2009), corrected for dilution. Any

metabolite not in any of these categories was set to 0 mM. Initial concentrations of gas

species were set to atmospheric conditions, and aqueous gases were set to equilibrium

concentrations via Henry's Law. See **Supplementary Table S7** for values for each

model metabolite.

The performance of each parameter set was quantified by the weighted root

mean squared errors (RMSE) of the simulated metabolite profiles to experimental

metabolite measurements across both experimental conditions. The RMSE values for

succinate were weighted 6-fold compared to other metabolites due to poor fit to

succinate measurements in early models, and the RMSE for butanol were weighted 20-

fold because butanol production was emphasized in this model (**Supplementary**

Methods S8). The ensemble was "screened" by keeping only the 100 parameter sets

with the lowest weighted RMSE, which were used for further optimization and analysis.

Because models generated from the initial parameter priors failed to adequately

fit the experimental data, sensitivity analysis was performed to adjust parameter priors.

The finite differences method was used to calculate the sensitivity of model fitness with

respect to each parameter value (**Supplementary Methods S6**) and was applied to all

models in the final screened ensemble. The priors for any parameters that were highly

sensitive across most models in the ensemble were then adjusted in the direction of

improved fitness to the data (**Figure 2d**).

After iterating this procedure to update parameter priors, the fit of all models to

the data remained poor. To remedy this, we applied a local parameter optimization to

each of the top 200 models in the screened ensemble (**Figure 2e**). In this procedure,

each screened model was used as the initial point in the MATLAB pattern search

algorithm, which attempts to find a local optimum fitness value by using heuristics to

traverse the parameter space. This algorithm showed better performance than both

particle swarm and genetic algorithm global search methods as well as simpler local

search methods, such as MATLAB's "fmincon". Individual parameter constraints were

applied to ensure rate constants, dissociation constants, and inhibition constants stayed

in physiologically realistic ranges, and parameters were log-transformed to allow the

Haldane relationship to be applied as a set of linear inequality constraints within the

optimization algorithm (see **Supplementary Methods S4** for details on optimization

constraints).

Local parameter optimization was carried out on MATLAB 2020b on the

Northwestern Quest High Performance Computing Cluster. Each of the 200 optimization

runs employed parallelization within the "pattern search" toolbox and ran on 250 CPUs

with a wall time of 4 hours. After optimization, each parameter set was re-ranked

according to the weighted RMSE fitness of the optimized point. The top 20 optimized

models were chosen for the final ensemble and further analysis, as these models

showed the strongest agreement to the experimental timecourse metabolite

measurements in addition to capturing a distinct shift in acetate and succinate

production in the butanol-negative condition.

2.4. METABOLIC CONTROL ANALYSIS

Metabolic control analysis (MCA) is a means of measuring how network features,

such as metabolite concentrations, reaction fluxes, and model parameters, impact each

other. This analysis can be either local, such as concentration elasticities (ϵ_{ji}^{x}), which

describe the change in reaction flux caused by those concentrations or parameters

within that reaction, or they can be global, such as flux control coefficients (C_{ij}^{J}),

concentration control coefficients (C_{ij}^{x}), and the Jacobian ($J_{ii'}$), which describe how

changes in each concentration or flux affect distant network features through shared

metabolites. The definitions of each normalized value, in terms of metabolite

concentrations x_i and $x_{i'}$ and reaction fluxes v_j and $v_{j'}$, as given in Equations 2-5 below:

$$\epsilon_{ji}^{x} = \frac{x_i}{v_j} \frac{\partial v_j}{\partial x_i}$$

324 [2]

$$C_{ij}^{x} = \frac{v_j}{x_i} \frac{\partial x_i}{\partial v_j}$$

325 [3]

$$C_{jj'}^{v} = \frac{v_{j'}}{v_j} \frac{\partial v_j}{\partial v_{j'}}$$

326 [4]

$$J_{ii'} = \frac{x_{i'}}{x_i} \frac{\partial}{\partial x_{i'}} \left(\frac{\partial x_i}{\partial t} \right)$$

[5]

The above MCA features were calculated analytically, as described in (Hofmeyr,

2001) and detailed in **Supplementary Methods S5**. Because the system in this model is

dynamic, these values were calculated at discrete timepoints for every 0.5 hours

between 0 and 24 hours in the model simulation. In all analyses presented in this work,

only values calculated between 0.5 hours and 7.5 hours are presented, since values

calculated at 0 hours caused numerical instability from initial concentrations of zero, and

values after 7.5 hours showed wide variation due to the lack of experimental data

between 8 and 20 hours to constrain model behavior.

3. RESULTS

3.1. DYNAMIC MODEL FITTING AND GOODNESS OF FIT

To train our dynamic model of cell-free metabolism, we iterated the process of

sampling around parameter priors to generate an ensemble, screening the ensemble

against the experimental data, and updating the parameter priors using parameter

sensitivity analysis fifteen times until the model adequately fit the metabolite data (see

Supplementary Table S9 for each iteration of parameter priors). From this iteration, the

top 200 screened parameter sets (of 10^8 sampled) showed significantly improved fit to

the nine measured metabolites in the butanol-positive condition, with a 32% reduction in

the median weighted RMSE compared to screened parameter sets from early iterations

(Supplementary Table S1). However, the experimental data from the butanol-negative

condition showed a large increase in acetate accumulation and a decrease in succinate

accumulation as compared to the butanol-positive condition, which were not captured by

any of the top 200 parameter sets **(Supplementary Fig. S3)**. These shifts in metabolism

were difficult to capture in part because the measured increase in acetate could not

simply be accounted for from the measured decrease in butanol. Additionally, previous

experimental data showed that changes in acetate and succinate titers – as well as

ethanol and lactate, to a lesser extent – were correlated with changes in butanol titers

(**Supplementary Fig. S2**), which implied that system-level effects were coupling the

production of all these metabolites.

We next employed automated parameter optimization algorithms to fine-tune

each parameter set and found that the local optimization "pattern search" algorithm (The

Mathworks, Inc.) significantly improved the weighted RMSE of nearly all models,

reducing the weighted RMSE a further 46% **(Supplementary Table S1)**. We found that

the top 20 parameter sets (in terms of post-optimization weighted RMSE) captured the

experimentally observed shifts in acetate and succinate as well as all other

experimentally measured metabolites **(Figure 3)**. Importantly, the combination of

ensemble screening and local parameter optimization was found to be necessary to

successfully fit both positive and negative butanol production conditions, as each

method alone was not sufficient to fit the data, even when sampling around the final

updated parameter priors (**Supplementary Fig. S4**).

3.2. ANALYSIS OF CELL-FREE METABOLISM AND THE BUTANOL PATHWAY

With a trained ensemble of 20 parameter sets, we wanted to understand how simulated metabolism was able to fit experimental metabolite measurements. The most interesting behaviors of this data were the shifts in acetate and succinate concentrations in the butanol-negative condition compared to the butanol-positive condition. Our models suggest that acetate accumulation in the butanol-negative condition is due to increased acetyl-CoA concentration (**Supplementary Fig. S5**), which itself is caused by the lack of acetyl-CoA flux into the butanol pathway; these conclusions are supported by the high control of acetyl-CoA concentration by butanol concentration in MCA, as well as pathway flux analysis (**Supplementary Fig. S6**; **Supplementary Methods S9**). Next, by decomposing the reactions for succinate production into "elementary" (forward and reverse) fluxes, we observed that the decrease in succinate production in the butanol-negative condition was not from lowered "forward" reaction fluxes but instead from increased "reverse" flux of succinyl-CoA synthase (in the direction of succinyl-CoA from succinate) (**Supplementary Fig. S7a**). This reverse flux – equivalent to product inhibition – was attributable to an increase in free Coenzyme A (CoA) (**Supplementary Fig. S7b**) and was confirmed with MCA showing strong negative control over this reaction by CoA (**Supplementary Fig. S7c**). The increase in CoA, in turn, is caused by the increase in phosphotransacetylase (PTA) flux in the acetate pathway from the butanol pathway knockout, which can be seen both in the high concentration control coefficient of CoA from PTA (**Supplementary Fig. S7c**) and the reactions that contribute to changes in acetyl-CoA concentration (**Supplementary Fig. S6b**).

Significantly, the increases in acetyl-CoA and CoA in the butanol-negative condition – which are responsible for the increase in acetate and decrease in succinate, respectively – are not trivial consequences of the butanol pathway knockout in the negative condition, as the butanol pathway uses acetyl-CoA and releases CoA in the

same stoichiometric ratios as the acetate pathway. Instead, our model analysis found

that these shifts are only possible due to a bottleneck in the butanol pathway wherein

intermediates were accumulated, causing butanol pathway flux to have a net decrease

in the quasi-steady state concentrations of both acetyl-CoA and CoA, as compared to

acetate pathway flux (**Supplementary Fig. S8a**). This prediction of butanol pathway

intermediate accumulation was conserved among all 20 parameter sets and will be

discussed further in the context of predictions for pathway optimization.

Beyond acetate and succinate shifts, carbon metabolism within the model

remains a highly connected, balanced network. Strong control of butanol production is

maintained among all models in the final ensemble, with consistent control by

acetaldehyde dehydrogenase (ACALD), NAD dehydrogenase (NADH16pp), malate

dehydrogenase (MDH), and NAD-dependent malic enzyme (ME1) (**Supplementary Fig.**

S8b). However, this control varies in direction (i.e., some models have parameters

where this control is positive and others where the control is negative) suggesting

several possible control mechanisms. Ethanol production is even more tightly controlled,

which allows the model to fit experimental observations of unchanged ethanol production

between conditions despite changing acetyl-CoA concentration. MCA shows that this

effect is due to saturation of the first step of ethanol production (ACALD) by acetyl-CoA,

as well as further distributed control over ethanol production across ACALD,

glyceraldehyde-3-phosphate dehydrogenase (GAPD), fructose bisphosphatase (FBP),

pyruvate dehydrogenase (PDH), and NADH16pp fluxes (**Supplementary Fig. S12**). In

contrast, the production of lactate is not strongly controlled by other reactions, as can be

seen in the very low flux control coefficients for lactate dehydrogenase (LDH)

(**Supplementary Fig. S9a**), but is instead more strongly controlled by NADH than by

pyruvate or other metabolites (**Supplementary Fig. S9b**). This finding suggesting that

lactate production stops at the same time as glycolytic flux not because of lack of

pyruvate, but instead due to lack of NADH regeneration in glycolysis, which thereby lowers the NADH:NAD$^+$ ratio and brings LDH flux into equilibrium. Lastly, the consumption of glucose itself shows a large amount of metabolic control from NADH16pp, GAPD, TPI, FBP, PTA, and HEX fluxes (**Supplementary Fig. S10**), which are important reactions in the control of the other measured metabolites discussed above. GAPD and TPI are both interesting cases, as they are thermodynamic bottlenecks with large positive Gibbs free energies (Beber et al., 2022). While some previous literature has suggested that glycolysis in *E. coli* is predominately controlled by the rate of ATP consumed outside of glycolysis itself (Solem et al., 2003), other work has identified GAPD as a key controlling and limiting step of glycolysis, both in *E. coli* (Centeno-Leija et al., 2013; Cho et al., 2012) and in other organisms (Shestov et al., 2014). In the context of metabolism in the cell-free system described here, this finding would indicate that GAPD flux, and the NADH generated by this reaction, is the dominant factor controlling glycolytic flux.

3.3. Model Predictions for Optimization of Butanol Production

Kinetic models offer advantages in their ability to extrapolate and make predictions about unseen experimental conditions (J. L. Greene et al., 2017; Strutz et al., 2019). Here, we simulate several new experimental conditions (e.g., altering butanol pathway enzyme concentrations and the initial concentrations of metabolites) that aim to increase butanol production. In the cell-free reaction, the concentrations of heterologous enzymes are easily perturbed while the enzymes in core metabolism are fixed. We therefore first used our model to adjust the levels of all five butanol pathway enzymes at 0.5x, 2x, 5x, and 10x fold-changes from the baseline butanol-positive condition (**Figure 4**). Within the model, these conditions were simulated by adjusting the forward and reverse rate constant parameters (V_{max}) of the corresponding reactions. When changing

all enzymes simultaneously, we found that final 20-hour butanol titers were predicted to

increase monotonically with increasing butanol pathway enzyme levels but all models

show diminishing returns at the highest levels of enzyme overexpression, implying that

heterologous pathway enzymes become no longer limiting. When increasing each

butanol pathway enzyme individually, all models in the final ensemble predicted that

increasing the bi-functional aldehyde-alcohol dehydrogenase, which carries out the final

two reactions in the butanol pathway, ADHE1 and ADHE2, gave a much larger

improvement in butanol titer than increasing any other butanol pathway enzyme (**Figure**

4).

As previously mentioned, all parameter sets predicted that a bottleneck to

accumulate intermediates in the butanol pathway was needed to fit the observed shifts in

acetate and succinate production. Interestingly, despite the lack of enzyme-specific

training data, all parameter sets further agreed that this bottleneck would accumulate

butyryl-CoA and butyraldehyde, the substrates to ADHE1 and ADHE2, respectively

(**Supplementary Fig. S8a**). To understand this model behavior, we performed a

thermodynamic analysis of the butanol pathway. This analysis showed that the first step

of the pathway, THL ($\Delta G'^m$ = +23 kJ/mol), and the second step, HBD ($\Delta G'^m$ = -20

kJ/mol), must be kinetically fast and therefore near equilibrium to establish flux into the

butanol pathway. Similarly, the third step of the pathway, CRT, is only slightly exergonic

($\Delta G'^m$ = -1.0 kJ/mol), and so the first three steps of the pathway are at quasi-equilibrium,

and the fourth step (TER) is again quite favorable ($\Delta G'^m$ = -61 kJ/mol). The quasi-

equilibrium of the first three reactions means that any bottleneck in these reactions or

the subsequent fourth reaction would not result in accumulation of intermediates but

would equilibrate back to acetyl-CoA, preventing the shifts seen in the data. Therefore,

bottlenecks responsible for accumulation were only possible in the last two steps of

ADHE1 and ADHE2 ($\Delta G'^m$ = -10 kJ/mol and $\Delta G'^m$ = -25 kJ/mol, respectively) once flux

was effectively irreversible. For this reason, the alcohol-aldehyde dehydrogenase

represents the primary enzyme bottleneck in the butanol pathway, and its

overexpression is consistently predicted to have the only significant impact on increased

butanol production, which is validated by previous experimental observations (Karim &

Jewett, 2016). These results provide valuable knowledge that can be leveraged

experimentally when expression and solubility of the enzyme are not limiting.

We next used the models to predict the effect of changing the initial cofactor and

substrate concentrations. First, titrations were simulated for the initial concentration of

acetyl-CoA, CoA, ATP, ADP, AMP, NADH, NAD$^+$, NADPH, NADP$^+$, pyruvate,

phosphoenolpyruvate, inorganic phosphate, and pyrophosphate (**Supplementary Fig.**

S11). While parameter sets did not have the same consensus as for enzyme level

predictions, the simulated initial concentrations of 50 mM NADH (compared to 0.1 mM in

the base condition of butanol-positive condition), 50 mM ATP (0.1 mM in the base

condition), 15 mM CoA (4.4 mM in the base condition), and 50 mM pyrophosphate (0

mM in the base condition) all predicted increases in butanol titer to varying degrees. We

then simulated butanol production with these new combinatorial conditions which

predicted the highest median butanol production (**Figure 5**). These cofactors have

previously been found experimentally to be the most important additions for controlling

cell-free metabolism (Karim et al., 2018). For high initial NADH, butanol titer is increased

both by competitively inhibiting PTA and providing reduction potential to relieve the

ADHE1 and ADHE2 bottleneck. The increase in CoA increases butanol titers primarily

by increasing pyruvate flux to acetyl-CoA via pyruvate dehydrogenase and indirectly

increasing NADH through increased flux in AKGDH. High pyrophosphate was predicted

to decrease acetyl-CoA synthase flux and minimize futile cycling. Lastly, high initial ATP

was predicted to increase butanol production by inhibiting PTA and LDH flux

(**Supplementary Fig. S11b**), though it should be noted that this trend for ATP in

particular was not observed in previous experimental studies of related cell-free systems

(Karim et al., 2018; Karim & Jewett, 2016). Next, we varied initial concentrations of

glucose, acetate, and glutamate (**Figure 5**). While glucose was seen to seen to have the

largest effect on final butanol titer, the correlation was not linear. For example, a 62%

decrease in initial glucose resulted in a 34% decrease in final butanol titer in the median

across parameter sets, with one parameter set even showing an increase in butanol titer

from this decreased glucose.

3.4. ANALYSIS OF PARAMETER OPTIMIZATION AND CONSERVATION OF PARAMETER

VALUES

Virtually all large-scale kinetic models are vastly underdetermined and individual

parameters are not identifiable, thus limiting the interpretability of individual parameter

values. Because we used an ensemble approach, we have an opportunity to study how

parameter values and network features are constrained – or unconstrained – as a result

of undertaking this type of model fitting. For a given parameter, the conserved prediction

of similar parameter values across the ensemble may support several conclusions,

including increased confidence in a true physiological parameter value, increased

structural identifiability due to network structure or reaction mechanisms, or increased

practical identifiability due to proximity to observed data. In all cases, these are important

factors in model building and analysis, and these results may help increase

understanding in these areas.

Here, we measure how tightly or loosely constrained individual parameter values

are between the parameter sets in the final ensemble to understand how the data might

constrain certain parameter values, or conversely, how different parameter values might

give similar model behaviors. However, because individual parameter values are highly

context-dependent, and the effect of a parameter on model responses cannot be

determined from the absolute parameter value alone, it is instead preferrable to use

proxy values for each parameter which are more representative of network behavior.

We first used as a proxy the metric of concentration elasticity, which relates the

response in reaction flux to the value of each dissociation constant parameter and

normalizes all parameters for better comparison. This metric further allows analysis of

the degree to which parameters control various reactions, as elasticity values near 0

signify saturated substrates, elasticities near 1 signify substrate concentrations far below

saturation in a first-order regime, and intermediate elasticities represent substrates with

more control over reaction rates. After performing this analysis on the ensemble of 20

models, we saw that there were several parameters whose corresponding elasticity

values were highly conserved. This result in summarized in **Figure 5**, which plots the

mean value of each elasticity against the standard deviation across the 20 models in the

final ensemble. Further discussion of specific examples of conserved elasticities is given

in **Supplementary Table S2**, and the full presentation of elasticities for all reactions

across all models are shown in **Supplementary Table S4** and **Supplementary Fig.**

S12.

Next, we used as a proxy the metric of parameter sensitivity with respect to

NADH oxidative flux (**Supplementary Methods S10**). Because this value is

physiologically relevant and yet was not directly constrained or favored by the model

fitting process, this provides an additional unbiased metric to analyze the extent to which

parameter values are constrained. Again, there were several parameters whose

sensitivity to this behavior were relatively conserved, which is summarized in

Supplementary Table S3, with all values given in **Supplementary Table S5** and shown

in **Supplementary Fig. S13**.

4. DISCUSSION

4.1. SIGNIFICANCE OF WORK

In this work, we successfully fit a large-scale dynamic model to multiple

conditions of experimental time course metabolite measurements. The resulting model

captured complex, unintuitive interactions between various branches of metabolism and

was able to further provide predictions as to the metabolic phenomena underlying these

effects. The ensemble approach was leveraged to provide quantitative measurements of

the uncertainty of these predictions; somewhat surprisingly, the predictions around both

the metabolic behaviors in the training data, as well as the predicted effects of

hypothetical enzyme level changes, were much more conserved within the ensemble

than initially expected, demonstrating high certainty in these predictions. The final model

produced several other optimization strategies with varying degrees of ensemble

uncertainty that may be used in future work to increase butanol production and further

retrain this model.

4.2. ADVANTAGES AND LIMITATIONS

The most significant advantage of the methodology of thiswork was the use of

the ensemble approach. First, this approach allows for a thorough sampling of

parameter space, which was seen to be necessary to provide adequate initial points for

subsequent optimization (**Supplementary Fig. S4**). Second, it allows quantification of

the uncertainty in both model parameters and model predictions, such as the increased

confidence in the predicted effect of butanol enzyme level changes. Third, the ensemble

approach allows for uncertain predictions to be hedged, such that even predictions

which vary between models can provide utility. For example, lower-confidence

predictions can be tested to improve butanol production, and in the worst-case scenario

will still provide valuable training data to refine the ensemble. This ease of re-

optimization by local optimization is also a significant advantage of this work which can

increase the speed of model-building in the design-build-test-learn cycle.

Another advantage of this work is the scalability and utility of additional

experimental data. The above analysis found that the most highly constrained

parameters were in reactions near experimentally measured metabolites, which implies

that currently under-constrained parameters can be fine-tuned simply by obtaining more

experimental measurements. The resulting tightening of parameter constraints should

give more accurate and conserved predictions across models. Simultaneously, a well-fit

model was obtained in this work despite measurements from relatively few metabolites,

so while additional measurements may have benefits, they should not be seen as a

reason not to undertake similar model-fitting efforts.

One primary limitation in this work, common to many large-scale models, is the

manual process of model selection. While there have been early efforts to use

automated model reconstruction tools on kinetic models (van Rosmalen et al., 2021),

their widespread use remains limited to constraint-based models with steady-state data

(Mendoza et al., 2019), and while parameter identifiability tools have seen great

improvements, they still scale poorly for larger models, especially those with complex

rate forms (Villaverde et al., 2019). Therefore, model selection remains an iterative

process which must balance reducing the number of parameters with leaving enough

reactions to allow the model sufficient flexibility to fit the data. For the "bottom-up"

parameterization in this work, the choice of reactions was further complicated by the

choice of parameter priors, especially where data could be fit in multiple ways. For

example, our model included both hexokinase (HEX) and the glucose

phosphotransferase system (PTS) as reactions to consume glucose. While we chose

higher rate constants for HEX, as we believed it unlikely that the multi-step, membrane-

bound PTS was highly active in cell-free extracts, a different choice may have led to a

different final model. Similarly, model predictions of succinate production from glutamate

versus from TCA flux may have been biased by parameter priors. Despite this potential

uncertainty, in this work we saw that these two decisions had minimal impact: in the first

case, the primary reaction for glucose consumption did not significantly affect model

predictions surrounding butanol optimization (**Supplementary Fib. S14**), and in the

second case, nearly all parameter sets, once optimized, utilized glutamate as the

primary source of succinate flux. Regardless, this fortunate result is not guaranteed for

future efforts, and so careful consideration of model selection, along with the potential

use of automated model construction or parameter identifiability methods, model

reduction methods (Strutz et al., 2019) and increasingly approximate rate forms (Du et

al., 2016; St John et al., 2019) should all be considered.

In conclusion, while there are certainly improvements to be made in future

modeling efforts, this work demonstrates the successful parameterization of a large-

scale dynamic kinetic model to capture complex metabolic data. The resulting model has

increased realism and confidence due to improved literature value parameter priors and

rigorously applied thermodynamic constraints. This model was used to gain new causal

understanding of metabolism in cell-free systems and how this metabolism interacts with

the engineered butanol pathway. This understanding was further leveraged to provide

strategies to optimize butanol titers, including the prediction that the alcohol/aldehyde

enzyme was the primary bottleneck of the butanol pathway, which itself was unique and

unanticipated due to its high conservation across models despite the lack of direct

training data. Lastly, the final trained model was used for a large-scale analysis of the

degree to which individual parameter values are constrained during model fitting, which

has broad implications in kinetic models in general. We hope that the success,

applicability, and general ease of use of these methods and results will inspire additional

experiments to measure dynamic behaviors around engineered pathways in metabolism

for the purpose of continued model building and improved metabolic understanding.

CONTRIBUTION STATEMENT

Jacob Martin: Conceptualization, Methodology, Software, Formal analysis, Writing –

Original Draft.

Blake J. Rasor: Data Collection, Writing – review & editing.

Jon DeBonis: Methodology, Software, Writing – review & editing.

Ashty Karim: Data Collection, Writing – review & editing.

Michael Jewett: Supervision, Funding acquisition, Writing – review & editing.

Keith Tyo: Conceptualization, Supervision, Writing – review & editing.

Linda Broadbelt: Conceptualization, Supervision, Writing – review & editing.

COMPETING INTERESTS

The authors claim no competing interests.

ACKNOWLEDGEMENTS

We would like to thank Prof. Niall Mangan, Prof. Matthew Plumlee, and Prof.

Andreas Waechter for project guidance and helpful discussions. Thanks to Jon Strutz

and Kevin Shebek for helpful discussions and assistance with software and eQuilibrator

3.0.

This research was supported in part through the computational resources and

staff contributions provided for the Quest high performance computing facility at

Northwestern University which is jointly supported by the Office of the Provost, the Office

for Research, and Northwestern University Information Technology. This work was

funded by the U.S. Department of Energy Office of Science, Biological and

Environmental Research Division (BER), Genomic Science Program (GSP) under

Contract No. DE-SC0018249. M.C.J. gratefully acknowledges the David and Lucile

Packard Foundation and the Camille Dreyfus Teacher–Scholar Program. J.P.M. was

supported by a National Institutes of Health Training Grant (T32GM008449) through

Northwestern University's Biotechnology Training Program. B.J.R. was supported by a

REFERENCES

Arvay, E., Biggs, B. W., Guerrero, L., Jiang, V., & Tyo, K. (2021). Engineering

Acinetobacter baylyi ADP1 for mevalonate production from lignin-derived aromatic

compounds. *Metabolic Engineering Communications*, *13*(May 2020), e00173.

https://doi.org/10.1016/j.mec.2021.e00173

Atsumi, S., Cann, A. F., Connor, M. R., Shen, C. R., Smith, K. M., Brynildsen, M. P.,

Chou, K. J. Y., Hanai, T., & Liao, J. C. (2008). Metabolic engineering of Escherichia

coli for 1-butanol production. *Metabolic Engineering*, *10*(6), 305–311.

https://doi.org/10.1016/j.ymben.2007.08.003

Beber, M. E., Gollub, M. G., Mozaffari, D., Shebek, K. M., Flamholz, A. I., Milo, R., &

Noor, E. (2022). eQuilibrator 3.0: a database solution for thermodynamic constant

estimation. *Nucleic Acids Research*, *50*(D1), D603–D609.

https://doi.org/10.1093/nar/gkab1106

Bennett, B., Kimball, E., Gao, M., Osterhout, R., Van Dien, S. J., & Rabinowitz, J. D.

(2009). Absolute metabolite concentrations and implied enzyme active site

occupancy in Escherichia coli. *Nature Chemical Biology*, *5*(8), 593–599.

https://doi.org/10.1038/nchembio.186.Absolute

Biggs, B. W., Alper, H. S., Pfleger, B. F., Tyo, K. E. J., Santos, C. N. S., Ajikumar, P. K.,

& Stephanopoulos, G. (2021). Enabling commercial success of industrial

biotechnology. *Science*, *374*(6575), 1563–1565.

https://doi.org/10.1126/science.abj5040

Bowie, J. U., Sherkhanov, S., Korman, T. P., Valliere, M. A., Opgenorth, P. H., & Liu, H.

(2020). Synthetic Biochemistry: The Bio-inspired Cell-Free Approach to Commodity

Chemical Production. *Trends in Biotechnology*, *38*(7), 766–778.

https://doi.org/10.1016/j.tibtech.2019.12.024

Buffing, M. F., Link, H., Christodoulou, D., & Sauer, U. (2018). Capacity for

instantaneous catabolism of preferred and non-preferred carbon sources in

Escherichia coli and Bacillus subtilis. *Scientific Reports*, *8*(1), 1–10.

https://doi.org/10.1038/s41598-018-30266-3

Bujara, M., Schümperli, M., Pellaux, R., Heinemann, M., & Panke, S. (2011).

Optimization of a blueprint for in vitro glycolysis by metabolic real-time analysis.

Nature Chemical Biology, *7*(5), 271–277. https://doi.org/10.1038/nchembio.541

Caspi, R., Altman, T., Billington, R., Dreher, K., Foerster, H., Fulcher, C. A., Holland, T.

A., Keseler, I. M., Kothari, A., Kubo, A., Krummenacker, M., Latendresse, M.,

Mueller, L. A., Ong, Q., Paley, S., Subhraveti, P., Weaver, D. S., Weerasinghe, D.,

Zhang, P., & Karp, P. D. (2014). The MetaCyc database of metabolic pathways and

enzymes and the BioCyc collection of Pathway/Genome Databases. *Nucleic Acids*

Research, *42*(D1), 459–471. https://doi.org/10.1093/nar/gkt1103

Centeno-Leija, S., Utrilla, J., Flores, N., Rodriguez, A., Gosset, G., & Martinez, A.

(2013). Metabolic and transcriptional response of Escherichia coli with a NADP +-

dependent glyceraldehyde 3-phosphate dehydrogenase from Streptococcus

mutans. *Antonie van Leeuwenhoek, International Journal of General and Molecular*

Microbiology, *104*(6), 913–924. https://doi.org/10.1007/s10482-013-0010-6

Chang, A., Jeske, L., Ulbrich, S., Hofmann, J., Koblitz, J., Schomburg, I., Neumann-

Schaal, M., Jahn, D., & Schomburg, D. (2021). BRENDA, the ELIXIR core data

resource in 2021: New developments and updates. *Nucleic Acids Research*,

49(D1), D498–D508. https://doi.org/10.1093/nar/gkaa1025

Cho, H. S., Seo, S. W., Kim, Y. M., Jung, G. Y., & Park, J. M. (2012). Engineering

glyceraldehyde-3-phosphate dehydrogenase for switching control of glycolysis in

escherichia coli. In *Biotechnology and Bioengineering* (Vol. 109, Issue 10, pp.

2612–2619). https://doi.org/10.1002/bit.24532

Chung, H., Yang, J. E., Ha, J. Y., Chae, T. U., Shin, J. H., Gustavsson, M., & Lee, S. Y.

705 (2015). Bio-based production of monomers and polymers by metabolically

706 engineered microorganisms. *Current Opinion in Biotechnology*, *36*, 73–84.

707 https://doi.org/10.1016/j.copbio.2015.07.003

708 Cornish-Bowden, A. (1979). Chapter 6 - Two-substrate reactions. In A. Cornish-Bowden

709 (Ed.), *Fundamentals of Enzyme Kinetics* (pp. 99–129). Butterworth-Heinemann.

710 https://doi.org/https://doi.org/10.1016/B978-0-408-10617-7.50011-0

711 Costa, R. S., Hartmann, A., & Vinga, S. (2016). Kinetic modeling of cell metabolism for

712 microbial production. *Journal of Biotechnology*, *219*, 126–141.

713 https://doi.org/10.1016/j.jbiotec.2015.12.023

714 Du, B., Zielinski, D. C., Kavvas, E. S., Dräger, A., Tan, J., Zhang, Z., Ruggiero, K. E.,

715 Arzumanyan, G. A., & Palsson, B. O. (2016). Evaluation of rate law approximations

716 in bottom-up kinetic models of metabolism. *BMC Systems Biology*, *10*(1), 40.

717 https://doi.org/10.1186/s12918-016-0283-2

718 Dudley, Q. M., Karim, A. S., & Jewett, M. C. (2015). Cell-free metabolic engineering:

719 Biomanufacturing beyond the cell. *Biotechnology Journal*, *10*(1), 69–82.

720 https://doi.org/10.1002/biot.201400330

721 Dudley, Q. M., Karim, A. S., Nash, C. J., & Jewett, M. C. (2020). In vitro prototyping of

722 limonene biosynthesis using cell-free protein synthesis. *Metabolic Engineering*,

723 *61*(April), 251–260. https://doi.org/10.1016/j.ymben.2020.05.006

724 Fackler, N., Heijstra, B. D., Rasor, B. J., Brown, H., Martin, J., Ni, Z., Shebek, K. M.,

725 Rosin, R. R., Simpson, S. D., Tyo, K. E., Giannone, R. J., Hettich, R. L.,

726 Tschaplinski, T. J., Leang, C., Brown, S. D., Jewett, M. C., & Kodiepke, M. (2021).

727 Stepping on the Gas to a Circular Economy: Accelerating Development of Carbon-

728 Negative Chemical Production from Gas Fermentation. *Annual Review of Chemical*

729 *and Biomolecular Engineering*, *12*, 439–470. https://doi.org/10.1146/annurev-

730 chembioeng-120120-021122

Galanie, S., Thodey, K., Trenchard, I. J., Interrante, M. F., & Smolke, C. D. (2015).

Complete biosynthesis of opiods in yeast. *Science*, *116*(2000), 1594–1602.

https://doi.org/10.1016/j.celrep.2014.10.048.Coupling

Gopalakrishnan, S., Dash, S., & Maranas, C. (2020). K-FIT: An accelerated kinetic

parameterization algorithm using steady-state fluxomic data. *Metabolic*

Engineering, *61*(March), 197–205. https://doi.org/10.1016/j.ymben.2020.03.001

Greene, J., Daniell, J., Köpke, M., Broadbelt, L., & Tyo, K. E. J. (2019). Kinetic ensemble

model of gas fermenting Clostridium autoethanogenum for improved ethanol

production. *Biochemical Engineering Journal*, *148*(May 2018), 46–56.

https://doi.org/10.1016/j.bej.2019.04.021

Greene, J. L., Wäechter, A., Tyo, K. E. J., & Broadbelt, L. J. (2017). Acceleration

Strategies to Enhance Metabolic Ensemble Modeling Performance. *Biophysical*

Journal, *113*(5), 1150–1162. https://doi.org/10.1016/j.bpj.2017.07.018

Grubbe, W. S., Rasor, B. J., Krüger, A., Jewett, M. C., & Karim, A. S. (2020). Cell-free

styrene biosynthesis at high titers. *Metabolic Engineering*, *61*(May), 89–95.

https://doi.org/10.1016/j.ymben.2020.05.009

Hofmeyr, J.-H. S. (2001). Metabolic control analysis in a nutshell. *Proceedings of the*

2nd International Conference on Systems Biology, *ii*, 291–300.

http://www.siliconcell.net/sica/NWO-CLS/CellMath/OiOvoer/Hofmeyr_nutshell.pdf

Horvath, N., Vilkhovoy, M., Wayman, J. A., Calhoun, K., Swartz, J., & Varner, J. D.

(2020). Toward a genome scale sequence specific dynamic model of cell-free

protein synthesis in Escherichia coli. *Metabolic Engineering Communications*,

10(November 2018), e00113. https://doi.org/10.1016/j.mec.2019.e00113

Jenior, M. L., Moutinho, T. J., Dougherty, B. V., & Papin, J. A. (2020). Transcriptome-

guided parsimonious flux analysis improves predictions with metabolic networks in

complex environments. *PLOS Computational Biology*, *16*(4), e1007099.

https://doi.org/10.1371/journal.pcbi.1007099

Jewett, M. C., Calhoun, K. A., Voloshin, A., Wuu, J. J., & Swartz, J. R. (2008). An

integrated cell-free metabolic platform for protein production and synthetic biology.

Molecular Systems Biology, *4*(220). https://doi.org/10.1038/msb.2008.57

Jia, G., Stephanopoulos, G., & Gunawan, R. (2012). Incremental parameter estimation

of kinetic metabolic network models. *BMC Systems Biology*, *6*.

https://doi.org/10.1186/1752-0509-6-142

Karim, A. S., Dudley, Q. M., Juminaga, A., Yuan, Y., Crowe, S. A., Heggestad, J. T.,

Garg, S., Abdalla, T., Grubbe, W. S., Rasor, B. J., Coar, D. N., Torculas, M., Krein,

M., Liew, F. M. (Eric), Quattlebaum, A., Jensen, R. O., Stuart, J. A., Simpson, S. D.,

Köpke, M., & Jewett, M. C. (2020). In vitro prototyping and rapid optimization of

biosynthetic enzymes for cell design. *Nature Chemical Biology*, *16*(8), 912–919.

https://doi.org/10.1038/s41589-020-0559-0

Karim, A. S., Heggestad, J. T., Crowe, S. A., & Jewett, M. C. (2018). Controlling cell-free

metabolism through physiochemical perturbations. *Metabolic Engineering*, *45*(May

2017), 86–94. https://doi.org/10.1016/j.ymben.2017.11.005

Karim, A. S., & Jewett, M. C. (2016). A cell-free framework for rapid biosynthetic

pathway prototyping and enzyme discovery. *Metabolic Engineering*, *36*, 116–126.

https://doi.org/10.1016/j.ymben.2016.03.002

Karim, A. S., Rasor, B. J., & Jewett, M. C. (2019). Enhancing control of cell-free

metabolism through pH modulation. *Synthetic Biology*, *5*(1), 1–7.

https://doi.org/10.1093/synbio/ysz027

Khodayari, A., Zomorrodi, A. R., Liao, J. C., & Maranas, C. D. (2014). A kinetic model of

Escherichia coli core metabolism satisfying multiple sets of mutant flux data.

Metabolic Engineering, *25*, 50–62. https://doi.org/10.1016/j.ymben.2014.05.014

Kim, O. D., Rocha, M., & Maia, P. (2018). A Review of Dynamic Modeling Approaches

and Their Application in Computational Strain Optimization for Metabolic Engineering. *Frontiers in Microbiology*, *9*(July), 1–22. https://doi.org/10.3389/fmicb.2018.01690

Kurata, H., & Sugimoto, Y. (2018). Improved kinetic model of Escherichia coli central carbon metabolism in batch and continuous cultures. *Journal of Bioscience and Bioengineering*, *125*(2), 251–257. https://doi.org/10.1016/j.jbiosc.2017.09.005

Lewis, N. E., Hixson, K. K., Conrad, T. M., Lerman, J. A., Charusanti, P., Polpitiya, A. D., Adkins, J. N., Schramm, G., Purvine, S. O., Lopez-Ferrer, D., Weitz, K. K., Eils, R., König, R., Smith, R. D., & Palsson, B. (2010). Omic data from evolved E. coli are consistent with computed optimal growth from genome-scale models. *Molecular Systems Biology*, *6*(390). https://doi.org/10.1038/msb.2010.47

Liew, F. E., Nogle, R., Abdalla, T., Rasor, B. J., Canter, C., Jensen, R. O., Wang, L., Strutz, J., Chirania, P., De Tissera, S., Mueller, A. P., Ruan, Z., Gao, A., Tran, L., Engle, N. L., Bromley, J. C., Daniell, J., Conrado, R., Tschaplinski, T. J., … Köpke, M. (2022). Carbon-negative production of acetone and isopropanol by gas fermentation at industrial pilot scale. *Nature Biotechnology*, *40*(3), 335–344. https://doi.org/10.1038/s41587-021-01195-w

Mahadevan, R., Edwards, J. S., & Doyle, F. J. (2002). Dynamic flux balance analysis of diauxic growth. *Biophysical Journal*, *83*(3), 1331–1340. https://pdfs.semanticscholar.org/8d56/a26c56704266b5d7c426ecdfc286962f9112.pdf

Mendoza, S. N., Olivier, B. G., Molenaar, D., & Teusink, B. (2019). A systematic assessment of current genome-scale metabolic reconstruction tools. *Genome Biology*, *20*(158). https://doi.org/10.1186/s13059-019-1769-1

Miguez, A. M., McNerney, M. P., & Styczynski, M. P. (2019). Metabolic Profiling of Escherichia coli-Based Cell-Free Expression Systems for Process Optimization.

Industrial and Engineering Chemistry Research, *58*(50), 22472–22482. https://doi.org/10.1021/acs.iecr.9b03565

Miguez, A. M., Zhang, Y., Piorino, F., & Styczynski, M. P. (2021). Metabolic Dynamics in Escherichia coli-Based Cell-Free Systems. *ACS Synthetic Biology*. https://doi.org/10.1021/acssynbio.1c00167

Min Lee, J., Gianchandani, E. P., Eddy, J. A., & Papin, J. A. (2008). Dynamic analysis of integrated signaling, metabolic, and regulatory networks. *PLoS Computational Biology*, *4*(5). https://doi.org/10.1371/journal.pcbi.1000086

Naseri, G., & Koffas, M. A. G. (2020). Application of combinatorial optimization strategies in synthetic biology. *Nature Communications*, *11*(1), 1–14. https://doi.org/10.1038/s41467-020-16175-y

Norsigian, C. J., Pusarla, N., McConn, J. L., Yurkovich, J. T., Dräger, A., Palsson, B. O., & King, Z. (2020). BiGG Models 2020: Multi-strain genome-scale models and expansion across the phylogenetic tree. *Nucleic Acids Research*, *48*(D1), D402–D406. https://doi.org/10.1093/nar/gkz1054

Orth, J. D., Palsson, B. Ø., & Fleming, R. M. T. (2010). Reconstruction and Use of Microbial Metabolic Networks: the Core Escherichia coli Metabolic Model as an Educational Guide. *EcoSal Plus*, *4*(1). https://doi.org/10.1128/ecosalplus.10.2.1

Orth, J. D., Thiele, I., & Palsson, B. O. (2010). What is flux balance analysis? *Nature Biotechnology*, *28*(3), 245–248. https://doi.org/10.1038/nbt.1614

Rasor, B. J., Vögeli, B., Jewett, M. C., & Karim, A. S. (2022). Cell-Free Protein Synthesis for High-Throughput Biosynthetic PathwayBiosynthetic pathways Prototyping. In A. S. Karim & M. C. Jewett (Eds.), *Cell-Free Gene Expression: Methods and Protocols* (pp. 199–215). Springer US. https://doi.org/10.1007/978-1-0716-1998-8_12

Rasor, B. J., Vögeli, B., Landwehr, G. M., Bogart, J. W., Karim, A. S., & Jewett, M. C. (2021). Toward sustainable, cell-free biomanufacturing. *Current Opinion in*

Biotechnology, *69*, 136–144. https://doi.org/10.1016/j.copbio.2020.12.012

Ro, D. K., Paradise, E. M., Quellet, M., Fisher, K. J., Newman, K. L., Ndungu, J. M., Ho,

K. A., Eachus, R. A., Ham, T. S., Kirby, J., Chang, M. C. Y., Withers, S. T., Shiba,

Y., Sarpong, R., & Keasling, J. D. (2006). Production of the antimalarial drug

precursor artemisinic acid in engineered yeast. *Nature*, *440*(7086), 940–943.

https://doi.org/10.1038/nature04640

Saa, P. A., & Nielsen, L. K. (2016). Construction of feasible and accurate kinetic models

of metabolism: A Bayesian approach. *Scientific Reports*, *6*.

https://doi.org/10.1038/srep29635

Saa, P. A., & Nielsen, L. K. (2017). Formulation, construction and analysis of kinetic

models of metabolism: A review of modelling frameworks. *Biotechnology Advances*,

35(8), 981–1003. https://doi.org/10.1016/j.biotechadv.2017.09.005

Sánchez, B. J., Zhang, C., Nilsson, A., Lahtvee, P.-J., Kerkhoven, E. J., & Nielsen, J.

(2017). Improving the phenotype predictions of a yeast genome-scale metabolic

model by incorporating enzymatic constraints. *Molecular Systems Biology*, *13*(8),

935. https://doi.org/10.15252/msb.20167411

Shestov, A. A., Liu, X., Ser, Z., Cluntun, A. A., Hung, Y. P., Huang, L., Kim, D., Le, A.,

Yellen, G., Albeck, J. G., & Locasale, J. W. (2014). Quantitative determinants of

aerobic glycolysis identify flux through the enzyme GAPDH as a limiting step. *ELife*,

3(July2014), 1–18. https://doi.org/10.7554/eLife.03342

Silverman, A. D., Karim, A. S., & Jewett, M. C. (2020). Cell-free gene expression: an

expanded repertoire of applications. *Nature Reviews Genetics*, *21*(3), 151–170.

https://doi.org/10.1038/s41576-019-0186-3

Solem, C., Koebmann, B. J., & Jensen, P. R. (2003). Glyceraldehyde-3-phosphate

dehydrogenase has no control over glycolytic flux in Lactococcus lactis MG1363.

Journal of Bacteriology, *185*(5), 1564–1571. https://doi.org/10.1128/JB.185.5.1564-

1571.2003

St John, P. C., Strutz, J., Broadbelt, L. J., Tyo, K. E. J., & Bomble, Y. J. (2019). Bayesian

inference of metabolic kinetics from genome-scale multiomics data. *PLoS*

Computational Biology, *15*(11), 1–23. https://doi.org/10.1371/journal.pcbi.1007424

Strutz, J., Martin, J., Greene, J., Broadbelt, L., & Tyo, K. (2019). Metabolic kinetic

modeling provides insight into complex biological questions, but hurdles remain.

Current Opinion in Biotechnology, *59*. https://doi.org/10.1016/j.copbio.2019.02.005

Suthers, P. F., Foster, C. J., Sarkar, D., Wang, L., & Maranas, C. D. (2021). Recent

advances in constraint and machine learning-based metabolic modeling by

leveraging stoichiometric balances, thermodynamic feasibility and kinetic law

formalisms. *Metabolic Engineering*, *63*(August 2020), 13–33.

https://doi.org/10.1016/j.ymben.2020.11.013

Tan, Y., Lafontaine Rivera, J. G., Contador, C. A., Asenjo, J. A., & Liao, J. C. (2011).

Reducing the allowable kinetic space by constructing ensemble of dynamic models

with the same steady-state flux. *Metabolic Engineering*, *13*(1), 60–75.

https://doi.org/10.1016/j.ymben.2010.11.001

Tran, L. M., Rizk, M. L., & Liao, J. C. (2008). Ensemble modeling of metabolic networks.

Biophysical Journal, *95*(12), 5606–5617.

https://doi.org/10.1529/biophysj.108.135442

Tsiantis, N., & Banga, J. R. (2020). Using optimal control to understand complex

metabolic pathways. *BMC Bioinformatics*, *21*(472). https://doi.org/10.1186/s12859-

020-03808-8

van Eunen, K., Kiewiet, J. A. L., Westerhoff, H. V., & Bakker, B. M. (2012). Testing

biochemistry revisited: How in vivo metabolism can be understood from in vitro

enzyme kinetics. *PLoS Computational Biology*, *8*(4).

https://doi.org/10.1371/journal.pcbi.1002483

van Rosmalen, R. P., Smith, R. W., Martins dos Santos, V. A. P., Fleck, C., & Suarez-

Diez, M. (2021). Model reduction of genome-scale metabolic models as a basis for

targeted kinetic models. *Metabolic Engineering*, *64*(January), 74–84.

https://doi.org/10.1016/j.ymben.2021.01.008

Villaverde, A., Tsiantis, N., & Banga, J. (2019). Full observability and estimation of

unknown inputs, states, and parameters of nonlinear biological models. *J. R. Soc.*

Interface.

Vogeli, B., Schulz, L., Garg, S., Tarasava, K., Clomburg, J. M., Lee, H., Gonnot, A.,

Moully, E. H., Kimmel, B. R., Tran, L., Brown, S. D., Mrksich, M., Karim, A. S., &

Jewett, M. C. (2022). *Cell-free prototyping enables implementation of optimized*

reverse β-oxidation pathways in heterotrophic and autotrophic bacteria.

Wu, G., Yan, Q., Jones, J. A., Tang, Y. J., Fong, S. S., & Koffas, M. A. G. (2016).

Metabolic Burden: Cornerstones in Synthetic Biology and Metabolic Engineering

Applications. *Trends in Biotechnology*, *34*(8), 652–664.

https://doi.org/10.1016/j.tibtech.2016.02.010

Figure 1: Metabolic Network Map. The utilized network includes *E. coli* core metabolism (gray arrows), as well as the 6-step butanol pathway (black arrows at the bottom of the map). Measured metabolites are labeled in black text and shown as red circles. Cofactors are shown as colored circles according to the legend, and water and protons (not kinetically included in the model) are shown as smaller circles. All other species are shown as unfilled circles.

Figure 2: Model parameterization pipeline. (a) Parameter priors are initially

constructed from literature values and similar previous models. (b) Individual models, or

parameter sets, are sampled from parameter priors while simultaneously applying

thermodynamic values as constraints. This is repeated to create ensembles of 10^8

models. (c) The initial ensemble is simulated via ODEs and then compared to

experimental data; the ensemble is then screened to the best ~10^2 models based on

agreement to this data. (d) When the agreement to experimental data after screening

was still insufficient, parameter sensitivity analysis was performed and used to update

parameter priors for additional rounds of ensemble sampling and screening. (e) Once

screened models showed adequate agreement to experimental data, each individual

model was further refined via local parameter optimization. (f) This full pipeline

generated an ensemble of 20 models which gave excellent fit to the experimental data

and were further used for analysis and predictions.

926
Figure 3: Final ensemble simulations compared to experimental data. Model

simulated metabolite time courses are shown for the nine metabolites with experimental

measurements, as well as predicted time courses for the ratios of NADH to all NAD

species, ATP to all adenosine species, and free CoA to all CoA species. Experimental

measurements are shown as circles. The simulations of the top 20 models in the

ensemble are shown as confidence intervals (CI); the dark region is a 50% CI and

therefore contains the middle 50% of models, and the lighter region is a 95% CI. Blue

data represent the butanol-positive condition, and red data represent the butanol-

negative condition.

Figure 4: Predictions for butanol optimization. (a) Enzyme level changes were simulated to optimize 24-hour butanol titer. The left-most column shows the simulated butanol at 24 hours in the base butanol-positive condition, which was fit to experimental data. Each light blue dot represents one of the 20 models in the final ensemble, and their distribution is summarized by each violin plot with medians given by unfilled dark blue circles and interquartile ranges given by dark blue lines. The horizontal dashed line shows the base condition median prediction as a baseline. The next three columns show the predicted butanol titer from simultaneously adjusting all five enzymes in the butanol pathway by either 0.5x, 2x, or 5x their respective levels in the base production condition. The next five columns show the predicted butanol titer from adjusting each of the five

enzymes one at a time by 5x their respective levels in the base production condition. (b)

Initial cofactor level changes were simulated to optimize 24-hour butanol titer. The left-

most column shows the simulated butanol at 24 hours in the base butanol-positive

condition and the base cofactor concentrations in gray. The next five columns show the

predicted butanol titer from raising the concentrations of the cofactors predicted to

improve butanol titers, which included NADH, ATP, CoA, and pyrophosphate (ppi), either

individually or in groups. (c) Predicted butanol titers from adjusting the initial

concentrations of either glucose, acetate, or glutamate are shown. Initial concentrations

of each species in the base condition (left-most column) are shown in gray.

Figure 5: Analysis of parameter variation across models shows small

number of highly conserved parameters. Elasticity values for each parameter, which

serve as a more informative proxy for absolute parameter values, are plotted with their

mean value across the top 20 models in the final ensemble (x axis) versus the standard

deviation across the top 20 models (y axis). Each point represents a single elasticity or

model parameter, and marginal histograms for both mean and standard deviation show

distributions across these elasticities. Elasticity values near 0 represent elasticities for

metabolites that are fully saturated (zeroth order kinetics), and values near 1 represent

metabolites far below saturation in a linear regime (first order kinetics). Those

parameters which a mean elasticity between 0.4 and 0.6, representing a high degree of

control over the reaction rate, and a standard deviation below 0.3, representing highly

conserved behavior among all models in the ensemble, are highlighted with names

given in the legend, denoted by the reaction name (i.e., ACKr for acetate kinase), then

the reaction direction ("_f" for forward or "_r" for reverse), then the reaction substrate

(i.e., "ac_c" for acetate). A complete list of reaction, substrate, and elasticity names is

given Elasticity values are calculated as the derivative of reaction rate with respect to

metabolite concentration and are normalized with respect to both rate and concentration,

as well as the stoichiometry of the metabolite in each reaction. To ensure all values were

between 0 and 1, metabolites acting as inhibitors are not shown in this figure.