The effect of metal remediation on the virulence and antimicrobial resistance of the opportunistic pathogen *Pseudomonas aeruginosa*

Luke Lear[1*], Elze Hesse[1], Laura Newsome[2], William Gaze[3], Angus Buckling[1] and Michiel Vos[3].

[1]College of Life and Environmental Science, University of Exeter, Penryn, Cornwall, United Kingdom, TR10 9FE

[2]College of Engineering, Mathematics and Physical Sciences, University of Exeter, Penryn, Cornwall, United Kingdom, TR10 9FE

[3]European Centre for Environment and Human Health, University of Exeter Medical School, Penryn, Cornwall, United Kingdom, TR10 9FE

*Corresponding author; email: L.M.Lear@exeter.ac.uk

Running title: Liming and bacterial pathogenicity

Keywords: liming, antibiotic resistance, metal pollution, siderophores, *Pseudomonas aeruginosa*, opportunistic pathogen

Abstract

Metal contamination poses both a direct threat to human health as well as an indirect threat through its potential to affect bacterial pathogens. Metals can not only co-select for antibiotic resistance, but also might affect pathogen virulence via increased siderophore production. Siderophores are extracellular compounds released to increase ferric iron uptake — a common limiting factor for pathogen growth within hosts – making them an important virulence factor. However, siderophores can also be positively selected for to detoxify non-ferrous metals, and consequently metal stress can potentially increase bacterial virulence. Anthropogenic methods to remediate environmental metal contamination commonly involve amendment with lime-containing materials, but whether this reduces *in situ* co-selection for antibiotic resistance and virulence remains unknown. Here, using microcosms containing metal-contaminated river water and sediment, we experimentally test whether metal remediation by liming reduces co-selection for these traits in the opportunistic pathogen *Pseudomonas aeruginosa* embedded within a natural microbial community. To test for the effects of environmental structure, which can impact siderophore production, microcosms were incubated under either static or shaking conditions. Evolved *P. aeruginosa* populations had greater fitness in the presence of toxic concentrations of copper than the ancestral strain, but this effect was reduced in the limed treatments. Evolved *P. aeruginosa* populations showed increased resistance to the clinically-relevant antibiotics apramycin, cefotaxime, and trimethoprim, regardless of lime addition or environmental structure. Although we found virulence to be significantly associated with siderophore production, neither virulence nor siderophore production significantly differed between the four treatments. We therefore demonstrate that although remediation via liming reduced the strength of selection for metal resistance mechanisms, it did not mitigate metal-imposed selection for antibiotic resistance or virulence in *P. aeruginosa*. Consequently, metal-contaminated environments may select for antibiotic resistance and virulence traits even when treated with lime.

<u>Graphical abstract</u>

<u>Introduction</u>

Metals are ubiquitous in the Earth's crust and many are essential for cellular processes

(1-3). However, agriculture and industry have resulted in toxic levels of metal

contamination in many environments via pesticide use, sewage sludge application,

atmospheric deposition and mining (1, 4). Human exposure to metals via

contaminated crops or water poses a serious health threat (5-9). Furthermore, metal

contamination can decrease microbial diversity, biomass and functionality, which in

turn can affect ecosystem function (10-12). Such are the potential health and

economic costs that metal remediation is common practice (13). Often, this involves

adding lime-containing materials to acidic metal-contaminated environments (14-16).

By raising the pH, liming causes metal ions to precipitate and become less soluble

and consequently less bioavailable (10, 13, 17-19). This decreases metal uptake by

plants which helps alleviate phytotoxicity and prevent metals entering the food chain

(20). Lime addition is the oldest and most widely used metal remediation method (19),

and has been used to treat contamination in lakes (14, 21), rivers (22) and soils (15,

16, 18).

Reducing metal bioavailability by liming will inevitably affect the microbial community

(18, 23, 24). However, how such remediation could affect the ability of environmental

pathogens to cause infection or to withstand antibiotic treatment remains largely

unexplored. A key microbial trait likely to change after liming is the production of

siderophore compounds (18). The canonical function of siderophores is to aid iron (Fe)

sequestration from the extra-cellular environment (25, 26). Fe is vital for microbial

growth as a cofactor for a number of essential enzymes (27, 28), but is most commonly

present as insoluble Fe^{3+} and therefore is of limited bioavailability, especially at near-

neutral pH (27-32). Siderophores are released by cells where they form extracellular

complexes with Fe^{3+}, these are then taken up by selective outer-membrane transport

proteins before Fe^{3+} is reduced to bioavailable Fe^{2+} and the siderophore made

available for reuse (33). Siderophores are important virulence factors as they allow

pathogens to grow within hosts that actively withhold iron (34, 35). Apart from iron,

siderophores can also chelate toxic metal ions, but these complexes cannot re-enter

the cell due to the selectivity of the outer-membrane transport proteins (28, 29). This

means siderophore production can be selected for as a detoxifying method in the

presence of bioavailable toxic metals (26, 29, 36). Consequently, toxic metal

concentrations can select for greater virulence by selecting for increased siderophore

production (37). Lime remediation of metal-contaminated environments thus could

potentially select either for the upregulation of siderophore production when it

predominantly results in decreased bioavailability of Fe, or for the downregulation of

siderophore production when it predominantly results in lower metal toxicity, with

concomitant expected changes in virulence. Previous work has shown siderophore

production to decrease as a consequence of liming at the level of whole microbial

communities (18), but whether this also occurs in environmental pathogens that rely

on siderophore-mediated iron uptake remains untested.

It is well established that some mechanisms that bacteria use to resist metal

contamination also confer resistance to antibiotics (38). This can occur through cross-

resistance when a single mechanism provides resistance to both types of stressors

(e.g. efflux pumps (38-44)), through co-resistance when metal and antibiotic

resistance genes are located on the same genetic element (45, 46)), or through co-

regulation when transcriptional and translational responses to both stressors are

linked (38, 43, 47-49). However, to our knowledge, it remains untested whether metal

remediation could decrease such co-selection for antibiotic resistance.

In this study, we use the opportunistic pathogen *Pseudomonas aeruginosa* to test whether liming alters virulence by influencing siderophore production, and whether it decreases co-selection by metals for antibiotic resistance. We applied an experimental evolution approach, utilising microcosms containing water and sediment and the resident microbial community from a river heavily contaminated with historical mine waste (50, 51). We embedded *P. aeruginosa* within this natural microbial community and quantified antibiotic resistance, siderophore production and virulence in this focal species after 14 days. *P. aeruginosa* is responsible for a significant proportion of nosocomial infections, particularly those in intensive care units and immunocompromised patients (52). This species is of significant clinical importance as it is resistant to many treatments, both intrinsically and due to its ability to rapidly evolve resistance (53). Outside of the clinical setting, *P. aeruginosa* is commonly found in soil and water (54). The production of siderophores by *P. aeruginosa* is well-studied as a virulence factor, metal resistance mechanism and public good (25, 27, 29, 55-57). Furthermore, the growing interest in its use along with other siderophore producing species to assist phytoremediation of metals using plants (28, 58), makes it an ideal focal species for this study.

The insect infection model *Galleria mellonella* (Greater Wax Moth larvae), a low-cost and ethically expedient alternative for mammalian virulence screens (59), is used here to quantify *P. aeruginosa* virulence (60). We quantified total siderophore production using a CAS assay (61) and pyoverdine production – the main siderophore produced by *P. aeruginosa* (62) – by measuring fluorescence; and tested whether these are correlated with virulence. Extracellular siderophore-metal complexes offer a fitness advantage not only to the producer but also to neighbouring cells, whether these are fellow-producers or not (63-65). Non-siderophore producing 'cheats' could gain a selective advantage as they benefit from siderophore production but do not carry the cost of production (25, 30, 31, 66). Cheat fitness is increased in spatially unstructured environments because the greater mixing increases the opportunity to take up siderophore-iron complexes and benefit from siderophores detoxifying the area (65). To take into account the effect of spatial structure on siderophore production, and consequently virulence, we performed our experiments in both static and shaken microcosms. We tested whether the addition of lime or a change in spatial structure affects *P. aeruginosa* resistance to the antibiotics apramycin, cefotaxime and

150 trimethoprim. Both apramycin and cefotaxime have been declared 'critically important'
151 for human medicine, and trimethoprim 'highly important' by the WHO (67). Moreover,
152 apramycin has been shown to be effective against highly drug resistant strains of *P.*
153 *aeruginosa* (68).
154
Methods
156
Collection of river samples and microcosm set-up
Water and sediment samples were collected from the metal-contaminated Carnon
River in Cornwall, U.K. (50°13'54.6' N, 5°07'48.7' W) (53). We chose this site as it is
polluted by multiple toxic metals (18, 50) and liming has been shown to reduce total
non-ferrous metal availability in samples collected in the near vicinity (18). The water
sample in reference (50) was taken at the same time as the samples used in this study,
and contains high concentrations of non-ferrous metals. Sediment was collected using
a sterile spatula and water was collected by filling a sterile 1000 mL duran bottle
(Schott Duran, Munich, Germany). Sediment (3 g +/-0.1 g) and river water (6 mL) was
added to each microcosm (25 mL, Kartell, Noviglio, Italy). The combined water and
sediment pH was measured using a Jenway 3510 pH meter (Jenway, Essex, UK).
168
Experimental design
Two treatments – liming (lime amendment/no amendment) and spatial structure
(shaken/unshaken) – were carried out in a full factorial design (Fig. 1); six replicates
were used per unique treatment combination resulting in a total of 24 microcosms. All
microcosms were incubated at 20°C. To raise the pH from 5.8 to ~7.0 to represent a
metal remediation scenario, 30 mg (+/- 1.0 mg) of undissolved hydrated lime (Verve
Garden lime, Eastleigh, U.K. (18)) was added to each relevant microcosm, then left
for 14 days to equilibrate. To observe differences between structured and non-
structured environments, microcosms were either kept static or were continuously
shaken at 210 rpm (Stuart orbital incubator S1600, Staffordshire, UK). Shaking began
on day 14 and ended on day 28 (Fig.1).

Figure 1 Timeline of experimental design. Lime addition and shaking treatments were performed factorially, with lime added on day 1 and shaking at 210 rpm starting on day 14. *Pseudomonas aeruginosa* was added to all microcosms on day 14. Microcosms were destructively sampled on day 28. Six replicates were used for all treatments (24 microcosms in total).

On day 14, 30 µL (7.3 x 10^8 colony forming units: cfu) of *Pseudomonas aeruginosa* (PAO1^R*lacZ*: (69)) was added to each microcosm. This lab-strain is both lacZ marked and gentamicin resistant allowing it to be easily distinguished from the rest of the community on agar containing X-gal (5-bromo-4-chloro-3-indolyl-β-D-galactopyranoside; 100 µg/L; VWR Chemicals) and gentamicin (30 µg/mL Sigma). *P. aeruginosa* was grown overnight in shaking microcosms containing 6 mL of King's medium B (KB; 10g glycerol, 20g proteose peptone no. 3, 1.5 g K_2HPO_4, 1.5 g $MgSO_4$, per litre). To remove residual nutrients, cultures were centrifuged at 3500 rpm (1233 g) for 30 minutes, after which supernatant was decanted, and the pellet resuspended in half the volume of M9 salt buffer (3 g KH_2PO_4, 6 g Na_2HPO_4, 5 g NaCl/L) followed by plating on KB agar to calculate the inoculation density. On day 28, all microcosms were destructively sampled by adding sterile glass beads and 12 mL of M9 buffer and vortexing for one minute. Samples were then aliquoted and stored in glycerol (25% final volume) at -80°C.

Iron analysis (ferrozine assay)

To determine if liming affected Fe speciation and therefore bioavailability, a ferrozine assay was used to measure relative concentrations of Fe^{2+} and total bioavailable iron (70, 71). The first step of this assay quantifies Fe^{2+} which is easily obtainable by bacteria and so does not require siderophores. The second step quantifies both Fe^{2+} and Fe^{3+} and therefore gives a measure of total bioavailable iron including that which requires scavenging mechanisms, such as siderophores. By dividing the first measurement by the second it is possible to estimate the proportion of iron in each treatment that is of relatively high bioavailability to *P. aeruginosa* (70, 71). The first measurement is given by digesting 100 µL of fresh sample (n=3) in 4.9 mL of 0.5 M hydrochloric acid for 1 hour before 50 µL was mixed with 2.45 mL of ferrozine solution (1 g ferrozine, 11.96 g (4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid/L; adjusted to pH 7) in a cuvette (n = 3 per replicate). This was left to stand for exactly one minute before absorbance at 562 nm was measured using a spectrophotometer (Jenway 7315, Essex, UK). To quantify total bioavailable Fe (step 2), 200 µL of 6.25 M hydroxylamine hydrochloride was added to the digested samples and left to stand for another hour. This was then added to ferrozine solution in cuvettes and measured as before. Standards of known concentrations of $FeSO_4.7H_2O$ were measured to allow conversion of absorbance to Fe concentrations.

Copper growth assay

To confirm that metal concentrations in our river water and sediment samples were sufficiently high to select for metal resistance mechanisms, and to test whether liming impacted this selection, we used a copper growth assay. Specifically, we added 20 µL of either the ancestral *P. aeruginosa* strain or defrosted samples of the evolved populations to a 96-well plate well containing 180 µL of plain Iso-Sensitest broth (Oxoid) and 20 µL to a well containing 180 µL of Iso-Sensitest broth at a concentration of 1 g/L of copper sulphate ($CuSO_4$; Alfa Aesar, Massachusetts, United States). The optical density OD_{600} was then read every 10 minutes for 18 hours using a Biotek Synergy 2 spectrophotometer. We used 1 g/L of copper sulphate as this equates to a copper concentration (6.26 mM) previously found in highly polluted environments (72, 73).

Siderophore (CAS) assay

Total siderophore production was quantified using the Chrome Azurol S (CAS) assay (74). Samples were plated onto tryptic soy agar (TSA: Oxoid) supplemented with nystatin (Sigma: 20 µg/mL) to suppress fungal growth and X-gal. After 48 hours, *P. aeruginosa* colonies were counted to quantify density, before 24 colonies per replicate were randomly picked using sterile toothpicks. Selected colonies were resuspended in 1 mL of KB media in a deep 96-well plate and grown overnight at 28 °C. These were mixed with glycerol (final concentration 25%) and frozen at -80 °C. A scraping from each frozen monoculture was then grown in 2 mL of iron-limited casamino acid (CAA: Fisher) medium overnight. Iron limitation was caused by the addition of human apotransferrin (100 mg/mL; BBI Solutions, Crumlin) and sodium bicarbonate (20 mM; Acros Organics) to induce the production of siderophores. Cultures were centrifuged, and the supernatant assayed using the liquid CAS assay to quantify total siderophores, whilst pyoverdine was quantified by measuring the fluorescence of each culture at 460 nm following excitation at 400 nm. By measuring the optical density of the pre-centrifuged cultures and quantifying absorbance of sterile media, siderophore production per clone was estimated using: $[1 - (A_i/A_{ref})] /[OD_i)]$, where OD_i = optical density 600 nm and A_i = absorbance at 630 nm of the assay mixture and A_{ref} = absorbance at 630 nm of reference mixture (CAA+CAS) (18, 75).

Galleria mellonella virulence assay

The insect infection model *Galleria mellonella* was used to quantify *P. aeruginosa* virulence (59, 60). Defrosted freezer stocks containing the whole sample microbiome were diluted 100-fold using M9 salt buffer, before 10 µL was injected into twenty final instar larvae per replicate using a 50 µL syringe (Hamilton, Nevada, USA). Injected larvae were incubated at 37 °C and mortality was monitored hourly after 13 hours for 12 hours with a final check at 42 hours. Larvae were classed as dead when mechanical stimulation of the head caused no response (60). M9-injected and non-injected controls were used to confirm mortality was not due to injection trauma or background *G. mellonella* mortality; >10% control death was the threshold for re-injecting (no occurrences). Prior to assays on microcosms containing *P. aeruginosa*, we confirmed the natural microbial community caused zero mortality by injecting replicates not inoculated with *P. aeruginosa* as described above.

 Antibiotic resistance assay

To test the evolved resistance of *P. aeruginosa* to the clinically relevant antibiotics

apramycin, cefotaxime and trimethoprim, we used the same *P. aeruginosa* colonies

isolated for the siderophore analysis. We first determined the minimum inhibitory

concentration of the three antibiotics for our ancestral strain, by growing the ancestral

strain for 24 hours (as described above), and plating it on TSA containing a range of

concentrations of the antibiotics that increased in 10 µg/mL increments from 0 - 60

µg/mL. The minimum inhibitory concentrations were found to be 12 µg/mL, 30 µg/mL

and 40 µg/mL for apramycin, cefotaxime and trimethoprim, respectively. Next, the

individual evolved clones were defrosted before 2 µL of each was plated onto either

plain TSA, TSA containing apramycin (Sigma: 15 µg/mL) cefotaxime (Molekula: 50

µg/mL) or TSA containing trimethoprim (Sigma: 60 µg/mL). Strains were considered

resistant if colonies could be observed after 48 hours. The ancestral strain was used

as a negative control.

Statistical analysis

The effect of liming and shaking, plus their interaction, on the final pH, density of *P.*

aeruginosa ($\log_{10}$(cfu mL^{-1})), and the proportion of total bioavailable iron (Fe^{2+} + Fe^{3+})

that was Fe^{2+} (Fe^{2+} / (Fe^{2+} + Fe^{3+})) was tested using linear models with liming and

shaking as explanatory variables. In general, model reduction was carried out by

sequentially removing terms from the full model and comparing model fits using *F*-

tests; we report parameter estimates of the most parsimonious model. The effect of

pH on the density of *P. aeruginosa* populations was tested using a linear model with

density (cfu mL^{-1}) $\log_{10}$ transformed.

To test whether evolved samples had greater resistance to copper than the ancestral

strain, we first calculated the relative fitness, *w*, of each population by dividing its

maximum optical density after 18 hours when grown with copper (OD_{maxC}) by its

maximum optical density when grown without copper (OD_{maxWC}), *i.e. w* = OD_{maxC} /

OD_{maxP}. We then carried out a one way ANOVA with *w* as the response variable and

treatment (including ancestor) as the explanatory variable. Secondly, we carried out a

Dunnett's test, using the '*DescTools'* R package (76), to test whether each treatment

differed to the ancestor. Finally, we tested the effect of liming and shaking on the metal

resistance of the final populations in a linear model, with log(w) as the response

variable, and liming, shaking and their interaction as the explanatory variables. In all

tests w was log transformed to normalise the residuals

To test liming and shaking effects on total siderophore and pyoverdine production,

linear mixed effects models (LMEM) were carried out using the '*lme4*' package (77)

with liming and shaking as explanatory variables, and random intercepts fitted for each

replicate to control for multiple clones being sampled from the same microcosm. For

these LMEMs, we used the '*DHARMa*' package (78) to check residual behaviour, after

which the most parsimonious model was arrived at by comparing models with and

without the liming-shaking interaction using χ-tests. Two samples had pyoverdine

values much lower than the rest, so a Grubbs test ('*outliers*' package; (79)) was used

to check if they were significant outliers. They were therefore were removed from this

and all further models to improve model fit. To test the association between copper

resistance and both total siderophore and pyoverdine production, we carried out two

linear models with log(w) as the dependent variable and either mean total siderophore

per microcosm or mean pyoverdine production per microcosm as the explanatory

variable.

Virulence was analysed in three separate models. First, we tested whether larvae that

died before 42 hours were injected with samples containing more siderophores and

pyoverdine than those that remained alive after 42 hours. This was done by carrying

out two separate binomial generalised linear mixed models (GLMM) using the '*lme4*'

package (77), with number of *G. mellonella* dead versus alive as the binomial

response variable, and either the production of total siderophores or pyoverdine as

the explanatory variable. In this model pyoverdine production was $\log_{10}$ transformed

to normalise residuals. Secondly, we tested whether the mean time it took deceased

larvae (20 per replicate) to die was associated with total siderophore and pyoverdine

production (both values taken from the mean of 24 clones) using a linear model.

Finally, we tested whether virulence differed between treatments. To do this, survival

curves were fitted using Bayesian regression in the R package '*rstanarm*' (80) and the

package '*tidybayes*' (81) was used to estimate parameters. A proportional hazards

model with an M-splines baseline hazard was fitted, with liming, shaking plus their

interaction as fixed effects. We additionally included random intercepts for each

sample to control for multiple (20) *G. mellonella* being inoculated with the same
sample. Models used three chains with uninformative priors and were run for 3000
iterations. Model convergence was assessed using Rhat values (all values were 1),
before we manually checked chain mixing.

The proportion of apramycin, cefotaxime, and trimethoprim resistance in each
treatment (number of resistant colonies out of 24 in total) was compared using Kruskal-
Wallace non-parametric tests, with resistance proportion as the response variable and
treatment as the explanatory variable. All analyses were carried out in R version 3.3.3
(82).

<u>Results and discussion</u>

Liming and shaking affected pH but not the relative abundance of Fe^{2+}

Here, we tested whether liming of metal-contaminated aquatic environments
decreases co-selection for virulence and antibiotic resistance in the opportunistic
pathogen *P. aeruginosa*. To do this, we evolved *P. aeruginosa* with or without lime in
microcosms containing a mixture of metal contaminated river water and sediment in
the presence of the natural microbial community. We employed both shaking and
static microcosms to represent turbulent and stagnant aquatic environments, in order
to test whether liming effects were dependent on environmental structure (Fig. 1).
As expected, liming significantly decreased the acidity of sediment and water from the
initial pH of 5.8. However, the extent of this effect was significantly greater in the
shaking treatments (liming-shaking interaction: $F_{1,20}=23.1$, $p<0.001$; Fig. 2), likely due
to increased mixing of lime and oxygen throughout the microcosms. The shaken-limed
treatment reached a pH of 7.2 (+/- 0.11 SD) whereas the static-limed treatment
reached a pH of 6.7 (+/- 0.25 SD). Both non-limed treatments had a final pH of 5.7 (+/-
0.19 SD). As pH is often a good predictor of iron speciation (83), we tested how the
treatments affected the relative proportions of Fe^{2+} and Fe^{3+}. We found the proportion
of more bioavailable Fe^{2+} to not significantly differ as a result of liming, shaking, nor
their interaction (lime main effect: $F_{1,9}=3.47$, $p=0.10$; shaking main effect: $F_{1,9}=3.00$,
$p=0.12$; lime-shaking interaction $F_{1,8}=0.73$, $p=0.42$; Fig.2), with Fe^{2+} making up 82% of
the total iron available on average across the treatments. Given that iron speciation
remained similar in all treatments, this indicates that the redox potential within the

microcosms did not change to become more anaerobic under static conditions (83).
Hence iron bioavailability was not significantly influenced by the different experimental
conditions and therefore iron limitation was unlikely to represent a significant driver for
siderophore production.
370

Figure 2 The final pH of microcosms containing river water and sediment after 28 days
incubation. We used a factorial design with limed and shaken treatments, each with
six replicates (each represented by a white circle). The starting pH was 5.8. The
significant effect of liming on pH (p<0.001) was increased through an interaction with
shaking (p<0.001).

P. aeruginosa populations incubated without lime had greater tolerance to copper
In order to test whether our river water and sediment samples selected for greater
metal resistance, we incubated the ancestral *P. aeruginosa* strain and final
populations in a medium containing a high concentration of copper (1 g/mL of copper
sulphate). We then compared the maximum optical density of each culture relative to

that of cultures grown without copper (w). Confirming that our samples contained toxic metals, the ancestral strain had a lower relative fitness (w) when grown with copper than all final populations (Dunnett's test: p=<0.013 for all contrasts; Fig. 3). Moreover, when comparing the effect of the different treatments on w, we found populations from the non-limed treatments to have greater relative fitness in a toxic copper environment than those from the limed treatments (liming main effect: $F_{1,21}$=4.44, p=0.047; Fig. 3), but shaking to not have an effect on relative fitness (shaking main effect: $F_{1,21}$=0.83, p=0.074; liming-shaking interaction: $F_{1,20}$=0.11, p=0.75). These results demonstrate that our river microcosms imposed strong selection on *P. aeruginosa* to evolve metal resistance, and that liming reduced this selection pressure.

Figure 3 The maximum optical density (OD_{600}) of *P. aeruginosa* populations after 18 hours incubation with toxic copper (1g/L copper sulphate) relative to their maximum OD_{600} when grown without copper (log w). Populations above the horizontal dashed line at 0 (log of 1) have a higher relative max OD_{600} when grown with copper, whereas those below the line have reduced maximum growth. The white box shows the

ancestral strain, whereas the grey boxes show populations incubated in microcosms
containing river water and sediment for 14 days. Circles show individual replicates
(n=6), and colours the different treatments.

Neither liming nor shaking affected P. aeruginosa density or siderophore production
Next, we tested the treatment effects on *P. aeruginosa* density and siderophore
production. The final density of *P. aeruginosa* after two weeks of evolution varied
substantially between samples ($1.1 \times 10^6 \pm 1.6 \times 10^6$ SD cfu/mL), but was not
significantly affected by liming, shaking, nor their interaction (liming main effect:
$F_{1,21}=1.96$, p=0.18; shaking main effect: $F_{1,21}=2.77$, p=0.11; liming-shaking interaction:
$F_{1,20}=0.70$, p=0.41). There was also no significant effect of pH on *P. aeruginosa*
density ($F_{1,22}=0.97$, p=0.36). Although pH can affect bacterial density (84), our finding
of no effect is consistent with previous results demonstrating that *P. aeruginosa*
densities are similar across an equivalent pH range as used here (85).

To test whether liming and shaking affected siderophore production, both total
siderophore production and the production of pyoverdine — the primary siderophore
produced by *P. aeruginosa* (56) — were measured for 24 clones per replicate (24 x
24 clones). Quantifying pyoverdine production in addition to total siderophores is
important, as it is a key virulence factor in *P. aeruginosa* but its production does not
necessarily correlate with that of other siderophores, such as pyochelin (62). We found
neither liming, shaking nor their interaction significantly affected mean total
siderophore production (liming main effect: $\chi^2=1.49$, d.f.=1, p=0.22; shaking main
effect: $\chi^2=0.49$, d.f.=1, p=0.48; liming-shaking interaction: $\chi^2=0.08$, d.f.=1, p=0.78) or
pyoverdine production (liming main effect: $\chi^2=0.56$, d.f.=1, p=0.46; shaking main
effect: $\chi^2=2.3$, d.f.=1, p=0.13; liming-shaking interaction: $\chi^2=1.14$, d.f.=1, p=0.29).
However, we note that there was a large variation in production between the 24 clones
used to represent each microcosm (mean production: total siderophores = 4.23;
pyoverdine = 766; $\sigma_{replicate}$: total siderophores = 1.94; pyoverdine = 69.3), and that two
pyoverdine values were significant outliers and consequently were removed from all
further analysis in order for model assumptions to be met (these were one from the
non-limed shaken treatment (pyoverdine production = 26.9, p<0.001) and one from

the limed-static treatment (pyoverdine production = 174, p<0.001); which were lower

than the pre-removed mean pyoverdine production of 710.6 and median of 789.8).

That siderophore production, which is regulated by iron availability and the presence

of toxic metals, did not significantly differ between treatments concurs with the non-

significant differences in Fe^{2+} availability between treatments. However, it is surprising

that siderophore production was not reduced by liming, given that *P. aeruginosa*

populations from the limed treatments were less tolerant to toxic copper. To explore

this further, we tested whether either total siderophore or pyoverdine production was

associated with copper tolerance, and found neither of them to be (Total siderophores:

$F_{1,20}=0.013$, p=0.91; Pyoverdine: $F_{1,20}=0.294$, p=0.59). This suggests other metal

resistance mechanisms, such as decreased outer membrane permeability and

increased induction of ATPase efflux transporters, could be responsible for the

increased copper tolerance of evolved populations (86). Our finding of no significant

differences in siderophore production contrasts with that of Hesse and co-workers

(18), who found that the addition of lime to soils collected in the near vicinity of our

locality significantly reduced community-wide siderophore production. This difference

is most likely due to shifts in siderophore production driven by changes in community

composition with liming selecting for non-producing isolates (18), whereas here we

solely focused on siderophore production by *P. aeruginosa*. This suggests that

although liming reduces community-wide siderophore production in metal-

contaminated acidic soils, this effect may not be seen in specific species. Interestingly,

P. aeruginosa has been proposed as a suitable siderophore-producing bacterium for

use in phytoremediation, which relies on the combined of use of microorganisms and

plants to aid toxic metal remediation (87, 88). It has been proposed that liming, by

reducing siderophore production, may hinder phytoremediation (18) as metal-uptake

by plants is often increased when metals are bound to bacterial siderophores. Given

that no significant effect of liming on siderophore production by *P. aeruginosa*, was

observed, we suggest that liming and *P. aeruginosa*-assisted phytoremediation could

be used simultaneously without compromise.

Virulence did not differ between treatments, but was positively associated with siderophore production

As we found a large variation in siderophore production, which is a known virulence factor in *P. aeruginosa* (64), we tested whether virulence, quantified using the *G. mellonella* infection assay, differed as a consequence of pyoverdine production, total siderophore production or treatment. Firstly, we tested whether *G. mellonella* larvae alive at the final time check (42 hours) had been injected with populations producing less total siderophores and pyoverdine compared with larvae that died before this point, and found that they were (total siderophores: χ^2=6.11, d.f.=1, p=0.013; pyoverdine: χ^2=6.98, d.f.=1, p=0.004). Next, we tested whether increased siderophore and pyoverdine production resulted in increased virulence. We found a significant positive association between virulence (mean time to death per population) and both total siderophore and pyoverdine production (total siderophores: $F_{1,22}$=8.9, p=0.007; Fig. 4A; pyoverdine: $F_{1,20}$=10.3, p=0.004; Fig. 4B), with every one unit increase in total siderophore production decreasing mean time to death by 1.42 hours (SE = 0.4745), and every one unit increase in pyoverdine production decreasing the mean time to death by 0.032 hours (SE = 0.0098). This positive association between siderophore production and virulence is supportive of previous work both in *G. mellonella* (37) and murine models (89). In addition to directly aiding iron uptake within a host, siderophores can increase virulence by triggering other virulence factors (90, 91). Finally, virulence was compared between treatments using survival curves (Fig. 4C). Virulence did not significantly differ as a function of treatment, with the credible intervals for liming, shaking and their interaction all crossing 1. No significant treatment effect on virulence is concurrent with the finding that the treatments did not significantly affect siderophore production. Finding virulence to not be significantly different between structured (static) and unstructured (shaking) environments contrasts with findings by Granato and co-workers (92), who found that pyoverdine-mediated virulence in *P. aeruginosa* was greater when grown in solid media than in liquid. The lack of changes detected in siderophore production and virulence between the experimental treatments might be due to the more subtle (and arguably more realistic) conditions under which spatial structure was varied in our study, as well as the presence of a resident microbial community.

Figure 4 Mean virulence of *Pseudomonas aeruginosa* evolved in metal-contaminated aquatic communities as a function of (A) mean total siderophore production and (B) mean pyoverdine production. Virulence was quantified using the *Galleria mellonella* infection model (n = 20 per replicate) and given as the mean time to death. Pyoverdine and total siderophore production were measured in standardised fluorescence units per OD_{600}. Individual circles show the mean production by 24 clones from each replicate. Colours and shapes represent different treatments: grey and □ = static, no lime, blue and + = static, limed, black and △ = shaken, no lime, and red and × = shaken, limed. Panel C shows the change in survival probability of larvae over time within each treatment. These do not significantly differ from one another. Shaded areas represent 95% confidence intervals.

Antibiotic resistance evolution

As metal pollution has been shown to co-select for antimicrobial resistance (38), we tested whether lime addition altered *P. aeruginosa* resistance to the clinically relevant antibiotics apramycin (15 µg/mL), cefotaxime (50 µg/mL) and trimethoprim (60 µg/mL) after evolution in metal-contaminated river sediments. Increased resistance was observed in all treatments (Fig. 5), with neither lime nor shaking affecting resistance to any of the antibiotics tested (apramycin: chi-squared=2.35 p=0.50 df=3; cefotaxime: chi-squared=2.98 p=0.40 df=3; trimethoprim: chi-squared=5.25 p=0.16 df=3; Fig. 5). Of note, one sample from the shaken, non-limed treatment consistently had the lowest resistance to all three antibiotics, with no isolates from this population being resistant to cefotaxime or trimethoprim, and fewer than 50% being resistant to apramycin. Our observation of rapid evolution of antibiotic resistance in the other replicates and treatments supports existing evidence that metal contamination can pose an important co-selective pressure for resistance (44, 49, 93), including in *P. aeruginosa* (42). That resistance did not differ significantly between treatments in our experiment demonstrates that liming to pH ~7 is not effective at remediating this co-selective effect, and neither was the loss of spatial structure via shaking. A plausible reason for this is that liming reduces metal bioavailability by precipitating ions from solution into the solid phase. This would mean cells in the sediment (the vast majority of the population) would still be exposed to metals where, although at a lower bioavailability, they can still be a cause of co-selection (48). This is supported by *P. aeruginosa* evolving greater tolerance to copper in the limed treatments. Although we did not determine the mechanistic basis of co-selection, we note that cross-resistance, co-resistance and co-regulation mechanisms have all been reported for Pseudomonads, and the altering of cellular targets is a mechanism commonly used by *P. aeruginosa* to tolerate metal, trimethoprim and beta-lactam antibiotics such as cefotaxime (42, 94). We are aware of a single study testing the effects of liming on antimicrobial resistance {Ramos, 1987 #4}. This study found liming decreased the susceptibility of Rhizobium species from soil to multiple antibiotics, and hypothesised this was due to a greater production of natural antibiotics at near-neutral pH selecting for resistance. Although we note that increasing soil pH will generally decreases the bioavailability of any metals present, the authors {Ramos, 1987 #4} stated that metal effects would not be operative in their study, suggesting no metal contamination was present.

Figure 5 The proportion of 24 *Pseudomonas aeruginosa* clones per replicate (n=6) resistant to (A) apramycin (15 µg/mL), (B) cefotaxime (50 µg/mL) and (C) trimethoprim (60 µg/mL) antibiotics. Clones were tested after two weeks of evolution in microcosms containing metal-contaminated river water and sediment while embedded in the resident microbial community. Circles show individual replicates; those with a red outline are from the same sample, which is the least resistant to all three antibiotics.

Conclusion

P. aeruginosa populations evolved metal resistance after two weeks, and liming reduced this effect. However liming and spatial structure (shaking) were observed to have little effect on *P. aeruginosa* pathogenic traits. Despite finding a positive association between siderophore production and virulence, neither siderophore production nor virulence systematically differed between treatments, suggesting that liming does not alter the effect of metals on siderophore-mediated virulence in *P. aeruginosa.* This finding also implies that concurrent use of liming and *P. aeruginosa*-assisted phytoremediation techniques is possible in scenarios were this bacterium can persist in a natural community. Moreover, we found *P. aeruginosa* rapidly evolved resistance to three clinically relevant antibiotics regardless of treatment. We therefore

show that a common metal remediation method did not reduce metal pollution-based co-selection for virulence or antibiotic resistance. Importantly, these findings further our understanding of how key determinants of pathogenicity evolve outside of the clinical setting, and further demonstrate that metal-polluted environments can select for them.

Funding

LL would like to thank NERC FRESH GW4 award no. NE/R011524/1, EH: UKRI Future Leaders Fellowship award MR/V022482/1, LN: NERC award NE/W006820/1, WG: NERC award NE/N019717/1, AB: NERC award NE/V012347/1, and MV: NERC award NE/T008083/1.

References

1. Sparks DL. Toxic metals in the environment: the role of surfaces. Elements. 2005;1(4):193-7.
2. Festa RA, Thiele DJ. Copper: an essential metal in biology. Current Biology. 2011;21(21):R877-R83.
3. Palmer LD, Skaar EP. Transition metals and virulence in bacteria. Annual review of genetics. 2016;50:67-91.
4. Kibria G, Hossain MM, Mallick D, Lau TC, Wu R. Monitoring of metal pollution in waterways across Bangladesh and ecological and public health implications of pollution. Chemosphere. 2016;165:1-9.
5. Paul D. Research on heavy metal pollution of river Ganga: A review. Annals of Agrarian Science. 2017;15(2):278-86.
6. Yunus FM, Khan S, Chowdhury P, Milton AH, Hussain S, Rahman M. A Review of Groundwater Arsenic Contamination in Bangladesh: The Millennium Development Goal Era and Beyond. Int J Environ Res Public Health. 2016;13:215.
7. Wu W, Zhang K, Jiang S, Liu D, Zhou H, Zhong R, et al. Association of co-exposure to heavy metals with renal function in a hypertensive population. Environment international. 2018;112:198-206.
8. Cui Y, Zhu Y-G, Zhai R, Huang Y, Qiu Y, Liang J. Exposure to metal mixtures and human health impacts in a contaminated area in Nanning, China. Environment International. 2005;31(6):784-90.
9. Tóth G, Hermann T, Da Silva M, Montanarella L. Heavy metals in agricultural soils of the European Union with implications for food safety. Environment international. 2016;88:299-309.

10.   Giller KE, Witter E, McGrath SP. Toxicity of heavy metals to microorganisms and
microbial processes in agricultural soils: a review. Soil biology and biochemistry. 1998;30(10-
11):1389-414.
11.   Wang Y, Shi J, Wang H, Lin Q, Chen X, Chen Y. The influence of soil heavy metals
pollution on soil microbial biomass, enzyme activity, and community composition near a
copper smelter. Ecotoxicology and environmental safety. 2007;67(1):75-81.
12.   Nwuche C, Ugoji EO. Effects of heavy metal pollution on the soil microbial activity.
International Journal of Environmental Science & Technology. 2008;5(3):409-14.
13.   Palansooriya KN, Shaheen SM, Chen SS, Tsang DC, Hashimoto Y, Hou D, et al. Soil
amendments for immobilization of potentially toxic elements in contaminated soils: A
critical review. Environment international. 2020;134:105046.
14.   Rogora M, Kamburska L, Mosello R, Tartari G. Lake Orta chemical status 25 years
after liming: problems solved and emerging critical issues. Journal of Limnology. 2016;75.
15.   Haimi J, Mätäsniemi L. Soil decomposer animal community in heavy-metal
contaminated coniferous forest with and without liming. European Journal of Soil Biology.
2002;38(2):131-6.
16.   Derome J, Saarsalmi A. The effect of liming and correction fertilisation on heavy
metal and macronutrient concentrations in soil solution in heavy-metal polluted scots pine
stands. Environmental Pollution. 1999;104(2):249-59.
17.   Chen Q, Luo Z, Hills C, Xue G, Tyrer M. Precipitation of heavy metals from
wastewater using simulated flue gas: sequent additions of fly ash, lime and carbon dioxide.
Water research. 2009;43(10):2605-14.
18.   Hesse E, Padfield D, Bayer F, Van Veen EM, Bryan CG, Buckling A. Anthropogenic
remediation of heavy metals selects against natural microbial remediation. Proceedings of
the Royal Society B. 2019;286(1905):20190804.
19.   Ruttens A, Adriaensen K, Meers E, De Vocht A, Geebelen W, Carleer R, et al. Long-
term sustainability of metal immobilization by soil amendments: cyclonic ashes versus lime
addition. Environmental Pollution. 2010;158(5):1428-34.
20.   Bolan NS, Adriano D, Mani P, Duraisamy A. Immobilization and phytoavailability of
cadmium in variable charge soils. II. Effect of lime addition. Plant and Soil. 2003;251(2):187-
98.
21.   Gunn JM, Kielstra BW, Szkokan-Emilson E. Catchment liming creates recolonization
opportunity for sensitive invertebrates in a smelter impacted landscape. Journal of
Limnology. 2016.
22.   Egeberg PK, Håkedal JT. The effect of river liming on the trace metal budgets of a
down stream lake. Water, Air, and Soil Pollution. 1998;104(1-2):57-75.
23.   Ramos MLG, Magalhães NF, Boddey RM. Native and inoculated rhizobia isolated
from field grown Phaseolus vulgaris: effects of liming an acid soil on antibiotic resistance.
Soil Biology and Biochemistry. 1987;19(2):179-85.
24.   Kelly JJ, Häggblom MM, Tate RL. Effects of heavy metal contamination and
remediation on soil microbial communities in the vicinity of a zinc smelter as indicated by
analysis of microbial community phospholipid fatty acid profiles. Biology and Fertility of
Soils. 2003;38(2):65-71.
25.   O'Brien S, Hodgson DJ, Buckling A. Social evolution of toxic metal bioremediation in
Pseudomonas aeruginosa. Proceedings of the Royal Society B: Biological Sciences.
2014;281(1787):20140858.

26.   Hesse E, O'Brien S, Tromas N, Bayer F, Luján AM, van Veen EM, et al. Ecological
selection of siderophore-producing microbial taxa in response to heavy metal
contamination. Ecology letters. 2018;21(1):117-27.
27.   Cuív PÓ, Clarke P, O'Connell M. Identification and characterization of an iron-
regulated gene, chtA, required for the utilization of the xenosiderophores aerobactin,
rhizobactin 1021 and schizokinen by Pseudomonas aeruginosa. Microbiology.
2006;152(4):945-54.
28.   Rajkumar M, Ae N, Prasad MNV, Freitas H. Potential of siderophore-producing
bacteria for improving heavy metal phytoextraction. Trends in biotechnology.
2010;28(3):142-9.
29.   Braud A, Geoffroy V, Hoegy F, Mislin G, Schalk I. The siderophores pyoverdine and
pyochelin are involved in Pseudomonas aeruginosa resistance against metals: another
biological function of these two siderophores. Environ Microbiol Rep. 2010;2:419-25.
30.   Kümmerli R, Jiricny N, Clarke L, West S, Griffin A. Phenotypic plasticity of a
cooperative behaviour in bacteria. Journal of evolutionary biology. 2009;22(3):589-98.
31.   Winkelmann G. Ecology of siderophores with special reference to the fungi.
Biometals. 2007;20(3-4):379.
32.   Cassat JE, Skaar EP. Iron in infection and immunity. Cell host & microbe.
2013;13(5):509-19.
33.   Hider RC, Kong X. Chemistry and biology of siderophores. Natural product reports.
2010;27(5):637-57.
34.   Skaar EP. The battle for iron between bacterial pathogens and their vertebrate hosts.
PLoS pathogens. 2010;6(8):e1000949.
35.   Harrison F, Browning LE, Vos M, Buckling A. Cooperation and virulence in acute
Pseudomonas aeruginosainfections. BMC biology. 2006;4(1):21.
36.   Braud A, Geoffroy V, Hoegy F, Mislin GL, Schalk IJ. Presence of the siderophores
pyoverdine and pyochelin in the extracellular medium reduces toxic metal accumulation in
Pseudomonas aeruginosa and increases bacterial metal tolerance. Environmental
Microbiology Reports. 2010;2(3):419-25.
37.   Lear L, Hesse E, Buckling A, Vos M. Copper selects for siderophore-mediated
virulence in Pseudomonas aeruginosa. 2021.
38.   Baker-Austin C, Wright MS, Stepanauskas R, McArthur J. Co-selection of antibiotic
and metal resistance. TRENDS in Microbiology. 2006;14(4):177.
39.   Hamzehpour MM, Pechere J-C, Plesiat P, Köhler T. OprK and OprM define two
genetically distinct multidrug efflux systems in Pseudomonas aeruginosa. Antimicrobial
agents and chemotherapy. 1995;39(11):2392-6.
40.   Teitzel GM, Parsek MR. Heavy Metal Resistance of Biofilm and Planktonic
Pseudomonas aeruginosa. Applied and Environmental Microbiology.
2003;69(4):2313-20.
41.   Köhler T, Kok M, Michea-Hamzehpour M, Plesiat P, Gotoh N, Nishino T, et al.
Multidrug efflux in intrinsic resistance to trimethoprim and sulfamethoxazole in
Pseudomonas aeruginosa. Antimicrobial Agents and Chemotherapy. 1996;40(10):2288-90.
42.   Sarma B, Acharya C, Joshi S. Pseudomonads: A versatile bacterial group exhibiting
dual resistance to metals and antibiotics. African Journal of Microbiology Research.
2010;4(25):2828-35.

43.   Perron K, Caille O, Rossier C, Van Delden C, Dumas J-L, Köhler T. CzcR-CzcS, a two-
component system involved in heavy metal and carbapenem resistance in Pseudomonas
aeruginosa. Journal of Biological Chemistry. 2004;279(10):8761-8.
44.   Seiler C, Berendonk T. Heavy metal driven co-selection of antibiotic resistance in soil
and water bodies impacted by agriculture and aquaculture. Frontiers in Microbiology.
2012;3(399).
45.   Ghosh A, Singh A, Ramteke P, Singh V. Characterization of large plasmids encoding
resistance to toxic heavy metals in Salmonella abortus equi. Biochemical and biophysical
research communications. 2000;272(1):6-11.
46.   Gullberg E, Albrecht LM, Karlsson C, Sandegren L, Andersson DI. Selection of a
multidrug resistance plasmid by sublethal levels of antibiotics and heavy metals. MBio.
2014;5(5):e01918-14.
47.   Allen D, Austin B, Colwell R. Antibiotic resistance patterns of metal-tolerant bacteria
isolated from an estuary. Antimicrobial Agents and Chemotherapy. 1977;12(4):545-7.
48.   Dickinson AW, Power A, Hansen M, Brandt K, Piliposian G, Appleby P, et al. Heavy
metal pollution and co-selection for antibiotic resistance: A microbial palaeontology
approach. Environment international. 2019;132:105117.
49.   Stepanauskas R, Glenn TC, Jagoe CH, Tuckfield RC, Lindell AH, King CJ, et al.
Coselection for microbial resistance to metals and antibiotics in freshwater microcosms.
Environmental Microbiology. 2006;8(9):1510-4.
50.   Vos M, Sibleyras L, Lo LK, Hesse E, Gaze WH, Klümper U. Zinc can counteract
selection for ciprofloxacin resistance. bioRxiv. 2019:780981.
51.   Pirrie D, Power MR, Rollinson G, Camm GS, Hughes SH, Butcher AR, et al. The spatial
distribution and source of arsenic, copper, tin and zinc within the surface sediments of the
Fal Estuary, Cornwall, UK. Sedimentology. 2003;50(3):579-95.
52.   Rice LB. Federal funding for the study of antimicrobial resistance in nosocomial
pathogens: no ESKAPE. The University of Chicago Press; 2008.
53.   Pachori P, Gothalwal R, Gandhi P. Emergence of antibiotic resistance Pseudomonas
aeruginosa in intensive care unit; a critical review. Genes & diseases. 2019;6(2):109-19.
54.   Rutherford V, Yom K, Ozer EA, Pura O, Hughes A, Murphy KR, et al. Environmental
reservoirs for exoS+ and exoU+ strains of Pseudomonas aeruginosa. Environmental
microbiology reports. 2018;10(4):485-92.
55.   O'Brien S, Hesse E, Luján A, Hodgson DJ, Gardner A, Buckling A. No effect of
intraspecific relatedness on public goods cooperation in a complex community. Evolution.
2018;72(5):1165-73.
56.   Granato E, Harrison F, Kümmerli R, Ross-Gillespie A. Do Bacterial "Virulence Factors"
Always Increase Virulence. A Meta-Analysis of Pyoverdine Production in Pseudomonas
aeruginosa As a Test Case Front Microbiol. 2016;7:1952.
57.   Kümmerli R, Jiricny N, Clarke LS, West SA, Griffin AS. Phenotypic plasticity of a
cooperative behaviour in bacteria. Journal of Evolutionary Biology. 2009;22(3):589-98.
58.   Braud A, Jézéquel K, Bazot S, Lebeau T. Enhanced phytoextraction of an agricultural
Cr-and Pb-contaminated soil by bioaugmentation with siderophore-producing bacteria.
Chemosphere. 2009;74(2):280-6.
59.   Guerrieri CG, Pereira MF, Galdino ACM, dos Santos ALS, Elias WP, Schuenck RP, et al.
Typical and atypical enteroaggregative Escherichia coli are both virulent in the Galleria
mellonella model. Frontiers in microbiology. 2019;10.

60.   Hernandez RJ, Hesse E, Dowling AJ, Coyle NM, Feil EJ, Gaze WH, et al. Using the wax
moth larva Galleria mellonella infection model to detect emerging bacterial pathogens.
PeerJ. 2019;7:e6150-e.
61.   Schwyn B, Neilands JB. Universal chemical assay for the detection and determination
of siderophores. Analytical biochemistry. 1987;160(1):47-56.
62.   Dumas Z, Ross-Gillespie A, Kümmerli R. Switching between apparently redundant
iron-uptake mechanisms benefits bacteria in changeable environments. Proceedings of the
Royal Society B: Biological Sciences. 2013;280(1764):20131055.
63.   Kümmerli R, Griffin AS, West SA, Buckling A, Harrison F. Viscous medium promotes
cooperation in the pathogenic bacterium Pseudomonas aeruginosa. Proceedings of the
Royal Society B: Biological Sciences. 2009;276(1672):3531-8.
64.   Buckling A, Harrison F, Vos M, Brockhurst MA, Gardner A, West SA, et al.
Siderophore-mediated cooperation and virulence in Pseudomonas aeruginosa. FEMS
microbiology ecology. 2007;62(2):135-41.
65.   Ross-Gillespie A, Gardner A, West SA, Griffin AS. Frequency dependence and
cooperation: theory and a test with bacteria. The American Naturalist. 2007;170(3):331-42.
66.   Cordero OX, Ventouras L-A, DeLong EF, Polz MF. Public good dynamics drive
evolution of iron acquisition strategies in natural bacterioplankton populations. Proceedings
of the National Academy of Sciences. 2012;109(49):20059-64.
67.   World Health Organization. Critically important antimicrobials for human medicine.
2019.
68.   Kang AD, Smith KP, Eliopoulos GM, Berg AH, McCoy C, Kirby JE. In vitro apramycin
activity against multidrug-resistant Acinetobacter baumannii and Pseudomonas aeruginosa.
Diagnostic microbiology and infectious disease. 2017;88(2):188-91.
69.   O'Brien S, Luján AM, Paterson S, Cant MA, Buckling A. Adaptation to public goods
cheats in Pseudomonas aeruginosa. Proceedings of the Royal Society B: Biological Sciences.
2017;284(1859):20171089.
70.   Lovley DR, Phillips EJ. Rapid assay for microbially reducible ferric iron in aquatic
sediments. Appl Environ Microbiol. 1987;53(7):1536-40.
71.   Lovley DR, Phillips EJ. Organic matter mineralization with reduction of ferric iron in
anaerobic sediments. Appl Environ Microbiol. 1986;51(4):683-9.
72.   UKWIR. The Chemical Investigations Programme Phase 2, 2015–2020 – Initial
Findings. 2018;1–4.
73.   Brun L, Maillet J, Richarte J, Herrmann P, Remy J. Relationships between extractable
copper, soil properties and copper uptake by wild plants in vineyard soils. Environmental
pollution. 1998;102(2-3):151-61.
74.   Schwyn B, Neilands J. Universal chemical assay for the detection and determination
of siderophores. Analytical biochemistry. 1987;160(1):47-56.
75.   Harrison F, Buckling A. Hypermutability Impedes Cooperation in Pathogenic Bacteria.
Current Biology. 2005;15:1968-71.
76.   Signorell A, Aho K, Alfons A, Anderegg N, Aragon T, Arppe A, et al. DescTools: Tools
for descriptive statistics. R package version 099. 2019;28:17.
77.   Douglas Bates MM, Ben Bolker, Steve Walker. Fitting Linear Mixed-Effects Models
Using lme4. Journal of Statistical Software. 2015;67(1):1-48.
78.   Hartig F. DHARMa: Residual Diagnostics for Hierarchical (Multi-Level / Mixed)
Regression Models. R package version 042 2021.
79.   Komsta L, Komsta ML. The outliers Package. 2007.

80.   Brilleman SL, Elci EM, Novik JB, Wolfe R. Bayesian survival analysis using the
rstanarm R package. arXiv preprint arXiv:200209633. 2020.
81.   Kay M. tidybayes: Tidy data and geoms for Bayesian models. R package version.
2019;1(0).
82.   R Core Team. R: A language and environment for statistical computing, . R
Foundation for Statistical Computing.
83.   Gotoh S, Patrick Jr W. Transformation of iron in a waterlogged soil as influenced by
redox potential and pH. Soil Science Society of America Journal. 1974;38(1):66-71.
84.   Dennis PG, Hirsch PR, Smith SJ, Taylor RG, Valsami-Jones E, Miller AJ. Linking
rhizoplane pH and bacterial density at the microhabitat scale. Journal of Microbiological
Methods. 2009;76(1):101-4.
85.   Moriarty T, Elborn J, Tunney M. Effect of pH on the antimicrobial susceptibility of
planktonic and biofilm-grown clinical Pseudomonas aeruginosa isolates. British journal of
biomedical science. 2007;64(3):101-4.
86.   Teitzel GM, Geddie A, Susan K, Kirisits MJ, Whiteley M, Parsek MR. Survival and
growth in the presence of elevated copper: transcriptional profiling of copper-stressed
Pseudomonas aeruginosa. Journal of bacteriology. 2006;188(20):7242.
87.   Rajkumar M, Sandhya S, Prasad MNV, Freitas H. Perspectives of plant-associated
microbes in heavy metal phytoremediation. Biotechnology advances. 2012;30(6):1562-74.
88.   Sinha S, Mukherjee SK. Cadmium–induced siderophore production by a high Cd-
resistant bacterial strain relieved Cd toxicity in plants through root colonization. Current
microbiology. 2008;56(1):55-60.
89.   Kang D, Revtovich AV, Chen Q, Shah KN, Cannon CL, Kirienko NV. Pyoverdine-
dependent virulence of Pseudomonas aeruginosa isolates from cystic fibrosis patients.
Frontiers in microbiology. 2019;10:2048.
90.   Lopez-Medina E, Fan D, Coughlin LA, Ho EX, Lamont IL, Reimmann C, et al. Candida
albicans inhibits Pseudomonas aeruginosa virulence through suppression of pyochelin and
pyoverdine biosynthesis. PLoS Pathog. 2015;11(8):e1005129.
91.   Lamont IL, Beare PA, Ochsner U, Vasil AI, Vasil ML. Siderophore-mediated signaling
regulates virulence factor production in Pseudomonas aeruginosa. Proceedings of the
National Academy of Sciences. 2002;99(10):7072-7.
92.   Granato ET, Ziegenhain C, Marvig RL, Kümmerli R. Low spatial structure and selection
against secreted virulence factors attenuates pathogenicity in Pseudomonas aeruginosa.
The ISME journal. 2018;12(12):2907-18.
93.   Song J, Rensing C, Holm PE, Virta M, Brandt KK. Comparison of metals and
tetracycline as selective agents for development of tetracycline resistant bacterial
communities in agricultural soil. Environmental science & technology. 2017;51(5):3040-7.
94.   Conejo MC, García I, Martínez-Martínez L, Picabea L, Pascual Á. Zinc eluted from
siliconized latex urinary catheters decreases OprD expression, causing carbapenem
resistance in Pseudomonas aeruginosa. Antimicrobial agents and chemotherapy.
2003;47(7):2313-5.
827