Generalization of contextual fear is sex-specifically affected by high salt intake

Short title: Salt sex-specifically influences fear generalization

Jasmin N. Beaver[1,2], Brady L. Weber[1,2], Matthew T. Ford[1], Anna E. Anello[1,2], Kaden M. Ruffin[1],

Sarah K. Kassis[1,2], T. Lee Gilman[1,2,3]*

[1]Department of Psychological Sciences, Kent State University, Kent, Ohio, United States of

America

[2]Brain Health Research Institute, Kent State University, Kent, Ohio, United States of America

[3]Healthy Communities Research Institute, Kent State University, Kent, Ohio, United States of

America

*Corresponding Author

Email: lgilman1@kent.edu (TLG)

Abstract: 242 words (max 300)

Abstract

A hallmark symptom of many anxiety disorders, and multiple neuropsychiatric disorders more broadly, is generalization of fearful responses to non-fearful stimuli. Anxiety disorders are often comorbid with cardiovascular diseases. One established, and modifiable, risk factor for cardiovascular diseases is salt intake. Yet, investigations into how excess salt consumption affects anxiety-relevant behaviors remains little explored. Moreover, no studies have yet assessed how high salt intake influences generalization of fear. Here, we used adult C57BL/6J mice of both sexes to evaluate the influence of two or six weeks of high salt consumption (4.0% NaCl), compared to controls (0.4% NaCl), on contextual fear acquisition, expression, and generalization. Further, we measured osmotic and physiological stress by quantifying serum osmolality and corticosterone levels, respectively. Consuming excess salt did not influence contextual fear acquisition nor discrimination between the context used for training and a novel, neutral context when training occurred 48 prior to testing. However, when a four week delay between training and testing was employed to induce natural fear generalization processes, we found that high salt intake selectively increases contextual fear generalization in females, but the same diet reduces contextual fear generalization in males. These sex-specific effects were independent of any changes in serum osmolality nor corticosterone levels, suggesting the behavioral shifts are a consequence of more subtle, neurophysiologic changes. This is the first evidence of salt consumption influencing contextual fear generalization, and adds information about sex-specific effects of salt that are largely missing from current literature.

Introduction

Excessive dietary salt intake occurs across nations and cultures [1,2]. While salt's practical uses include food preservation and flavor enhancement, consumption of too much salt has been associated with increased risk for cardiovascular diseases [3–6]. Indeed, average estimates of global salt intake indicate people consume 3,950 mg of sodium (the component of NaCl that has been associated with multiple negative health outcomes) [3,7,8]. When put in the context that, according to the World Health Organization, adults should consume less than 2,000 mg of sodium, this overconsumption of salt becomes quantifiably apparent [9]. It is estimated that reducing consumption of salt can attenuate risk for cardiovascular disease by ~20% [5,10], meaning dietary salt consumption has a profound physiological influence.

Globally, cardiovascular diseases and neuropsychiatric disorders, particularly disorders involving anxiety symptoms, are often comorbid [11–15] (see also reviews [16,17]). Increasing literature suggests dietary components such as excess sugar and/or fat are associated with both metabolic conditions like insulin resistance and type II diabetes, as well as with neuropsychiatric conditions (e.g., depression) and diseases (e.g., Alzheimer's) [18–22]. Trait anxiety appears largely unaffected by excess salt consumption [23–27] (reviewed in [28]). Far less understood is how excess salt intake might affect neuropsychiatric symptoms like generalized (i.e., non-specific, state) anxiety, though some researchers have begun to recognize this possibility [29,30]. We hypothesized that excessive consumption of dietary salt would enhance fear generalization in rodents, a possibility that has not yet been explored.

Fear generalization is an established neurobehavioral phenomenon that occurs across species, and studies are making considerable progress in identifying brain regions critical to this process [31–34]. These include specific subregions of the cingulate and prefrontal cortices, hippocampus, amygdala, and brainstem nuclei (e.g., locus coeruleus). Growing evidence indicates the hippocampus and amygdala are affected by excess salt consumption in rodents

[24,26,35–41], including: reductions in nicotinic receptor levels [35]; increased markers of

oxidative stress [24,26,37,41], neuronal activation [42], and neuroinflammation [42]; reduced

long-term potentiation [40]; and increased microvasculature leakiness [36]. Other brain regions

remain underexplored. This evidence, combined with human evidence of comorbidity between

anxiety disorders and cardiovascular diseases, supports our hypothesis – at least based upon

male data – that fear generalization is augmented by high salt intake. Unfortunately, both the

scant literature on behavioral effects of increased salt intake, and the larger body of work on the

neurocircuitry of fear generalization, have focused almost exclusively on male rodents. Here,

we included both sexes in our studies, but given the current state of the literature we could not

formulate data-based *a priori* hypotheses specifically regarding how female fear generalization

behavior might differ from male fear generalization under conditions of excess salt intake.

Materials and methods

Animals

Mice of both sexes on a C57BL/6J background were purchased from Jackson Laboratory (Bar Harbor, ME) and bred in-house. All mice were at least 9 weeks old prior to commencing experiments. At the start of diet manipulation, mice were singly housed to facilitate accurate quantification of diet and water consumption relative to the mouse's body weight. Prior to experiments, all mouse cages contained Nestlets (Ancare, Bellmore, NY) and huts; at the time of singly housing for diet manipulation, two additional pieces of enrichment (e.g., swing, wood block, etc.) were added to each cage. Cages contained 7090 Teklad Sani-chip bedding (Envigo, East Millstone, NJ) and were housed in a 12:12 light/dark cycle room, with lights on at 07:00, and temperature maintained at 22 ± 2°C.

Prior to diet manipulation, mice were fed LabDiet 5001 rodent laboratory chow (LabDiet, Brentwood, MO) *ad libitum*. At the time of diet manipulation, mice were pseudorandomly assigned to either a control diet (0.4% NaCl w/w; D17012, Research Diets, Inc., New Brunswick, NJ) or a high salt diet (4.0% NaCl; D17013; Fig. 1A). These diets were available *ad libitum*, and water was always available *ad libitum*. Twice weekly, measurements of diet and water consumption plus body weight were taken. Refer to Supporting Information for statistics and graphs on average water consumed per day (S15 and S19 Tables; S9 and S13 Figs.); average food consumed per day (S16 and S20 Tables; S10 and S13 Figs.); weekly body weight changes (S17 and S21 Tables; S11 and S14 Figs.); raw body weights (S18 and S22 Tables; S12 and S14 Figs.); average NaCl consumed per day (S23 and S28 Tables; S15 and S20 Figs.); average NaCl consumed as percentage of body weight (S24 and S29 Tables; S16 and S20 Figs.); weekly ratio between water:NaCl consumption (S25 and S30 Tables; S17 and S21

Figs.); average kcal consumed per day (S26 and S31 Tables; S18 and S22 Figs.); and average

kcal consumed as percentage of body weight (S27 and S32 Tables; S19 and S22 Figs.).

Fig. 1. ***Study timeline and fear conditioning procedures.***

A) Mice were assigned to either a control (0.4% NaCl) or high salt (4.0% NaCl) diet for two or

six weeks. Experiment 1 involved diet manipulation for two weeks, at which time mice

underwent context fear training, followed 48 h later by context fear testing. Experiment 2 (grey

shading) involved diet manipulation for six weeks, at which time mice underwent context fear

training, followed 48 h later by context fear testing. Experiment 3 involved diet manipulation for

two weeks, at which time mice underwent context fear training, followed four weeks later (while

continuing the same diet manipulation) by context fear testing. B) Mice trained with a mild foot

shock in the Training Context were tested 48 h (Experiments 1, 2) or four weeks (Experiment 3)

later in either the Training Context or the Neutral Context. The Training Context included a

metal grid floor, visible illumination, patterned background, and 70% ethanol scent. The Neutral

Context included a smooth acrylic floor, infrared illumination, no background, and Windex scent.

C) Control no shock mice were exposed to the Training Context, but were never administered a

foot shock. These mice were tested at the same timelines for Experiments 1-3 in the Training

Context, to assess for any baseline influences of diet consumption on fear behavior.

Diet manipulations lasted for two weeks (Experiment 1; Fig. 1A) or six weeks (Experiments 2

and 3; Fig. 1A). All procedures were approved by the Kent State University Institutional Animal

Care and Use Committee (protocol 536 LG 22-14), and adhered to the National Research

Council's Guide for the Care and Use of Laboratory Animals, 8th Ed. [43]. Every measure was

taken to minimize suffering.

Experimental timelines

Experimental timelines are graphically illustrated in Fig. 1A. For Experiment 1, context fear training occurred after two weeks of diet manipulation, and context fear testing occurred 48 h after training. For Experiment 2, training occurred after six weeks of diet manipulation, and testing occurred 48 h after training. For Experiment 3, training occurred after two weeks of diet manipulation, and testing occurred four weeks after training (a standard time frame for natural development of fear generalization in mice [44]).

Context fear conditioning

Females and males always underwent behavior separately. Fear conditioning took place in identical Coulbourn Instruments chambers (7 in D × 7 in W × 12 in H; Allentown, PA) composed of two opposing aluminum walls and two Plexiglas walls surrounding a stainless-steel shock grid floor. Each chamber was located within its own sound-attenuating enclosure, and had a camera mounted to the top for real-time measurements of freezing behavior (absence of all movement except that necessary for breathing) using FreezeFrame (v. 5.201, Coulburn Instruments).

Context fear training

The Training Context (Fig. 1B) consisted of the shock grid floor, a patterned background, visible illumination, and scent cue (70% ethanol). Training occurred at either the two week or six week time point (Fig. 1A). Training involved exposure to five, 1 s, 0.8 mA scrambled foot shocks pseudorandomly administered during a 6 min training session at 137, 186, 229, 285, and 324 s. Training with five foot shocks is optimal for eliciting time-dependent contextual fear generalization in rodents [44–46]. Freezing behavior was quantified as the average percent freezing across the first six 5 s bins (i.e., 30 s) that occurred immediately after the 5 s bin containing the foot shock.

Context fear testing

Testing occurred at either the two (Experiment 1) or six week (Experiments 2 and 3) time point

(Fig. 1A). Mice were tested either in the Training Context, or in a Neutral Context which

consisted of a smooth floor, no background, only infrared illumination, and a different scent cue

(Windex®, SC Johnson, Racine, WI; Fig. 1B). All mice were only ever tested in a single

context, and testing lasted for 10 minutes, with the average percentage freezing occurring

during minutes two through six being quantified and analyzed [46]. Foot shocks were never

administered during testing.

No shock controls

160   Control 'no shock' mice were run concurrently in every cohort of animals (Fig. 1C) for every

161   Experiment. These no shock mice were exposed to the Training Context for 'training', but no

162   foot shock was ever administered. Likewise, control no shock mice were tested in the Training

163   Context, to evaluate how diet manipulation and the Experiment timelines influenced freezing

164   behavior in the absence of any contextual fear conditioning.

Serum measurements

Forty-eight hours after testing for context fear expression, mice were briefly anesthetized with

isoflurane then rapidly decapitated for collection of trunk blood. Trunk blood was allowed to clot

at room temperature for 30 min, then was spun at 3500 rpm for 1 h at 4°C. Serum was then

removed from tubes and aliquoted into two vials – one for measuring serum osmolality, the

other for measuring serum corticosterone. Serum osmolality was measured immediately using

an Osmo1 single-sample micro-osmometer (Advanced Instruments, Norwood, MA). Prior to

every serum batch, a Clinitrol™ 290 reference solution (Advanced Instruments) from a new

ampule was measured to confirm accurate readings. Samples for serum corticosterone were

stored immediately at −80°C until analysis with a corticosterone ELISA kit (Enzo Life Sciences,

Inc., Farmingdale, NY) using their small volume protocol. Serum corticosterone levels were log-

transformed to account for positive skewness [27,47,48]. Log-transformed serum corticosterone

levels are, for brevity, hereafter referred to as 'cort' levels.

Statistical analyses

Data were analyzed with GraphPad Prism 9.4.0 (GraphPad Software, San Diego, CA) and

IBM SPSS Statistics 28.0.0.0 (IBM, Armonk, NY). Our *a priori* significance threshold was set at

$p<0.05$. Analyses were made within each Experiment. Repeated measures data (diet/water

consumption, salt or kcal calculations, body weight, context fear training) were analyzed using

3-way repeated measures ANOVAs within individual sexes ([repeated measure, e.g., body

weight] × diet × context) and pairwise comparisons with Bonferroni correction. Greenhouse-

Geisser corrections were employed for within-subjects analyses. Measurements of contextual

fear expression, serum osmolality, and serum corticosterone across contexts were analyzed

with a 3-way ANOVA (diet × sex × context) and Holm-Šídák's post-hocs. Control no shock fear

behavior data were analyzed separately, given these were purposely included as negative

controls and not intended to be analyzed in comparison to experimental groups. These no

shock control data were analyzed on their own with 3-way ANOVAs (diet × sex × context) and

Holm-Šídák's post-hocs, or with repeated measures 3-way ANOVAs within each sex ([repeated

measure, e.g., body weight] × diet × sex) plus pairwise comparisons with Bonferroni correction,

and also using Greenhouse-Geisser corrections for within-subjects analyses. All ANOVA

statistics are reported in Supporting Information, in S1-S32 Tables. The criterion to exclude

outliers was *a priori* assigned as being >5 standard deviations ± mean. Numbers of mice per

sex/shock/diet/context condition varied, and exact numbers are present in the respective figure

legends for each group of data. Broadly: no shock fear behavior n=7-9; shock fear behavior

n=7-10; shock serum osmolality n=8-9; shock serum corticosterone n=7-9. Details of outliers

that were excluded (training baseline, n=1; no shock testing, n=1; shock testing, n=4; osmolality,

n=1; log-transformed serum corticosterone, n=2; water:NaCl consumption ratio across weeks,

n=8) are provided in S33 Table. Graphs were created with GraphPad Prism, and show the

mean ± the 95% confidence interval (CI), with non-repeated measures graphs also showing

individual data points.

Results

No shock controls were built into every Experiment to evaluate how consumption of a high salt

(4.0% NaCl) diet, relative to the control (0.4% NaCl) diet, might affect fear behavior in the

absence of contextual fear conditioning. For 'training' of no shock controls, no three-way

interactions of time × sex × diet were observed in any of the three experiments, nor were two-

way interactions of sex × diet nor time × diet (S1 Table). The only two-way interaction

evaluated that reached the significance threshold was for a time × sex interaction in training for

Experiment 3 (S1 Table), indicating that during 'training' one sex behaved differently over time

than the other. In Experiments 1 and 2, no significant main effects of sex nor diet were

detected, though both had significant main effects of time (S1 Table), meaning behavior of mice

changed over the course of 'training'. This is to be expected when mice are initially exploring a

new environment, then become complacent in the absence of any apparent threat. Pairwise

comparisons indicated a few time points at which significant differences in freezing behavior

between males consuming control and high salt diets were detected (S1A Fig., left and right

panels), but these did not exhibit a consistent pattern, and were absent in Experiment 2 (S1A

Fig., middle panel), suggesting they were not meaningful.

As with 'training', testing of no shock control mice revealed no significant interactions involving

diet, and no main effect of diet (S2 Table), illustrating that diet did not impact freezing behavior

in the absence of any fear conditioning. A significant interaction between Experiment × sex was

observed, but Holm-Šídák's post-hoc tests revealed no significant differences across the sexes

within any Experiment, nor within the sexes across any Experiments (S1B Fig.), indicating at a

granular level, there were no Experiment-specific sex differences.

Acquisition of contextual fear resulted in no significant three- nor two-way interactions in any

of the three Experiments within either sex (S3-S5 Tables). Likewise, no main effects of Context

nor diet were detected (S3-S5 Tables), meaning diet condition did not affect fear learning, and

there were no inherent differences in mice to be tested in one Context or the other. In fact, the

only significant main effect was that of time, detected across all three Experiments (S3-S5

Tables). This illustrates all mice that underwent contextual fear conditioning with foot shocks

present during training exhibited increasing levels of freezing (i.e., fear behavior) over the

course of the training session (Fig. 2). Pairwise comparisons between male mice to be tested in

the Neutral Context indicated that, in Experiment 2, mice consuming high salt exhibited elevated

freezing levels following the second and third of the five training foot shocks, relative to mice

consuming control diet (Fig. 2B, right panel). All other acquisition curves lacked significant

differences across diets and future testing contexts (Fig. 2), indicating that a high salt diet does

not impact acquisition of contextual fear after two (Experiments 1, 3) or six (Experiment 2)

weeks of consumption.

Fig. 2. ***Context fear training across Experiments***.

Mice assigned to 0.4% NaCl represented by blue symbols, mice assigned to 4.0% NaCl

represented by red symbols; mice to be tested in Training Context represented by squares and

solid lines, mice to be tested in Neutral Context represented by circles and dotted lines. All

mice were trained in the Training Context with five, pseudorandomized mild (1 s, 0.8 mA) foot

shocks during a 6 min training session. Percent freezing during training of mice in A)

Experiment 1, B) Experiment 2 (grey shading), and C) Experiment 3. Baseline percent freezing

averaged across the first two minutes of the training session and plotted along x-axis as 0 time

point. Average percent freezing for each 30 sec period following each of the five mild foot

shocks are plotted along x-axis as Post-shock Periods 1-5. Experiment 1: 0.4% NaCl females

Training Context, n=7; 0.4% NaCl females Neutral Context, n=9; 4.0% NaCl females Training

Context, n=8; 4.0% NaCl females Neutral Context, n=9; 0.4% NaCl males Training Context,

n=8; 0.4% NaCl males Neutral Context, n=9; 4.0% NaCl males Training Context, n=7; 4.0%

NaCl males Neutral Context, n=8. Experiment 2: 0.4% NaCl females Training Context, n=9;

0.4% NaCl females Neutral Context, n=8; 4.0% NaCl females Training Context, n=8; 4.0% NaCl

females Neutral Context, n=8; 0.4% NaCl males Training Context, n=9; 0.4% NaCl males

Neutral Context, n=8; 4.0% NaCl males Training Context, n=8; 4.0% NaCl males Neutral

Context, n=10. Experiment 3: 0.4% NaCl females Training Context, n=8; 0.4% NaCl females

Neutral Context, n=9; 4.0% NaCl females Training Context, n=9; 4.0% NaCl females Neutral

Context, n=8; 0.4% NaCl males Training Context, n=7; 0.4% NaCl males Neutral Context, n=8;

4.0% NaCl males Training Context, n=8; 4.0% NaCl males Neutral Context, n=8. Data are

graphed as mean ± 95% confidence interval. [+]indicates $p<0.05$ difference between males to be

tested in Neutral Context and consuming 0.4% and 4.0% NaCl diets at indicated time points.

Unlike context fear training, where diet was without significant effect on acquisition, high salt

diet consumption did significantly affect context fear expression. Moreover, these dietary effects

were often sex-selective. Across all three Experiments, there were no significant three-way

interactions, nor were there any significant diet × context interactions (S7 Table). However,

sex-specific fear behaviors between contexts were significant for Experiments 2 and 3, and fear

behaviors across diet and sex combinations were significant for Experiment 3 (S7 Table).

Further, diet alone had a significant effect in Experiment 2, and sex and context independently

had significant effects on fear behavior in Experiment 1 (S7 Table). In other words, diet was

without a significant effect in Experiment 1, but diet did influence the outcomes of Experiments 2

and 3, and for all three Experiments, sex had a significant – and mostly interactional – impact as

well.

Holm-Šídák's post-hoc testing helped indicate directionality of high salt consumption's effects

on contextual fear expression and generalization across sexes for the three different Experiment

timelines employed. Experiment 1 involved training mice after two weeks of diet manipulation,

and testing 48 h after training. The expected low (average ~20% or less) freezing levels in the

Neutral Context, with higher (average ~40% or more) freezing levels in the Training Context,

were observed across nearly all the sex and diet combinations (Fig. 3A). A non-significant trend

was noted for contextual fear expression in the Training Context between females and males

consuming the high salt diet (p=0.0542; Fig. 3A). Excepting this, Experiment 1 suggests that

contextual fear expression after two weeks of high salt diet consumption is unaffected across

contexts and sexes.

Fig. 3. *Context fear expression across contexts and Experiments*.

Mice assigned to 0.4% NaCl represented by blue symbols, mice assigned to 4.0% NaCl

represented by red symbols; mice tested in Training Context represented by squares and solid

lines, mice tested in Neutral Context represented by circles and dotted lines. Percent freezing

during minutes two through six of the 10 min testing session are graphed for all mice. Testing

occurred A) 48 h after training in Experiment 1, during which mice underwent two weeks of diet

manipulation; B) 48 h after training in Experiment 2 (grey shading), during which mice

underwent six weeks of diet manipulation; and C) four weeks after training in Experiment 3,

during which mice underwent six total weeks of diet manipulation (training occurred after two

weeks of diet manipulation). Experiment 1: 0.4% NaCl females Training Context, n=7; 0.4%

NaCl females Neutral Context, n=9; 4.0% NaCl females Training Context, n=8; 4.0% NaCl

females Neutral Context, n=9; 0.4% NaCl males Training Context, n=8; 0.4% NaCl males

Neutral Context, n=9; 4.0% NaCl males Training Context, n=7; 4.0% NaCl males Neutral

Context, n=8. Experiment 2: 0.4% NaCl females Training Context, n=9; 0.4% NaCl females

Neutral Context, n=8; 4.0% NaCl females Training Context, n=8; 4.0% NaCl females Neutral

Context, n=8; 0.4% NaCl males Training Context, n=9; 0.4% NaCl males Neutral Context, n=8;

4.0% NaCl males Training Context, n=8; 4.0% NaCl males Neutral Context, n=10. Experiment

 3: 0.4% NaCl females Training Context, n=8; 0.4% NaCl females Neutral Context, n=9; 4.0%

NaCl females Training Context, n=9; 4.0% NaCl females Neutral Context, n=8; 0.4% NaCl

males Training Context, n=7; 0.4% NaCl males Neutral Context, n=8; 4.0% NaCl males Training

Context, n=8; 4.0% NaCl males Neutral Context, n=8. Data are graphed as mean ± 95%

confidence interval. *p<0.05, **p<0.01, ***p<0.001 indicate difference between Training Context

and Testing Context within same sex and diet. +++indicates p<0.001 difference between

females and males on same diet and tested in same context.

Experiment 2 involved training mice after six weeks of diet manipulation, then testing 48 h

following training. Similar to Experiment 1, contextual fear expression was largely unaffected by

diet or sex, with one exception. Male mice consuming the control diet exhibited reduced

contextual fear expression in the Training Context as compared to female mice consuming the

control diet (Fig. 3B); this sex difference in Training Context fear expression was absent in mice

of both sexes consuming the high salt diet. As anticipated when testing for contextual fear

expression relatively soon after training (in this case, 48 h), freezing behavior averaging ≤ 20%

indicated that mice did not exhibit fear behavior in the Neutral Context (see S1 Fig. for no shock

comparisons).

Experiment 3 evaluated how high salt intake affects contextual fear generalization by training

mice after two weeks of diet manipulation, then testing for contextual fear expression four weeks

later (i.e., after 6 wks of diet manipulation). When doing this, contextual fear generalization

(freezing ~40% or more in Neutral Context) was observed in all but males consuming a high salt

diet (Fig. 3C). In fact, contextual fear expression in females consuming a high salt diet tested in

the Neutral Context was so high that it was statistically indifferent from females consuming

either diet and tested in the Training Context (Fig. 3C). On the contrary, males tested in the

Neutral Context after consuming high salt exhibited significantly lower contextual fear than

females tested in the Neutral Context after consuming high salt (Fig. 3C). This indicates a

bidirectional, sex-selective effect of high salt consumption on contextual fear generalization.

Regardless of diet condition, all mice had *ad libitum* access to drinking water. Nonetheless, we

evaluated if serum osmolality was affected across diet condition, sex, or testing Context in mice

trained with foot shocks, as well as across diet, sex, and Experiment in control no shock mice.

In control no shock mice, no three-way interactions were observed, but a significant two-way

interaction where serum osmolality across the sexes differed between Experiments (S11 Table).

Despite this interaction, Holm-Šídák's post-hoc testing revealed no significant directional

differences when the data were examined between sexes within each Experiment for control no

shock mice. Analyses of mice trained with foot shocks revealed zero three- or two-way

interactions, and no significant main effects (S12 Table). Serum osmolality was not affected by

diet, neither in control no shock mice nor in fear conditioned mice. Taken together, these

findings suggest serum osmolality was unaltered in all mice, and thus the behavioral changes

observed were not the consequence of osmotic stress. For food, water, salt, and kcal

consumption data, plus body weight records, readers are referred to the Supporting Information.

Fig. 4. ***Serum osmolality in context fear conditioned mice across Experiments***.

Mice assigned to 0.4% NaCl represented by blue symbols, mice assigned to 4.0% NaCl

represented by red symbols; mice tested in Training Context represented by squares and solid

lines, mice tested in Neutral Context represented by circles and dotted lines. Serum osmolality

in context fear conditioned mice from A) Experiment 1, B) Experiment 2 (grey shading), and C)

Experiment 3. Experiment 1: 0.4% NaCl females Training Context, n=8; 0.4% NaCl females

Neutral Context, n=9; 4.0% NaCl females Training Context, n=8; 4.0% NaCl females Neutral

Context, n=9; 0.4% NaCl males Training Context, n=8; 0.4% NaCl males Neutral Context, n=9;

4.0% NaCl males Training Context, n=8; 4.0% NaCl males Neutral Context, n=8. Experiment 2:

0.4% NaCl females Training Context, n=9; 0.4% NaCl females Neutral Context, n=9; 4.0% NaCl females Training Context, n=8; 4.0% NaCl females Neutral Context, n=8; 0.4% NaCl males Training Context, n=9; 0.4% NaCl males Neutral Context, n=9; 4.0% NaCl males Training Context, n=9; 4.0% NaCl males Neutral Context, n=9. Experiment 3: 0.4% NaCl females Training Context, n=8; 0.4% NaCl females Neutral Context, n=9; 4.0% NaCl females Training Context, n=9; 4.0% NaCl females Neutral Context, n=8; 0.4% NaCl males Training Context, n=7; 0.4% NaCl males Neutral Context, n=8; 4.0% NaCl males Training Context, n=7; 4.0% NaCl males Neutral Context, n=7. Data are graphed as mean ± 95% confidence interval.

In addition to serum osmolality, serum corticosterone levels were also measured in all mice 48 h after completion of context fear testing. Similar to serum osmolality, no three-way interactions were detected in log-transformed serum corticosterone (cort) levels of no shock control mice, but we did find a significant interaction between sexes across Experiments (S13 Table). Unlike with serum osmolality, Holm-Šídák's post-hoc tests revealed directional, diet-specific differences in cort levels within sexes across Experiments, and across sexes within an Experiment (S6 Fig.). Specifically, control-diet consuming females had lower cort levels in Experiment 2 relative to Experiment 1 (p=0.0423). Similarly, high salt-consuming males had lower cort levels in Experiment 3 relative to Experiment 1 (p=0.0109). Within Experiment 3, females consuming high salt had significantly higher cort levels relative to high salt males (p=0.0143). Considering these changes were observed in mice that were not exposed to a foot shock at any point, these data suggest that six weeks of high salt consumption could sex-selectively reduce cort levels in males, but not females. However, the absence of a significant difference (p=0.2369) between cort levels of high salt males between Experiments 1 and 2 challenges this interpretation.

Cort levels in mice that underwent context fear conditioning with foot shocks had no significant three- nor two-way significant interactions (S14 Table). Sex alone had a significant effect on cort levels in Experiments 1 and 3, whereas context specifically influenced cort levels in

Experiment 2. Drilling down to directional differences, there were largely no significant

differences detected by Holm-Šídák's post-hoc tests (Fig. 5A, 5B). The one exception was that

in Experiment 3, males had lower cort levels than females (p=0.0183), specifically between

those consuming control diet and tested in the Training Context (Fig. 5C). Similar, but non-

significant, trends were noted between females and males tested in the Neutral Context

consuming control diet (p=0.0549) or consuming high salt (p=0.0859; Fig. 5C) for Experiment 3.

Generally, cort levels did not consistently vary as a result of sex, diet, testing context, nor their

interaction, indicating that shifts in circulating corticosterone are not responsible for the

observed changes in contextual fear expression nor contextual fear generalization. Circulating

cort levels were largely not correlated with contextual fear expression, except in females

consuming high salt and tested in the Training Context in Experiment 3 (p=0.039; see S23 Fig.).

Fig. 5. *Log-transformed serum corticosterone levels in fear conditioned mice across*

Experiments.

Mice assigned to 0.4% NaCl represented by blue symbols, mice assigned to 4.0% NaCl

represented by red symbols; mice tested in Training Context represented by squares and solid

lines, mice tested in Neutral Context represented by circles and dotted lines. Log-transformed

serum corticosterone levels in context fear conditioned mice in A) Experiment 1, B) Experiment

2 (grey shading), and C) Experiment 3. Experiment 1: 0.4% NaCl females Training Context,

n=6; 0.4% NaCl females Neutral Context, n=7; 4.0% NaCl females Training Context, n=7; 4.0%

NaCl females Neutral Context, n=7; 0.4% NaCl males Training Context, n=8; 0.4% NaCl males

Neutral Context, n=9; 4.0% NaCl males Training Context, n=8; 4.0% NaCl males Neutral

Context, n=8. Experiment 2: 0.4% NaCl females Training Context, n=8; 0.4% NaCl females

Neutral Context, n=9; 4.0% NaCl females Training Context, n=7; 4.0% NaCl females Neutral

Context, n=7; 0.4% NaCl males Training Context, n=9; 0.4% NaCl males Neutral Context, n=9;

4.0% NaCl males Training Context, n=8; 4.0% NaCl males Neutral Context, n=10. Experiment

3: 0.4% NaCl females Training Context, n=7; 0.4% NaCl females Neutral Context, n=8; 4.0%

NaCl females Training Context, n=8; 4.0% NaCl females Neutral Context, n=7; 0.4% NaCl

males Training Context, n=8; 0.4% NaCl males Neutral Context, n=8; 4.0% NaCl males Training

Context, n=8; 4.0% NaCl males Neutral Context, n=7. Data are graphed as mean ± 95%

confidence interval. [+]indicates $p<0.05$ difference between females and males on same diet and

tested in same context.

Discussion

This is the first investigation into how context fear training, and subsequent expression and

generalization, are affected across the sexes by a high salt diet. Our findings suggest that a

relatively short (two week) period of high salt consumption might reduce expression of recently

learned context fear in males, whereas a longer (six week) consumption period of high salt

prevents decreases in expression of recently learned context fear in males. Moreover, context

fear generalization was enhanced in females, but attenuated in males, by consumption of a high

salt diet. Despite being contrary to our hypothesis regarding males, these data indicate that

excess salt consumption affects behavior in a sex-selective manner. Further, these behavioral

changes occurred independent of any changes in serum osmolality, and likewise largely did not

map onto serum corticosterone levels, indicating that neither osmotic nor physiologic stress

states were responsible.

Previous studies assessing how excess salt intake affects context fear behavior exclusively

used male rats [25] or male mice [26]. These studies both used higher salt diets of 8.0% NaCl,

compared to the 4.0% NaCl diet used here [25,26]. Both teams reported attenuated context

fear expression, indexed by freezing behavior. This freezing behavior was measured during

testing that occurred in the Training Context either 24 h [26], or both 2 and 24 h [25], after

training. Their observations differ from ours, which did not observe significant effects of high

salt diet on contextual fear expression in the Training Context. A similarity between the present

study and that of Ge and colleagues is the absence of any diet influence on fear acquisition to

the auditory cue [26], or in our case, to the context. However, this raises an important caveat

regarding both prior investigations - rodents were trained using a cued fear paradigm with an

auditory tone [25,26], rather than training the rodents as the present study did with a traditional

context fear conditioning protocol that lacks discrete cues. These different training procedures

engage distinct brain regions [49,50], and likewise elicit distinguishable patterns of neuronal

activation [51–53]. Thus, in the present study we assessed contextual fear expression

specifically following contextual fear conditioning, rather than cued fear conditioning.

Generalization of contextual fear is a hallmark symptom of many anxiety disorders, and is a

conserved cross-species neurobehavioral phenomenon that is useful for understanding how

neurophysiological changes influence behaviors [32,54]. The majority of contextual fear

generalization investigations in rodents have been performed in males [33], though studies have

recently begun to include females. Such sex-inclusive studies indicate that rodents of both

sexes can and do exhibit generalized contextual fear, at timepoints both days [55] and weeks

[56] after context fear training. Sex differences in the influence of various contextual modalities

have been reported (see review [33]), and in our study contextual differences included tactile,

olfactory, and visual modalities. We observed robust discrimination between contexts across

the sexes when training was temporally proximal to testing, as well as convincing evidence of

contextual fear generalization in both sexes consuming control diet when testing occurred four

weeks following training. In contrast to our hypothesis that high salt intake would enhance

contextual fear generalization in males, we instead found an attenuation of this measure in

males, but an enhancement in females. Given the near absence of investigations into how

excess salt intake affects female behaviors (see review [28]), plus the relatively scant literature

on contextual fear generalization in females (see review [33]), we did not have enough

information to formulate an *a priori* hypothesis for females consuming high salt. Some

researchers have suggested excess salt intake exacerbates neuropsychiatric symptoms and

conditions [29,30]. Our data support this relationship, at least in females. However, the

relationship between high salt consumption and anxiety-relevant behaviors was not supported in

our male mice, and in fact our data suggest an inverse relationship. Given frequent comorbidity

between cardiovascular diseases and pathological anxiety [11–15] (see also reviews [16,17]),

and the adverse impact of overconsumption of salt on cardiovascular disease risk [5,10], this

finding in males was surprising. Nonetheless, it may be that in males, high salt-mediated

cardiovascular pathology is necessary for emergence of increased anxiety-related behaviors,

whereas in females excess salt is sufficient. This would parallel findings indicating greater

metabolic disruptions in men relative to women with comorbid obesity and mental health

disorders [57–59] (see review [60]), greater incidence of depression and anxiety in men versus

women with multiple sclerosis [61], and increased incidence of a biomarker of cardiovascular

disease in men versus women with depression [62]. However, some literature suggests this is

not a consistent finding across disease conditions [63].

Along these lines, we evaluated if mice consuming excess salt exhibited peripheral indicators

of osmotic (serum osmolality) or physiological (serum corticosterone) stress. Changes in these

measures could suggest an overall disrupted state of homeostasis that might confound

contextual fear learning and/or expression. For example, osmotic stress can reduce sensitivity

to pain [64]. Though the shock level (0.8 mA) and duration (1 sec) that we use are intended to

only elicit discomfort/distress, it remains possible that osmotic stress could skew sensory

perception of this brief aversive experience. Similarly, endogenous corticosterone levels can

influence temporal development of contextual fear generalization as well as the persistence of

contextual discrimination [65]. The absence of any sex-, diet-, or context-specific differences in

serum osmolality indicates that osmotic stress was not present in our experimental animals. No

significant correlations between cort levels and fear expression in the Neutral Context were

detected for either sex in Experiment 3. Indeed, of all the mice trained in the Training Context

with foot shocks, only Experiment 3 females consuming high salt and tested in the Training

Context displayed a significant (negative) correlation between context fear expression and cort

levels (S23 Fig.). The absence of any significant correlations between cort levels and any other

sex/diet/context combination across Experiments indicates that cort levels overall probably did

not confound observed context fear expression and generalization.

Limitations

Limitations of this study include singly housing of mice to allow for tracking of individual

food/water/salt/kcal consumption and changes in body weight, meaning that mice were

excluded from social stimulation. Some data suggests behavioral responses to single housing

differs across the sexes in mice and rats (see review [66]), so this might have influenced sex-

specific behaviors observed in the present study. However, a meta-analysis of 293 male and

female mouse studies indicates that singly housing mice reduces coefficients of variation in trait

variability by 37% in both sexes [67], suggesting our housing condition may have reduced the

variability of our data. Diet manipulations were relatively short (two or six weeks), restricted to

non-breeding adults, and contained no other accompanying unhealthy dietary components (e.g.,

low fiber, high fat). Only a single shock level and context fear conditioning paradigm was used,

and mice were only tested in a single context. Thus, for future studies it will be important to

evaluate behavior in group housed mice, assess longer diet manipulation periods and include

additional unhealthy dietary components, and study how consumption of excess salt by younger

mice affects their fear behaviors. Testing of behavior during the natural activity period of mice

would be informative, as would assessing generalization and concurrent cort levels during

different stages of the circadian cycle [68].

509 Conclusions

Our study provides foundational evidence of sex-selective consequences of excess salt intake

on context fear expression and generalization. Additional studies will be essential to flesh out

these initial findings and determine what brain cell populations are most impacted by excess salt

intake, and how the effects of high salt might be mitigated. One logical starting point is

evaluation of microglia, given converging evidence that both fear processing [69,70] and high

salt intake [42,71,72] affect microglial activation. Moreover, oral administration of the brain-

penetrant anti-inflammatory drug minocycline can mitigate microglial activation and behavioral

changes in rodents consuming high salt [42], and the same drug attenuated fear measures in

human participants [73]. Future studies should also assess how excess salt intake affects cued

fear expression and discrimination, and in so doing begin identifying how specific limbic brain

regions might differentially be affected in their functionality and/or connectivity following

extended periods of excess salt consumption. The evidence presented here, in conjunction with

past literature focused on salt's effects on behaviors outside of the fear domain(reviewed in

[28]), indicates that salt is an underappreciated, non-caloric influence on conserved mammalian

behaviors.

525   **Acknowledgements**

526   We gratefully acknowledge the mice used in this study, and the expert care thereof by our

527   caretakers. Figures were created with Biorender.com (Toronto ON) and GraphPad Prism (San

528   Diego, CA).

References

1. Powles J, Fahimi S, Micha R, Khatibzadeh S, Shi P, Ezzati M, et al. Global, regional and
national sodium intakes in 1990 and 2010: a systematic analysis of 24 h urinary sodium
excretion and dietary surveys worldwide. Bmj Open. 2013;3: e003733. doi:10.1136/bmjopen-
2013-003733

2. Santos JA, Tekle D, Rosewarne E, Flexner N, Cobb L, Al-Jawaldeh A, et al. A Systematic
Review of Salt Reduction Initiatives Around the World: A Midterm Evaluation of Progress
Towards the 2025 Global Non-Communicable Diseases Salt Reduction Target. Adv Nutr.
2021;12: 1768–1780. doi:10.1093/advances/nmab008

3. Bibbins-Domingo K, Chertow GM, Coxson PG, Moran A, Lightwood JM, Pletcher MJ, et al.
Projected Effect of Dietary Salt Reductions on Future Cardiovascular Disease. New Engl J
Medicine. 2010;362: 590–599. doi:10.1056/nejmoa0907355

4. Cogswell ME, Mugavero K, Bowman BA, Frieden TR. Dietary Sodium and Cardiovascular
Disease Risk — Measurement Matters. New Engl J Medicine. 2016;375: 580–586.
doi:10.1056/nejmsb1607161

5. Adler AJ, Taylor F, Martin N, Gottlieb S, Taylor RS, Ebrahim S. Reduced dietary salt for the
prevention of cardiovascular disease. Cochrane Db Syst Rev. 2014; CD009217.
doi:10.1002/14651858.cd009217.pub3

6. He FJ, Tan M, Ma Y, MacGregor GA. Salt Reduction to Prevent Hypertension and
Cardiovascular Disease. J Am Coll Cardiol. 2020;75: 632–647. doi:10.1016/j.jacc.2019.11.055

7. Mozaffarian D, Fahimi S, Singh GM, Micha R, Khatibzadeh S, Engell RE, et al. Global
Sodium Consumption and Death from Cardiovascular Causes. New Engl J Medicine. 2014;371:
624–634. doi:10.1056/nejmoa1304127

8. O'Donnell M, Mente A, Yusuf S. Sodium Intake and Cardiovascular Health. Circ Res.
2015;116: 1046–1057. doi:10.1161/circresaha.116.303771

9. World Health Organization. WHO global sodium benchmarks for different food categories.
Geneva: World Health Organization; 2021. Available:
https://apps.who.int/iris/rest/bitstreams/1344511/retrieve

10. He FJ, MacGregor GA. Salt reduction lowers cardiovascular risk: meta-analysis of outcome
trials. Lancet. 2011;378: 380–382. doi:10.1016/s0140-6736(11)61174-4

11. Scherrer JF, Salas J, Cohen BE, Schnurr PP, Schneider FD, Chard KM, et al. Comorbid
Conditions Explain the Association Between Posttraumatic Stress Disorder and Incident
Cardiovascular Disease. J Am Hear Assoc Cardiovasc Cerebrovasc Dis. 2019;8: e011133.
doi:10.1161/jaha.118.011133

12. Szlejf C, Suemoto CK, Santos IS, Brunoni AR, Nunes MA, Viana MC, et al. Poorer
cardiovascular health is associated with psychiatric comorbidity: results from the ELSA-Brasil
Study. Int J Cardiol. 2019;274: 358–365. doi:10.1016/j.ijcard.2018.06.037

13. Tran J, Norton R, Conrad N, Rahimian F, Canoy D, Nazarzadeh M, et al. Patterns and
temporal trends of comorbidity among adult patients with incident cardiovascular disease in the
UK between 2000 and 2014: A population-based cohort study. Plos Med. 2018;15: e1002513.
doi:10.1371/journal.pmed.1002513

14. Dhital PS, Sharma K, Poudel P, Dhital PR. Anxiety and Depression among Patients with
Coronary Artery Disease Attending at a Cardiac Center, Kathmandu, Nepal. Nurs Res Pract.
2018;2018: 4181952. doi:10.1155/2018/4181952

15. Vidal C, Polo R, Alvarez K, Falgas-Bague I, Wang Y, Cook BL, et al. Co-Occurrence of
Posttraumatic Stress Disorder and Cardiovascular Disease Among Ethnic/Racial Groups in
the United States. Psychosom Med. 2018;80: 680–688. doi:10.1097/psy.0000000000000601

16. Meuret AE, Tunnell N, Roque A. Anxiety Disorders, Rethinking and Understanding Recent
Discoveries. Adv Exp Med Biol. 2020;1191: 237–261. doi:10.1007/978-981-32-9705-0_15

17. Johnson HM. Anxiety and Hypertension: Is There a Link? A Literature Review of the
Comorbidity Relationship Between Anxiety and Hypertension. Curr Hypertens Rep. 2019;21: 66.
doi:10.1007/s11906-019-0972-5

18. Walker JM, Harrison FE. Shared Neuropathological Characteristics of Obesity, Type 2
Diabetes and Alzheimer's Disease: Impacts on Cognitive Decline. Nutrients. 2015;7: 7332–
7357. doi:10.3390/nu7095341

19. Tumminia A, Vinciguerra F, Parisi M, Frittitta L. Type 2 Diabetes Mellitus and Alzheimer's
Disease: Role of Insulin Signalling and Therapeutic Implications. Int J Mol Sci. 2018;19: 3306.
doi:10.3390/ijms19113306

20. Subba R, Sandhir R, Singh SP, Mallick BN, Mondal AC. Pathophysiology linking depression
and type 2 diabetes: Psychotherapy, physical exercise, and fecal microbiome transplantation as
damage control. Eur J Neurosci. 2021;53: 2870–2900. doi:10.1111/ejn.15136

21. Joseph JJ, Golden SH. Cortisol dysregulation: the bidirectional link between stress,
depression, and type 2 diabetes mellitus. Ann Ny Acad Sci. 2017;1391: 20–34.
doi:10.1111/nyas.13217

22. Silva N de ML e, Lam MP, Soares CN, Munoz DP, Milev R, Felice FGD. Insulin Resistance
as a Shared Pathogenic Mechanism Between Depression and Type 2 Diabetes. Frontiers
Psychiatry. 2019;10: 57. doi:10.3389/fpsyt.2019.00057

23. Ernsberger P, Azar S, Iwai J. Open-field behavior in two models of genetic hypertension and
the behavioral effects of salt excess. Behav Neural Biol. 1983;37: 46–60. doi:10.1016/s0163-
1047(83)91061-0

24. Chugh G, Asghar M, Patki G, Bohat R, Jafri F, Allam F, et al. A High-Salt Diet Further
Impairs Age-Associated Declines in Cognitive, Behavioral, and Cardiovascular Functions in
Male Fischer Brown Norway Rats. J Nutrition. 2013;143: 1406–1413.
doi:10.3945/jn.113.177980

25. Guo C, Wei Z, Huang F, Qin M, Li X, Wang Y, et al. High salt induced hypertension leads to
cognitive defect. Oncotarget. 2017;5: 95780–95790. doi:10.18632/oncotarget.21326

26. Ge Q, Wang Z, Wu Y, Huo Q, Qian Z, Tian Z, et al. High salt diet impairs memory-related
synaptic plasticity via increased oxidative stress and suppressed synaptic protein expression.
Mol Nutr Food Res. 2017;61: 1700134. doi:10.1002/mnfr.201700134

27. Gilman TL, George CM, Andrade MA, Mitchell NC, Toney GM, Daws LC. High Salt Intake
Lowers Behavioral Inhibition. Front Behav Neurosci. 2019;13: 271.
doi:10.3389/fnbeh.2019.00271

28. Beaver JN, Gilman TL. Salt as a non-caloric behavioral modifier: A review of evidence from
pre-clinical studies. Neurosci Biobehav Rev. 2022;135: 104385.
doi:10.1016/j.neubiorev.2021.10.007

29. Abdoli A. Hypothesis: High salt intake as an inflammation amplifier might be involved in the
pathogenesis of neuropsychiatric disorders. Clin Exp Neuroimmunol. 2017;8: 146–157.
doi:10.1111/cen3.12389

30. Jones AYM, Dean E, Lo SK. Interrelationships Between Anxiety, Lifestyle Self-reports and
Fitness in a Sample of Hong Kong University Students. Ann Ny Acad Sci. 2002;5: 65–71.
doi:10.1080/102538902900012350

31. Webler RD, Berg H, Fhong K, Tuominen L, Holt DJ, Morey RA, et al. The Neurobiology of
Human Fear Generalization: Meta-Analysis and Working Neural Model. Neurosci Biobehav Rev.
2021;128: 421–436. doi:10.1016/j.neubiorev.2021.06.035

32. Dunsmoor JE, Paz R. Fear Generalization and Anxiety: Behavioral and Neural Mechanisms.
Biol Psychiat. 2015;78: 336–343. doi:10.1016/j.biopsych.2015.04.010

33. Asok A, Kandel ER, Rayman JB. The Neurobiology of Fear Generalization. Front Behav
Neurosci. 2019;12: 329. doi:10.3389/fnbeh.2018.00329

34. Jasnow AM, Lynch JF, Gilman TL, Riccio DC. Perspectives on fear generalization and its
implications for emotional disorders. J Neurosci Res. 2017;95: 821–835. doi:10.1002/jnr.23837

35. Terry AV, Hernandez CM, Buccafusco JJ. Dahl salt-sensitive and salt-resistant rats:
examination of learning and memory performance, blood pressure, and the expression of
central nicotinic acetylcholine receptors. Neuroscience. 2001;103: 351–363. doi:10.1016/s0306-
4522(00)00569-8

36. Pelisch N, Hosomi N, Ueno M, Nakano D, Hitomi H, Mogi M, et al. Blockade of AT1
Receptors Protects the Blood–Brain Barrier and Improves Cognition in Dahl Salt-Sensitive
Hypertensive Rats. Am J Hypertens. 2011;24: 362–368. doi:10.1038/ajh.2010.241

37. Liu Y-Z, Chen J-K, Li Z-P, Zhao T, Ni M, Li D-J, et al. High-salt diet enhances hippocampal
oxidative stress and cognitive impairment in mice. Neurobiol Learn Mem. 2014;114: 10–15.
doi:10.1016/j.nlm.2014.04.010

38. Faraco G, Brea D, Garcia-Bonilla L, Wang G, Racchumi G, Chang H, et al. Dietary salt
promotes neurovascular and cognitive dysfunction through a gut-initiated TH17 response. Nat
Neurosci. 2018;21: 240–249. doi:10.1038/s41593-017-0059-z

39. Mitchell NC, Gilman TL, Daws LC, Toney GM. High Salt Intake Enhances Swim Stress-
Induced PVN Vasopressin Cell Activation and Active Stress Coping. Psychoneuroendocrino.
2018;93: 29–38. doi:10.1016/j.psyneuen.2018.04.003

40. Ge Q, Hu X, Ma N, Sun M, Zhang L, Cai Z, et al. Maternal high-salt diet during pregnancy
impairs synaptic plasticity and memory in offspring. Faseb J. 2021;35: e21244.
doi:10.1096/fj.202001890r

41. Leal PE de PT, Silva AA da, Rocha-Gomes A, Riul TR, Cunha RA, Reichetzeder C, et al.
High-Salt Diet in the Pre- and Postweaning Periods Leads to Amygdala Oxidative Stress and
Changes in Locomotion and Anxiety-Like Behaviors of Male Wistar Rats. Front Behav Neurosci.
2022;15: 779080. doi:10.3389/fnbeh.2021.779080

42. Gilman TL, Mitchell NC, Daws LC, Toney GM. Neuroinflammation contributes to high salt
intake-augmented neuronal activation and active coping responses to acute stress. Int J
Neuropsychoph. 2018;22: pyy099-. doi:10.1093/ijnp/pyy099

43. National Research Council. Guide for the Care and Use of Laboratory Animals. 8th ed.
Washington, D.C.: National Academies Press (US); 2011. doi:10.17226/12910

44. Cullen PK, Gilman TL, Winiecki P, Riccio DC, Jasnow AM. Activity of the anterior cingulate
cortex and ventral hippocampus underlie increases in contextual fear generalization. Neurobiol
Learn Mem. 2015;124: 19–27. doi:10.1016/j.nlm.2015.07.001

45. Poulos AM, Mehta N, Lu B, Amir D, Livingston B, Santarelli A, et al. Conditioning- and time-
dependent increases in context fear and generalization. Learn Memory. 2016;23: 379–385.
doi:10.1101/lm.041400.115

46. Lynch JF, Winiecki P, Gilman TL, Adkins JM, Jasnow AM. Hippocampal GABAB(1a)
Receptors Constrain Generalized Contextual Fear. Neuropsychopharmacol. 2017;42: 914–924.
doi:10.1038/npp.2016.255

47. Uarquin DG, Meyer JS, Cardenas FP, Rojas MJ. Effect of Overcrowding on Hair
Corticosterone Concentrations in Juvenile Male Wistar Rats. J Am Assoc Laboratory Animal Sci
Jaalas. 2016;55: 749–755. Available: https://pubmed.ncbi.nlm.nih.gov/27931312/

48. Teilmann AC, Kalliokoski O, Sørensen DB, Hau J, Abelson KSP. Manual versus automated
blood sampling: impact of repeated blood sampling on stress parameters and behavior in male
NMRI mice. Lab Anim. 2014;48: 278–291. doi:10.1177/0023677214541438

49. Phillips RG, LeDoux JE. Differential Contribution of Amygdala and Hippocampus to Cued
and Contextual Fear Conditioning. Behav Neurosci. 1992;106: 274–285. doi:10.1037/0735-
7044.106.2.274

50. Curzon P, Rustay NR, Browman KE. Cued and Contextual Fear Conditioning for Rodents.
CRC Press/Taylor & Francis, Boca Raton (FL); 2011. Available:
https://www.ncbi.nlm.nih.gov/books/NBK5223

51. Trogrlic L, Wilson YM, Newman AG, Murphy M. Context fear learning specifically activates
distinct populations of neurons in amygdala and hypothalamus. Learn Memory. 2011;18: 678–
687. doi:10.1101/lm.2314311

52. Butler CW, Wilson YM, Gunnersen JM, Murphy M. Tracking the fear memory engram:
discrete populations of neurons within amygdala, hypothalamus, and lateral septum are
specifically activated by auditory fear conditioning. Learn Memory. 2015;22: 370–384.
doi:10.1101/lm.037663.114

53. Chaaya N, Jacques A, Belmer A, Richard DJ, Bartlett SE, Battle AR, et al. Localization of
Contextual and Context Removed Auditory Fear Memory within the Basolateral Amygdala
Complex. Neuroscience. 2019;398: 231–251. doi:10.1016/j.neuroscience.2018.12.004

54. Cooper SE, Dis EAM van, Hagenaars MA, Krypotos A-M, Nemeroff CB, Lissek S, et al. A
meta-analysis of conditioned fear generalization in anxiety-related disorders.
Neuropsychopharmacol. 2022;47: 1652–1661. doi:10.1038/s41386-022-01332-2

55. Keiser AA, Turnbull LM, Darian MA, Feldman DE, Song I, Tronson NC. Sex Differences in
Context Fear Generalization and Recruitment of Hippocampus and Amygdala during Retrieval.
Neuropsychopharmacol. 2017;42: 397–407. doi:10.1038/npp.2016.174

56. Asok A, Hijazi J, Harvey LR, Kosmidis S, Kandel ER, Rayman JB. Sex Differences in
Remote Contextual Fear Generalization in Mice. Front Behav Neurosci. 2019;13: 56.
doi:10.3389/fnbeh.2019.00056

57. Udo T, McKee SA, White MA, Masheb RM, Barnes RD, Grilo CM. Menopause and
metabolic syndrome in obese individuals with binge eating disorder. Eat Behav. 2014;15: 182–
185. doi:10.1016/j.eatbeh.2014.01.003

58. Bjerkeset O, Romundstad P, Evans J, Gunnell D. Association of Adult Body Mass Index and
Height with Anxiety, Depression, and Suicide in the General PopulationThe HUNT Study. Am J
Epidemiol. 2008;167: 193–202. doi:10.1093/aje/kwm280

59. Staiano AE, Marker AM, Martin CK, Katzmarzyk PT. Physical activity, mental health, and
weight gain in a longitudinal observational cohort of nonobese young adults. Obesity. 2016;24:
1969–1975. doi:10.1002/oby.21567

60. Tronieri JS, Wurst CM, Pearl RL, Allison KC. Sex Differences in Obesity and Mental Health.
Curr Psychiat Rep. 2017;19: 29. doi:10.1007/s11920-017-0784-8

61. Marrie RA, Patten SB, Tremlett H, Wolfson C, Warren S, Svenson LW, et al. Sex differences
in comorbidity at diagnosis of multiple sclerosis A population-based study. Neurology. 2016;86:
1279–1286. doi:10.1212/wnl.0000000000002481

62. Garcia RG, Zarruk JG, Guzman JC, Barrera C, Pinzon A, Trillos E, et al. Sex differences in
cardiac autonomic function of depressed young adults. Biol Psychol. 2012;90: 179–185.
doi:10.1016/j.biopsycho.2012.03.016

63. Basso L, Boecking B, Neff P, Brueggemann P, Cederroth CR, Rose M, et al. Sex
Differences in Comorbidity Combinations in the Swedish Population. Biomol. 2022;12: 949.
doi:10.3390/biom12070949

64. Chatoui H, Abbaoui A, Hiba OE, Draoui A, Gamrani H. Neurobehavioral and
neurophysiological effects of prolonged osmotic stress in rats: A focus on anxiety state and pain
perception. J Chem Neuroanat. 2020;106: 101789. doi:10.1016/j.jchemneu.2020.101789

65. Corrêa M dos S, Vaz B dos S, Grisanti GDV, Paiva JPQ de, Tiba PA, Fornari RV.
Relationship between footshock intensity, post-training corticosterone release and contextual
fear memory specificity over time. Psychoneuroendocrino. 2019;110: 104447.
doi:10.1016/j.psyneuen.2019.104447

66. Ma L, Xu Y, Wang G, Li R. What do we know about sex differences in depression: A review
of animal models and potential mechanisms. Prog Neuro-psychopharmacology Biological
Psychiatry. 2019;89: 48–56. doi:10.1016/j.pnpbp.2018.08.026

67. Prendergast BJ, Onishi KG, Zucker I. Female mice liberated for inclusion in neuroscience
and biomedical research. Neurosci Biobehav Rev. 2014;40: 1–5.
doi:10.1016/j.neubiorev.2014.01.001

68. Kalsbeek A, Spek R van der, Lei J, Endert E, Buijs RM, Fliers E. Circadian rhythms in the
hypothalamo–pituitary–adrenal (HPA) axis. Mol Cell Endocrinol. 2012;349: 20–29.
doi:10.1016/j.mce.2011.06.042

69. Chaaya N, Jacques A, Belmer A, Beecher K, Ali SA, Chehrehasa F, et al. Contextual Fear
Conditioning Alter Microglia Number and Morphology in the Rat Dorsal Hippocampus. Front Cell
Neurosci. 2019;13: 214. doi:10.3389/fncel.2019.00214

70. Enomoto S, Kato TA. Involvement of microglia in disturbed fear memory regulation: Possible
microglial contribution to the pathophysiology of posttraumatic stress disorder. Neurochem Int.
2021;142: 104921. doi:10.1016/j.neuint.2020.104921

71. Hu L, Zhu S, Peng X, Li K, Peng W, Zhong Y, et al. High Salt Elicits Brain Inflammation and
Cognitive Dysfunction, Accompanied by Alternations in the Gut Microbiota and Decreased
SCFA Production. J Alzheimer's Dis. 2020;77: 629–640. doi:10.3233/jad-200035

72. Zhang T, Wang D, Li X, Jiang Y, Wang C, Zhang Y, et al. Excess salt intake promotes M1
microglia polarization via a p38/MAPK/AR-dependent pathway after cerebral ischemia in mice.
Int Immunopharmacol. 2020;81: 106176. doi:10.1016/j.intimp.2019.106176

73. Chan SY, Capitão L, Probert F, Klinge C, Hoeckner S, Harmer CJ, et al. A single
administration of the antibiotic, minocycline, reduces fear processing and improves implicit
learning in healthy volunteers: analysis of the serum metabolome. Transl Psychiat. 2020;10:
148. doi:10.1038/s41398-020-0818-6

Supporting Information

Tables

S1 Table. Three-way repeated measures ANOVAs on context fear 'training' for control no shock mice across Experiments.

S2 Table. Three-way ANOVAs on context fear testing of control no shock mice across Experiments.

S3 Table. Three-way repeated measures ANOVAs on context fear training for context fear conditioned mice of both sexes in Experiment 1.

S4 Table. Three-way repeated measures ANOVAs on context fear training for context fear conditioned mice of both sexes in Experiment 2.

S5 Table. Three-way repeated measures ANOVAs on context fear training for context fear conditioned mice of both sexes in Experiment 3.

S6 Table. Three-way repeated measures ANOVAs on full 10 min time course of context fear testing for control no shock mice across Experiments.

S7 Table. Three-way ANOVAs on context fear expression (min 2-6) for context fear conditioned mice across Experiment.

S8 Table. Three-way repeated measures ANOVAs on full 10 min time course of context fear testing for mice of both sexes in Experiment 1.

S9 Table. Three-way repeated measures ANOVAs on full 10 min time course of context fear testing for mice of both sexes in Experiment 2.

S10 Table. Three-way repeated measures ANOVAs on full 10 min time course of context fear testing for mice of both sexes in Experiment 3.

S11 Table. Three-way ANOVAs on serum osmolality in control no shock mice across Experiments.

S12 Table. Three-way ANOVAs on serum osmolality in context fear conditioned mice across Experiments.

S13. Three-way ANOVAs on log-transformed serum corticosterone levels in control no shock mice across Experiments.

S14 Table. Three-way ANOVAs on log-transformed serum corticosterone in context fear conditioned mice across Experiments.

S15 Table. Three-way repeated measures ANOVAs on weekly average water consumption per day for context fear conditioned mice across Experiments.

S16 Table. Three-way repeated measures ANOVAs on weekly average food consumption per day for context fear conditioned mice across Experiments.

S17 Table. Three-way repeated measures ANOVAs on weekly body weight change for

context fear conditioned mice across Experiments.

S18 Table. Three-way repeated measures ANOVAs on twice weekly body weight

measurements of context fear conditioned mice across Experiments.

S19 Table. Three-way repeated measures ANOVAs on weekly average water

consumption per day for control no shock mice across Experiments.

S20 Table. Three-way repeated measures ANOVAs on weekly average food consumption

per day for control no shock mice across Experiments.

S21 Table. Three-way repeated measures ANOVAs on weekly body weight changes for

control no shock mice across Experiments.

S22 Table. Three-way repeated measures ANOVAs on twice weekly body weights of

control no shock mice across Experiments.

S23 Table. Three-way repeated measures ANOVAs on weekly average NaCl consumed

per day by context fear conditioned mice across Experiments.

S24 Table. Three-way repeated measures ANOVAs on weekly average NaCl consumed

as a percentage of body weight in context fear conditioned mice across Experiments.

S25 Table. Three-way repeated measures ANOVAs on weekly ratio of water to NaCl

consumed in context fear conditioned mice across Experiments.

S26 Table. Three-way repeated measures ANOVAs on weekly average kcal consumed per day by context fear conditioned mice across Experiments.

S27 Table. Three-way repeated measures ANOVAs on weekly kcal consumed as a percentage of body weight in context fear conditioned mice across Experiments.

S28 Table. Three-way repeated measures ANOVAs on weekly average NaCl consumed per day by control no shock mice across Experiments.

S29 Table. Three-way repeated measures ANOVAs on weekly NaCl consumed as a percentage of body weight by control no shock mice across Experiments.

S30 Table. Three-way repeated measures ANOVAs on weekly water to NaCl ratio consumed by control no shock mice across Experiments.

S31 Table. Three-way repeated measures ANOVAs on weekly average kcal consumed per day by control no shock mice across Experiments.

S32 Table. Three-way repeated measures ANOVAs on weekly average kcal consumed as a percentage of body weight by control no shock mice across Experiments.

S33 Table. Details of confirmed outliers (> 5 standard deviations (SD) ± average of data set sans suspected outlier) within respective data sets. Outliers are marked with red font.

Figures

S1 Fig. Fear behavior in no shock control mice across Experiments.

S2 Fig. Fear behavior in no shock control mice across Experiments with shortened Y

axes.

S3 Figure. Time course of context fear expression testing in no shock groups.

S4 Fig. Time course of context fear expression testing in shock groups.

S5 Fig. Serum osmolality in no shock control mice across Experiments.

S6 Fig. Log-transformed serum corticosterone levels of no shock control mice across

Experiments.

S7 Fig. Raw (pre-transformed) serum corticosterone levels of no shock control mice

across Experiments.

S8 Fig. Raw (pre-transformed) serum corticosterone levels in fear conditioned mice

across Experiments.

S9 Fig. Average water consumed per day by context fear conditioned mice across

Experiments.

S10 Fig. Average food consumed per day by context fear conditioned mice across

Experiments.

S11 Fig. Weekly body weight change in context fear conditioned mice across

Experiments.

S12 Fig. Body weight of context fear conditioned mice across Experiments.

S13 Fig. Average water and food consumed per day by control no shock mice across

Experiments.

S14 Fig. Body weights and body weight changes in control no shock mice across

Experiments.

S15 Fig. Average NaCl consumed per day by context fear conditioned mice across

Experiments.

S16 Fig. NaCl consumed as a percentage of body weight by context fear conditioned

mice across Experiments.

S17 Fig. Water to NaCl consumed ratio for context fear conditioned mice across

Experiments.

S18 Fig. Average kcal consumed per day by context fear conditioned mice across

Experiments.

S19 Fig. Consumed kcal as a percentage of body weight by context fear conditioned

mice across Experiments.

S20 Fig. Average NaCl consumed per day and NaCl consumed as a percentage of body

weight by control no shock mice across Experiments.

S21 Figure. Ratio of water to NaCl consumed by control no shock mice across

Experiments.

S22 Fig. Average kcal consumed per day and kcal consumed as a percentage of body

weight by control no shock mice across Experiments.

S23 Fig. Correlations between context fear expression and log-transformed serum

corticosterone levels across contexts and Experiments.

S24 Figure. Correlations between context fear expression and log-transformed serum

corticosterone levels in control no shock mice across Experiments.

Key

- 0.4% NaCl Training Context
- 4.0% NaCl Training Context
- 0.4% NaCl Neutral Context
- 4.0% NaCl Neutral Context

A **Experiment 1**

Females

Males

B **Experiment 2**

Females

Males

C **Experiment 3**

Females

Males

Post-shock Period

A · Experiment 1

Training after 2 wks diet ➜ Testing 48 h later

B · Experiment 2

Training after 6 wks diet ➜ Testing 48 h later

C · Experiment 3

Training after 2 wks diet ➜ Testing 4 wks later

Key

☐ 0.4% NaCl Training Context
☐ 4.0% NaCl Training Context
○ 0.4% NaCl Neutral Context
○ 4.0% NaCl Neutral Context

A

Experiment 1

Training after 2 wks diet → Testing 48 h later

Log transformed (Corticosterone, ng/mL)

Females   Males

B

Experiment 2

Training after 6 wks diet → Testing 48 h later

Log transformed (Corticosterone, ng/mL)

Females   Males

C

Experiment 3

Training after 2 wks diet → Testing 4 wks later

Log transformed (Corticosterone, ng/mL)

Females   Males