Title: A STIM dependent dopamine-insulin axis maintains the larval

drive to feed and grow in Drosophila

Authors: Nandashree Kasturacharya[1,2], Jasmine Kaur Dhall[1] and Gaiti

Hasan[1,2,3]

1. National Centre for Biological Sciences, TIFR, Bellary Road,

Bengaluru 560065

2. The University of Trans-Disciplinary Health Sciences and

Technology (TDU), Bengaluru – 560064

3. Corresponding author

Abstract:

Appropriate nutritional intake is essential for organismal survival. In

holometabolous insects such as *Drosophila melanogaster*, the quality and

quantity of food ingested as larvae determines adult size and fecundity.

Here we have identified a subset of dopaminergic neurons (THD') that

maintain the larval motivation to feed. Dopamine release from these

neurons requires the ER Ca^{2+} sensor STIM. Larvae with loss of STIM stop

feeding and growing, whereas expression of STIM in THD' neurons

rescues feeding, growth and viability of STIM null mutants. Moreover STIM

is essential for maintaining excitability and release of dopamine from THD'

neurons. Optogenetic stimulation of THD' neurons activated

neuropeptidergic cells, including median neuro secretory cells that secrete

insulin-like peptides. Loss of STIM in THD' cells alters the developmental

profile of specific insulin-like peptides including ilp3. Loss of ilp3 partially

rescues STIM null mutants and inappropriate expression of ilp3 in larvae

affects development and growth. In summary we have identified a novel

STIM-dependent function of dopamine neurons that modulates

developmental changes in larval feeding behaviour and growth.

Author summary

The ability to feed appropriately when hungry is an essential feature for

organismal survival and is under complex neuronal control. An array of

neurotransmitters and neuropeptides integrate external and internal

signalling cues to initiate, maintain and terminate feeding. In adult

vertebrates and invertebrates dopamine serves as a reward cue for motor

actions, including feeding. Larvae of holometabolous insects, including

Drosophila melanogaster, feed and grow constantly followed by gradual

cessation of feeding, once sufficient growth is achieved for transition to the

next stages of development. Here we identified a subset of larval

dopaminergic neurons in *Drosophila melanogaster*, activity in which

maintains continuous feeding in larvae. By analysis of a null mutant we

show that these neurons require the Stromal Interaction Molecule (STIM)

an ER Ca^{2+} sensor, to maintain excitability. In turn they modulate activity of

certain neuropeptidergic cells. Among these are the median neurosecretory

cells (MNSc) that synthesize and secrete insulin-like peptides including ilp3.

The identified dopaminergic neurons dysregulate the normal pattern of

larval ilp3 expression leading to premature cessation of feeding and growth.

Overall, our study identified a simple dopamine modulated mechanism for

feeding and growth whose manipulation could be useful for model organism

studies related to feeding disorders, obesity and diabetes.

Introduction:

Animal growth occurs primarily during the juvenile stage of
development. In holometabolous insects, including *Drosophila*, larval
development is considered equivalent to the juvenile stage(1,2). Steady
and appropriate nutritional intake in larvae, is essential for growth and
development, and ultimately determines both survival and fecundity of the
animals. In *Drosophila*, the feeding rate increases in the second instar
larval stage and larvae continue to feed voraciously till the wandering stage
of third instar larvae(3). Increase in feeding rate is accompanied by
acceleration of cell division and cell growth(4).

Feeding behaviour and its modulation in *Drosophila* larvae has been
studied primarily in the third instar larval stage, where it is regulated by
multiple neurotransmitters and neuropeptides that respond to both external
cues and the internal metabolic state (5). Neuropeptide F (NPF; a human
NPY homolog) serves as a motivational signal for foraging in larvae in
response to appetitive odours. The activity of NPF neurons appears
dependent on inputs from two pairs of central dopaminergic neurons that
receive tertiary olfactory inputs (6,7). Mutants for the short neuropeptide F
(sNPF), encoded by an independent gene from NPF, affect body size by
regulating food intake in larvae (8). When food is restricted octopaminergic
circuits regulate feeding independent of NPF signalling in 3^{rd} instar larvae
(9). Neurons that secrete the Hugin neuropeptide respond to averse
gustatory signals and their activation suppresses larval feeding (10). Where
essential amino acids are imbalanced a subset of dopaminergic neurons
are required for rejection of food by larvae (11). In addition serotonergic
neurons from the brain project to the gut where they potentially regulate
feeding-related muscle movements (12). Insulin like peptides (ilps), are

secreted by the medial neurosecretory cells (MNSc) that access the

internal metabolic state and release ilps into neurohaemal sites for

circulation. In adults ilps terminate feeding based on the energy state of the

organism (13). Analysis of the recently concluded larval connectome

demonstrates that MNSc receive both direct and indirect inputs from the

enteric nervous system found in the larval gut (14). The larval MNSc also

receive synaptic inputs from central neurons that release the Hugin

neuropeptide (10). Ilps released through neurohaemal sites from the MNSc

circulate through the body and regulate energy metabolism (15), synthesis

and release of the steroid hormone 20-OH ecdysone from the prothoracic

gland which drives larval instar progression (16,17).

A key difference between larval and adult feeding behaviour is that

adult *Drosophila* feed sporadically, driven by hunger and satiety signals

(18) whereas *Drosophila* larvae accelerate feeding as second instar larvae

and feed continuously till the wandering stage of third instar larvae to

optimise growth. They stop feeding for a few hours prior to pupariation

(19,20). Despite studies identifying several neurotransmitters and

neuropeptides in feeding regulation and their cognate neurons as part of

the feeding connectome in larvae (5) mechanisms that initiate and maintain

persistent feeding in early larval stages are not fully understood. In this

study, whilst characterizing the cellular and molecular phenotypes of null

mutants for the ER-Calcium sensor protein STIM (**St**romal **I**nteraction

Molecule) (21) we identified a novel dopaminergic-neuropeptide connection

that drives persistent feeding and growth in early larvae. Growth deficits

and lethality in STIM mutants appeared to have a focus in dopaminergic

cells (22). However, the cellular basis of STIM function and systemic

phenotypes arising from loss of STIM in dopaminergic neurons remained to

be understood. Here, we show that STIM function is required in a subset of

central dopaminergic neurons for their excitability and dopamine release.

These dopaminergic neurons impact larval growth by providing the

motivation for persistent larval feeding and modulating neuropeptidergic

cells, including the MNSc, to regulate levels of insulin-like peptides.

Results

Reduced food intake and growth deficits in STIM mutant larvae

STIM^{KO} larvae appear normal after hatching but their transition from

first to second instar stages is slower than wild-type animals (**S1A and S1B**

Fig) (22) and as second instars they die gradually between 86h to 326h

after egg laying (**AEL; S1B Fig**). To identify the precise time window when

STIM^{KO} larvae become sickly they were observed over 6h time intervals

from 36h to 90h AEL. Whereas, wild type (*Canton-S* or CS) larvae

transition from 1st to 2nd instar between 42-54h AEL, the same transition in

STIM^{KO} larvae occurs between 60-72h AEL, indicating a delay of 18h (**Fig**

1A and 1B). The delay is followed by an inability to transition to 3rd instar

(**Fig 1C**). *STIM^{KO}* larvae also exhibit retarded growth. At 72h they appear

similar to CS larvae of 60h (**Fig 1D)**. After 72h however, there is a complete

cessation of growth in *STIM^{KO}* larvae (**Fig 1D and 1E**), followed by gradual

loss of viability after 80-86h (**S1B Fig**). From these results, it became

evident that cessation of growth precedes loss of viability in *STIM^{KO}* larvae.

The momentum of larval growth is maintained primarily by cell

growth in the endoreplication tissues (23,24). In a few organs like the brain

and the imaginal discs growth is accompanied by constant cell division.

Normally, at the end of embryonic development, mitotic cells such as a

majority of neuroblasts (NBs) and imaginal disc cells enter a quiescent

state (4,25,26). Postembryonic larval development is initiated in the late

first instar and early second instar stages by cell growth and renewed cell proliferation in the brain and imaginal discs, where it is nutrient dependent (26). Cessation of growth in $STIM^{KO}$ larvae (**Fig 1D and 1E**), suggested a deficit in cell growth and/or cell division. To investigate the status of cell proliferation in $STIM^{KO}$ larvae we chose the well-characterized system of thoracic neuroblasts (27). Upon comparison of thoracic segments of WT and $STIM^{KO}$ larvae at 70-74h (**Fig 1F,** first two columns) it was evident that NBs exited from quiescence and entered the proliferative state in both genotypes. Both the NB marker Deadpan (red) and the post-mitotic cell marker Prospero (blue) appeared normal in $STIM^{KO}$ larvae of 70-74h AEL. Subsequently, at 82-86h, the number of postmitotic cells (Prospero positive) decreased significantly in $STIM^{KO}$ animals as compared to the controls but the number of thoracic neuroblasts remained unchanged (**Fig 1F,** compare third and fourth columns). Upon quantification, the ratio of dividing neuroblasts (Deadpan surrounded by Prospero positive cells) to non-dividing neuroblasts (Deadpan with either no or few Prospero positive cells) changed significantly in 86h aged $STIM^{KO}$ larvae (**Fig 1G and 1H**). To identify the cause underlying the reduced number of postmitotic cells we analyzed different phases of the cell cycle of thoracic neuroblasts in $STIM^{KO}$ larval brains. For this, we used the genetically encoded FUCCI system (28). Here, the G1, S, and G2 phases of interphase are marked by green, red and green+red (yellow) fluorescent tags respectively. At 72-76h, both control and $STIM^{KO}$ showed an asynchronous pattern of division. But at 82-86h, control larvae persisted with the asynchronous pattern, whereas a majority of thoracic neuroblasts in $STIM^{KO}$ animals remained in the G2/M state (**S1C Fig**).

Among other factors, the exit of quiescence and maintenance of neuroblast proliferation at the early second instar stage depends on

nutritional intake (26,29,30). The slow growth and delayed exit from

quiescence suggested that $STIM^{KO}$ larvae may lack adequate nutritional

inputs. As a first step staged larvae were placed on yeast, mixed with a

blue dye and tested for ingestion of food. Even as early as 40-44h AEL

there was a significant reduction of food intake in $STIM^{KO}$ larvae (**Fig 1I,**

quantification in **Fig 1J**). By 80-84h AEL two classes of $STIM^{KO}$ larvae were

evident. One with reduced food intake and others with no food intake (**Fig**

1I and S1D Fig). The proportion of $STIM^{KO}$ larvae with no food intake

reached ~70% by 82-86h AEL (**S1E Fig**). The ability to feed was further

quantified in $STIM^{KO}$ animals by measuring mouth hook contractions

through larval development (31). Control larvae (*CS*) exhibit a steady

increase in mouth hook contractions with age, except prior to and during

larval molts, indicating greater nutrient intake with age. In contrast, increase

in mouth hook movements of $STIM^{KO}$ larvae follows a slower

developmental trajectory, with minimal increase as they progress from first

to second instar larvae and a cessation at 74h AEL, that is further retarded

at 86h (**Fig 1K; videos 1-4)**. Thus, the acceleration of mouth hook

movements observed in *CS* larvae from first to third instar is retarded in

$STIM^{KO}$ larvae before the appearance of growth deficits (**Fig 1I-K**).

suggesting that the consequent nutritional deficits prevent normal growth.

STIM function is required in a subset of larval dopaminergic neurons

To understand how the loss of STIM might affect larval feeding we

identified specific cells that require STIM function for larval growth and

viability. From previous work, we know that knock out of *STIM* in

dopaminergic neurons marked by *THGAL4* (32) leads to larval lethality

(22). Over-expression of a wildtype $UASSTIM^{+}$ transgene, henceforth

referred to as $STIM^{+}$, in dopaminergic neurons marked by *THGAL4* rescued

larval lethality of $STIM^{KO}$ animals to a significant extent (**S2A Fig**). Further

to identify specific dopaminergic neurons that require STIM function for

growth and viability we tested rescue by overexpression of STIM$^+$ in two

non-overlapping subsets of dopaminergic neurons marked by *THC'GAL4*

and *THD'GAL4* (33) and henceforth referred to as *THD'* and *THC'*. Rescue

of $STIM^{KO}$ larvae from 2nd to 3rd instar (~90%) was evident upon over-

expression of *STIM$^+$* in THD' marked neurons (**Fig 2A and 2B**). The rescue

by THD' was equivalent to the rescue by *THGAL4* (20$\pm$2 and 18$\pm$1.5 viable

adults eclosed respectively from batches of 25 larvae; **S2A Fig;** dark and

light green arrows). In contrast rescue by expression of *STIM$^+$* in

THC'GAL4 marked neurons was considerably less (out of batches of 25

animals 5$\pm$1 adults eclosed**;   S2A Fig;** blue arrow), indicating a greater

requirement for STIM function in THD' marked dopaminergic neurons.

Next, we analysed *THD'* driven expression of mCD8GFP and

identified two classes of GFP positive cells in the larval brain. All GFP

expressing cells in the central brain (three cells of DL1 and two cells of DL2

clusters (11,34,35)), appear positive for Tyrosine Hydroxylase (TH) but a

pair of THD' cells in the ventral ganglion (VG) appear TH negative (TH^{-ve})

(**Fig 2C**). To understand the relative contribution of the VG-localised TH^{-ve}

cells to *THD'>STIM$^+$* rescue of $STIM^{KO}$ animals, we restricted *THD'GAL4*

expression to VG localised THD' neurons using *THGAL80* (36) (**Fig 2D**).

Expression of *STIM$^+$* in the VG localised THD' neurons (*THD'GAL4,*

THGAL80) reduced the rescue of $STIM^{KO}$ larvae significantly (**S2A Fig and**

Fig 2E) and was absent in adults (**Fig 2F**). Thus, the rescue of viability in

$STIM^{KO}$ animals derives to a significant extent from brain-specific THD'

dopaminergic neurons.

Ingestion of food and frequency of mouth hook contractions is also

rescued by *THD'>STIM$^+$* expression in $STIM^{KO}$ larvae (**Fig 2G – 2I**). To

further confirm the relevance of STIM function in THD' marked

dopaminergic neurons we knocked down STIM using a previously

characterized STIM RNAi (*dsSTIM*) (37). *THD'>dsSTIM* animals exhibit

delayed larval growth (**Fig 2J and S2B Fig)** and reduced feeding (**Fig 2K**

and 2L) but no larval lethality. Adults however exhibit reduced body weight

(**S2C Fig**). Taken together these data identified a CNS-specific subgroup of

dopaminergic neurons that require STIM function for persistent feeding

during the early stages of larval growth.

Neuronal excitability and dopamine release requires STIM

The status of THD' marked central brain dopaminergic cell clusters,

DL1 and DL2 was investigated next in $STIM^{KO}$ larval brains at 80-84h,

when a few viable organisms are still present despite cessation of growth

and feeding (**Fig 2G**). THD' cells were marked with GFP in controls,

$STIM^{KO}$ and $STIM^{KO}$ animals with STIM$^+$ rescue and the brains were stained

with anti-TH sera. *THD'>mCD8GFP* cells appeared no different in $STIM^{KO}$

as well as STIM$^+$ rescued $STIM^{KO}$ animals at 80-84h AEL when compared

to controls at either 58-62h or 80-84h AEL (**S3A Fig**). Moreover, the

numbers of THD' GFP cells and TH positive cells in the CNS also appeared

identical (**S3B Fig**). Therefore, the loss of STIM does not lead to the loss of

dopaminergic neurons in the larval brain.

In order to test if reduced feeding in $STIM^{KO}$ larvae is indeed due to

a loss in dopamine signalling we measured larval feeding with knockdown

of a key dopamine synthesising enzyme Tyrosine Hydroxylase (TH), in

THD' neurons (*THD'>dsTH*). Knockdown of *TH* led to significantly fewer

mouth hook contractions in larvae at 80-86h AEL (**Fig 3A, Video 5-7**),

indicating reduced feeding, This was accompanied by slower progression

through larval moults and some mortality at each larval stage. Finally out of

a total of 25 just 20±1.2 3rd instar larvae pupated, of which 15 adults

emerged (**S3C Fig)**. In agreement with lower nutrient intake during larval

stages, third instar larvae were smaller in size **(Fig 3B-C**), and gave rise to

adults with significantly reduced body weight (**Fig 3D**).

To understand how loss of STIM in THD' marked neurons might

affect their neuronal function, we investigated properties of excitation. For

this purpose, Potassium Chloride (KCl, 70mM) evoked cytosolic calcium

transients were measured from THD' neurons using the genetically

encoded Ca^{2+} sensor GCaMP6m in *ex vivo* preparations of similarly staged

control (58-62h AEL) and *STIM*^{KO} (70-74h AEL) larvae. We chose these

time points because at 72h *STIM*^{KO} larvae appear healthy and

developmentally similar to control larvae at 60h (**Fig 1D**). THD' cells

responded with similar changes in GCaMP intensity, in control and *STIM*^{KO}

larvae at these time points (**Fig 3E-3G**). However, the ability to evoke and

maintain cytosolic Ca^{2+} transients upon KCl depolarisation was lost in THD'

neurons from the DL1 and DL2 clusters of *STIM*^{KO} larvae at 76-80h AEL

(**Fig 3E-3G**). Overexpression of *STIM*$^{+}$ in THD' cells of *STIM*^{KO} larvae

rescued the KCl evoked Ca^{2+} response in larvae as late as 80-84h AEL (**Fig**

3E-3G).

STIM requirement for maintaining excitability of THD' neurons was

tested further by KCl stimulation of THD' neurons with STIM knockdown

(*THD'>dsSTIM)* from 2nd instar larvae aged 58-62h. Two classes of

responses to depolarisation by KCl were observed. Most cells (70%)

responded with normal or greater changes in intensity as compared to

control cells, whereas in 30% of cells KCl did not evoke a Ca^{2+} transient

(**S3D-S3F Fig**). We attribute this heterogeneous response to differential

STIM knockdown by the RNAi in individual THD' cells and a potential effect

of STIM knock-down on ER-Ca^{2+} homeostasis (see below).

These data suggest that loss of STIM affects membrane excitability

properties and the ability of THD' neurons to respond to stimuli. This idea

was tested directly by the expression of transgenes that alter membrane

potential. Over-expression of an inward rectifier K^+ channel, Kir2.1 in THD'

neurons, that is known to hyperpolarise the plasma membrane (38),

resulted in developmental delays followed by the lethality of second and

third instar larvae (**S4A Fig**). Further, overexpression of a bacterial Na^+

channel NaChBac (39) in THD' neurons of $STIM^{KO}$ larvae evinced a weak

rescue of developmental deficits, including the transition to third instar

larvae (4.4$\pm$0.4) and adult viability (2.4$\pm$0.6 from of batches of 15 animals**;**

Fig. 3H-3K). Though weak, NaChBac's rescue was consistent. We attribute

the variability to a stochastic effect of NaChBac in THD' cells. This is also

evident from the variable extent of rescue in growth observed in $STIM^{KO}$;

THD'>NaChBac larvae (**Fig 3I and 3J**). Alternatively, in addition to

neuronal excitability, STIM might affect other cellular functions such as ER

stress, that are not alleviated by expression of NaChBac, resulting in the

weak rescue. Control animals with overexpression of *NaChBac* in THD'

neurons exhibit delayed pupariation (**S4B Fig**).

Neuronal excitability is required for neurotransmitter release at

presynaptic terminals. We hypothesized that dopamine release from THD'

neurons might be affected in $STIM^{KO}$ larvae. To test this idea, we identified

the pre-synaptic (green) and post-synaptic (red) terminal regions of THD'

neurons by marking them with *SyteGFP* and *Denmark* respectively (**Fig

4A)** (40)]**.** Analysis of pre-synaptic regions (SyteGFP expression) identified

three distinct areas in the CNS. One at the centre of the CNS

corresponding to the mushroom body (**Fig 4A;** asterisk), the second as a

branched form in the basomedial protocerebrum of the CNS (**Fig 4A;**

arrowhead) and the third one consisting of punctae spread across the

oesophageal regions where brain-gut interactions take place (**Fig 4A;**
hash). Based on the projection patterns observed we speculate that cells
marked by THD' correspond to DL1-2, DL1-5, DL1-6 from the DL1 cluster
and DL2-2 and DL2-3 from the DL2 cluster (35).

Next, dopamine release was measured in the most prominent
presynaptic areas of THD' neurons, corresponding to the MB and the
basomedial protocerebrum, by change in fluorescence of the genetically
encoded fluorescent GPCR-activation-based-DopAmine sensor (GRAB$_{DA}$**)**
(41)**.** Dopamine release in THD' neurons of $STIM^{KO}$ larvae at 76-80h is
significantly attenuated as compared with controls (**Fig 4B and 4C**).
Importantly, overexpression of STIM$^+$ in THD' neurons rescued dopamine
release, though with altered dynamics from control animals (**Fig 4B and**
4C; see discussion). We chose to measure dopamine release in 76-80h
$STIM^{KO}$ larvae because THD' neurons in their brains no longer responded
to KCl evoked depolarization (**Fig 3B**) even though the larvae appear
normal (**Fig 1D**). Dopamine release was stimulated by Carbachol (CCh), an
agonist for the muscarinic acetylcholine receptor (mAChR), that links to
Ca^{2+} release from ER-stores through the ER-localised IP$_3$ receptor (42) and
is expressed on THD' neurons (43). CCh-induced changes in ER-Ca^{2+} were
tested directly by introducing an ER-Ca^{2+} sensor (44) in THD' neurons (**Fig**
4D and E). Though ER-Ca^{2+} release, in response to CCh could be
measured in just 7 out of 23 cells, the subsequent step of ER-store refilling,
presumably after Store-operated Ca^{2+} entry into the cytosol through the
STIM/Orai pathway, could be observed in all control THD' cells (58-62h
AEL), whereas it was absent in THD' neurons from $STIM^{KO}$ brains (76-80h
AEL**; Fig 4E**). The ER-Ca^{2+} response was rescued by over-expressing
STIM$^+$ in THD' neurons (**Fig 4D, and E**). Taken together our data establish
an important role for STIM-dependent ER-Ca^{2+} homeostasis in maintaining

optimal dopamine release from THD' neurons in turn required for the larval

drive to feed constantly.

Because late third instar larvae stop feeding (3,19,20,45) we

hypothesized that CCh-stimulated Ca^{2+} responses in THD' cells might

change in wandering stage third instar larvae. To test this idea we

monitored carbachol-stimulated GCaMP activity in THD' neurons from wild-

type larvae at 96h AEL (mid 3^{rd} instar), 120h AEL (early wandering stage)

and 124h AEL (late wandering stage). A robust GCaMP response was

observed at 96h, whereas at the beginning of the wandering stage (118-

122h AEL), the peak response was both reduced and delayed. A further

delay in the peak response time was observed in late wandering stage

larvae (122-126h AEL) (**S4E and S4F Fig**). Thus with gradual cessation of

feeding in late third instar larvae, the dynamics of CCh-stimulated Ca^{2+}

responses in THD' neurons also undergo changes (see discussion).

THD' dopaminergic neurons both activate and inhibit specific

neuropeptidergic cells

A role for neuropeptides in modulating larval feeding has been

identified previously (5–7) Based on these findings we postulated that

dopamine release from THD' cells might modulate neuropeptidergic cells in

the larval brain. This idea was tested by optogenetic stimulation of red-

shifted Channelrhodopsin (*CsChrimson*) (46) expressing in THD'

(*THD'>UAS-Chrimson*) cells followed by Ca^{2+} imaging from GCaMP6f

expressing neuropeptide cells *(C929LexA>LexAopGCaMP6f)* (**Fig 5A**).

Upon optogenetic activation of THD' neurons, a change in cellular Ca^{2+}

signals was observed in a total of 64 peptidergic neurons from 9 brains,

including some lateral neurosecretory cells (LNCs), median neurosecretory

cells (MNCs), and regions of the suboesophageal zone (SEZ) (**Fig 5B and**

5C). Elevated Ca^{2+} signals were observed in a subset of neuropeptidergic

cells (**Fig 5B - 5D,** yellow asterisks, n=24), whereas in some cells Ca^{2+}

signals were reduced (**Fig 5B, 5C and 5E,** red asterisks, n=40**)**. There was

no change in cellular Ca^{2+} upon optogenetic stimulation in the remaining

cells with visible GCaMP6f fluorescence (n=35). Upon cessation of

optogenetic activation of THD' neurons Ca^{2+} levels returned to baseline (**Fig**

5D and 5E). Individual cells exhibit either activatory or inhibitory responses

upon repeated optogenetic stimulation with pulses of red light **(Fig 5F (i)**

and (ii)), indicating that THD' cells evoke specific responses of either

stimulation or inhibition based on the class of neuropeptide cells. Specificity

of the *THD'>Chrimson* evoked response was further confirmed by testing

brains from larvae that were reared without all *trans*-Retinal (ATR, a

cofactor for the function of Channelrhodopsin) (47) and by imaging in the

absence of light (**Fig 5G**).

STIM expression in THD' neurons regulates the expression of insulin-

like peptides

Though optogenetic stimulation of THD' cells shows dopaminergic

modulation of neuropeptidergic neurons (**Fig 5A and 5B**) it does not allow

identification of specific neuropeptides that function downstream of the

THD' neurons. Optogenetic activation of THD' and Ca^{2+} responses in

neuropeptidergic cells (marked by *C929LexA>LexAopmCherry*) helped

identify three clusters of neuropeptidergic cells that are downstream of

THD' neurons and include the well-characterised ilp expressing MNC

cluster (48), as a putative target of THD' neurons (**Fig 5B**). Analysis of an

RNAseq experiment, conducted independently to identify changes in gene

expression in brains from second instar *STIM^{KO}* larvae (72-76h AEL) with

developmentally   comparable   CS   brains   (58-62h;

https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE202098;   to   be

published elsewhere), found significant changes in expression of *ilp2*, *ilp3*

and *ilp5*. Whereas *ilp2* and *ilp5* were significantly downregulated, *ilp3* was

upregulated more than 5 fold (**Fig 6A**). The differential regulation of *ilp2, -3*

and *-5* expression in *STIM^{KO}* brains was further validated by qPCRs (**Fig**

6B). Importantly, expression of *ilp3* and *ilp5* were restored back to normal

in brains from *STIM^{KO}* larvae, rescued by overexpression of *STIM^{+}* in THD'

neurons (**Fig 6B**). During normal larval development *ilp3* transcript levels

are low in actively feeding larvae (L2 and L3, 12h) and are gradually

upregulated in later stages of L3 as larvae stop feeding (**Fig 6C**; DGET (49)

). Based on the up-regulation of *ilp3* in *STIM^{KO}* larvae (**Fig 6A and 6B**), we

hypothesised that knock down of *ilp3* in MNSc might rescue *STIM^{KO}* larvae.

Indeed, partial rescue of larval lethality in 2^{nd} instar larvae followed by their

transition to 3^{rd} instar larvae (5 ± 0.5) was observed (**Fig 6D - G**). A few of

the rescued 3^{rd} instar larvae grew to full size, and pupariated (**Fig 6E**, L3b

type larvae) and some even eclosed as adults (out of 3 batches of 25

animals 2.3 ± 0.3; **Fig 6D   and 6E**). The partial rescue observed by *ilp3*

knockdown may be due to roles of additional dopamine-modulated

neuropeptides plus the lower expression of ilp2 and ilp5 in *STIM^{KO}* animals

(**Fig 6A and 6B**) because both ilps are growth signals in 2^{nd} and 3^{rd} instar

larvae. A small proportion of *STIM^{KO}* larvae rescued by *ilp3* knockdown

appear significantly larger than control larvae (**Fig 6E and 6F,** L3b),

indicating that loss of ilp3 can rescue growth in some animals. This idea is

further supported by knock-down of *ilp3* in MNSc from wildtype animals that

resulted in slower transition to pupariation and larger pupae, and over-

weight adults (**Fig 6H-J; S5A-B Fig**). Conversely, over-expression of *ilp3* in

MNSc resulted in delayed larval transition from L2 to L3 and smaller sized

larvae (**Fig 6K-M)** similar to delayed larval development in ilp5 knockdown

animals (**S5A Fig**). However, overexpression of ilp3 had no effect on

feeding as indicated by measurement of larval mouth hook movements (**Fig**

6N) even though adults were of reduced body weight (**S5B Fig**). Taken

together, these data show that dopamine signals from THD' cells are

required to maintain normal expression of growth promoting ILPs (*ilp2* and

ilp5) and repress expression of *ilp3* that appears to function as an anti-

growth signal.

A direct synaptic connection between central dopaminergic neurons

and the MNSc has not been reported (5,50) and is likely not supported by

data that mapped synaptic connections in the larval brain(35,50–52).

However, a neuromodulatory role for dopamine is documented where it can

effect a larger subset of neurons, beyond direct synaptic partners, by

means of diffusion aided volumetric transmission (53–56). To test this

possibility we began by measuring larval developmental transitions in

animals with knockdown of three dopamine receptors, *Dop1R1*, *Dop2R2*

and *DopEcR* in the MNSc. Among these, knockdown of *Dop1R1* delayed

development, reduced larval and adult viability (**Fig 7A**) and negatively

impacted growth **(Fig 7B and 7C)**. A weaker phenotype of delayed

development and loss of viability was observed with knockdown of

DopEcR, whereas larvae with knockdown of Dop2R2 appeared normal (**Fig**

7A). The response of MNSc to dopamine was tested next. Brains

expressing a Ca^{2+} sensor in the MNSc (*MNSc>GCaMP6*m)   were

stimulated with dopamine in the presence of a Na_v blocker Tetrodotoxin

(57), to prevent extraneous neuronal inputs. Of the seven targeted MNSc in

one hemisphere, we observed consistent activation of one cell whereas

dopamine addition inhibited the Ca^{2+} response in three to four cells (**Fig 7D**;

quantified in **Fig 7E**). No changes in the Ca^{2+} responses of MNSc were

observed in the absence of dopamine (**Fig 7D and 7E**).

Discussion

In *Drosophila*, as in other holometabolous insects, growth is restricted to the larval stages. In early stages of larval development cells exit mitotic quiescence and re-enter mitosis resulting in organismal growth (1,25,58).This change is accompanied by an increase in the feeding rate of the organism so as to provide sufficient nutrition for the accompanying growth in organismal size. In $STIM^{KO}$ larvae we observed a loss of this ability to feed persistently starting from early second instar larvae. The focus of this feeding deficit lies in a subset of central dopaminergic neurons that require STIM function to maintain excitability. Importantly, these dopaminergic neurons communicate with multiple neuropeptidergic cells in the brain including the MNSc (**Fig 5D**) to regulate appropriate changes in larval feeding behaviour and growth through development.

STIM and excitability of dopaminergic neurons

We identified the THD' cells as critical for larval feeding from their inability to function in the absence of the SOCE regulator STIM. Loss of excitability and the absence of dopamine release from THD' cells in $STIM^{KO}$ larvae (**Fig. 3A-3C and Fig 4B-4D**) suggests that voltage-dependent receptor activity is required to maintain growth in early 2nd instar larvae. Changes in expression of ion channels and presynaptic components have been observed earlier upon knockdown of STIM in *Drosophila* and mammalian neurons (59,60). Moreover, loss of STIM-dependent SOCE in *Drosophila* neurons effects their synaptic release properties (61). Partial rescue of viability in $STIM^{KO}$ organisms by over-expression of a bacterial sodium channel NaChBac (**Fig 3D-3G**) and restoration of dopamine

471 release upon rescue by STIM$^+$ (**Fig. 4B and 4C**) supports the idea that

472 STIM-dependant SOCE maybe required for appropriate function and/or

473 expression of ion channels and synaptic components in THD' neurons.

474 Changes in ER-Ca^{2+} (**Fig 4D and 5E**) suggest that STIM is also required to

475 maintain neuronal Ca^{2+} homeostasis.

Dopaminergic control of larval food seeking

Whilst mechanisms that regulate developmental progression of

Drosophila larvae have been extensively studied, neural control of

essential changes in feeding behaviour that need to accompany each larval

developmental stage have not been identified previously. Artificial

manipulation of activity in the central dopaminergic neuron subset studied

here (THD'), either by expression of an inward rectifying potassium channel

(Kir2.1) (**S4A Fig**) or the bacterial sodium channel NaChBac (**S4B Fig**),

suggests an important role for THD' neurons during larval development.

This idea is supported by the altered dynamics of muscarinic acetylcholine

receptor (mAChR) stimulated Ca^{2+} release observed in THD' neurons

between early, mid and late third instar larvae when larval feeding slows

down and ultimately stops (**S4C and S4D Fig**) and re-iterates that

signaling in and from these neurons drives larval feeding whereas lower

carbachol-induced Ca^{2+} responses signal cessation of feeding. A weaker

rescue of *STIM^{KO}* larvae is also obtained from *STIM$^+$* expression in the

THC' neuron subset. Taken together these observations suggest a

neuromodulatory role for dopamine, where DA release from THD' neurons

has a greater influence on feeding than the DA release from THC' neurons,

possibly due to the DL1 and DL2 cluster (among THD' marked neurons)

receiving more feeding and metabolic inputs (7,35,58). A role for cells

other than THD', in maintaining kinetics of dopamine release required for

feeding behaviour are also indicated because expression of STIM$^+$ in THD'

neurons did not revert kinetics of dopamine release to wild type levels (**Fig**

4C). The prolonged dopamine release observed in wild-type THD' neurons

may arise from synaptic/modulatory inputs to THD' neurons from other

neurons that require STIM function.

Though the cells that provide cholinergic inputs to THD' cells have

not been identified it is possible that such neurons sense the nutritional

state. In this context, two pairs of cells in the THD' subset also motivate the

search for food in hungry adult *Drosophila* (62). Starved flies with knock

down of the mAChR on THD' neurons exhibit a decrease in food seeking

behaviour (43). Cholinergic inputs to THD' neurons for sensing nutritional

state/hunger may thus be preserved between larval and adult stages.

Interestingly, dopamine is also required for reward-based feeding,

initiation, and reinforcement of feeding behaviour in mice (63). These

findings parallel past studies where prenatal mice genetically deficient for

dopamine (DA$^{-/-}$), were unable to feed and died from starvation. Feeding

could however be initiated upon supplementation with L-DOPA (64)

allowing them to survive. More recent findings show that dopaminergic

neurons in the ventral tegmental area (VTA), and not the substantia nigra,

drive motivational behaviour and facilitate action initiation for feeding in

mice (65).

Dopaminergic control of neuropeptide signaling

Both activation and inhibition of specific classes of neuropeptidergic

cells by optogenetic activation of THD' cells suggests a dual role for

dopamine possibly due to the presence of different classes of DA

receptors. The *Drosophila* genome encodes four DA receptors referred to

as Dop1R1, Dop1R2, DD2R and a non-canonical DopEcR (66). Dop1R1,

Dop1R2 and DopEcR activate adenylate cyclase and stimulate cAMP

signaling whereas DD2R is inhibitory (66). Cell specific differences among

dopamine receptors have been observed in adults. Down regulation of

Dop1R1 on AstA and NPF cells shifted preference towards sweet food

whereas down regulation of *DopEcR* in DH44 cells shifted preference

towards bitter food (67). In third instar larvae a dopaminergic-NPF circuit,

arising from central dopaminergic DL2 neurons, two cells of which are

marked by *THD'GAL4* (**Fig 2E**), motivates feeding in presence of appetitive

odours (6,45). The dopamine-neuropeptide axis identified here

demonstrates a broader role for dopamine in regulating neuropeptide

release and/or synthesis, in the context of larval feeding behaviour,

perhaps similar to the mammalian circuit described above.

Of specific interest is the untimely upregulation of *ilp3* transcripts in

$STIM^{KO}$ larvae. Rescue of lethality in $STIM^{KO}$ larvae either by bringing back

activity to THD' neurons or by reducing ilp3 levels suggests an

interdependence of Dopamine-Insulin signaling that is likely conserved

across organisms (68–73). Our data suggest that ilp3 expression is

suppressed during the feeding and growth stages of larvae (**Fig 6H-6M**),

and once enough nutrition accumulates expression of ilp3 is up-regulated,

concurrent with a reduction in carbachol-induced Ca^{2+} signals in THD'

neurons, possibly followed by upregulation and release of ilp3. The idea of

ilp3 as a metabolic signal whose expression is antagonistic to larval growth

is supported by the observation that knock-down of ilp3 in the MNSc leads

to larger pupae in wild type animals and larger larvae in $STIM^{KO}$ (**Fig 6F**

and G). To our knowledge this is the first report of ilp3 as a larval signal

that is antagonistic to growth. Given that *Drosophila* encode a single Insulin

receptor for ilp2, ilp3 and ilp5 (74) the cellular mechanism of ilp3 action

remains to be elucidated. Possibly, ilps with different affinity for the insulin

receptor stimulate different cellular subsets and/or different intracellular

signaling mechanisms, including ecdysone signaling that is essential for

larval transition to pupae (74). Interestingly, in $STIM^{KO}$ larval brains there is

a significant increase in expression of the Insulin Receptor (**S1 Table**).

Expression of other neuropeptides did not show significant changes

in $STIM^{KO}$ larval brains (**S1 Table**), suggesting that for neuropeptidergic

cells in the LNC and SOG dopamine signals alter release properties rather

than synthesis. However, we were unable to identify specific neuropeptides

for cells in the LNC and the SOG that responded upon activation of THD'.

The importance of dopamine for multiple aspects of feeding

behaviour is well documented in mice (63,64). Of interest are more recent

findings linking dysregulation of dopamine-insulin signaling with the

regulation of energy metabolism (75) and the induction of binge eating

(76,77). The identification of a simple neuronal circuit where dopamine-

insulin signaling regulates feeding and growth could serve as a useful

model for investigating new therapeutic strategies targeted towards the

treatment of psychological disorders for obesity and metabolic syndrome

(75,76).

Material and methods

Fly rearing and Stocks:

Drosophila strains were reared on standard cornflour agar media

consisting of 80 g corn flour, 20 g glucose, 40 g sugar, 15 g yeast extract, 4

ml propionic acid, 5 ml *p*-hydroxybenzoic acid methyl ester in ethanol, 5 ml

ortho butyric acid in 1l at 25°C, unless otherwise specified, under a 12:12

hr light : dark cycle. In all studies the *Canton S* (*CS*) strain was used as a

wild-type control and CRISPR-Cas9 generated deficiency for *STIM* referred

to as *STIM^{KO}* served as a null mutant for the *Drosophila STIM* gene (22).

Details of other fly lines used are provided in Table 1 below.

Staging:

Synchronized larvae of the appropriate ages as described below

were collected and transferred to agar less media containing yeast (4gm),

sucrose (8gm), cornflour (16gm), Propionic acid (1ml) and 0.05gm of

Benzoic acid in 1ml of absolute alcohol. The number of viable organisms

and the developmental stage were scored at specific time points as

mentioned below and in the figures and figure legends.

Larval staging experiments were performed to obtain lethality and

developmental profiles of the indicated genotypes as described previously

(78). Depending on the experiment, timed and synchronized egg-laying was

done either for 6h to allow development profiling at 60-66h, 80-86h, 128-

136h, 176-182h and 320-326h after egg laying or for 2h at 35-37h(36h), 41-

43h(42h), 47-49h(48h), 53-55h(54h), 59-61h(60h), 65-67h(66h), 71-

73h(72h), 83-85h(84h), and 89-91h(90h) after egg laying for identifying a

lethality window between 36-90h. Larvae were collected at either 60–66h or

35-37h after egg laying (AEL) in batches of 25 (for developmental profile) or

10 (for lethality window). They were screened and staged subsequently.

Heterozygous larvae were identified using dominant markers (*FM7iGFP*,

TM6Tb, and *CyOGFP*) and removed. Each batch of larvae was placed in a

separate vial and minimally three vials containing agar-less media were

tested for every genotype at each time point. The larvae were screened at

the indicated time points for the number of survivors and stage of

development, determined by the morphology of the anterior spiracles and

mouth hooks (79). Experiments to determine the viability of experimental

genotypes and their corresponding genetic controls were performed

simultaneously in all cases. Larval images were taken on the MVX10

Olympus stereo microscope using an Olympus DP71 camera.

Feeding assay:

Feeding assay was performed at specific developmental time points

in larvae (40-44h, 58-62h, 80-84h AEL) of the specified genotypes. Larvae

were placed in a 35mm punched dish with coverslip at the base thus

creating a small depression in the centre of the coverslip. In this depression

a coin sized cotton swab was placed containing 4.5.% of yeast solution with

0.25% eriogluasin dye (blue dye). For scoring the number of larvae that fed,

30 larvae per plate were taken and incubated in the feeding plate for 4hrs at

25^{o}C. Larvae were removed from the paste, washed, collected and scored

for presence of blue dye (Dye^{+ve}) and absence of blue dye within the gut

(Dye^{-ve}).

For quantification of ingested blue dye, 12-15 larvae were incubated

for 2h in yeast paste with the blue dye. Larvae were removed from the yeast

paste and 10 larvae with blue dye in the gut were washed, and

homogenized in 50µl of cold 1xPBS. The homogenate was spun at 5k for 2

minutes in a table top Eppendorf centrifuge. The supernatant (2µl) was

taken for quantification of protein using the Thermo scientific Pierce Protein

assay kit, Cat#23227. Optical density (OD) at 625nm as a measure of

ingested blue dye was measured from 30µl of the homogenate. Due to

variation in larval sizes between control and experimental samples, the OD

was normalized to whole larval protein concentration (µg/µl). OD was

obtained using the SkanIt Software 6.1.1 RE for Microplate Readers RE,

ver. 6.1.1.7. Larval imaging and processing was performed on the Olympus

MVX10 stereo microscope using FIJI software.

Larval imaging and measurement:

Staged larvae from specified genotypes were collected at specific development time points, anesthetized on ice for 1h and mounted with ice cold HL3 buffer. The mounted larvae were imaged immediately using an Olympus MVX10 stereo microscope. For measurement of larval length from mouth hook to tip of the posterior spiracle FIJI software was used. A minimum of 10-15 larvae were taken per genotype for length analysis.

Quantification of larval mouth hook contractions:

Mouth hook contractions were measured by placing 1-3 appropriately staged larvae in a drop of 2% yeast solution in a petridish. Videos were taken for 30 seconds on an Olympus MVX10 stereo microscope. For each genotype a minimum of 10 animals were imaged. Mouth hook contractions were counted manually from the visualised videos.

Pupal volume measurement:

Pupal volume was measured by obtaining the width and height of each pupal image from the Olympus MVX10 stereo microscope. A formula for obtaining the volume of a cylinder, ($\pi r^2 h$) was applied to calculate the volume (80).

Adult fly weight measurement:

For weight measurement of adult flies, 10 flies (5 females and 5 males) of the appropriate genotype were taken 6 – 10 hr post-eclosion and weighed after placing them in a small Eppendorf tube. Thereafter, the weight of the same empty tube was measured. Fly weights were

calculated by subtracting the weight of the tube from the total weight of

flies + tube. A minimum of five such measurements were performed for

each genotype.

Immunohistochemistry:

Larval brains were dissected in ice-cold 1xPBS and fixed with 4%

paraformaldehyde in 1xPBS on the shaker for 20mins at room

temperature. Fixed brains were washed with PBTx (0.3% TritonX-100 in

1XPBS) 3-4 times at 10minutes intervals, blocked with 5% normal goat

serum (NGS) in PBTx for 2hrs at room temperature, and incubated with

primary antibodies diluted in 5%NGS+PBTx at the appropriate

concentration as mentioned below, overnight at 4^{o}C. Antibody solution was

removed and re-used upto three times. Brains were washed with PBTx 3

times at 10 minute intervals followed by incubation with secondary

antibodies at the dilutions described below, for 2hrs at room temperature

and three washes in PBTx of 10minute intervals each. Brains were

mounted in 70% glycerol diluted in 1xPBS. Confocal images were

acquired by using FV3000 LSM and the Fluoview imaging software.

Following primary antibodies were used: Chick anti-GFP (1:8000,

Abcam Cat#13970 RRID: AB_300798), mouse anti-Prospero (1:100, DSHB

Cat# Prospero (MR1A), RRID:AB_528440), Rabbit anti-Deadpan (1:400,

Abcam– ab195172), Mouse anti-TH (1:5, ImmunoStar Cat#22941; RRID:

AB_572268), Mouse anti-bruchpilot (NC82) (dilution1:50, monoclonal, Gift

from Eric Buchner, University of Wuerzburg, Germany).

Following secondary antibodies were used :Goat anti-Chicken IgY

(H+L), Alexa Fluor 488 (Thermo Fischer Scientific, Cat#A-11039; RRID:

AB_2534096), Goat anti-Rabbit IgG (H+L), Alexa Fluor 568 (Thermo

Fischer Scientific Cat# A-11011; RRID: AB_143157), Goat anti-Mouse IgG

(H+L) Alexa Fluor 633 (Thermo Fischer Scientific Cat#A-21052; RRID:

AB_2535719), Goat anti-Rabbit IgG (H+L) Alexa Fluor 594 (Thermo Fischer

Scientific Cat#A-11037; RRID: AB_2534095).

Ex-vivo imaging of the larval brain:

GCaMP signals were obtained from appropriately aged larval brains

dissected from the specified genotypes and dissected in hemolymph like

saline (HL3) (70mM NaCl, 5mM KCl, 20mM MgCl2, 10mM NaHCO3, 5mM

trehalose, 115mM sucrose, 5 mM HEPES, 1.5mM Ca2+, pH 7.2).

Dissected brains were transferred to a 35mm punched dish with a cover

slip adhered to the bottom. Brains were embedded in ~5µl of 0.8% low

melt agarose (Invitrogen, Cat#16520-100) and bathed in 86µl of HL3.

Images were acquired as a time series on an XY plane at an interval of

2sec using a 20X-oil objective on an Olympus FV3000 inverted confocal

microscope (Olympus Corp., Japan). For KCl stimulation, at the 40^{th} frame,

7µl of HL3 was added and at the 80^{th} frame 7µl of 1M KCl was added. The

final concentration of KCl in the solution surrounding the brain was 70mM.

For stimulation with Carbachol (Sigma Aldrich Cat# C4382), 10µl of HL3

was added at the 40^{th} frame followed by 10µl of 100mM Carbachol at the

80^{th} frame. Final carbachol concentration was maintained at 0.5mM Ca^{2+}

responses were imaged till the 300^{th} frame (600sec).

Changes in ER-Ca^{2+} were measured using an ER-GCaMP-210

strain (44). The brain sample was prepared as above. Images were

acquired as a time series on an XY plane at an interval of 1 sec using a 20X

oil objective on an Olympus FV3000 inverted confocal microscope

(Olympus Corp., Japan). For Carbachol stimulation, 10µl of HL3 was added

at the 50^{th} frame and 10µl of 100mM of Carbachol was added at the 100^{th}

frame. Final carbachol concentration was maintained at 0.5mM. Images

were obtained for 600 frames (600 secs).

Larval brain expressing *MNSc>GCaMP* is used for Dopamine (DA)

(Sigma, Cat#H8502) stimulation. Dissected brains were transferred to a

35mm punched dish with a cover slip adhered to the bottom. Brains were

embedded in ~5μl of 0.8% low melt agarose (Invitrogen, Cat#16520-100)

and bathed in 80μl of HL3 having 1uM of TTX. Images were acquired as a

time series on an XY plane at an interval of 1.5sec using a 20X-oil objective

on an Olympus FV3000 inverted confocal microscope (Olympus Corp.,

Japan). At the 30th frame, 10μl of HL3 was added and at the 60th frame 10μl

of 1mM DA was added. Ca^{2+} responses were imaged till the 250th frame

(450sec).

Analysis of Optogenetic signals:

Larvae from the specified genotypes were reared in fly media

containing 0.2mM ATR (All-trans retinal Sigma-Aldrich Cat#R2500). Brain

samples were prepared as mentioned above. Images were acquired as a

time series on an XY plane at an interval of 2 sec/frame using a 20X oil

objective on an Olympus FV3000 inverted confocal microscope (Olympus

Corp., Japan). For optogenetic stimulation of *CsChrimson*, a 633nm LED

(from Thor labs) was used and GCaMP6f fluorescent images were obtained

simultaneously using a 488nm laser line so as to measure changes in

cytosolic Ca^{2+} upon CsChrimson activation. Image acquired till 200th frames

(400 secs).

A minimum of 6 independent brain preparations were used for all live

imaging experiments and the exact number of cells imaged are indicated in

the figures. Raw fluorescence data were extracted from the marked ROIs

using a time series analyser plugin in Fiji. ΔF/F was calculated using the

following formula for each time point (t): ΔF/F = $(F_t/F_0)/F_0$, where F_0 is the

average basal fluorescence obtained from the first 40 frames.

RNA isolation and quantitative PCR

Larval brains from appropriate genotypes and their respective age

were dissected in ice cold 1XPBS. Larval brains (10-15) were transferred to

a microtube containing 500µl TRIzol (Life Technologies), and vortexed

immediately for 10-15secs. After vortexing the samples were either

processed within 30 min or stored at −80 °C for processing for up to 2

weeks. RNA was isolated following manufacturer's protocol for TRIzol.

Concentration and integrity of RNA was determined on a bioanalyzer

(Agilent 2100). RNA (1µg) was taken for cDNA synthesis. DNAse treatment

and first strand synthesis was performed as previously described (81).

Reverse Transcription followed by PCR (RT-PCR) was performed in a

reaction mixture of 25 µl with 1 µl of the cDNA. Quantitative real time PCRs

(qPCRs) were performed in a total volume of 10µl with Kapa SYBR Fast

qPCR kit (KAPA Biosystems) on a QuantStudio 3 Real-Time PCR system.

Duplicates were performed for each qPCR reaction. Minimum of three

biological replicates were taken for each qPCR reaction. *rp49* was used as

the internal control. The fold change of gene expression in any experimental

condition relative to wild-type was calculated as $2^{-\Delta\Delta Ct}$, where ΔΔCt = (Ct

(target gene) − Ct (rp49))$_{Expt.}$ − (Ct (target gene) − Ct (rp49))$_{Control.}$

Sequences of PCR primers used are as follows:

rp49

F_5'CGGATCGATATGCTAAGCTGT3'

R_5'GCGCTTGTTCGATCCGTA3'

dilp2

F_5'CCATGAGCAAGCCTTTGTCC3'

R_5'TTCACTGCAGAGCGTTCCTTG3'

dilp3

F_5'ACTCGACGTCTTCGGGATG3'

R_5'CGAGGTTTACGTTCTCGGCT3'

dilp5

F_5'ACTCACTGTCGAGCATTCGG3'

R_5'GAGTCGCAGTATGCCCTCAA3'

Quantification and Statistical Analysis

All bar graphs and line plots show the means and standard error of

means. In boxplots, horizontal lines in the box indicate median, box limits

are from 25th-75th percentiles, and individual data points are represented by

closed circles (unless otherwise specified in the figure legends). Unpaired

student t-Test (for two genotypes) and one way ANOVA followed by post-

hoc Tukey's significance test (for data with multiple genotypes) was

performed to calculate P values, given for all figures in **S2 Table.** All

graphs were plotted using Origin 8.0 software. Origin 7.5 MicroCal, Origin

Lab, Northampton, MA, USA N/A, Fiji Open access (RRID:SCR_002285).

Diagrammatic representative images are made with help of Biorender

website (https://app.biorender.com).

Table 1: Fly strains

Fly line	Description	Source
Canton S	Wild type	
STIM^{KO}	Null mutant for STIM gene generated with help of	Generated in the lab

	CRISPR-Cas9 gene editing technique	
THGAL4	Dopaminergic GAL4 driver marks TH (Tyrosine Hydroxylase) positive cells in brain and hypoderma	Serge Birman CNRS, ESPCI Paris Tech, France
THGAL80	Inhibits Gal4 expression in Dopaminergic cells	Toshihiro Kitamoto, University of Iowa, Carver College of Medicine.
THD'GAL4	Marks subgroup of Dopaminergic cells in larval brain	Mark N Wu, Johns Hopkins University, Baltimore
THC'GAL4	Marks dopaminergic cells of larval brain	
C929GAL4	Marks neuropeptidergic cells	P. H. Taghert, Washington University
c929LexA (DimmLexA)	Marks neuropeptidergic cells	Michael Texada, Janelia Farms, USA
transTango	Marks Post synaptic neurons with RFP	Michael Rosbash, Brandeis University, USA
UASSTIM+	*Stim* wildtype cDNA under UAS control	Generated in the lab (82)
UASGRAB$_{DA}$	A Genetically Encoded Fluorescent Sensor for	Yulong Li, Peking University School of

	Dopamine	Life Sciences, Beijing, China
LexAopmCherry	Expresses mCherry under LexAop control	Claude Desplan, New York University, USA
UASmCD8GFP	Expresses membrane tagged GFP under UAS control	RRID: BDSC_5130
UASKir2.1	Prevents membrane depolarization	RRID: BDSC_6595
UASGCaMP6m	Ca^{2+} Sensor with intermediate kinetics expresses under UAS	RRID: BDSC_42748 from Douglas Kim
LexAopGCaMP6f	Ca^{2+} Sensor with faster kinetics expresses under LexAop	RRID: BDSC_44277
UASSyteGFP,Denmark	Syt-GFP marks presynaptic terminal and Denmark (RFP) marks post synaptic part of the neuron.	RRID: BDSC_33065
UASCsChrimson	Optogenetically encoded cation channel that is activated at 625nm	RRID: BDSC_55136
InscGAL4	Marks neuroblasts of larval brain with GAL4	RRID: BDSC_8751
UASFUCCI	Marks different phases of cell cycle with fluorescent markers	RRID: BDSC_55100
UASNaChBac	Increases sodium conductance activates the neuron	RRID: BDSC_9468

UASERGCaMP-210	ER specific Ca^{2+} sensor	Cahir O'Kane, Cambridge University, UK
dsSTIM (III)	UAS-RNAi against Stim gene	VDRC_47073
dsTH (III)	UAS-RNAi against *Tyrosine hydroxylase* (TH) gene	RRID: BDSC_25796
UASDicer2(X)	Enhancer for RNAi	RRID: BDSC_24648
MNScGAL4 (Dilp2GAL4)	Marks ilp producing MNSc cells of larval brain	RRID: BDSC_37516
UASilp3$^+$	Over expression of wildtype ilp3 gene	Ernst Hafen's lab
UASdsilp3	RNAi for ilp3 gene	RRID: BDSC_33681
UASDop1R1RNAi	RNAi for Dop1R1 receptor	RRID: BDSC_62193
UASDopEcR RNAi	RNAi for DopEcR receptor	RRID: BDSC_31981
UASDop2R2RNAi	RNAi for Dop2R receptor	RRID: BDSC_36824
THD'GAL4, THGAL80/CyO	Recombinant line made for this study	
THD'GAL4, UASGCaMP6m/CyO		
c929LexA, LexAopGCaMP6f, LexAopmCherry/TM6Tb		
THD'GAL4, UASCsChrimson/CyO		
STIM^{KO}; +; THGAL4/TM3SerG	Strains made for this study	
STIM^{KO};		

THD'GAL4/CyOG; +	
STIM^{KO}; *THC'GAL4/CyOG; +*	
STIM^{KO}; *UASSTIM/CyOG; +*	
STIM^{KO}; *UASGCaMP6m/CyOG; +*	
STIM^{KO}; *UASmCD8GFP/CyOG; +*	
STIM^{KO}; *UASdsilp3/CyOG; +*	
STIM^{KO}; *UASMNScGAL4/CyOG; +*	
UAS-STIM+/CyOG; *UASGRAB$_{DA}$/Tb*	
UAS-STIM+/CyOG; *UASERGCaMP-210/Tb*	

Acknowledgements

This study was supported by core funds from NCBS (TIFR), a SERB

distinguished fellowship to GH (SB/S9/YSCP/SERB-DF/2018-1), a DBT

grant (BT/PR28450/MED/122/166/2018) to GH and a DST-WOS-A grant

(SR/WOS-A/LS-703/2016) to NK. We thank Fly base for access to various

databases, Bloomington and VDRC fly stock centres, Fly Facility at NCBS

for fly lines and NCBS Central Imaging and Flow Facility (CIFF) for confocal

imaging and NCBS lab support for timely help.

References:

1. Tennessen JM, Thummel CS. Coordinating growth and maturation - Insights from drosophila. Curr Biol. 2011;21(18):R750–7. PMID: 21959165

2. Delanoue R, Romero NM. Growth and Maturation in Development: A Fly's Perspective. Int J Mol Sci. 2020 Feb 13;21(4):1260. PMID: 32070061

3. Sewell D, Burnet B, Connolly K. Genetic analysis of larval feeding behaviour in Drosophila melanogaster. Genet Res. 1974 Oct;24(2):163–73. PMID: 4217753

4. Homem CCF, Knoblich JA. *Drosophila* neuroblasts: a model for stem cell biology. Development. 2012 Dec 1;139(23):4297–310. PMID: 23132240

5. Miroschnikow A, Schlegel P, Pankratz MJ. Making Feeding Decisions in the Drosophila Nervous System. Vol. 30, Curr Biol. 2020 Jul 20; 30(14):R831-R840. PMID: 32693083

6. Pu Y, Zhang Y, Zhang Y, Shen P. Two Drosophila Neuropeptide Y-like Neurons Define a Reward Module for Transforming Appetitive Odor Representations to Motivation. Sci Rep. 2018 Aug 3;8(1):11658. PMID: 30076343

7. Wang Y, Pu Y, Shen P. Neuropeptide-Gated Perception of Appetitive Olfactory Inputs in Drosophila Larvae. Cell Rep. 2013 Mar 30;3(3):820–30. PMID: 23453968

8. Lee KS, You KH, Choo JK, Han YM, Yu K. Drosophila short neuropeptide F regulates food intake and body size. J Biol Chem. 2004 Dec 3;279(49):50781–9. PMID: 15385546

9.   Youn H, Kirkhart C, Chia J, Scott K. A subset of octopaminergic

neurons that promotes feeding initiation in Drosophila melanogaster.

PLoS One. 2018 Jun 27;13(6):e0198362. PMID: 29949586

10.   Hückesfeld S, Peters M, Pankratz MJ. Central relay of bitter taste to

the protocerebrum by peptidergic interneurons in the Drosophila

brain. Nat Commun. 2016 Sep 13;7(1):12796. PMID: 27619503

11.   Bjordal M, Arquier N, Kniazeff J, Pin JP, Léopold P. Sensing of Amino

Acids in a Dopaminergic Circuitry Promotes Rejection of an

Incomplete Diet in Drosophila. Cell. 2014 Jan 30;156(3):510–21.

PMID: 24485457

12.   Schoofs A, Hückesfeld S, Pankratz MJ. Serotonergic network in the

subesophageal zone modulates the motor pattern for food intake in

Drosophila. J Insect Physiol. 2018;106:36–46. PMID: 28735009

13.   Erion R, Sehgal A. Regulation of insect behavior via the insulin-

signaling pathway. Front Physiol. 2013 Dec 3;4:353.

PMID: 24348428

14.   Schlegel P, Texada MJ, Miroschnikow A, Schoofs A, Hückesfeld S,

Peters M, et al. Synaptic transmission parallels neuromodulation in a

central food-intake circuit. Elife. 2016 Nov 15;5:e16799.

PMID: 27845623

15.   Chatterjee N, Perrimon N. What fuels the fly: Energy metabolism in

Drosophila and its application to the study of obesity and diabetes.

Sci Adv. 2021 Jun 9;7(24):4336–45. PMID: 34108216

16.   Koyama T, Rodrigues MA, Athanasiadis A, Shingleton AW, Mirth CK.

Nutritional control of body size through FoxO-Ultraspiracle mediated

ecdysone biosynthesis. Elife. 2014 Nov 25;3:e03091.

PMID: 25421296

17. Kannangara JR, Mirth CK, Warr CG. Regulation of ecdysone

production in *Drosophila* by neuropeptides and peptide hormones.

Open Biol. 2021 Feb 17;11(2):200373. PMID: 33593157

18. Lin S, Senapati B, Tsao CH. Neural basis of hunger-driven behaviour

in Drosophila. Open Biol. 2019 Mar 29;9(3):180259. PMID: 30914005

19. Almeida de Carvalho MJ, Mirth CK. Food intake and food choice are

altered by the developmental transition at critical weight in Drosophila

melanogaster. Anim Behav. 2017 Apr 1;126:195–208.

20. Wegman LJ, Ainsley JA, Johnson WA. Developmental timing of a

sensory-mediated larval surfacing behavior correlates with cessation

of feeding and determination of final adult size. Dev Biol. 2010

Sep;345(2):170–9. PMID: 206304

21. Zhang SL, Yu Y, Roos J, Kozak JA, Deerinck TJ, Ellisman MH, et al.

STIM1 is a Ca2+ sensor that activates CRAC channels and migrates

from the Ca2+ store to the plasma membrane. Nature 2005 Oct

6;437(7060):902–5. PMID: 16208375

22. Pathak T, Trivedi D, Hasan G. CRISPR-Cas-Induced Mutants Identify

a Requirement for dSTIM in Larval Dopaminergic Cells of *Drosophila*

melanogaster. G3 (Bethesda). 2017 Mar 10;7(3):923–33.

PMID: 28131984

23. Fox DT, Duronio RJ. Endoreplication and polyploidy: insights into

development and disease. Development. 2013 Jan 1;140(1):3–12.

PMID: 23222436

24. Edgar BA, Orr-Weaver TL. Endoreplication Cell Cycles: more or less.

Cell. 2001 May;105(3):297–306. PMID: 11348589

25. Doe CQ. Neural stem cells: balancing self-renewal with

differentiation. Development. 2008 May;135(9):1575–87.

PMID: 18356248

26. Britton JS, Edgar BA. Environmental control of the cell cycle in
Drosophila: Nutrition activates mitotic and endoreplicative cells by
distinct mechanisms. Development. 1998 Jun;125(11):2149–58.
PMID: 9570778

27. Lacin H, Truman JW. Lineage mapping identifies molecular and
architectural similarities between the larval and adult Drosophila
central nervous system. Elife. 2016 Mar 15;5:e13399.
PMID: 26975248

28. Zielke N, Korzelius J, Straaten M Van, Bender K, Schuhknecht GFP,
Dutta D, et al. Resource Fly-FUCCI□: A Versatile Tool for Studying
Cell Proliferation in Complex Tissues. Cell Reports. 2014 Apr
24;7(2):588–598. PMID: 24726363

29. Sousa-Nunes R, Yee LL, Gould AP. Fat cells reactivate quiescent
neuroblasts via TOR and glial insulin relays in Drosophila. Nature.
2011 Mar 24;471(7339):508–12. PMID: 21346761

30. Chell JM, Brand AH. Nutrition-Responsive Glia Control Exit of Neural
Stem Cells from Quiescence. Cell. 2010 Dec 23;143(7):1161–73.
PMID: 21183078

31. Shen P. Analysis of Feeding Behavior of *Drosophila* Larvae on Liquid
Food. Cold Spring Harb Protoc. 2012 May 1;2012(5):pdb.prot069310.
PMID: 22550301

32. Friggi-Grelin F, Coulom H, Meller M, Gomez D, Hirsh J, Birman S.
Targeted gene expression in Drosophila dopaminergic cells using
regulatory sequences from tyrosine hydroxylase. J Neurobiol.
2003;54(4):618–27. PMID: 12555273

33. Xie T, Ho MCW, Liu Q, Horiuchi W, Lin CC, Task D, et al. A Genetic
Toolkit for Dissecting Dopamine Circuit Function in Drosophila. Cell
Rep. 2018 Apr 10;23(2):652–665. PMID: 29642019

34.   Hartenstein V, Cruz L, Lovick JK, Guo M. Developmental analysis of

the dopamine-containing neurons of the Drosophila brain. J Comp

Neurol. 2017 Feb 1;525(2):363–379. PMID: 27350102

35.   Selcho M, Pauls D, Han KA, Stocker RF, Thum AS. The Role of

Dopamine in Drosophila Larval Classical Olfactory Conditioning.

Giurfa M, editor. PLoS One. 2009 Jun 12;4(6):e5897.

PMID: 19521527

36.   Sitaraman D, Zars M, LaFerriere H, Chen YC, Sable-Smith A,

Kitamoto T, et al. Serotonin is necessary for place memory in

Drosophila. Proc Natl Acad Sci U S A. 2008 Apr 8;105(14):5579–84.

PMID: 18385379

37.   Venkiteswaran G, Hasan G. Intracellular Ca2+ signaling and store-

operated Ca2+ entry are required in Drosophila neurons for flight.

Proc Natl Acad Sci U S A. 2009;106(25):10326–31. PMID: 19515818

38.   Nitabach MN, Blau J, Holmes TC. Electrical Silencing of Drosophila

Pacemaker Neurons Stops the Free-Running Circadian Clock. Cell.

2002 May 17;109(4):485–95. PMID: 12086605

39.   Nitabach MN, Wu Y, Sheeba V, Lemon WC, Strumbos J, Zelensky

PK, et al. Electrical hyperexcitation of lateral ventral pacemaker

neurons desynchronizes downstream circadian oscillators in the fly

circadian circuit and induces multiple behavioral periods. J Neurosci.

2006 Jan 11;26(2):479–89. PMID: 16407545

40.   Nicolaï LJJ, Ramaekers A, Raemaekers T, Drozdzecki A, Mauss AS,

Yan J, et al. Genetically encoded dendritic marker sheds light on

neuronal connectivity in Drosophila. Proc Natl Acad Sci U S A. 2010

Nov 3;107(47):20553–8. PMID: 21059961

41.   Sun F, Zeng J, Jing M, Zhou J, Feng J, Owen SF, et al. A Genetically

Encoded Fluorescent Sensor Enables Rapid and Specific Detection

of Dopamine in Flies, Fish, and Mice. Cell. 2018 Jul 12;174(2):481-
496.e19. PMID: 30007419

42. Luo D, Broad LM, Bird GS, Putney JW. Signaling Pathways
Underlying Muscarinic Receptor-induced [Ca 2+]i Oscillations in
HEK293 Cells. J Biol Chem. 2001 Feb 23;276(8):5613–21.
PMID: 11096083

43. Sharma A, Hasan G. Modulation of flight and fee ding behaviours
requires presynaptic IP3Rs in dopaminergic neurons. Elife. 2020 Nov
6;9:e622967. PMID: 33155978

44. Oliva MK, Pérez-Moreno JJ, O'Shaughnessy J, Wardill TJ, O'Kane
CJ. Endoplasmic Reticulum Lumenal Indicators in Drosophila Reveal
Effects of HSP-Related Mutations on Endoplasmic Reticulum Calcium
Dynamics. Front Neurosci. 2020 Aug 10;14:816. PMID: 32903680

45. Wu Q, Wen T, Lee G, Park JH, Cai HN, Shen P. Developmental
Control of Foraging and Social Behavior by the Drosophila
Neuropeptide Y-like System. Neuron. 2003 Jul 3;39(1):147–61.
PMID: 12848939

46. Klapoetke NC, Murata Y, Kim SS, Pulver SR, Birdsey-Benson A, Cho
YK, et al. Independent optical excitation of distinct neural populations.
Nat Methods. 2014 Mar;11(3):338–46. PMID: 24509633

47. Fenno L, Yizhar O, Deisseroth K. The development and application of
optogenetics. Annu Rev Neurosci. 2011;34(1):389–412.
PMID: 21692661

48. Nässel DR, Zandawala M. Recent advances in neuropeptide
signaling in Drosophila, from genes to physiology and behavior. Vol.
179, Prog Neurobiol. 2019 Aug;179:101607. PMID: 30905728

49. Hu Y, Comjean A, Perrimon N, Mohr SE. The Drosophila Gene

Expression Tool (DGET) for expression analyses. BMC

Bioinformatics. 2017 Feb 10;18(1):98. PMID: 28187709

50. Eschbach C, Fushiki A, Winding M, Schneider-Mizell CM, Shao M,

Arruda R, et al. Recurrent architecture for adaptive regulation of

learning in the insect brain. Nat Neurosci. 2020 Apr 23;23(4):544–55.

PMID: 32203499

51. Saumweber T, Rohwedder A, Schleyer M, Eichler K, Chen YC, Aso

Y, et al. Functional architecture of reward learning in mushroom body

extrinsic neurons of larval Drosophila. Nat Commun. 2018 Dec

1;9(1):1104. PMID: 29549237

52. Hückesfeld S, Schlegel P, Miroschnikow A, Schoofs A, Zinke I,

Haubrich AN, et al. Unveiling the sensory and interneuronal pathways

of the neuroendocrine connectome in Drosophila. Elife. 2021 Jun

4;10;e65745. PMID: 34085637

53. Leopold A V., Shcherbakova DM, Verkhusha V V. Fluorescent

Biosensors for Neurotransmission and Neuromodulation: Engineering

and Applications. Front Cell Neurosci. 2019 Oct 23;13:474.

PMID: 31708747

54. Andretic R, van Swinderen B, Greenspan RJ. Dopaminergic

Modulation of Arousal in Drosophila. Curr Biol. 2005 Jul

12;15(13):1165–75. PMID: 16005288

55. Wiencke K, Horstmann A, Mathar D, Villringer A, Neumann J.

Dopamine release, diffusion and uptake: A computational model for

synaptic and volume transmission. PLoS Comput Biol. 2020 Nov

30;16(11):e1008410. PMID: 33253315

56. Birman S. Arousal Mechanisms: Speedy Flies Don't Sleep at Night.

Curr Biol. 2005 Jul 12;15(13):R511–3. PMID: 16005286

57. Ganetzky B, Wu C F. Drosophila Mutants With Opposing Effects on

Nerve Excitability: Genetic and Spatial Interactions in Repetitive

Firing. J Physiol. 1982 Mar;47(3):501-14. PMID: 6279790

58. Colombani J, Raisin S, Pantalacci S, Radimerski T, Montagne J,

Léopold P. A Nutrient Sensor Mechanism Controls Drosophila

Growth. Cell. 2003 Sep 19;114(6):739–49. PMID: 14505573

59. Mitra R, Hasan G. Store-operated Ca2+ entry regulates neuronal

gene expression and function. Curr Opin Neurobiol. 2022 Apr

1;73:102520. PMID: 35220059

60. Dhanya SK, Hasan G. Purkinje Neurons with Loss of STIM1 Exhibit

Age-Dependent Changes in Gene Expression and Synaptic

Components. J Neurosci. 2021 Apr 28;41(17):3777–98.

PMID: 33737457

61. Richhariya S, Jayakumar S, Abruzzi K, Rosbash M, Hasan G. A

pupal transcriptomic screen identifies Ral as a target of store-

operated calcium entry in Drosophila neurons. Sci Rep. 2017 Feb

14;7:42586. PMID: 28195208

62. Tsao CH, Chen CC, Lin CH, Yang HY, Lin S. Drosophila mushroom

bodies integrate hunger and satiety signals to control innate food-

seeking behavior. Elife. 2018 Mar 16;7:e35264. PMID: 29547121

63. Wise RA. Role of brain dopamine in food reward and reinforcement.

Philos Trans R Soc Lond B Biol Sci. 2006 Jul 29;361(1471):1149–58.

PMID: 16874930

64. Szczypka MS, Rainey MA, Kim DS, Alaynick WA, Marck BT,

Matsumoto AM, et al. Feeding behavior in dopamine-deficient mice.

Proc Natl Acad Sci U S A. 1999 Oct 12;96(21):12138–43.

PMID: 10518589

65.   Boekhoudt L, Roelofs TJM, De Jong JW, De Leeuw AE, Luijendijk

MCM, Wolterink-Donselaar IG, et al. Does activation of midbrain

dopamine neurons promote or reduce feeding? Int J Obes (Lond).

2017 Jul;41(7):1131–40. PMID: 28321131

66.   Karam CS, Jones SK, Javitch JA. Come Fly with Me: An overview of

dopamine receptors in *Drosophila melanogaster*. Basic Clin

Pharmacol Toxicol. 2020 Jun 9;126(S6):56–65. PMID: 31219669

67.   Sareen PF, McCurdy LY, Nitabach MN. A neuronal ensemble

encoding adaptive choice during sensory conflict in Drosophila. Nat

Commun. 2021 Jul 5;12(1):4131. PMID: 34226544

68.   Kleinridders A, Pothos EN. Impact of Brain Insulin Signaling on

Dopamine Function, Food Intake, Reward, and Emotional Behavior.

Curr Nutr Rep. 2019 Jun 18;8(2):83–91. PMID: 31001792

69.   Fiory F, Perruolo G, Cimmino I, Cabaro S, Pignalosa FC, Miele C, et

al. The Relevance of Insulin Action in the Dopaminergic System.

Front Neurosci. 2019 Aug 16;13:868. PMID: 31474827

70.   Tavares G, Martins FatimaO, Melo BernardeteF, Matafome P, Conde

Silvia v. Peripheral Dopamine Directly Acts on Insulin-Sensitive

Tissues to Regulate Insulin Signaling and Metabolic Function. Front

Pharmacol. 2021 Sep 9;12:713418.

71.   Stouffer MA, Woods CA, Patel JC, Lee CR, Witkovsky P, Bao L, et al.

Insulin enhances striatal dopamine release by activating cholinergic

interneurons and thereby signals reward. Nat Commun. 2015 Oct

27;6(1):8543. PMID: 26503322

72.   Labouèbe G, Liu S, Dias C, Zou H, Wong JCY, Karunakaran S, et al.

Insulin induces long-term depression of ventral tegmental area

dopamine neurons via endocannabinoids. Nat Neurosci. 2013 Mar

27;16(3):300–8. PMID: 23354329

73. Li Y, Ren L, Fu H, Yang B, Tian J, Li Q, et al. Crosstalk between dopamine and insulin signaling in growth control of the oyster. Gen Comp Endocrinol. 2021 Nov 1;313:113895. PMID: 34480943

74. Chowański S, Walkowiak-Nowicka K, Winkiel M, Marciniak P, Urbański A, Pacholska-Bogalska J. Insulin-Like Peptides and Cross-Talk With Other Factors in the Regulation of Insect Metabolism. Front Physiol. 2021 Jun 29;12:701203. PMID: 34267679

75. Nash AI. Crosstalk between insulin and dopamine signaling: A basis for the metabolic effects of antipsychotic drugs. J Chem Neuroanat. 2017 Oct 1;83–84:59–68. PMID: 27480675

76. Volkow ND, Wang GJ, Baler RD. Reward, dopamine and the control of food intake: Implications for obesity. Trends Cogn Sci. 2011 Jan;15(1):37–46. PMID: 21109477

77. Wang GJ, Geliebter A, Volkow ND, Telang FW, Logan J, Jayne MC, et al. Enhanced striatal dopamine release during food stimulation in binge eating disorder. Obesity (Silver Spring). 2011 Aug;19(8):1601–8. PMID: 21350434

78. Joshi R, Venkatesh K, Srinivas R, Nair S, Hasan G. Genetic Dissection of *itpr* Gene Function Reveals a Vital Requirement in Aminergic Cells of Drosophila Larvae. Genetics. 2004 Jan 1;166(1):225–36. PMID: 15020420

79. Ashburner, M. (1989) Drosophila. A Laboratory Hand- book. Cold Spring Harbor Laboratory Press, Cold Spring Harbor.

80. Delanoue R, Slaidina M, Léopold P. The Steroid Hormone Ecdysone Controls Systemic Growth by Repressing dMyc Function in Drosophila Fat Cells. Dev Cell. 2010 Jun 15;18(6):1012–21. PMID: 20627082

81.   Pathak T, Agrawal T, Richhariya S, Sadaf X, Hasan XG. Store-

Operated Calcium Entry through Orai Is Required for Transcriptional

Maturation of the Flight Circuit in Drosophila. J. Neurosci.

2015;35(40):13784–99. PMID: 26446229

82.   Agrawal N, Venkiteswaran G, Sadaf S, Padmanabhan N, Banerjee S,

Hasan G. Inositol 1,4,5-trisphosphate receptor and dSTIM function in

Drosophila insulin-producing neurons regulates systemic intracellular

calcium homeostasis and flight. J Neurosci. 2010 Jan 27;30(4):1301–

13. PMID: 20107057

1089

Figure legends:

Fig 1. $STIM^{KO}$ larvae exhibit reduced feeding followed by systemic growth arrest.

(A-C) Number of 1st instar (L1), 2nd instar (L2) and 3rd instar (L3) larvae from *CS* (grey) and $STIM^{KO}$ (magenta) measured at 6h intervals after egg laying (AEL) at the specified time points (mean ± SEM). Number of sets (N) = 3, number of larvae per replicate (n) = 10. *P < 0.05, Student's *t*-test with unequal variances. P values are given in **S2 Table.**

(D) Representative images of larvae from *CS* and $STIM^{KO}$ at the indicated time. Scale bar = 1 mm

(E) Measurement of larval length (mean ± SEM) from *CS* (grey) and $STIM^{KO}$ (magenta) larvae at the specified time points. Number of larvae per genotype per time point is (n) ≥ 12. *P < 0.05 for all time points, Student's *t*-test with unequal variances. P values are given in **S2 Table.**

(F) Representative images of thoracic neuroblasts marked with *Insc>mCD8GFP* (green), a neuroblast marker (anti-Deadpan, red) and a marker for post-mitotic cells (anti-Prospero, blue) from control

(*Insc>mCD8GFP*) and *STIM^{KO}; Insc>mCD8GFP* animals at the indicated

ages. Similar images were obtained from four or more animals. Scale bar =

20µm.

(G) Diagrammatic summary of neuroblast proliferation in control

(*Insc>mCD8GFP*) and *STIM^{KO}; Insc>mCD8GFP* animals.

(H) Stack bar graph showing number of dividing neuroblasts to non-dividing

neuroblasts. *P < 0.05, Student's *t*-test with unequal variances, n = 4

animals from each genotype. P values are given in **S2 Table.**

(I) Representative images of dye-fed larvae from *CS* and *STIM^{KO}* at the

indicated times AEL, scale bar = 200µm except for *CS* (80-84h) where

scale = 1mm.

(J) Quantification (mean ± SEM) of ingested blue dye in *CS* and *STIM^{KO}*

larvae at the indicated ages by normalizing optical density (OD) of the dye

at 655nm to concentration of protein. Number of feeding plates per time

point (N) = 6, number of larvae per plate (n) =10. *P < 0.05, Student's *t*-test

with unequal variances. P values are given in **S2 Table.**

(K) Line graph with quantification of larval mouth hook contractions per 30

seconds (mean ± SEM) from *CS* and *STIM^{KO}* at indicated developmental

time points. Number of larvae per genotype per time point is (n) ≥ 10. *P <

0.05 at all time points, Student's *t*-test with unequal variances. All P values

are given in **S2 Table.**

1128

Fig 2. Feeding and growth deficits of *STIM^{KO}* larvae arise from central

dopaminergic neurons

(A) Number (mean ± SEM) of 3^{rd} instar larvae (L3) are restored close to

wildtype (CS) levels by expression of *STIM^{+}* in *THD'* cells of *STIM^{KO}* larvae

(rescue). Larvae were monitored at 6h intervals from 66h to 96h AEL.

Number of sets (N) = 3, number of larvae per set (n) = 10. Letters represent

statistically similar groups for the 90h and 96h time point. Also see **S2A Fig**

for a complete developmental profile of CS, $STIM^{KO}$ and rescue $(STIM^{KO}$;

THD'>STIM⁺) animals with appropriate genetic controls.

(B) Stack bars with the number of adults (mean ± SEM) that eclosed at 320

to 326h AEL from the indicated genotypes. The genotype of rescue larvae is

$STIM^{KO}$*; THD'>STIM*⁺. Number of sets (N) = 3, number of organisms per set

(n) = 25.

(C) Representative confocal images of the larval brain from animals of the

genotype *THD'>mCD8GFP*. Anti-GFP (green) indicates the expression of

THD'GAL4 and anti-TH (magenta) marks all dopaminergic cells. Asterisks

mark TH⁺ᵛᵉ cells in CNS whereas arrowheads mark non-TH positive cells in

ventral ganglia of larval brain (i and ii). DL1 and DL2 clusters in the central

brain of three and two dopaminergic cells respectively are marked (iii).

Scale bars = 20µm.

(D) Representative confocal images of the larval brain from animals of the

genotype *THGAL80,THD'>mCD8GFP*. *THD'GAL4* driven GFP expression

(green) is suppressed in DL1 and DL2 clusters in the CNS (asterisk) by

THGAL80 but not in the ventral ganglia (arrowheads). Scale bar = 20µm.

(E) Line graph shows the number (mean ± SEM) of 3ʳᵈ instar larvae from

CS, $STIM^{KO}$, $STIM^{KO}$;THD'>STIM⁺ and $STIM^{KO}$;THGAL80,THD'>STIM⁺ at 6h

intervals between 60 to 96h AEL. Number of sets (N) = 3, number of larvae

per set (n) = 10. Different alphabet represent statistically significant groups

for 84h, 90h and 96h. Also see **S2A Fig** for complete larval staging profile

and additional genetic controls.

(F) Stack bar graph showing the number of adults eclosed (mean ± SEM) at

320 to 326h AEL from *CS (*wildtype*), $STIM^{KO}$* (mutant), $STIM^{KO}$*;THD'>STIM*⁺

(rescue) and $STIM^{KO}$*;THGAL80,THD'>STIM*⁺ (THGAL80 - restrictive rescue)

genotypes. Different alphabet represent statistically significant groups.

Number of sets (N) = 3, number of organisms per set (n) = 25.

(G) Representative images of dye-fed larvae of *CS, STIM^{KO}* and rescue

(*STIM^{KO}; THD'>STIM$^+$*) genotypes at 80-84h AEL Scale bar = 1mm.

(H) Bar graph with quantification of ingested food containing a blue dye in

larvae of the indicated genotypes (*CS, STIM^{KO}* and *STIM^{KO}; THD'>STIM$^+$*

rescue) at the indicated developmental times. Mean (mean ± SEM) optical

density (655nM) of blue dye in larval lysates after normalizing to larval

protein concentration (OD/Protein concentration X 10^{-2}) was obtained from 6

feeding plates (N) each containing 10 larvae (n). Different alphabet

represent statistically significant groups.

(I) Expression of *STIM$^+$* in THD' cells rescues the feeding behaviour deficit

of *STIM^{KO}* larvae. Box graph with quantification of larval mouth hook

contractions of the indicated genotypes (*CS, STIM^{KO}* and *STIM^{KO};*

THD'>STIM$^+$ rescue). Circles represent single larvae in the box graph of 25th

and 75th percentiles with the median (bar), and mean (square). Number of

larvae per genotype per time point is (n) ≥10. Different alphabet represent

statistically significant groups.

(J) Number of 3r^{d} instar larvae (mean ± SEM) from RNAi knockdown of

STIM$^+$ in THD' neurons (*THD'>dsSTIM*) along with control genotypes

THD'/+ and *dsSTIM/+* at 6h intervals between 66h to 96h AEL (After Egg

laying). Number of sets (N) = 3, number of larvae per set (n) = 10. Also see

S2B and **S2C Figs** for complete larval staging profile and adult weights.

Different alphabet represent statistically significant groups.

(K) Representative images of dye-fed larvae of the indicated genotypes at

58-62h and 82-86h AEL. Scale bar = 200µm.

(L) Quantification (mean ± SEM) of blue dye containing ingested food

(OD/Protein concentration X 10^{-2}; similar to panel D above) in larvae of the

indicated genotypes at 58-62h and 82-86h AEL. No. of plates for each time

point, (N) = 3, number of larvae per plate (n) = 10.

In all graphs and box plots, different alphabets represent distinct

statistical groups as calculated by one way ANOVA followed by post-hoc

Tukey's test. P values for individual panels are given in **S2 Table.**

1195

Fig 3. **Dopamine synthesis and excitability in THD' neurons is**

important for larval growth.

(A) Larval mouth hook movements that correlate with feeding are reduced in

larvae with reduced dopamine synthesis in THD' neurons. Box graph with

quantification of mouth hook contractions in larvae with knockdown of

Tyrosine Hydroxylase (*dsTH*) in THD' neurons and appropriate control

genotypes. Number of larvae per genotype per time point is (n) ≥10.

(B) Representative images of larvae with knockdown of Tyrosine

Hydroxylase (*THD'>dsTH*) and controls (*THD'/+, dsTH/+*) at 82-86h. Scale

bar = 1mm.

(C) Quantification of larval length from the indicated genotypes. n≥15.

(D) Quantification of weight of 10 flies from the indicated genotypes.. Each

circle represents one set of adult flies consisting of 5 females and 5 males,

6-8h post-eclosion. A minimum of 5 sets were measured for each genotype.

(E) Representative images of the central brain (left panels) indicating the

region of focus (boxed), followed by images of DL1 and DL2 clusters of

THD' cells from two lobes of same brain with Ca^{2+} transients at the

indicated time points before and after addition of a depolarizing agent (KCl –

70mM). Ca^{2+} transients were measured by changes in the intensity of

GCaMP6m fluorescence from *THD'>GCaMP6m* (control), *STIM^{KO};*

THD'>GCaMP6m, *STIM^{KO};THD'>GCaMP6m, STIM^{+}* (rescue). Scale bar

20µm

(F) Changes in GCaMP6m fluorescence (mean ± SEM of $\Delta F/F$) from THD'

neurons of the indicated genotypes. Number of brains, (N) $\geq$ 5, number of

cells, (n) $\geq$ 15.

(G) Peak intensities of GCaMP6m fluorescence (ΔF) in THD' cells from the

indicated genotypes. Box plots show 25^{th} and 75^{th} percentiles, the median

(bar), and mean (square) of ΔF of each cell (small circles).

(H) Number of 3^{rd} instar larvae of *CS, STIM^{KO}* and *STIM^{KO};THD'>NaChBac*

genotypes at the indicated times AEL (mean ± SEM). Number of sets (N) =

3, number of larvae per set (n) = 15.

(I) Representative images of larvae from *CS, STIM^{KO}* and *STIM^{KO};*

THD'>NaChBac at 94-98h AEL, scale bar = 1mm.

(J) Quantification of larval length from the indicated genotypes. n=10.

(K) Number of adults eclosed at 320h AEL (mean ± SEM) from wildtype

(*CS*), mutant (*STIM^{KO}*) and NaChBac rescue (*STIM^{KO}; THD'>NaChBac*)

animals. Numbers were obtained from three experiments (N = 3), number of

organisms per experiment, n = 15.

In all box plots, circles represent single larvae or flies. The box plots

span 25^{th} and 75^{th} percentiles with the median (bar), and mean (square).

Alphabets represent distinct statistical groups as calculated by one way

ANOVA followed by post-hoc Tukey's test. P values are given in **S2 Table.**

1238

Fig 4. STIM is necessary for dopamine release and ER-Ca^{2+}

homeostasis

(A) Axonal and dendritic projections of THD' neurons visualized by

expression of SyteGFP (green) and Denmark (anti-RFP, magenta)

respectively in a representative brain immunostained for anti-Brp (blue).

Panels marked as (d) contain magnified images from (c) of presynaptic

terminals (green) located at the base of the mushroom body in the CNS

(asterisk), as branches extruding into basomedial protocerebrum of the CNS

(arrowhead) and as punctae near the oesophageal region (hash).

(B) Representative images of dopamine release before and after addition of

Carbachol (CCh) as measured by changes in intensity of $GRAB_{DA}$ at the

presynaptic terminals of THD' neurons of control (*THD'>GRAB_{DA}*), *STIM^{KO}*

(*STIM^{KO};THD'>GRAB_{DA}*) and rescue (*STIM^{KO};THD'>GRAB_{DA}, STIM^+)*

genotypes. Scale bar = 20μm

(C) Normalized changes in fluorescence ($\Delta F/F$) of $GRAB_{DA}$ measuring

dopamine release from THD' neurons of control (*THD'>GRAB_{DA}*), *STIM^{KO}*

(*STIM^{KO};THD'>GRAB_{DA}*) and rescue (*STIM^{KO};THD'>GRAB_{DA}, STIM^+)*

genotypes. Traces show the average change in $GRAB_{DA}$ florescence (mean

± SEM of ($\Delta F/F$)) measured from individual presynaptic regions of interest

($\geq$10) taken from N $\geq$ 6 brains of each genotype.

(D) Representative time series images of ER calcium transients as

measured by changes in intensity of ER-GCaMP in THD' neurons of control

(*THD'>ER-GCaMP-210*), *STIM^{KO}* (*STIM^{KO};THD'> ER-GCaMP-210)* and

rescue (*STIM^{KO};THD'> ER-GCaMP-210, STIM^+*) genotypes.

(E) i) Traces of normalized ER-GCaMP ($\Delta F/F$) responses (mean ± SEM)

from THD' neurons of the indicated genotypes. Control (*THD'>ER-GCaMP-*

210) Type 1, indicates cells that exhibit ER-Ca^{2+} release upon CCh addition

followed by refilling of ER-stores. Control Type 2 indicates cells where ER-

Ca^{2+} release was not evident. Rescue indicates *STIM^{KO}; THD'>ERGCaMP-*

210,STIM$^+$. For each genotype, number of cells (n) $\geq$ 15 and N $\geq$ 5 brains.

(ii) Quantification of control cells that exhibit ER Ca^{2+} release (Type 1) and

cells where ER Ca^{2+} release was not observed (Type 2). (iii) Boxed region

from (i) enlarged to show ER Ca^{2+} response of Type 1 and Type 2 control

cells.

1273

Fig 5. Differential regulation of neuropeptidergic cell activity by THD'

neurons

(A)   Pictorial representation of an experimental setup for testing

functional connectivity of dopaminergic (THD') and peptidergic (C929)

cells   in   organisms   of   the   genotype   *THD'>Chrimson,*

C929LexA>LexAopGCaMP6m. The lower panel shows the peptidergic

neurons containing regions 1-3 in which change in Ca^{2+} transients were

observed upon activation of THD' cells.

(B)   Ca^{2+} transients in peptidergic Median Neurosecretory cells (MNSc,

top), lateral neurosecretory cells (LNSc, middle) and suboesophageal

region (SEZ, bottom) as measured by the intensity of GCaMP6m before

Chrimson activation (100s) and after Chrimson activation (300s and 400s)

of THD' cells. Activated cells (red asterisks) and inhibited cells (yellow

asterisks) are marked. Scale bar = 20um.

(C)   Quantification of total C929 marked cells from 9 brains that were

either optogenetically activated or inhibited.

(D)   Average GCaMP6m responses (mean ± SEM of ΔF/F) across time

in C929 marked cells that were activated upon Chrimson activation of

THD' neurons at 633nm. The box plot shows 25th and 75th percentiles of

peak responses measured by change in GCaMP6m of individual activated

peptidergic cells (circles), with the median (bar), and mean (square). n =

24 cells from (N) = 9 brains

(E)   Average GCaMP6m responses (mean ± SEM of ∆F/F) across time

in C929 marked cells that were inhibited upon Chrimson activation of THD'

neurons at 633nm. The box plot shows 25^{th} and 75^{th} percentiles of peak

responses measured by change in GCaMP6m of individual inhibited

peptidergic cells (circles), with the median (bar), and mean (square). n =

40 cells from (N) = 9

(F)   (i) Pulsed activation of THD' neurons produce corresponding pulses

of activation in certain peptidergic neurons as evident from their GCaMP6m

responses (mean ± SEM). Red panels indicate the time intervals (50s each)

of *THD'>Chrimson* activation. n = 21 cells from N = 11 brains.

(ii) Pulsed activation of THD' neurons produce corresponding pulses of

inhibition in certain peptidergic neurons as evident from their GCaMP6m

responses (mean ± SEM). Red panels indicate the time intervals (50s

each) of *THD'>Chrimson* activation. n = 19 cells from N = 9 brains.

(G)   GCaMP6m responses (mean ± SEM) are absent in C929 cells in

the absence of red light and in larvae that were not fed with All-trans

Retinol (ATR). n = 50 cells from N=9 brains.

1313

Fig 6. Developmental expression of insulin-like peptide 3 (ilp3) is

regulated by STIM in THD' neurons

(A)   FPKM values obtained for transcripts encoding the indicated insulin

like peptides (*ilp*) obtained from an RNAseq experiment performed from

CS brains (58-62h, mid 2^{nd} instar stage) and *STIM*^{KO} brains (72-76h, mid

2^{nd} instar stage).

(B)   Quantification of *ilp2, ilp3* and *ilp5* transcripts in *CS*, *STIM*^{KO} and

STIM^{KO}; *THD'>STIM*$^{+}$ (rescue) brains. Expression changes of *ilp3* and *ilp5*

transcripts in $STIM^{KO}$ brains were rescued upon over-expression of $STIM^+$

in THD' neurons. Transcripts were measured relative to controls. Data

show (mean ± SEM) of fold changes of gene expression (N=3).

(C)   Heatmap of *ilp3* expression from larval to pupal stages. FPKM

values   were   obtained   from   the   DGET   database

(https://www.flyrnai.org/tools/dget/web/).

(D)   Knockdown of *ilp3* partially rescues viability in $STIM^{KO}$ larvae. Stack

bar graph showing the number of viable organisms and their developmental

stage at the specified hours after egg laying for *CS, $STIM^{KO}$; MNScGAL4/+,*

$STIM^{KO}$;   *dsilp3/+*   (control)   genotypes   followed   by

$STIM^{KO}$;MNScGAL4>dsilp3. Data are from three experiments, each with 25

organisms per genotype.

(E)   Representative images of larvae from *CS* and *$STIM^{KO}$; MNSc>dsilp3*

at 118-122h AEL, scale bar = 1mm.

(F)   Quantification of larval length from *CS* and *$STIM^{KO}$; MNSc>dsilp3*

(rescue, L3b) genotypes. Box plots show the 25th and 75th percentiles with

median (bar), mean (square) and sizes of individual larvae (circles), n=10.

*$P<0.05$ (unpaired t-test).

(G)   Representative image of an adult $STIM^{KO}$ mutant fly rescued by

knockdown of *ilp3* (*$STIM^{KO}$; MNScGAL4>dsilp3*). Scale bar = 1mm.

(H)   Number of 3rd instar larvae at the indicated times AEL (mean ±

SEM) from the indicated genotypes with knockdown of *ilp3* in the MNSc

(*MNSc>dsilp3*) and of controls (*MNSc/+ and dsilp3/+*). Alphabets indicate

statistically different groups at 132h and 144h. Number of sets (N) = 3 and

number of larvae per set (n) = 10.

(I)   Representative images of pupae with knock-down of *ilp3* (*MNSc>*

dsilp3) at 144-156h AEL and appropriate controls (*MNSc/+* and *dsilp3/+*).

Scale bar = 1mm.

(J) Quantification of pupal volume from the indicated genotypes. Box

plots show the 25^{th} and 75^{th} percentiles with median (bar), mean (square)

and sizes of individual pupa (circles), n=30

(K) Number of 3^{rd} instar larvae obtained by over-expression of *ilp3*

(*MNSc>ilp3⁺)* and in appropriate control genotypes (*MNSc/+, ilp3⁺/+*) at

the indicated time AEL (mean $\pm$ SEM). (N) = 3 and (n) = 10. Alphabets

indicate statistically different groups.

(L) Representative images of larvae with over-expression of *ilp3*

(*MNSc>ilp3⁺*) and appropriate controls (*MNSc/+* and *ilp3⁺/+*) at 94-98h

AEL, scale bar = 1mm.

(M) Overexpression of ilp3 results in smaller sized larvae.

Quantification of larval length from the indicated genotypes. Box plots

show the 25^{th} and 75^{th} percentiles with median (bar), mean (square) and

sizes of individual larvae (circles), n $\geq$24.

(N) Overexpression of ilp3 does not affect feeding. Box graph showing

25^{th} and 75^{th} percentiles, the median (bar), and mean (square) of number

of larval mouth hook contraction per 30 seconds, where each circle

represents a single larva of the indicated genotype. Number of larvae per

genotype is (n)$\geq$10.

For all graphs and box plots, alphabets represent distinct statistical

groups as calculated by one way ANOVA followed by post-hoc Tukey's

test, n.s. indicates non-significant groups. P values are given in **S2 Table.**

1372

Fig 7. MNSc respond differentially to Dopamine stimulation

(A) Development profiles of larvae with RNAi mediated knockdown of

dopamine receptors in the MNSc. Bar graphs represent number of viable

organisms and their developmental stages at the specified hours after egg

laying for the indicated genotypes (*CS, MNScGAL4/+, dsDop1R1/+,*

dsDopEcR/+,   dsDop2R1/+   (Controls),   *MNSc>dsDop1R1,*

MNSc>dsDopEcR, and *MNSc>dsDop2R2*). N=3 batches of n=25 larvae

for every genotype. P values of all genotypes compared with appropriate

controls were calculated by one way ANOVA followed by post-hoc Tukey's

test and are given in **S2 Table.**

(B) Representative images of larvae from specified genotypes at 128-

134hr. Scale bar = 1mm

(C) Quantification of weight of 10 flies from the indicated genotypes. Box

plot shows the 25^{th} and 75^{th} percentiles with median (bar) and mean

(square). Each circle represents one set of adult flies consisting of 5

females and 5 males, 6-8h post-eclosion. A minimum of 5 sets were

measured for each genotype. Alphabets represent distinct statistical

groups as calculated by one way ANOVA followed by post-hoc Tukey's

test. P values are given in **S2 Table.**

(D) Representative images of larval brains expressing the calcium indicator

GCaMP6 in MNSc. Brains were stimulated with either dopamine (DA) in

presence of tetrodotoxin (top panel) or a control solution (HL3; lower panel)

and imaged over time. Scale bar = 20μm

(E) Changes in GCaMP6m fluorescence (mean ± SEM of ΔF/F) from MNSc

cells with addition of either dopamine (DA) and Tetrodotoxin (TTX) (7 brains

and 12-14 cells) or HL3 (No DA+ No TTX) (5 brains and 14 cells).

1399

Supplementary figure legends:

Supplementary fig 1.

(A-B) Number of viable organisms (mean ± SEM) at the indicated

developmental stage of *CS* and *STIM^{KO}* after egg laying. Number of sets (N)

= 3, number of organisms per set (n) = 25.

(C) Confocal images of larval brains at the indicated time points expressing

the FUCCI marker. Control genotype is *Insc>FUCCI* (top row) and the

mutant genotype is *STIM^{KO}*; *Insc>FUCCI* (bottom row). Here late mitosis/G1

phase, S-phase and G2/early mitosis are marked by green, red, and yellow

fluorescent indicators respectively. Scale bar = 20mm; n = 5 larval brains.

(D) Representative image of dye-fed larvae from *CS* and *STIM^{KO}* at 80-

84h AEL. Scale bar: 2mm

(E) Bar graph showing the average number of Dye^{+ve} (presence of blue dye

in the gut) and Dye^{-ve} (absence of blue dye in the gut) *CS* and *STIM^{KO}*

larvae at the indicated ages (mean ± SEM). Number of feeding plates per

time point (N) = 3, number of larvae per plate (n) = 30. $^*P < 0.05$, Student's

t-test with unequal variances. P values are given in **S2 Table**.

1417

Supplementary fig 2.

(A)   Stack bar graph showing the number of viable organisms (mean ±

SEM) and their developmental stage at the specified hours after egg laying

for the indicated genotypes. Coloured arrows mark bars that exhibit rescue

of *STIM^{KO}* upon expression of STIM$^+$ driven by *THGAL4* (dark green)*,*

THC'GAL4 (blue)*, THD'GAL4* (green) and restricted rescue in presence *of*

THGAL80 (red) with all appropriate genetic controls as indicated. Number

of sets (N) = 3, number of organisms per set (n) = 25.

(B)   Stack bar graph showing the number of viable organisms (mean ±

SEM) and their developmental stage at specified hours after egg laying for

the indicated genotypes; *THD'/+, dicer;+;dsSTIM/+* (controls) and

THD'>dicer,dsSTIM. Number of sets (N)=3, number of organisms per set

(n)=25.

(C)   Quantification of weight of 10 flies from indicated genotypes. Box

plots show the 25^{th} and 75^{th} percentiles with median (bar), mean (square)

and each circle represents one set. Each set consists 10 flies of which 5

are females and 5   are male adult flies collected at   6-8h after eclosion.

Number of set (N) $\geq$5. Alphabets indicate different statistical groups.

In all panels significant changes between relevant genotypes at the

indicated stages were calculated by one way ANOVA followed by post-hoc

Tukey's test. P values are given in **S2 Table.**

1439

Supplementary fig 3.

(A)   Representative confocal images of larval brains showing THD'

neurons marked with anti-GFP (green) and anti-TH (red) from control

(*THD'>mCD8GFP*), *STIM^{KO}* (*STIM^{KO}*; *THD'>mCD8GFP)* and rescue

(*STIM^{KO};THD'>mCD8GFP, STIM^+*) animals at the indicated developmental

time points. Scale bars = 20μm.

(B)   Numbers of TH positive cells (magenta) and THD' cells (green) in

the larval CNS of the indicated genotypes. Cells were quantified from

THD'>mCD8GFP (control) at 58-62h and 80-84h and from

STIM^{KO};THD'>mCD8GFP (*STIM^{KO}*) and *STIM^{KO};THD'>mCD8GFP, STIM^+*

(rescue) at 80-84h. The numbers of TH^{+ve} and GFP^{+ve} cells were counted

manually and were no different among four hemi-lobes from four brains of a

single genotype and among all brain hemi-lobes of all genotypes. Hence the

absence of error bars.

(C) Stack bar graph showing the number of viable organisms (mean ± SEM)

and their developmental stage at specified hours after egg laying for the

indicated genotypes *THD'/+, dsTH/+* (controls), and *THD'>dsTH*. Number of

sets (N)=3, number of organisms per set (n)=25. Significant changes were

calculated by one way ANOVA followed by post-hoc Tukey's test. P values

are given in **S2 Table.**

(D) Representative images of the central brain (left panels) indicating the

region of focus (boxed), followed by images of THD' cells with Ca^{2+}

transients at the indicated time points after addition of a depolarizing agent

(KCl, 70mM). Ca^{2+} transients were measured in the indicated genotypes:

THD'>GCaMP6m and *THD'>dicer;dsSTIM,GCaMP6m* by measuring

changes in the intensity of GCaMP6m fluorescence. Scale bar = 20µm

(E) Quantified changes in GCaMP6m fluorescence (mean ± SEM of ∆F/F)

from THD' neurons of the indicated genotypes. Number of brains, (N) $\geq$ 6,

number of cells, (n) $\geq$11.

(F) Peak intensities of GCaMP6m fluorescence (∆F) in THD' cells from the

indicated genotypes. Box plots show 25^{th} and 75^{th} percentiles, the median

(bar), and mean (square) of ∆F of each cell (small circles).

1472

Supplementary fig 4.

(A-B) Altered excitability in THD' neurons by expression of either a

mammalian inward rectifying potassium channel (Kir2.1) or a bacterial

sodium channel (NachBac) affects larval developmental progression and

viability. Stack bar graphs show the number of viable organisms (mean $\pm$

SEM) and their developmental stage at the indicated hours after egg laying

(top) for the indicated genotypes. Number of sets (N) = 3, number of

organisms per set (n) = 25. P values calculated after one way ANOVA

followed by post-hoc Tukey's test are given for relevant stages and

genotypes in **S2 Table.**

(C) Representative images of the central brain (left panels) indicating the

region of focus (boxed), followed by images of THD' cells with Ca^{2+}

transients at the indicated time points after addition of mAChR agonist (CCh

100μM). Ca^{2+} transients were measured in the indicated genotype by

measuring the changes in the intensity of GCaMP6m fluorescence. Scale

bar = 20μm

(D) Quantified changes in GCaMP6m fluorescence (mean ± SEM of ∆F/F)

from THD' neurons of the indicated genotypes. Number of brains, (N) $\geq$ 4,

number of cells, (n) $\geq$12.

1492

Supplementary fig 5.

(A) Larval developmental profile with overexpression of ilp3 in MNSc

(red arrow) appears similar to knockdown of ilp5 (green arrow). Stack bar

graph showing the number of viable organisms and their developmental

stage at the specified hours after egg laying for the indicated genotypes.

N=3 sets with 25 larvae in each set.

(B) Knockdown and overexpression of ilp3 in the MNSc affects adult

weights in opposing directions. Quantification of weight of 10 flies from the

indicated genotypes. Box plot shows the 25th and 75th percentiles with

median (bar), mean (square). Each circle represents one set consisting of

10 flies in each of which 5 females and 5 males were collected at 6-8h

post- eclosion. A minimum of 5 sets were quantified for each genotype.

Alphabets indicate statistically different groups based on P values

calculated after one way ANOVA followed by post-hoc Tukey's test, given

for relevant stages and genotypes in **S2 Table.**

1508

Supplementary video 1: Mouth hook contractions in *CS* larvae (70-74h).

Supplementary video 2: Mouth hook contractions in $STIM^{KO}$ larvae (70-

74h).

Supplementary video 3: Mouth hook contractions in *CS* larvae (82-84h).

Supplementary video 4: Mouth hook contractions in $STIM^{KO}$ larvae (82-

84h).

Supplementary video 5: Mouth hook contractions in *THD'>dsTH* larvae

(82-86h).

Supplementary video 6: Mouth hook contractions in *THD'/+* (control)

larvae (82-86h).

Supplementary video 7: Mouth hook contractions in *dsTH/+* (control)

larvae (82-86h).

1521

1522   **Supplementary Table 1:** Expression status of genes encoding

1523   neuropeptides in $STIM^{KO}$ brains.

1524   **Supplementary Table 2:** P values for main and supplementary figures.

1525

A *THD'> SyteGFP, Denmark*

a — anti-RFP, anti-GFP, antiNC82

b — anti-RFP Post synaptic

c — anti-GFP Pre Synaptic

d

B *THD' > GRAB_DA*

Control at 58-62h   HL3 150s   CCh 260s

STIM^KO at 76-80h

Rescue at 80-84h

C

Control   STIM^KO   Rescue

HL3   CCh

ΔF/F

Seconds

D *THD'>ERGCaMP6-210*

Control at 58-62h   HL3 90s   CCh 300s

STIM^KO at 76-80h

Rescue at 80-84h

E

(i)

Control (Type-1)   STIM^KO
Control (Type-2)   Rescue

HL3   CCh

ΔF/F

Seconds

(ii)

No.of cells

Release of ER Ca^{2+}   No release of ER Ca^{2+}

Type 1   Type 2

(iii)

Seconds

A

633nm LED

● Dopaminergic cells
● Neuropeptide cells

C929 > mCherry

1 2

1. Median Neuro secretory cells (MNSc)

2. Lateral Neuro secretory cells (LNSc)

3. Sub Oesophageal Zone (SEZ)

B *THD'> Chrimson; C929 > GCaMP6f*

Chrimson activation

100s   300s   400s

MNSc

LNSc

SEZ

★ Activation   ★ Inhibition

C

Number of cells

━ Total cells
━ Activated cells
━ Inhibited cells

99   24   40

D

ΔF/F

Activation

ΔF/F$_{Max}$

Activation

Seconds

E

ΔF/F

Inhibition

100 200 300 400

ΔF/F$_{Min}$

Inhibition

Seconds

F *THD' > Chrimson; C929 > GCaMP6f*

(i) Activation

ΔF/F

Seconds

(ii) Inhibition

ΔF/F

Seconds

G *THD' > Chrimson; C929 >GCaMP6f*

ΔF/F

Without ATR and light

seconds

A

CS 58-62h | STIM^KO 72-76h

ilp2
ilp3
ilp5
ilp6
ilp7

2.5 ——— 260
Normalised gene counts

B

■ CS (58-62h: mid L2)
■ STIM^KO (72-76h:mid L2)
■ THD' rescue (56-62h:mid L2)

Relative fold change

ilp2   ilp3   ilp5

C

ilp3

L1
L2
L3(12h)
L3 puffstage(1-2)
L3 puffstage(3-6)
L3 puffstage(7-9)
White prepupae(new)
White prepupae(12h)
white prepupae (24h)

0 ——— 5
Normalised gene counts

D

No. of viable organisms

60-66h   80-86h   128-136h   320-326

□ I instar
□ II instar
□ III instar
□ Pupa
□ Adult

CS
STIM^KO;MNSc/+
STIM^KO;dsilp3/+

E

at 118-122hh

L3   L2   L3a   L3b

CS   STIM^KO; MNSc>dsilp3

F

at 118-122h

Larval length (mm)

CS   dsilp3 rescue (L3b)

G

STIM^KO; MNSc>dsilp3

H

■ MNSc/+   ● dsilp3/+
▲ MNSc>dsilp3

No. of larvae(L3)

96 108 120 132 144 156
hAEL

I

at 144-156h

MNSc/+   dsilp3/+   MNSc>dsilp3

J

at 144-156h

Pupal volume (mm³)

MNSc/+   dsilp3/+   MNSc>dsilp3

K

■ MNSc/+   ● ilp3⁺/+
▲ MNSc>ilp3⁺

No. of larvae(L3)

60   72   84   96
hAEL

L

at 94-98h

MNSc/+   ilp3⁺/+   MNSc>ilp3⁺

M

at 94-98h

Larval length (mm)

MNSc/+   ilp3⁺/+   MNSc>ilp3⁺

N

at 82-86h

No. of contraction per 30s

n.s.

MNSc/+   ilp3⁺/+   MNSc>ilp3⁺

A

60-66h 80-86h 128-134h 168-174h 320-326h

No. of viable organisms /25

I instar
II instar
III instar
Pupa
Adult

CS
MNSc/+
dsDop1R1/+
dsDopEcR/+
dsDop2R2/+

B

at 128-134h

MNSc/+   dsDop1R1/+   MNSc>dsDop1R1

C

Average weight of 10 flies (mg)

a   a

b

MNSc/+   dsDop1R1/+   MNSc>dsDop1R1

D

MNSc>GCaMP6

HL3   100uM DA+ 1uMTTX

50s   150s   250s

HL3

Activation   Inhibition

E

MNSc>GCaMP6m

— DA+TTX (Inhibition)   — DA+TTX (Activation)

— No DA+No TTX

ΔF/F

HL3   100uM DA

seconds