RESEARCH ADVANCE

TITLE: Ventral striatal dopamine encodes unique properties of visual stimuli in mice

AUTHORS: L. Sofia Gonzalez[1,2], Austen A. Fisher[1,2], Evelin M. Cotella[1], and J. Elliott Robinson*[1]

[1] Division of Experimental Hematology and Cancer Biology, Department of Pediatrics, Cincinnati

Children's Hospital Medical Center, University of Cincinnati College of Medicine, Cincinnati, OH,

United States

[2]These authors contributed equally.

*To whom correspondence should be addressed:

Dr. J. Elliott Robinson

Division of Experimental Hematology and Cancer Biology

Department of Pediatrics

Cincinnati Children's Hospital Medical Center

3333 Burnet Ave.

Cincinnati, OH 45229

Phone: (513) 803-5122

elliott.robinson@cchmc.org

ABSTRACT:

Previously, we used the genetically encoded dopamine sensor dLight1 to show that mice modeling the neurodevelopmental disorder neurofibromatosis type 1 (NF1) have enhanced dopaminergic responses to salient visual stimuli in the lateral nucleus accumbens/ventral striatum (LNAc), as well as more vigorous behavioral responses to visual threats (Robinson et al., 2019). Here, we sought to better understand the nature of this phenotype by testing the ability of striatal dopamine release to encode the properties of sensory stimuli, including threatening looming discs, in typically developing mice. In this Research Advance, we report that ventral striatum dopamine release encodes the rate and magnitude of environmental luminance changes rather than the perceived threat intensity of the stimulus. These visually-evoked dopamine responses are wavelength-dependent at low irradiances, independent of the circadian cycle, and relatively robust to previous exposure history. Thus, we have further elaborated the mesolimbic dopamine system's ability to encode visual information in mice, which may be altered in neurodevelopmental disease models.

INTRODUCTION:

The mesolimbic dopamine system is an evolutionarily conserved set of circuits that plays a role in approach and avoidance, appetitive behavior, and reward processing (Wise, 2004; Everitt and Robbins, 2005; Alcantara et al., 2022). In this circuitry, ascending dopaminergic projections from the ventral midbrain, including the ventral tegmental area (VTA), innervate targets throughout the limbic forebrain, such as the ventral striatum/nucleus accumbens (NAc). Dopaminergic signaling in the NAc has been widely studied for its involvement in motivational salience, behavioral reinforcement, and reward prediction error encoding (Schultz et al., 2015; Berridge and Robinson, 2016; Watabe-Uchida et al., 2017; Berke, 2018). Less well characterized is the role of dopaminergic neurotransmission in the response to unpredicted or alerting sensory events, which may encourage investigation or prime motivated behavioral responses to these stimuli (Horvitz, 2000; Bromberg-Martin et al., 2010a; Schultz, 2010). While many previous studies have reported phasic firing of dopaminergic neurons in response to light flashes in laboratory animals (Horvitz et al., 1997; Comoli et al., 2003; Dommett et al., 2005), it is unclear how NAc dopamine release encodes the properties and/or emotional valence of arousing visual stimuli, such as visual threats.

Across a range of species (Ball and Tronick, 1971; Sun and Frost, 1998; Maier et al., 2004; Nakagawa and Hongjian, 2010; Yilmaz and Meister, 2013; Temizer et al., 2015), rapidly approaching objects or looming visual threats elicit automatic defensive or avoidance responses. In mice, presentation of an expanding, overhead, black disc that simulates aerial predator approach (a looming stimulus) promotes rapid escape to an available shelter, followed by long periods of freezing (Yilmaz and Meister, 2013). In our previous work published in eLife (Robinson et al., 2019), mice modeling cognitive dysfunction associated with neurofibromatosis type 1 (NF1) exhibited more vigorous escape in response to looming stimulus presentation. Additionally, NAc dopamine release evoked by a white light stimulus was enhanced in NF1 model mice, which was correlated with behavioral conditioning abnormalities. Despite the demonstration that white light

can induce NAc dopamine release (Robinson et al., 2019; Kutlu et al., 2021), the striatal dopamine response to visual threats is not well characterized in mice. Additionally, it is unknown what visual stimulus characteristics – if any – are encoded by NAc dopamine. Thus, one cannot fully interpret the significance of aberrant responses in NF1 mice without a more thorough understanding of visual stimulus encoding by mesolimbic dopamine release in typically developing subjects.

In this Research Advance, we sought to probe ventral striatal dopaminergic responses to arousing visual stimuli, including looming visual threats. Given the ability of dopaminergic neurons to signal stimulus saliency (Bromberg-Martin et al., 2010b), we hypothesized that looming discs would induce 'alerting' NAc dopamine release whose magnitude would scale proportionately with perceived threat intensity. To test this hypothesis, we utilized the genetically-encoded sensor dLight1 (Patriarchi et al., 2019) to monitor dopamine release in the NAc lateral shell (LNAc) of freely moving adult C57Bl/6J mice with fiber photometry, as performed previously (Robinson et al., 2019). The LNAc was chosen because dopamine release and/or VTA dopaminergic axon terminal activity encode both stimulus valence and prediction errors in this region (de Jong et al., 2018; Robinson et al., 2019; Yuan et al., 2019). Here, we report that LNAc dopamine release reliably reads out unique visual stimulus properties in mice, a phenomenon that is likely relevant to a wide body of researchers employing light sources or optical methods in behavioral research.

RESULTS:

Dopaminergic responses to looming visual threats.

To explore the encoding of visual threats by LNAc dopamine, we first measured dLight1 signals evoked by looming discs (Figure 1A-D) using a custom Bonsai-controlled (Lopes et al., 2015) setup for programmable visual stimulus presentation on an overhead liquid crystal display (LCD) within a light and sound-attenuating chamber (see *Materials and Methods* for details). During photometry recordings, mice were exposed to trains of five overhead, black, looming discs

on a light gray background that we empirically determined to produce short-latency escape in

C57Bl/6J mice (Figure 1E, Video 1), consistent with previous studies (Evans et al., 2018; Yilmaz

and Meister, 2013). As controls, we presented mice with trains of black discs that do not reliably

evoke defensive responses (Figure 1E), such as a static disc (a fixed 30.5 cm black disc on a

light gray background), a receding disc (a black disc that contracted from 30.5 cm to 0 cm on a

light gray background), and contrast inverted discs (light gray static, looming, or receding discs

on a black background). We observed that looming discs induced low-amplitude dopamine

transients at the onset of the first stimulus in each train that – contrary to our hypothesis – were

not significantly different from the dLight1 responses to non-threatening static and receding discs

(Figure 1B-C). Surprisingly, repeating these experiments with contrast-inverted discs that do not

induce escape (Figure 1E) evoked ~3 to 6-fold greater dopamine release than black discs (Figure

1B-C). This raised the possibility that LNAc dopamine release tracks stimulus brightness rather

than threat intensity.

Dopaminergic responses to rapid changes in environmental lighting conditions.

Because inverted looming discs, in which the number of bright overhead pixels ramps as

the disc expands, produced lower amplitude (Figure 1C), longer latency (Figure 1D) dLight1

responses than static or receding inverted discs with an instantaneous pixel change, we

hypothesized that LNAc dopamine may encode the rate of change of dark-to-light transitions. To

test this possibility, we exposed mice to full screen, instantaneous transitions from black to light

gray during dLight1 recordings, which eliminated disc edge motion as a contributing visual

stimulus property. We found that instantaneous dark-to-light transitions produced a high

amplitude (10.38 ± 0.43 z-score), short duration (full width a half maximal amplitude: 143 ± 9.7

ms) dopamine transient that peaked 434 ± 3.3 ms after transition onset (Figure 2A). Lengthening

the dark-to-light transition time (i.e. the fade-in time) to full screen illumination (Figure 2B) non-

linearly decreased the magnitude of the dLight1 peak and increased the peak latency (Figure 2C).

For transition times less than ~500 ms, the dopamine peak latency closely matched the fade-in

time, above which peak response occurred hundreds of milliseconds to seconds before full field

illumination was reached (Figure 2C). When transition times were greater than 1 s, evoked

dLight1 transients were often too small to accurately resolve from the fluorescent baseline for

individual mice. However, averaging the fluorescence trace from all mice prior to peak detection

allowed signals to be resolved for longer transition times. Thus, results are presented as both the

fluorescence peak(s) derived from the photometry trace averaged across all mice (Figure 2C) and

individual mice (Figure 2-Figure Supplement 1), which showed high concordance for transition

times of 1 second or less (Figure 2-Figure Supplement 1). No dLight1 response was reliably

evoked by a ten-second dark-to-light transition despite the stimulus ramping to the same number

of bright pixels as trials with shorter transition times (Figure 2B).

Next, we examined whether LNAc dopamine release also reads out the magnitude of

environmental lighting changes by measuring the dLight1 response to ten-second, instantaneous

exposures to white light across a range of intensities (0.2 nW/cm^2 – 5.0 μW/cm^2, measured at

mouse level) generated by a light emitting diode (LED) (Figure 2D-F). High irradiance LED

illumination (5 μW/cm^2) evoked a dLight1 transient at stimulus onset (Figure 2D) that was similar

to transients evoked by the LCD monitor (Figure 2A-B; irradiance at mouse level: 11 μW/cm^2).

This response did not depend on the time of testing within the vivarium day-night cycle (Figure

2G) and was larger than the response to auditory tones (80 dB; 1 – 16 kHz; Figure 2-Figure

Supplement 1). When LED intensity was reduced, we observed an intensity-dependent decrease

in the magnitude of the dLight1 peak and increase in the peak response latency (Figure 2E-F).

Significant dopaminergic responses were observed at all irradiances tested, including 0.2

nW/cm^2, which was not perceptible to the human experimenter. As a point of reference, the lock

screen of a Samsung S21 smart phone on the lowest brightness setting had an irradiance of 20

nW/cm^2 when placed in the same position as the white LED. Likewise, time-locked dopamine

release could be evoked by simply uncovering the enclosure peephole that allows users to

observe mouse behavior (irradiance: 17 nW/cm^2; Figure 2-Figure Supplement 1). These results

were not likely caused by mouse movement, as illumination of a white LED that was 1000-fold

more intense (5 mW/cm^2) than the highest irradiance tested had no appreciable effect on behavior

when freely exploring mice entered a target zone within a dark arena (Video 2). Thus, LNAc

dopamine release is sensitively evoked by ambient light and reliably encodes the speed of these

lighting transitions over short timescales.

Dopaminergic responses to repeated light stimuli.

Previous literature suggests that dopaminergic neuron firing in response to sensory events

habituates as the novel stimulus becomes familiar (Schultz, 1998). In order to test if the dLight1

response to white light is affected by a history of previous exposures, we exposed mice to twenty

consecutive one-second white light pulses over five trials (100 pulses total) across a range of

interstimulus intervals (ISIs; 10 ms to 10 s; Figure 3A). We found that light-evoked dopamine

transient magnitude decayed logarithmically as a function of the ISI duration (Figure 3B). When

the ISI was short (e.g. 10 – 100 ms), dLight1 responses habituated rapidly. This is exemplified by

the dopaminergic response to 40 Hz light flicker, which is used therapeutically to enhance neural

activity in the context of Alzheimer's disease (Singer et al., 2018). Presentation of a sixty-second

40 Hz white LED flicker (5.0 μW/cm^2 irradiance, 50% duty cycle) induced a dopamine transient

only at stimulus onset (Figure 3C) that was indistinguishable from the response to constant

illumination (Figure 2D). Conversely, in shorter experiments when the ISI was long (100 s), no

habituation of the dopamine response to light was observed from stimulus-to-stimulus (Figure

3D). Repeating this experiment immediately after three hundred consecutive one-second

exposures with a one-second ISI produced an ~20% reduction in the dLight1 response to 100 s

ISI light; however, this effect did not reach statistical significance (p = 0.08) (Figure 3D). Therefore,

habituation of the dopamine response to repeated light stimuli is much more strongly influenced

by stimulus frequency than the total number of previous exposures.

During repeated stimulus experiments, the greatest reduction in the peak LNAc dopamine

response to light stimuli occurred between the first and second light pulse in each stimulus train.

In order to further characterize this phenomenon, we varied the duration of the first stimulus to

determine if the total amount of initial light exposure modulates the dopaminergic response to a

subsequent stimulus (Figure 3E). We found that the dopaminergic response to a 1 second white

LED test stimulus was not significantly different when preceded by either a 300 second or 1

second preconditioning light stimulus one second earlier (Figure 3F). No difference in dLight1

response to the preconditioning stimulus was observed between conditions (Figure 3F). We did

observe, however, that the 300-second preconditioning stimulus produced a dopaminergic

response at light offset, whereas the 1-second preconditioning stimulus did not (Figure 3G).

Taken together, our findings indicate that ISI is a more significant determinant of stimulus-to-

stimulus dopamine release habituation than light stimulus duration.

Dopaminergic responses to individual wavelengths across the visual spectrum

In these and previous experiments (Robinson et al., 2019), we employed a white LED light

to induce striatal dopamine release; however, this light source is composed of multiple

wavelengths throughout the visible spectrum. Therefore, we next investigated if light-evoked

dopamine release exhibits wavelength specificity. This is additionally germane given the

widespread use of molecular and optical technologies in rodents that require delivery of specific

wavelengths of visible light in order to probe neural activity, structure, or biology (Fenno et al.,

2011; Resendez and Stuber, 2015; Sabatini and Tian, 2020). In order to determine if the dLight1

response varied by wavelength, we measured dopamine release induced by ten-second

exposures to environmental ultraviolet (UV; 360 nm), blue (475 nm), green (555 nm), red (635

nm), and far-red (730 nm) light across a 100,000-fold range of irradiances (1 nW/cm^2 to 100

μW/cm^2). These experiments revealed broad sensitivity of LNAc dopamine release to light across

the visual spectrum (Figure 4A-B). The dopamine response was least sensitive to UV and red

light when the irradiance was low (1 nW/cm^2), and far-red light (730 nm) only induced dopamine

release when the irradiance was very high (100 μW/cm^2). The ability of red light to induce

dopamine release at intensities as low as 0.1 μW/cm^2 is consistent with research that rodents are

better at perceiving red wavelengths than is commonly acknowledged (Danskin et al., 2015;

Nikbakht and Diamond, 2021). Whereas the dLight1 response to UV and red light was irradiance-

dependent (Figure 4C), the response to blue and green light remained robust across the entire

irradiance range (Figure 4A). These experiments suggest that the mesolimbic dopamine system

is responsive to all visible wavelengths yet is most sensitive to blue and green light.

DISCUSSION

In these investigations, we used the genetically-encoded dopamine sensor dLight1 to

demonstrate that LNAc dopamine release can encode rapid changes in luminance but not

looming threat intensity. We found that rapid dark-to-light transitions evoked time-locked

dopamine responses at stimulus onset at irradiances as low as 0.2 nW/cm^2, which is consistent

with findings that mice see over a 100 million-fold range of light intensity beginning at ~4 μcd/m^2

(Umino et al., 2008). The magnitude of these dopaminergic responses were highly dependent on

light stimulus frequency and transition rate rather than duration or novelty. In fact, high amplitude

LNAc dLight1 responses to a white LED persisted after hundreds of exposures and months of

regular testing. Although mesolimbic dopamine systems regulate wakefulness (Eban-Rothschild

et al., 2016) and exhibit circadian oscillation (Korshunov et al., 2017), the time of testing did not

appear to be a significant contributor to our findings. Sudden dark-to-light transitions are highly

salient to nocturnal rodents that must avoid detection by visual predators (Thompson et al., 2010),

so it is possible that the dopaminergic response to light represents a specialized saliency signal.

Previously, phasic dopamine responses to unexpected, unconditioned sensory events have been

conceptualized in the context of novel action discovery (Redgrave and Gurney, 2006), the need to alert to stimuli that require motivated responses (Schultz and Romo, 1990; Horvitz, 2000), mechanisms of associative learning (Lisman and Grace, 2005), etc. Thus, further studies will be needed to firmly establish the ethological and neurobiological importance of dopaminergic responses to environmental light.

We also found that LNAc dopamine is broadly evoked by wavelengths across the visual spectrum. The mouse visual system utilizes numerous opsin proteins for image-forming and non-image forming phototransduction with unique wavelength sensitivities. These include the rod opsin rhodopsin for scoptopic (low-light) vision (λmax = 498 nm) and short (λ_{max} ~360 nm) and medium/long wavelength (λ_{max} = 508 nm) cone opsins for photopic vision (Peirson et al., 2018). Additionally, melanopsin (λ_{max} ~480 nm) (Lucas et al., 2001; Peirson et al., 2018) is expressed in intrinsically photosensitive retinal ganglion cells (ipRGCs) that are involved in circadian entrainment, the pupillary light reflex, and light-regulated changes in mood (Panda et al., 2003; Hattar et al., 2003; Fernandez et al., 2018). Given the high proportion of rods in the mouse retina (~97% of photoreceptors) (Jeon et al., 1998) and the reduced sensitivity of dLight1 responses to 360 and 635 nm light at lower irradiances, it is probable that rod-based phototransduction is primarily responsible for visually-evoked dopamine release under dim lighting conditions. Conversely, multiple opsin types likely contributed to dLight1 signals when LED irradiance was higher. While it has been hypothesized that ipRGCs engage VTA outputs via a disynaptic circuit involving the preoptic area (Zhang et al., 2021), the relative insensitivity (~1000-fold lower than cone-based vision) (Lall et al., 2010) and slow speed of melanopsin phototransduction (Berson et al., 2002) may reduce the likelihood that it significantly influenced the observed findings. Repeating dLight1 investigations in mutant mice lacking essential components of rod-, cone-, and non-imaging forming phototransduction will help parse the role of these pathways in dopaminergic responses to light.

Visual information is conveyed from the retina to the brain via the axons of retinal ganglion

cells that synapse in downstream nuclei to mediate image processing, circadian entrainment, pupillary reflexes, gaze orientation, etc. (Peirson et al., 2018). While thalamocortical visual pathways are required for conscious visual perception, neither the primary visual cortex (V1) nor the visual thalamus (e.g. lateral geniculate nucleus) significantly innervate ventral midbrain dopamine neurons (Watabe-Uchida et al., 2012). Previous work by Redgrave and colleagues suggest that dopaminergic responses to light are driven by the superior colliculus (SC) (Comoli et al., 2003; Dommett et al., 2005; Takakuwa et al., 2017), which receives direct input from retinal ganglion cells (Dhande and Huberman, 2014) in its superficial layers and promotes motivated behavior via deep motor-output layers (Branco and Redgrave, 2020). SC glutamatergic projection neurons directly synapse onto VTA (Solié et al., 2022) and substantia nigra pars compacta dopamine neurons (Huang et al., 2021), both of which project to the LNAc (Beier et al., 2015; Poulin et al., 2018). Likewise, optogenetic stimulation of SC neuron somata is sufficient to evoke LNAc dopamine release *in vivo* (Robinson et al., 2019). While these observations support a role for the SC in dopaminergic responses to light, the relative contribution of different visual processing centers to our findings is an important area of future study.

Mesolimbic dopaminergic circuits are thought to play a role in the pathophysiology of several neuropsychiatric conditions, including disorders of impulse control, schizophrenia, and neurodevelopmental disorders (Li et al., 2006; Purper-Ouakil et al., 2011; Maia and Frank, 2017; Robinson and Gradinaru, 2018), including NF1 (Brown et al., 2010; Diggs-Andrews et al., 2013; Anastasaki et al., 2015). Patients with NF1 exhibit high rates of attention deficit/hyperactivity disorder (Mautner et al., 2015; Miguel et al., 2015), in which difficulties with attentional orientation are associated with a diminished ability to suppress distractive stimuli (Aboitiz et al., 2014) such that irrelevant environmental cues are assigned exaggerated stimulus salience (Tegelbeckers et al., 2015). Previously in eLife, we showed that dopaminergic responses to light are enhanced in NF1 model mice and correlate with disruptions in the expression of conditioned behavior (Robinson et al., 2019). Our current findings suggest that these responses reflect changes in the

encoding of environmental lighting conditions and, given their correlation with phenotypic expression, may reflect altered stimulus saliency. Alternatively, it has been recently reported that cultured retinal ganglion cells from *Nf1* mutant mice display increased network activity *in vitro* (Anastasaki et al., 2022). As such, enhanced dopaminergic responses to light may reflect gain of function within retinal microcircuits. Going forward, it will be imperative to more comprehensively examine visual system structure and function in NF1 mouse models *in vivo* and *in situ*.

MATERIALS AND METHODS

Key Resources Table				
Reagent type (species) or resource	**Designation**	**Source or reference**	**Identifiers**	**Additional information**
Recombinant DNA reagent	pAAV-hSyn-dLight1.2	Addgene	Cat#: 111068 RRID:Addgene_111068	Produced by Addgene in the AAV5 serotype
Software, Algorithm	Python 3.8	Python Software Foundation	RRID:SCR_008394	
Software, Algorithm	Fiber Photometry Trace Processing	Tucker-Davis Technologies		https://www.tdt.com/docs/sdk/offline-data-analysis/offline-data-python/FibPhoEpocAveraging/
Software, Algorithm	Bonsai 2.6.3	Bonsai Foundation CIC	RRID:SCR_017218	
Software, Algorithm	Looming Visual Stimulus Generation	Austen Fisher, Robinson Lab		https://github.com/jelliottrobinson/Robinson_Lab
Software, Algorithm	ABET II Software for	Lafayette Instrument	Model 89501	

	Operant Control	Company		
Software, Algorithm	Ethovision XT 17	Noldus Information Technology	RRID:SCR_0 00441	
Software, Algorithm	GraphPad Prism 9	GraphPad Software, Inc.	RRID:SCR_0 02798)	
Software, Algorithm	Data Science Workbench 14.0.0.15	TIBCO Software, Inc.	RRID:SCR_0 14213	
Other	Mono Fiber-Optic Cannula	Doric Lenses, Inc.	Cat#: MFC_400/430 - 0.66_6mm_M F1.25_FLT	OD: 400 μm, Length: 6 mm
Other	Mono Fiber-Optic Patch Cable	Doric Lenses, Inc.	Cat#: MFP_400/430 /1100- 0.57_1m_FC M- MF1.25_LAF, Doric Lenses Inc.	OD: 400 μm, Length: 1 m

Experimental Animals

Experimental subjects were adult, male and female C57Bl/6J mice (the Jackson

Laboratory Stock No: 000664) that were greater than 12 weeks of age. Animals were pair or group

housed (3-4 per group) throughout the duration of the experiment in a vivarium on a 14-hour/10-

hour light/dark cycle (lights on at 0600 hrs, lights off at 2000 hrs) with *ad libitum* access to food

and water. All experiments were performed during the light phase of the vivarium light/dark cycle,

except when white LED exposure was performed 2-3 hours into the dark phase, as shown in

Figure 2G. Animal husbandry and experimental procedures involving animal subjects were

conducted in compliance with the Guide for the Care and Use of Laboratory Animals of the

National Institutes of Health and approved by the Institutional Animal Care and Use Committee

(IACUC) and by the Department of Veterinary Services at Cincinnati Children's Hospital Medical

Center (CCHMC) under IACUC protocol 2020-0058. Mice were only excluded from studies if they

could not complete an entire experiment due to health concerns or loss of brain implant, if there

was no dynamic photometry signal six weeks after surgery, or if the location of the photometry

fiber tip was histologically determined to be outside the lateral nucleus accumbens (NAc).

Surgical Procedures

Stereotaxic viral vector injections and optical fiber implantation surgeries for dLight1 were

performed as previously described (Robinson et al., 2019). This procedure was similar to the

published protocol of Tian and colleagues (Patriarchi et al., 2019). In brief, mice were

anesthetized with isoflurane (1 – 3% in 95% O_2/5% CO_2 provided via nose cone at 1 L/min), the

scalp was shaved and sterilized with chlorhexidine surgical scrub, the skull surface was exposed,

and a craniotomy hole was drilled over the lateral NAc (antero-posterior: 1.2 mm, medio-lateral:

1.6 mm relative to Bregma). 800-1000 nL of a AAV5-hSyn-dLight1.2 vector (~1 × 10^{13} viral

genomes/mL, obtained from Addgene; catalog #AAV5-111068) was delivered into the LNAc

(antero-posterior: 1.2 mm, medio-lateral: 1.6 mm, dorso-ventral: -4.2 mm relative to Bregma)

using a blunt or beveled 34 or 35-gauge microinjection needle within a 10 uL microsyringe

(NanoFil, World Precision Instruments) controlled by a microsyringe pump with SMARTouch

Controller (UMP3T-1, World Precision Instruments) over ten minutes. Following viral injection, a

6 mm long, 400 µm outer diameter mono fiber-optic cannula (MFC_400/430-

0.66_6mm_MF1.25_FLT, Doric Lenses Inc.) with a metal ferrule was lowered to the same

stereotaxic coordinates and affixed to the skull surface with C&B Metabond (Parkel Inc.) and

dental cement. Mice were given 5mg/kg carprofen (s.c.) intraoperatively and for two days

postoperatively for pain. Mice were allowed a minimum of five weeks for surgical recovery and

virus expression prior to participation in behavioral studies.

Fiber Photometry

Fluorescent signals were monitored using an RZ10x fiber photometry system from Tucker-Davis Technologies, which allowed for dLight1 excitation and emission light to be delivered and collected via the same implanted optical fiber. Our system employed a 465-nm LED for sensor excitation and a 405-nm LED for isosbestic excitation. Light was filtered and collimated using a six channel fluorescent MiniCube [FMC6_IE(400-410)_E1(460-490)_F1(500-540)_E2(555-570)_F2(580-680)_S] from Doric Lenses, Inc., which was coupled to the implanted optical fiber via a one-meter, low autofluorescence fiber optic patch cable (MFP_400/430/1100-0.57_1_FCM-MF1.25LAF, Doric Lenses Inc.). The emission signal from 405 nm isosbestic excitation was used as a reference signal to account for motion artifacts and photo-bleaching. A first order polynomial fit was applied so that the 465 nm signal to be aligned with the 405 nm signal. Then, the polynomial fitted model was subtracted from the 465 nm channel to calculate ΔF values. The code for performing this function was provided by Tucker-Davis Technologies, Dr. David Root (University of Colorado, Boulder), and the laboratory of Dr. Marisela Morales (NIDA); it is available at: https://www.tdt.com/docs/sdk/offline-data-analysis/offline-data-python/FibPhoEpocAveraging/.

During behavioral experiments, the ΔF time-series trace was z-scored within epochs to account for data variability across animals and sessions, as described by Morales and colleagues (Barker et al., 2017). When fiber photometry was performed during sensory stimulus exposure experiments, dLight1 signals were synchronized to stimulus onset via delivery of TTL pulses to the photometry system. Peak data (magnitude, latency, and full width at half-maximal intensity) was analyzed using Python.

Visual Stimulus Exposure

Visual stimuli were delivered to mice during fiber photometry recordings within a custom setup that featured a 24 inch liquid crystal display (LCD) mounted 25.4 cm above mouse level in a light and sound attenuating chamber (Model 83018DDP, Lafayette Instrument Company).

Stimuli (looming, static, and receding discs; full screen fades; etc.) were generated on the LCD

display using Bonsai (Lopes et al., 2015), which also controlled delivery of a TTL pulse to the

photometry system via a BNC cable to timestamp stimulus onset. The TTL pulse was generated

with an Arduino Uno Rev3 microcontroller. During each experiment, mice were placed within the

bottom of a clean shoebox cage with a thin layer cob bedding in the light and sound attenuating

chamber underneath the LCD. Looming discs expanded from 0 cm to 30.5 cm over 0.84 s and

froze at full expansion for 0.26 s, encompassing 61.9 degrees of visual angle, as previously

described (Evans et al., 2018; Yilmaz and Meister, 2013). Receding discs shrunk from 30.5 cm

to 0 cm over 0.84 s. Static discs maintained their 30.5 cm diameter throughout the duration of the

stimulus. During each stimulus train, five discs were shown consecutively with a 0.5 ms

interstimulus interval (ISI). Mice were exposed to five stimulus trains with a 600 s inter-trial interval

(ITI). Single full field fades from black to light gray (0 – 10 s fade duration) were delivered via the

LCD screen across 5 trials with a 120 s ITI.

White LED exposures were delivered via the house light of a modular conditioning

chamber (Model 80015NS, Lafayette Instruments Company) placed within the light and sound

attenuating box and controlled by ABET II software (Lafayette Instrument Company). A TTL

breakout adapter (Model 81510) was used to synchronize stimulus delivery with the photometry

recording. Single ten-second light stimuli were delivered across ten trials with a randomized ITI

between 90 and 180 s. Glass neutral density filters were used to attenuate the irradiance when

necessary (0.1 – 3.0 OD, HOYA Filter USA and/or Edmund Optics TECHSPEC filters). Trains of

20 one-second light stimuli with variable ISIs (10 ms – 10 s) were delivered across five trials (100

total exposures) with a 300 s ITI. One-second light stimuli with a 100 s ISI were delivered before

and after 300 one-second light stimuli with a one-second ISI; 100 s separated the 300 one-second

stimuli and each 100 s ISI stimulus train. One minute of 40 Hz flicker exposure (50% duty cycle)

was repeated across 5 trials with a 120 s ITI. For paired light stimuli experiments, a one-second

white LED stimulus was delivered one second after a 300 s or 1 s light stimulus across five trials

with a 300 s ITI.

Individual wavelength light stimuli were generated with a Lumencor Aura III LED light engine, which was triggered via TTL inputs from the Lafayette Instruments TTL breakout adapter and controlled by ABET II. The liquid light guide that delivered the visual stimulus was positioned in the approximate location of the white LED within the testing chamber. LED light power (measured at mouse level with a Thor Labs PM100D optical power meter with S130VC photodiode sensor) was modulated using the onboard Lumencor graphical user interface and, when necessary, attenuated via the use of glass neutral density filters (0.1 – 3.0 OD, HOYA Filter USA and/or Edmund Optics TECHSPEC filters) placed in front of the liquid light guide outlet within a custom housing. Ten-second single wavelength stimuli were delivered in random order with a randomized ITI (140 – 200 s) to achieve five total exposures per color per mouse.

Auditory Stimulus Exposure

Auditory stimulus exposures were performed in the modular testing chamber within the light and sound attenuating enclosure similarly to single white LED exposures. A ten-second 80 dB tone (1 – 16 kHz; generated via Lafayette Instruments 7 Tone Generator Model 81415M) was presented via a speaker (0.25 – 16 kHz; Model 80135M14, Lafayette Instrument Company) across 5 trials with a randomized ITI (140 – 200 s).

Looming Stimulus Assay

The looming stimulus assay was performed as previously described (Yilmaz and Meister, 2013) using an apparatus built to the specifications of Evans et al. (2018). The apparatus featured a 20.3 cm (w) x 61 cm (l) x 40.6 cm (h) clear, open, rectangular acrylic arena with a dark, infrared (IR) light-transmitting shelter at one end and a 'threat zone' at the opposite end that housed a 9 cm clear plastic petri dish to encourage exploration outside of the shelter. A 15.6 inch monitor was mounted above the arena so that discs (19.5 cm maximum diameter encompassing 27

degrees of visual angle) could be presented to the mice when they entered the threat zone. The

arena floor was backlit with an infrared light (880 nm back-lit collimated backlight, Advanced

Illumination) to improve mouse tracking under dim light conditions. The entire apparatus was

placed inside a custom light-attenuating enclosure for testing. During testing, mice were recorded

with a Basler acA2040-120 um camera with a Edmunds Optics TECHSPEC 6mm C Series fixed

focal length lens, and real time position tracking was performed with Bonsai. This allowed for

presentation of the overhead looming, receding, or static disc stimulus to be automatically

triggered when the animal was in the threat zone following a ten-minute habituation period. Mouse

position and velocity data was analyzed *post hoc* using Ethovision XT software (Noldus

Information Technology) and Python. Note: In Video 1, the clear, circular pedestals that separated

the infrared backlight from the apparatus base can be seen with the IR camera; they were below

the arena floor and inaccessible to the mouse. The setup was modified for spotlight experiments

so that the pedestals would not be visible in the captured videos.

Spotlight experiments were performed in the same apparatus using the same procedure

described above except that a high intensity white LED (5 mW/cm^2 measured at mouse level)

positioned to illuminate the threat zone replaced the LCD monitor.

Statistical Analysis

Statistical analysis was performed using Python, GraphPad Prism 9 (GraphPad Software,

Inc.), and/or Data Science Workbench 14 (for 3-way repeated measures ANOVA; TIBCO

Software, Inc.). All statistical tests performed on data presented in the manuscript are stated in

the figure captions and provided in detail with the corresponding source data in the Supplemental

Data and Statistical Analysis file. For each experiment, statistical tests were chosen based on the

structure of the experiment and data set. No outliers were removed during statistical analysis.

Parametric tests were used throughout the manuscript. Sample size estimates were based on

studies in Robinson et al. (2019) and power analysis performed using the sampsizepwr() function

in Matlab (MathWorks). When analysis of variance (ANOVA; 1-way, 2-way, 3-way, and/or repeated measures) was performed, multiple comparisons were corrected using the Bonferroni correction. When repeated measures ANOVA could not be performed due to missing values (Figure 2-Figure Supplement 1, panel C), data was analyzed by fitting a mixed model in GraphPad Prism 9; this approach uses a compound symmetry covariance matrix and is fit using restricted maximum likelihood (REML). When results were compared to a pre-stimulus baseline, this value was defined as the amplitude of the dLight1 peak that occurred 500 ms prior to stimulus delivery. When results were compared to a 'null' stimulus, the value was defined as the dLight1 peak that occurred at the onset of a TTL that timestamped a trial in which no stimulus was delivered.

Data and Materials Availability

Viral vector plasmids used in this study are available on Addgene. Codes used for fiber photometry signal extraction and analysis are available at https://www.tdt.com/docs/sdk/offline-data-analysis/offline-data-python/FibPhoEpocAveraging/. Codes used for visual stimulus generation are available at https://github.com/jelliottrobinson/Robinson_Lab. Source data is available in the provided Supplemental Data and Statistical Analysis file.

ACKNOWLEDGEMENTS

We would like to acknowledge Mary Claire Casper and Shiva Senthilkumar for assistance with histological sample preparation and preliminary data analysis, respectively. We would also like to thank Dr. Richard Lang and Shane D'Souza at Cincinnati Children's Hospital Medical Center, Dr. Gregory Schwartz at Northwestern University Feinberg School of Medicine, and Dr. Diego Fernandez at the National Institute of Mental Health for helpful discussions regarding technical considerations and/or interpretation of the experimental findings. This work was funded by a Cincinnati Children's Research Foundation Trustee Award, a Simons Foundation Autism Research Initiative (SFARI) Bridge to Independence Award (663007), a SFARI Supplement to

Enhance Equity and Diversity (SEED) Award, and a Gilbert Family Foundation Neurofibromatosis

Gene Therapy Initiative Team Science Award to JER.

Competing interests: The authors have no competing interests to declare.

REFERENCES:

Aboitiz F, Ossandón T, Zamorano F, Palma B, Carrasco X. 2014. Irrelevant stimulus processing in ADHD: catecholamine dynamics and attentional networks. *Front Psychol* **5**:183. doi:10.3389/fpsyg.2014.00183

Alcantara IC, Tapia APM, Aponte Y, Krashes MJ. 2022. Acts of appetite: neural circuits governing the appetitive, consummatory, and terminating phases of feeding. *Nat Metab* **4**:836–847. doi:10.1038/s42255-022-00611-y

Anastasaki C, Mo J, Chen J-K, Chatterjee J, Pan Y, Scheaffer SM, Cobb O, Monje M, Le LQ, Gutmann DH. 2022. Neuronal hyperexcitability drives central and peripheral nervous system tumor progression in models of neurofibromatosis-1. *Nat Commun* **13**:2785. doi:10.1038/s41467-022-30466-6

Anastasaki C, Woo AS, Messiaen LM, Gutmann DH. 2015. Elucidating the impact of neurofibromatosis-1 germline mutations on neurofibromin function and dopamine-based learning. *Hum Mol Genet* **24**:3518–3528. doi:10.1093/hmg/ddv103

Ball W, Tronick E. 1971. Infant responses to impending collision: optical and real. *Science* **171**:818–820. doi:10.1126/science.171.3973.818

Barker DJ, Miranda-Barrientos J, Zhang S, Root DH, Wang H-L, Liu B, Calipari ES, Morales M. 2017. Lateral Preoptic Control of the Lateral Habenula through Convergent Glutamate and GABA Transmission. *Cell Rep* **21**:1757–1769. doi:10.1016/j.celrep.2017.10.066

Beier KT, Steinberg EE, DeLoach KE, Xie S, Miyamichi K, Schwarz L, Gao XJ, Kremer EJ, Malenka RC, Luo L. 2015. Circuit Architecture of VTA Dopamine Neurons Revealed by Systematic Input-Output Mapping. *Cell* **162**:622–634. doi:10.1016/j.cell.2015.07.015

Berke JD. 2018. What does dopamine mean? *Nat Neurosci* **21**:787–793. doi:10.1038/s41593-018-0152-y

Berridge KC, Robinson TE. 2016. Liking, wanting, and the incentive-sensitization theory of addiction. *Am Psychol* **71**:670–679. doi:10.1037/amp0000059

Berson DM, Dunn FA, Takao M. 2002. Phototransduction by retinal ganglion cells that set the circadian clock. *Science* **295**:1070–1073. doi:10.1126/science.1067262

Branco T, Redgrave P. 2020. The Neural Basis of Escape Behavior in Vertebrates. *Annu Rev Neurosci* **43**:417–439. doi:10.1146/annurev-neuro-100219-122527

Bromberg-Martin ES, Matsumoto M, Hikosaka O. 2010a. Distinct tonic and phasic anticipatory activity in lateral habenula and dopamine neurons. *Neuron* **67**:144–155. doi:10.1016/j.neuron.2010.06.016

Bromberg-Martin ES, Matsumoto M, Hikosaka O. 2010b. Dopamine in motivational control: rewarding, aversive, and alerting. *Neuron* **68**:815–834. doi:10.1016/j.neuron.2010.11.022

Brown JA, Emnett RJ, White CR, Yuede CM, Conyers SB, O'Malley KL, Wozniak DF, Gutmann DH. 2010. Reduced striatal dopamine underlies the attention system dysfunction in neurofibromatosis-1 mutant mice. *Hum Mol Genet* **19**:4515–4528. doi:10.1093/hmg/ddq382

Comoli E, Coizet V, Boyes J, Bolam JP, Canteras NS, Quirk RH, Overton PG, Redgrave P. 2003. A direct projection from superior colliculus to substantia nigra for detecting salient visual events. *Nat Neurosci* **6**:974–980. doi:10.1038/nn1113

Danskin B, Denman D, Valley M, Ollerenshaw D, Williams D, Groblewski P, Reid C, Olsen S, Blanche T, Waters J. 2015. Optogenetics in Mice Performing a Visual Discrimination Task: Measurement and Suppression of Retinal Activation and the Resulting Behavioral Artifact. *PLoS One* **10**:e0144760. doi:10.1371/journal.pone.0144760

de Jong JW, Afjei SA, Dorocic IP, Peck JR, Liu C, Kim CK, Tian L, Deisseroth K, Lammel S. 2018. A Neural Circuit Mechanism for Encoding Aversive Stimuli in the Mesolimbic Dopamine System. *Neuron* **0**. doi:10.1016/j.neuron.2018.11.005

Dhande OS, Huberman AD. 2014. Retinal ganglion cell maps in the brain: implications for visual processing. *Curr Opin Neurobiol* **24**:133–142. doi:10.1016/j.conb.2013.08.006

Diggs-Andrews KA, Tokuda K, Izumi Y, Zorumski CF, Wozniak DF, Gutmann DH. 2013. Dopamine deficiency underlies learning deficits in neurofibromatosis-1 mice. *Ann Neurol* **73**:309–315. doi:10.1002/ana.23793

Dommett E, Coizet V, Blaha CD, Martindale J, Lefebvre V, Walton N, Mayhew JEW, Overton PG, Redgrave P. 2005. How visual stimuli activate dopaminergic neurons at short latency. *Science* **307**:1476–1479. doi:10.1126/science.1107026

Eban-Rothschild A, Rothschild G, Giardino WJ, Jones JR, de Lecea L. 2016. VTA dopaminergic neurons regulate ethologically relevant sleep-wake behaviors. *Nat Neurosci* **19**:1356–1366. doi:10.1038/nn.4377

Evans DA, Stempel AV, Vale R, Ruehle S, Lefler Y, Branco T. 2018. A synaptic threshold mechanism for computing escape decisions. *Nature* **558**:590–594. doi:10.1038/s41586-018-0244-6

Everitt BJ, Robbins TW. 2005. Neural systems of reinforcement for drug addiction: from actions to habits to compulsion. *Nat Neurosci* **8**:1481–1489. doi:10.1038/nn1579

Fenno L, Yizhar O, Deisseroth K. 2011. The development and application of optogenetics. *Annu Rev Neurosci* **34**:389–412. doi:10.1146/annurev-neuro-061010-113817

Fernandez DC, Fogerson PM, Lazzerini Ospri L, Thomsen MB, Layne RM, Severin D, Zhan J, Singer JH, Kirkwood A, Zhao H, Berson DM, Hattar S. 2018. Light Affects Mood and Learning through Distinct Retina-Brain Pathways. *Cell* **175**:71-84.e18. doi:10.1016/j.cell.2018.08.004

Hattar S, Lucas RJ, Mrosovsky N, Thompson S, Douglas RH, Hankins MW, Lem J, Biel M, Hofmann F, Foster RG, Yau K-W. 2003. Melanopsin and rod-cone photoreceptive systems account for all major accessory visual functions in mice. *Nature* **424**:76–81. doi:10.1038/nature01761

Horvitz JC. 2000. Mesolimbocortical and nigrostriatal dopamine responses to salient non-reward events. *Neuroscience* **96**:651–656.

Horvitz JC, Stewart T, Jacobs BL. 1997. Burst activity of ventral tegmental dopamine neurons is elicited by sensory stimuli in the awake cat. *Brain Res* **759**:251–258. doi:10.1016/s0006-8993(97)00265-5

Huang M, Li D, Cheng X, Pei Q, Xie Z, Gu H, Zhang X, Chen Z, Liu A, Wang Y, Sun F, Li Y, Zhang J, He M, Xie Y, Zhang F, Qi X, Shang C, Cao P. 2021. The tectonigral pathway regulates appetitive locomotion in predatory hunting in mice. *Nat Commun* **12**:4409. doi:10.1038/s41467-021-24696-3

Jeon CJ, Strettoi E, Masland RH. 1998. The major cell populations of the mouse retina. *J Neurosci* **18**:8936–8946.

Korshunov KS, Blakemore LJ, Trombley PQ. 2017. Dopamine: A Modulator of Circadian Rhythms in the Central Nervous System. *Front Cell Neurosci* **11**:91. doi:10.3389/fncel.2017.00091

Kutlu MG, Zachry JE, Melugin PR, Cajigas SA, Chevee MF, Kelly SJ, Kutlu B, Tian L, Siciliano CA, Calipari ES. 2021. Dopamine release in the nucleus accumbens core signals perceived saliency. *Curr Biol* **31**:4748-4761.e8. doi:10.1016/j.cub.2021.08.052

Lall GS, Revell VL, Momiji H, Al Enezi J, Altimus CM, Güler AD, Aguilar C, Cameron MA, Allender S, Hankins MW, Lucas RJ. 2010. Distinct contributions of rod, cone, and melanopsin photoreceptors to encoding irradiance. *Neuron* **66**:417–428. doi:10.1016/j.neuron.2010.04.037

Li D, Sham PC, Owen MJ, He L. 2006. Meta-analysis shows significant association between dopamine system genes and attention deficit hyperactivity disorder (ADHD). *Hum Mol Genet* **15**:2276–2284. doi:10.1093/hmg/ddl152

Lisman JE, Grace AA. 2005. The hippocampal-VTA loop: controlling the entry of information into long-term memory. *Neuron* **46**:703–713. doi:10.1016/j.neuron.2005.05.002

Lopes G, Bonacchi N, Frazão J, Neto JP, Atallah BV, Soares S, Moreira L, Matias S, Itskov PM, Correia PA, Medina RE, Calcaterra L, Dreosti E, Paton JJ, Kampff AR. 2015. Bonsai: an event-based

framework for processing and controlling data streams. *Front Neuroinform* **9**:7. doi:10.3389/fninf.2015.00007

Lucas RJ, Douglas RH, Foster RG. 2001. Characterization of an ocular photopigment capable of driving pupillary constriction in mice. *Nat Neurosci* **4**:621–626. doi:10.1038/88443

Maia TV, Frank MJ. 2017. An Integrative Perspective on the Role of Dopamine in Schizophrenia. *Biol Psychiatry* **81**:52–66. doi:10.1016/j.biopsych.2016.05.021

Maier JX, Neuhoff JG, Logothetis NK, Ghazanfar AA. 2004. Multisensory integration of looming signals by rhesus monkeys. *Neuron* **43**:177–181. doi:10.1016/j.neuron.2004.06.027

Mautner V-F, Granström S, Leark RA. 2015. Impact of ADHD in adults with neurofibromatosis type 1: associated psychological and social problems. *J Atten Disord* **19**:35–43. doi:10.1177/1087054712450749

Miguel CS, Chaim-Avancini TM, Silva MA, Louzã MR. 2015. Neurofibromatosis type 1 and attention deficit hyperactivity disorder: a case study and literature review. *Neuropsychiatr Dis Treat* **11**:815–821. doi:10.2147/NDT.S75038

Nakagawa H, Hongjian K. 2010. Collision-sensitive neurons in the optic tectum of the bullfrog, Rana catesbeiana. *J Neurophysiol* **104**:2487–2499. doi:10.1152/jn.01055.2009

Nikbakht N, Diamond ME. 2021. Conserved visual capacity of rats under red light. *Elife* **10**:e66429. doi:10.7554/eLife.66429

Panda S, Provencio I, Tu DC, Pires SS, Rollag MD, Castrucci AM, Pletcher MT, Sato TK, Wiltshire T, Andahazy M, Kay SA, Van Gelder RN, Hogenesch JB. 2003. Melanopsin is required for non-image-forming photic responses in blind mice. *Science* **301**:525–527. doi:10.1126/science.1086179

Patriarchi T, Cho JR, Merten K, Marley A, Broussard GJ, Liang R, Williams J, Nimmerjahn A, von Zastrow M, Gradinaru V, Tian L. 2019. Imaging neuromodulators with high spatiotemporal resolution using genetically encoded indicators. *Nat Protoc* **14**:3471–3505. doi:10.1038/s41596-019-0239-2

Peirson SN, Brown LA, Pothecary CA, Benson LA, Fisk AS. 2018. Light and the laboratory mouse. *J Neurosci Methods* **300**:26–36. doi:10.1016/j.jneumeth.2017.04.007

Poulin J-F, Caronia G, Hofer C, Cui Q, Helm B, Ramakrishnan C, Chan CS, Dombeck DA, Deisseroth K, Awatramani R. 2018. Mapping projections of molecularly defined dopamine neuron subtypes using intersectional genetic approaches. *Nat Neurosci* **21**:1260–1271. doi:10.1038/s41593-018-0203-4

Purper-Ouakil D, Ramoz N, Lepagnol-Bestel A-M, Gorwood P, Simonneau M. 2011. Neurobiology of attention deficit/hyperactivity disorder. *Pediatr Res* **69**:69R-76R. doi:10.1203/PDR.0b013e318212b40f

Redgrave P, Gurney K. 2006. The short-latency dopamine signal: a role in discovering novel actions? *Nat Rev Neurosci* **7**:967–975. doi:10.1038/nrn2022

Resendez SL, Stuber GD. 2015. In vivo calcium imaging to illuminate neurocircuit activity dynamics underlying naturalistic behavior. *Neuropsychopharmacology* **40**:238–239. doi:10.1038/npp.2014.206

Robinson JE, Coughlin GM, Hori AM, Cho JR, Mackey ED, Turan Z, Patriarchi T, Tian L, Gradinaru V. 2019. Optical dopamine monitoring with dLight1 reveals mesolimbic phenotypes in a mouse model of neurofibromatosis type 1. *Elife* **8**. doi:10.7554/eLife.48983

Robinson JE, Gradinaru V. 2018. Dopaminergic dysfunction in neurodevelopmental disorders: recent advances and synergistic technologies to aid basic research. *Curr Opin Neurobiol* **48**:17–29. doi:10.1016/j.conb.2017.08.003

Sabatini BL, Tian L. 2020. Imaging Neurotransmitter and Neuromodulator Dynamics In Vivo with Genetically Encoded Indicators. *Neuron* **108**:17–32. doi:10.1016/j.neuron.2020.09.036

Schultz W. 2010. Dopamine signals for reward value and risk: basic and recent data. *Behav Brain Funct* **6**:24. doi:10.1186/1744-9081-6-24

Schultz W. 1998. Predictive reward signal of dopamine neurons. *J Neurophysiol* **80**:1–27. doi:10.1152/jn.1998.80.1.1

Schultz W, Carelli RM, Wightman RM. 2015. Phasic dopamine signals: from subjective reward value to formal economic utility. *Curr Opin Behav Sci* **5**:147–154. doi:10.1016/j.cobeha.2015.09.006

Schultz W, Romo R. 1990. Dopamine neurons of the monkey midbrain: contingencies of responses to stimuli eliciting immediate behavioral reactions. *J Neurophysiol* **63**:607–624. doi:10.1152/jn.1990.63.3.607

Singer AC, Martorell AJ, Douglas JM, Abdurrob F, Attokaren MK, Tipton J, Mathys H, Adaikkan C, Tsai L-H. 2018. Noninvasive 40-Hz light flicker to recruit microglia and reduce amyloid beta load. *Nat Protoc* **13**:1850–1868. doi:10.1038/s41596-018-0021-x

Solié C, Contestabile A, Espinosa P, Musardo S, Bariselli S, Huber C, Carleton A, Bellone C. 2022. Superior Colliculus to VTA pathway controls orienting response and influences social interaction in mice. *Nat Commun* **13**:817. doi:10.1038/s41467-022-28512-4

Sun H, Frost BJ. 1998. Computation of different optical variables of looming objects in pigeon nucleus rotundus neurons. *Nat Neurosci* **1**:296–303. doi:10.1038/1110

Takakuwa N, Kato R, Redgrave P, Isa T. 2017. Emergence of visually-evoked reward expectation signals in dopamine neurons via the superior colliculus in V1 lesioned monkeys. *Elife* **6**. doi:10.7554/eLife.24459

Tegelbeckers J, Bunzeck N, Duzel E, Bonath B, Flechtner H, Krauel K. 2015. Altered salience processing in attention deficit hyperactivity disorder. *Hum Brain Mapp* **36**:2049–2060. doi:10.1002/hbm.22755

Temizer I, Donovan JC, Baier H, Semmelhack JL. 2015. A Visual Pathway for Looming-Evoked Escape in Larval Zebrafish. *Curr Biol* **25**:1823–1834. doi:10.1016/j.cub.2015.06.002

Thompson S, Recober A, Vogel TW, Kuburas A, Owens JA, Sheffield VC, Russo AF, Stone EM. 2010. Light aversion in mice depends on nonimage-forming irradiance detection. *Behav Neurosci* **124**:821–827. doi:10.1037/a0021568

Umino Y, Solessio E, Barlow RB. 2008. Speed, spatial, and temporal tuning of rod and cone vision in mouse. *J Neurosci* **28**:189–198. doi:10.1523/JNEUROSCI.3551-07.2008

Watabe-Uchida M, Eshel N, Uchida N. 2017. Neural Circuitry of Reward Prediction Error. *Annu Rev Neurosci* **40**:373–394. doi:10.1146/annurev-neuro-072116-031109

Watabe-Uchida M, Zhu L, Ogawa SK, Vamanrao A, Uchida N. 2012. Whole-brain mapping of direct inputs to midbrain dopamine neurons. *Neuron* **74**:858–873. doi:10.1016/j.neuron.2012.03.017

Wise RA. 2004. Dopamine, learning and motivation. *Nat Rev Neurosci* **5**:483–494. doi:10.1038/nrn1406

Yilmaz M, Meister M. 2013. Rapid innate defensive responses of mice to looming visual stimuli. *Curr Biol* **23**. doi:10.1016/j.cub.2013.08.015

Yuan L, Dou Y-N, Sun Y-G. 2019. Topography of Reward and Aversion Encoding in the Mesolimbic Dopaminergic System. *J Neurosci* **39**:6472–6481. doi:10.1523/JNEUROSCI.0271-19.2019

Zhang Z, Beier C, Weil T, Hattar S. 2021. The retinal ipRGC-preoptic circuit mediates the acute effect of light on sleep. *Nat Commun* **12**:5115. doi:10.1038/s41467-021-25378-w

1   **VIDEO TITLES:**

2   **Video 1. Behavioral response to presentation of black looming discs on a light**
3   **background when mice entered the threat zone of a rectangular arena.**
4
5   **Video 2. Behavioral response to illumination of a spotlight when mice entered the target**
6   **zone of a rectangular arena.**
7

FIGURES AND CAPTIONS

Figure 1. LNAc dopaminergic encoding of visual threats.

A. Fluorescent dopamine signals were recorded with dLight1 and fiber photometry in the nucleus

accumbens lateral shell (LNAc) during presentation of looming and control discs. **B.** Average

dLight1 response to trains of five black or contrast inverted discs ± standard error of the mean

(SEM). **C.** The dLight1 response to black or inverted discs was dependent on disc

color/background, disc type (static vs. looming vs. receding), and stimulus number (n = 11; 3-way

repeated measures ANOVA; $F_{8,240}$ = 2.02, $p_{disc\ background\ x\ disc\ type\ x\ stimulus\ number}$ = 0.045; $F_{1,240}$ = 143.92,

$p_{background}$ < 0.001; $F_{2,240}$ = 150.32, $p_{disc\ type}$ < 0.001; $F_{4,240}$ = 192.64, $p_{stimulus\ number}$ < 0.001). Bonferroni

post hoc tests revealed that contrast inverted discs evoked more dopamine than black discs.

Contrast inverted looming discs evoked less dopamine than inverted static and receding discs

after the first presentation. **D.** dLight1 transient peak latency was dependent on disc

color/background and disc type (n = 11; 2-way repeated measures ANOVA; $F_{2,20}$ = 64.78, p_{disc}

$_{\text{background x disc type}} < 0.001$; $F_{1,20} = 25.69$, $p_{\text{background}} < 0.001$; $F_{2,20} = 7.58$, $p_{\text{disc type}} = 0.01$). Bonferroni

post hoc tests showed that contrast inverted static and receding discs evoked transients with

shorter latency compared to black discs. Additionally, transients evoked by contrast inverted

receding discs had shorter latency than contrast inverted looming discs. **E.** Escape velocity

following overhead disc presentation was dependent on disc color/background and disc type (n =

12; 2-way repeated measures ANOVA; $F_{2,22} = 49.28$, $p_{\text{disc background x disc type}} < 0.001$; $F_{1,22} = 18.38$,

$p_{\text{background}} = 0.001$; $F_{2,22} = 28.89$, $p_{\text{disc type}} < 0.001$). Bonferroni *post hoc* tests showed that black

looming discs induced greater escape velocity than all other overhead discs. For panels C and D,

* indicates p < 0.05 vs. black disc of the same type (e.g. black static disc vs. contrast inverted

static disc); + indicates p < 0.05 vs looming disc of the same color (e.g. black looming vs. black

receding disc). For panel E, * indicates p < 0.05 vs. other overhead discs.

Figure 2. Dopaminergic responses to rapid dark to light transitions.

A. Instantaneous LCD screen transitions from dark to light evoked rapid dopamine release at stimulus onset when compared to the pre-stimulus baseline (inset: baseline and stimulus-induced dLight1 peak values for individual mice; n = 11; paired t-test; t_{10} = 7.01, p < 0.001). **B.** dLight1 responses to the onset of LCD screen dark to light transitions at different transition lengths (0.05 – 2.0 s) ± SEM. **C.** The magnitude (*pink*) and latency (*teal*) of dopaminergic responses to dark-to-light transitions varied non-linearly depending on transition speed (peak amplitude: one-phase exponential decay, y_0 = 8.84 z-score, plateau = 1.36 z-score, tau = 0.43 s, R^2 = 0.90; peak latency: one-phase exponential association, y_0 = 0.43 ms, plateau = 0.59 ms, tau = 0.54 s, R^2 = 0.98). **D.** dLight1 responses to the onset ten-second white LED stimuli across a range of irradiances (0 $\mu W/cm^2$ – 0.01 $\mu W/cm^2$) ± SEM. **E.** The magnitude of the dopaminergic response to 10s white LED stimuli was dependent on the stimulus irradiance (n = 7; 1-way repeated measures ANOVA; $F_{7,42}$ = 38.79, p < 0.001). Data shown with a one-phase exponential association fit (y_0 = 1.20 z-

score, plateau = 0.84 z-score, tau = 0.00074 μW/cm^2, R^2 = 0.63). **F.** The latency of the

dopaminergic response to 10 s white LED stimuli was dependent on the stimulus irradiance (n =

7; 1-way repeated measures ANOVA; $F_{6,36}$ = 47.35, p < 0.001). Data shown with a one-phase

exponential decay fit (y_0 = 0.42 ms, plateau = 0.35 ms, tau = 0.0079 μW/cm^2, R^2 = 0.76). **G.** The

dopaminergic response to 5.0 μW/cm^2 white light was not different (*right*) if measured at the

beginning of the vivarium dark (*left*) or light (*center*) phase of the day-night cycle (n = 11; paired

t-test; t_{10} = 1.27, p = 0.23). **G.** In all panels, * indicates p < 0.05.

Figure 3. Dopaminergic responses to repeated light stimuli.

A. *(Left)* Dopamine release evoked by 20 one-second white LED stimuli was reduced with

repeated exposures and was dependent on the interstimulus interval (ISI; 10 ms – 10 s; n = 9; 2-

way repeated measures ANOVA; $F_{57,456}$ = 9.54, $p_{stimulus\ number\ x\ interstimulus\ interval}$ < 0.001; $F_{19,456}$ = 72.98,

$p_{stimulus\ number}$ < 0.001; $F_{3,456}$ = 63.91, $p_{interstimulus\ interval}$ < 0.001). *(Right)* Averaged dLight1 fluorescent

traces showing the dopaminergic response to 20 one-second white LED light pulses with a 100

ms ISI (*purple*) or 10 s ISI (*orange*). **B.** Total habituation of the peak dLight1 response to repeated

stimuli (shown as the peak response to the 20[th] stimulus as a percentage of the 1[st] stimulus) is

dependent on the duration of the interstimulus interval (n = 9; 1-way repeated measures ANOVA;

$F_{3,24}$ = 104.0, p < 0.001). Data shown with a semi-log fit (y-intercept: 48.92%, slope: 22.58 %s^{-1},

R^2 = 0.86). **C.** *(Left)* Averaged dLight1 trace showing LNAc dopamine evoked by a 60 second

presentation of 40 Hz white LED flicker (*inset:* response during the first second after stimulus

onset) ± SEM. *(Right)* 40 Hz flicker only evoked significant dopamine release at stimulus onset (n

= 10; 1-way repeated measures ANOVA; $F_{2,18}$ = 100.4, p < 0.001). Bonferroni *post hoc* tests

confirmed that the dLight1 peak at LED onset was greater than baseline and offset responses, which did not differ from each other ($p = 0.09$). **D.** (*Left*) No habituation in the peak dLight1 response to repeated 1 s white LED stimuli was observed when the ISI was sufficiently long (100 s) – both before and after presentation of 300 one-second LED stimuli with a one-second ISI ($n = 7$; 2-way repeated measures ANOVA; $F_{4,24} = 0.73$, $p_{\text{stimulus number x exposure history}} = 0.52$; $F_{4,24} = 1.01$, $p_{\text{stimulus number}} = 0.40$; $F_{1,24} = 4.61$, $p_{\text{exposure history}} = 0.08$). *(Right)* Averaged dLight1 fluorescent traces showing the dopaminergic response to 5 one-second white LED light pulses with a 100 s ISI before (*black*) or after (*pink*) 300 one-second LED stimuli. **E.** Averaged dLight1 fluorescent traces showing the dopaminergic response to a 1 s white LED stimulus one second after a 1 s (*left*) or 300 s (*right*) preconditioning stimulus ± SEM. **F.** The dLight1 response to a 1 s white LED test stimulus was not dependent on the length of the preconditioning stimulus ($n = 11$; 2-way repeated measures ANOVA; $F_{1,10} = 0.27$, $p_{\text{initial stimulus length x stimulus Number}} = 0.61$; $F_{1,10} = 3.83$, $p_{\text{initial stimulus length}} = 0.08$; $F_{1,10} = 55.10$, $p_{\text{stimulus number}} < 0.001$). Bonferroni *post hoc* tests revealed that the dLight1 response to the test stimulus onset was significantly smaller than the response to the onset of the preconditioning stimulus, regardless of its duration. There was no difference between the dLight1 response to the onset of the preconditioning ($p = 0.11$) or test stimulus ($p = 0.40$) between experiments. **G.** The dLight1 response to light offset was larger for a 300 s light stimulus compared to a 1 s light stimulus ($n = 11$; paired t-test; $t_{10} = 4.91$, $p < 0.001$). In all panels, * indicates $p < 0.05$.

Figure 4. Dopaminergic responses to individual wavelengths across the visual spectrum.

A. The dopaminergic response to UV (360 nm), blue (475 nm), green (555 nm), and red (635 nm)

LED light was wavelength and irradiance-dependent (n = 8; 2-way repeated measures ANOVA;

$F_{12,84}$ = 9.63, $p_{wavelength \times irradiance}$ < 0.001; $F_{3,84}$ = 37.59, $p_{wavelength}$ < 0.001; $F_{4,84}$ = 10.08, $p_{irradiance}$ =

0.004). Bonferroni *post hoc* tests revealed that dopamine evoked by UV and red light was smaller

than blue and green wavelengths at the lowest irradiance tested (0.001 μW/cm^2). The dLight1

response to the red LED was also significantly lower than blue and green LEDs at irradiances

0.01 μW/cm^2 and 0.1 μW/cm^2. For comprehensive reporting of all significant *post hoc* tests across

irradiances and wavelengths, see the provided Supplemental Data and Statistical Analysis file.

B. Averaged dLight1 trace showing LNAc dopamine evoked by either 0.001 μW/cm^2 (1 nW/cm^2)

or 10 μW/cm^2 UV, blue, green, or red LEDs ± SEM. **C.** *(Left)* 100 μW/cm^2 far-red (730 nm) LED

light evoked significant dopamine release compared to 10 μW/cm^2 light (n = 8; paired t-test; t_7 =

2.54, p = 0.04). *(Right)* Averaged dLight1 trace showing LNAc dopamine evoked by a 100 μW/cm^2

far-red LED ± SEM. In all panels, * indicates p < 0.05.

Figure 2-Figure Supplement 1.

A. The magnitude of the dopaminergic response to dark-to-light transitions was dependent on the

stimulus transition time (n = 11; 1-way repeated measures ANOVA; $F_{7,70}$ = 17.76, p < 0.001). **B.**

The latency of the peak dopaminergic response to dark-to-light transitions was dependent on the

stimulus transition time (n = 11; 1-way repeated measures ANOVA; $F_{7,70}$ = 11.03, p = 0.007). In

A and B, pink circles (mean ± SEM) and gray dots (individual values) represent data derived from

fluorescent dLight1 traces averaged across trials for each individual mice. Shown for comparison,

black circles represent data derived from fluorescence traces averaged across all mice prior to

peak detection, which yielded a single value for each transition time. **C.** The dopaminergic

response to 80 dB tones was dependent on their frequency (n = 10; 1-way repeated measures

ANOVA; $F_{5,45}$ = 10.11, p < 0.001) and was less than the response to a 5 μW/cm^2 LED light stimulus

(Mixed-effects ANOVA; F = 26.62, $p_{stimulus}$ < 0.001). Bonferroni *post hoc* tests revealed that the

dLight1 response to the light stimulus was significantly larger than all tone responses. **D.** (*Left*)

Opening the enclosure peephole evoked significant dopamine release at stimulus onset when

compared to the pre-stimulus baseline (n = 6; paired t-test; t_5 = 5.72, p = 0.002). (*Right*) Averaged

dLight1 trace showing LNAc dopamine evoked by opening the peephole on the door of the

behavioral testing chamber enclosure to observe mouse behavior ± SEM. The irradiance

associated with this manipulation was 17 nW/cm^2 measured at mouse level. In all panels, *

indicates p < 0.05.