Genetic drift promotes and recombination hinders speciation on a holey fitness landscape

Ata Kalirad[1,2], Christina L. Burch[3], Ricardo B. R. Azevedo[1*],

1 Department of Biology and Biochemistry, University of Houston, Houston, Texas, USA
2 Department for Integrative Evolutionary Biology, Max Planck Institute for Biology Tübingen, Tübingen, Germany
3 Department of Biology, University of North Carolina, Chapel Hill, North Carolina, USA

* razevedo@uh.edu

Abstract

Dobzhansky and Muller proposed a general mechanism through which microevolution, the substitution of alleles within populations, can cause the evolution of reproductive isolation between populations and, therefore, macroevolution. As allopatric populations diverge, many combinations of alleles differing between them have not been tested by natural selection and may thus be incompatible. Such genetic incompatibilities often cause low fitness in hybrids between species. Furthermore, the number of incompatibilities grows with the genetic distance between diverging populations. However, what determines the rate and pattern of accumulation of incompatibilities remains unclear. We investigate this question by simulating evolution on a holey fitness landscape. We find that genetic incompatibilities accumulate more slowly among genetically robust populations and identify two determinants of the accumulation rate: recombination rate and population size. In large populations with abundant genetic variation, recombination selects for increased genetic robustness and, consequently, incompatibilities accumulate more slowly. In small populations, genetic drift interferes with this process and promotes the accumulation of genetic incompatibilities. Our

results suggest a novel mechanism by which genetic drift promotes and recombination hinders speciation.

Introduction

The rates of origination of new species vary extensively across the tree of life. For example, some lineages of cichlids are estimated to produce new species approximately $70\times$ faster than some lineages of hawks [1]. A major challenge for evolutionary biology is to explain this macroevolutionary pattern through the action of microevolutionary processes operating within populations [2,3]. Stanley [4] speculated that speciation is "a largely random process" and that, therefore, "macroevolution is decoupled from microevolution". Here we explore the alternative: that macroevolution and microevolution are, in fact, mechanistically coupled.

Speciation results from the build up of reproductive isolation (RI) between populations. Thus, variation in speciation rates should be caused by variation in the rates at which RI increases between populations as they diverge. As allopatric populations diverge, genetic incompatibilities accumulate between them and generate postzygotic RI [5,6], one of the causes of speciation. Orr [7] developed a combinatorial model to describe this process. He proposed that any pair of alleles at two divergent loci, not previously tested by natural selection, has a probability p of being incompatible. If two populations evolving in allopatry have diverged at k loci, then $k(k-1)/2$ combinations of alleles at pairs of these loci will not have been tested by natural selection (Fig 1a). Combining these two terms, the number of Dobzhansky–Muller incompatibilities (DMIs) between pairs of loci is expected to increase with k according to:

$$I(k) = \frac{pk(k-1)}{2} \approx \frac{pk^2}{2} \tag{1}$$

Orr [7] termed this pattern of faster than linear accumulation "snowballing". Empirical tests of this prediction confirmed that genetic incompatibilities accumulated faster than linearly for inviability in species of *Drosophila* [8,9], male sterility in subspecies of *Mus musculus* [10], and both seed and pollen sterility in species of *Solanum* [11,12].

Computational studies using models of RNA folding [13], transcription factor-DNA binding [14], and spatial gene regulation during development [15] also found that genetic incompatibilities snowballed. These empirical and computational studies were all conducted over relatively short evolutionary distances. Surprisingly, a study conducted over much longer genetic distances discovered that genetic incompatibilities did not snowball for animal proteins [16]; rather, they accumulated approximately linearly with genetic distance. F. Kondrashov et al. [16] suggested that a linear accumulation of genetic incompatibilities is not well explained by Orr's combinatorial model. We note, however, that the snowballing prediction in Eq 1 relies on certain assumptions [7, 13, 17], in particular, that the probability of incompatibility p does not, itself, evolve as populations diverge [13]. Relaxing this assumption could explain the linear accumulation pattern. For example, if p declines with k according to the relationship $p(k) = C/k$ (where C is a constant) then DMIs will accumulate linearly: $I(k) = C(k - 1)$. The extent to which evolved differences in p account for the linear accumulation of genetic incompatibilities among animal proteins is unknown. Nonetheless, the possibility raises the question of what might cause p, a determinant of the rate and pattern of accumulation of genetic incompatibilities, to evolve.

Here we use a computational model of RNA secondary structure to investigate whether and why populations differ in the rate at which they accumulate genetic incompatibilities. We chose the folding of RNA sequences as a simple model of a genotype-phenotype map grounded in biophysical reality [18]. Crucially, this genotype-phenotype map naturally incorporates epistasis and has been used to study its evolutionary consequences (e.g., robustness [19, 20], evolvability [21, 22], molecular evolution [23]). As in Kalirad and Azevedo [13], we conduct evolution experiments that monitor the evolution of RNA secondary structures on a holey fitness landscape [24] (also known as a neutral network [25, 26]), in which genotypes are either viable or inviable (Fig 1c). Inviable genotypes carry combinations of alleles displaying DMIs (Fig 1b). Thus, holey fitness landscapes provide the simplest modeling framework to study the evolution of reproductive isolation caused by genetic incompatibilities.

We follow the approach taken in the classic empirical tests of the combinatorial model [8, 11] and monitor the accumulation of incompatible introgressions, IIs, rather than DMIs [13]. Because introgression is a perturbation of the genotype, we predict that

Fig 1. Populations evolving on a holey fitness landscape accumulate genetic incompatibilities. **(a)** Two populations diverge in allopatry. Both populations are initially fixed for lowercase alleles (open circles) at five loci, A–E. Derived alleles are indicated by uppercase letters (closed circles). The first two substitutions (closed circles) were of alleles *A* and *B* in population 1 (left, green); the next three substitutions were of alleles *C*, *D*, and *E* in population 2 (right, blue). **(b)** Genetic incompatibilities can be inferred by introgressing individual divergent alleles from one population into the other and counting the number of incompatible introgressions (IIs). This is illustrated by the introgression of the divergent alleles from population 1 (*ABcde*) into population 2 (*abCDE*) to create the four combinations that have not been tested by natural selection. We ignore the introgression of the *e* allele because it results in the transitional *abCDe* genotype in population 2, which is known to be viable (red arrow in (a)). Two alleles from population 1 (*A* and *c*) are deleterious in the genetic background of population 2 (grey boxes). We refer to these as IIs. Each II implies the existence of one or more DMIs. **(c)** Holey fitness landscape of the 32 genotypes at loci A–E. Nodes (circles) represent genotypes. Lines connect genotypes that can be reached by a mutation at a single locus. Each genotype has 5 mutational neighbors. 7 of the 32 genotypes (22%) are inviable, shown in light grey. The genotypes generated by the two IIs shown in (b) are marked by arrows. The remaining 25 genotypes (78%) are viable, shown in colors from yellow to blue, and define a neutral network. The different colors indicate the proportion ν of the mutational neighbors of a genotype that are viable, a measure of mutational robustness. The evolutionary trajectories of the populations shown in (a) are indicated by thick blue and green lines.

the probability that an introgression is incompatible will be inversely related to the recipient's genetic robustness, its ability to maintain viability in the face of perturbations to the genotype. We test this prediction by experimentally manipulating population genetic parameters known to promote the evolution of genetic robustness.

Materials and methods

Experimental design

Our experimental design capitalises on prior findings regarding how populations evolve on a holey fitness landscape (Figs 1a and 1c). On a holey fitness landscape, all viable genotypes have the same viability but, in the presence of mutation and recombination, they do not all have the same fitness. Mutations are either lethal or neutral with respect to viability, so viable individuals can differ in their expected number of viable offspring depending on the proportion ν of their mutational neighbors that are viable, a measure of mutational robustness (Fig 1c). Similarly, recombination between different pairs of genotypes may produce varying proportions of viable recombinants. Thus, populations evolving on a holey fitness landscape experience selection for higher genetic robustness [19, 27–29]. The ability of populations to respond to this selection depends on factors that determine the relative strengths of selection on genetic robustness and genetic drift: mutation rate, recombination rate, and population size (Fig 2; U, r, $N \to \nu$).

To test our prediction that the rate at which IIs accumulate between populations will be inversely related to their genetic robustness, we first experimentally evolved populations of RNA secondary structures using combinations of population sizes and recombination rates expected to result in different equilibrium levels of genetic robustness. Populations evolved under an individual-based Wright-Fisher model with a selection-recombination-mutation life cycle, with a constant population size (between $N = 10$ and $N = 500$) and recombination rate (between no recombination and free recombination). Our use of the Wright-Fisher model differs from most theoretical research into the combinatorial model [7, 13, 30, 31] in that it does not impose a strong selection, weak mutation (SSWM) regime [32]. Unlike SSWM, in which mutations arise

Fig 2. Evolutionary causes of the accumulation of genetic incompatibilities between populations in our model. Effects are summarized for asexual **(a)** and sexual **(b)** populations. Ellipses contain the following properties of populations: U, genomic mutation rate; N, population size; r, recombination probability or rate; π, nucleotide diversity; ν, mutational robustness; $\mathbf{\Pi}_W$ and $\mathbf{\Pi}_B$, genetic incompatibilities within and between populations, respectively. Light gray ellipses in **(a)** indicate that in asexual populations recombination does not operate ($r = 0$). Red lines ending in arrows indicate positive effects; blue lines ending in bars indicate negative effects. The thickness of the line indicates the strength of the effect. Light gray lines in a indicate that the effect does not operate.

and transit to fixation or loss sequentially, our implementation of the Wright-Fisher model uses a genome-wide mutation rate of $U = 0.1$, ensuring that multiple mutations segregate simultaneously ($NU \geqslant 1$ in all of our experimental populations). This difference in approach has two important consequences for this work. It enables 1) recombination and 2) selection for genetic robustness, neither of which occur in the SSWM regime [19].

Once these populations reached a mutation-recombination-selection-drift equilibrium, we made exact copies of each population to obtain pairs of identical ancestral populations. These population pairs were then allowed to diverge allopatrically, again evolving under a Wright-Fisher model with a selection-recombination-mutation life cycle and using the same population size and recombination rate used to generate the ancestral populations. We then investigated the effect that the differences in genetic robustness among the ancestral populations had on the rate at which genetic incompatibilities accumulated between the diverging population pairs.

Genotype and phenotype

The genotype is an RNA sequence of length L. In all simulations summarized in this paper we used $L = 100$ nucleotides and assumed that genotypes are haploid. The phenotype associated with a particular genotype is its minimum free-energy secondary structure, as determined using the ViennaRNA package version 2.1.9 with default parameters [33].

Selection

The viability of sequence i is defined as in Kalirad and Azevedo [13]

$$
w_i = \begin{cases} 1 & \text{if} \quad \beta_i > \alpha \quad \text{and} \quad \delta_i \leqslant \alpha \\ \\ 0 & \text{otherwise} \end{cases} \tag{2}
$$

where β_i is the number of base pairs in the secondary structure of sequence i, δ_i is the base-pair distance between the secondary structure of sequence i and a reference secondary structure, and α is an arbitrary threshold. In all simulations summarized in this paper we used $\alpha = 12$.

Recombination

From a population of size N, we randomly sample with replacement two sets of N viable genotypes, S_1 and S_2. Genotypes from S_1 and S_2 are paired, such that the ith genotypes from each set recombine.

Free recombination   In the free recombination regime, every offspring is recombinant. Crossovers occur independently with probability 0.5 at each of the $L - 1$ internucleotide sites in the genome.

Limited recombination   In the limited (not free) recombination regime, offspring are recombinant with probability r resulting from a single crossover located at random at one of the $L - 1$ internucleotide sites in the genome.

No recombination   If there is no recombination, the offspring is a copy of a randomly selected parent.

Mutation

Mutations arise according to a binomial process where each site mutates with probability equal to the per-site mutation rate μ in each generation. All types of point mutations occur with equal probability. Insertions and deletions are not considered. In all simulations summarized in this paper we used $\mu = 0.001$, resulting in a genome-wide mutation rate of $U = \mu L = 0.1$.

Evolution

Ancestor   We start by generating a random RNA sequence with $L = 100$ nucleotides and defining its minimum free-energy secondary structure as the reference provided it is viable (see Eq 2). This initial sequence is not representative of the holey fitness landscape it defines because it has the reference structure (i.e., $\delta = 0$, see Eq 2), whereas most viable sequences are $\delta \approx \alpha$ base pairs away from the reference structure [13]. To obtain a representative sequence, we subject the initial sequence to a random walk of 200 steps, allowing multiple hits at individual sites. We define the resulting sequence as the ancestor.

Initial evolution   The ancestor is cloned N times to create a population. We allow this population to evolve for t_1 generations to reach an approximate mutation-recombination-selection-drift equilibrium ($t_1 = 1000$ for all regimes except $N = 500$ with free-recombination, which required $t_1 = 4000$ to reach equilibrium). We made an exact copy of the resulting population to obtain two identical ancestral populations.

Divergence   The two ancestral populations are allowed to evolve for t_2 generations under a Wright-Fisher model following a selection-recombination-mutation life cycle ($t_2 = 500$ for all regimes except $N = 500$ with free-recombination, which required $t_2 = 2000$ to reach similar levels of mean genetic divergence).

Incompatible introgressions

Two viable genotypes, i and j, differ at k sites. The number of incompatible introgressions [13] (IIs) from genotype i to genotype j, $\mathbf{II}_{ij}$, was measured by introgressing each allele at a divergent site from genotype i to genotype j one at a time and counting the number of inviable introgressed genotypes ($w = 0$, see Eq 2). $\mathbf{II}_{ij}$ is not necessarily equal to $\mathbf{II}_{ji}$. As populations diverged, we periodically estimated $\mathbf{II}_{ij}$ in a particular direction (e.g., from population 1 to population 2; Fig 2, $\mathbf{II}_\mathrm{B}$) from 10 randomly chosen pairs of individuals from the two populations, sampled with replacement (except when $N = 1$, where the assay was conducted between the single individuals).

Segregating IIs

We also counted IIs within populations as they diverged (Fig 2, $\mathbf{II}_\mathrm{W}$). We used the same approach as for IIs between populations, except that the 10 pairs of individuals were sampled randomly with replacement from the same population.

Mutational robustness

The mutational robustness of a genotype was measured as the proportion ν of its $3L$ single mutations that did not render the genotype inviable (Fig 1c). The robustness of each replicate population was estimated as the average ν of 10 individuals sampled at random with replacement (except when $N = 1$, where it was simply the ν of the individual).

Reproductive isolation

The level of postzygotic reproductive isolation (RI) between two populations is defined as:

$$\mathrm{RI} = 1 - \frac{1}{N} \sum_{i=1}^{N} w_i \quad , \tag{3}$$

where w_i is the fitness of a genotype resulting the from recombination between the ith genotypes of sets S_1 and S_2, where set S_j contains N genotypes drawn randomly with replacement from diverging population j. Recombinant sequences were produced by

either a single crossover or free recombination.

Statistical analysis

Snowball exponent   The model in Eq 4 was fit by least-squares regression. For a particular regime (e.g., SSWM in Fig 3a) or combination of regimes (e.g., all populations summarized in Figs 3 and 4) the snowball exponent c was estimated by maximum likelihood (ML) while allowing each replicate simulation run to have different values of a and b. ML estimation was performed using the `mle2` function of the R package `bbmle` version 1.0.25 (`https://github.com/bbolker/bbmle`).

Rate of II accumulation   Once a value of c was estimated, we fit the model in Eq 4 for each replicate simulation run by linear regression ($\mathbf{II} = a + bx$, where $x = k^c$ for a fixed value of c). The rates of II accumulation summarized in Figs 3c and 4c were calculated by averaging estimates of b from individual replicate runs.

Phylogenetic analyses   Phylogenetic generalized least-squares regression analyses were done using the R package `caper` [34] version 1.0.1. Phylogenetic signal was tested using the R packages `phytools` [35] version 1.0-3 and `geiger` [36] version 2.0.10.

Data and code availability

The software used to run all simulations was written in Python 2.7 with NumPy version 1.14.5 [37]. The code used to analyze the resulting data was written in Python 3.7 with NumPy version 1.21.0 and R 4.1.0 [38]. All code and data are available at `https://github.com/Kalirad/RNA_pop_snowball`.

Results

Genetic drift promotes the accumulation of genetic incompatibilities in asexual populations

We allowed asexual populations of different sizes—between 10 and 500 individuals—to evolve until they reached an equilibrium genetic robustness. We then made a copy of the resulting population and allowed the two populations to diverge in allopatry. Both

populations experienced the same selection regime (see Methods). Thus, this is a ₂₀₂ mutation-order speciation [39] scenario. Fig 3a shows the pattern of accumulation of IIs ₂₀₃ following the population split. The strength of selection for genetic robustness in these ₂₀₄ populations is of the order of the deleterious mutation rate [40] $U_d = U(1 - \nu)$ (Fig 2a: ₂₀₅ $U \to \nu$). For comparison, we subjected "populations" of $N = 1$ individual to a random ₂₀₆ walk, which is equivalent to the SSWM regime and does not impose selection for genetic ₂₀₇ robustness [19] (Fig 3a, gray circles). ₂₀₈

We modelled the accumulation of IIs using the following phenomenological equation ₂₀₉

$$\mathbf{II} = a + bk^c \tag{4}$$

where a is the initial number of IIs within populations ($a = 0$ when $N = 1$), b is the rate ₂₁₀ of accumulation of IIs with genetic divergence, and c is a snowball exponent. Values of ₂₁₁ $c = 1$ and 2 correspond to linear and quadratic patterns of II accumulation, respectively. ₂₁₂ The SSWM simulations showed an overall snowball exponent of $c = 1.619$ (maximum ₂₁₃ likelihood, ML, estimate; 95% confidence interval, CI: 1.510, 1.736). Thus, IIs ₂₁₄ accumulated according to a snowball pattern ($c > 1$). ₂₁₅

Increasing population size caused the accumulation of IIs to slow down in asexual ₂₁₆ populations. Fitting Eq 4 to the data at each population size while keeping the snowball ₂₁₇ exponent constant at $c = 1.479$ (the ML estimate of c taking into account all simulated ₂₁₈ populations summarized in Figs 3 and 4, from $N = 1$ to $N = 500$) resulted in a ₂₁₉ reduction of the slope b with population size (Fig 3c, blue circles). For example, ₂₂₀ populations of $N = 10$ and 500 individuals accumulated IIs at 90.6% and 81.1% of the ₂₂₁ rate of populations of $N = 1$ individual, respectively. Increasing population size reduced ₂₂₂ the importance of genetic drift and was expected to allow selection for genetic robustness ₂₂₃ to operate more efficiently (Fig 2a: $N \to \nu$). Indeed, larger populations evolved higher ₂₂₄ genetic robustness, ν, before populations began diverging from each other (Fig 3d, blue ₂₂₅ circles). The level of genetic robustness in diverging populations explains $R^2 = 75.0\%$ of ₂₂₆ the variation in the rate of II accumulation in populations of different sizes. ₂₂₇

Fig 3. Genetic drift promotes and recombination hinders the accumulation of genetic incompatibilities. Incompatible introgressions (IIs) accumulate with genetic distance in asexual **(a)** and sexual **(b)** populations of different sizes ($N = 1, 10, \ldots 500$). Genetic distance (k) between divergent populations was measured by Jost's D [41]. In all simulations, genome length was $L = 100$ nucleotides and the mutation rate was $U = 0.1$ per genome per generation. Sexual populations experienced random mating and free recombination. **(c)** Relative rates of II accumulation (b_N/b_1) at different population sizes. For each N, the rate b_N was estimated by fitting Eq 4 to each simulation run, as described in the Methods. The average value of b_N was then divided by the average value under the SSWM regime (b_1). **(d)** Genetic robustness at different population sizes. The genetic robustness of a genotype was measured as the proportion ν of its mutational neighbors that are viable. Note that these populations are at equilibrium before divergence begins, so that robustness does not evolve over the period shown. Plotted values are means of 200 replicate simulation runs at each population size calculated every 100 ($N = 1, 10, 50, 100$) or 400 ($N = 500$) generations. Error bars are 95% CIs.

Genetic drift promotes the accumulation of genetic incompatibilities in sexual populations

Next, we repeated the evolutionary simulations but allowed genotypes to mate at random and recombine freely. Fig 3b shows the accumulation of IIs in populations of

Fig 4. Recombination hinders the accumulation of genetic incompatibilities.
(a) IIs accumulate with genetic distance in populations of $N = 100$ individuals
experiencing different recombination probabilities. Sexual populations $(r > 0)$
experienced random mating. The values of the recombination probability
$(r = 0, 0.1, 0.5, 1)$ indicate the expected proportion of matings resulting in a
recombinant progeny containing one crossover event. Under free recombination, every
progeny is a recombinant with an expected 49.5 crossover events. **(b)** Genetic
robustness at different recombination probabilities. **(c)** Relative rates of II
accumulation (b_r/b_0) at different recombination probabilities. For each r, the rate b_r
was estimated by fitting Eq 4 to each simulation run. The average value of b_r was then
divided by the average value under the asexual regime (b_0). **(d)** Number of IIs
segregating within populations at different recombination probabilities (see Methods).
Values are means of 200 replicate simulation runs at each recombination probability
calculated every 100 generations. Error bars are 95% CIs.

the same size as in Fig 3a but with free-recombination. In populations of a single   232
individual (i.e., the SSWM regime), recombination cannot operate. For populations   233
with $N > 1$, the effect of population size on the accumulation of IIs was stronger than   234
among the asexual populations (Fig 3c, orange circles). Larger populations accumulated   235
IIs much more slowly than smaller populations. Again, this pattern is well explained by   236

variation in the level of genetic robustness in free-recombining populations of different sizes ($R^2 = 97.9\%$; Fig 3d, orange circles).

What is the mechanism underlying the differences in genetic robustness among populations of different sizes? One possibility is that more robust genotypes are further from the holes in the fitness landscape. In all simulations reported here, a sequence is inviable if it is at a base-pair distance $\delta > 12$ from a reference structure (see Methods). As expected, genotypes from a population of $N = 10$ individuals were at an average distance of $\bar{\delta} = 11.32 \pm 0.10$ (95% CI); living on the edge of a hole made them more fragile. More abundant populations were further from the edge and, therefore, more robust: $\bar{\delta} = 10.18 \pm 0.16$ for $N = 50$ and $\bar{\delta} = 6.84 \pm 0.23$ for $N = 500$ individuals.

Recombination hinders the accumulation of genetic incompatibilities

Changing the recombination probability in populations of the same size ($N = 100$) allows us to isolate the causal role of recombination in determining the rate of II accumulation (Fig 4a). Populations experiencing more recombination both evolved higher genetic robustness and accumulated IIs more slowly (Fig 2b: $r \to \nu$; Figs 4b and 4c). For example, populations where $r = 10\%$ and 100% of matings experienced one crossover event accumulated IIs at 83.1% and 50.4% of the rate of asexual populations, respectively (again, assuming $c = 1.479$). Increasing the *amount* of recombination from one crossover to free-recombination caused a further reduction in the rate of II accumulation to 27.5% of the rate of asexual populations. The level of genetic robustness in diverging populations explains $R^2 = 98.8\%$ of the variation in the rate of II accumulation in populations experiencing different probabilities and amounts of recombination (Fig 4b).

Recombination hinders the accumulation of genetic incompatibilities within populations

The previous results indicate that recombination hinders the accumulation of genetic incompatibilities. But how does recombination do that? To begin to answer that question we return to our expectation for the accumulation of IIs in the absence of

recombination (Fig 2a). In sufficiently large populations, genetic variation builds up. Because of epistastic interactions, some of these variants constitute within-population IIs (Fig 2a: $\pi \rightarrow \mathbf{II}_W$). In the absence of recombination, however, these segregating genetic incompatibilities will only rarely be expressed—and, therefore, be exposed to natural selection—because they can only occur in the same individual through multiple mutations, which are rare. Therefore, cryptic IIs are expected to accumulate *within* asexual populations. We observed this in our simulations: asexual populations of $N = 100$ and 500 individuals had accumulated 1.36 ± 0.17 and 5.13 ± 0.34 segregating IIs on average *before* divergence began (95% CIs; Fig 3a, genetic distance of 0; Fig 4d, $r = 0$).

We hypothesize that the extent to which a mutation is involved in an II *within* a population is an important determinant of whether that mutation will cause an II *among* populations if it spreads in the population (Fig 2a: $\mathbf{II}_W \rightarrow \mathbf{II}_B$). To test this hypothesis we isolated from each asexual population the first mutation to rise to high frequency and cause an II among populations and the first mutation to rise to high frequency and not cause an II. We then went back to the generations in which these mutations arose, placed them in every individual present in the population at the time, and noted the proportion of these "backcrossed" individuals that remained viable (Fig 5a). As predicted, we found that the proportion of viable backcrossed individuals was much lower for mutations that caused IIs (0.71 ± 0.04, mean and 95% CI; Fig 5b, blue filled circle) than for mutations that did not (0.96 ± 0.01; Fig 5b, blue open circle). In other words, mutations that caused IIs among populations were likely to have started out causing IIs within populations.

We propose that recombination slows down the accumulation of IIs *between* populations because it suppresses the accumulation of IIs *within* populations. In the presence of recombination, segregating IIs can be expressed in recombinant individuals. This suppresses the accumulation of IIs within populations in two ways. First, it causes selection against mutations involved in IIs within populations (Fig 2b: $r \dashv \mathbf{II}_W$). Second, it generates recombination load, which in turn selects for genetic robustness [27, 42–45]. Genetically robust individuals are less likely to express IIs (Fig 2b: $r \rightarrow \nu \dashv \mathbf{II}_W$). We observe this suppression of segregating IIs by recombination in our simulations (Fig 4d). We believe the reduced supply of segregating IIs slows down the accumulation of IIs

Fig 5. Recombination selects for alleles that are more compatible with the genetic backgrounds of other individuals in the population. (a) Illustration of the experimental design used to generate data in (b). Two populations of $N = 4$ individuals diverge in allopatry over the holey fitness landscape shown in Fig 1c. After some time, population 2 (blue) fixes derived allele D. Introgressing D from population 2 to population 1 (arrow I) results in an inviable genotype ($ABcDe$), rendering 50% of introgressed sequences in population 2 inviable (gray box). Backcrossing D into every genotype present in population 2 (arrow II) when the D allele arose by mutation reveals an inviable genotype ($abcDE$), and renders 50% of the sequences in that population inviable too (gray box). Thus, D is involved in IIs both within and among populations. (b) We repeated the simulations summarized in Fig 4 and isolated the first mutation to occur in a population, rise to a frequency of at least 50%, and be involved in an II with the other population. We then went back to the generation in which the mutation arose and backcrossed it into every individual present within its own population at that time. Panel (b) shows the proportion of viable backcrossed individuals made from populations experiencing different recombination probabilities (colored filled circles). As a control, we did the same calculation for the first mutation to rise to high frequency that does not cause an II among populations (open circles). Black and gray circles show the mean genetic robustness of the populations when the mutations arise and when they reach high frequency, respectively. Values are means of 200 replicate simulation runs at each recombination probability. Error bars are 95% CIs.

among populations.

We tested this hypothesis by conducting backcross simulations at different recombination rates (Fig 5b). As recombination rate and robustness increased, we observed a higher proportion of viable backcrossed individuals for mutations that caused IIs among populations (Fig 5b, orange filled circles). Thus, in the presence of recombination, even mutations that are ultimately destined to participate in genetic incompatibilities among populations are less likely to result in within-population genetic incompatibilities (Fig 2b). That is because recombination selects for genotypes that are more compatible with the genetic backgrounds of other individuals in the population and, therefore, with mutations currently segregating in the population.

The number of genetic incompatibilities between populations is correlated with the amount of postzygotic reproductive isolation

In the presence of recombination, genetic incompatibilities between populations cause postzygotic reproductive isolation (RI) between them. Populations evolved at different population sizes under free-recombination (Fig 6a) or at a population size of $N = 100$ under different recombination probabilities (Fig 6b) evolve RI proportional to the number of IIs between them. For the data shown in Fig 6, adding an II between populations causes an increase in postzygotic RI of approximately 5% (one-crossover assay).

Fig 6. Genetic drift promotes and recombination hinders speciation. (a) Postzygotic reproductive isolation (RI) in populations of different sizes evolving under free-recombination (summarized in Fig 3b). **(b)** Postzygotic RI in populations of $N = 100$ individuals evolving under different recombination probabilities (summarized in Fig 4a). RI assays were conducted at the end of the simulations with recombination probability $r = 1$ and either one crossover (filled circles) or free-recombination (open circles). Values are means of 200 replicate simulation runs per regime. Error bars are 95% CIs.

Discussion

Rosenzweig [46] noted that "no one denies that population size influences speciation rate. However, the direction of its effect is in doubt." In his classic neutral theory of biodiversity, Hubbell [47] assumed that all species have the same speciation rate per capita and that, therefore, more abundant species will have higher speciation rates.

Similarly, Marzluff and Dial [48] proposed that a broad geographic distribution, a correlate of population size, promotes speciation. In contrast, Stanley [49] proposed that species whose populations contract and fragment due to, for example, predation or environmental deterioration, tend to experience high rates of speciation.

The few attempts to study this issue using explicit theoretical models have reached the same conclusion as ours, that low abundance promotes speciation. However, the models they considered differ from ours in important aspects. Mayr [50] proposed that speciation often results from "genetic revolutions" precipitated by founder events. Others have developed similar models [51]. Most of them modeled ecological speciation and did not consider genetic incompatibilities as the basis for RI. Gavrilets and Hastings [52] did consider several DMI scenarios and showed that founder effect speciation is "plausible" under some of them, but did not consider the accumulation of multiple DMIs of arbitrary complexity. One question raised by founder effect speciation models is whether fluctuations of population size, such as those caused by founder events, may have a different effect on the accumulation of genetic incompatibilities when compared to constant (low) population size, as in our model.

Khatri and Goldstein showed that genetic incompatibilities accumulate more quickly in smaller populations in several models of gene regulation [14, 15, 53]. Like our model, their models included strong incompatibilities that create holes in the fitness landscape. However, the fitness of a viable genotype is a continuous function of the phenotype in their models. Khatri and Goldstein have argued that their result arises because smaller populations evolve a greater drift load and occupy regions of genotype space closer to the holes in the fitness landscape. We believe that the distance of a genotype to a hole plays the same role in our model because it determines the mutational robustness of the genotype. One limitation of the Khatri and Goldstein studies is that they considered only the SSWM regime and, therefore, were not able to evaluate the effect of recombination. Furthermore, under SSWM, neither genetic drift nor recombination would have any effect on the rate of accumulation of incompatibilities in our model. Therefore, our results must also involve evolutionary mechanisms not captured in the models studied by Khatri and Goldstein. We believe the ingredient they missed is segregating variation at loci involved in genetic incompatibilities. Such variation has been known to exist in natural populations of several species [54, 55]. Corbett-Detig et

al. [55] found evidence that multiple pairwise DMIs are currently segregating within natural populations of *D. melanogaster*. They surveyed a large panel of recombinant inbred lines (RILs) and found 22 incompatible pairs of alleles at unlinked loci in the RILs; of the 44 alleles, 27 were shared by two or more RILs, indicating that multiple DMIs are polymorphic within natural populations. They also found evidence for multiple DMIs in RIL panels in *Arabidopsis* and maize [55]. Corbett-Detig et al. did not attempt to identify DMIs among linked loci or higher-order DMIs and therefore are likely to have underestimated the actual number of segregating DMIs in the RILs. These observations suggest that the conditions for the effects reported here may occur in nature.

Maya-Lastra and Eaton [56] modelled genetically variable populations evolving in the presence of multiple pairwise DMIs like those envisioned by Orr and, like us, observed that smaller populations accumulated more DMIs. They incorporated recombination in their model but did not investigate its effect. They assumed that mutations involved in DMIs were beneficial and that the DMIs were weak. When DMIs were caused by neutral mutations, the effect of population size disappeared. Selection also played a decisive role in our model—recombination generated selection for genetic robustness. Therefore, higher recombination probabilities and rates made selection stronger and larger population sizes made selection more efficient. If every genotype had the same genetic robustness in our model—i.e., if all mutations were truly neutral—, we would also expect the effect of population size on the accumulation of genetic incompatibilities to vanish.

These earlier studies, when combined with ours, lead to a clear prediction that postzygotic RI should build up more quickly in smaller populations. There have been multiple attempts to test this prediction experimentally by subjecting populations to founder events, sometimes repeatedly. In one of these studies, Ringo et al. [57] subjected 8 experimental populations of *D. simulans* to repeated bottlenecks of a single pair of individuals every 6 months. Between bottlenecks these "drift" lines were allowed to expand to large size. Another 8 populations were maintained at a larger size and subjected to a variety of artificial selection regimes. The drift lines experienced stronger inbreeding than the "selection" lines. An ancestral "base" population was kept at larger size than either treatment group throughout. Ringo et al. found that after 6

bottleneck/expansion cycles the drift lines evolved stronger postzygotic RI relative to the base population than did the selection lines (28% vs 4% reduction in hybrid fitness), in agreement with our prediction. However, the results from three similar experiments on other species of *Drosophila* were inconclusive [58–61].

There is also comparative evidence supporting our prediction. Huang et al. [62] tested for correlations between a measure of the level of neutral polymorphism within species—a proxy for effective population size, N_e—and speciation rates estimated from phylogenies in three genera of lichen-forming fungi. They found a significantly negative correlation in one genus, in agreement with our prediction, but no significant correlations in the other two genera. A larger phylogenetic study found significant negative correlations between speciation rates and a different proxy for N_e—geographic range size—in both birds and mammals [56]. These results, while encouraging, have the limitation that they have used overall speciation rate, which may not be driven primarily by postzygotic RI in all lineages. Furthermore, the estimates of speciation rate may themselves be problematic because many different temporal patterns of speciation and extinction rates can have equal probability for a given phylogeny [63,64]. To address these limitations we tested for a correlation between the level of neutral polymorphism within species and the rate with which postzygotic RI accumulates with genetic distance in *Drosophila* [65]. We found a significant negative correlation (Fig 7b). Note, however, that these comparative correlations could have alternative explanations. For example, reductions in population size might be a *consequence* of speciation, rather than its cause [66].

Another prediction from our work is that differences in the size of diverging populations will cause asymmetric accumulation of genetic incompatibilities: the expected number of IIs will be greater in introgressions from a large population to a small population than in the opposite direction. Asymmetric postzygotic RI has been observed between reciprocal crosses of many closely related species—a phenomenon termed Darwin's corollary to Haldane's rule [71]. We propose that this kind of asymmetry might be explained by differences in genetic robustness between the recipient populations to asymmetric introgression of cytoplasmic organelles and of the X chromosome in male offspring. Yukilevich [68] surveyed RI asymmetry between many *Drosophila* species. We were able to determine the level of neutral polymorphism within

Fig 7. *Drosophila* species with higher levels of neutral polymorphism and higher recombination accumulate postzygotic RI more slowly. (a) Phylogenetic relationships between 8 *Drosophila* species groups [65] (*ananassae, immigrans, melanogaster, montium, obscura, repleta, virilis, willistoni*). **(b)** The rates with which postzygotic RI accumulates with genetic distance in the species groups shown in **(a)** were estimated by Rabosky & Matute [65] based on the proportion of sterile or inviable F_1 offspring in interspecific crosses [67, 68]. Positive and negative values represent faster- and slower-than-average RI velocities. The levels of neutral polymorphism in individual species were measured by the synonymous nucleotide diversity (π_s) at the *Adh* locus (see S1 Table). Values are the mean π_s for the species or subspecies in each *Drosophila* species group. Multiple estimates for a single species were averaged before calculating the overall mean for the species group. The line shows a fit by linear ordinary least-squares regression of RI velocity against π_s (OLS : slope and 95% CI = -105 ± 67, $R^2 = 0.71$, $P = 0.008$). The results of a phylogenetic generalized least-squares regression analysis assuming Brownian motion were almost identical (PGLS : $R^2 = 0.82$, $P = 0.002$). **(c)** Residuals in RI velocity from the regression in **(b)** in the species groups with either shorter or longer total map lengths than average. The point areas in **(b)** and **(c)** are proportional to the mean total map length in cM in the species group (see S2 Table). Adding map length to the regression model did not improve fit in a statistically significant manner (multiple regression OLS : coefficient = -0.0016 ± 0.0021, $R^2 = 0.84$, $P = 0.11$; PGLS : $R^2 = 0.89$, $P = 0.13$). None of the traits showed a significant phylogenetic signal based on either the K or λ statistics [69, 70] (see S3 Table).

both species of 8 species pairs in which he found asymmetric postzygotic RI. The 418
pattern of asymmetry in RI between these species pairs showed a statistically significant 419
trend in the direction we predicted (Fig 8). 420

Environmental stochasticity selects directly for environmental robustness and 421
indirectly for genetic robustness [20, 40, 73]. Therefore, we predict that temporal 422
variation in environmental conditions should hinder speciation. This prediction is 423
consistent with the observation that speciation rates are higher in tropical lineages than 424

Fig 8. Differences in the level of neutral polymorphism of two species explain asymmetry in postzygotic RI between them. Postzygotic RI asymmetry for the $sp1/sp2$ pair was estimated [72] by taking the RI score from a ♀ $sp1$ × ♂ $sp2$ cross and subtracting the RI score from the reciprocal cross. The x-axis shows the difference in the synonymous nucleotide diversity ($\pi_{s1} - \pi_{s2}$) at the Adh locus (except the rec/sub pair for which we used the $Adhr$ locus; see S4 Table). The colors of the points match the species groups shown in Fig 7a except that the $quinaria$ group (sister group to $immigrans$) is represented by orange: $montium$ group: $\underline{au}raria$, $\underline{sub}auraria$, $\underline{tri}auraria$; $obscura$ group: $\underline{bog}otana$, $\underline{pse}udoobscura$; $quinaria$ group: $\underline{rec}ens$, $\underline{sub}quinaria$; $repleta$ group: $\underline{ari}zonae$, $\underline{moj}avensis$; $willistoni$ group: $\underline{equ}inoxialis$, $\underline{pau}listorum$, $\underline{wil}listoni$. The line shows a fit by linear ordinary least-squares regression through the origin (slope $= 20.9 \pm 17.8$, $R^2 = 0.52$, $P = 0.028$).

in temperate lineages of amphibians [74], birds [75], and mammals [76]. Furthermore, Yukilevich [72] found that postzygotic RI (but not prezygotic RI) accumulated more than twice as quickly between tropical species pairs of *Drosophila* than between temperate and subtropical species pairs of *Drosophila*. However, many other hypotheses have been proposed to explain these patterns [77, 78].

As with population size, no one denies that recombination influences speciation rate. But, again, the direction of its effect is disputed. Butlin [79] argued that "speciation can be viewed as the evolution of restrictions on the freedom of recombination". In general, low recombination is found to promote speciation in the *presence* of gene flow between diverging populations because it restricts introgression. For example, chromosomal rearrangements often suppress recombination along a large region of the genome. Navarro and Barton [80] showed that such chromosomal rearrangements facilitate the accumulation of DMIs in the presence of gene flow. However, these results are not universal. Bank et al. studied the effect of recombination on the maintenance of

a DMI involving two loci in a haploid with gene flow from a continental to an island $_{439}$
population [81]. They showed that recombination can have opposing effects on the $_{440}$
maintenance of the DMI depending on the fitnesses of the genotypes at the DMI loci. In $_{441}$
some scenarios, the DMI is maintained more easily through selection against immigrants $_{442}$
when recombination is low. But in other scenarios, a DMI is maintained more easily $_{443}$
through selection against hybrids when recombination is high. Founder effect speciation $_{444}$
models have also tended to conclude that high recombination should promote $_{445}$
speciation [82]. $_{446}$

There is some comparative evidence for the prediction that recombination slows $_{447}$
down the accumulation of postzygotic RI. We found a negative correlation between the $_{448}$
genome-wide recombination rate and the rate with which postzygotic RI accumulates $_{449}$
with genetic distance in *Drosophila* species groups [65], after removing the effect of the $_{450}$
level of neutral polymorphism (Fig 7c). Including recombination rate in the ordinary $_{451}$
least-squares regression model increased the proportion of variance in RI velocity $_{452}$
explained by the predictors from 71% to 84%. However, this improvement was not $_{453}$
statistically significant ($P = 0.11$). $_{454}$

In contrast to the results from earlier models, we found that recombination within $_{455}$
populations can hinder speciation in the *absence* of gene flow. Our results indicate that $_{456}$
recombination suppresses the accumulation of genetic incompatibilities within $_{457}$
populations, which in turn slows down their accumulation between populations. In other $_{458}$
words, we found that microevolution and macroevolution are *coupled* in our model. This $_{459}$
coupling is also a feature of our result that genetic drift promotes speciation. In both $_{460}$
cases, processes acting within populations to alter genetic robustness explain variation in $_{461}$
speciation rates, contrary to Stanley's [4] contention that speciation is largely random. $_{462}$

The size of a population is also predicted to be inversely related to its probability of $_{463}$
extinction [83–86]. Thus, if our results are correct, depending on the balance of the $_{464}$
effects of population size on speciation and extinction, random genetic drift within $_{465}$
species could be a driver of deterministic species selection [2], the process championed $_{466}$
by Stanley [4]. $_{467}$

Our results lead to the prediction that recombination in large populations impedes $_{468}$
speciation; such populations would maintain extensive genetic variation but even highly $_{469}$
divergent genotypes would remain compatible with each other. This prediction is $_{470}$

consistent with the observation that recombination between highly divergent genotypes is common in viruses, including influenza A [87,88], HIV-1 [89,90], hepatitis B [91] and hepatitis E [92] viruses, rhinoviruses [93,94], coronaviruses [95–101], and geminiviruses [102]. Such recombination events have often contributed to the emergence of viral diseases. For example, the virus that caused the 1918 influenza pandemic is believed to have arisen when a human virus containing an H1 hemagglutinin acquired an N1 neuraminidase from an avian virus [88].

The extent to which our results can be generalized to real organisms requires further investigation. Our model makes several strong assumptions, such as, that all viable genotypes have the same viability (although they can differ in overall fitness), all genetic incompatibilities result in complete inviability, strong epistatic interactions are common, haploidy, and populations of constant size diverge in allopatry under the same selection regime (i.e., mutation-order speciation [39]). The effect of each of these assumptions can be investigated within the framework of our model. Such an exploration has the potential to serve as the foundation for a new theory of the coupling between microevolution and macroevolution.

Supporting information

S1 Table.   Synonymous nucleotide diversity (π_s) at the Adh locus for different species of *Drosophila*.

S2 Table.   Total recombination map length for different species of *Drosophila*.

S3 Table.   Phylogenetic signal for each trait analyzed in Fig. 7.

S4 Table.   Asymmetry in synonymous nucleotide diversity (π_s) at the Adh locus and in postzygotic reproductive isolation (RI) for different species of *Drosophila*.

Acknowledgments 496

The authors thank D. Wiernasz and M. Servedio for helpful discussions. We used the 497
Opuntia cluster from the Hewlett Packard Enterprise Data Science Institute at the 498
University of Houston. We thank the advanced support from the Research Computing 499
Data Core at the University of Houston. This work was funded by grants from the 500
National Science Foundation DEB-2014566 to RBRA and DEB-2014943 to CB. All 501
authors conceived the research and designed the analyses. AK and RBRA conducted the 502
analyses. All authors wrote the manuscript. The authors declare no competing interests. 503

References

1. Cooney CR, Thomas GH. Heterogeneous relationships between rates of speciation and body size evolution across vertebrate clades. Nat Ecol Evol. 2021;5(1):101–110. doi:10.1038/s41559-020-01321-y.

2. Jablonski D. Species Selection: Theory and Data. Annu Rev Ecol Evol Syst. 2008;39:501–524.

3. Harvey MG, Singhal S, Rabosky DL. Beyond Reproductive Isolation: Demographic Controls on the Speciation Process. Annu Rev Ecol Evol Syst. 2019;50(1):75–95. doi:10.1146/annurev-ecolsys-110218-024701.

4. Stanley SM. A theory of evolution above the species level. Proc Natl Acad Sci U S A. 1975;72(2):646–650. doi:10.1073/pnas.72.2.646.

5. Dobzhansky T. Genetics and the Origin of Species. New York: Columbia Univ. Press; 1937.

6. Muller HJ. Isolating mechanisms, evolution and temperature. Biol Symp. 1942;6:71–125.

7. Orr HA. The population genetics of speciation: the evolution of hybrid incompatibilities. Genetics. 1995;139(4):1805–1813.

8. Matute DR, Butler IA, Turissini DA, Coyne JA. A Test of the Snowball Theory for the Rate of Evolution of Hybrid Incompatibilities. Science. 2010;329(5998):1518–1521. doi:10.1126/science.1193440.

9. Matute DR, Gavin-Smyth J. Fine Mapping of Dominant X-Linked Incompatibility Alleles in *Drosophila* Hybrids. PLoS Genet. 2014;10(4):e1004270. doi:10.1371/journal.pgen.1004270.

10. Wang RJ, White MA, Payseur BA. The Pace of Hybrid Incompatibility Evolution in House Mice. Genetics. 2015;201(1):229–242. doi:10.1534/genetics.115.179499.

11. Moyle LC, Nakazato T. Hybrid Incompatibility "Snowballs" Between *Solanum* Species. Science. 2010;329(5998):1521–1523. doi:10.1126/science.1193063.

12. Sherman NA, Victorine A, Wang RJ, Moyle LC. Interspecific Tests of Allelism Reveal the Evolutionary Timing and Pattern of Accumulation of Reproductive Isolation Mutations. PLoS Genet. 2014;10(9):e1004623. doi:10.1371/journal.pgen.1004623.

13. Kalirad A, Azevedo RBR. Spiraling Complexity: A Test of the Snowball Effect in a Computational Model of RNA Folding. Genetics. 2017;206(1):377–388. doi:10.1534/genetics.116.196030.

14. Khatri BS, Goldstein RA. Simple Biophysical Model Predicts Faster Accumulation of Hybrid Incompatibilities in Small Populations Under Stabilizing Selection. Genetics. 2015;201(4):1525–1537. doi:10.1534/genetics.115.181685.

15. Khatri BS, Goldstein RA. Biophysics and population size constrains speciation in an evolutionary model of developmental system drift. PLoS Comput Biol. 2019;15(7):e1007177. doi:10.1371/journal.pcbi.1007177.

16. Kondrashov AS, Sunyaev S, Kondrashov FA. Dobzhansky–Muller incompatibilities in protein evolution. Proc Natl Acad Sci U S A. 2002;99(23):14878–14883. doi:10.1073/pnas.232565499.

17. Welch JJ. Accumulating Dobzhansky-Muller Incompatibilities: Reconciling Theory and Data. Evolution. 2004;58(6):1145–1156. doi:10.2307/3449212.

18. Mathews DH, Sabina J, Zuker M, Turner DH. Expanded sequence dependence of thermodynamic parameters improves prediction of RNA secondary structure. J Mol Biol. 1999;288(5):911–940. doi:10.1006/jmbi.1999.2700.

19. van Nimwegen E, Crutchfield JP, Huynen M. Neutral evolution of mutational robustness. Proc Natl Acad Sci U S A. 1999;96(17):9716–9720. doi:10.1073/pnas.96.17.9716.

20. Ancel LW, Fontana W. Plasticity, evolvability, and modularity in RNA. J Exp Zool (Mol Dev Evol). 2000;288(3):242–283. doi:10.1002/1097-010X(20001015)288:3<242::AID-JEZ5>3.0.CO;2-O.

21. Wagner A. Robustness and evolvability: a paradox resolved. Proc R Soc B. 2008;275(1630):91–100. doi:10.1098/rspb.2007.1137.

22. Draghi JA, Parsons TL, Wagner GP, Plotkin JB. Mutational robustness can facilitate adaptation. Nature. 2010;463(7279):353–355. doi:10.1038/nature08694.

23. Draghi JA, Parsons TL, Plotkin JB. Epistasis increases the rate of conditionally neutral substitution in an adapting population. Genetics. 2011;187(4):1139–1152. doi:10.1534/genetics.110.125997.

24. Gavrilets S. Fitness Landscapes and the Origin of Species. Princeton, N.J.: Princeton Univ. Press; 2004.

25. Lipman DJ, Wilbur WJ. Modelling Neutral and Selective Evolution of Protein Folding. Proc R Soc B. 1991;245(1312):7–11. doi:10.1098/rspb.1991.0081.

26. Schuster P, Fontana W, Stadler PF, Hofacker IL. From sequences to shapes and back: a case study in RNA secondary structures. Proc R Soc B. 1994;255(1344):279–284. doi:10.1098/rspb.1994.0040.

27. Azevedo RBR, Lohaus R, Srinivasan S, Dang KK, Burch CL. Sexual reproduction selects for robustness and negative epistasis in artificial gene networks. Nature. 2006;440(7080):87–90. doi:10.1038/nature04488.

28. Szöllősi GJ, Derényi I. The effect of recombination on the neutral evolution of genetic robustness. Math Biosci. 2008;214(1-2):58–62. doi:10.1016/j.mbs.2008.03.010.

29. Klug A, Park SC, Krug J. Recombination and mutational robustness in neutral fitness landscapes. PLoS Comput Biol. 2019;15(8):e1006884. doi:10.1371/journal.pcbi.1006884.

30. Orr HA, Turelli M. The evolution of postzygotic isolation: accumulating Dobzhansky-Muller incompatibilities. Evolution. 2001;55(6):1085–1094. doi:10.1111/j.0014-3820.2001.tb00628.x.

31. Livingstone K, Olofsson P, Cochran G, Dagilis A, MacPherson K, Seitz Jr KA. A stochastic model for the development of Bateson–Dobzhansky–Muller incompatibilities that incorporates protein interaction networks. Math Biosci. 2012;238(1):49–53. doi:10.1016/j.mbs.2012.03.006.

32. Gillespie JH. A simple stochastic gene substitution model. Theor Popul Biol. 1983;23(2):202–215.

33. Lorenz R, Bernhart SH, Höner zu Siederdissen C, Tafer H, Flamm C, Stadler PF, et al. ViennaRNA Package 2.0. Algorithms for Molecular Biology. 2011;6:26. doi:10.1186/1748-7188-6-26.

34. Orme D, Freckleton R, Thomas G, Petzoldt T, Fritz S, Isaac N, et al.. caper: Comparative Analyses of Phylogenetics and Evolution in R.; 2018.

35. Revell LJ. phytools: an R package for phylogenetic comparative biology (and other things). Meth Ecol Evol. 2012;3(2):217–223. doi:10.1111/j.2041-210X.2011.00169.x.

36. Pennell MW, Eastman JM, Slater GJ, Brown JW, Uyeda JC, FitzJohn RG, et al. geiger v2.0: an expanded suite of methods for fitting macroevolutionary models to phylogenetic trees. Bioinformatics. 2014;30(15):2216–2218. doi:10.1093/bioinformatics/btu181.

37. Harris CR, Millman KJ, van der Walt SJ, Gommers R, Virtanen P, Cournapeau D, et al. Array programming with NumPy. Nature. 2020;585(7825):357–362. doi:10.1038/s41586-020-2649-2.

38. Team RC. R: A Language and Environment for Statistical Computing; 2021. Available from: https://www.R-project.org/.

39. Schluter D. Evidence for Ecological Speciation and Its Alternative. Science. 2009;323(5915):737–741. doi:10.1126/science.1160006.

40. de Visser JAGM, Hermisson J, Wagner GP, Meyers LA, Bagheri-Chaichian H, Blanchard JL, et al. Perspective: Evolution and detection of genetic robustness. Evolution. 2003;57(9):1959–1972.

41. Jost L. G_{ST} and its relatives do not measure differentiation. Mol Ecol. 2008;17(18):4015–4026. doi:10.1111/j.1365-294X.2008.03887.x.

42. Gardner A, Kalinka AT. Recombination and the evolution of mutational robustness. J Theor Biol. 2006;241(4):707–715.

43. Misevic D, Ofria C, Lenski RE. Sexual reproduction reshapes the genetic architecture of digital organisms. Proc R Soc B. 2006;273(1585):457–464.

44. Martin OC, Wagner A. Effects of recombination on complex regulatory circuits. Genetics. 2009;183:673–684. doi:10.1534/genetics.109.104174.

45. Lohaus R, Burch CL, Azevedo RBR. Genetic architecture and the evolution of sex. J Hered. 2010;101(suppl. 1):S142–S157. doi:10.1093/jhered/esq013.

46. Rosenzweig ML. Loss of speciation rate will impoverish future diversity. Proc Natl Acad Sci U S A. 2001;98(10):5404–5410. doi:10.1073/pnas.101092798.

47. Hubbell SP. The Unified Neutral Theory of Biodiversity and Biogeography. Princeton, N.J.: Princeton Univ. Press; 2001.

48. Marzluff JM, Dial KP. Life History Correlates of Taxonomic Diversity. Ecology. 1991;72(2):428–439.

49. Stanley SM. Population Size, Extinction, and Speciation: The Fission Effect in Neogene Bivalvia. Paleobiology. 1986;12(1):89–110.

50. Mayr E. Change of genetic environment and evolution. In: Huxley J, Hardy AC, Ford EB, editors. Evolution as a Process. Allen & Unwin; 1954. p. 157–180.

51. Carson HL, Templeton AR. Genetic revolutions in relation to speciation phenomena: the founding of new populations. Annu Rev Ecol Syst. 1984;15:97–131. doi:10.1146/annurev.es.15.110184.000525.

52. Gavrilets S, Hastings A. Founder Effect Speciation: A Theoretical Reassessment. Am Nat. 1996;147(3):466–491. doi:10.2307/2463218.

53. Khatri BS, Goldstein RA. A coarse-grained biophysical model of sequence evolution and the population size dependence of the speciation rate. J Theor Biol. 2015;378:56–64. doi:10.1016/j.jtbi.2015.04.027.

54. Cutter AD. The polymorphic prelude to Bateson–Dobzhansky–Muller incompatibilities. Tr Ecol Evol. 2012;27(4):209–218. doi:10.1016/j.tree.2011.11.004.

55. Corbett-Detig RB, Zhou J, Clark AG, Hartl DL, Ayroles JF. Genetic incompatibilities are widespread within species. Nature. 2013;504(7478):135–137. doi:10.1038/nature12678.

56. Maya-Lastra CA, Eaton DAR. Genetic incompatibilities do not snowball in a demographic model of speciation. bioRxiv. 2021;doi:10.1101/2021.02.23.432472.

57. Ringo J, Wood D, Rockwell R, Dowse H. An Experiment Testing Two Hypotheses of Speciation. Am Nat. 1985;126(5):642–661. doi:10.1086/284445.

58. Powell JR. The founder-flush speciation theory: an experimental approach. Evolution. 1978;32(3):465–474. doi:https://doi.org/10.1111/j.1558-5646.1978.tb04589.x.

59. Kilias G, Alahiotis SN, Pelecanos M. A multifactorial genetic investigation of speciation theory using *Drosophila melanogaster*. Evolution. 1980;34(4):730–737. doi:10.1111/j.1558-5646.1980.tb04012.x.

60. Dodd DMB, Powell JR. Founder-flush speciation: an update of experimental results with *Drosophila*. Evolution. 1985;39(6):1388–1392. doi:10.1111/j.1558-5646.1985.tb05704.x.

61. Galiana A, González-Candelas F, Moya A. Postmating isolation analysis in founder-flush experimental populations of *Drosophila pseudoobscura*. Evolution. 1996;50(2):941–944. doi:10.1111/j.1558-5646.1996.tb03905.x.

62. Huang JP, Leavitt SD, Lumbsch HT. Testing the impact of effective population size on speciation rates—a negative correlation or lack thereof in lichenized fungi. Sci Rep. 2018;8:5729. doi:10.1038/s41598-018-24120-9.

63. Louca S, Pennell MW. Extant timetrees are consistent with a myriad of diversification histories. Nature. 2020;580(7804):502–505. doi:10.1038/s41586-020-2176-1.

64. Louca S, Pennell MW. Why extinction estimates from extant phylogenies are so often zero. Curr Biol. 2021;31(14):3168–3173. doi:10.1016/j.cub.2021.04.066.

65. Rabosky DL, Matute DR. Macroevolutionary speciation rates are decoupled from the evolution of intrinsic reproductive isolation in *Drosophila* and birds. Proc Natl Acad Sci U S A. 2013;110(38):15354–15359. doi:10.1073/pnas.1305529110.

66. Glazier DS. Toward a predictive theory of speciation: The ecology of isolate selection. J Theor Biol. 1987;126(3):323–333. doi:10.1016/S0022-5193(87)80239-4.

67. Coyne JA, Orr HA. Patterns of Speciation in *Drosophila*. Evolution. 1989;43(2):362–381. doi:10.2307/2409213.

68. Yukilevich R. Asymmetrical patterns of speciation uniquely support reinforcement in *Drosophila*. Evolution. 2012;66(5):1430–1446. doi:10.1111/j.1558-5646.2011.01534.x.

69. Pagel M. Inferring the historical patterns of biological evolution. Nature. 1999;401(6756):877–884. doi:10.1038/44766.

70. Blomberg SP, Garland T Jr, Ives AR. Testing for phylogenetic signal in comparative data: behavioral traits are more labile. Evolution. 2003;57(4):717–745. doi:10.1111/j.0014-3820.2003.tb00285.x.

71. Turelli M, Moyle LC. Asymmetric Postmating Isolation: Darwin's Corollary to Haldane's Rule. Genetics. 2007;176(2):1059–1088. doi:10.1534/genetics.106.065979.

72. Yukilevich R. Tropics accelerate the evolution of hybrid male sterility in *Drosophila*. Evolution. 2013;67(6):1805–1814. doi:10.1111/evo.12056.

73. Meir E, von Dassow G, Munro E, Odell GM. Robustness, flexibility, and the role of lateral inhibition in the neurogenic network. Curr Biol. 2002;12(10):778–786.

74. Pyron RA, Wiens JJ. Large-scale phylogenetic analyses reveal the causes of high tropical amphibian diversity. Proc R Soc B. 2013;280(1770):20131622. doi:10.1098/rspb.2013.1622.

75. Ricklefs RE. Estimating diversification rates from phylogenetic information. Tr Ecol Evol. 2007;22(11):601–610. doi:10.1016/j.tree.2007.06.013.

76. Rolland J, Condamine FL, Jiguet F, Morlon H. Faster Speciation and Reduced Extinction in the Tropics Contribute to the Mammalian Latitudinal Diversity Gradient. PLoS Biol. 2014;12(1):e1001775. doi:10.1371/journal.pbio.1001775.

77. Mittelbach GG, Schemske DW, Cornell HV, Allen AP, Brown JM, Bush MB, et al. Evolution and the latitudinal diversity gradient: speciation, extinction and biogeography. Ecol Lett. 2007;10(4):315–331. doi:https://doi.org/10.1111/j.1461-0248.2007.01020.x.

78. Schluter D. Speciation, Ecological Opportunity, and Latitude. Am Nat. 2016;187(1):1–18. doi:10.1086/684193.

79. Butlin RK. Recombination and speciation. Mol Ecol. 2005;14(9):2621–2635. doi:10.1111/j.1365-294X.2005.02617.x.

80. Navarro A, Barton NH. Accumulating postzygotic isolation genes in parapatry: A new twist on chromosomal speciation. Evolution. 2003;57(3):447–459. doi:10.1554/0014-3820(2003)057[0447:APIGIP]2.0.CO;2.

81. Bank C, Bürger R, Hermisson J. The Limits to Parapatric Speciation: Dobzhansky–Muller Incompatibilities in a Continent–Island Model. Genetics. 2012;191(3):845–863. doi:10.1534/genetics.111.137513.

82. Templeton AR. The reality and importance of founder speciation in evolution. BioEssays. 2008;30(5):470–479. doi:10.1002/bies.20745.

83. MacArthur RH, Wilson EO. The theory of island biogeography. Princeton, N.J.: Princeton University Press; 1967.

84. Richter-Dyn N, Goel NS. On the extinction of a colonizing species. Theor Popul Biol. 1972;3(4):406–433. doi:10.1016/0040-5809(72)90014-7.

85. Pimm SL, Jones HL, Diamond J. On the Risk of Extinction. Am Nat. 1988;132(6):757–785. doi:10.1086/284889.

86. Lande R. Risks of Population Extinction from Demographic and Environmental Stochasticity and Random Catastrophes. Am Nat. 1993;142(6):911–927. doi:10.1086/285580.

87. Lindstrom SE, Cox NJ, Klimov A. Genetic analysis of human H2N2 and early H3N2 influenza viruses, 1957-1972: evidence for genetic divergence and multiple reassortment events. Virology. 2004;328(1):101–119. doi:10.1016/j.virol.2004.06.009.

88. Worobey M, Han GZ, Rambaut A. Genesis and pathogenesis of the 1918 pandemic H1N1 influenza A virus. Proc Natl Acad Sci U S A. 2014;111(22):8107–8112. doi:10.1073/pnas.1324197111.

89. Robertson DL, Sharp PM, McCutchan FE, Hahn BH. Recombination in HIV-1. Nature. 1995;374(6518):124–126. doi:10.1038/374124b0.

90. Robertson DL, Hahn BH, Sharp PM. Recombination in AIDS viruses. J Mol Evol. 1995;40(3):249–259. doi:10.1007/BF00163230.

91. Simmonds P, Midgley S. Recombination in the genesis and evolution of hepatitis B virus genotypes. J Virol. 2005;79(24):15467–15476. doi:10.1128/JVI.79.24.15467-15476.2005.

92. van Cuyck H, Fan J, Robertson DL, Roques P. Evidence of recombination between divergent hepatitis E viruses. J Virol. 2005;79(14):9306–9314. doi:10.1128/JVI.79.14.9306-9314.2005.

93. Huang T, Wang W, Bessaud M, Ren P, Sheng J, Yan H, et al. Evidence of Recombination and Genetic Diversity in Human Rhinoviruses in Children with

Acute Respiratory Infection. PLoS ONE. 2009;4(7):e6355.
doi:10.1371/journal.pone.0006355.

94. Wisdom A, Kutkowska AE, Leitch ECM, Gaunt E, Templeton K, Harvala H,
et al. Genetics, Recombination and Clinical Features of Human Rhinovirus
Species C (HRV-C) Infections; Interactions of HRV-C with Other Respiratory
Viruses. PLoS ONE. 2009;4(12):e8518. doi:10.1371/journal.pone.0008518.

95. Lau SKP, Feng Y, Chen H, Luk HKH, Yang WH, Li KSM, et al. Severe Acute
Respiratory Syndrome (SARS) Coronavirus ORF8 Protein Is Acquired from
SARS-Related Coronavirus from Greater Horseshoe Bats through
Recombination. J Virol. 2015;89(20):10532–10547. doi:10.1128/JVI.01048-15.

96. Sabir JSM, Lam TTY, Ahmed MMM, Li L, Shen Y, Abo-Aba SEM, et al.
Co-circulation of three camel coronavirus species and recombination of
MERS-CoVs in Saudi Arabia. Science. 2016;351(6268):81–84.
doi:10.1126/science.aac8608.

97. Dudas G, Rambaut A. MERS-CoV recombination: implications about the
reservoir and potential for adaptation. Virus Evolution. 2016;2(1):vev023.
doi:10.1093/ve/vev023.

98. Tang X, Wu C, Li X, Song Y, Yao X, Wu X, et al. On the origin and continuing
evolution of SARS-CoV-2. Natl Sci Rev. 2020;7(6):1012–1023.
doi:10.1093/nsr/nwaa036.

99. Boni MF, Lemey P, Jiang X, Lam TTY, Perry BW, Castoe TA, et al.
Evolutionary origins of the SARS-CoV-2 sarbecovirus lineage responsible for the
COVID-19 pandemic. Nat Microbiol. 2020;5(11):1408–1417.
doi:10.1038/s41564-020-0771-4.

100. Bobay LM, O'Donnell AC, Ochman H. Recombination events are concentrated
in the spike protein region of *Betacoronaviruses*. PLoS Genet.
2020;16(12):e1009272. doi:10.1371/journal.pgen.1009272.

101. Zhou H, Ji J, Chen X, Bi Y, Li J, Wang Q, et al. Identification of novel bat coronaviruses sheds light on the evolutionary origins of SARS-CoV-2 and related viruses. Cell. 2021;184(17):4380–4391. doi:10.1016/j.cell.2021.06.008.

102. Padidam M, Sawyer S, Fauquet CM. Possible emergence of new geminiviruses by frequent recombination. Virology. 1999;265(2):218–25. doi:10.1006/viro.1999.0056.

Table S1. Synonymous nucleotide diversity (πS) at the *Adh* locus for different species of *Drosophila* .

group	species	πS	sequences	reference	notes
ananassae	*ananassae*	0.015822	10	1	
obscura	*bogotana*	0.004841	8	2	
obscura	*miranda*	0.003703	12	3	
obscura	*persimilis*	0.013317	6	4	
obscura	*pseudoobscura*	0.016660	139	2, 5, 6	
obscura	*subobscura*	0.023465	16	7	
virilis	*americana*	0.034155	19	8	G96X
virilis	*americana*	0.023301	19	8	G96Y
virilis	*borealis*	0.059733	2	9, 10	
virilis	*lummei*	0.000000	2	9, 11	
virilis	*texana*	0.029706	10	8	
virilis	*virilis*	0.005490	2	11	
repleta	*arizonae*	0.013155	13	12	Adh-1
repleta	*arizonae*	0.052620	11	12	Adh-2
repleta	*buzzatii*	0.016323	4	13	
repleta	*mojavensis*	0.013141	13	12	Adh-1
repleta	*mojavensis*	0.012597	15	14	Adh-1
repleta	*mojavensis*	0.021064	13	12	Adh-2
repleta	*mojavensis*	0.009859	15	14	Adh-2
repleta	*hydei*	0.017564	9	13	
repleta	*mulleri*	0.019467	6	13	
immigrans	*albomicans*	0.031127	16	15	
immigrans	*hypocausta*	0.031060	2	16	
immigrans	*nasuta*	0.003687	16	15	
immigrans	*siamana*	0.038144	2	16, 17	
melanogaster	*lutescens*	0.009242	4	18–20	
melanogaster	*mauritiana*	0.002228	6	21	
melanogaster	*melanogaster*	0.024036	10	22	
melanogaster	*melanogaster*	0.027122	4	23	
melanogaster	*melanogaster*	0.030007	11	24	
melanogaster	*sechellia*	0.000000	2	21	
melanogaster	*simulans*	0.027407	7	25	
melanogaster	*yakuba*	0.007072	12	25	
melanogaster	*yakuba*	0.015629	36	26	
willistoni	*equinoxialis*	0.005490	2	27	
willistoni	*nebulosa*	0.011506	4	27, 28	
willistoni	*paulistorum*	0.018475	3	27	
willistoni	*willistoni*	0.018035	18	29, 30	
montium	*auraria*	0.045728	2	31	

montium	*birchii*	0.051745	2	31
montium	*kikkawai*	0.034314	21	32
montium	*leontia*	0.027858	2	31
montium	*lini*	0.062418	2	31
montium	*rufa*	0.000000	2	31
montium	*serrata*	0.022432	2	31
montium	*triauraria*	0.011194	2	31
montium	*triauraria*	0.019847	3	33

Methods: Sequences were obtained from GenBank. Synonymous nucleotide diversity (πS) was calculated using the Nei-Gojobori method in MEGA-CC version 11.0.13 (34).

References: 1) Shih & Jones 2008 Genetics 180: 1261–1263. 2) Shaeffer & Miller 1991 PNAS 88: 6097–6101. 3) Yi et al. 2003 Genetics 164: 1369–1381. 4) Wang et al. 1997 Genetics 147: 1091–1106. 5) Schaeffer & Miller 1993 Genetics 135: 541–552. 6) Schaeffer 2002 Genet. Res. 80: 163–175. 7) Jones et al. 2005 Genetics 170: 207–219. 8) McAllister & Charlesworth 1999 Genetics 153: 221–233. 9) Nurminsky et al. 1996 Mol. Biol. Evol. 13: 132–149. 10) Morales-Hojas 2011 Mol. Phyl. Evol. 60: 249–258. 11) Wang et al. 2006 Mol. Phyl. Evol. 40: 484–500. 12) Matzkin & Eanes 2003 Genetics 163: 181–194. 13) Begun 1997 Genetics 145: 375–382. 14) Matzkin 2004 Mol. Biol. Evol. 21: 276–285. 15) Satomura & Tamura 2016 Mol. Biol. Evol. 33: 367–374. 16) Katoh et al. 2007 Zool. Sci. 24: 913–921. 17) Rice et al. 2018 Evol. Dev. 20: 78–88. 18) Katoh et al. 2000 J. Mol. Evol. 51: 122–130. 19) O'Grady & Kidwell 2002 Mol. Phyl. Evol. 22: 442–453. 20) Katoh & Watada, unpublished; Ko, David & Akashi, unpublished. 21) Kliman et al. 2000 Genetics 156: 1913–1931. 22) Begun et al. 1999 Mol. Biol. Evol. 16: 1816–1819. 23) Laurie et al. 1991 Genetics 129: 489–499. 24) Kreitman 1983 Nature 304: 412–417. 25) McDonald & Kreitman 1991 Nature 351: 652–654. 26) Siddiq et al. 2017 Nat. Ecol. Evol. 1: 0025. 27) Gleason et al. 1998 Evolution 52: 1093–1103. 28) Gao et al. 2011 Mol. Phyl. Evol. 60: 98–107. 29) Anderson et al. 1993 Mol. Biol. Evol. 10: 605–618. 30) Griffith & Powell 1997 J. Mol. Evol. 45: 232–237. 31) Watada, Katoh, Yoshioka & Miyake, unpublished. 32) Goto et al. 2004 Genes Genet. Syst. 19–26. 33) Dai, Lu, Lv, Chen, Cheng & Zhang, unpublished. 34) Kumar et al. 2012 Bioinformatics 28: 2685–2686.

Table S2. Total recombination map length for different species of *Drosophila*.

group	species	map length (cM)	reference	mapping function	corrected†
ananassae	*ananassae*	350.1	1	Kosambi	yes
immigrans-tripunctata	*funebris*	895.6	1	Kosambi	yes
immigrans-tripunctata	*mediopunctata*	609.6	1	Kosambi	yes
melanogaster	*mauritiana*	488.5	2	Foss	yes
melanogaster	*melanogaster*	294.9	1	Kosambi	yes
melanogaster	*melanogaster*	287.3	3	N/A*	no
melanogaster	*simulans*	426.2	4	Kosambi	yes
montium	*serrata*	282.3	5	Kosambi	yes
obscura	*persimilis*	605.1	1	Kosambi	yes
obscura	*pseudoobscura*	557.1	1	Kosambi	yes
obscura	*subobscura*	1007.6	1	Kosambi	yes
repleta	*buzzatii*	696.5	1	Kosambi	yes
repleta	*hydei*	655.2	1	Kosambi	yes
virilis	*montana*	632.7	6	Kosambi	yes
virilis	*virilis*	732.3	7	N/A*	no
willistoni	*willistoni*	320.4	8	Kosambi	yes

* High resolution mapping.

† The length of each chromosome was multiplied by $(n + 1) / (n - 1)$, where n is the number of markers per chromosome.

References: 1) Cáceres et al. 1999 Genetics 153: 251–259. 2) True et al. 1996 Genetics 142: 507–523. 3) Comeron et al. 2012 PLoS Genetics 8: e1002905. 4) Barker & Moth 2001 Dros. Inf. Serv. 84: 205–206. 5) Stocker et al. 2012 G3 2: 287–297. 6) Schäfer et al. 2010 J. Evol. Biol. 23: 518–527. 7) Hemmer et al. 2020 Mobile DNA 11: 10. 8) Spassky & Dobzhansky 1950 Heredity 4: 201–215.

Table S3. Phylogenetic signal for each trait analyzed in Fig. 7.

trait	Blomberg's K	P-value	Pagel's λ	$P(\lambda=0)$	$P(\lambda=1)$
nucleotide diversity	0.430	0.928	0.000	1.000	0.018
map length	1.124	0.056	1.155	0.156	0.175
postzygotic RI velocity	0.427	0.957	0.000	1.000	0.017

Methods: The phylogeny shown in Fig. 6a was used for all analyses. Blomberg's K and corresponding P-value were calculated using the R package `phytools` version 1.0-3 (1). The P-value was calculated from 100,000 randomizations.
Pagel's λ was estimated by maximum likelihood using `phytools`. P values were calculated by likelihood ratio tests. The log-likelihood for $\lambda=0$ was obtained from `phytools`; the log-likelihood for $\lambda=1$ was calculated using the R package `geiger` version 2.0.10 (2).

References: 1) Revell 2012 Meth. Ecol. Evol. 3: 217–223. 2) Pennell et al. 2014 Bioinformatics 30: 2216–2218.

Table S4. Asymmetry in synonymous nucleotide diversity (πS) at the *Adh* locus and in postzygotic reproductive isolation (RI) for different species of *Drosophila* .

species 1	species 2	group	πS1	πS2	πS asym	RI asym	RI12*	RI21*	references	notes
arizonae	*mojavensis*	*repleta*	0.032887	0.014165	0.018722	0.334	0.500	0.166		mean of 4 estimates
auraria	*triauraria*	*montium*	0.045728	0.015520	0.030208	1.000	1.000	0.000		
auraria	*subauraria*	*montium*	0.045728	0.011091	0.034637	0.500	1.000	0.500	1	
equinoxialis	*paulistorum*	*willistoni*	0.005489	0.018475	-0.012986	-0.500	0.500	1.000		
lini	*ogumai*	*montium*	0.062418	0.022183	0.040235	0.000	0.500	0.500	1	
lini	*ohnishii*	*montium*	0.062418	0.027833	0.034585	0.000	0.500	0.500	1	
mauritiana	*sechellia*	*melanogaster*	0.002228	0.000000	0.002228	0.000	0.500	0.500		
ogumai	*ohnishii*	*montium*	0.022183	0.027833	-0.005650	0.000	0.500	0.500	1	
paulistorum	*willistoni*	*willistoni*	0.018475	0.018035	0.000440	0.250	1.000	0.750		
persimilis	*pseudoobscura*	*obscura*	0.013317	0.016660	-0.003343	0.000	0.500	0.500		
bogota	*pseudoobscura*	*obscura*	0.004841	0.016660	-0.011819	0.500	0.500	0.000		
recens	*subquinaria*	*quinaria*	0.063736	0.066094	-0.002358	-0.450	0.500	0.950	2	*Adhr* not *Adh*
simulans	*sechellia*	*melanogaster*	0.027407	0.000000	0.027407	0.000	0.500	0.500		
simulans	*mauritiana*	*melanogaster*	0.027407	0.002228	0.025179	0.000	0.500	0.500		
triauraria	*quadraria*	*montium*	0.015520	0.023228	-0.007708	-0.500	0.000	0.500	3	

* 12 = species 1 ♀ ✕ species 2 ♂; 21 = species 2 ♀ ✕ species 1 ♂. Data from Table S2 of Yukilevich 2012 Evolution 66: 1430–1446.

References for estimates of πS: 1) Watada, Katoh, Yoshioka & Miyake, unpublished. 2) Ginsberg et al. 2019 J. Evo. Biol. 32: 1093–1105.
3) Dai, Lu, Lv, Chen, Cheng & Zhang, unpublished. All other estimates of πS are summarized in Supplementary Data 1.