**Host abundance and heterogeneity in infectiousness determine extent of the environmental**

**reservoir**

Nichole A. Laggan[1*], Katy L. Parise[2], J. Paul White[3], Heather M. Kaarakka[3], Jennifer A. Redell[3],

John E. DePue[4], William H. Scullon[5], Joseph Kath[6], Jeffrey T. Foster[2], A. Marm Kilpatrick[7],

Kate E. Langwig[1], Joseph R. Hoyt[1]

[1]Department of Biological Sciences, Virginia Polytechnic Institute, Blacksburg, VA, 24060 USA

[2]Pathogen and Microbiome Institute, Northern Arizona University, Flagstaff, AZ, 86011, USA

[3]Wisconsin Department of Natural Resources, Madison, WI, 53707, USA

[4]Michigan Department of Natural Resources, Baraga, MI 49870, USA

[5]Michigan Department of Natural Resources, Norway, MI 49908, USA

[6]Illinois Department of Natural Resources, Springfield, IL, 62702, USA

[7]Department of Ecology and Evolutionary Biology, University of California, Santa Cruz, CA,

95064, USA

**\*Correspondence to**: Nichole A. Laggan: nicholelaggan@gmail.com

**Abstract**

Environmental pathogen reservoirs exist for many globally important diseases and can fuel disease outbreaks, affect pathogen evolution, and increase the threat of host extinction. Differences in pathogen shedding among hosts can create mosaics of infection risk across landscapes by increasing pathogen contamination in high use areas. However, how the environmental reservoir establishes in multi-host communities and the importance of factors like host-specific infection and abundance in environmental contamination and transmission remain important outstanding questions. Here we examine how *Pseudogymnoascus destructans,* the fungal pathogen that causes white-nose syndrome in bats, invades and establishes in the environment. We quantified dynamic changes in pathogen shedding, infection intensities, host abundance, and the subsequent propagule pressure imposed by each species within the community. We find that the initial establishment of the pathogen reservoir is driven by different species within the community than those that are responsible for maintaining the reservoir over time. Our results also show that highly shedding species do not always contribute the most to pathogen reservoirs. More broadly, we demonstrate how individual host shedding rates scale to influence landscape-level pathogen contamination.

**Introduction**

Emerging infectious diseases threaten efforts to conserve global biodiversity (Daszak et al. 2000, Taylor et al. 2001, Jones et al. 2008, Fisher et al. 2012). In some disease systems, pathogens may survive for long periods of time in the environment in the absence of a living host (Turner et al. 2016, Plummer et al. 2018, Islam et al. 2020). Pathogen persistence in the environment allows for transmission independent of infected hosts, can exacerbate disease impacts, and increase the risk of host extinction (de Castro and Bolker 2005, Mitchell et al. 2008, Almberg et al. 2011, Hoyt et al. 2020). However, pathogen contamination in the environment is not homogenous, rather, variation in the extent of the environmental reservoir is likely driven by a complex process of pathogen shedding from hosts within the community that lead to subsequent transmission events.

Infected hosts can vary in the amount of pathogen they shed into the environment with some hosts producing disproportionately high amounts of pathogen, independent of host contacts (Sheth et al. 2006, Chase-Topping et al. 2008, Lawley et al. 2008, Direnzo et al. 2014). Research has shown that inherent variation in host shedding can be driven by differences in behavior (Godfrey 2013, Rushmore et al. 2013, VanderWaal and Ezenwa 2016), innate susceptibility (Searle et al. 2011, Gervasi et al. 2013), space use (Brooks-Pollock et al. 2014), and infection severity (Lloyd-Smith et al. 2005, Munywoki et al. 2015). In multi-host disease systems, the variation produced through community composition and species abundance play key roles that determine transmission dynamics (Kilpatrick et al. 2006, Paull et al. 2012).

Host abundance within a community can interact with host shedding to moderate transmission from a particular species (Lloyd-Smith et al. 2005, Paull et al. 2012, Kilonzo et al. 2013). Species that have low rates of shedding, but are highly abundant, may contribute more to

transmission than might be expected based on the amount of pathogen they shed (Peterson and

McKenzie 2014, Scheele et al. 2017). Conversely, a species that has high rates of shedding or is

highly infectious, but at low abundance, may contribute less to transmission than other species

(Lloyd-Smith et al. 2005, Kilpatrick et al. 2006). In addition, for some wildlife infectious

diseases, infectiousness, shedding, and impacts are positively correlated (Langwig et al. 2016,

Brannelly et al. 2020) such that hosts that are initially important in transmission suffer from high

impacts and become less important contributors to the environmental reservoir over time.

Heterogeneity in pathogen shedding and how it influences disease dynamics is important for

many disease systems (Sheth et al. 2006, Chase-Topping et al. 2008, Henaux and Samuel 2011,

Brooks-Pollock et al. 2014, Direnzo et al. 2014, Slater et al. 2016), but how differences in host

shedding scale to a community-level and influence the environmental reservoir is rarely linked

together.

White-nose syndrome (WNS) is an emerging infectious disease caused by the fungal

pathogen *Pseudogymnoascus destructans* (Lorch et al. 2011, Warnecke et al. 2012*), that has had*

devastating effects on bat populations (Langwig et al. 2012, Frick et al. 2015, Langwig et al.

2016). White-nose syndrome exhibits seasonal infection dynamics that are driven by the

environmental reservoir and host-pathogen ecology (Langwig et al. 2015a, Hoyt et al. 2021,

Langwig et al. 2021). *Pseudogymnoascus destructans* can persist for long periods of time in the

environment, which results in widespread infection when hosts return to hibernacula

(subterranean sites where bats hibernate in the winter) in the fall (Lorch et al. 2013, Hoyt et al.

2015, Langwig et al. 2015a, Campbell et al. 2019, Hoyt et al. 2020, Hicks et al. 2021). During

this time, susceptible bats become infected or reinfected by *P. destructans* when they come into

contact with the environmental reservoir upon entering hibernacula (Langwig et al. 2015a). Over

the winter hibernation period, *P. destructans* grows into the skin tissue, causing deleterious

physiological changes, including increased arousals from hibernation, weight loss, dehydration,

and ultimately death (Warnecke et al. 2013, Verant et al. 2014, McGuire et al. 2017, Hoyt et al.

2021). During hibernation, susceptible bat species vary greatly in their infection intensities and

three species have suffered declines that exceed 95% (Langwig et al. 2012, Langwig et al. 2016,

Hoyt et al. 2020, Hoyt et al. 2021). While species vary in infection intensities, species abundance

also varies greatly within bat communities (Langwig et al. 2012, Frick et al. 2015), which may

also influence pathogen contamination in the environment.

Contamination of *P. destructans* in the environment increases after the first year of

invasion (Hoyt et al. 2020) and the extent of the environmental reservoir has been linked to

increased pathogen prevalence and increased pathogen loads for bats across the globe (Hoyt et

al. 2020, Hoyt et al. 2021). As a result, bat mortality also increases with higher levels of

environmental contamination (Hoyt et al. 2018, Hoyt et al. 2020, Hicks et al. 2021). However,

how the environmental pathogen reservoir becomes established in these multi-host communities

has not been investigated.

Environmental transmission is an important driver of infectious disease dynamics and

understanding the factors that lead to pathogen establishment in the environment is crucial for

disease control and prevention. Here we use a unique dataset that encompasses the stages of *P.*

*destructans* invasion and establishment to capture pathogen accumulation in the environmental

reservoir across 21 sites in the Midwestern, United States. Using these data, we explored

potential differences in pathogen shedding among species. We also assessed the relationship

between bat infections and the amount of pathogen shed into the environment by each species

present in the community. Because we hypothesized that bat species abundance would play a key

role in site-level contamination and pathogen establishment of the environmental reservoir, we

also explored how differential pathogen shedding among species and their abundance influences

the pathogen pressure within the community.

**Methods**

*Sample collection and quantification*

We quantified *P. destructans* fungal loads on bats and from hibernacula substrate

throughout bat hibernation sites in the Midwestern, United States. Samples were collected from

21 sites in Wisconsin, Illinois, and Michigan over seven years from pathogen invasion to

establishment. Hibernacula were visited twice yearly, once during early hibernation (November

to December) and once during late hibernation (March to April) to capture differences in

infection and environmental contamination at the beginning and end of hibernation. During each

visit, we counted the total number of bats within each site by species (*Eptesicus fuscus, Myotis*

*lucifugus, Myotis septentrionalis* and *Perimyotis subflavus*). We collected epidermal swab

samples from bats within sites to quantify bat infections (fungal loads) and determine infection

prevalence. Samples were collected using previously established protocols that consisted of

rubbing a polyester swab dipped in sterile water over the muzzle and forearm of the bat five

times (Langwig et al. 2015a, Hoyt et al. 2016).

We also collected environmental substrate swabs from hibernacula walls and ceilings

directly under hibernating bats to measure the amount of *P. destructans* shed into the

environment. To capture site-level environmental contamination, *P. destructans* was collected

from the environment in locations more than two meters away from any roosting bats, but in

areas that bats might roost. These environmental substrate samples were taken by swabbing an

area of substrate equal to the length of a bat's forearm (36-40 mm) five times back and forth, as

described previously (Langwig et al. 2015b). We preserved *P. destructans* DNA samples by

storing all swabs in salt preservation buffer (RNAlater; Thermo Fisher Scientific) directly after

collection. DNA was extracted from all samples with a modified Qiagen DNeasy Blood &

Tissue Kit (Langwig et al. 2015b). The presence and quantity of *P. destructans* was determined

by quantitative Polymerase Chain Reaction (qPCR) (Muller et al. 2013).

To verify that fungal loads measured using qPCR accurately reflected viable fungal

spores in the environment, that would be capable of infecting a host, we collected additional

substrate swab samples from a subset of locations that were paired with swabs used for qPCR.

These samples were cultured by streaking the substrate swab across a plate containing Sabouraud

Dextrose Agar treated with chloramphenicol and gentamicin. The plates were stored at 4 °C and

colony forming units (CFU's) of *P. destructans* were quantified within six weeks of initial

inoculation. We paired substrate samples analyzed using qPCR to determine $\log_{10}$ *P. destructans*

loads for comparison with colony forming units obtained from culture samples to validate

viability. There was a significant relationship between quantity of DNA measured through qPCR

and the number of *P. destructans* CFU's (Figure S1). This suggests that qPCR was a valid

method to estimate the extent of *P. destructans* in the environment and supports that qPCR

results are reflective of the number of infectious propagules in the environment and not relic

DNA (Figure S1).

All research was approved through Institutional Animal Care and Use Committee

protocols: Virginia Polytechnic Institute: 17-180; University of California, Santa Cruz:

Kilpm1705; Wisconsin Endangered/Threatened Species Permit 882 & 886; Michigan

Department of Natural Resources permit SC-1651; Illinois Endangered/ Threatened Species

Permit 5015, 2582 and Scientific Collections permit NH20.5888; US Fish and Wildlife Service

Threatened & Endangered Species Permit TE64081B-1.

*Data analysis*

We separated invasion stage into two distinct categories: "invasion" which included the

first year the pathogen arrived, as described previously (Langwig et al. 2015b, Hoyt et al. 2020),

and "establishment" which included the second and third years of *P. destructans* presence in a

site when species declines begin to occur, which corresponds to the epidemic phase as has been

previously noted (Langwig et al. 2015b, Hoyt et al. 2020). We used these stages to capture

pathogen shed into the environment before and after pathogen accumulation in the environment

occurred (invasion and establishment, respectively) and to examine the dynamic changes

between stages of pathogen invasion and establishment.

We first examined the presence and quantity of *P. destructans* on each species and the

amount each species shed into the environment. We used mixed effect models with $\log_{10}$

environmental fungal load collected under each bat as our response variable, bat species

interacting with pathogen invasion stage as our predictor and site as a random effect. We

similarly compared infection intensity among species using the same model as described above,

but with $\log_{10}$ fungal loads on bats as the response variable. We examined differences in

infection prevalence by species using a generalized linear mixed effects model with a binomial

distribution and a logit link with species as our predictor and bat infection prevalence (0|1) as our

response and site as a random effect.

To examine how differences in infection on each bat contributed to contamination of the

environment under each individual, we used a linear mixed effects model to explore the

relationship between infection on each individual bat and pathogen shed into the environment

under each individual. In this analysis, we used paired $\log_{10}$ environmental *P. destructans* loads

under a bat as our response variable with $\log_{10}$ bat fungal loads interacting with species as our

predictor and site as a random effect. We combined the invasion and established stages since the

amount of pathogen shed into the environment was hypothesized to be a product of how infected

the host was, and therefore, comparable across years.

We investigated the role of species abundance in environmental contamination of *P.*

*destructans*. We tested for differences among species abundance within each site by using a

linear mixed effects model with species as our predictor and $\log_{10}$ average abundance during

early hibernation (before over-winter declines occur) as our response and included site as a

random effect for both invasion stages. To explore how species abundance influenced the degree

of pathogen contamination within sites, we used a linear mixed effect model with $\log_{10}$

environmental fungal loads collected > 2m from any bat during late hibernation as our response

variable and $\log_{10}$ average abundance interacting with species identity as our predictors with site

as a random effect.

Finally, we calculated differences in propagule pressure among species by multiplying

infection prevalence by the species abundance within each site to get the number of infected

individuals. We then multiplied the number of infected individuals by the average amount of

fungal spores (conidia) shed into the environment by each species in each site. We used a

generalized linear mixed effect model with a negative binomial distribution to account for

dispersed counts with an interaction between invasion stage and species as our predictor and

propagule pressure as our response. We performed this analysis for both invasion and

establishment during late hibernation to investigate how pathogen pressure may differ across

stages of pathogen invasion when bats are heavily shedding into the environment. To assess if

propagule pressure was a suitable metric for pathogen invasion of the environmental reservoir,

we examined the relationship between site-level environmental contamination, as described

above, and sum propagule pressure for each site using a linear mixed effects model with site

included as a random effect. All analyses were conducted in R v.4.2.1 (R Core Team 2022).

Mixed-effects models were run using the package "lme4" (Bates et al. 2015) except for analyses

using a negative binomial distribution which were ran using "glmmTMB" (Brooks et al. 2017)

***Results***

We found that pathogen shedding (the amount of pathogen under bats) varied among the

four species (Figure 1). During initial pathogen invasion into bat communities, we found that on

average individual *M. septentrionalis* (Figure 1A; Table S1; intercept =-3.79 ± 0.23) contributed

more pathogen into the environment than *E. fuscus* (coeff = -0.77 ± 0.37, P = 0.04) and *M.*

*lucifugus* (coeff= -0.58 ± 0.27, P = 0.03), but did not differ statistically from individual *P.*

*subflavus* (coeff =-0.21 ± 0.38, P = 0.58). In the established stage (Figure 1B), we found support

for higher shedding in *M. lucifugus* (intercept = -3.38 ± 0.11) than *E. fuscus* (coeff = -0.80 ±

0.14, P < 0.0001) and *P. subflavus* (coeff = -0.62 ± 0.13, P < 0.0001), but similar shedding to *M.*

*septentrionalis* (coeff = -0.38 ± 0.24, P = 0.11).

Our results showed consistent support that host infection intensity predicted the amount

of pathogen shed into the environment for all species (Figure 2C; Table S4; host infection and

shedding relationship: *M. lucifugus* slope = 0.38 ± 0.05, P < 0.001; *E. fuscus* slope = 0.33 ± 0.07,

P < 0.00001; *M. septentrionalis* slope = 0.41 ± 0.10, P = 0.00003; *P. subflavus* slope = 0.33 ±

0.08, P = 0.00006). Importantly, we found no support for differences among species in the slope

between environmental pathogen shedding and host infection level (Figure 2C; Table S4; all P >

0.05), suggesting that differences in shedding were primarily driven by the extent of infection on

individual bats. *Eptesicus fuscus* (intercept = -3.37 ± 0.16) had significantly lower infection

levels than all other species within the community (Figure 2B; Table S3; *M. lucifugus* coeff =

1.38 ± 0.15, P < 0.0001; *M. septentrionalis* coeff = 1.08 ± 0.21 , P < 0.0001; *P. subflavus* coeff =

1.30 ± 0.17, P < 0.0001) In addition, we found that *M. lucifugus* (intercept = 2.19 ± 0.27) had the

highest infection prevalence, with 87% of hosts infected on average across the invasion and

established stages (Figure 2A; Table S2; *E. fuscus* coeff = -0.79 ± 0.31, P < 0.01; *M.*

*septentrionalis* coeff = -0.83 ± 0.35, P < 0.05; *P. subflavus* coeff = -1.10 ± 0.24, P < 0.0001).

In addition to differences in infection intensity, species also varied in abundance, with *M.*

*lucifugus* (intercept = 1.63 ± 0.19) being the most abundant species across sites, with an average

population size of 39.81 ± 7.76 before declines from WNS (invasion) (Figure 3; Table S5; *E.*

*fuscus* coeff = -0.65 ± 0.24, P < 0.01; *M. septentrionalis* coeff = -0.60 ± 0.21, P < 0.01; *P.*

*subflavus* coeff = -0.41 ± 0.23, P < 0.08). During the established stage, while population declines

were occurring, *M. lucifugus* (intercept = 1.28 ± 0.13) remained the most abundant species

within the community with an average population size of 18.20 ± 7.41 (Figure 3; Table S6; *E.*

*fuscus* coeff = -0.54 ± 0.16, P < 0.001; *M. septentrionalis* coeff = -0.76 ± 0.14, P < 0.0001; *P.*

*subflavus* coeff = -0.38 ± 0.15, P < 0.01). *M. septentrionalis* had the lowest abundance in the

community with an average colony size of 11.22 ± 5.50 individuals before disease impacts and

declined to an average of 3.52 ± 4.12 individuals in a site during the established stage (Figure 3;

Table S5-7). Furthermore, host abundance was important in influencing the degree of site-level

environmental pathogen contamination (Figure S2; Table S7; *M. lucifugus* slope = 0.40 ± 0.14, P

= 0.03).

Finally, when we combined infection prevalence, species abundance, and pathogen

shedding into the metric of propagule pressure, we found that during the first year of invasion,

*M. septentrionalis* (Figure4A; Table S8; intercept = 7.85 ± 0.66) had higher propagule pressure

than *P. subflavus* (coeff = -2.81 ± 1.01, P < 0.01) and *E. fuscus* (-4.28 ± 0.85, P < 0.0001), but

not *M. lucifugus* (coeff = 0.43 ± 0.73, P = 0.55). In years following invasion, once *P. destructans*

is established, *M. lucifugus* (Figure 4B; intercept = 11.08 ± 0.56) had consistently higher

propagule pressure and contributed more pathogen to the environmental reservoir than all other

species (Figure 4B; *M. septentrionalis* coeff = -4.14 ± 0.96, P < 0.0001; *E. fuscus* coeff = -4.18 ±

0.68, P < 0.0001; *P. subflavus* coeff = -2.53 ± 0.68, P = 0.002). *Myotis septentrionalis* (Figure

4B; intercept = 6.94 ± 0.95) no longer contributed more than *P. subflavus* (coeff = 1.62 ± 1.03, P

= 0.11) and *E. fuscus* (coeff = -0.04 ± 1.02, P = 0.97) due to their rarity following disease-

induced declines (Figure 4). Mean environmental contamination in areas > 2m from bats

increased with total propagule pressure at a site (summed propagule pressure among species)

(Figure 4C; relationship between environmental contamination and propagule pressure intercept

= -5.48 ± 0.14, slope = 0.12, P = 0.002).

***Discussion***

Host heterogeneity in pathogen shedding and subsequent environmental transmission can

have dramatic impacts on community-level disease outcomes. Our findings demonstrate how

host heterogeneity in infection intensity can lead to increased pathogen shedding, which is

consistent with research in other systems (Lloyd-Smith et al. 2005, Matthews et al. 2006, Chase-

Topping et al. 2008, Direnzo et al. 2014, Munywoki et al. 2015, Maguire et al. 2016,

VanderWaal and Ezenwa 2016). In addition, our study demonstrates that host heterogeneities in

abundance and infection interact to influence the extent of the environmental reservoir and the

contribution of species, which varied over time. We identified a shift in the species responsible

for the greatest environmental reservoir contamination between disease stages, suggesting that

reservoir contamination is a dynamic process that varies over time.

Our findings suggest that the observed heterogeneities in pathogen shedding are primarily

due to variation in infection intensities among species within the community. The relationship

between individual host shedding into the environment and host infection intensity did not differ

among species, suggesting that individuals of different species with the same infection intensity

are equally efficient at depositing pathogen into the environment. Instead, the severity of

infection, which varied among species, determined how much pathogen is shed per individual.

Some species such as *E. fuscus* had low level infections, while the other species such as *M.*

*septentrionalis* and *M. lucifugus* were much more infected and contributed more pathogen to the

environmental reservoir (Figure 2B).

Disease-caused declines in several species (Figure 3) resulted in large changes in species

abundance and community composition. We found evidence that species abundance predicted

site-level contamination (Figure S2), and this supported the need to account for both species'

infection and abundance when determining the contribution to the environmental reservoir. To

combine these factors, we used propagule pressure or "introduction effort" which is a

fundamental ecological metric that determines whether a biological invasion will be successful in

reaching establishment (Lockwood et al. 2005), and is especially applicable to emerging

infectious diseases. Our results indicated that during the initial invasion of *P. destructans*, both

*M. lucifugus* and *M. septentrionalis* had higher propagule pressure than the other two species

(Figure 4A), which was not apparent through examination of only pathogen shedding. The high

infection levels of *M. septentrionalis* resulted in a large contribution to the initial establishment

of the environmental reservoir despite their low relative abundance (Figure 2B; Figure 3).

However, in subsequent years, as this species declined precipitously (Figure 3), their

contribution was greatly reduced and equivalent to less infected species (e.g. *E. fuscus* and *P.*

*subflavus*, Figure 4B).

Our results suggest that host abundance, shedding, and infection prevalence are

collectively required to describe pathogen contamination in the environment. We found that

environmental contamination in the sampled locations that were not associated with bats

increased as propagule pressure increased (Figure 4C), suggesting that propagule pressure is an

accurate metric for measuring species contributions to the environmental reservoir. For example,

while *E. fuscus*, *M. septentrionalis*, and *P. subflavus* eventually contributed similar propagule

pressures during disease establishment, this was likely driven by different factors that need to be

considered when evaluating their influence on the reservoir. *Eptesicus fuscus* had low infection

intensity but had moderate prevalence (Figure 2A-B) and moderate abundance (Figure 3) which

elevated its contribution. *Perimyotis subflavus* had high infection levels, but reduced prevalence

(Figure 2A-B) compared to the other highly infected species and was only present in moderate

abundances (Figure 3). Finally, *M. septentrionalis* was highly infected (Figure2B), but at low

abundance across communities (Figure 3), which reduced its overall importance in years

following initial pathogen invasion. *Myotis lucifugus* maintained high propagule pressure

through the invasion and establishment of *P. destructans* (Figure 4A-B), suggesting that the

presence of *M. lucifugus* within a community will result in rapid establishment of *P. destructans*

in the environment and maintenance of environmental contamination within a site over time. Our

results highlight how both species infection intensity and abundance must be considered to

evaluate the influence of pathogen shedding on environmental reservoir establishment,

subsequent environmental transmission and how this can change dynamically across disease

stages.

Understanding environmental reservoir dynamics is crucial for many globally important

disease systems. Discerning how variation across species contributes to maintenance of the

environmental reservoir is important in determining the extent of epidemics, predicting long-

term impacts on host communities, and developing control strategies. We demonstrate that the

establishment and maintenance of the environmental reservoir is strongly influenced by both

heterogeneity in pathogen infection and species abundance. Evaluating these effects together

through the metric of propagule pressure allowed us to capture which species within the

community contributed to pathogen invasion success and ultimately the maintenance of indirect

transmission which is an important driver of infection and mortality (Hoyt et al. 2018). Broadly,

our results demonstrate that multiple factors of species heterogeneity can scale to influence

environmental reservoir dynamics within communities.

**Acknowledgements:**

The research was funded by the joint NSF-NIH-NIFA Ecology and Evolution of Infectious

Disease award DEB-1911853, NSF DEB-1115895, USFWS (F17AP00591), and by the NSF

GRFP. We would like to acknowledge landowners for access to sites, and M.J. Kailing for

assisting culture collection and B. Heeringa and D. Kirk for field assistance.

**Author Contributions:**

NAL and JRH wrote the original draft of the manuscript. JRH, KEL, and AMK designed

methodology; NAL, JRH, KEL, AMK, JPW, HMK, JAR, JED, WHS, and JK collected the data;

NAL conducted the laboratory experiment; JTF and KLP supervised and performed sample

testing; NAL analyzed the data with assistance from KEL and JRH; All authors contributed

critically to draft revision.

**Conflict of Interest:** The authors declare no conflicts of interest

**Figure captions:**

**Figure 1**: **Differences in pathogen shedding among species during reservoir invading and established stages.** $\log_{10}$ *P. destructans* environmental loads (ng DNA) among species during pathogen invasion (A) and establishment (B) in late hibernation. Each point represents the $\log_{10}$ environmental *P. destructans* loads from under an individual bat. Black points represent the estimated mean and bars indicate ± standard error for each species.

**Figure 2: Bat infection prevalence, intensity, and the relationship between host infection and pathogen shedding.** For all panels, bat species are displayed by color. Black points indicate mean bat infections by species and bars represent ± standard error. (A) Bat infection prevalence for each species during pathogen invasion in late hibernation. (B) Bat $\log_{10}$ *P. destructans* loads (ng DNA) for each species during pathogen invasion in late hibernation. (C) The relationship between $\log_{10}$ *P. destructans* loads (ng DNA) on an individual bat and the amount of $\log_{10}$ *P. destructans* (ng DNA) in the environment directly underneath each individual during pathogen invasion and establishment in late hibernation. The gray solid line shows the 1:1 line, and points along this line would indicate that the amount of *P. destructans* on bats was equivalent to the amount shed into the environment. Solid colored lines indicate statistical support for a positive relationship. Each individual bat is represented by a point.

**Figure 3. Changes in host abundance between invading and established years.** Species within the community are differentiated by color and points represent a species population at an individual site. Species abundance in the invasion (light gray box) and established years (dark gray box) in late hibernation within each site. Circular points indicate the mean abundance for each year and bars denote ± standard error.

**Figure 4**. **Differences in propagule pressure among species during invading and established stages and the relationship between propagule pressure and environmental contamination.** (A-B) Propagule pressure (# of pathogen particles) by species during late hibernation in (A) invading and (B) established stages. Colored points represent populations of species within a site, black points represent the mean and bars indicate ± standard error for each species. Point size is weighted by species population abundance. (C) Relationship between $\log_{10}$ propagule

pressure and $\log_{10}$ mean site-level contamination. Site-level contamination was assessed in

samples > 2m away from roosting bats within sites during late hibernation in the established

stage. Points indicate the sum propagule pressure at a site for each year and shape denotes

invasion stage. The total environmental contamination increased as propagule pressure increased

within a site (Environmental contamination: intercept = -5.48 ± 0.14, slope = 0.12, relationship

between environmental contamination and propagule pressure P = 0.002).

**Figure 1**

**Figure 2**

**Figure 3**

 **Figure 4**

## References

Almberg, E. S., P. C. Cross, C. J. Johnson, D. M. Heisey, and B. J. Richards. 2011. Modeling routes of chronic wasting disease transmission: environmental prion persistence promotes deer population decline and extinction. PLoS One **6**:e19896.

Bates, D., M. Mächler, B. M. Bolker, and S. C. Walker. 2015. Fitting linear mixed-effects models using lme4. Journal of Statistical Software **67**:1-48.

Brannelly, L. A., H. I. McCallum, L. F. Grogan, C. J. Briggs, M. P. Ribas, M. Hollanders, T. Sasso, M. Familiar López, D. A. Newell, and A. M. Kilpatrick. 2020. Mechanisms underlying host persistence following amphibian disease emergence determine appropriate management strategies. Ecology Letters.

Brooks, M. E., K. Kristensen, K. J. van Benthem, A. Magnusson, C. W. Berg, A. Nielson, H. J. Skaug, M. Maechler, and B. M. Bolker. 2017. glmmTMB Balances Speed and Flexibility Among Packages for Zero-inflated Generalized Linear Mixed Modeling. The R Journal **9**:378-400.

Brooks-Pollock, E., G. O. Roberts, and M. J. Keeling. 2014. A dynamic model of bovine tuberculosis spread and control in Great Britain. Nature **511**:228-231.

Campbell, L. J., D. P. Walsh, D. S. Blehert, and J. M. Lorch. 2019. Long-term survival of Pseudogymnoascus destructans at elevated temperatures. Journal of Wildlife Diseases.

Chase-Topping, M., D. Gally, C. Low, L. Matthews, and M. Woolhouse. 2008. Super-shedding and the link between human infection and livestock carriage of Escherichia coli O157. Nat Rev Microbiol **6**:904-912.

Daszak, P., A. A. Cunningham, and A. D. Hyatt. 2000. Emerging infectious diseases of wildlife - threats to biodiversity and human health. Science **287**:443.

de Castro, F., and B. Bolker. 2005. Mechanisms of disease-induced extinction. Ecology Letters **8**:117-126.

Direnzo, G. V., P. F. Langhammer, K. R. Zamudio, and K. R. Lips. 2014. Fungal infection intensity and zoospore output of Atelopus zeteki, a potential acute chytrid supershedder. PLoS One **9**:e93356.

Fisher, M. C., D. A. Henk, C. J. Briggs, J. S. Brownstein, L. C. Madoff, S. L. McCraw, and S. J. Gurr. 2012. Emerging fungal threats to animal, plant and ecosystem health. Nature **484**:186-194.

Frick, W. F., S. J. Puechmaille, J. R. Hoyt, B. A. Nickel, K. E. Langwig, J. T. Foster, K. E. Barlow, T. Bartonicka, D. Feller, A. J. Haarsma, C. Herzog, I. Horacek, J. van der Kooij, B. Mulkens, B. Petrov, R. Reynolds, L. Rodrigues, C. W. Stihler, G. G. Turner, and A. M. Kilpatrick. 2015. Disease alters macroecological patterns of North American bats. Global Ecology and Biogeography **24**:741-749.

Gervasi, S. S., J. Urbina, J. Hua, T. Chestnut, R. A. Relyea, and A. R. Blaustein. 2013. Experimental evidence for American bullfrog (Lithobates catesbeianus) susceptibility to chytrid fungus (Batrachochytrium dendrobatidis). Ecohealth **10**:166-171.

Godfrey, S. S. 2013. Networks and the ecology of parasite transmission: A framework for wildlife parasitology. Int J Parasitol Parasites Wildl **2**:235-245.

Henaux, V., and M. D. Samuel. 2011. Avian influenza shedding patterns in waterfowl: implications for surveillance, environmental transmission, and disease spread. J Wildl Dis **47**:566-578.

Hicks, A. C., S. Darling, J. Flewelling, R. von Linden, C. U. Meteyer, D. Redell, J. P. White, J. Redell, R. Smith, D. Blehert, N. Rayman, J. R. Hoyt, J. C. Okoniewski, and K. E. Langwig. 2021. Environmental transmission of Pseudogymnoascus destructans to hibernating little brown bats. biorxiv.

Hoyt, J. R., A. M. Kilpatrick, and K. E. Langwig. 2021. Ecology and impacts of white-nose syndrome on bats. Nature Reviews Microbiology:1-15.

Hoyt, J. R., K. E. Langwig, J. Okoniewski, W. F. Frick, W. B. Stone, and A. M. Kilpatrick. 2015. Long-term persistence of Pseudogymnoascus destructans, the causative agent of white-nose syndrome, in the absence of bats. Ecohealth **12**:330-333.

Hoyt, J. R., K. E. Langwig, K. Sun, G. Lu, K. L. Parise, T. Jiang, W. F. Frick, J. T. Foster, J. Feng, and A. M. Kilpatrick. 2016. Host persistence or extinction from emerging infectious disease: insights from white-nose syndrome in endemic and invading regions. Proceedings of the Royal Society B: Biological Sciences **283**.

Hoyt, J. R., K. E. Langwig, K. Sun, K. L. Parise, A. Li, Y. Wang, X. Huang, L. Worledge, H. Miller, J. P. White, H. M. Kaarakka, J. A. Redell, T. Görföl, S. A. Boldogh, D. Fukui, M. Sakuyama, S. Yachimori, A. Sato, M. Dalannast, A. Jargalsaikhan, N. Batbayar, Y. Yovel, E. Amichai, I. Natradze, W. F. Frick, J. T. Foster, J. Feng, and A. M. Kilpatrick. 2020. Environmental reservoir dynamics predict global infection patterns and population impacts for the fungal disease white-nose syndrome. Proceedings of the National Academy of Sciences:201914794.

Hoyt, J. R., K. E. Langwig, J. P. White, H. M. Kaarakka, J. A. Redell, A. Kurta, J. E. DePue, W. H. Scullon, K. L. Parise, and J. T. Foster. 2018. Cryptic connections illuminate pathogen transmission within community networks. Nature:1.

Islam, M. S., M. H. Zaman, M. S. Islam, N. Ahmed, and J. D. Clemens. 2020. Environmental reservoirs of Vibrio cholerae. Vaccine **38 Suppl 1**:A52-A62.

Jones, K. E., N. G. Patel, M. A. Levy, A. Storeygard, D. Balk, J. L. Gittleman, and P. Daszak. 2008. Global trends in emerging infectious diseases. Nature **451**:990-U994.

Kilonzo, C., X. Li, E. J. Vivas, M. T. Jay-Russell, K. L. Fernandez, and E. R. Atwill. 2013. Fecal shedding of zoonotic food-borne pathogens by wild rodents in a major agricultural region of the central California coast. Appl Environ Microbiol **79**:6337-6344.

Kilpatrick, A. M., P. Daszak, M. J. Jones, P. P. Marra, and L. D. Kramer. 2006. Host heterogeneity dominates West Nile virus transmission. Proceedings of the Royal Society B-Biological Sciences **273**:2327-2333.

Langwig, K. E., W. F. Frick, J. T. Bried, A. C. Hicks, T. H. Kunz, and A. M. Kilpatrick. 2012. Sociality, density-dependence and microclimates determine the persistence of populations suffering from a novel fungal disease, white-nose syndrome. Ecol Lett **15**.

Langwig, K. E., W. F. Frick, J. R. Hoyt, K. L. Parise, K. P. Drees, T. H. Kunz, J. T. Foster, and A. M. Kilpatrick. 2016. Drivers of variation in species impacts for a multi-host fungal disease of bats. Philosophical Transactions of the Royal Society B: Biological Sciences **10.1098/rstb.2015.0456**.

Langwig, K. E., W. F. Frick, R. Reynolds, K. L. Parise, K. P. Drees, J. R. Hoyt, T. L. Cheng, T. H. Kunz, J. T. Foster, and A. M. Kilpatrick. 2015a. Host and pathogen ecology drive the seasonal dynamics of a fungal disease, white-nose syndrome. Proceedings of the Royal Society B: Biological Sciences **282**:20142335.

Langwig, K. E., J. R. Hoyt, K. L. Parise, J. Kath, D. Kirk, W. F. Frick, J. T. Foster, and A. M. Kilpatrick. 2015b. Invasion Dynamics of White-Nose Syndrome Fungus, Midwestern United States, 2012-2014. Emerging Infectious Disease **21**:1023-1026.

Langwig, K. E., J. P. White, K. L. Parise, H. M. Kaarakka, J. A. Redell, J. E. DePue, W. H. Scullon, J. T. Foster, A. M. Kilpatrick, and J. R. Hoyt. 2021. Mobility and infectiousness in the spatial spread of an emerging fungal pathogen. J Anim Ecol **90**:1134-1141.

Lawley, T. D., D. M. Bouley, Y. E. Hoy, C. Gerke, D. A. Relman, and D. M. Monack. 2008. Host transmission of Salmonella enterica serovar Typhimurium is controlled by virulence factors and indigenous intestinal microbiota. Infect Immun **76**:403-416.

Lloyd-Smith, J. O., S. J. Schreiber, P. E. Kopp, and W. M. Getz. 2005. Superspreading and the effect of individual variation on disease emergence. Nature **438**:355-359.

Lockwood, J. L., P. Cassey, and T. Blackburn. 2005. The role of propagule pressure in explaining species invasions. Trends Ecol Evol **20**:223-228.

Lorch, J. M., C. U. Meteyer, M. J. Behr, J. G. Boyles, P. M. Cryan, A. C. Hicks, A. E. Ballmann, J. T. H.
Coleman, D. N. Redell, D. M. Reeder, and D. S. Blehert. 2011. Experimental infection of bats with
Geomyces destructans causes white-nose syndrome. Nature **480**:376-378.
Lorch, J. M., L. K. Muller, R. E. Russell, M. O'Connor, D. L. Lindner, and D. S. Blehert. 2013. Distribution
and environmental persistence of the causative agent of white-nose syndrome, Geomyces
destructans, in bat hibernacula of the eastern United States. Applied and Environmental
Microbiology **79**:1293-1301.
Maguire, C., G. V. DiRenzo, T. S. Tunstall, C. R. Muletz-Wolz, K. R. Zamudio, and K. R. Lips. 2016. Dead or
alive? Viability of chytrid zoospores shed from live amphibian hosts. Diseases of Aquatic
Organisms **199**:179-187.
Matthews, L., J. C. Low, D. L. Gally, M. C. Pearce, D. J. Mellor, J. A. Heesterbeek, M. Chase-Topping, S. W.
Naylor, D. J. Shaw, S. W. Reid, G. J. Gunn, and M. E. Woolhouse. 2006. Heterogeneous shedding
of Escherichia coli O157 in cattle and its implications for control. Proc Natl Acad Sci U S A
**103**:547-552.
McGuire, L. P., H. W. Mayberry, and C. K. Willis. 2017. White-nose syndrome increases torpid metabolic
rate and evaporative water loss in hibernating bats. American Journal of Physiology-Regulatory,
Integrative and Comparative Physiology **313**:R680-R686.
Mitchell, K. M., T. S. Churcher, T. W. Garner, and M. C. Fisher. 2008. Persistence of the emerging
pathogen Batrachochytrium dendrobatidis outside the amphibian host greatly increases the
probability of host extinction. Proc Biol Sci **275**:329-334.
Muller, L. K., J. M. Lorch, D. L. Lindner, M. O'Connor, A. Gargas, and D. S. Blehert. 2013. Bat white-nose
syndrome: a real-time TaqMan polymerase chain reaction test targeting the intergenic spacer
region of Geomyces destructans. Mycologia **105**:253-259.
Munywoki, P. K., D. C. Koech, C. N. Agoti, N. Kibirige, J. Kipkoech, P. A. Cane, G. F. Medley, and D. J.
Nokes. 2015. Influence of age, severity of infection, and co-infection on the duration of
respiratory syncytial virus (RSV) shedding. Epidemiol Infect **143**:804-812.
Paull, S. H., S. Song, K. M. McClure, L. C. Sackett, A. M. Kilpatrick, and P. T. J. Johnson. 2012. From
superspreaders to disease hotspots: linking transmission across hosts and space. Frontiers in
Ecology and the Environment **10**:75-82.
Peterson, A. C., and V. J. McKenzie. 2014. Investigating differences across host species and scales to
explain the distribution of the amphibian pathogen Batrachochytrium dendrobatidis. PLoS One
**9**:e107441.
Plummer, I. H., C. J. Johnson, A. R. Chesney, J. A. Pedersen, and M. D. Samuel. 2018. Mineral licks as
environmental reservoirs of chronic wasting disease prions. PLoS One **13**:e0196745.
Rushmore, J., D. Caillaud, L. Matamba, R. M. Stumpf, S. P. Borgatti, and S. Altizer. 2013. Social network
analysis of wild chimpanzees provides insights for predicting infectious disease risk. J Anim Ecol
**82**:976-986.
Scheele, B. C., D. A. Hunter, L. A. Brannelly, L. F. Skerratt, and D. A. Driscoll. 2017. Reservoir-host
amplification of disease impact in an endangered amphibian. Conserv Biol **31**:592-600.
Searle, C. L., S. S. Gervasi, J. Hua, J. I. Hammond, R. A. Relyea, D. H. Olson, and A. R. Blaustein. 2011.
Differential Host Susceptibility to Batrachochytrium dendrobatidis, an Emerging Amphibian
Pathogen. Conservation Biology **25**:965-974.
Sheth, P. M., A. Danesh, A. Sheung, A. Rebbapragada, K. Shahabi, C. Kovacs, R. Halpenny, D. Tilley, T.
Mazzulli, K. MacDonald, D. Kelvin, and R. Kaul. 2006. Disproportionately high semen shedding of
HIV is associated with compartmentalized cytomegalovirus reactivation. J Infect Dis **193**:45-48.
Slater, N., R. M. Mitchell, R. H. Whitlock, T. Fyock, A. K. Pradhan, E. Knupfer, Y. H. Schukken, and Y.
Louzoun. 2016. Impact of the shedding level on transmission of persistent infections in
Mycobacterium avium subspecies paratuberculosis (MAP). Vet Res **47**:38.

Taylor, L. H., S. M. Latham, and M. E. Woolhouse. 2001. Risk factors for human disease emergence.
Philos Trans R Soc Lond B Biol Sci **356**:983-989.
Turner, W. C., K. L. Kausrud, W. Beyer, W. R. Easterday, Z. R. Barandongo, E. Blaschke, C. C. Cloete, J.
Lazak, M. N. Van Ert, and H. H. Ganz. 2016. Lethal exposure: An integrated approach to
pathogen transmission via environmental reservoirs. Scientific reports **6**:27311.
VanderWaal, K. L., and V. O. Ezenwa. 2016. Heterogeneity in pathogen transmission: mechanisms and
methodology. Funct. Ecol. **30**:1606-1622.
Verant, M. L., M. U. Carol, J. R. Speakman, P. M. Cryan, J. M. Lorch, and D. S. Blehert. 2014. White-nose
syndrome initiates a cascade of physiologic disturbances in the hibernating bat host. BMC
physiology **14**:10.
Warnecke, L., J. M. Turner, T. K. Bollinger, J. M. Lorch, V. Misra, P. M. Cryan, G. Wibbelt, D. S. Blehert,
and C. K. R. Willis. 2012. Inoculation of bats with European Geomyces destructans supports the
novel pathogen hypothesis for the origin of white-nose syndrome. Proceedings of the National
Academy of Sciences of the United States of America **109**:6999-7003.
Warnecke, L., J. M. Turner, T. K. Bollinger, V. Misra, P. M. Cryan, D. S. Blehert, G. Wibbelt, and C. K. R.
Willis. 2013. Pathophysiology of white-nose syndrome in bats: a mechanistic model linking wing
damage to mortality. Biology Letters **9**.