Cis-regulatory variation in relation to sex and sexual dimorphism in *Drosophila melanogaster*

Prashastha Mishra[1], Tania S. Barrera[1], Karl Grieshop[1,2,3] and Aneil F. Agrawal[1]

Affiliations:
[1]Department of Ecology and Evolutionary Biology, University of Toronto, Toronto, ON M5S 3B2, Canada
[2]Department of Molecular Biosciences, The Wenner-Gren Institute, Stockholm University, Stockholm SE-10691, Sweden
[3]School of Biological Sciences, University of East Anglia, Norwich Research Park, Norwich, NR4 7TJ, United Kingdom

ORCIDs:
https://orcid.org/0000-0001-8925-5066 (KG)

Keywords:
sexual dimorphism, sex biased gene expression, allelic imbalance

Short title:
Sex differences in *cis*-regulatory variation

Author Contributions: PM, KG, TB performed dissections and RNA extractions. PM performed bioinformatic data processing. PM and AA did the data analysis. PM, KG, and AA did the writing of the manuscript. AA conceived the experiment.

Data availability: Upon acceptance, RNAseq data will be submitted to the Sequence Read Archive (NIH) and analysis scripts will be made available via GitHub.

Conflict of Interest: Authors have no conflicts of interest.

Acknowledgements:
We thank Ina Anreiter for help with dissections, Baharul Choudhury for help with RNA extractions, Santiago Sánchez-Ramírez for bioinformatics advice, and Stephen Wright for helpful discussion. We thank John Pool and Yuheng Huang (U Madison) for providing the SP159N line.

Funding:
This work was supported by the Natural Sciences and Engineering Research Council of Canada (AFA).

Abstract

Much of sexual dimorphism is likely due to sex-biased gene expression, which results from differential regulation of a genome that is largely shared between males and females. Here we use allele-specific expression to explore *cis*-regulatory variation in *Drosophila melanogaster* in relation to sex. We develop a Bayesian framework to infer the transcriptome-wide joint distribution of *cis*-regulatory effects across the sexes. We use this approach to quantify transcriptome-wide sex differences in *cis*-regulatory effects as well as examine patterns of *cis*-regulatory variation with respect to two other levels of variation in sexual dimorphism: (i) across genes varying in their degree of sex-biased expression, and (ii) among tissues that vary in their degree of dimorphism (e.g., relatively low dimorphism in heads vs high dimorphism in gonads). We uncover evidence of widespread *cis*-regulatory variation in all tissues examined, with female-biased genes being especially enriched for this variation. A sizeable proportion of *cis*-regulatory variation is inferred to have sex-specific effects, with sex-dependent *cis* effects being much more frequent in gonads than in heads. Finally, we detect some genes with reversed allelic imbalance between the sexes. Such variants could provide a mechanism for sex-specific dominance reversals, a phenomenon important for sexually antagonistic balancing selection.

Introduction

Sexual dimorphism is a universal feature of sexually reproducing species, with males and females differing in their appearance, physiology, life history and behaviour (Fairbairn et al. 2007). Males and females carry similar genetic material (other than differentiated sex chromosomes in some species), and these extensive differences in phenotype are largely the result of differential expression of a shared genome (Parsch and Ellegren 2013).

Transcriptional gene regulation is central to the evolution of sex-biased gene expression. Regulation of gene expression can occur through *cis*-acting regulatory elements linked to their target alleles or through *trans*-acting factors. *Cis*-regulatory variants have been reported to be enriched between species relative to their frequency within species in *Drosophila* (Wittkopp et al. 2004; Wittkopp et al. 2008), yeast (Metzger et al. 2017) and mice (Goncalves et al. 2012), among others. This suggests that *cis*-regulatory factors are preferentially used to respond to natural selection for divergent expression. Additionally, *cis*-acting variants are more likely to have additive effects on expression than *trans*-acting variants (Wray 2007; Lemos et al. 2008), which may influence their ability to respond to selection. In contrast, *trans* factors affect expression in multiple genes, and mutations in *trans* may be constrained by pleiotropic side effects (Stern 2000). Lastly, at a practical level, *cis* variants are easier to study. Here, we focus entirely on regulatory variation in *cis*.

Studies indicate that *cis*-regulatory variation is highly pervasive in a wide range of taxa, including humans (Pastinen and Hudson 2004; Stranger et al. 2012), yeast (Kita et al. 2017), mice (Campbell et al. 2008) and *Arabidopsis* (Cubillos et al. 2014). As in other species, *cis*-regulatory variation is common within *Drosophila melanogaster*, with a higher abundance of variation being detected in *cis* than in *trans* (Genissel et al. 2007; Gruber and Long 2009; Osada et al. 2017). Here we examine *cis*-regulatory variation in *D. melanogaster* by measuring differential expression between alleles, termed "allelic imbalance" (henceforth, AI). We investigate its relationship to sexual dimorphism in three ways.

First, we examine whether *cis*-regulatory variation is more common at genes with dimorphic expression (i.e., sex-biased genes). *Cis*-regulatory variation is expected to be enriched in loci undergoing sexual conflict over gene expression. Under the some conditions, sexual conflict can generate balancing selection that stably maintains polymorphisms (Kidwell et al. 1977; Connallon and Clark 2014). If sex-biased genes are more likely to experience sexual conflict than unbiased genes, then one might predict *cis*-regulatory variants to be more common at sex-biased genes. However, even though sexually divergent selection in the past may be responsible for creating the dimorphic expression observed in the present, it

is unclear to what extent sexual conflict may persist in genes with sex-biased expression. Sex-biased

genes could represent instances of resolved or ongoing conflicts (Bonduriansky and Chenoweth 2009;

Rowe et al. 2018). Similarly, unbiased genes could either lack sexual conflict entirely or simply lack the

appropriate variation to evolve sex-biased expression to mitigate conflict (Parsch and Ellegren 2013).

Based on population genetic analyses of human and fly data, Cheng & Kirkpatrick (2016) concluded that

sexual conflict is the most intense in moderately sex-biased genes; this may lead one to predict *cis*-

regulatory variation being most common in genes with an intermediate sex bias. Two previous studies of

cis-regulatory variation in *D. melanogaster* found different patterns from one another and neither matched

this prediction (Osada et al. 2017; Puixeu et al. 2023).

Regulatory variants that affect expression similarly in both sexes would not allow for the resolution of

sexual conflict via sex-biased gene expression. Thus, our second goal is to examine the extent to which

cis-regulatory effects differ between the sexes, measured as 'sex-dependent' allelic imbalance. In terms of

the transcriptome-wide joint distribution of *cis*-regulatory effects across the sexes, we can consider sex

differences from two related perspectives: the frequency with which AI differs between sexes or the

intersexual correlation in AI. The latter harkens to the intersexual genetic correlation in expression (often

represented as r_{MF}), which governs the extent to which expression can evolve independently between the

sexes.

The most extreme form of sex-dependent AI occurs when there is a reversal of the dominant allele in the

sexes ("sex-reversed AI"), whereby heterozygous males have >50% expression from one allele, while

heterozygous females have >50% expression from the alternative allele. We perform additional analyses

to find instances of sex-reversed AI. Cases of sex-reversed AI hold interest because they provide a

potential mechanism for sex-specific dominance reversals in fitness, whereby heterozygotes of each sex

could dominantly express the preferred allele for a locus under sexually antagonistic selection.

Theoretical models indicate that sexually antagonistic variants exhibiting dominance reversals are

preferentially maintained by balancing selection, due to partially mitigating sexual conflict (Kidwell et al.

1977; Spencer and Priest 2016; Connallon and Chenoweth 2019; Grieshop et al. 2024).

Our third goal was to examine *cis*-regulatory variation across tissues varying in their degree of sexual

dimorphism. We studied expression data from whole body, head, and gonad samples. Gonads and heads

offer an interesting comparison because gonads are highly sexually dimorphic whereas heads are much

less so. We compare the frequency of AI and particularly, sex-dependent AI between these two tissues to

discern if patterns in AI reflect the underlying levels of sexual dimorphism.

Methods

<u>Sample preparation and sequencing</u>

We crossed two inbred lines of *D. melanogaster* to obtain F_1 individuals for the study. DGRP-177

belongs to the Drosophila Genetic Reference Panel, a suite of genotypes derived from natural populations

of *D. melanogaster* in North Carolina (Mackay et al. 2012). SP159N was derived from a South African

population, as part of the Drosophila Population Genomics Project (Pool et al. 2012; Lack et al. 2016).

We crossed DGRP-177 males with SP159N females ("main cross"), and also performed a reciprocal

cross. All flies were reared and maintained at 25°C on a 12-hr light-dark cycle. After eclosion, virgin F_1

males and females were collected and kept in separate vials for 3 days. Vials that were observed to have

eggs laid in them were discarded. The F_1 individuals from the main cross (DGRP-177 males × SP159N

females) were used to obtain samples of whole body flies, heads, and gonads, with three replicate samples

of each sex*tissue-type combination. From the reciprocal cross, only whole body samples were collected

(three replicates of each sex). Testes and ovaries were obtained by dissection of anaesthetised whole flies,

performed over ice, with the flies placed in PBS buffer. Each replicate for gonads consisted of 10-12

individuals, while each replicate for whole flies and heads consisted of 5-7 individuals. Heads and whole

flies were flash-frozen in liquid nitrogen and then stored at -80°C as they awaited RNA extraction;

gonads were stored at -80°C upon dissection. RNA extraction was performed using ThermoFisher

PicoPure RNA Isolation Kit. Paired-end sequencing at a read length of 200 bp was performed using

Illumina NovaSeq6000. In addition, DNA was extracted from the parents and F_1 individuals using the

Qiagen DNeasy Blood and Tissue Kit. Paired-end whole genome sequences were obtained at a read

length of 100 bp using Illumina HiSeqX.

<u>Genotype-specific references</u>

Genotype-specific references are constructed from the reference genome by incorporating single

nucleotide polymorphisms (SNPs) for each parental genotype separately to reduce mapping bias when

analysing allele-specific expression (Rozowsky et al. 2011; Graze et al. 2012). We called variants (SNPs

and indels) for both DGRP-177 and SP159N using the GATK Best Practices workflow (McKenna et al.

2010; Auwera et al. 2013). The whole genome sequences for DGRP-177 and SP159N were aligned to the

D. melanogaster Release 6 reference genome (dos Santos et al. 2015) using BWA (Li and Durbin 2009)

with default parameters. The resulting alignment file was processed using GATK tools to sort reads and

mark PCR/optical duplicates. Variants were called by using GATK HaplotypeCaller and the resulting

VCF file was separated into two VCFs, one each for indels and SNPs. We obtained a BED file of

coordinates corresponding to heterozygous SNP calls and regions around indels for each parental genome.

Depending on the length of the indels, the following coordinates were included in the BED file: 10 bp

around indels of length ≤ 6 bp, 20 bp around indels of length > 6 bp but ≤ 12 bp, and 100 bp around indels

> 12 bp. The coordinates were then masked in the reference genome with N's. We masked sites without

high certainty of a homozygous SNP call in both parents, i.e., any sites where the major allele was < 95%.

We also masked sites where the depth of coverage fell below 10. Following recommendations by GATK,

the remaining variants—the homozygous SNPs—were subject to hard-filtering based upon quality

measures (Table S1). The filtered SNPs corresponding to DGRP-177 and SP159N were incorporated

separately into the masked reference genome, to create genotype-specific references for each parent.

<u>Competitive mapping</u>

For each sample, RNA-seq reads were aligned to both genotype-specific references using STAR v2.7

(Dobin et al. 2013) with default parameters. Following a similar method to Sánchez-Ramírez & Cutter

(2021), a Python-based competitive read mapping approach was used to obtain allele-specific read counts

from RNA-seq data for the F_1 heterozygotes. The two BAM alignments yielded by STAR were sorted and

indexed by read names. The alignment score (AS) and number of mismatches (nM) of each read when

aligned to the two references were used to assign each read's origin as follows: a) if a read had a higher

AS when aligned to a given parent's genotype-specific reference, it was ascribed as belonging to that

genotype, b) if the read has equal AS for both alignments, the read was assigned to the genotype that

yielded a lower nM, and c) if AS and nM do not differ between the two alignments for a read, it was

assigned to neither genotype, i.e., an "ambiguous" read. (We also tried the competitive alignment using

only nM to ascribe parental origin and obtained results similar to that from using both AS and nM.)

Ambiguous reads were excluded from further analysis. The percentage of such informative (i.e., non-

ambiguous) reads ranged from 18% to 43% for the samples examined (Table S2). Competitive read

mapping yielded two alignment files, corresponding to allele-specific reads of each parent. Finally, allele-

specific read counts were obtained from those BAM alignments using HTSeq-count v0.12 (Anders et al.

2015), with secondary or chimeric alignments being ignored when aggregating reads.

Bias in mapping can occur for multiple reasons and lead to errors in estimation of allele-specific

expression (Degner et al. 2009; León-Novelo et al. 2014). Genomic DNA from F_1s should have 50% of

reads mapping to each parent in the absence of mapping bias. As described in the Supplementary

Material, we used genomic DNA from male and female F_1 samples to conservatively filter out genes with

possible mapping bias (Supplementary Text S1, Figures S1-S3, Tables S3-S5).

Parental effects in AI

Parental effects on allele-specific gene expression in *Drosophila* are believed to be rare (Wittkopp et al. 2006; Coolon et al. 2012; Puixeu et al. 2023). Still, certain forms of mapping bias could manifest as apparent parental effects. To detect real or apparent "parental" effects, and then remove such genes from further analyses, we applied generalised linear models to the whole body samples from the main and reciprocal crosses utilising the R package *lme4* v.1 (Bates et al. 2007). The first test was applied separately to each sex. Only genes that had read counts >30 in all the 6 replicates were considered. For each gene, we analysed the proportion of reads assigned to the DGRP-177 genome with a quasibinomial model, including cross direction (main versus reciprocal) as a fixed effect. The intercept term of this model indicates allelic imbalance in the absence of an effect of cross direction. Additionally, we analysed the data for both sexes in a single model, including sex and "cross direction" as fixed effects, but without an interaction term. This test was applied to all genes with a total read count >20 in all 6 male and female replicates and average read count >30. All genes that had a significant cross-direction effect ($p < 0.05$) were excluded from subsequent analyses.

Inference of allelic imbalance and sex differences in allelic imbalance

For our analyses examining AI and sex-depenent AI (hereafter "SD-AI"), the data for each "tissue" type (gonads, heads, whole bodies) were analysed separately. Whole bodies from the main and reciprocal crosses were also analysed separately for two reasons. First, as we expect similar patterns in the whole body analyses from each cross direction, any major discrepancies could serve as a cause for concern. Second, the whole body results are comparable to results for gonads and heads because all the analyses are based on three replicates per sex for each tissue type. Only genes with at least 30 assignable reads in each of the replicates were included in the analysis for a given tissue. For each gene, we analysed the proportion of reads assigned to the DGRP-177 genome with a quasibinomial model, including sex as a fixed effect and using sum contrasts. In this model, the intercept term indicates the sex-averaged AI effect and the "sex" term indicates the difference in AI between sexes (i.e., a significant "sex" term is evidence of sex-dependent AI).

Inference of tissue effects on allelic imbalance

We performed analyses contrasting heads and gonads, parallel to our analyses of the two sexes. Separately for each sex, we first analysed the gonad and head data with a quasibinomial model including a tissue term. Significance of the intercept term is taken as evidence of (tissue-averaged) AI. A significant tissue term ($p < 0.05$) indicates tissue-dependent AI. Unexpectedly, we noted substantial differences in

the frequency of AI and tissue-dependent AI in the male and female analyses. Subsequently, we examined

those genes for which there was sufficient data in both sexes using a quasibinomial model that included

sex, tissue, and their interaction. The interaction term indicates sex differences in tissue-dependent AI.

<u>Inference of sex- and tissue-reversed allelic imbalance</u>

The strongest form of sex-dependent AI—sex-reversed AI—occurs when one allele is responsible for

greater than 50% expression in males, while the other allele is more highly expressed in females. To

maximize our power, we jointly analysed the whole body data from the main and reciprocal crosses. All

genes with read counts <20 in any of the replicates were excluded from analysis. Additionally, we

excluded genes where the average read count across all the 12 replicates was <30. We then tested each

sex separately for AI, using a quasibinomial model that only fit an intercept term, which signified the AI

effect. A gene was said to have sex-reversed AI if it fulfilled the following conditions: (i) it had a

significant AI effect in females, (ii) it had a significant AI effect in males, and (iii) the AI effect in the

sexes were in opposite directions. The number of false positives expected from this procedure depends on

the assumptions one makes for a null hypothesis. We considered several possibilities to arrive at a

conservative estimate of the number of false positive cases of dominance reversal (Supplementary Text

S1).

Likewise, we used data from gonads and heads to detect cases of tissue-reversed AI in males and females

separately. Again, after excluding genes with low read counts as above, we tested each tissue separately

for AI using a quasibinomial model where the intercept term represented AI. A gene was said to have

tissue-reversed AI if it showed significant AI in both heads and gonadal tissue, and the direction of the AI

was opposite between the two tissues. The number of false positive cases of tissue-reversed AI were

estimated in a manner analogous to that for sex-reversed AI (Supplementary Text S1).

<u>Modeling the distribution of AI and SD-AI</u>

In the earlier section, we described analyzing each gene individually to test for the presence of AI and

SD-AI. As with any set of frequentist analyses, the results will be subject to both false positives and false

negatives. Moreover, estimates of the average magnitudes of AI and SD-AI using "significant" genes will

be subject to bias due to the "winner's curse". As an alternative, we used a Bayesian approach to model

the joint distribution of AI effects across sexes, separately for each tissue.

As the true features of this distribution are unknown, there many ways one could choose to model it. Our

goal was to approximate the distribution using relative simple models framed with respect to our key

biological interests. To this end, we developed two models, each characterized by four parameters. The

first model framework characterizes the distribution in terms of (i) the frequency of genes with non-zero

sex-averaged AI, F_{AI}; (ii) the standard deviation in sex-averaged AI, σ_{AI}; (iii) the frequency of genes with

SD-AI among those with non-zero sex-averaged AI, $F_{SD\text{-}AI}$; and (iv) the standard deviation in the sex

difference in AI, $\sigma_{SD\text{-}AI}$. We model AI as a deviation in expression of allele 1 from 50% and sex

differences in AI as the difference of allele 1 expression in males from that in females. Because we

assume there is no net directionality to AI or SD-AI effects (i.e., there is no bias across genes, on average,

of which parental strain allele is more highly expressed), the average values of AI and SD-AI are zero;

thus, σ_{AI} and $\sigma_{SD\text{-}AI}$ represent the average magnitudes of AI and SD-AI, respectively. The second model

framework is similar but instead characterizes the distribution in terms of (i) the frequency of genes with

non-zero sex-averaged AI, F_{AI} (as in the first model framework); (ii) the standard deviation in AI in

females, $\sigma_{AI,F}$; (iii) the standard deviation in AI in males, $\sigma_{AI,M}$; and (iv) the intersexual correlation in AI

effects, ρ_{MF}. The first model framework is motivated primarily by an interest in estimating the fraction of

AI effects that differ between the sexes ($F_{SD\text{-}AI}$) whereas the second is motivated primarily by a related but

different property, the intersexual correlation in AI effects, ρ_{MF}.

For both model frameworks, the likelihood function relating the observed read counts to the distribution

incorporates a fifth (nuisance) parameter to allow for overdispersion. Because genes for which we have a

high amount of data (e.g., highly expressed genes) could have a disproportionate influence on the inferred

distribution of AI effects, we performed the analysis on a filtered data set for each tissue type. We

removed genes in the bottom quartile of total read count for either sex and then down-sampled the read

counts to equal levels for all genes (separately by sex). It is worth remembering that this (and other)

filtering steps non-randomly excludes genes from analysis (e.g., genes with low expression in either or

both sexes) and, thus, the inferred distribution may not be representative of such genes. We used the R

package *BayesianTools* (Hartig et al. 2023) to estimate a posterior distribution. Further details about the

models are provided in the Supplementary Text S3.

<u>Differential gene expression analysis</u>

We obtained gene-level read counts by using *htseq-count* on BAM alignments of each RNA-seq sample

to the *D. melanogaster* reference genome. We then estimated differential gene expression between the

sexes using the R package *DESeq2* (Love et al. 2014). In estimating differential expression, we removed

any genes averaging fewer than 50 reads across all the replicates for a given tissue. Differential

expression at each gene is estimated as the $\log_2$ fold change in male-to-female gene expression ($\log_2$FC).

Excluding a small number of genes where the $\log_2$FC had a standard error greater than 1.5, all other genes

were assigned to one of five sex-bias bins, as follows: highly female-biased ($-\infty < \log_2FC \leq -2$),

moderately female-biased ($-2 < \log_2FC \leq -0.5$), unbiased ($-0.5 \leq \log_2FC \leq 0.5$), moderately male-biased

($0.5 < \log_2FC \leq 2$), highly male-biased ($2 < \log_2FC < \infty$). Genes with relatively high uncertainty (i.e.,

standard error for $\log_2FC > 1.5$) were not assigned to any of these bins . The four tissue types—heads,

gonads, whole bodies (main cross) and whole bodies (reciprocal cross) —were separately analysed for

sex-biased expression.

Results

We crossed two inbred lines of *Drosophila melanogaster* (DGRP-177 and SP159N) and measured allele

specific expression in heads, gonads, and whole bodies of F_1s of both sexes. For whole body samples, we

had F_1s from both cross directions. A series of filters based on parental and F1 genomic data were applied

to remove genes with possible mapping biases (Methods).

<u>Apparent signals of parental effects</u>

Studies suggest that parental effects on allelic expression rarely occur in *Drosophila melanogaster*

(Wittkopp et al. 2006; Coolon et al. 2012; Chen et al. 2015). Moreover, patterns consistent with parental

effects may be due to residual mapping bias. We performed three tests using whole body samples from

the main and reciprocal crosses to identify genes with evidence of "parental effects," seeking to remove

these genes from subsequent analysis. We analyzed the data from each sex separately, and also performed

an analysis using combined data from the sexes. 346 out of 7716 (4.5%; X genes included) genes tested in

females and 829 out of 8817 (9.4%; X and Y genes excluded) genes tested in males showed a significant

effect of cross direction on allelic imbalance ($p < 0.05$). (The substantially higher incidence of "parental

effects" in F_1 males relative to females could be due to males of the two crosses having *trans*-regulatory

differences attributable to complementary XY combinations.) In the combined data from both sexes, 532

of 6371 (8.4%; X and Y genes excluded) genes had a significant effect of cross direction. A total of

10522 genes were analyzed across these three tests, with 1375 (13.1%) showing evidence of parental

effects in at least one of the tests. While many of these genes are expected to be false positives (5% of all

genes in each test), our primary goal here is to reduce the possibility of mapping bias influencing our

analyses of AI. We therefore excluded all these genes showing significant parental effects from further

analysis.

Sex-dependent effects on allelic imbalance

Allelic imbalance appears to be pervasive in the *D. melanogaster* transcriptome, with more than 30% of

the genes tested for each tissue showing significant AI (Table 1). Given our p-value threshold, we expect

~5% of genes to falsely appear as showing significant AI (i.e., false positives); the observed fraction is

much greater than expected under the null hypothesis. We use a p-value criterion for the ease of

comparisons among tissues, which would be complicated when applying a false discovery rate; false

discovery rate q-values depend not only on the data from the focal genes but also the distribution of p-

values within a group, which could cause genes with similar evidence for AI to have different q-values in

each group. Nevertheless, a version of Table 1 using FDR cut-offs is shown in Table S8. As expected,

using the more conservative FDR requirement reduces the overall frequency of detection of AI but the

main patterns are unchanged.

In each tissue, numerous genes exhibit significant sex-dependent allelic imbalance, SD-AI (Table 1).

About 25% of genes exhibited significant SD-AI in gonads, the most sexually dimorphic tissue in our

dataset. In contrast, only 10% of genes show SD-AI in heads. We considered the possibility that our

inferences of AI or SD-AI may be a consequence of mapping bias, despite our efforts to minimize such

errors. If mapping bias persists in the dataset despite the filters, we would expect patterns in genomic data

to be reflected in RNA-seq. Firstly, we would expect a positive correlation in fraction of DGRP-177 reads

between F1 genomic and F1 transcriptomic datasets in both the sexes. However, the correlations are very

close to zero (Figures S2 & S3; Tables S4 & S5), indicating that mapping bias is unlikely to be a

widespread problem among genes inferred to have AI. Secondly, we would expect a significant positive

correlation in the sex difference in the fraction of DGRP-177 reads between genomic and transcriptomic

datasets. These correlations are also close to zero, remaining so even as we consider only genes with

significant sex-dependent AI (Figure S4; Table S6). Thus, our inferences of AI or SD-AI are unlikely to

be substantially marred by mapping bias.

Table 1. Frequency of AI and sex-dependent AI.

Tissue	Number of genes tested	% genes with AI	% genes with sex-dependent AI
gonads	2547	31.6%	25.3%
heads	4062	45.0%	10.1%
whole bodies (main)	3627	36.5%	19.6%
whole bodies (reciprocal)	3901	35.7%	18.3%

347
348

<u>Estimating the distribution of AI and SD-AI</u>

The preceding section summarizes the results from assessing the statistical significance of each gene individually. As such, Table 1 reflects the frequencies of genes for which there was statistical power to detect evidence of AI and SD-AI. Thus, the values in Table 1 not only depend on the aspects of biology that are our real interest (frequency and effect sizes of AI and SD-AI), but also the number of samples, sequencing depth, and the choice of statistical threshold for significance. As an alternative to the approach used to generate Table 1, we employed a Bayesian analysis to model the underlying joint distribution of AI across the sexes in each tissue type. The results are summarized in Table 2. Across all tissues, the inferred frequency of genes with AI was similar (F_{AI} = 38-48%) as was the magnitude of the (sex-averaged) effect (~5%). However, there were major differences between tissues with respect to the frequency of sex-differences in AI. In gonads, almost all genes with AI were inferred to have sex differences (F_{SD-AI} = 94%), whereas in heads very few were (F_{SD-AI} = 5%). In gonads, the effect sizes of SD-AI tend to be quite large relative to the sex-average effect size of AI. (This is related to a result shown in a later section: AI effects tend to be larger in male than female gonads.) The values in Table 2 represent, to our knowledge, the only available estimates of the distribution of AI/SD-AI effects but, as with any statistical model, some caution is warranted as these estimates are contingent on the assumptions underlaying the framework of the modelled distribution.

369　**Table 2.** Parameter estimates [with 95% high posterior density interval] for the joint distribution of AI

370　effects across sexes.

	Frequency of genes with AI, F_{AI}	Average magnitude of sex-averaged AI[1], σ_{AI}	Frequency of genes with SD-AI among genes with AI, F_{SD-AI}	Average magnitude of sex difference in AI[2], σ_{SD-AI}	Overdispersion parameter, υ
Gonads	0.48	0.053	0.94	0.105	0.006
	[0.45, 0.52]	[0.0496, 0.056]	[0.90, 1.00]	[0.099, 0.114]	[0.005, 0.006]
Heads	0.42	0.050	0.05	0.074	0.001
	[0.40, 0.45]	[0.054, 0.060]	[0.01, 0.08]	[0.051, 0.106]	[0.001, 0.001]
Whole Body (main)	0.45	0.049	0.75	0.059	0.001
	[0.44, 0.47]	[0.048, 0.051]	[0.71, 0.79]	[0.058, 0.061]	[0.001, 0.002]
Whole Body (reciprocal)	0.38	0.055	0.66	0.065	0.003
	[0.35, 0.40]	[0.053, 0.058]	[0.60, 0.71]	[0.060, 0.070]	[0.002, 0.003]

371　[1] σ_{AI} is the standard deviation of sex-averaged effects, which is modelled as approximately
372　normally distributed with mean 0.
373　[2] σ_{SD-AI} is the standard deviation of sex differences in AI effects, which is modelled as
374　approximately normally distributed with mean 0.
375
376

<u>AI in relation to sex-biased gene expression</u>

We examined how the patterns of AI vary with the degree of sex bias in expression by fitting a Bayesian

model separately to gene sets binned by sex bias. For this section, we used an alternative framework for

the distribution of AI effects, parameterizing the distribution in terms of the frequency with which AI

occurs, F_{AI} (as in the previous section), as well as the correlation of AI effects between males and

females, ρ_{MF}. There is heterogeneity in the frequency of AI with respect to sex bias (Fig. 1A; Tables S10-

12). In heads, F_{AI} declines from female-biased to unbiased to male-biased genes. There is a similar trend

in the whole body samples, though here it is noteworthy that the inferred average effect sizes also vary

with lower estimates for female- than male-biased genes (Fig. 1A; Table S10-11).

390
Figure 1. Model estimates of (A) the frequency of AI, F_{AI}, and (B) the intersexual correlation of AI
effects, ρ_{MF}. The leftmost section in each panel shows results irrespective of sex-bias; the remaining
sections show results stratified by sex bias: highly female-biased ($-\infty < \log2FC \leq -2$), moderately female-
biased ($-2 < \log2FC \leq -0.5$), unbiased ($-0.5 \leq \log2FC \leq 0.5$), moderately male-biased ($0.5 < \log2FC \leq 2$),
highly male-biased ($2 < \log2FC < \infty$). No analysis was performed for high sex-bias categories for heads
as there were too few genes. Error bars represent 95% high density interval of the posterior. See Table
S11 for other model parameters.
398

The intersexual correlation of AI effects is high in heads across all sex bias categories ($\rho_{MF} > 0.9$; Fig.

1B) whereas it is much lower in gonads ($\rho_{MF} < 0.25$). In the whole body samples, there is a striking

change in ρ_{MF} across sex-bias categories, going from low values for female-biased genes to high values

for male-biased ones. (Applying the other model parameterization to these data, a qualitatively similar

compatible pattern is manifest in that SD-AI is common among female-biased genes and much rarer for

male-biased ones; Table S10).

<u>Sex-dependent reversals of allelic imbalance</u>

Instances of sex-dependent allelic imbalance may include cases where the direction of imbalance is

reversed between the sexes (i.e., male expression of the DGRP-177 allele is >50%, while female

expression is <50%). Cases of sex-reversed AI are of special interest but are expected to be rare as this

scenario presumably involves more complex regulation. While the Bayesian models in the previous

sections are designed to make inferences about the distribution using information from a large number of

genes, they are not well-suited to clearly identify individual genes exhibiting sex-reversed AI. To detect

cases of sex-reversed AI, we analyzed genes individually, requiring significant AI effects in both sexes

but of opposite direction. We examined whole body samples, combining data from both main and

reciprocal crosses to maximize power. Of the 3796 genes tested in this fashion, 176 (4.6%) genes showed

significantly sex-reversed AI. Using relatively conservative assumptions, we expect to find ~26 cases by

chance (i.e., false positives; see Supplementary Material).

We further examined these 176 genes using the head and gonad data to see if similar patterns can be

observed in narrower tissue samples, i.e., gonads or heads. In the gonads, 155 of 176 genes had sufficient

data to be examined for AI, whereas 160 genes could be examined in the heads. Of the 155 genes

examined in the gonads, the majority (109) had sex-reversed point estimates of AI in the same direction

as observed with the whole bodies whereas only 5 showed sex-reversed point estimates in the opposite

direction relative to that found in the whole bodies. Thus, for the putative set of genes with sex-reversed

AI, there exists a fair degree of concordance in the number and direction of sex-reversed AI between the

gonads and whole bodies. In contrast, in heads, only 35 out of the 160 genes tested had sex-reversed point

estimates of AI, and roughly half (only 17) of these were in the same direction as in the whole bodies,

with the remaining half (18) in the opposite direction relative to whole bodies, i.e., not different from that

expected by chance.

<u>Tissue-dependent patterns of allelic imbalance</u>

In our previous analyses, we noted marked differences in the frequency of AI between heads and gonads.

Though the analyses used different sets of genes for each tissue, the results suggest that AI may differ

between tissues. To investigate tissue differences in AI more formally, we applied to the Bayesian model

to estimate the joint distribution of AI effects in heads and gonads, separately for each sex (Table 3). In

both sexes, we infer that ~40% of genes exhibit AI and > 80% of these have effects that differ between

gonads and heads. Using the alternative Bayesian model to characterize the distribution in terms of the

between-tissue correlation in effects (Table S13), we find this correlation is quite low ($\rho_{head\text{-}gonad} \approx 0.25$)

and that the average magnitude of effect sizes is larger in gonads than heads (gonads ≈ 0.09; heads $\approx$

0.05).

442
443 **Table 3.** Parameter estimates [with 95% high posterior density interval] for the joint distribution of AI
444 effects across heads and gonads.
445

	Frequency of genes with AI, F_{AI}	Average magnitude of tissue-averaged AI[1], σ_{AI}	Frequency of genes with Tissue-Dependent AI among genes with AI, $F_{TD\text{-}AI}$	Average magnitude of tissue difference in AI[2], $\sigma_{TD\text{-}AI}$	Overdispersion parameter, v
Female	0.37 [0.33, 0.41]	0.053 [0.049, 0.057]	0.94 [0.88, 1]	0.094 [0.086, 0.101]	0.0039 [0.0036, 0.0042]
Male	0.44 [0.41, 0.47]	0.050 [0.048, 0.053]	0.84 [0.78, 0.91]	0.099 [0.094, 0.105]	0.0012 [0.001, 0.0014]

446 [1] σ_{AI} is the standard deviation of sex-averaged effects, which is modelled as approximately normally
447 distributed with mean 0.
448 [2] $\sigma_{TD\text{-}AI}$ is the standard deviation of tissue differences in AI effects, which is modelled as approximately
449 normally distributed with mean 0.

450
Analogous to our earlier test for sex-reversed AI, we tested for tissue-reversed AI, separately in each sex.

We find 33 instances of possible tissue-reversed AI between gonads and heads in females, and 94 such

instances in males. We estimate that 11 of these genes are likely false positives in females and 9 in males

(see Supplementary Material). The much higher number of putative tissue-reversed AI in males may be a

result of tissue differences in AI being larger in males, leading to easier detection of reversals in AI.

Discussion

This study focuses on *cis*-regulatory variation in relation to sexual dimorphism in *D. melanogaster*. We

utilise a divergent cross between North American (DGRP-177) and South African (SP159N) genotypes to

maximize the number of polymorphic sites in the F_1, thus increasing our power to detect allele-specific

expression. To some extent, the patterns of allelic imbalance observed here may reflect inter-population

differences in expression or may be idiosyncratic to the specific lines used here. As with any study of

genetic variation, our results will be influenced by the interplay between the capacity of mutation to
generate regulatory variation and selection to filter it. Our goal was to explore sex-specific patterns of *cis*-
regulatory variation but our study was not designed to assess the relative importance of different
evolutionary factors (e.g., mutation-selection-drift balance, balancing selection within populations,
divergent selection among populations) in causing these patterns.
468
When analyzing individual genes, allelic imbalance was detected in 30-45% of genes analyzed (Table 1).
These values are considerably higher than the values (6-17%) reported by Puixeu et al. (2023). The
greater divergence of haplotypes used in our study may have contributed to this difference; we crossed fly
lines from different continents whereas they crossed lines from within a single population. However,
experimental and statistical differences are likely major contributors to this discrepancy. Rather than
relying on the proportion of genes that reach significance in individual gene tests which depend heavily
on statistical power, we also developed a Bayesian approach to directly estimate the frequency of genes
with AI, finding that the frequency is quite high (38-48%, Table 2).
477
In heads and whole body samples, there is an enrichment of *cis*-regulatory variants in female-biased
genes. Osada et al. (2017) and Puixeu et al. (2023) also examined *cis*-regulatory variation in relation to
sex bias. Some of the patterns reported in those two studies appear similar while others appear different
from ours or from each other. However, it is not possible to directly compare those studies to our own or
to each other because of differences in methodology use to investigate these patterns. An advantage of our
approach is that patterns can be investigated with respect to biologically interpretable properties such as
the frequency of AI and average effect sizes.
485
The higher frequency of AI among female-biased genes that we detected may suggest that these genes are
more frequently under balancing selection or tend to experience weaker purifying selection. Sexually
antagonistic selection is one possible reason genes could experience balancing selection or weaker
purifying selection. Cheng and Kirkpatrick (2016) suggested that genes with intermediate levels of sex-
biased expression—both male- and female-biased—would be most likely to experience sexually
antagonistic selection. As we do not observe an elevated frequency of AI for genes with intermediate
levels of sex-bias, our results would seem to indicate that either that sexually antagonistic selection does
not drive the pattern we observe or that the sexually antagonistic selection is not distributed across the
genome as Cheng and Kirkpatrick (2016) suggested. While it is most intuitive to think of sexually
antagonistic selection causing balancing selection or weak purifying selection, it could cause strong
purifying selection if selection in one sex is much stronger than the other. Thus, the suggestion of Cheng

and Kirkpatrick (2016) is not at necessarily at odds with our results if, for example, sexually antagonistic

selection results in weak purifying selection for intermediately female-biased genes but strong purifying

selection for intermediately male-biased genes.

Purifying selection (regardless of whether it involves sexual antagonism) is believed to be the most

common form of selection and genes with *cis*-regulatory variation have previously been shown to be

subject to weaker purifying selection in *Capsella* (Steige et al. 2017). Strongly female-biased genes in *D.*

melanogaster have somewhat elevated levels of non-synonymous diversity (Singh and Agrawal 2023; but

see details within), suggesting they may tend to be under weaker purifying selection and providing a

potential explanation for the increased frequency of AI among these genes. A non-selective alternative

explanation for our observed pattern is mutational bias; female-biased genes could have greater scope for

cis-regulatory mutations (e.g., if such genes tend to have larger or more complex regulatory regions). We

cannot ascertain which of these possibilities explain the observed pattern.

In general, *cis* effects are small in magnitude, with a large majority of genes exhibiting allelic expression

differences below 5-10%. This finding is general to all the tissues examined, though average effects were

notably higher in gonads than heads (Table S13). Others have also found frequent, weak *cis*-regulatory

effects in other species, including *Capsella* (Steige et al. 2017), mice (Crowley et al. 2015) and

Saccharomyces (Zhang and Emerson 2019). However, there have been reports of strong *cis* effects in *D.*

melanogaster (León-Novelo et al., 2018), interspecific *Drosophila* hybrids (Graze et al. 2012) and natural

populations of flycatchers (Wang et al. 2017), though the detection of low-effect variants may be

encumbered by the lack of statistical power in some of these studies. An advantage of the Bayesian model

approach we used is that it should allow us to infer the average effect size without the bias ("winner's

curse") from gene-by-gene analyses.

Using gene-by-gene analyses, we found numerous instances of sex-dependent *cis*-regulatory effects in all

tissues, though these were much more common in gonads than heads (Table 1). The inference from the

Bayesian analysis indicated that almost all (>94%) *cis*-regulatory variants affect the sexes differently in

gonads. Puixeu et al. (2023) also found a substantially higher frequency genes with sex differences in *cis*-

regulatory effects in gonads than heads. The distinct regulatory architecture of the gonads, manifest in the

high frequency of *cis*-regulatory variants with sex-dependent effects, should allow expression in gonads

to evolve relatively independently between the two sexes whenever there is selective pressure to do so.

By contrast, in heads we estimated that only a small fraction (~5%) of *cis*-regulatory effects have sex-

dependent effects, restricting the potential for independent evolution of expression between the two sexes.

It is unsurprising that the frequency of sex-dependent AI in whole bodies is intermediate between the

estimates inferred for gonads and heads. Relatedly, the intersexual correlation of AI effects, ρ_{MF}, in whole

bodies is intermediate between gonads and heads. However, there is a striking pattern of ρ_{MF} in relation to

sex bias in whole bodies that is not apparent in either gonads or heads: ρ_{MF} steadily increases from

strongly female-biased to strongly male-biased gene categories. This pattern seems similar to a pattern

observed for the intersexual genetic correlation (r_{MF}) in whole body gene expression (Singh and Agrawal

2023) in which the average r_{MF} increases from strongly female-biased to strongly male-biased gene

categories. It should be noted ρ_{MF} and r_{MF} are not defined at the same "level"; ρ_{MF} is a correlation in AI

effects between sexes estimated from a set of genes whereas each gene has its own r_{MF} value—the

intersexual correlation in expression level—estimated among a set of genotypes. Nonetheless, one would

expect that if ρ_{MF} was low (high) for a class of genes, then the average r_{MF} for that class of genes would

also be low (high) if *cis* regulatory effects were a major source of the genetic variation in expression. It is

unclear why this pattern in ρ_{MF} occurs in whole bodies but not heads or gonads and it is also unknown if

r_{MF} is related to sex-bias in tissues other than whole bodies.

Reversals in the direction of AI between the sexes are the most extreme form of sex-dependent AI and we

detect such reversals at 176 loci in whole body samples. Puixeu et al. (2023) also documented some

examples of sex-reversed AI (though using less stringent criteria to assess false positives). These are

intriguing because heterozygotes of each sex will predominantly express alternative alleles for these

genes. If the coding regions of the associated alleles were under sexually antagonistic selection, this type

of *cis*-regulatory variation provides a potential mechanism by which sex-specific dominance reversals

could occur (Grieshop et al. 2024). Dominance reversals have been invoked in the theoretical literature to

explain the maintenance of genetic variation (Curtsinger et al., 1994; Kidwell et al., 1977; reviewed in

Grieshop et al. 2024), though it has been unclear whether dominance reversals would be biologically

plausible. Recently, this idea has gathered increasing empirical support with the discovery of sex-specific

dominance reversal for traits likely under sexually divergent selection in salmonids (Barson et al. 2015;

Pearse et al. 2019) as well as signals of dominance reversals for fitness in seed beetles (Grieshop and

Arnqvist 2018). Sex reversals in AI potentially provide a mechanism, but it remains to be seen whether

such *cis*-regulatory effects underlie any cases of dominance reversals for fitness.

In this study, we focussed on sex-dependent aspects of *cis*-regulatory variation because of its implications

for sexual dimorphism and sexual conflict. However, 'sex' is one of many dimensions in which the

context for a gene's expression can differ. *Cis*-regulatory effects can vary within other contexts, such as

temperature (Chen et al. 2015; Li and Fay 2017), tissue (Pinter et al. 2015) , or metabolic state (Shih and

Fay 2021). Of the two dimensions we examined—sex and tissue—neither stood out as harbouring

dramatically more variation than the other (i.e., sex-dependent AI was very common in gonads as was

tissue-dependent AI in both sexes). We were motivated to look for sex-dependent AI, at least in part,

because of the possibility that sexually antagonistic selection might help maintain such variants.

However, the qualitatively similar frequency of tissue-dependent AI suggests that either sexual

antagonism is playing little role in maintaining regulatory variation genome-wide or that antagonistic

pleiotropy across tissues is similarly common. Regardless, our results add to the existing literature that

cis-regulatory effects are often context dependent in magnitude and sometimes even in direction.

Moreover, different dimensions of context can interact to affect allelic expression, e.g., we find a sizeable

number of genes with tissue-by-sex interaction effects on AI. It is plausible that this thinking extends to

other contexts (e.g., sex-dependent *cis*-regulatory effects may be sensitive to temperature).

Allele-specific expression analysis can be plagued by inaccurate mapping of reads to the reference. We

attempted minimize the influence of mapping bias in several ways. We constructed genotype-specific

references and ascertained their efficacy by competitively mapping parental genomic reads to the

references. We tested F_1 genomic data for significant deviation from the expected read count ratio of 1:1

and removed any such gene from further consideration. We also tested for apparent 'parental effects' on

allelic expression, which may ensue from any residual mapping bias, and excluded any such genes.

Among our retained genes, the percentage of reads assigned to the DGRP-177 allele are uncorrelated

between genomic and transcriptomic data, whereas a positive correlation would be expected if mapping

bias was a substantial problem. The correlation remains negligible even if we only consider those genes

with significant AI in expression. These steps are designed to allay the effects of mapping bias on our

inferences, with evidence suggesting they have been largely effective (Supplementary Material).

Nonetheless, we cannot entirely rule out the possibility that mapping bias may persist in the dataset within

a small number of retained genes. While we believe our major findings are robust, the results for

individual genes would benefit from further experimental validation.

As we infer *cis*-regulatory polymorphisms from allelic imbalance (i.e., relative expression) in

heterozygotes, an additional note of caution is needed with respect to total expression. If allele *A* is

expressed more than *B* in an *A/B* heterozygote, it is reasonable to assume that total expression of this gene

would be greater in *A/A* than *B/B* homozygotes. However, this may not always be the case owing to

negative feedback loops in transcription. Further experimental testing—including both homozygotes—

would be needed to evaluate how often observed AI causes the predicted effects on total expression,

though this is extremely difficult to do at scale while controlling for genetic background effects.

In summary, our data indicate that *cis*-regulatory variants are common and their effects are often

dependent on sex and tissue. The frequency of *cis*-regulatory variants varies non-randomly with respect to

sex-biased gene expression as well as across tissues. Further work remains to be done to more thoroughly

examine variants at individual loci as well as to understand the evolutionary forces shaping patterns of

variation genome-wide.

Literature Cited

Anders S, Pyl PT, Huber W. 2015. HTSeq--a Python framework to work with high-throughput

sequencing data. Bioinformatics. 31(2):166–169. doi:10.1093/bioinformatics/btu638.

Auwera GA, Carneiro MO, Hartl C, Poplin R, del Angel G, Levy-Moonshine A, Jordan T, Shakir K,

Roazen D, Thibault J, et al. 2013. From FastQ Data to High-Confidence Variant Calls: The

Genome Analysis Toolkit Best Practices Pipeline. Current Protocols in Bioinformatics. 43(1).

doi:10.1002/0471250953.bi1110s43. [accessed 2021 Aug 20].

https://onlinelibrary.wiley.com/doi/10.1002/0471250953.bi1110s43.

Barson NJ, Aykanat T, Hindar K, Baranski M, Bolstad GH, Fiske P, Jacq C, Jensen AJ, Johnston SE,

Karlsson S, et al. 2015. Sex-dependent dominance at a single locus maintains variation in age at

maturity in salmon. Nature. 528(7582):405–408. doi:10.1038/nature16062.

Bates, D., Maechler, M., Bolker, B. and S. Walker. 2015. Fitting Linear Mixed-Effects Models Using

lme4. Journal of Statistical Software, 67(1), 1-48. doi:10.18637/jss.v067.i01.

Bonduriansky R, Chenoweth SF. 2009. Intralocus sexual conflict. Trends in Ecology & Evolution.

24(5):280–288. doi:10.1016/j.tree.2008.12.005.

Campbell CD, Kirby A, Nemesh J, Daly MJ, Hirschhorn JN. 2008. A survey of allelic imbalance in F1

mice. Genome Res. 18(4):555–563. doi:10.1101/gr.068692.107.

Chen J, Nolte V, Schlötterer C. 2015. Temperature Stress Mediates Decanalization and Dominance of

Gene Expression in Drosophila melanogaster. PLoS Genet. 11(2):e1004883.

doi:10.1371/journal.pgen.1004883.

Cheng C, Kirkpatrick M. 2016. Sex-specific selection and sex-biased gene expression in humans and

flies. PLoS Genet. 12(9):e1006170. doi:10.1371/journal.pgen.1006170.

Connallon T, Chenoweth SF. 2019. Dominance reversals and the maintenance of genetic variation for

fitness. PLoS Biol. 17(1):e3000118. doi:10.1371/journal.pbio.3000118.

Connallon T, Clark AG. 2014. Balancing Selection in Species with Separate Sexes: Insights from Fisher's

Geometric Model. Genetics. 197(3):991–1006. doi:10.1534/genetics.114.165605.

Coolon JD, Stevenson KR, McManus CJ, Graveley BR, Wittkopp PJ. 2012. Genomic imprinting absent

in Drosophila melanogaster adult females. Cell Reports. 2(1):69–75.

doi:10.1016/j.celrep.2012.06.013.

Crowley JJ, Zhabotynsky V, Sun W, Huang S, Pakatci IK, Kim Y, Wang JR, Morgan AP, Calaway JD,

Aylor DL, et al. 2015. Analyses of allele-specific gene expression in highly divergent mouse

crosses identifies pervasive allelic imbalance. Nat Genet. 47:353–360. doi:10.1038/ng.3222.

Cubillos FA, Stegle O, Grondin C, Canut M, Tisné S, Gy I, Loudet O. 2014. Extensive *cis* -Regulatory
Variation Robust to Environmental Perturbation in *Arabidopsis*. Plant Cell. 26(11):4298–4310.
doi:10.1105/tpc.114.130310.

Curtsinger JW, Service PM, Prout T. 1994. Antagonistic pleiotropy, reversal of dominance, and genetic
polymorphism. American Naturalist. 144:210–228. doi:10.1086/285671.

Degner JF, Marioni JC, Pai AA, Pickrell JK, Nkadori E, Gilad Y, Pritchard JK. 2009. Effect of read-
mapping biases on detecting allele-specific expression from RNA-sequencing data. Bioinformatics.
25(24):3207–3212. doi:10.1093/bioinformatics/btp579.

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR.
2013. STAR: ultrafast universal RNA-seq aligner. Bioinformatics. 29(1):15–21.
doi:10.1093/bioinformatics/bts635.

Fairbairn DJ, Blanckenhorn WU, Székely T. 2007. Sex, size and gender roles: evolutionary studies of
sexual size dimorphism. Oxford University Press.

Genissel A, McIntyre LM, Wayne ML, Nuzhdin SV. 2007. Cis and Trans Regulatory Effects Contribute
to Natural Variation in Transcriptome of Drosophila melanogaster. Molecular Biology and
Evolution. 25(1):101–110. doi:10.1093/molbev/msm247.

Goncalves A, Leigh-Brown S, Thybert D, Stefflova K, Turro E, Flicek P, Brazma A, Odom DT, Marioni
JC. 2012. Extensive compensatory *cis-trans* regulation in the evolution of mouse gene expression.
Genome Res. 22(12):2376–2384. doi:10.1101/gr.142281.112.

Graze RM, Novelo LL, Amin V, Fear JM, Casella G, Nuzhdin SV, McIntyre LM. 2012. Allelic
Imbalance in Drosophila Hybrid Heads: Exons, Isoforms, and Evolution. Molecular Biology and
Evolution. 29(6):1521–1532. doi:10.1093/molbev/msr318.

Grieshop K, Arnqvist G. 2018. Sex-specific dominance reversal of genetic variation for fitness. PLoS
Biol. 16:e2006810. doi:10.1371/journal.pbio.2006810.

Grieshop K, Ho EKH, Kasimatis KR. 2024. Dominance reversals and the maintenance of genetic
variation. Proc B. accepted.

Gruber JD, Long AD. 2009. *Cis* -regulatory variation is typically polyallelic in Drosophila. Genetics.
181(2):661–670. doi:10.1534/genetics.108.098459.

Hartig F., Minunno F., S. Paul. 2023. BayesianTools: General-Purpose MCMC and SMC Samplers and
Tools for Bayesian Statistics. R package version 0.1.8, <https://CRAN.R-
project.org/package=BayesianTools>.

Kidwell JF, Clegg MT, Stewart FM, Prout T. 1977. Regions of stable equilibria for models of differential
selection in the two sexes under random mating. Genetics. 85(1):171–183.
doi:10.1093/genetics/85.1.171.

Kita R, Venkataram S, Zhou Y, Fraser HB. 2017. High-resolution mapping of *cis* -regulatory variation in

budding yeast. Proc Natl Acad Sci USA. 114(50):E10736–E10744. doi:10.1073/pnas.1717421114.

Lack JB, Lange JD, Tang AD, Corbett-Detig RB, Pool JE. 2016. A Thousand Fly Genomes: An

Expanded *Drosophila* Genome Nexus. Mol Biol Evol. 33(12):3308–3313.

doi:10.1093/molbev/msw195.

Lemos B, Araripe LO, Fontanillas P, Hartl DL. 2008. Dominance and the evolutionary accumulation of

cis- and trans-effects on gene expression. Proceedings of the National Academy of Sciences.

105(38):14471–14476. doi:10.1073/pnas.0805160105.

León-Novelo L, Gerken AR, Graze RM, McIntyre LM, Marroni F. 2018. Direct testing for allele-specific

expression differences between conditions. G3 Genes|Genomes|Genetics. 8(2):447–460.

doi:10.1534/g3.117.300139.

León-Novelo LG, McIntyre LM, Fear JM, Graze RM. 2014. A flexible Bayesian method for detecting

allelic imbalance in RNA-seq data. BMC Genomics. 15:920. doi:10.1186/1471-2164-15-920.

Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows-Wheeler transform.

Bioinformatics. 25(14):1754–1760. doi:10.1093/bioinformatics/btp324.

Li XC, Fay JC. 2017. Cis-Regulatory Divergence in Gene Expression between Two Thermally Divergent

Yeast Species. Genome Biology and Evolution. 9(5):1120–1129. doi:10.1093/gbe/evx072.

Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for RNA-seq

data with DESeq2. Genome Biol. 15(12):550. doi:10.1186/s13059-014-0550-8.

Mackay TFC, Richards S, Stone EA, Barbadilla A, Ayroles JF, Zhu D, Casillas S, Han Y, Magwire MM,

Cridland JM, et al. 2012. The Drosophila melanogaster Genetic Reference Panel. Nature.

482(7384):173–178. doi:10.1038/nature10811.

McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, Garimella K, Altshuler D,

Gabriel S, Daly M, et al. 2010. The Genome Analysis Toolkit: A MapReduce framework for

analyzing next-generation DNA sequencing data. Genome Research. 20(9):1297–1303.

doi:10.1101/gr.107524.110.

Metzger BPH, Wittkopp PJ, Coolon JosephD. 2017. Evolutionary Dynamics of Regulatory Changes

Underlying Gene Expression Divergence among Saccharomyces Species. Genome Biology and

Evolution. 9(4):843–854. doi:10.1093/gbe/evx035.

Osada N, Miyagi R, Takahashi A. 2017. *Cis* - and *Trans* -regulatory effects on gene expression in a

natural population of *Drosophila melanogaster*. Genetics. 206:2139–2148.

doi:10.1534/genetics.117.201459.

Parsch J, Ellegren H. 2013. The evolutionary causes and consequences of sex-biased gene expression. Nat

Rev Genet. 14(2):83–87. doi:10.1038/nrg3376.

Pastinen T, Hudson TJ. 2004. Cis-Acting Regulatory Variation in the Human Genome. Science.

306(5696):647–650. doi:10.1126/science.1101659.

Pearse DE, Barson NJ, Nome T, Gao G, Campbell MA, Abadía-Cardoso A, Anderson EC, Rundio DE,

Williams TH, Naish KA, et al. 2019. Sex-dependent dominance maintains migration supergene in

rainbow trout. Nat Ecol Evol. 3(12):1731–1742. doi:10.1038/s41559-019-1044-6.

Pinter SF, Colognori D, Beliveau BJ, Sadreyev RI, Payer B, Yildirim E, Wu C, Lee JT. 2015. Allelic

imbalance is a prevalent and tissue-specific feature of the mouse transcriptome. Genetics.

200(2):537–549. doi:10.1534/genetics.115.176263.

Pool JE, Corbett-Detig RB, Sugino RP, Stevens KA, Cardeno CM, Crepeau MW, Duchen P, Emerson JJ,

Saelao P, Begun DJ, et al. 2012. Population genomics of sub-Saharan Drosophila melanogaster:

African diversity and non-African admixture. PLoS Genet. 8(12):e1003080.

doi:10.1371/journal.pgen.1003080.

Puixeu, G, Macon, A, and B Vicoso. 2023. Sex-specific estimation of *cis* and *trans* regulation of gene

expression in heads and gonads of *Drosophila melanogaster*. *G3: Genes, Genomes, Genetics*

jkad121.

Rozowsky J, Abyzov A, Wang J, Alves P, Raha D, Harmanci A, Leng J, Bjornson R, Kong Y,

Kitabayashi N, et al. 2011. AlleleSeq: analysis of allele-specific expression and binding in a

network framework. Mol Syst Biol. 7(1):522. doi:10.1038/msb.2011.54.

Rowe L, Chenoweth SF, Agrawal AF. 2018. The genomics of sexual conflict. Amer Nat. 192:274-86.

Sánchez-Ramírez S, Cutter AD. 2021. CompMap: an allele-specific expression read-counter based on

competitive mapping. Bioinformatics. [accessed 2021 Aug 18].

http://biorxiv.org/lookup/doi/10.1101/2021.02.12.431019.

dos Santos G, Schroeder AJ, Goodman JL, Strelets VB, Crosby MA, Thurmond J, Emmert DB, Gelbart

WM, the FlyBase Consortium. 2015. FlyBase: introduction of the Drosophila melanogaster Release

6 reference genome assembly and large-scale migration of genome annotations. Nucleic Acids

Research. 43(D1):D690–D697. doi:10.1093/nar/gku1099.

Schärer L, Rowe L, Arnqvist G. 2012. Anisogamy, chance and the evolution of sex roles. Trends in

Ecology & Evolution. 27(5):260–264. doi:10.1016/j.tree.2011.12.006.

Shih C-H, Fay J. 2021. Cis-regulatory variants affect gene expression dynamics in yeast. eLife.

10:e68469. doi:10.7554/eLife.68469.

Singh A, Agrawal AF. 2023. Two forms of sexual dimorphism in gene expression in *Drosophila*

melanogaster: their coincidence and evolutionary genetics. Molecular Biology and Evolution.

40(5):msad091.

Spencer HG, Priest NK. 2016. The evolution of sex-specific dominance in response to sexually

antagonistic selection. Amer Nat 187:658-66.

Steige KA, Laenen B, Reimegård J, Scofield DG, Slotte T. 2017. Genomic analysis reveals major

determinants of *cis*- regulatory variation in *Capsella grandiflora*. Proc Natl Acad Sci USA.

114(5):1087–1092. doi:10.1073/pnas.1612561114.

Stern DL. 2000. Perspective: Evolutionary developmental biology and the problem of variation.

Evolution. 54(4):1079–1091. doi:10.1111/j.0014-3820.2000.tb00544.x.

Stranger BE, Montgomery SB, Dimas AS, Parts L, Stegle O, Ingle CE, Sekowska M, Smith GD, Evans

D, Gutierrez-Arcelus M, et al. 2012. Patterns of Cis Regulatory Variation in Diverse Human

Populations. Barsh GS, editor. PLoS Genet. 8(4):e1002639. doi:10.1371/journal.pgen.1002639.

Wang M, Uebbing S, Ellegren H. 2017. Bayesian Inference of Allele-Specific Gene Expression Indicates

Abundant Cis-Regulatory Variation in Natural Flycatcher Populations. Genome Biology and

Evolution. 9(5):1266–1279. doi:10.1093/gbe/evx080.

Wittkopp PJ, Haerum BK, Clark AG. 2004. Evolutionary changes in cis and trans gene regulation.

Nature. 430(6995):85–88. doi:10.1038/nature02698.

Wittkopp PJ, Haerum BK, Clark AG. 2006. Parent-of-Origin Effects on mRNA Expression in *Drosophila*

melanogaster Not Caused by Genomic Imprinting. Genetics. 173(3):1817–1821.

doi:10.1534/genetics.105.054684.

Wittkopp PJ, Haerum BK, Clark AG. 2008. Regulatory changes underlying expression differences within

and between Drosophila species. Nat Genet. 40(3):346–350. doi:10.1038/ng.77.

Wray GA. 2007. The evolutionary significance of cis-regulatory mutations. Nat Rev Genet. 8(3):206–

216. doi:10.1038/nrg2063.

Zhang X, Emerson JJ. 2019. Inferring the genetic architecture of expression variation from replicated

high throughput allele-specific expression experiments. Genetics. [accessed 2021 Dec 17].

http://biorxiv.org/lookup/doi/10.1101/699074.