1   **MalariaSED: a deep learning framework to decipher the regulatory contributions of**
2   **noncoding variants in malaria parasites**
3

4   Chengqi Wang[1]*, Yibo Dong[1], Jenna Oberstaller[1], Chang Li[1], Min Zhang[1], Justin Gibbons[1],
5   Camilla Valente Pires[1], Lei Zhu[4], Rays H.Y. Jiang[1], Kami Kim[2], Jun Miao[2], Thomas D. Otto[3],
6   Liwang Cui[2], John H. Adams[1], Xiaoming Liu[1]
7
8   [1]Center for Global Health and Infectious Diseases Research and USF Genomics Program,
9   College of Public Health, University of South Florida, Tampa, FL, USA.
10
11   [2]Department of Internal Medicine, Morsani College of Medicine, University of South Florida,
12   Tampa, FL, USA
13
14   [3]Institute of Infection, Immunity and Inflammation, MVLS, University of Glasgow, Glasgow, UK.
15
16   [4]School of Biological Sciences, Nanyang Technological University, Singapore, Singapore
17
18
19   *To whom correspondence should be addressed. Email: chengqi@usf.edu

Abstract

Malaria remains one of the deadliest infectious diseases. Transcriptional regulation effects of noncoding variants in this unusual genome of malaria parasites remain elusive. We developed a sequence-based, *ab initio* deep learning framework, MalariaSED, for predicting chromatin profiles in malaria parasites. The MalariaSED performance was validated by published ChIP-qPCR and TF motifs results. Applying MalariaSED to ~1.3 million variants showed that geographically differentiated noncoding variants are associated with parasite invasion and drug resistance. Further analysis revealed chromatin accessibility changes at *P. falciparum* rings are partly associated with artemisinin resistance. MalariaSED illuminates the potential functional roles of noncoding variants in malaria parasites.

Introduction

Malaria remains one of the largest global public health challenges, with an estimated ~241 million cases worldwide in 2020 despite remarkable achievements in reducing this deadly disease over the past decade. It has been reported that some noncoding regions in *P. falciparum* are associated with artemisinin resistance (ART-R)[1,2] or regulate a group of genes related to crucial biological processes, such as antigen variation[3,4], merozoite invasion[5-7] and gametocytogenesis[6]. Therefore, understanding the effects of genomic variants in malaria parasites is critical for monitoring the spread of drug resistance and the escape of the host immune response. Unlike genomic variants in protein-coding regions whose impacts can often be deduced from resultant codon changes, functional effects associated with noncoding variants cannot be so easily assessed. Despite the large number of noncoding genomic variants recorded in the public variant-tracking resource MalariaGEN[8] (~1.31 million noncoding vs. ~1.35 million coding), detecting the contribution of mutations located at noncoding regions is still challenging, and establishing the role of the vast noncoding space in the genetic basis of crucial parasite phenotypes remains difficult.

Genome-wide association studies (GWASs) have been widely used to identify variants in the *P. falciparum* genome significantly associated with drug resistance[1,9-18]. However, GWAS is an a posteriori approach requiring many historical samples that take time to acquire. Although functional screens have been implemented previously[15,19], these had limited throughput for testing the whole repertoire of noncoding variants, and these procedures are also time-consuming and expensive[20].

Sequence-based prediction models provide a new perspective on the effects of genomic variants on gene regulation. However, all currently available models were developed in model

organisms[21-24], which are not suitable for malaria parasites due to several unusual features of

their genome architecture[25]. The most striking trait of the *P. falciparum* genome is its high AT

content (~80% on average, rising to ~92% in intergenic regions). Additional features

complicating the applicability of existing predictive models include the apparent absence of

some subunits of the preinitiation complex[26], as well as the outsized contributions of a relatively

small 27-member sequence-specific transcription factor (TF) family known as ApiAP2s to

transcriptional regulation[27]. In comparison, the similarly sized genome of the yeast

Saccharomyces cerevisiae contains ~170 sequence-specific TFs[28]. A sequence-based model

accounting for these peculiarities is required for accurate prediction of gene expression profiles

associated with noncoding variants specifically for malaria parasites.

In this work, we have leveraged recent advancements in deep learning (DL) to develop a

sequence-based computational framework, MalariaSED, for noncoding variant effect prediction

in malaria parasites. To our knowledge, this is the first time in malaria parasites using DL to

learn regulatory sequence patterns, including TF binding, chromatin accessibility and histone

modification profiles. This framework is complementary to experimental approaches at higher

throughput and with lower cost. MalariaSED predicts epigenetic effects based upon sequence

alterations at single-nucleotide resolution, enabling us to trace the associated biological process

naturally-selected on regulatory sequences and to improve the prioritization of functional

variants associated with important phenotypes.

Results

MalariaSED accurately predicts chromatin profiles from genomic sequences in malaria

parasites

The unusual genome property of malaria parasites requires the development of a novel

computational framework to understand the regulatory sequence code from parasite noncoding

sequences. To this end, we developed the first sequence-based deep learning framework,

MalariaSED (Supplement table 1), for malaria parasites, which learns the sequence code of

transcriptional regulatory elements by training in large-scale epigenetic experimental data,

including open chromatin accessibility (ATAC-Seq)[29], TF binding, and histone-mark profile

(ChIP-Seq)[5-7,30,31]. MalariaSED computes a chromatin score $f(s)$ for a sequence s using three

stages:

$$f(s) = net(LSTM(Conv(s)))$$

Here, *s* is an input sequence represented as a $4 \times l$ matrix, where each row represents a

nucleotide, and each column indicates a position in the length l sequence. '1' represents a

specific nucleotide present at a specific position of the sequence, and '0' does not (Fig. 1A). Our

results indicate that the 1 kb sequence context substantially improves the performance of

MalariaSED (Supplement figure 1). The convolution layer (Conv) is a motif scanner analogizing

the position weight matrix. The bidirectional long short-term memory (LSTM) uses forward and

reverse complement features as input to capture the relationship between a set of sequence

information outputted from the convolution layer. The LSTM is crucial since TF binding sites

may have a group of sequence patterns separated from each other by a long distance. We used

the flatten layer and sigmoid function ('net' in the equation) to generate chromatin prediction

scores. The high predictive performance for eight chromatin profiles at multiple parasite stages

in both *P. falciparum* and *P. berghei* demonstrates that the MalariaSED architecture can capture

important patterns in DNA sequences (average Area Under the Receiver Operating

Characteristic curve (auROC) > 0.96, Fig. 1B, Supplement table 2). Fig. 2 illustrates an

agreement between the predicted PfAP2-G binding profile and ChIP-seq/Input ratio[6] across a

genomic region of chromosome 10 throughout commitment and early gametocytogenesis. We

also tested the previously DL model trained in the mammalian genome[21], which can only obtain

an average auROC of 0.81 (Supplement figure 2). We reasoned that the outperformance of

MalariaSED is due to the structural discrepancy between the malaria parasite genome and that

of model organisms, such as the number of TFs and highly skewed AT content.

CRISPR/Cas9 mutations in ApiAP2 binding sites demonstrate the capacity of MalariaSED

to identify damaging genetic mutations

Unidirectional flow from sequence information to consequent epigenetic status in MalariaSED

framework provides a novel technique for estimating the noncoding variant effects in *malaria*

genomes. We tested this capability in predicting ChIP profile changes at multiple TF binding

motif mutations generated by the CRISPR–Cas9 system. These ChIP-qPCR profiles are

currently available for three TFs, PfAP2-G5[32], PfAP2-G[6] and PfAP2-I[5,] in *P. falciparum*. Four

PfAP2-G5 motif mutations, S1-mutation (mut), S2-mut and S3-mut and S3-deletion (del), were

generated at the *ap2-g* gene upstream regulatory region[32]. Their surrounding [-100, +100] bp

regions are illustrated in Fig. 3A as S1, S2 and S3. The PfAP2-G5 ChIP-Seq results indicate a

strong binding affinity of PfAP2-G5 at S1 and S2, whereas S3 presents a moderate level. The

mutations, S1-mut and S2-mut from the original ChIP-qPCR profiles showed minor binding

effects, but S3-mut and S3-del strongly weakened PfAP2-G5 binding sites. Further comparison

between S3-del and S3-mut indicated higher binding effects at S3-mut. We validated the

predictions from MalariaSED by these ChIP profiles. The estimated PfAP2-G5 binding

probability from MalariaSED, represented as S1-WT, S2-WT and S3-WT in Fig 3B, is correlated

with the ChIP-Seq profiles. In contrast, the binding effects shown as S1-mut, S2-mut, S3-mut

and S3-del in Fig.3B are in concordance with the original ChIP-qPCR results. We also used

MalariaSED to predict the PfAP2-G binding change at a region containing eight GTAC motifs

(S4 in Fig. 3C). A previous study showed that mutation of the three motifs by CRISPR/Cas9

significantly reduced binding of PfAP2-G[6]. This result supports the predictions from MalariaSED,

which shows a strong decrease in PfAP2-G binding probability in Fig. 3D. An exceptional case

for MalariaSED prediction occurred at the promoter region of the *pfmsp5* gene, where ChIP-

qPCR showed PfAP2-I binding but was not identified by the original ChIP-Seq signal[5]. Since

MalariaSED was trained by the ChIP-Seq results, it only predicted a PfAP2-I binding probability

lower than 0.1 for both the WT and ATGCA motif mutations.

Single-nucleotide substitutions at the TF motif weaken TF binding

The successful validations of TF binding changes in CRISPR/Cas9-generated motif mutations

imply potential DNA motif contributions to MalriaSED prediction. To validate this globally, we

used MalariaSED to interrogate the average TF binding effects of all possible single nucleotide

substitutions (SNSs) at the reported motifs (Supplement Table 3, see Methods for definition).

Basically, for each site of a reported motif, the reference nucleotide was in turn changed by all

other three nucleotides as motif mutation sets (e.g., a DNA motif comprising 5 nucleotides

generates 5 x 4 = 20 motif mutations). Each generated sequence in the mutation sets was

individually selected to replace the TF motifs that appeared in all the predicted TF binding

regions. The average binding effect of all these SNSs was used to evaluate the motif

contribution to TF binding. All other k-mer sequences with the same length as the investigated

motif were used as the control sets (shown as 'ALL' in Fig. 4). We used this strategy to check

the reported motifs of the parasite-specific TF PfAP2-G, a key determinant of sexual

commitment in *P. falciparum*. We observed a significantly reduced binding affinity of PfAP2-G

after incorporating SNSs at its binding motif GTRC (committed schizonts) and GTAC (sexual

rings and stage I gametocytes) (Fig. 4A, Wilcoxon test p < 2.2e-16). Similarly, for PfAP2-I, a key

regulator of red blood cell invasion, SNSs at the reported motif (GTGCA) significantly weakened

its binding affinity at the schizont stage (Fig. 4B, Wilcoxon test p < 2.2e-16). Previous results

indicate that two motifs (GGTCG and CTTGC) at other DNA-binding domains of PfAP2-I are not

required for PfAP2-I binding during the RBC stage[5]. MalariaSED prediction also showed that

SNSs in these two motifs did not significantly reduce PfAP2-I binding (Fig. 4B). When another

invasion regulator, PfBDP1, was examined, SNSs introduced into the PfAP2-I GTGCA motif led

to a significantly reduced binding level in schizonts (Fig. 4C, Wilcoxon test p < 2.2e-16),

supporting the conclusion that PfBDP1 is likely to form a protein complex with PfAP2-I to

regulate invasion genes[5,7,33]. For the parasite trophozoite stage, we also checked the DNA

binding change of PfBDP1 by introducing SNSs at the PfAP2-I motif GTGCA (Fig. 4C) and did

not observe significant changes compared with other 5-mer sequences. This result supports

that PfBDP1 binds many gene promoters not bound by PfAP2-I in trophozoites and are in line

with the conclusion that PfBDP1 and PfAP2-I are involved in a protein complex in trophozoites

but not in schizonts[7]. Furthermore, the predicted results from MalaraSED do not show

significant binding effects of the motifs TAACT and ACAAC on PfBDP1 binding sites

(Supplement Figure 3). It has been shown that TAACT, the second motif of PfBDP1 binding, co-

occurs with the micronemal DNA motif ACAAC in the promoters of several micronemal and

rhoptry neck genes[5]. The limited contributions of these two motifs suggest that some other

unidentified TFs may recruit PfBDP1 to regulate micronemal gene expression. We also tested

MalariaSED on two AP2 family TFs, PbAP2-G2 and PbAP2-O, in *P. berghei*[30,31]. PbAP2-G2

represses asexual genes to support conversion to the sexual stage, while AP2-O plays a critical

role in regulating mosquito midgut invasion. MalariaSED identifies SNSs in two binding motifs,

GTTG for PbAP2-G2 and GCTA for PbAP2-O, which are more likely to abolish their binding (Fig.

4D, Wilcoxon test p < 2.2e-16). Taken together, these results suggest that MalariaSED can

capture crucial DNA sequences responsible for TF binding. Critically, MalariaSED predicted

disrupted TF binding in mutants with mutations at TF binding motifs from sequence data alone,

demonstrating the utility of sequence-based models for interpreting functional variants in malaria

parasites.

Geographically differentiated variants at noncoding regions are more likely to alter their

surrounding chromatin environments

Approximately six million curated genomic variants (including SNPs and short indels) from 7,000

P. falciparum samples recently made available on MalariaGEN show the enrichment of

geographically differentiated SNPs in nonsynonymous variants[8]. Further analysis indicates that

genes harboring the top geographically variable nonsynonymous SNPs tend to be involved in

the process of parasite transmission by the mosquito vector and are associated with drug

resistance. Despite noncoding variants showing the second-highest geographic differentiation

after nonsynonymous variants, the limited functional annotation of malaria parasite noncoding

regions restricts our understanding of noncoding-variant contributions to this process. Our initial

analysis of ~1.3 million noncoding variants[8] (trimming off ~1.4 noncoding loci with low

Low_VQSLOD provided by MalariaGEN) indicates that noncoding variants with high levels of

geographic differentiation (global $F_{ST} > 0.1$) are located closer to their nearest gene (Fig. 5A,

Wilcoxon test $p < 2.2e-16$ compared with genomic variants showing global $F_{ST} < 0.1$),

suggesting that the differentiation process at the noncoding region may partly be due to the

selection of regulatory activity. We used MalariaSED to predict the impact of chromatin profiles

of noncoding variants (Supplement Figure 4, Supplement Table 4). We observed significant

alteration of chromatin profiles in the group of genomic variants presenting strong geographic

differentiation (Fig 5B, Wilcoxon test $p < 2.2e-16$ compared with noncoding variants showing

limited geographic differentiation). The higher chromatin effects observed at these

geographically differentiated variants indicate that *P. falciparum* may use epigenetic

mechanisms to locally regulate the expression of genes of certain biological pathways to benefit

its survival.

Drug resistance and red blood cell invasion-related pathways are associated with

geographic differentiation in *P. falciparum* noncoding regions

We next investigated the biological pathways associated with strong geographic differentiation

at noncoding regions by assigning variants to their nearest genes ($F_{ST} > 0.1$, distance to the

nearest gene < 3 kb). Only the noncoding variants leading to significantly predicted chromatin

profile changes (the top 1% of chromatin effects) were selected. Pathway enrichment analysis

indicated that oxidative stress defense (S-nitrosylation and S-glutathionylation), proteasome

degradation (ubiquitin-protein conjugates) and food vacuoles were associated with noncoding

variants predicted to cause significant chromatin accessibility changes in the ring and early

trophozoite stages (Fig. 6A, B, Supplement table 5 for the complete list of enriched pathways).

Other enriched pathways included protein transport (PTEX export), epigenetic modifications and

the palmitome. Notably, most of these pathways have been previously demonstrated to be

associated with altered transcription profiles in ART-R parasites[34,35]. Our results suggest that

geographically diverse noncoding variants may enable the parasite to modulate the transcription

of key pathways underlying the artemisinin response, ultimately allowing survival under drug

pressure. Interrogating enriched genes around each geographically differentiated variant

leading to binding affinity changes for the TFs PfBDP1, PfAP2-G and PfAP2-I (Fig. 6C-E), we

identified a group of genes related to merozoite invasion, including *EBA181, AARP, RhopH2,*

RhopH3 and *SRA, RON2, MSP9, RAP1*. PfBDP1 is a chromatin-associated protein responsible

for regulating the expression of genes involved in red blood cell (RBC) invasion[7]. TF PfAP2-I

has been shown to associate with PfBDP1 and positively regulate the transcription of invasion-

related genes[5]. As a known pivotal activator of early gametocyte genes, PfAP2-G was also

recently reported to have a potential role in regulating genes important for RBC invasion[6]. *P.*

falciparum strains present broad diversity in invasion phenotypes and gene expression across

populations [36]. The TF binding affinity changes surrounding noncoding variants upstream of

invasion-related genes may explain how parasites maintain this diversity of gene expression.

Significant chromatin accessibility changes surround artemisinin resistance-related

eQTLs

The large-scale Expression quantitative trait loci (eQTL) and Transcriptome-wide association

studies (TWAS) study conducted by Tracking Resistance to Artemisinin Collaboration I (TRACI)

identified more than 13,000 SNS-expression linkages in 773 parasite isolates from the Greater

Mekong subregion (GMS) [1]. The result provides a new perspective to understand ART-R

associated with SNSs through the unidirectional flow of information from sequence changes to

consequent gene expression changes and, ultimately, phenotype alteration. However, the

extensive gaps in our understanding of genomic changes and the resulting transcriptional

alterations in malaria parasites impede the use of TWAS. Considering the correlations between

chromatin accessibility and neighboring gene activity[29], we reasoned that the surrounding

chromatin profile changes resulting from eQTLs would lead to gene transcription changes. To

test the associations between the eQTL and its surrounding chromatin profiles, we investigated

the capacity of different chromatin profile effects to distinguish local intergenic eQTLs from SNS

reported in MalariaGEN[8]. Our results show that chromatin accessibility in rings outperforms

other epigenetic markers in different parasite stages as an indicator/predictor of eQTLs (auROC

= 0.7, Fig. 7A, Supplement table 6), consistent with skewness toward the ring-stage parasites

collected in the TRACI study[1]. We further tested this on a group of local eQTLs linked with ART-

R by regulating ART-R-related gene expression (see Methods). Ring-stage chromatin

accessibility also presented the highest predictive performance in identifying ART-R-associated

eQTLs (auROC = 0.73, Fig. 7B), indicating that transcriptional regulatory activity changes may

lead to neighboring gene transcriptional alterations associated with ART-R phenotypes. Taken

together, these results demonstrate that MalariaSED is informative for predicting the

transcriptional impact of common noncoding variants in *P. falciparum*, such as eQTLs.

Multiple chromatin profile alterations at the schizont stage show signatures of intergenic

selection

Previous analysis of single-cell sequencing data revealed strong evidence that *de novo*

mutations are under selection in *P. falciparum* populations[37]. This evidence includes the

skewness toward protein-coding regions, high nonsynonymous vs. synonymous substitution

ratio, and elevated expression levels of *de novo* mutations-targeted genes. Because of limited

functional annotation of *P. falciparum* intergenic regions, this analysis was focused on coding

regions. We used MalariaSED to investigate the chromatin effects of the 35 noncoding *de novo*

mutations reported in the SCS study. Compared with the genome-wide background distribution

generated from all SNSs reported in MalariaGEN[8], significant changes were observed in

schizonts, including chromatin accessibility, H3K9ac and two TFs, PfAP2-I and PfAP2-G binding

affinities (Fig 8, Wilcoxon test p < 0.05). Our results show signatures of selection for intergenic

variants in schizonts, which could result in neighboring gene expression changes.

Discussion

This work presents the first sequence-based computational framework, MalariaSED, that learns

regulatory sequence codes in malaria parasites. In malaria parasite genomes, MalariaSED

outperforms two DL models previously developed for mammalian genomes, Enrofmer[21] and

Basenji2[23]. The DL-based framework enables us to predict the chromatin effects of sequence

alterations with single-nucleotide sensitivity, thereby providing a means to assess the functional

relevance of noncoding variants. We demonstrated the accuracy of this approach using

previously published ChIP–qPCR data from CRISPR/Cas9-generated mutations in AP2 TF-

family binding motifs[6,32]. One failure case occurred at the *pfmsp5* promoter, which was not

identified in PfAP2-I ChIP-seq but detected in the ChIP–qPCR experiment[5]. MalariaSED could

not predict PfAP2-I binding in either the wild type or parasites carrying mutations at the PfAP2-I

binding motif at the *pfmsp5* promoter. We reason that some DNA sequence patterns may be

related to discrepancies between the ChIP–qPCR and ChIP-Seq results. This DNA sequence

pattern is reflected at the *pfmsp5* promoter and cannot be captured by MalariaSED trained in

ChIP-seq data. We also demonstrated the significant contributions of previously identified DNA

motifs to known TF binding, such as the members in the PfAP2 family, PfBDP1, PbAP2-O and

PbAP2-G2[5-7,30,31]. We are not clear the reasons for MalariaSED not capturing previously

reported PfAP2-G5 binding motifs, although it achieves prediction performance auROC > 0.95

and validates previous CRISPR/Cas9 results. One explanation is that MalariaSED may use

DNA patterns where nucleotides are intermittently arranged for AP2-G5 binding predictions.

We used MalariaSED to explore the epigenetic effects of large amounts of noncoding

variants released from MalariaGEN partner studies[38]. The results indicate that noncoding

variants that acquire high levels of geographic differentiation significantly likely impact the

surrounding epigenetic profiles. This suggests that some noncoding regions have been at least

in part subjected to natural selection, which is consistent with the previous analysis results on

coding regions. It is unclear why all epigenetic markers present significant changes surrounding

geographic differentiation variants. Further study is needed to understand the dependence of

each chromatin profile. We speculate that other histone modifications may recruit some TFs and

that the binding affinity alterations may result from these marker changes. Our reanalysis of

~1.3 million noncoding variants also provides new insight into parasite local geographical

adaptation driven by noncoding variations. The associated potential transcriptional change in

protein trafficking, epigenetic and translational machinery, and the alternations of many PfAP2-

G binding affinities provide new evidence that parasites trade off reproduction (transmission to

new hosts) and growth (within-host replication) to adapt to different local regions where

parasites present diverse transmission intensity[39]. We need to point out that the PfAP2-G

binding change regions tend to be located upstream of genes important for RBC invasion. This

is further demonstrated by the overlapping gene list with PfAP2-I and PfBDP1, as well as the

limited functional enrichment in sexual rings and stage I gametocytes for PfAP2-G binding site

changes. We also observed that a group of geographic differentiation variants located at

noncoding regions are likely to regulate gene functions in food vacuoles, oxidative stress

defense, and proteasome degradation. This reflects the evolution of genetically distinct *P.*

falciparum subpopulations that are thought to occur due to high drug pressure and oxidative

stress[1,40].

To test the epigenetic contributions resulting from intergenic variants to transcriptional

change, we used chromatin effects to discriminate reported eQTLs[1] against the nontrait-

associated SNSs collected in MalariaGEN partner studies[38]. We observed high concordance

between the chromatin accessibility change at the ring stage and the local eQTL discovered in

P. falciparum clinical isolates. This result is also shown up in the subset of eQTLs linked with

ART-R via transcriptional profiles. The agreement between the clinical isolate stage (high

skewness toward rings) and the ring stage of the chromatin profile supports a previous study

showing that the chromatin accessibility pattern correlates with neighboring gene expression

values[29]. Furthermore, MalariaSED bridges the gap in our understanding between DNA

sequence changes and the resulting epigenetic profile alterations in malaria parasites. Building

on the known association between gene transcription and parasite phenotype, the directional

flow of information from sequence variation to consequent chromatin profile change to gene

transcription alternation and further to ART-R effect will allow us to extract the causality of

variant(s). Based on currently available chromatin profiles, we are not able to identify any

chromatin profiles associated with distal eQTLs (auROC < 0.65 for all 13 chromatin profiles

here). This may be due to limited epigenetic data available in *P. falciparum* and the fact that

only one chromatin profile in rings can be implemented in MalariaSED. It will be interesting to

study these distal eQTL hotspots when new epigenetic data are available for MalariaSED

training.

MalariaSED facilitates the functional explanations of *de novo* mutations discovered in

SCS studies. The significant *de novo* mutations enrichment in coding regions and the

disproportionate genes targeted by *de novo* mutations suggested strong selection in the *P.*

falciparum population collected in Chikhwawa, Malawi[37,41]. This includes the ApiAP2 gene family,

implying selection in the epigenetic regulation mechanism. We used *de novo* mutations to

investigate the chromatin effects of intergenic *de novo* mutations and identified significant

changes in schizonts, including H3K9ac, PfAP2-G and PfAP2-I binding affinities, as well as

chromatin accessibility. However, due to the limited number of intergenic *de novo* mutations in

the single cell sequencing study, we are hesitant to make conclusions of potential functions

associated with these 35 noncoding *de novo* mutations. Further study is required when more

single cell sequencing datasets are generated.

In the long run, we expected that sequence-based regulation analysis would become a

crucial part of malaria parasite research, especially to help unveil the regulatory activity of

noncoding genomic regions, which are currently poorly understood in malaria parasites. Such

analyses could, in the future, be implemented to prioritize noncoding variants potentially

perturbing gene expression contributing to parasite immune escape and drug resistance. With

the increasing data availability of functional variants, the approach can be readily trained,

adapted, and further improved.

Methods

Datasets for MalariaSED development

The previously published ATAC-Seq datasets[29], nine ChIP-seq datasets, including PfAP2-I[5],

PfAP2-G[6], PfAP2-G5[32] and PfBDP1[7], from *P. falciparum* RBCs to the gametocyte stage and two

ChIP-Seq datasets for PbAP2-O and PbAP2-G2[30,31] in *P. berghei* were used to develop the

MalariaSED framework. The whole genome is divided into bins of 200 bp, yielding more than

50,000 bins. A bin is labeled "positive" if more than 60% of it overlaps with ChIP or ATAC peaks,

while the remaining bins are used as negative datasets. We used the ratio 7/1.5/1.5 to divide all

extracted bins into training, test, and validation sets. The number of bins used for MalariaSED

training for different epigenetic markers is listed in supplement table 2.

Deep neural network models with long short-term memory (LSTM) layers

For each bin 200 bp in length, we extended it by 400 bp in both directions, resulting in a 1 kb

input sequence. We used one-hot encoding to convert each 1 kb sequence into a 1000 x 4 bit

matrix, where '1' represents a specific nucleotide present and '0' does not. The architecture of

MalariaSED framework is shown in Fig. 1A. Two 1D convolutional layers followed up with max

pooling and drop-out layers are combined with bidirectional LSTM to capture dependencies of

DNA sequence patterns contributing to specific epigenetic markers. Regularization and

constraint are applied to all layers to minimize overfitting. After the flatten layer, the final

predictions are computed through sigmoid activation in dense layers. The Adam optimizer was

used to search for model parameters based on weighted binary cross-entropy. We use early

stopping to seize the parameter optimization process if there is no improvement in validation

auPRC for ten consecutive epochs. Bayesian optimization[42] was used to optimize the

hyperparameters.

Sequence pattern contribution to MalariaSED

To estimate the sequence contributions to MalariaSED predictions, we predict chromatin profile

changes by generating all possible SNSs along with each site of the k-mer sequence. Each 200

bp region predicted as positive from MalariaSED was extended by 400 bp in both directions,

resulting in a 1 kb sequence window. SNS generation was then performed for every positive 1

kb bp region covering all 4-mer or 5-mer sequences. For each 4/5-mer sequence, we took the

average chromatin effects resulting from all SNSs as the contribution to MalariaSED. The effect

of an SNS on a specific chromatin marker was measured by the absolute value of log2-fold

change of odds predicted from MalariaSED. This measurement was also successfully utilized to

evaluate the chromatin effects of genomic variants in the human genome[24,43].

Functional enrichment for geographic differential variants at intergenic regions

The intergenic variants presenting high geographic differential (Fst > 0.1) from MalariaGEN

partner studies[38] were selected for this analysis. We further picked up the geographic differential

variants predicted to change the particular chromatin profile at the top 1% level to explore the

potential biological function change. A gene is used for functional enrichment analysis if its

distance to at least one selected variant uses the abovementioned criteria shorter than 3 kb. To

identify GO terms enriched in the gene list of interest, Fisher's exact test was used to test

whether more genes with this GO term were observed compared with the background

comprising the whole genome.

Chromatin profile prediction surrounding eQTLs

We downloaded 13,257 eQTLs from a previous TRACI study[1]. Only local intergenic eQTLs

(15 kb between an eQTL and its regulated target(s)) were selected for the positive dataset in

DL predictions. We strengthened the ART-R-associated eQTLs by the cutoff of an absolute

value of 0.1 for the adjusted Spearman correlation between the eQTL-regulated transcriptional

profile and parasite clearance time. All SNSs gathered in MalariaGene[8] are used as the

negative dataset. Each reported eQTL or SNS is extended to include the surrounding 1 kb DNA

sequence to fit the input requirement of MalariaSED. The chromatin effects of each variant were

measured by the log2-fold change difference of odds, which was previously used in human

genetic studies[43].

Data and code availability

The data supporting the findings of the study are available within the paper and its

supplementary information files. The MalariaSED web portal for 15 epigenetic profile predictions

for malaria parasites is at malariased.charleywanglab.org. The source code for running and

training MalariaSED models is available at https://github.com/CharleyWang/MalariaSED.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Funding

This work was supported by the National Institutes of Health grant R01 AI117017 (J.H.A.) and
U19 AI089672. This work was supported by the internal award of College of Public Health at
University of South Florida.
427
Acknowledgements
We thank the USF Genomics Program Omics Hub for the productive discussion. We appreciate
MalariaGEN for providing support for using their collected large-scale genomic variants data.
We thank Dr. Derek Wildman and Dr. Monica Uddin for the long-time support. We appreciate Dr.
Ian Cheeseman from Texas Biomedical Research Institute for providing the noncoding de novo
mutations data. We appreciate Dr. Michael Duffy from The University of Melbourne for the
guidance of using their published ChIP-seq data.
435

FIGURE LEGENDS

Figure 1. MalaiaSED accurately predicts chromatin profiles in malaria parasites. A. The
general framework of the deep learning model. The input DNA sequence is encoded into the
four rows matrix by one hot encoder, where each element represents the appearance of a
nucleotide at a specific location. The following two convolutional layers capture cumulative
effects of short sequence patterns by analogy to motif scan. The bidirectional LSTM layer
follows up to summarize long-term dependencies between captured DNA patterns. Outputs are
fed into the flatten and full connection layers. We calculate the final score by dense layer with
sigmoid activation. **B.** Performances of the deep learning framework are measured by AUC in
multiple experimental epigenetic profiles in *P. falciparum* and *P. berghei*.

**Figure 2. MalariaSED can predict the binding dynamic of PfAP2-G throughout
commitment and early gametocytogenesis.** High agreement between log2-transformed
PfAP2-G ChIP/input ratio and predicted genomic tracks from MalariaSED over a region of
chromosome 10. The PfAP2-G ChIP intensity covers parasite stage committed schizonts,
sexual ring and stage I gametocyte.

**Figure 3. PfAP2-G5 and PfAP2-G binding affinity predicted from MalariaSED are
consistent with previous published ChIP-qPCR results. A**. PfAP2-G5 ChIP-Seq intensity
along the promoter region of *ap2-g* during Trophozoits stage. ChIP-seq results come from a
recent study[32]. The flanking region [-100,+100] bp of each mutation on PfAP2-G5-binding sites
is highlighted as S1 (dark red), S2 (Black), and S3 (Brown). **B**. The binding affinity predicted

from MalariaSED is consistent with the recently published ChIP-qPCR results[32].The height of

each bar indicates the predicted binding affinity of PfAP2-G5 in three mutants highlighted in A.

C. PfAP2-G ChIP-Seq intensity[6] along the promoter region of *ap2-g* in early sexual rings. The

S4 (Dark yellow) indicates flanking region [-100,+100]bp of the mutation of the PfAP2-G motif.

D. Similar to B, but showing the predicted binding affinity of the mutant S4 in C. The results are

in concordance with the published ChIP-qPCR result[6] that PfAP2-G is not able to bind in the

mutant.

Figure 4. TFs binding effects of genomic variants predicted by the DL framework

demonstrate the contribution of previously identified sequence motif. A. Potential single-

nucleotide substitutions (SNSs) at reported pfAP2-G motif, GTRC (committed schizonts), GTAC

(sexual rings and stage I gametocytes) significantly reduce the binding affinity of PfAP2-G.

Comparisons are between reported PfAP2-G motifs and all 4-mers (ALL) appeared on predicted

ApAP2-G binding sites. **B.** The PfAP2-I binding effects of base substitutions at the reported

motif GTGCA is significant compared with all 5-mers (ALL) in *P. falciparum* genome during

schizonts. The other two motifs (GGTCG and CTTGC) from PfAP2-I binding domains 1 and 2

are not able to show significantly reduced PfAP2-I affinity, which is consistent with experimental

results[5]. **C.** The significantly reduced binding affinity of SNSs at reported PfBDP1 binding motif

GTGCA in schizonts support the previous conclusion that PfAP2-I and PfBDP1 may form a

protein complex to bind DNA. However, SNSs at GTGCA motif are not likely to disrupt PfBDP1

binding at trophozoite stage. **D,E.** Single-nucleotide substitutions at two reported binding motifs

in *P. Berghei* significantly weaken PbAP2-G2 and PbAP2-O binding sites. '***' indicates

Wilcoxon test compared with whole-genome background p < 2.2e-16.

Figure 5. Noncoding variants with high levels of geographic differentiation tend to alter

their surrounding chromatin profiles. A. Non-coding variants strongly differentiated at the

global level are closer to their neighboring genes. We used global Fst > 0.1 to identify variants

with high levels of geographic differentiation. '***' indicates Wilcoxon test compared with

remaining non-coding variants, p < 2.2e-16. **B.** Non-coding variants acquired high geographic

differentiation (Fst > 0.1) are predicted to bring in significant epigenetic changes at their +/-

100bp surrounding regions.

Figure 6. The enriched functions or pathways upon noncoding variants with strong

geographic differentiation. The height of yellow bar indicates the number of genes assigned

to noncoding variants that are strongly geographically differentiated and significantly change the

chromatin profile. To estimate the number of genes expected to observe for each function or

pathway term, the gene fraction in *P. falciparum* genome presenting the specific functional term

is multiplied by the number of assigned genes to geographic differential variants exhibiting the

high chromatin effects. (Hypergeometric test, FDR-corrected P-value < 0.1, check supplement

table X for full list).

Figure 7. Chromatin accessibility alternation is surrounding previously identified eQTLs

in the TRACI study. A. The chromatin effects of reported eQTLs in the TRACI study are used

to discriminate them from all SNPs gathered in Pf3K. **B.** The eQTLs linked with ART-R

associated transcriptions were also investigated. The eQTL prediction performance of different

chromatin profiles is measured by AUC.

Figure 8. The results from MalariaSED indicates de novo mutations at non-coding

regions discovered from the single cell study have higher chance to alter their

surrounding chromatin profiles in schizonts. '*' indicates Wilcoxon test compared with non-

coding SNPs collected from MalariaGEN, p < 0.05.

SUPPLEMENTARY FIGURE LEGENDS

Supplement figure S1. The same DL architecture as MalariaSED was trained on 200, 700,

1000 and 1400 bp DNA sequences, respectively. We present the auROC of each model as a

point and use a box plot to show the distribution of the models inputting the same DNA

sequence length. The results indicate 1000bp input outperforms other length input ('***'

indicates Wilcoxon test p < 0.01 compared between the 1kb input sequence and any other

length listed here).

Supplement figure S2. The prediction performance between MalariaSED and Enformer. We

present the auROC of each model as a point and use a box plot to show the distribution of the

models. We used 1280bp DNA sequence as the input to match the requirement of Enformer.

('***' indicates Wilcoxon test p < 0.01).

Supplement figure S3. TFs binding effects of single nucleotide substitutions at 5-mers

sequences predicted by MalariaSED in *P. falciparum.* '***' indicates Wilcoxon test compared

with whole-genome background p < 2.2e-16.

Supplement figure S4. MalariaSED prediction for epigenetic markers of ~1.3 millions variants

gathered by Pf3K. The horizontal and vertical direction represent, respectively, the predicted

probabilities that the sequences carrying the reference allele and alternative allele.

Supplement figure S5. The MalariaSED prediction results for de novo mutations at non-coding

regions discovered from the single cell study. '*' indicates Wilcoxon test compared with non-

coding SNPs collected from MalariaGEN, p < 0.05.

SUPPLEMENTARY TABLES LEGENDS

Supplement table S1. MalariaSED model configuration.

Supplement table S2. The prediction performance of MalariaSED in the independent test sets.

Supplement table S3. The average SNS effects at each DNA motif through all predicted TF

binding sites.

Supplement table S4. The chromatin effects prediction from MalariaSED for all reported

genomic variants collected by malariaGEN.

Supplement table S5. The GO and MPMP enrichment for geographically differentiated variants

presenting high chromatin effects.

Supplement table S6. The chromatin effects prediction from MalairaSED for local eQTLs

reported in TRACKI study.

References

1   Zhu, L. *et al.* The origins of malaria artemisinin resistance defined by a genetic and
       transcriptomic background. *Nat Commun* **9**, 5158, doi:10.1038/s41467-018-07588-x
       (2018).
2   Gibbons, J. *et al.* Altered expression of K13 disrupts DNA replication and repair in
       Plasmodium falciparum. *BMC Genomics* **19**, 849, doi:10.1186/s12864-018-5207-7 (2018).
3   Wang, C., Adapa, S. R., Gibbons, J., Sutton, S. & Jiang, R. H. Punctuated chromatin states
       regulate Plasmodium falciparum antigenic variation at the intron and 2 kb upstream
       regions. *BMC Genomics* **17**, 652, doi:10.1186/s12864-016-3005-7 (2016).
4   Swamy, L., Amulic, B. & Deitsch, K. W. Plasmodium falciparum var gene silencing is
       determined by cis DNA elements that form stable and heritable interactions. *Eukaryot
       Cell* **10**, 530-539, doi:10.1128/EC.00329-10 (2011).
5   Santos, J. M. *et al.* Red Blood Cell Invasion by the Malaria Parasite Is Coordinated by the
       PfAP2-I Transcription Factor. *Cell Host Microbe* **21**, 731-741 e710,
       doi:10.1016/j.chom.2017.05.006 (2017).

6   Josling, G. A. *et al.* Dissecting the role of PfAP2-G in malaria gametocytogenesis. *Nat*
Commun **11**, 1503, doi:10.1038/s41467-020-15026-0 (2020).
7   Josling, G. A. *et al.* A Plasmodium Falciparum Bromodomain Protein Regulates Invasion
Gene Expression. *Cell Host Microbe* **17**, 741-751, doi:10.1016/j.chom.2015.05.009
(2015).
8   MalariaGen *et al.* An open dataset of Plasmodium falciparum genome variation in 7,000
worldwide samples. *Wellcome Open Res* **6**, 42, doi:10.12688/wellcomeopenres.16168.2
(2021).
9   Cerqueira, G. C. *et al.* Longitudinal genomic surveillance of Plasmodium falciparum
malaria parasites reveals complex genomic architecture of emerging artemisinin
resistance. *Genome Biol* **18**, 78, doi:10.1186/s13059-017-1204-4 (2017).
10   Wang, Z. *et al.* Genome-wide association analysis identifies genetic loci associated with
resistance to multiple antimalarials in Plasmodium falciparum from China-Myanmar
border. *Sci Rep* **6**, 33891, doi:10.1038/srep33891 (2016).
11   Takala-Harrison, S. *et al.* Independent emergence of artemisinin resistance mutations
among Plasmodium falciparum in Southeast Asia. *The Journal of infectious diseases* **211**,
670-679, doi:10.1093/infdis/jiu491 (2015).
12   Wendler, J. P. *et al.* A genome wide association study of Plasmodium falciparum
susceptibility to 22 antimalarial drugs in Kenya. *PLoS One* **9**, e96486,
doi:10.1371/journal.pone.0096486 (2014).
13   Borrmann, S. *et al.* Genome-wide screen identifies new candidate genes associated with
artemisinin susceptibility in Plasmodium falciparum in Kenya. *Sci Rep* **3**, 3318,
doi:10.1038/srep03318 (2013).
14   Miotto, O. *et al.* Multiple populations of artemisinin-resistant Plasmodium falciparum in
Cambodia. *Nat Genet* **45**, 648-655, doi:10.1038/ng.2624 (2013).
15   Van Tyne, D. *et al.* Identification and functional validation of the novel antimalarial
resistance locus PF10_0355 in Plasmodium falciparum. *PLoS Genet* **7**, e1001383,
doi:10.1371/journal.pgen.1001383 (2011).
16   Mu, J. *et al.* Plasmodium falciparum genome-wide scans for positive selection,
recombination hot spots and resistance to antimalarial drugs. *Nat Genet* **42**, 268-271,
doi:10.1038/ng.528 (2010).
17   Miotto, O. *et al.* Genetic architecture of artemisinin-resistant Plasmodium falciparum.
Nat Genet **47**, 226-234, doi:10.1038/ng.3189 (2015).
18   Mok, S. *et al.* Drug resistance. Population transcriptomics of human malaria parasites
reveals the mechanism of artemisinin resistance. *Science* **347**, 431-435,
doi:10.1126/science.1260403 (2015).
19   Pradhan, A. *et al.* Chemogenomic profiling of Plasmodium falciparum as a tool to aid
antimalarial drug discovery. *Sci Rep* **5**, 15930, doi:10.1038/srep15930 (2015).
20   Visscher, P. M. *et al.* 10 Years of GWAS Discovery: Biology, Function, and Translation.
Am J Hum Genet **101**, 5-22, doi:10.1016/j.ajhg.2017.06.005 (2017).
21   Avsec, Z. *et al.* Effective gene expression prediction from sequence by integrating long-
range interactions. *Nat Methods* **18**, 1196-1203, doi:10.1038/s41592-021-01252-x
(2021).

22 Zrimec, J. *et al.* Deep learning suggests that gene expression is encoded in all parts of a
co-evolving interacting gene regulatory structure. *Nat Commun* **11**, 6141,
doi:10.1038/s41467-020-19921-4 (2020).
23 Kelley, D. R. *et al.* Sequential regulatory activity prediction across chromosomes with
convolutional neural networks. *Genome Res* **28**, 739-750, doi:10.1101/gr.227819.117
(2018).
24 Zhou, J. *et al.* Deep learning sequence-based ab initio prediction of variant effects on
expression and disease risk. *Nat Genet* **50**, 1171-1179, doi:10.1038/s41588-018-0160-6
(2018).
25 Hollin, T. & Le Roch, K. G. From Genes to Transcripts, a Tightly Regulated Journey in
Plasmodium. *Front Cell Infect Microbiol* **10**, 618454, doi:10.3389/fcimb.2020.618454
(2020).
26 Callebaut, I., Prat, K., Meurice, E., Mornon, J. P. & Tomavo, S. Prediction of the general
transcription factors associated with RNA polymerase II in Plasmodium falciparum:
conserved features and differences relative to other eukaryotes. *BMC Genomics* **6**, 100,
doi:10.1186/1471-2164-6-100 (2005).
27 Campbell, T. L., De Silva, E. K., Olszewski, K. L., Elemento, O. & Llinas, M. Identification
and genome-wide prediction of DNA binding specificities for the ApiAP2 family of
regulators from the malaria parasite. *PLoS Pathog* **6**, e1001165,
doi:10.1371/journal.ppat.1001165 (2010).
28 Templeton, T. J. *et al.* Comparative analysis of apicomplexa and genomic diversity in
eukaryotes. *Genome Res* **14**, 1686-1695, doi:10.1101/gr.2615304 (2004).
29 Toenhake, C. G. *et al.* Chromatin Accessibility-Based Characterization of the Gene
Regulatory Network Underlying Plasmodium falciparum Blood-Stage Development. *Cell*
Host Microbe **23**, 557-569 e559, doi:10.1016/j.chom.2018.03.007 (2018).
30 Kaneko, I., Iwanaga, S., Kato, T., Kobayashi, I. & Yuda, M. Genome-Wide Identification of
the Target Genes of AP2-O, a Plasmodium AP2-Family Transcription Factor. *PLoS Pathog*
11, e1004905, doi:10.1371/journal.ppat.1004905 (2015).
31 Yuda, M., Iwanaga, S., Kaneko, I. & Kato, T. Global transcriptional repression: An initial
and essential step for Plasmodium sexual development. *Proc Natl Acad Sci U S A* **112**,
12824-12829, doi:10.1073/pnas.1504389112 (2015).
32 Shang, X. *et al.* A cascade of transcriptional repression determines sexual commitment
and development in Plasmodium falciparum. *Nucleic Acids Res* **49**, 9264-9279,
doi:10.1093/nar/gkab683 (2021).
33 Oehring, S. C. *et al.* Organellar proteomics reveals hundreds of novel nuclear proteins in
the malaria parasite Plasmodium falciparum. *Genome Biol* **13**, R108, doi:10.1186/gb-
2012-13-11-r108 (2012).
34 Rocamora, F. *et al.* Oxidative stress and protein damage responses mediate artemisinin
resistance in malaria parasites. *PLoS Pathog* **14**, e1006930,
doi:10.1371/journal.ppat.1006930 (2018).
35 Mok, S. *et al.* Artemisinin-resistant K13 mutations rewire Plasmodium falciparum's intra-
erythrocytic metabolic program to enhance survival. *Nat Commun* **12**, 530,
doi:10.1038/s41467-020-20805-w (2021).

36 Bowyer, P. W. *et al.* Variation in Plasmodium falciparum erythrocyte invasion
phenotypes and merozoite ligand gene expression across different populations in areas
of malaria endemicity. *Infect Immun* **83**, 2575-2582, doi:10.1128/IAI.03009-14 (2015).
37 Dia, A. *et al.* Single-genome sequencing reveals within-host evolution of human malaria
parasites. *Cell Host Microbe* **29**, 1496-1506 e1493, doi:10.1016/j.chom.2021.08.009
(2021).
38 MalariaGen *et al.* An open dataset of Plasmodium falciparum genome variation in 7,000
worldwide samples. *Wellcome Open Res* **6**, 42, doi:10.12688/wellcomeopenres.16168.1
(2021).
39 Rono, M. K. *et al.* Adaptation of Plasmodium falciparum to its transmission environment.
Nat Ecol Evol **2**, 377-387, doi:10.1038/s41559-017-0419-9 (2018).
40 Amaratunga, C. *et al.* Artemisinin-resistant Plasmodium falciparum in Pursat province,
western Cambodia: a parasite clearance rate study. *Lancet Infect Dis* **12**, 851-858,
doi:10.1016/S1473-3099(12)70181-0 (2012).
41 Nkhoma, S. C. *et al.* Co-transmission of Related Malaria Parasite Lineages Shapes Within-
Host Parasite Diversity. *Cell Host Microbe* **27**, 93-103 e104,
doi:10.1016/j.chom.2019.12.001 (2020).
42 Snoek, J., Larochelle, H. & Adams, R. Practical bayesian optimization of machine learning
algorithms. *arXiv.*, 1206.2944 (2012).
43 Zhou, J. & Troyanskaya, O. G. Predicting effects of noncoding variants with deep
learning-based sequence model. *Nat Methods* **12**, 931-934, doi:10.1038/nmeth.3547
(2015).
670

Fig. 1

Fig. 2

Fig. 3

Fig. 4

A PfAP2-G Committed schizonts / PfAP2-G Sex rings / PfAP2-G Stage I Gametocytes

D *P. Berghei* PbAP2-G2 Gametocytes

B PfAP2-I Schizonts

C PfBDP1 — Trophozoites / Schizonts

E PbAP2-O Ookinetes

Binding effects of single-nucleotide substitutions

Fig. 5

Fig. 6

Fig. 7

A

AUC=0.545, H3K9ac_Trophozoites
AUC=0.568, H3K9ac_Schizonts
AUC=0.522, BDP1_Trophozoites
AUC=0.516, BDP1_Schizonts
AUC=0.541, AP2–I_Schizonts
AUC=0.530, AP2–G_Schizonts
AUC=0.512, AP2–G_Sex_rings
AUC=0.553, AP2–G_Gametocytes
AUC=0.524, AP2–G5_Trophozoites
AUC=0.700, ATAC–Seq_Rings
AUC=0.644, ATAC–Seq_Early_Trophozoites
AUC=0.549, ATAC–Seq_Late_Trophozoites
AUC=0.543, ATAC–Seq_Schizonts

B

AUC=0.503, H3K9ac_Trophozoites
AUC=0.416, H3K9ac_Schizonts
AUC=0.602, BDP1_Trophozoites
AUC=0.506, BDP1_Schizonts
AUC=0.603, AP2–I_Schizonts
AUC=0.633, AP2–G_Schizonts
AUC=0.581, AP2–G_Sex_rings
AUC=0.591, AP2–G_Gametocytes
AUC=0.626, AP2–G5_Trophozoites
AUC=0.735, ATAC–Seq_Rings
AUC=0.668, ATAC–Seq_Early_Trophozoites
AUC=0.643, ATAC–Seq_Late_Trophozoites
AUC=0.660, ATAC–Seq_Schizonts

Fig. 8