Susceptibility of *Pseudomonas aeruginosa* and *Acinetobacter baumannii* biofilms grown on denture acrylic to denture-cleaning agents and sonication

Marwah Behbehani [1], Ailbhe McDonald [1], Sean P Nair [3], Ingrid M Green [2,3] *

[1] Department of Prosthodontics, UCL Eastman Dental Institute, 21 University Street, London, WC1E 6DE

[2] Microbiology Unit, Eastman Dental Hospital, UCLH NHS Foundation Trust, UCL Royal Free Campus, Rowland Hill Street, London NW3 2PF

[3] Department of Microbial Diseases, UCL Eastman Dental Institute, Royal Free Campus, Rowland Hill Street, London NW3 2PF

Corresponding author: e-mail address: i.green@ucl.ac.uk

Acknowledgments

This work was undertaken at University College London Hospitals/University College Hospital (UCLH/UCL), which received a proportion of funding from the Department of Health's National Institute for Health Research (NIHR) Biomedical Research Centres Funding Scheme, UK.

Abstract

Background

Dentures can act as reservoirs for microorganisms that cause local and systemic diseases, including opportunistic pathogens *Pseudomonas aeruginosa* and *Acinetobacter spp*, leading causes of hospital-acquired pneumonia.

Objectives

To determine the efficacy of a brief cleaning protocol with four commonly used denture cleaning agents combined with ultra-sonic cleaning on *P. aeruginosa* and *A. baumannii* biofilms grown on denture acrylic. The effectiveness of a commercial sonicator on bacterial biofilm reduction was also evaluated.

Materials and Methods

P. aeruginosa and *A. baumannii* 48 h biofilms were grown in artificial saliva on polymethyl methacrylate (PMMA) discs, exposed to four disinfectants, with or without ultra-sonication, for 1 min and the number of viable bacteria post-exposure to the treatment determined. *P. aeruginosa* biofilms were exposed for one or five mins to two disinfectants plus a commercial sonicator.

Results

Ultra-sonication alone reduced total viable biofilm from 67.3% (*A. baumannii)* to 94.8% (*P. aeruginosa).* Without ultra-sonication, 0.012% sodium hypochlorite, Steradent® Active Plus and Ecolab® 0.2% chlorhexidine reduced *P. aeruginosa* and *A. baumannii* biofilms viable counts by between 99.3% and >99.99%. Sonication further reduced 99.3% kill to undetectable levels (>99.99% kill). Colgate®Total Pro-Gum Health Daily mouthwash reduced *P. aeruginosa* by 17% and *A. baumannii*

46 biofilms by 2.1 log. These number were further reduced by >99.99% with sonication

47 (p=<0.001). Domestic sonicators, such as those for cleaning jewellery at home, were

48 effective when used in conjunction with denture cleaning agents.

49 **Conclusion**

50 Rapid reduction of *P. aeruginosa* and *A. baumannii* biofilms on denture acrylic is

51 possible with denture-cleaning agents and sonication

Keywords:

Pseudomonas aeruginosa; *Acinetobacter baumannii*; denture; biofilm; ultrasonic cleaning; disinfection

Abbreviations:

CHX, chlorohexidine digluconate; NaOCl, sodium hypochlorite; AP, alkaline peroxide; CPC, cetylpyridinium chloride; PMMA, polymethylmethacrylate; PBS, phosphate-buffered saline; CFU, colony forming units.

Highlights

- Four antimicrobial agents tested against *P. aeruginosa* and *A. baumannii* biofilms
- 0.012% NaOCl and Steradent Active Plus were the most effective agents tested
- Sonication improved chlorhexidine and cetylpyridinium chloride-based agents

Introduction

Pseudomonas aeruginosa and Acinetobacter baumannii are two important Gram-negative nosocomial pathogens, which pose an increasing challenge to healthcare providers [1,2]. They are leading causes of hospital acquired pneumonia and bacteraemia [1,3], and are associated with increased mortality rates [4]. P. aeruginosa and A. baumannii are adept at biofilm formation on skin, soft tissue and wounds, as well as abiotic surfaces including endotracheal tubes [5,6] Bacteria within biofilms are protected from hostile environments, such as antimicrobials, via a number of means such the biofilm neutralising antimicrobials by binding to them, the role of the exopolysaccharide matrices of a biofilm in limiting the rate of diffusion of the antimicrobial into the biofilm, and, as a consequence of oxygen deprivation within biofilms, a lower metabolic activity of bacteria can reduce the uptake of antimicrobial agents [7].

Dentures, worn by 20% of the total UK population [8], have been shown to harbour biofilms that can act as reservoirs of respiratory pathogens such as P. aeruginosa and Acinetobacter spp [9,10]. P. aeruginosa and A. baumannii have secretion systems and efflux pumps capable of the removal of a wide variety of antibiotics, biocides detergents and antiseptics from the bacterial cell [11,12] which contributes to the challenge of biofilm removal. Moreover, P. aeruginosa has the ability to actively detach individual cells from its biofilm, a method called dispersal [13] implying that denture biofilms are a potential rich source for cross-infection, auto-infection and re-infection [14], with potential serious implications, particularly when patients are hospitalised and intubated.

Due to the topography of denture fitting surfaces, there is evidence that the use of both mechanical and chemical methods of disinfection is required for adequate

cleaning and the removal of pathogenic or opportunistic bacteria from dentures [15].

Mechanical cleaning with an ultrasonic device can be useful for physically and

cognitively impaired patients [16] and sonication has been demonstrated to be an

effective remover of Pseudomonas biofilms [17]. However, laboratory grade sonicators

are expensive, potentially hindering their use. Domestic sonicators, such as those

used for home cleaning of jewellery, are more affordable and could be beneficial for

denture cleaning both within hospitals and nursing homes, as well as for personal

use.

The aim of this study was to determine whether commonly used denture cleaning

agents, when combined with the mechanical effect of sonication, are effective at

removing *P. aeruginosa* and *A. baumannii* biofilms grown on a commonly used

denture material, polymethylmethacrylate (PMMA).

Materials and Methods

Denture Cleaning Agents and Sonication

P. aeruginosa (FR01) and *A. baumannii* (NCTC 1256) were grown in artificial saliva

[18] at 37°C for 16 h with shaking at 225 rpm. The bacterial inoculums were diluted

with artificial saliva to obtain an optical density (OD) of 0.05 at 600_{nm}.

Polymethylmethacrylate (PMMA) heat-cured denture resin discs (Lucitone 199®,

Denstply, UK), 10 mm in diameter were fabricated (cured for 70°C for 6 h, 100°C for

4 h in a water-bath), and then polished (350-Grit, SiC Paper, Struers, UK; 500-Grit,

Chaperline & Jacobs Ltd, Sutton, UK) to simulate the fit surface of the denture.

Surface roughness (R_a μm) was verified by measuring 5 randomly selected points on

10 randomly selected discs, using a Proscan 1000 scanning laser profilometer

(Scantron Industrial Products Ltd, England), using the displacement probe (KL131A)

at 780_{nm} with a beam diameter of 15 - 35 μm and resolution of 0.02 μm. The R_a

value was measured in both X and Y directions.

The discs were sterilised by autoclaving at 121.5°C for 15 min and positioned

aseptically on light body silicone (Aquasil Ultra LV, Dentsply, USA) in sterile 12-well

plates, exposing only the polished surface of the discs. To each well, 3 mL of the *P.*

aeruginosa or the *A. baumannii* inoculum (OD 0.05) was added and incubated for 48

h at 37°C. Discs were aseptically removed after 48 h and dipped briefly in sterile

PBS (Sigma-Aldrich, Merck Lifesciences Ltd, Dorset, UK), to remove any planktonic

bacteria after which they were placed into 3 mL of one of the test agents: Steradent®

Active Plus tablets (Reckitt Benckiser Ltd, Swindon, UK); 0.012% sodium

hypochlorite (NaOCl) (Evans, Vanodine International PLC, UK); Ecolab®

Chlorhexidine mouthwash (0.2% w/v Chlorhexidine digluconate) (CHX) (Ecolab®

Ltd, Swindon, UK); Colgate® Total Pro Gum Health Daily Mouthwash, (0.075% w/v

Cetylpyridinium chloride) (CPC) (Colgate-Palmolive, UK) or sterile phosphate-

buffered saline (PBS). Using sterile distilled water, Steradent® Active Plus was

prepared according to the manufacturer's instructions and Evans bleach was diluted

from the 4% w/v stock solution to achieve 0.012% w/v. Ecolab® CHX mouthwash

and the Colgate® Total CPC mouthwash were used as supplied by the

manufacturer. Half of the discs were sonicated in the active agents for 1 min using

an ultra-sonic water-bath (Fisher Scientific, Germany), the remainder were not

subjected to sonication but were exposed to the cleaning agents alone for 1 min. All

discs were subsequently placed into 1 mL of neutralising buffer (Becton Dickinson,

Detroit, MI, USA) and vortexed with glass beads (2.5 - 3.5 mm; VWR, Leicestershire,

UK) for 1 min to disrupt the biofilm. The samples were serial diluted, the *A.*

baumannii plated onto nutrient agar (Oxoid Ltd, Basingstoke, UK) and *P. aeruginosa*

plated on to pseudomonas plates made with pseudomonas agar powder (Oxoid Ltd,

Basingstoke, UK), supplemented with 1% glycerol (Sigma-Aldrich, Germany) and

1.2% agar technical (Oxoid Ltd, UK). The plates were incubated for 24 h at 37°C and

the colonies counted allowing the number and percentage of bacteria killed by

different antimicrobial agents and sonication regimes to be calculated.

Domestic Sonicator Efficacy

The efficacy of a domestic sonicator (Ultra 7000, James Products, UK) in reducing

viable biofilms, was compared to a laboratory ultra-sonic water-bath (FB 15047,

Fisher Scientific, Germany) in conjunction with anti-microbial denture agents. To do

this, *P. aeruginosa* 48 h biofilms were grown on PMMA discs as above. The discs

were then removed aseptically and dipped briefly in sterile PBS to remove any

planktonic bacteria prior to being immersed in 3 mL of one of the following agents:

Ecolab® 0.2% CHX mouthwash, Colgate® Total Pro Mouthwash (CPC) or sterile

PBS whilst being sonicated for 1 min or 5 min in either the laboratory ultrasonic

water bath or the domestic sonicator. Control discs were subjected to the anti-

microbial agents but not sonicated. The discs were then submersed in neutralising

broth with glass beads, vortexed for 1 min and then serial diluted prior to being

plated on to pseudomonas agar plates, and incubated for 24 h. The colonies were

counted, and the number and percentage of bacteria killed was calculated. All

experiments were performed in triplicate with three biological repeats.

Statistical Analysis

Data were analysed using SPSS version 24 (SPSS Inc., Chicago, IL) using non-

parametric tests, Kruskal-Wallis and Mann-Witney, to determine the number of viable

bacteria post exposure to the antimicrobial test agents and sonication. The 5% level

of statistical significance was used throughout the analyses.

Results

Surface roughness (Ra) and biofilm formation on PMMA discs

The arithmetic mean surface roughness of 10 acrylic discs was 5.2 μm (4.3 to 7.3

μm). See Fig 1A. *P. aeruginosa* and *A. baumannii* biofilms had an average CFU mL$^-$

^{1} per disc of 2.6 x 10^6 and 3.87 x 10^5 respectively. See Fig 1B.

Susceptibility of biofilms to denture cleaning agents, with or without

sonication

The 48 h biofilms were exposed to denture cleaning agents for 1 min. When

compared to the PBS control, 0.012% NaOCl and Steradent® were the most

effective anti-microbial agents evaluated, significantly reducing the median viable

number of both *P. aeruginosa* and *A. baumannii* CFUs mL^{-1} by over 99.9% or 5.22

and 3.59 log respectively (p = <0.001). Ecolab® CHX mouthwash reduced viable *P.*

aeruginosa biofilm numbers from 3.35 x 10^6 to 1 x 10^4 CFU mL^{-1} although this was

not significant (p = 0.269), however it was able to significantly reduce *A. baumannii*

viable counts from 7.95 x 10^4 to <2 x 10^1 CFU mL^{-1} (p = 0.001). Colgate® Total CPC

mouthwash reduced the number of viable *P. aeruginosa* CFUs mL^{-1} by 17% (3.35 x

10^6 to 1.5 x 10^6 CFU mL^{-1}) which was not statistically significant (*p* = 0.971), nor was

its 2.1 log reduction of viable bacteria of the *A. baumannii* biofilms (*p*=0.945). See

Fig 2.

Sonication alone with the laboratory ultra-sonicator reduced biofilm viability counts

by greater than 1 log for *P. aeruginosa* (p = <0.001) (3.35 x 10^6 to 1.75 x 10^5 CFU

mL^{-1}) and half a log for *A. baumannii* (7.95 x 10^4 to 2.60 x 10^4 CFU mL^{-1}; p = 0.04).

When sonication was used in conjunction with an antimicrobial, this significantly

reduced the number of viable *P. aeruginosa* (3.33×10^6 to 1.2×10^2 CFU mL^{-1}; p =

<0.001) and *A. baumannii* CFU mL^{-1} (5.8×10^2 to $< 2 \times 10^1$ CFU mL^{-1}; p = <0.001)

when compared to the cleaning agent only.

Sonication with Ecolab® CHX of *P. aeruginosa* biofilms led to a significant reduction

in viable numbers compared to use of Ecolab® CHX only (1.0×10^4 to $<2 \times 10^1$ CFU

mL^{-1}; p = <0.001). However no significant reduction was observed for *A. baumannii*

biofilms as numbers were already very low with the Ecolab® CHX mouthwash alone

(p = 0.258). Sonication did not significantly enhance NaOCl's nor Steradent®'s kill

rate for *P. aeruginosa* (p = 1.00; p = 0.666 respectively) or *A. baumannii* (p = 0.730;

p = 0.436 respectively) biofilms as numbers were already very low or undetectable

from use of the cleaning agent alone.   See Fig 2.

On comparison of a domestic sonicator with the laboratory standard sonicator, there

was no significant difference in the numbers of *P. aeruginosa* killed in biofilms when

sonicated in PBS for either 1 min or 5 min (p = 0.190 and 0.113 respectively) by

either machine. No difference between the sonicators could be determined with the

Ecolab® CHX mouthwash as the number of viable CFU mL^{-1} were below the

detection limit for both sonicators. When Colgate® Total CPC mouthwash was used

in conjunction with 1 min sonication, the laboratory sonicator was significantly more

effective than the domestic sonicator (p = <0.0001), however when the exposure

time was increased to 5 min, no significant difference (p = 0.297) was observed. See

Figure 3

212 Discussion

P. aeruginosa and *A. baumannii*, potential respiratory pathogens that are proficient

at biofilm formation, have been isolated from denture biofilms [10] and predispose

denture users to aspirational pneumonia and systemic inflammation [19] Controlling

denture plaque through an appropriate cleaning procedure is therefore important to

reduce this risk [20].

A combination of chemical and mechanical denture cleaning methods is widely

believed to be required for adequate denture cleaning [15,21], although there is a lack

of consensus in the literature and from randomized controlled clinical trials on which

of the available methods is best for disinfecting dentures [15]. There is little data on the

effectiveness of chemical disinfectants and/or ultrasonic cleaning against *P.*

aeruginosa denture biofilms and no published data available on disinfection of *A.*

baumannii biofilms from denture acrylic. This study aimed to determine how effective

short exposure times to denture cleaning agents and sonication would be against *P.*

aeruginosa and *A. baumannii* biofilms grown on denture acrylic, given that

healthcare and social-care staff are under increased pressure to deliver care with

limited resources. In addition, as the majority of denture wearers are elderly, often

with reduced manual dexterity affecting their ability to brush their dentures

thoroughly, a quick, accessible, and affordable 'hands-off' chemical and physical

cleaning method would encourage perseverance with cleaning routines to attain

thorough cleaning. This is particularly relevant, as up to 40% of denture wearers do

not remove their dentures at night increasing the risk of pneumonia 2.3-fold when

compared to those who do [19].

In this report, we tested four commonly used denture cleaning agents: 0.012%

NaOCl, Steradent® Active Plus, Ecolab® 0.2% chlorhexidine mouthwash (CHX) and

Colgate® Total Mouthwash 0.075% cetylpyridinium chloride (CPC) against 48 h *P.*

aeruginosa and *A. baumannii* biofilms grown on denture acrylic resin with and

without ultrasonic cleaning. NaOCl (0.012%) and Steradent® were very effective

(p=<0.001) in reducing the numbers of both pathogens below the level of detection

without sonication. Our results demonstrated that a short exposure time (1 min) to a

0.012% concentration of NaOCl led to significant reduction of viable CFU mL^{-1} of

biofilms formed on denture acrylic resin, results that concurred with studies that used

higher concentrations of NaOCl and longer exposure times. Salles et al. exposed *P.*

aeruginosa biofilms grown for 48 h on acrylic resin to 0.25% and 0.5% w/v NaOCl for

20 min, resulting in undetectable CFU counts [22]and Orsi et al. used even higher

concentrations of NaOCl (1% and 2% w/v) for between 5 to 15 min on *P. aeruginosa*

biofilms grown on acrylic resin, albeit with less mature (24 h) biofilms [23]. Kohler et al.

found that the *P. aeruginosa* MIC for NaOCl was significantly higher at 0.2% (w/v)

than for *A. baumannii* (0.1% w/v) and demonstrated that *P. aeruginosa* was less

susceptible to NaOCl than *A. baumannii* with 1 min contact time [24]. However, as their

work involved suspensions rather than biofilms, comparison with our study is difficult.

Our results demonstrated that short exposure times of 1 min to a lower concentration

of NaOCl (0.012%) led to a significant reduction of viable biofilms on denture acrylic

resin.

Steradent® Active Plus showed comparable kills to 0.012% NaOCl against *P.*

aeruginosa biofilms. This agrees with Mojarad et al. who showed that an alkaline

peroxide (AP) based denture cleaner resulted in complete disinfection of acrylic

dentures contaminated with *P. aeruginosa* 48 h biofilms, although their study had a

significantly longer exposure time of 15 mins [25]. Paranhos et al. also demonstrated

that a 5 min exposure to an effervescent alkaline peroxide-based tablet was effective

at significantly reducing *P. aeruginosa* CFU in 48 h biofilms grown on denture acrylic

resin [21]. Our study is the first to demonstrated that Steradent® Active Plus is capable

of significantly reducing *A. baumannii* biofilm CFUs on denture acrylic PMMA.

The Ecolab® 0.25% chlorhexidine mouthwash reduced the viable count of the *A.*

baumannii biofilms to levels comparable to 0.012% NaOCl and Steradent® but not

when use against *P. aeruginosa* biofilms. This contrasts with a study by Masadeh et

al, where they found that chlorhexidine mouthwashes, with and without ethanol,

showed some activity against *P. aeruginosa* biofilms but no activity against *A.*

baumannii biofilms that had been grown on polypropylene tubes [26]. The high Mg2+

content in the outer membrane of *P. aeruginosa*, compared to other organisms, may

confer intrinsic resistance of *P. aeruginosa* to chlorhexidine as chlorhexidine works

by binding to negatively charged sites on the bacterial cell wall [27] and could explain

the poor performance of these antimicrobial agents. The enhanced survival of *A.*

baumannii has been demonstrated to be due to upregulation of genes for the major

multi-drug efflux system, AdeAB, as well genes coding for an active chlorhexidine

efflux protein, AceI, when *A. baumannii* is exposed to chlorhexidine [28].

Although, ethanol, an ingredient in Ecolab® 0.2% Chlorhexidine mouthwash, is

known to be bactericidal against *P. aeruginosa* and *A. baumannii* in concentrations

of greater than 30% v/v [29,30] it has been shown to increase the virulence of *A.*

baumannii [31] and encourage biofilm formation in *P. aeruginosa* [32], particularly when

present in lower concentrations. However, resistance of *P. aeruginosa* to

antimicrobials has been related to several mechanisms, including enzymatic

inhibition, efflux systems, alteration of outer membrane permeability, and genetic

mutation [27].

The least effective agent in significantly reducing viable numbers of *P. aeruginosa*

and *A. baumannii* biofilms when compared with the control group (PBS) was

Colgate® Total Mouthwash, in which the active ingredient, CPC, an amphoteric

surfactant, works in a similar way to chlorhexidine. Similar results were

demonstrated by Masadeh et al who showed that a CPC based mouth wash

exhibited poor activity against *A. baumannii* 48 h biofilms [26]. There were no studies

in the literature that have tested antimicrobial activity of CHX mouthwashes and CPC

mouthwashes on *P. aeruginosa* or *A. baumannii* biofilms grown on dentures or

denture materials such as PMMA.

Despite being effective antimicrobial agents, NaOCl and Alkaline Peroxide (AP), an

active agent in Steradent, and CHX have been associated with several

disadvantages, depending on the concentration used and treatment time of the

denture materials. AP has also been shown to modify the colour of the acrylic

significantly, whereas NaOCl has been shown to change the surface hardness and

roughness, and noticeably alter the colour of the acrylic resin [33,34]. A low NaOH

concentration of 0.012% would have the benefit of reduced side effects whilst still

efficiently disinfecting dentures, although it would be unsuitable for use on partial

dentures, in which certain components are constructed from cobalt chrome.

CHX, a frequently used antimicrobial agent and a common ingredient in

mouthwashes, has been recognised as an increasing cause of anaphylaxis [35] and

penetrates denture acrylic when used over a prolonged period of time [36]. This

causes staining of the denture materials but also, severe cytotoxicity to human

gingival fibroblasts [36]. A study by Röhner et al. compared 0.05% CHX with a 0.08%

solution of NaOCl and found that not only was it less effective at eradicating

Staphylococcus aureus, S. epidermidis and *Pseudomonas aeruginosa* biofilms from

polyethlyene, but it also caused more cytotoxicity to human chondrocytes than

NaOCl [37]. In addition, an active ingredient of many mouthwashes, including Ecolab®

0.2% chlorhexidine, is ethanol, posing a hypothetical risk from repeated use of the

development of oral cancer [38].

Ultrasonic cleaning is the use of high energy ultrasound waves in an aqueous

solution to clean different objects. In this study, 1 min sonication with the control

solution (PBS) was not as effective as chemical cleaning alone; however, when

combined with Ecolab® CHX and Colgate® Total CPC mouthwashes, sonication

significantly enhanced the ability of both agents to reduce viable counts (p =

<0.0001). This has been supported by a previous study by Muqbil et al. which

demonstrated the superiority of ultrasonic cleaning when combined with chemical

cleaning compared to isolated treatment on *P. aeruginosa* biofilms [39]. Our results

support those of Nishi et al. who demonstrated the insufficiency of ultrasonic

cleaning with water against Candida and bacterial spp. However, when combined

with a chemical disinfectant, the resultant killing was higher than chemical cleaning

alone [40], suggesting that it is not only the mechanical action of the ultrasonic waves

that are responsible for killing but the combination of chemical and mechanical

cleaning.

One of the limitations of laboratory or industrial sonicators is their cost. A less

powerful domestic sonicator, such as those used for cleaning jewellery and personal

appliances would be a more economical alternative, especially for personal home

use. We tested a domestic sonicator, on its own and combined with Ecolab® CHX or

Colgate® Total CPC mouthwashes, against *P. aeruginosa* biofilms. When combined

with Ecolab® CHX there was no difference between the domestic and laboratory

sonicator in the reduction of viable counts. However, with Colgate® Total CPC, a

longer treatment time (5 min) was necessary to achieve a significant reduction in

CFU ml^{-1}. The domestic sonicator could therefore be an alternative to the laboratory

sonicator when combined with chemical agents and longer treatment times,

especially where loss of dexterity results in poor denture brushing and cleaning.

345 Immersion in 0.12% NaOCl and Steradent® was statistically as effective as

346 combined ultrasonic cleaning techniques, this indicates that the use of immersion

347 alone with these denture cleaning agents is adequate for removal of *P. aeruginosa*

348 and *A. baumannii* biofilms on denture acrylic.

349 Conclusion

We have demonstrated that *P. aeruginosa* and *A. baumannii* biofilms are efficiently

removed with low concentrations of NaOCl as well as Steradent® Active Plus and

Ecolab® 0.2% CHX in conjunction with ultrasonic cleaning, in a short time frame of

just 1 min. This regime could be particularly effective in an in-patient environment.

With advancing age, a person's sight and dexterity tend to deteriorate making

brushing dentures challenging, thus the use of a domestic sonicator would aid

denture cleaning in an increasingly elderly population. As certain individuals are

reluctant to follow standard cleaning advice or remove their dentures overnight [19],

this short efficient cleaning regime would be particularly useful.

359 Conflict of interests

360 None declared.

361 Ethical approval

362 Not required.

References

References

1.   Gaynes R, Edwards JR. Overview of nosocomial infections caused by Gram-negative

bacilli. *Clinical Infectious Diseases*. 2005;41:848-854. Accessed August 24, 2020.

https://academic.oup.com/cid/article/41/6/848/2022258

2.   Roca I, Espinal P, Vila-Farrés X, Vila J, Stefani S, Santos Sanches I. The *Acinetobacter*

baumannii oxymoron: commensal hospital dweller turned pan-drug-resistant

menace. *Front Microbiol*. 2012;3(148):1-30. doi:10.3389/fmicb.2012.00148

3.   European Centre for Disease Prevention and Control. *Annual Epidemiological Report*

2016 – Healthcare-Associated Infections Acquired in Intensive Care Units.; 2018.

Accessed August 24, 2020. https://ecdc.europa.eu/en/publications-data/healthcare-

associated-infections-acquired-intensive-care-units-annual

4.   Wuerth BA, Bonnewell JP, Wiemken TL, Arnold FW. Trends in pneumonia mortality

rates and hospitalizations by organism, United States, 2002–2011. *Emerg Infect Dis*.

2016;22(9):1624-1627. doi:http://dx.doi.org/10.3201/eid2209.150680

5.   Drenkard E, Ausubel FM. Pseudomonas biofilm formation and antibiotic resistance

are linked to phenotypic variation. *Nature*. 2002;416:740-743.

doi:https://doi.org/10.1038/416740a

6.   Espinal P, Martí S, Vila J, Child JA. Effect of biofilm formation on the survival of

Acinetobacter baumannii on dry surfaces. *Journal of Hospital Infection*. 2012;80:56-

60. doi:10.1016/j.jhin.2011.08.013

7.   Walters III MC, Roe F, Bugnicourt A, Franklin MJ, Stewart PS. Contributions of

antibiotic penetration, oxygen limitation, and low metabolic activity to tolerance of

Pseudomonas aeruginosa biofilms to ciprofloxacin and tobramycin. *Antimicrobial*

Agent and Chemotherapy. 2003;47(1):317-323. doi:10.1128/AAC.47.1.317-323.2003

8.   The NHS Information Centre for Health and Social, Chenery V, Cooke P. *Adult Dental*

Health Survey 2009: England Key Findings. (O'Sullivan I, Lader D, eds.).; 2011.

9.   O'Donnell LE, Smith K, Williams C, et al. Dentures are a reservoir for respiratory

pathogens. *Journal of Prosthodontics*. 2016;25(2):99-104. doi:10.1111/jopr.12342

10.   Sumi Y, Miura H, Sunakawa M, Michiwaki Y, Sakagami N. Colonization of denture

plaque by respiratory pathogens in dependent elderly. *Gerodontology*. 2002;19(1):25-

29. doi:10.1111/j.1741-2358.2002.00025.x

11.   Levy SB. Active efflux, a common mechanism for biocide and antibiotic resistance.

Journal of Applied Microbiology Symposium Supplement. 2002;92(1):65-71.

doi:10.1046/j.1365-2672.92.5s1.4.x

12.   Eze EC, Chenia HY, el Zowalaty ME. *Acinetobacter baumannii* biofilms: effects of

physicochemical factors, virulence, antibiotic resistance determinants, gene

regulation, and future antimicrobial treatments. *Infect Drug Resist*. 2018;11:2277-

2299. doi:10.2147/IDR.S169894

13.   Kim SK, Lee JH. Biofilm dispersion in *Pseudomonas aeruginosa*. *Journal of*

Microbiology. 2016;54(2):71-85. doi:10.1007/s12275-016-5528-7

14.   Ramage G, Jose A, Coco B, et al. Commercial mouthwashes are more effective than

azole antifungals against *Candida albicans* biofilms in vitro. *Oral Surg Oral Med Oral*

Pathol Oral Radiol Endod. 2011;111(4):456-460. doi:10.1016/j.tripleo.2010.10.043

15.   Souza de R, Freitas Oliveira Paranhos de H, da Silva LC, et al. Cochrane Library

Cochrane Database of Systematic Reviews Interventions for cleaning dentures in

adults (Review). Published online 2009. doi:10.1002/14651858.CD007395.pub2

16.   Duyck J, Vandamme K, Krausch-Hofmann S, et al. Impact of denture cleaning method

and overnight storage condition on denture biofilm mass and composition: A cross-

over randomized clinical trial. *PLoS One*. 2016;11(1):e0145837.

doi:10.1371/journal.pone.0145837

17.   Karbysheva S, Luca M di, Butini ME, Winkler T, Schü Tz M, Id AT. Comparison of

sonication with chemical biofilm dislodgement methods using chelating and reducing

agents: Implications for the microbiological diagnosis of implant associated infection.

PLoS One. 2020;15(4). doi:10.1371/journal.pone.0231389

18.   Badawi H, Evans RD, Wilson M, Ready D, Noar JH, Pratten J. The effect of orthodontic

bonding materials on dental plaque accumulation and composition in vitro.

Biomaterials. 2003;24(19):3345-3350.

19.   Iinuma T, Arai Y, Abe Y, et al. Denture wearing during sleep doubles the risk of

pneumonia in the very elderly. *JDR Clinical Research*. 2015;93:28S-36S.

doi:10.1177/0022034514552493

20.   Abe S, Ishihara K, Okuda K. Prevalence of potential respiratory pathogens in the

mouths of elderly patients and effects of professional oral care. *Arch Gerontol*

Geriatr. 2001;32(1):45-55. doi:10.1016/S0167-4943(00)00091-1

21.   Paranhos HFO, Silva-Lovato CH, de Souza RF, et al. Effect of three methods for

cleaning dentures on biofilms formed in vitro on acrylic resin. *Journal of*

Prosthodontics. 2009;18(5):427-431. doi:10.1111/j.1532-849X.2009.00450.x

22.   Salles MM, Oliveira VDC, Souza RF de, Silva-Lovato CH, de Freitas Oliveira Paranhos H.

Antimicrobial action of sodium hypochlorite and castor oil solutions for denture

cleaning-in vitro evaluation. *Original Research Prosthesis*. 2015;29(1):1-6.

doi:10.1590/1807-3107BOR-2015.vol29.0104

23.   Orsi IA, Junior AG, Villabona CA, Fernandes FHCN, Ito IY. Evaluation of the efficacy of

434   chemical disinfectants for disinfection of heat-polymerised acrylic resin.

435   *Gerodontology*. 2011;28(4):253-257. doi:10.1111/j.1741-2358.2010.00400.x

436   24.   Köhler AT, Rodloff AC, Labahn M, Reinhardt M, Truyen U, Speck S. Efficacy of sodium

437   hypochlorite against multidrug-resistant Gram-negative bacteria. *Journal of Hospital*

438   *Infection*. 2018;100:e40-e46. doi:10.1016/j.jhin.2018.07.017

439   25.   Mojarad N, Khalili Z, Aalaei S. A comparison of the efficacy of mechanical, chemical,

440   and microwave radiation methods in disinfecting complete dentures. *Dent Res J*

441   *(Isfahan)*. 2017;14(2):131-136. doi:10.4103/1735-3327.205793

442   26.   Masadeh MM, Gharaibeh SF, Alzoubi KH, Al-Azzam SI, Obeidat WM. Antimicrobial

443   activity of common mouthwash solutions on multidrug-resistance bacterial biofilms.

444   *Original Article J Clin Med Res •*. 2013;5(5):389-394. doi:10.4021/jocmr1535w

445   27.   McDonnell G, Russell AD. Antiseptics and disinfectants: activity, action, and

446   resistance. *Clin Microbiol Rev*. 1999;12(1):147-179. Accessed June 30, 2020.

447   http://cmr.asm.org/

448   28.   Hassan KA, Jackson SM, Penesyan A, et al. Transcriptomic and biochemical analyses

449   identify a family of chlorhexidine efflux proteins. *Proceedings of the National*

450   *Academy of Sciences*. 2013;110(50):20254-20259. doi:10.1073/pnas.1317052110

451   29.   Centers for Disease Control and Prevention (CDC). Guideline for disinfection and

452   sterilization in healthcare facilities. Infection Control Guidelines, CDC. Published 2008.

453   https://www.cdc.gov/infectioncontrol/guidelines/disinfection/disinfection-

454   methods/chemical.html

455   30.   Chiang SR, Jung F, Tang HJ, et al. Desiccation and ethanol resistances of multidrug

456   resistant *Acinetobacter baumannii* embedded in biofilm: The favorable antiseptic

efficacy of combination chlorhexidine gluconate and ethanol. *Journal of Microbiology,*

Immunology and Infection. 2018;51(6):770-777. doi:10.1016/j.jmii.2017.02.003

31.   Nwugo CC, Arivett BA, Zimbler DL, Gaddy JA, Richards AM. Effect of Ethanol on

Differential Protein Production and Expression of Potential Virulence Functions in the

Opportunistic Pathogen *Acinetobacter baumannii. PLoS One*. 2012;7(12):51936.

doi:10.1371/journal.pone.0051936

32.   Tashiro Y, Inagaki A, Ono K, et al. Bioscience, biotechnology, and biochemistry low

concentrations of ethanol stimulate biofilm and pellicle formation in *Pseudomonas*

aeruginosa. 2014;78(1):178-181. doi:10.1080/09168451.2014.877828

33.   Paranhos HDFO, Peracini A, Pisani MX, et al. Color stability, surface roughness and

flexural strength of an acrylic resin submitted to simulated overnight immersion in

denture cleansers. *Braz Dent J*. 2013;24(2):152-156. doi:10.1590/0103-

6440201302151

34.   Kurt A, Erkose-Genc G, Uzun M, Sarı T, Isik-Ozkol G. The effect of cleaning solutions

on a denture base material: Elimination of *Candida albicans* and alteration of physical

properties. *Journal of Prosthodontics*. 2018;27(6):577-583. doi:10.1111/jopr.12539

35.   Odedra KM, Farooque S. Chlorhexidine: an unrecognised cause of anaphylaxis.

Postgrad Med J. 2014;90(1070):709-714. doi:10.1136/postgradmedj-2013-132291

36.   Procópio ALF, da Silva RA, Maciel JG, et al. Antimicrobial and cytotoxic effects of

denture base acrylic resin impregnated with cleaning agents after long-term

immersion. *Toxicology in Vitro*. 2018;52:8-13. doi:10.1016/j.tiv.2018.05.012

37.   Röhner E, Jacob B, Böhle S, et al. Sodium hypochlorite is more effective than

chlorhexidine for eradication of bacterial biofilm of staphylococci and *Pseudomonas*

aeruginosa. *Knee Surgery, Sports Traumatology, Arthroscopy*. Published online 2020.

doi:10.1007/s00167-020-05887-9

38. Werner CWDA, Seymour RA. Are alcohol containing mouthwashes safe? *Br Dent J*.

2009;207(10):1-4. doi:10.1038/sj.bdj.2009.1014

39. Muqbil I, Burke FJT, Miller CH, Palenik CJ. Antimicrobial activity of ultrasonic cleaners.

Journal of Hospital Infection. 2005;60(3):249-255. doi:10.1016/j.jhin.2004.11.017

40. Nishi Y, Seto K, Kamashita Y, Kaji A, Kurono A, Nagaoka E. Survival of microorganisms

on complete dentures following ultrasonic cleaning combined with immersion in

peroxide-based cleanser solution. *Gerodontology*. 2014;31(3):202-209.

doi:10.1111/ger.12027

Figures

Fig. 1A and 1B

493   *Figure 1A and 1B*

494   Fig. 2A and 2B

496   *Figure 2A and 2B*

497   Fig. 3

499   *Figure 3*

Figure Legends

Figure 1 (A)Surface roughness (Ra) of 10 PMMA acrylic discs. The bold line represents the median, the box represents the 1st and 3rd quartiles, and the whiskers represent 95% of observations. (B) Colony forming units (CFU) per mL of 48 h P. aeruginosa and A. baumannii biofilms counted on 10 PMMA discs. Values represent the mean of 10 biofilms. Error bars are the ± SD.

Figure 2 Survival of 48 h (A) *P. aeruginosa* and (B) *A. baumannii* biofilms after exposure to sodium hypochlorite (NaOCl), Ecolab chlorhexidine (Ecolab CHS), Colgate cetylpryidinium chloride (Colgate CPC), Steradent (STD) and PBS (control) with or without sonication, for 1 min. Light grey boxes represent exposure to denture cleaning agents only and dark grey boxes represent exposure to denture cleaning plus sonication. Each experiment was performed in triplicate with three biological repeats.

Figure 3 Survival of 48 h *P. aeruginosa* biofilms after exposure to Ecolab chlorhexidine (Ecolab CHS), Colgate cetylpryidinium chloride (Colgate CPC), and PBS (control) with sonication in either a commercial (COM) or laboratory (Lab) based sonicator for 1 min and 5 min. Dark grey boxes represent 1 min and light grey boxes represent 5 mins sonication times. The white box represents PBS without sonication as a control group. Each experiment was performed in triplicate with three biological repeats