Susceptibility of *Pseudomonas aeruginosa* and *Acinetobacter baumannii* biofilms grown on denture acrylic to denture-cleaning agents and sonication

Marwah Behbehani [1], Ailbhe McDonald [1], Sean P Nair [3], Ingrid M Green [2,3] *

[1] Department of Prosthodontics, UCL Eastman Dental Institute, 21 University Street, London, WC1E 6DE

[2] Microbiology Unit, Eastman Dental Hospital, UCLH NHS Foundation Trust, UCL Royal Free Campus, Rowland Hill Street, London NW3 2PF

[3] Department of Microbial Diseases, UCL Eastman Dental Institute, Royal Free Campus, Rowland Hill Street, London NW3 2PF

Corresponding author: e-mail address: i.green@ucl.ac.uk

Abstract

Dentures can harbour microorganisms that cause local and systemic diseases in denture wearers. Opportunistic pathogens, *Pseudomonas aeruginosa* and Acinetobacter *spp*, are amongst the bacterial species isolated from denture biofilms and are a leading cause of hospital-acquired pneumonia. This study investigated the disinfecting efficacy of four cleaning agents and their combined use with ultrasonic cleaning on *P. aeruginosa* and *A. baumannii* biofilms on denture acrylic. Biofilms were grown for 48 hr in artificial saliva on polymethyl methacrylate (PMMA) discs and then exposed to a range of disinfectants, with or without sonication, for 1 min. The disinfecting efficacy was determined by enumerating the number of viable bacteria post-exposure to the treatment regime. Sonication alone resulted in a 94.8% and 67.3% reduction of microorganisms for *P. aeruginosa* (p=<0.001) and *A. baumannii* (p=0.04) biofilms respectively. Without sonication, 0.012% sodium hypochlorite and Steradent® Active Plus reduced viable counts by >99.99% (p=<0.001) for *P. aeruginosa* and *A. baumannii* biofilms. As Ecolab® 0.2% chlorhexidine mouthwash reduced *A. baumannii* biofilms counts below detectable levels, no further reduction of bacteria was observed with sonication. Sonication reduced viable *P. aeruginosa* biofilms by Ecolab® 0.2% chlorhexidine from 1×10^{4} (99.3% kill) to undetectable levels (>99.99% kill) (p=<0.001). Colgate®Total Pro-Gum Health Daily mouthwash reduced *P. aeruginosa* by 17% and *A. baumannii* biofilms by 2.1 log ($p = 0.971$; $p=0.945$ respectively) but with sonication, these number were further reduced by >99.99% (p=<0.001). Domestic sonicators, such as those for cleaning jewellery at home, were effective when used in conjunction with denture cleaning agents.

Keywords:

Pseudomonas aeruginosa; *Acinetobacter baumannii*; denture; biofilm; ultrasonic cleaning; disinfection

Abbreviations:

CHX, chlorohexidine digluconate; NaOCl, sodium hypochlorite; AP, alkaline peroxide; CPC, cetylpyridinium chloride; PMMA, polymethylmethacrylate; PBS, phosphate-buffered saline; CFU, colony forming units.

Highlights

- Four antimicrobial agents tested against *P. aeruginosa* and *A. baumannii* biofilms
- 0.012% NaOCl and Steradent Active Plus were the most effective agents tested
- Sonication improved chlorhexidine and cetylpyridinium chloride-based agents

Introduction

Pseudomonas aeruginosa and *Acinetobacter baumannii* are two important Gram-negative nosocomial pathogens, which pose an increasing challenge to healthcare providers [1,2]. They are leading causes of hospital acquired pneumonia and bacteraemia [1,3], and are associated with increased mortality rates [4]. *P. aeruginosa* and *A. baumannii* are adept at biofilm formation on skin, soft tissue and wounds, as well as abiotic surfaces including endotracheal tubes [5,6] Bacteria within biofilms are protected from hostile environments, such as antimicrobials, via a number of means such the biofilm neutralising antimicrobials by binding to them, the role of the exopolysaccharide matrices of a biofilm in limiting the rate of diffusion of the antimicrobial into the biofilm, and, as a consequence of oxygen deprivation within biofilms, a lower metabolic activity of bacteria can reduce the uptake of antimicrobial agents [7].

Dentures, worn by 20% of the total UK population [8], have been shown to harbour biofilms that can act as reservoirs of respiratory pathogens such as *P. aeruginosa* and *Acinetobacter spp* [9,10]. *P. aeruginosa* and *A. baumannii* have secretion systems and efflux pumps capable of the removal of a wide variety of antibiotics, biocides detergents and antiseptics from the bacterial cell [11,12] which contributes to the challenge of biofilm removal. Moreover, *P. aeruginosa* has the ability to actively detach individual cells from its biofilm, a method called dispersal [13] implying that denture biofilms are a potential rich source for cross-infection, auto-infection and re-infection [14], with potential serious implications, particularly when patients are hospitalised and intubated.

Due to the topography of denture fitting surfaces, there is evidence that the use of both mechanical and chemical methods of disinfection is required for adequate

cleaning and the removal of pathogenic or opportunistic bacteria from dentures [15].

Mechanical cleaning with an ultrasonic device can be useful for physically and

cognitively impaired patients [16] and sonication has been demonstrated to be an

effective remover of Pseudomonas biofilms [17]. However, laboratory grade

sonicators are expensive, potentially hindering their use. Domestic sonicators, such

as those used for home cleaning of jewellery, are more affordable and could be

beneficial for denture cleaning both within hospitals and nursing homes, as well as

for personal use.

The aim of this study was to determine whether commonly used denture cleaning

agents, when combined with the mechanical effect of sonication, are effective at

removing *P. aeruginosa* and *A. baumannii* biofilms grown on a commonly used

denture material, polymethylmethacrylate (PMMA).

92 Materials and Methods

93 Denture Cleaning Agents and Sonication

P. aeruginosa (FR01) and *A. baumannii* (NCTC 1256) were grown in artificial saliva

[18] at 37°C for 16 hr with shaking at 225 rpm. The bacterial inoculums were diluted

with artificial saliva to obtain an optical density (OD) of 0.05 at 600_{nm}.

Polymethylmethacrylate (PMMA) heat-cured denture resin discs (Lucitone 199®,

Denstply, UK), 10 mm in diameter were fabricated (cured for 70°C for 6 hr, 100°C for

4 hr in a water-bath), and then polished (350-Grit, SiC Paper, Struers, UK; 500-Grit,

Chaperline & Jacobs Ltd, Sutton, UK) to simulate the fit surface of the denture.

Surface roughness (R_a μm) was verified by measuring 5 randomly selected points on

10 randomly selected discs, using a Proscan 1000 scanning laser profilometer

(Scantron Industrial Products Ltd, England), using the displacement probe (KL131A)

at 780_{nm} with a beam diameter of 15 - 35 µm and resolution of 0.02 µm. The R_a

value was measured in both X and Y directions.

The discs were sterilised by autoclaving at 121.5°C for 15 min and positioned

aseptically on light body silicone (Aquasil Ultra LV, Dentsply, USA) in sterile 12-well

plates, exposing only the polished surface of the discs. To each well, 3 mL of the *P.*

aeruginosa or the *A. baumannii* inoculum (OD 0.05) was added and incubated for 48

hr at 37°C. Discs were aseptically removed after 48 hr and dipped briefly in sterile

PBS (Sigma-Aldrich, Merck Lifesciences Ltd, Dorset, UK), to remove any planktonic

bacteria after which they were placed into 3 mL of one of the test agents: Steradent®

Active Plus tablets (Reckitt Benckiser Ltd, Swindon, UK); 0.012% sodium

hypochlorite (NaOCl) (Evans, Vanodine International PLC, UK); Ecolab®

Chlorhexidine mouthwash (0.2% w/v Chlorhexidine digluconate) (CHX) (Ecolab®

Ltd, Swindon, UK); Colgate® Total Pro Gum Health Daily Mouthwash, (0.075% w/v

Cetylpyridinium chloride) (CPC) (Colgate-Palmolive, UK) or sterile phosphate-

buffered saline (PBS). Using sterile distilled water, Steradent® Active Plus was

prepared according to the manufacturer's instructions and Evans bleach was diluted

from the 4% w/v stock solution to achieve 0.012% w/v. Ecolab® CHX mouthwash

and the Colgate® Total CPC mouthwash were used as supplied by the

manufacturer. Half of the discs were sonicated in the active agents for 1 min using

an ultra-sonic water-bath (Fisher Scientific, Germany), the remainder were not

subjected to sonication but were exposed to the cleaning agents alone for 1 min. All

discs were subsequently placed into 1 mL of neutralising buffer (Becton Dickinson,

Detroit, MI, USA) and vortexed with glass beads (2.5 - 3.5 mm; VWR, Leicestershire,

UK) for 1 min to disrupt the biofilm. The samples were serial diluted, the *A.*

baumannii plated onto nutrient agar (Oxoid Ltd, Basingstoke, UK) and *P. aeruginosa*

plated on to pseudomonas plates made with pseudomonas agar powder (Oxoid Ltd,

Basingstoke, UK), supplemented with 1% glycerol (Sigma-Aldrich, Germany) and

1.2% agar technical (Oxoid Ltd, UK). The plates were incubated for 24 hr at 37°C

and the colonies counted allowing the number and percentage of bacteria killed by

different antimicrobial agents and sonication regimes to be calculated.

Domestic Sonicator Efficacy

The efficacy of a domestic sonicator (Ultra 7000, James Products, UK) in reducing

viable biofilms, was compared to a laboratory ultra-sonic water-bath (FB 15047,

Fisher Scientific, Germany) in conjunction with anti-microbial denture agents. To do

this, *P. aeruginosa* 48 hr biofilms were grown on PMMA discs as above. The discs

were then removed aseptically and dipped briefly in sterile PBS to remove any

planktonic bacteria prior to being immersed in 3 mL of one of the following agents:

Ecolab® 0.2% CHX mouthwash, Colgate® Total Pro Mouthwash (CPC) or sterile

PBS whilst being sonicated for 1 min or 5 min in either the laboratory ultrasonic

water bath or the domestic sonicator. Control discs were subjected to the anti-

microbial agents but not sonicated. The discs were then submersed in neutralising

broth with glass beads, vortexed for 1 min and then serial diluted prior to being

plated on to pseudomonas agar plates, and incubated for 24 hr. The colonies were

counted, and the number and percentage of bacteria killed was calculated. All

experiments were performed in triplicate with three biological repeats.

Statistical Analysis

151   Data were analysed using SPSS version 24 (SPSS Inc., Chicago, IL) using non-

152   parametric tests, Kruskal-Wallis and Mann-Witney, to determine the number of viable

153 bacteria post exposure to the antimicrobial test agents and sonication. The 5% level

154 of statistical significance was used throughout the analyses.

Results

156 Surface roughness (Ra) and biofilm formation on PMMA discs

157 The arithmetic mean surface roughness of 10 acrylic discs was 5.2 µm (4.3 to 7.3

158 µm). See Fig 1A. *P. aeruginosa* and *A. baumannii* biofilms had an average CFU mL^{-1}

159 per disc of 2.6 x 10^6 and 3.87 x 10^5 respectively. See Fig 1B.

161 Susceptibility of biofilms to denture cleaning agents, with or without

162 sonication

The 48 hr biofilms were exposed to denture cleaning agents for 1 min. When

compared to the PBS control, 0.012% NaOCl and Steradent® were the most

effective anti-microbial agents evaluated, significantly reducing the median viable

number of both *P. aeruginosa* and *A. baumannii* CFUs mL^{-1} by over 99.9% or 5.22

and 3.59 log respectively (p = <0.001). Ecolab® CHX mouthwash reduced viable *P.*

aeruginosa biofilm numbers from 3.35 x 10^6 to 1 x 10^4 CFU mL^{-1} although this was

not significant (p = 0.269), however it was able to significantly reduce *A. baumannii*

viable counts from 7.95 x 10^4 to <2 x 10^1 CFU mL^{-1} (p = 0.001). Colgate® Total CPC

mouthwash reduced the number of viable *P. aeruginosa* CFUs mL^{-1} by 17% (3.35 x

10^6 to 1.5 x 10^6 CFU mL^{-1}) which was not statistically significant (*p* = 0.971), nor was

its 2.1 log reduction of viable bacteria of the *A. baumannii* biofilms (*p*=0.945). See

Fig 2.

Sonication alone with the laboratory ultra-sonicator reduced biofilm viability counts

by greater than 1 log for *P. aeruginosa* (p = <0.001) (3.35×10^6 to 1.75×10^5 CFU

mL^{-1}) and half a log for *A. baumannii* (7.95×10^4 to 2.60×10^4 CFU mL^{-1}; p = 0.04).

When sonication was used in conjunction with an antimicrobial, this significantly

reduced the number of viable *P. aeruginosa* (3.33×10^6 to 1.2×10^2 CFU mL^{-1}; p =

<0.001) and *A. baumannii* CFU mL^{-1} (5.8×10^2 to $< 2 \times 10^1$ CFU mL^{-1}; p = <0.001)

when compared to the cleaning agent only.

Sonication with Ecolab® CHX of *P. aeruginosa* biofilms led to a significant reduction

in viable numbers compared to use of Ecolab® CHX only (1.0×10^4 to $<2 \times 10^1$ CFU

mL^{-1}; p = <0.001). However no significant reduction was observed for *A. baumannii*

biofilms as numbers were already very low with the Ecolab® CHX mouthwash alone

(p = 0.258). Sonication did not significantly enhance NaOCl's nor Steradent®'s kill

rate for *P. aeruginosa* (p = 1.00; p = 0.666 respectively) or *A. baumannii* (p = 0.730;

p = 0.436 respectively) biofilms as numbers were already very low or undetectable

from use of the cleaning agent alone.   See Fig 2.

On comparison of a domestic sonicator with the laboratory standard sonicator, there

was no significant difference in the numbers of *P. aeruginosa* killed in biofilms when

sonicated in PBS for either 1 min or 5 min (p = 0.190 and 0.113 respectively) by

either machine. No difference between the sonicators could be determined with the

Ecolab® CHX mouthwash as the number of viable CFU mL^{-1} were below the

detection limit for both sonicators. When Colgate® Total CPC mouthwash was used

in conjunction with 1 min sonication, the laboratory sonicator was significantly more

effective than the domestic sonicator (p = <0.0001), however when the exposure

time was increased to 5 min, no significant difference (p = 0.297) was observed. See

Figure 3

Discussion

P. aeruginosa and *A. baumannii*, potential respiratory pathogens that are proficient at biofilm formation, have been isolated from denture biofilms [10] and predispose denture users to aspirational pneumonia and systemic inflammation [19] Controlling denture plaque through an appropriate cleaning procedure is therefore important to reduce this risk [20].

A combination of chemical and mechanical denture cleaning methods is widely believed to be required for adequate denture cleaning [15,21], although there is a lack of consensus in the literature and from randomized controlled clinical trials on which of the available methods is best for disinfecting dentures [15]. There is little data on the effectiveness of chemical disinfectants and/or ultrasonic cleaning against *P. aeruginosa* denture biofilms and no published data available on disinfection of *A. baumannii* biofilms from denture acrylic. This study aimed to determine how effective short exposure times to denture cleaning agents and sonication would be against *P. aeruginosa* and *A. baumannii* biofilms grown on denture acrylic, given that healthcare and social-care staff are under increased pressure to deliver care with limited resources. In addition, as the majority of denture wearers are elderly, often with reduced manual dexterity affecting their ability to brush their dentures thoroughly, a quick, accessible, and affordable 'hands-off' chemical and physical cleaning method would encourage perseverance with cleaning routines to attain thorough cleaning. This is particularly relevant, as up to 40% of denture wearers do not remove their dentures at night increasing the risk of pneumonia 2.3-fold when compared to those who do [19].

In this report, we tested four commonly used denture cleaning agents: 0.012%

NaOCl, Steradent® Active Plus, Ecolab® 0.2% chlorhexidine mouthwash (CHX) and

Colgate® Total Mouthwash 0.075% cetylpyridinium chloride (CPC) against 48 hr *P.*

aeruginosa and *A. baumannii* biofilms grown on denture acrylic resin with and

without ultrasonic cleaning. NaOCl (0.012%) and Steradent® were very effective

(p=<0.001) in reducing the numbers of both pathogens below the level of detection

without sonication. Our results demonstrated that a short exposure time (1 min) to a

0.012% concentration of NaOCl led to significant reduction of viable CFU mL^{-1} of

biofilms formed on denture acrylic resin, results that concurred with studies that used

higher concentrations of NaOCl and longer exposure times. Salles et al. exposed *P.*

aeruginosa biofilms grown for 48 hr on acrylic resin to 0.25% and 0.5% w/v NaOCl

for 20 min, resulting in undetectable CFU counts [22]and Orsi et al. used even higher

concentrations of NaOCl (1% and 2% w/v) for between 5 to 15 min on *P. aeruginosa*

biofilms grown on acrylic resin, albeit with less mature (24 hr) biofilms [23]. Kohler et

al. found that the *P. aeruginosa* MIC for NaOCl was significantly higher at 0.2% (w/v)

than for *A. baumannii* (0.1% w/v) and demonstrated that *P. aeruginosa* was less

susceptible to NaOCl than *A. baumannii* with 1 min contact time [24]. However, as

their work involved suspensions rather than biofilms, comparison with our study is

difficult.

Our results demonstrated that short exposure times of 1 min to a lower concentration

of NaOCl (0.012%) led to a significant reduction of viable biofilms on denture acrylic

resin.

Steradent® Active Plus showed comparable kills to 0.012% NaOCl against *P.*

aeruginosa biofilms. This agrees with Mojarad et al. who showed that an alkaline

peroxide (AP) based denture cleaner resulted in complete disinfection of acrylic

dentures contaminated with *P. aeruginosa* 48 hr biofilms, although their study had a

significantly longer exposure time of 15 minutes [25]. Paranhos et al. also

demonstrated that a 5 min exposure to an effervescent alkaline peroxide-based

tablet was effective at significantly reducing *P. aeruginosa* CFU in 48 hr biofilms

grown on denture acrylic resin [21]. Our study is the first to demonstrated that

Steradent® Active Plus is capable of significantly reducing *A. baumannii* biofilm

CFUs on denture acrylic PMMA.

The Ecolab® 0.25% chlorhexidine mouthwash reduced the viable count of the *A.*

baumannii biofilms to levels comparable to 0.012% NaOCl and Steradent® but not

when use against *P. aeruginosa* biofilms. This contrasts with a study by Masadeh et

al, where they found that chlorhexidine mouthwashes, with and without ethanol,

showed some activity against *P. aeruginosa* biofilms but no activity against *A.*

baumannii biofilms that had been grown on polypropylene tubes [26]. The high

Mg2+ content in the outer membrane of *P. aeruginosa*, compared to other

organisms, may confer intrinsic resistance of *P. aeruginosa* to chlorhexidine as

chlorhexidine works by binding to negatively charged sites on the bacterial cell wall

[27] and could explain the poor performance of these antimicrobial agents. The

enhanced survival of *A. baumannii* has been demonstrated to be due to upregulation

of genes for the major multi-drug efflux system, AdeAB, as well genes coding for an

active chlorhexidine efflux protein, AceI, when *A. baumannii* is exposed to

chlorhexidine [28].

Although, ethanol, an ingredient in Ecolab® 0.2% Chlorhexidine mouthwash, is

known to be bactericidal against *P. aeruginosa* and *A. baumannii* in concentrations

of greater than 30% v/v [29,30] it has been shown to increase the virulence of *A.*

baumannii [31] and encourage biofilm formation in *P. aeruginosa* [32], particularly

when present in lower concentrations. However, resistance of *P. aeruginosa* to

antimicrobials has been related to several mechanisms, including enzymatic

inhibition, efflux systems, alteration of outer membrane permeability, and genetic

mutation [27].

The least effective agent in significantly reducing viable numbers of *P. aeruginosa*

and *A. baumannii* biofilms when compared with the control group (PBS) was

Colgate® Total Mouthwash, in which the active ingredient, CPC, an amphoteric

surfactant, works in a similar way to chlorhexidine. Similar results were

demonstrated by Masadeh et al who showed that a CPC based mouth wash

exhibited poor activity against *A. baumannii* 48 hr biofilms [26]. There were no

studies in the literature that have tested antimicrobial activity of CHX mouthwashes

and CPC mouthwashes on *P. aeruginosa* or *A. baumannii* biofilms grown on

dentures or denture materials such as PMMA.

Despite being effective antimicrobial agents, NaOCl and Alkaline Peroxide (AP), an

active agent in Steradent, and CHX have been associated with several

disadvantages, depending on the concentration used and treatment time of the

denture materials. AP has also been shown to modify the colour of the acrylic

significantly, whereas NaOCl has been shown to change the surface hardness and

roughness, and noticeably alter the colour of the acrylic resin [33,34]. A low NaOH

concentration of 0.012% would have the benefit of reduced side effects whilst still

efficiently disinfecting dentures, although it would be unsuitable for use on partial

dentures, in which certain components are constructed from cobalt chrome.

CHX, a frequently used antimicrobial agent and a common ingredient in

mouthwashes, has been recognised as an increasing cause of anaphylaxis [35] and

penetrates denture acrylic when used over a prolonged period of time [36]. This

causes staining of the denture materials but also, severe cytotoxicity to human

gingival fibroblasts [36]. A study by Röhner et al. compared 0.05% CHX with a

0.08% solution of NaOCl and found that not only was it less effective at eradicating

Staphylococcus aureus, S. epidermidis and *Pseudomonas aeruginosa* biofilms from

polyethlyene, but it also caused more cytotoxicity to human chondrocytes than

NaOCl [37]. In addition, an active ingredient of many mouthwashes, including

Ecolab® 0.2% chlorhexidine, is ethanol, posing a hypothetical risk from repeated use

of the development of oral cancer [38].

Ultrasonic cleaning is the use of high energy ultrasound waves in an aqueous

solution to clean different objects. In this study, 1 min sonication with the control

solution (PBS) was not as effective as chemical cleaning alone; however, when

combined with Ecolab® CHX and Colgate® Total CPC mouthwashes, sonication

significantly enhanced the ability of both agents to reduce viable counts ($p =$

<0.0001). This has been supported by a previous study by Muqbil et al. which

demonstrated the superiority of ultrasonic cleaning when combined with chemical

cleaning compared to isolated treatment on *P. aeruginosa* biofilms [39]. Our results

support those of Nishi et al. who demonstrated the insufficiency of ultrasonic

cleaning with water against Candida and bacterial spp. However, when combined

with a chemical disinfectant, the resultant killing was higher than chemical cleaning

alone [40], suggesting that it is not only the mechanical action of the ultrasonic

waves that are responsible for killing but the combination of chemical and

mechanical cleaning.

One of the limitations of laboratory or industrial sonicators is their cost. A less

powerful domestic sonicator, such as those used for cleaning jewellery and personal

appliances would be a more economical alternative, especially for personal home

use. We tested a domestic sonicator, on its own and combined with Ecolab® CHX or

Colgate® Total CPC mouthwashes, against *P. aeruginosa* biofilms. When combined

with Ecolab® CHX there was no difference between the domestic and laboratory

sonicator in the reduction of viable counts. However, with Colgate® Total CPC, a

longer treatment time (5 min) was necessary to achieve a significant reduction in

CFU ml^{-1}. The domestic sonicator could therefore be an alternative to the laboratory

sonicator when combined with chemical agents and longer treatment times,

especially where loss of dexterity results in poor denture brushing and cleaning.

Immersion in 0.12% NaOCl and Steradent® was statistically as effective as

combined ultrasonic cleaning techniques, this indicates that the use of immersion

alone with these denture cleaning agents is adequate for removal of *P. aeruginosa*

and *A. baumannii* biofilms on denture acrylic.

We have demonstrated that *P. aeruginosa* and *A. baumannii* biofilms are efficiently

removed with low concentrations of NaOCl as well as Steradent® Active Plus and

Ecolab® 0.2% CHX in conjunction with ultrasonic cleaning, in a short time frame of

just 1 min. This regime could be particularly effective in an in-patient environment.

With advancing age, a person's sight and dexterity tend to deteriorate making

brushing dentures challenging, thus the use of a domestic sonicator would aid

denture cleaning in an increasingly elderly population. As certain individuals are

reluctant to follow standard cleaning advice or remove their dentures overnight [19],

this short efficient cleaning regime would be particularly useful.

351 Funding

352 This work was undertaken at University College London Hospitals/University College

353 Hospital (UCLH/UCL), which received a proportion of funding from the Department of

354 Health's National Institute for Health Research (NIHR) Biomedical Research Centres

355 Funding Scheme, UK.

356 Conflict of interests

357 None declared.

358 Ethical approval

359 Not required.

360 Acknowledgments

References

[1]   Gaynes R, Edwards JR. Overview of nosocomial infections caused by Gram-negative bacilli. Clinical Infectious Diseases 2005;41:848–54.

[2]   Roca I, Espinal P, Vila-Farrés X, Vila J, Stefani S, Santos Sanches I. The *Acinetobacter baumannii* oxymoron: commensal hospital dweller turned pan-drug-resistant menace. Front Microbiol 2012;3:1–30. https://doi.org/10.3389/fmicb.2012.00148.

[3]   European Centre for Disease Prevention and Control. Annual epidemiological report 2016 – Healthcare-associated infections acquired in intensive care units. Stockholm: 2018.

[4]   Wuerth BA, Bonnewell JP, Wiemken TL, Arnold FW. Trends in pneumonia mortality rates and hospitalizations by organism, United States, 2002–2011. Emerg Infect Dis 2016;22:1624–7. https://doi.org/http://dx.doi.org/10.3201/eid2209.150680.

[5]   Drenkard E, Ausubel FM. Pseudomonas biofilm formation and antibiotic resistance are linked to phenotypic variation. Nature 2002;416:740–3. https://doi.org/https://doi.org/10.1038/416740a.

[6]   Espinal P, Martí S, Vila J, Child JA. Effect of biofilm formation on the survival of *Acinetobacter baumannii* on dry surfaces. Journal of Hospital Infection 2012;80:56–60. https://doi.org/10.1016/j.jhin.2011.08.013.

[7]   Walters III MC, Roe F, Bugnicourt A, Franklin MJ, Stewart PS. Contributions of antibiotic penetration, oxygen limitation, and low metabolic activity to tolerance of *Pseudomonas aeruginosa* biofilms to ciprofloxacin and tobramycin. Antimicrobial Agent and Chemotherapy 2003;47:317–23. https://doi.org/10.1128/AAC.47.1.317-323.2003.

[8]   The NHS Information Centre for Health and Social, Chenery V, Cooke P. Adult dental health survey 2009: England key findings. 2011.

[9]   O'Donnell LE, Smith K, Williams C, Nile CJ, Lappin DF, Bradshaw D, et al. Dentures are a reservoir for respiratory pathogens. Journal of Prosthodontics 2016;25:99–104. https://doi.org/10.1111/jopr.12342.

[10]   Sumi Y, Miura H, Sunakawa M, Michiwaki Y, Sakagami N. Colonization of denture plaque by respiratory pathogens in dependent elderly. Gerodontology 2002;19:25–9. https://doi.org/10.1111/j.1741-2358.2002.00025.x.

[11]   Levy SB. Active efflux, a common mechanism for biocide and antibiotic resistance. Journal of Applied Microbiology Symposium Supplement 2002;92:65–71. https://doi.org/10.1046/j.1365-2672.92.5s1.4.x.

[12]   Eze EC, Chenia HY, el Zowalaty ME. *Acinetobacter baumannii* biofilms: effects of physicochemical factors, virulence, antibiotic resistance determinants, gene regulation, and future antimicrobial treatments. Infect Drug Resist 2018;11:2277–99. https://doi.org/10.2147/IDR.S169894.

[13]   Kim S-K, Lee J-H. Biofilm dispersion in *Pseudomonas aeruginosa*. Journal of Microbiology 2016;54:71–85. https://doi.org/10.1007/s12275-016-5528-7.

[14]   Ramage G, Jose A, Coco B, Rajendran R, Rautemaa R, Murray C, et al. Commercial mouthwashes are more effective than azole antifungals against *Candida albicans* biofilms in vitro. Oral Surg Oral Med Oral Pathol Oral Radiol Endod 2011;111:456–60. https://doi.org/10.1016/j.tripleo.2010.10.043.

[15]   Souza de R, Freitas Oliveira Paranhos de H, da Silva LC, Freitas de Souza R, de Freitas Oliveira Paranhos H, Lovato da Silva CH, et al. Cochrane Library

Cochrane Database of Systematic Reviews Interventions for cleaning dentures
in adults (Review) 2009. https://doi.org/10.1002/14651858.CD007395.pub2.

[16] Duyck J, Vandamme K, Krausch-Hofmann S, Boon L, de Keersmaecker K,
Jalon E, et al. Impact of denture cleaning method and overnight storage
condition on denture biofilm mass and composition: A cross-over randomized
clinical trial. PLoS One 2016;11:e0145837.
https://doi.org/10.1371/journal.pone.0145837.

[17] Karbysheva S, Luca M di, Butini ME, Winkler T, Schü Tz M, Id AT.
Comparison of sonication with chemical biofilm dislodgement methods using
chelating and reducing agents: Implications for the microbiological diagnosis of
implant associated infection. PLoS One 2020;15.
https://doi.org/10.1371/journal.pone.0231389.

[18] Badawi H, Evans RD, Wilson M, Ready D, Noar JH, Pratten J. The effect of
orthodontic bonding materials on dental plaque accumulation and composition
in vitro. Biomaterials 2003;24:3345–50.

[19] Iinuma T, Arai Y, Abe Y, Takayama M, Fukumoto M, Fukui Y, et al. Denture
wearing during sleep doubles the risk of pneumonia in the very elderly. JDR
Clinical Research 2015;93:28S-36S.
https://doi.org/10.1177/0022034514552493.

[20] Abe S, Ishihara K, Okuda K. Prevalence of potential respiratory pathogens in
the mouths of elderly patients and effects of professional oral care. Arch
Gerontol Geriatr 2001;32:45–55. https://doi.org/10.1016/S0167-
4943(00)00091-1.

[21] Paranhos HFO, Silva-Lovato CH, de Souza RF, Cruz PC, de Freitas-Pontes
KM, Watanabe E, et al. Effect of three methods for cleaning dentures on
biofilms formed in vitro on acrylic resin. Journal of Prosthodontics
2009;18:427–31. https://doi.org/10.1111/j.1532-849X.2009.00450.x.

[22] Salles MM, Oliveira VDC, Souza RF de, Silva-Lovato CH, de Freitas Oliveira
Paranhos H. Antimicrobial action of sodium hypochlorite and castor oil
solutions for denture cleaning-in vitro evaluation. Original Research Prosthesis
2015;29:1–6. https://doi.org/10.1590/1807-3107BOR-2015.vol29.0104.

[23] Orsi IA, Junior AG, Villabona CA, Fernandes FHCN, Ito IY. Evaluation of the
efficacy of chemical disinfectants for disinfection of heat-polymerised acrylic
resin. Gerodontology 2011;28:253–7. https://doi.org/10.1111/j.1741-
2358.2010.00400.x.

[24] Köhler AT, Rodloff AC, Labahn M, Reinhardt M, Truyen U, Speck S. Efficacy
of sodium hypochlorite against multidrug-resistant Gram-negative bacteria.
Journal of Hospital Infection 2018;100:e40–6.
https://doi.org/10.1016/j.jhin.2018.07.017.

[25] Mojarad N, Khalili Z, Aalaei S. A comparison of the efficacy of mechanical,
chemical, and microwave radiation methods in disinfecting complete dentures.
Dent Res J (Isfahan) 2017;14:131–6. https://doi.org/10.4103/1735-
3327.205793.

[26] Masadeh MM, Gharaibeh SF, Alzoubi KH, Al-Azzam SI, Obeidat WM.
Antimicrobial activity of common mouthwash solutions on multidrug-resistance
bacterial biofilms. Original Article J Clin Med Res • 2013;5:389–94.
https://doi.org/10.4021/jocmr1535w.

[27] McDonnell G, Russell AD. Antiseptics and disinfectants: activity, action, and
resistance. Clin Microbiol Rev 1999;12:147–79.

[28] Hassan KA, Jackson SM, Penesyan A, Patching SG, Tetu SG, Eijkelkamp BA,
et al. Transcriptomic and biochemical analyses identify a family of
chlorhexidine efflux proteins. Proceedings of the National Academy of
Sciences 2013;110:20254–9. https://doi.org/10.1073/pnas.1317052110.
[29] Centers for Disease Control and Prevention (CDC). Guideline for disinfection
and sterilization in healthcare facilities. Infection Control Guidelines, CDC
2008. https://www.cdc.gov/infectioncontrol/guidelines/disinfection/disinfection-
methods/chemical.html.
[30] Chiang SR, Jung F, Tang HJ, Chen CH, Chen CC, Chou HY, et al. Desiccation
and ethanol resistances of multidrug resistant *Acinetobacter baumannii*
embedded in biofilm: The favorable antiseptic efficacy of combination
chlorhexidine gluconate and ethanol. Journal of Microbiology, Immunology and
Infection 2018;51:770–7. https://doi.org/10.1016/j.jmii.2017.02.003.
[31] Nwugo CC, Arivett BA, Zimbler DL, Gaddy JA, Richards AM. Effect of Ethanol
on Differential Protein Production and Expression of Potential Virulence
Functions in the Opportunistic Pathogen *Acinetobacter baumannii*. PLoS One
2012;7:51936. https://doi.org/10.1371/journal.pone.0051936.
[32] Tashiro Y, Inagaki A, Ono K, Inaba T, Yawata Y, Uchiyama H, et al.
Bioscience, biotechnology, and biochemistry low concentrations of ethanol
stimulate biofilm and pellicle formation in *Pseudomonas aeruginosa*
2014;78:178–81. https://doi.org/10.1080/09168451.2014.877828.
[33] Paranhos HDFO, Peracini A, Pisani MX, Oliveira VDC, Freitas De Souza R,
Silva-Lovato H, et al. Color stability, surface roughness and flexural strength of
an acrylic resin submitted to simulated overnight immersion in denture
cleansers. Braz Dent J 2013;24:152–6. https://doi.org/10.1590/0103-
6440201302151.
[34] Kurt A, Erkose-Genc G, Uzun M, Sarı T, Isik-Ozkol G. The effect of cleaning
solutions on a denture base material: Elimination of *Candida albicans* and
alteration of physical properties. Journal of Prosthodontics 2018;27:577–83.
https://doi.org/10.1111/jopr.12539.
[35] Odedra KM, Farooque S. Chlorhexidine: an unrecognised cause of
anaphylaxis. Postgrad Med J 2014;90:709–14.
https://doi.org/10.1136/postgradmedj-2013-132291.
[36] Procópio ALF, da Silva RA, Maciel JG, Sugio CYC, Soares S, Urban VM, et al.
Antimicrobial and cytotoxic effects of denture base acrylic resin impregnated
with cleaning agents after long-term immersion. Toxicology in Vitro 2018;52:8–
13. https://doi.org/10.1016/j.tiv.2018.05.012.
[37] Röhner E, Jacob B, Böhle S, Rohe S, Löffler B, Matziolis G, et al. Sodium
hypochlorite is more effective than chlorhexidine for eradication of bacterial
biofilm of staphylococci and *Pseudomonas aeruginosa*. Knee Surgery, Sports
Traumatology, Arthroscopy 2020. https://doi.org/10.1007/s00167-020-05887-9.
[38] Werner CWDA, Seymour RA. Are alcohol containing mouthwashes safe? Br
Dent J 2009;207:1–4. https://doi.org/10.1038/sj.bdj.2009.1014.
[39] Muqbil I, Burke FJT, Miller CH, Palenik CJ. Antimicrobial activity of ultrasonic
cleaners. Journal of Hospital Infection 2005;60:249–55.
https://doi.org/10.1016/j.jhin.2004.11.017.
[40] Nishi Y, Seto K, Kamashita Y, Kaji A, Kurono A, Nagaoka E. Survival of
microorganisms on complete dentures following ultrasonic cleaning combined
with immersion in peroxide-based cleanser solution. Gerodontology
2014;31:202–9. https://doi.org/10.1111/ger.12027.

Figures

Fig. 1A and 1B

Figure 1 (A)Surface roughness (Ra) of 10 PMMA acrylic discs. The bold line

represents the median, the box represents the 1st and 3rd quartiles, and the whiskers

represent 95% of observations. (B) Colony forming units (CFU) per mL of 48 *hr P.*

aeruginosa and *A. baumannii* biofilms counted on 10 PMMA discs. Values represent

the mean of 10 biofilms. Error bars are the ± SD.

Fig. 2A and 2B

Figure 2 Survival of 48-h (A) *P. aeruginosa* and (B) *A. baumannii* biofilms after

exposure to sodium hypochlorite (NaOCl), Ecolab chlorhexidine (Ecolab CHS),

Colgate cetylpryidinium chloride (Colgate CPC), Steradent (STD) and PBS (control)

with or without sonication, for 1 min. Dark grey boxes represent exposure to denture

cleaning agents only and light grey boxes represent exposure to denture cleaning

plus sonication. Each experiment was performed in triplicate with three biological

repeats.

Fig. 3

Figure 3 Survival of 48-h *P. aeruginosa* biofilms after exposure to Ecolab

chlorhexidine (Ecolab CHS), Colgate cetylpryidinium chloride (Colgate CPC), and

PBS (control) with sonication in either a commercial (COM) or laboratory (Lab)

based sonicator for 1 min and 5 min. Dark grey boxes represent 1 min and light grey

boxes represent 5 mins sonication times. The white box represents PBS without

sonication as a control group. Each experiment was performed in triplicate with

three biological repeats