Mechanistic insights into zearalenone-accelerated colorectal cancer in mice using integrative multi-omics approaches

Emily Kwun Kwan Lo[1†], Xiuwan Wang[2†], Pui-Kei Lee[3], Ho-Ching Wong[3], Jetty Chung-Yung Lee[1], Carlos Gómez-Gallego[4], Danyue Zhao[3,5*], Hani El-Nezami[1,4] and Jun Li[2,6*]

[1]School of Biological Sciences, University of Hong Kong, Pokfulam 999077, Hong Kong, China
[2]Department of Infectious Diseases and Public Health, Jockey Club College of Veterinary Medicine and Life Sciences, City University of Hong Kong, Kowloon Tong, Hong Kong 999077, China
[3]Department of Applied Biology and Chemical Technology, The Hong Kong Polytechnic University, Hong Kong
[4]Institute of Public Health and Clinical Nutrition, School of Medicine, University of Eastern Finland, FI-70211 Kuopio, Finland
[5]Research Institute for Future Food, The Hong Kong Polytechnic University, Hong Kong
[6]School of Data Science, City University of Hong Kong, Hong Kong, China

† These authors have contributed equally to this work and share first authorship
*Correspondence: jun.li@cityu.edu.hk (J.L.), daisydy.zhao@polyu.edu.hk (D.Z.)

Summary

Zearalenone (ZEA), a secondary metabolite from *Fusarium* fungi found in cereal-based foods, promotes the growth of colon, breast, and prostate cancers *in vitro*. However, the lack of animal studies hinders a deeper mechanistic understanding of the cancer promotive effect of ZEA. The aim of this study was to unveil the effect of ZEA on colon cancer progression and its underlying mechanism. Through integrative analyses of transcriptomics, metabolomics, metagenomics, and host phenotypes, we investigated the impact of a 4-week ZEA intervention on colorectal cancer in xenograft mice. Our results showed that a 4-week ZEA intervention increased the tumor weight twofold. ZEA exposure significantly increased the mRNA and protein levels of BEST4, DGKB and KI67, and the phosphorylation levels of ERK1/2 and AKT. Metabolomics analysis of the serum revealed the levels of amino acids, including histidine, arginine, citrulline and glycine decreased significantly in the ZEA group. Furthermore, ZEA lowered the alpha diversity of the gut microbiota and reduced the abundance of 9 genera, such as *Tuzzerella* and *Rikenella*, etc. Further association analysis indicated that *Tuzzerella* was negatively associated with the expression of BEST4 and DGKB genes, serum uric acid level and tumor weight. In addition, circulatory hippuric acid level was positively correlated with the tumor weight and the expression of oncogenic genes, including ROBO3, JAK3 and BEST4. Altogether, our results indicated that ZEA promoted colon cancer tumor progression through enhancing the BEST4/AKT/ERK1/2 pathway, lowering the circulatory amino acid concentration, alternating the gut microbiota composition, and suppressing the SCFA production.

Keywords

Zearalenone, colon cancer, RNA-Seq, gut microbiota, metabolomics, multi-omics

Introduction

Colorectal cancer (CRC) is one of the most common cancers and the third leading cause of cancer death [1], while the pathogenesis of CRC is still undergoing active exploration. Exogenous estrogen exposure has been classified as an environmental hazard, contributing to CRC progression through binding to estrogen receptors [2]. Zearalenone (ZEA) is a mycotoxin commonly found in our diet, mainly in cereals and grains. ZEA is a secondary metabolite produced by *Fusarium* species. Due to its structural similarity to estrogen, studies on ZEA carcinogenicity have therefore focused on hormone-dependent cancers. A recent epidemiological study discovered a link between increased urinary ZEA metabolites and the development of breast cancer [3]. Furthermore, ZEA has been implicated in promoting cancer cell growth and progression in breast [4], prostate [5] and colon cancer cells [6, 7] *in vitro*. These investigations, however, disregarded the three-dimensional growth, microenvironment of tumors, and immunoregulatory role of the gut microbiota in living animals [8]. To the best of our knowledge, there is no *in vivo* study evaluating the impact of ZEA on CRC.

In recent years, increasing evidence suggests that dysbiosis in the gut microbiome may contribute to CRC development [9-11]. The microenvironment of CRC is a complex community composed of cancer cells and microorganisms [12]. Previous studies have shown that the gut microbiota shapes immune responses and promotes tumor growth [13]. Since ZEA is mainly digested and absorbed through the gastrointestinal tract, the gut microbiota is directly exposed to ZEA and potentially regulates the metabolism and carcinogenesis of ZEA [14]. Indeed, studies have indicated that ZEA disrupts the composition and function of the gut microbiota in healthy animals [15-17].

In this study, using combined analyses of the tumor transcriptome, targeted and untargeted metabolomics and 16S rDNA sequencing, we comprehensively evaluated the impact of ZEA exposure on the development and progression of CRC in a colon cancer xenograft mouse model. We aimed to investigate the underlying mechanisms related to accelerated tumor growth and identify the key transcriptional signature, gut microbes, and metabolites that potentially contribute to ZEA-induced colon cancer development.

Materials and Methods

Animal study

Nude BALB/cAnN mice were purchased from the Laboratory Animal Unit of the University of Hong Kong. Five-week-old male mice were housed in individual ventilated cages in a sterile environment with regulated temperature (23-24ºC) and relative humidity (60-70%) on a 12-hour light cycle. The mice were fed a phytoestrogen-free diet (Open Standard diet, D11112201, Research Diets Inc., New Brunswick, NJ) and water provided *ad libitum*. After one week of acclimatization, the mice were injected in the right flank with $1x10^6$ SW480 human colon cancer cells mixed with PBS/Matrigel (1:1). When the tumor size reached 70-100 mm^3, the mice were randomly divided into two treatment groups (n=9 per group): control [oral gavage vehicle (olive oil) thrice weekly] or ZEA (0.5 mg/kg body weight (BW) in olive oil thrice weekly) (Figure 1A). The dosage in this study was chosen based on an animal study in which 1 mg/kg BW ZEA promoted DNA adduct formation in healthy mice, and the reported levels of ZEA in grain-based food in humans (3.0-33.0 µg/kg bw/day) [14, 18], in which 33.0 µg/kg bw/day was converted through body surface area conversion from humans to mice [19]. The measurement of tumor size was made as previously described with modification [20]. The sizes of tumors were measured by

an electric caliper thrice weekly. Tumor volumes were calculated based on the formula: volume

$(mm^3) = (\pi \times length (mm) \times width (mm)^2)/6$. The length represents the longest tumor diameter,

and the width represents the perpendicular tumor diameter. BW and food consumption were

recorded thrice weekly throughout the study. After 28 days, the mice were sacrificed with

pentobarbital (250 mg/kg, intraperitoneal injection), and the tumors and caecal contents were

removed, weighed and snap frozen in liquid nitrogen. Animal handling for the experimentation

was approved by the Committee on the Use of Live Animals in Teaching and Research (CULATR

No. 4785-18), the University of Hong Kong.

RNA extraction and RNA-seq

Total RNA was isolated from the tumor tissue using an RNAspin Mini kit (GE Healthcare,

UK) according to the manufacturer's instructions. After passing the quality check, the samples

were fragmented and reversely transcribed to cDNA. The final cDNA library was then built by

Novogene (Novogene, China) through the following procedure: purification, terminal repair, A-

tailing, ligation of sequencing adapters, size selection and PCR enrichment. Finally, reads were

generated using Illumina HiSeq 2500.

Quality control of RNA-Seq reads and quantification of gene expression

We mapped all raw reads to the GRCm38 (Genome Reference Consortium Mouse Build 38,

NCBI BioProject Accession: PRJNA20689) genome using BWA v0.7.17 and removed those with

more than 95% mapping coverage. The ribosomal RNA (rRNA) reads were filtered with

SortMeRNA [21]. Raw reads were further processed for quality control using the following

procedures [22] (available at

https://github.com/TingtZHENG/metagenomics/blob/master/scripts/fqc.pl): (i) Illumina

primers/adaptors/linker sequences were removed; (ii) paired-end reads with 25 bp consecutively

exact matches from both ends were removed to avoid PCR duplicates; and (iii) terminal regions

with continuous Phred-based quality less than 20 were removed. High-quality remaining reads

were aligned using TopHat2 v2.1.1 employing the GRCh38 (Genome Reference Consortium

Human Build 38, NCBI BioProject Accession: PRJNA31257) assembly as the human reference

genome. Thereafter, alignment outputs from TopHat2 were formatted and sorted with SAMtools

v1.9. SAMtools outputs were used to perform gene counts using the htseq-count function from

HTseq framework v0.11.2. In addition, RSEM v1.3.1 was executed to estimate fragments per

kilobase of transcript per million mapped reads (FPKM) based on the alignment results. Based on

the read count of genes for each sample, we first applied centered log-ratio (clr) transformation.

Analysis of differentially expressed genes and enriched pathways

Subsequently, differentially expressed genes were identified by DESeq2 v1.34.0, which

calculated the significance of gene expression profile alternations in comparison groups. The

adjusted p-value < 0.1 were used as the cutoff values for significantly differentially expressed

genes. Next, differentially expressed pathways were identified by Generally Applicable Gene-set

Enrichment (GAGE, v2.44.0) analysis using gene sets derived from KEGG pathways. An adjusted

p-value < 0.1 was used to define significantly differentially expressed pathways in the comparison

groups. In addition, Gene set enrichment analysis (GSEA, v4.0.3) was performed on gene

expression profiles using gene sets from the Molecular Signatures Database (MSigDB, v7.1): (1)

Cancer Hallmarks, (2) Oncogenic Signatures and (3) Chemical and Genetic Perturbations. |NES

score| > 1 and FDR < 0.25 were used as the cutoff values for significantly enriched gene sets.

Hematoxylin-eosin (H&E) staining, immunohistochemistry (IHC) and TUNEL assay

A part of the tumor tissue was washed, fixed immediately with formalin and embedded in

paraffin for sectioning into 5 mm slices. The sections were then deparaffinized and used to conduct

H&E staining, IHC, or TUNEL fluorescent staining. H&E staining was performed using H&E kit

from Baso (Zhuhai, China) according to the manufacturer's recommendations. The TUNEL assay

was performed using In Situ Cell Death Detection Kit (Roche Diagnostics) according to the

manufacturer's instructions with modification [23]. In brief, the slides were incubated in 0.1 M

citrate buffer in a 70°C water bath for 30 minutes and then blocked with CAS block (Thermo

Fisher, USA). The slides were washed and then incubated with TUNEL reaction mixture for 60

min at 37°C. For IHC, the sections were rehydrated in graded alcohols and distilled water. It was

then boiled in Tris-EDTA followed by quenching endogenous peroxidase activities using 3%

H_2O_2. The sections were immersed in CAS-Bloc Histochemical Reagent (Thermo Fisher, USA)

and incubated with primary antibodies against pERK1/2, pAKT, Ki67, BEST4, (1:100, Abcam)

and DGKB (Invitrogen, USA) at 4°C overnight. It was then washed thoroughly and incubated with

anti-mouse or anti-rabbit HRP-conjugated secondary antibody (Bio-Rad) or anti-goat HRP-

conjugated secondary antibody (Bio-Rad). Positive signals were visualized using 3,3'-

diaminobenzidine (Abcam, USA). The nuclei were counterstained with hematoxylin. Expression

in IHC images was quantified by NIH ImageJ v1.50 software as described in [24].

Non-targeted metabolomics analysis

One hundred microliters of serum sample were added to 1 mL ice-cold 100% methanol, spiked with 4-chloro-phenylalanine as the internal standard at a final concentration of 100 ng/mL. The mixture was vortexed for 30 s and kept at -20°C for 20 min. The cold mixture was then centrifuged at 17,000 x g at 4°C for 15 min. The supernatant was collected and dried under a gentle flow of nitrogen. The dried samples were reconstituted with 100 µL 60% methanol and sonicated for 1 min before centrifugation at 17,000 x g for 10 min. The supernatant collected was subjected to LC–MS/MS analysis using Agilent 6540 UHPLC-QTOF-MS/MS (Agilent Technologies, Santa Clara, CA, USA) with a Waters ACQUITY HSS T3 column, 2.1 mm × 100 mm, 1.7 µm (Milford, MA). The column temperature was set at 40°C and the injection volumes for positive mode and negative mode were set at 10 µL and 15 µL, respectively. The mobile phases consisted of water with 0.1% formic acid (A) and acetonitrile with 0.1% formic acid (B) at a flow rate of 0.3 mL/min. The gradient (B%) started with 1% from 0 to 1 min and increased to 99% from 1 min to 10 min, holding at 99% until 13 min and then returned to 1% at 13.5 min and maintained until 16 min. The MS parameters were as follows: capillary voltage 4000 V, nozzle voltage 1500 V, skimmer voltage 65 V, drying gas temperature 300°C, sheath gas temperature 320°C, fragmentor voltage 140 V, drying gas flow rate 8 L/min, sheath gas flow rate 11 L/min, and nebulizer pressure 40 psi. The metabolites were identified by comparing the data to the HMDB and METLIN databases using Progenesis QI software version 3.0 (Nonlinear Dynamics, Newcastle, UK).

Targeted metabolomics for amino acid profiling

Serum samples (50 µL) were mixed with 50 µL internal standard (Kairos Amino Acid Internal Standard Set, Waters USA) and water according to the manufacturer's instructions. The mixture

was then centrifuged at 9000 x g for 15 min. The mixture (10 μL) was then mixed with 70 μL of

borate buffer and 10 μL of AccQ•Tag reagent. After vortexing for 5 s, the mixture was allowed to

stand at room temperature for 1 min and heat for 10 min at 55°C. The mixture was then analyzed

using Agilent 6460 UHPLC-QqQ-MS/MS (Agilent Technologies, Santa Clara, CA, USA) with a

Waters Cortecs C18 column, 2.1 mm × 150 mm, 1.6 μm (Milford, MA). For chromatographic

separation, a binary mobile phase system was used including 0.1% formic acid in water (A) and

0.1% formic acid in acetonitrile (B) with a flow rate of 0.5 mL/min. The gradient elution program

(B%) for each run started at 1%, lasted for 1 min, and then increased to 13% at 2 min, 13% to 15%

from 2 to 5.5 min, 15% to 95% from 5.5 to 6.5 min, kept at 95% until 7.5 min, then decreased to

1% from 7.5 min to 7.6 min and maintained at 1% until 9 min. The column temperature was 55°C,

and the injection volume was 2 μL. Data processing was achieved using Agilent MassHunter

Workstation software (version B.06.00).

Metabolomics data analysis

Pathway analysis was performed to integrate the targeted amino acid results using the

Ingenuity Pathway Analysis Software (IPA) version 70750971 (QIAGEN, Redwood City, USA).

Metabolites strongly correlated with tumor weight were identified by Spearman's rank correlation

analysis. The |correlation| > 0.6 and *p-value* < 0.05 were used as the cutoff values of statistical

significance.

 Total microbial DNA extraction and 16S rDNA gene quantitative analysis

The DNA content of the caecal samples was extracted using a DNeasy Powersoil Kit (Qiagen,

Germany) according to the manufacturer's instructions. The quality of extracted DNA samples

was examined through 1% agarose gel electrophoresis. The DNA samples were then sent to

Novogene Co. Ltd (Beijing, China) for sequencing. After quality check, the hypervariable region

V3-V4 (341F&534R) of the 16S rDNA was amplified. PCR was then performed by Phusion High-

Fidelity PCR master mix (New English Biolabs, England). The PCR product was quantified with

2% agarose electrophoresis gel. The products between 400 and 450 bp were extracted by Qiagen

Gel Extraction Kit (Qiagen, Germany). Samples were quantified through Qubit and prepared for

sequence library generation by the NEBNext Ultra DNA library prep kit. The resulting library was

then sequenced on the Illumina HiSeq 2500 platform. The R package dada2 v1.22.0 was used to

perform read quality control and resolve amplicon sequence variants (ASVs). Thirty low-quality

3' bases of the forward and reverse reads were removed following inspection of QC plots. Reads

with any Ns or a maximum expected error of > 2 following truncation and mapping to phiX were

also removed. Error estimation was calculated on all samples (pooled) following dereplication.

After that, dada2 merged overlapping forward and reverse reads, and then removed chimeric

sequences. Following chimera removal, taxonomy was assigned independently to the SILVA

v138.1 reference database. Next, Shannon diversity at the genus level was calculated with the R

package vegan v2.6.2. Furthermore, the alpha diversity indexes were compared between the

experimental and control groups using independent-sample t-test. The R package vegan and

compositions v2.0.4 [25] were used to calculate beta diversity based on robust Aitchison distance.

The Wilcoxon rank sum test was used to evaluate whether there was significant difference for pair-

wise distance within groups between comparison groups. We then evaluated beta diversity in

multiple dimension scale (MDS) ordinations with vegan. Moreover, permutational multivariate

analysis of variances (PERMANOVA) of the distance matrices between groups was performed to

calculate R values (0 = no difference; 1 =maximum dissimilarity) to determine dissimilarities

among groups. Differential abundance of taxa at the genus level between comparison groups was

completed using Analysis of Compositions of Microbiomes with Bias Correction (ANCOM-BC)

implemented in R package ANCOMBC v1.4.0 [26]. An adjusted *p value* < 0.05 was used as the

cutoff to define significantly differentially abundant genera between groups.

Gene expression, microbiota, and metabolome integrative analysis

We performed an integrative analysis to reveal the associations among multi-omics and

phenotypic data. First, Spearman's correlation coefficients between the differentially expressed

genes, differentially abundant microbial taxa and differentially abundant metabolites between the

experimental and control groups were calculated. Statistical significance was set as FDR < 0.05

after Bonferroni correction, and a strong correlation was defined as |correlation|>0.8. Furthermore,

we built the significant strong correlations into a visualization network using Cytoscape software

v3.9.1.

Short chain fatty acids (SCFAs) analysis of caecal contents

The concentration of SCFAs in the caecum was measured by gas chromatography–mass

spectrometry (GC–MS) as previously described with modification [27, 28]. In brief, the caecal

content samples were homogenized using a blade homogenizer (T25, ULTRA-TUREAX, IKA) in

an extraction buffer containing 0.005 M sodium hydroxide with internal standard (10 µg/ml

deuterated acetic acid, acetic acid-d4). The homogenates were centrifuged at 13,200 x g for 20 min

at 4°C. The supernatant was mixed with 0.5 ml of 1-propanol/pyridine (3:2, v:v) and 0.1 ml of

propyl chloroformate. The mix was vortexed for 1 min and incubated at 60°C to derivatize the

SCFAs. The derivatized samples were then mixed with 0.5 ml hexane and centrifuged at 2000 x g

for 4 min. Thereafter, 400 µl of the upper layer was transferred to a glass vial for GC–MS analysis

(Agilent 6890N-5973 GC–MS, USA) set according to Zheng et al [27] report. The concentration

of SCFAs was quantified using the calibration curves constructed using acetic acid, propionic acid,

and butyric acid response ratio against acetic acid-d4.

Statistical analysis

Data are presented as the means ± SDs. Analysis was performed by using independent-sample

t-test on GraphPad Prism 9.0 (GraphPad Software, CA, USA), unless specified. A *p-value* of <0.05

was considered statistically significant. Heatmap visualization was realized using the R package

heatmap v1.0.12 and other figures were generated using the R package ggplot2 v3.3.5.

Results

Body weight and feed intake were unaffected by ZEA

We found that the administration of ZEA was well tolerated by the mice with a steady increase

in body weight (Suppl. Fig. 1A) and no evidence of reduced motor function. There was no

significant difference in the weight of the major organs including liver, spleen, lungs, heart and

kidney (Suppl. Fig. 1C) among the treatment groups. Interestingly, the body weight of the ZEA

group peaked after 24 days of treatment and dropped thereafter; it is anticipated that the increase

in tumor burden affected them. To determine whether the drop in body weight was due to a

potential reduced feed consumption, we analyzed the food intake throughout the treatment period.

The result revealed no significant difference in the feed consumption (Suppl. Fig. 1B). In

summary, these results indicated that body weight and feed intake were unaffected by the ZEA

intervention.

ZEA promoted tumor growth

We identified nearly linear increases in the tumor burden of the control (vehicle) and ZEA

groups over time (Fig. 1B). The tumor volume was significantly larger than that of the control (p

< 0.001) at the end of the intervention (Fig. 1B). The weight of the tumor in the ZEA group tripled

significantly ($p = 0.036$ with independent sample t-test) compared with the control group (Fig.

1C). The TUNEL assay showed the inhibition of tumor apoptosis by ZEA compared with the

control (Fig. 1D-F). Consistently, IHC staining showed that Ki67 positive cells in tumor tissues

were significantly increased two-fold ($p = 0.002$ with independent sample t-test) in the ZEA group

compared with the control (Fig. 1D-F). Together, we showed that ZEA reduced cancer cell death

and promoted tumor cell proliferation.

Fig. 1. ZEA accelerated tumor growth in a colon cancer xenograft mouse model. (A) Study design. Male 6-week-old Balb/c nude mice (n=9) were treated with vehicle (olive oil) or ZEA (0.5 mg/kg BW) thrice per week after subcutaneous injection of the human colon cancer cell line SW480. (B) Tumor volume variation during 28 days of monitoring, n=9. (C) Weight of tumors on day 28 and representative photograph of excised tumors imaged after ZEA treatment. (D) Representative microscopic photographs of H&E staining, Ki67 IHC staining, TUNEL staining (200x). (E) Quantification of TUNEL labelling was quantified as the total pixel area of the tumor. (F) while the expression of Ki67 was quantified as the relative Ki67-positive nuclei compared to the control. Signal normalized to the total pixels of the tumor (n=6-7). * $p<0.05$, ** $p<0.01$, compared to control.

Changes in gene expression associated with ZEA-induced tumor growth

DESeq2 identified 12 significantly up-regulated genes and 9 down-regulated genes (Table 1 and Fig. 2A) in the ZEA group compared with the control. According to the overall expression profile across different samples, these differentially expressed genes (DEGs) can be grouped into 3 clusters (Suppl. Fig. 2). Cluster 3 contains most of the significantly up-regulated genes, such as DGKB, EBF2, PTGIR, and BEST4. A over 20-fold expression increase ($\log_2$ fold change = 4.37) was observed in the DGKB gene (Table 1), which was previously reported as one of the gene signatures in the prognostic prediction of CRC [29]. In addition, another up-regulated gene worth mentioning is BEST4, which was previously described to be overexpressed in CRC patients [30].

Fig. 2. Transcriptional changes in tumor cells in response to ZEA treatment. (A) Boxplot of log-transformed expression levels (in the form of FPKM values) of all differentially expressed genes identified by DESeq2 at adjusted *p-value* <0.1; (B) The histogram represents KEGG pathway enrichment result for tumor sample DEmRNAs in ZEA group compared with the control. The DE gene PTGIR was found to be involved in neuroactive ligand-receptor interaction pathways; (C)

Normalized enrichment scores (NES) of the positively and negatively enriched gene sets (FDR < 0.25) from Molecular Signatures Database (MSigDB) hallmark gene sets significantly enriched in ZEA treated mice compared to control mice.

Table 1. The 21 differentially expressed genes (adjusted *p-value*, a positive/negative $\log_2$ fold change ($\log_2$FC) value indicates up-regulation/down-regulation) for the comparison of the ZEA treatment versus the control, ranked by the $\log_2$FC.

Gene symbol	Gene name	$\log_2$ fold change	Adjusted *p-value*
DGKB	diacylglycerol kinase beta	4.37	0.039
EBF2	EBF transcription factor 2	2.92	0.054
PTGIR	prostaglandin I2 receptor	1.71	0.066
CYP4F29P	cytochrome P450 family 4 subfamily F member 29, pseudogene	1.37	0.013
LINC02593	long intergenic non-protein coding RNA 2593	1.36	0.054
BEST4	bestrophin 4	1.32	0.054
ROBO3	roundabout guidance receptor 3	0.84	0.022
NFATC4	nuclear factor of activated T cells 4	0.67	0.044
ENTPD2	ectonucleoside triphosphate diphosphohydrolase 2	0.63	0.054
JAK3	Janus kinase 3	0.63	0.045
ABTB1	ankyrin repeat and BTB domain containing 1	0.52	0.041
LPCAT1	lysophosphatidylcholine acyltransferase 1	0.38	0.005
TOX3	TOX high mobility group box family member 3	-3.00	0.045
SMIM13	small integral membrane protein 13	-0.56	0.018
CXADR	CXADR Ig-like cell adhesion molecule	-0.46	0.066
GCLM	glutamate-cysteine ligase modifier subunit	-0.45	0.027
TMEM167A	transmembrane protein 167A	-0.43	0.013
RP2	RP2 activator of ARL3 GTPase	-0.41	0.068
TBRG1	transforming growth factor beta regulator 1	-0.40	0.094
C4orf46	chromosome 4 open reading frame 46	-0.39	0.041
EIF1AX	eukaryotic translation initiation factor 1A X-linked	-0.34	0.025

We then performed an enrichment analysis for KEGG pathways and MSigDB gene sets, including (1) cancer hallmarks, (2) oncogenic signatures and (3) chemical and genetic perturbations. The result revealed three up-regulated KEGG pathways (Fig. 2B), including the calcium signaling pathway (p_{adj} = 0.055), protein digestion and absorption (p_{adj} = 0.026) and neuroactive ligand-receptor interaction (p_{adj}=0.025). Down-regulated pathways included ribosome

(p_{adj} =3.20 x 10^{-12}), oxidative phosphorylation (p_{adj} = 3.51 x 10^{-4}) and thermogenesis (p_{adj} = 0.0128)

(Fig. 2B). Recent studies reported that the KEGG pathway "ribosome" (ko03010) was

downregulated in colon cancer [31, 32]. Furthermore, the DE gene PTGIR was involved in the DE

up-regulated pathway neuroactive ligand-receptor interaction. Regarding the MSigDB gene sets,

the enrichment analysis showed that ZEA intervention promoted gene signatures representing

oncogenic signalling, tumor invasiveness and metastasis, such as epithelial mesenchymal

transition (EMT), gain of stemness, KRAS pathway and cancer invasive signature (|NES score| >

1 and FDR < 0.25, Fig. 2C and Suppl. Fig. 3).

327   Changes in serum metabolic profiles

To explore the potential metabolic consequences of the ZEA intervention, both targeted and

non-targeted metabolomics analyses of serum samples were performed. Overall, the metabolic

profiles of the ZEA intervention and control groups were separated into two clusters (Fig. 3A-B,

Suppl. Fig. 4A-B). A total of 41 targeted metabolites were detected, among which 19 metabolites

were significantly different ($p < 0.05$ with independent sample t-test) between the ZEA and control

groups (Fig. 3A, Suppl. Fig. 4A, Suppl. Table 1). The concentration of serum amino acid (AA)

was lower in the ZEA intervention group than in the control group, which was consistent with two

previous studies in healthy rat and porcine intestinal cell lines [33, 34]. Among the AA metabolites

surveyed, phosphoethanolamine, taurine, 1-methyl histidine and kynurenine had the most obvious

changes (|log2FC| > 0.8). In addition, a total of 21 metabolites were identified with significant

differences ($p < 0.05$ with independent sample t-test) between the ZEA and control groups (Fig.

3B, Suppl. Table 2). Fourteen of them were upregulated, including organic acids and derivatives,

indolic compounds, purine derivatives, and ethanolamine, etc., whereas the remaining were

downregulated, mainly belonging to lipids and quinazoline. Among the identified metabolites,

coumarinic acid, uric acid and malic acid showed the largest fold changes ($|\log_2 FC| > 1$).

To investigate the significantly altered metabolic pathways, the log2FC of each metabolite was

calculated using Ingenuity Pathway Analysis for canonical pathway analysis. The top 20 canonical

pathways were identified and ranked according to their FDRs (Bonferroni corrected *p-values*) (Fig.

3C). Among these pathways, 4 activated pathways, including tRNA charging, superpathway of

citrulline metabolism, citrulline biosynthesis and purine nucleotides de novo biosynthesis II

pathway, were identified with positive z-scores in the ZEA treatment group compared with the

control group. The enriched citrulline metabolism and biosynthesis pathways indicated that the

ZEA intervention influenced the amino acid metabolism in mice, consistent with one previous

study using a breast cancer cell line [35].

Fig. 3. Effect of ZEA intervention on metabolites in serum samples. (A) Heatmap with log-transformed levels of 19 differentially abundant metabolites between the ZEA and control groups following targeted metabolomics analysis. (B) Heatmap with log-transformed expression levels of 21 identified differentially abundant metabolites between the ZEA and control groups following non-targeted metabolomics profiling. (C) Top 20 canonical pathways identified with the -log(FDR) and z-score listed. The threshold was set at 1.3 for -log(FDR) (equivalent to FDR < 0.05).

Changes in gut microbiota and microbiota-derived SCFAs

The alpha diversity (Shannon index) of the gut microbiota was estimated based on the taxonomic profile at the OTU level. The result indicated a significantly lower ($p = 0.007$ with independent-sample t-test) alpha diversity in the ZEA treatment group than in the control group (Fig. 4A). Moreover, multiple dimension scale (MDS) ordination based on beta diversity (robust Aitchison distance) showed a difference between the two groups to a certain extent (R = 0.1254, $p = 0.096$ with ANOSIM) (Fig. 4B). The Aitchison distances between samples in the ZEA treated group were significantly larger than those in the control group ($p = 0.006$ with Wilcoxon test), which suggests that the microbial communities became more diversified with the ZEA intervention. Nine genera were significantly differentially abundant (DA) between ZEA treatment and control groups based on ANCOM-BC (Fig. 4C). The relative abundances of *Alistipes, Candidatus Saccharimonas, Helicobacter, Lachnospiraceae UCG-010, Paludicola, Rikenella, Rikenellaceae RC9 gut group* and *Tuzzerella* were significantly lower in the ZEA treatment group than in the control group (Fig. 4C, Suppl. Fig. 5). Among the differentially abundant genera, *Rikenella* is an SCFA producer and *Candidatus Saccharimonas* is related to anti-inflammation [36]. Together, these results suggested that ZEA possibly contributed to tumor growth through repressing the SCFA-producing bacteria or anti-inflammatory bacteria.

We further assessed the impact of ZEA treatment on the major gut microbiota-derived SCFAs (acetic acid, propionic acid, isobutyric acid, butyric acid and valeric acid) by GC-MS. Overall, ZEA treatment reduced the SCFA production compared to that in the control group (Suppl. Fig. 6). The concentrations of total SCFAs, acetic acid, propionic acid, and butyric acid in the ZEA treatment group were significantly reduced ($p <0.05$ with independent-sample t-test) compared to the control (Suppl. Fig. 6).

Fig. 4. Altered caecal microbiome community upon ZEA treatment. (A) Boxplot of alpha-diversity indices. Shannon index represent the community diversity of the microbiota in the caecal content. Boxes represent the interquartile range (IQR) between the first and third quartiles (25th and 75th percentiles, respectively), and the horizontal line inside the box defines the median. ** $p < 0.01$ ($p = 0.007$, independent-sample t-test). (B) Multiple dimension scale (MDS) plots showing the beta diversity results of caecal microbial profiles between the control and ZEA groups based on robust Aitchison distance. (ANOSIM statistic R 0.1254, $p = 0.096$, based on 999 permutations). (C) Boxplot of log-transformed relative abundance of significantly (FDR <0.05) different genera between the control and ZEA detected by ANCOM-BC.

 Associations among gene expression, metabolites and gut microbes

A total of 13 metabolites were significantly correlated with tumor weight using Spearman's

correlation (Suppl. Table 3). Among them, six identified metabolites, including indolic compounds

and derivatives (serotonin and indoleacetaldehyde), purine derivatives (uric acid), and organic

acids derivatives (p-cresol sulfate, succinylacetone and hippuric acid), were positively correlated

with tumor weight (correlation > 0.6, p < 0.05). In contrast, other metabolites, including two amino

acids (1-methyl histidine and β-alanine) and lipid-related molecules (HDoHE and HETE), etc,.

were negatively correlated (correlation < -0.6, p < 0.05) with tumor weight.

To establish links among differentially expressed genes, differentially abundant metabolites

and differentially abundant gut microbes, Spearman's correlation coefficients were calculated. A

total of 12 significantly and strongly (FDR < 0.05 and |correlation|>0.8) positive and 21

significantly and strongly negative correlations were identified for integration analysis (Fig. 5 A,

Suppl. Table 4). These 33 associations consisted of 36 features (12 differentially expressed genes,

19 differentially abundant metabolites and five differentially abundant genera) (Fig. 5 A & B).

Majority (25 out 33, 75.8%) of the associations are between metabolites and genes expression.

Among the 19 differentially abundant metabolites, ten metabolites were significantly correlated

with tumor weight. More specifically, indolic compounds and derivatives (serotonin and

indoleacetaldehyde), purine derivatives (uric acid), and organic acids derivatives (p-cresol sulfate,

succinylacetone, hippuric acid and 2-hydroxycinnamic acid), were positively correlated with

tumor weight, while two amino acids (1-methyl histidine and β-alanine) and one lipid-related

molecule (HETE) were negatively correlated with tumor weight. Our results revealed strong

connections between host metabolism and the metabolic potential of the gut microbes.

Fig. 5. Integrative analysis of the differentially expressed genes, differentially abundant genera, and differentially abundant metabolites using Pearson correlation. (A) Association network

showing the Pearson correlation analysis of differentially expressed genes (red nodes), differentially abundant genera (orange nodes), and differentially abundant metabolites (dark green nodes). The significant correlations (Bonferroni corrected FDR<0.05) are shown in the network. Red lines represent positive correlations, and blue lines represent negative correlations; (B) Corrplot depicting the correlations among differentially expressed genes, differentially abundant genera and differentially abundant metabolites. Only features showing the correlations of > 0.8 or < −0.8 in any of the pairwise associations were included for visualization. The significance level was set at an FDR < 0.05; (C) Boxplot showing correlation of seven nodes whose degree ≥ 3 in Fig. 5(A). DGKB, diacylglycerol kinase beta; ENTPD2, ectonucleoside triphosphate diphosphohydrolase 2; ROBO3, roundabout guidance receptor 3; Hippuric acid; Phosphoethanolamine; *Tuzzerella*; *Rikenella*.

Seven nodes had at least three associations with other features (Fig. 5A & C). Among these nodes, the differentially expressed gene DGKB was strongly (|correlation|>0.8) positively correlated with uric acid (correlation = 0.909), p-Cresol sulfate (correlation = 0.902) and sinapic acid (correlation = 0.902), negatively correlated with *Tuzzerella* (correlation = -0.881) (Fig. 5 A-C), indicating that the interplay among host metabolites, gut microbiota and tumor gene expression may contribute to the ZEA-induced tumor growth. Among the metabolite features, hippuric acid was frequently associated with multiple differentially expressed genes with oncogenic potential (positive correlation with BEST4 (correlation = 0.895), CYP4F29P (correlation = 0.888), JAK3 (correlation = 0.888) and ROBO3 (correlation = 0.881)) (Fig. 5 A-C). Additionally, *Tuzzerella* showed the largest number of associations with other genes/metabolites. *Tuzzerella* were negatively associated with BEST4 (correlation = -0.86), DGKB (correlation = -0.881) and uric acid (correlation = -0.937) (Fig. 5 A-C). In summary, these identified gene-metabolite-genus crosslinks provide a mechanistic insight into the zearalenone-aggravated tumor growth in CRC.

Discussion

In the present study, we performed a comprehensive and integrative multi-omics study to unveil the underlying molecular mechanisms of the effect of ZEA on CRC. Through an integrative multi-omics analysis, we revealed that ZEA promoted CRC tumor growth through promoting BEST4/AKT/ERK1/2 pathway, modifying serum metabolites and modulating gut microbiota composition.

Among the 21 differentially expressed genes in the tumor section, two ZEA-promoted genes (bestrophin 4, BEST4; diacylglycerol kinase beta, DGKB) in the ZEA group were reported to be related to the malignant growth of colon cancer. BEST4 was found to be upregulated in CRC patients, and its high expression was correlated with a more advanced TNM stage, lymph node metastasis and worse survival rate by activating the BEST4/PI3K/AKT signaling pathway [30]. Our further RT–PCR analysis confirmed that BEST4 was significantly up-regulated in the ZEA group ($p = 0.024$ with independent-sample t-test) (Suppl. Fig. 7). IHC staining showed increased protein levels of BEST4 and DGBK (Suppl. Fig. 8A-B). In the following IHC staining, we confirmed that the levels of phosphorylated ERK1/2 and AKT were significantly increased (Suppl. Fig. 8A-B). We found an up-regulated BEST4 level which could contribute to the activation of AKT and ERK1/2, consistent with He et al. study [30]. DGKB was previously found to be one of the gene signatures of prognostic prediction in CRC [29]. However, the exact role of DGKB in colon cancer pathology is still unclear.

Through KEGG pathway enrichment analysis, we found a strong suppression in the "ribosome" (KEGG pathway 03010). Although pathway ribosome was previously linked to cancer progression through providing metabolic needs to the actively growing cancer cells, recent studies from five independent datasets have shown that ribosome biogenesis was one of the downregulated

KEGG pathways in colon cancer [31, 32]. Furthermore, it is known that cancer promotes the shift

from an aerobic to anaerobic metabolic pathway to survive the highly hypoxic nature of a

hyperproliferating tumor. Our findings showed that ZEA downregulated the oxidative

phosphorylation pathway while enriching protein digestion and absorption. The suppression in

oxidative phosphorylation is consistent with a lower level of TCA metabolites, such as malic acid

and citric acid (Suppl. Fig. 4), in serum.

We conducted non-targeted and targeted metabolomics analyses of serum samples to reveal

the metabolic influence of the ZEA intervention. We found that the serum amino acid (AA)

concentration was lower in the ZEA-treated group (Suppl. Table 1-2). The finding is in line with

previous findings in healthy rat and porcine intestinal cell lines, in which ZEA altered amino acid

metabolism [33, 34]. Cancer cells proliferate at a higher rate than healthy cells; hence their

metabolic demand is larger. Amino acid is a class of essential nutrients for healthy cells, and

plasma amino acid concentrations were previously found to be generally lower in lung, gastric,

colorectal, breast, or prostate cancer patients [37-39]. Patients with a more advanced stages of

CRC were found to have a lower circulatory citrulline, histidine and glutamine levels [39, 40].

Indeed, a lower serum histidine, 1-methyl histidine, 3-methyl histidine, citrulline and glutamine

levels, and enrichment in the citrulline metabolism and biosynthesis pathways in the ZEA group

were observed. A negative correlation between tumor weight and histidine derivative, 1-methyl

histidine, were found in our study. Such a reduction of histidine is suggested to be connected to

the increased histidine decarboxylate activity which was found to be correlated with tumor stages

as histidine could act as an angiogenic factor [41, 42]. In addition, we found a strong negative

correlation between β-alanine and tumor weight, which was consistent with the findings from Bi

et al. β-alanine was reported to have a negative correlation with the tumor invasiveness score, a

measurement of tumor angiogenesis [43]. On the other hand, glycine was reported as a biomarker

for colon cancer whose levels were significantly lower in CRC patients than in healthy individuals

[44]. In agreement with the finding from Leichtle et al., we found a significantly lower serum

glycine level in the ZEA group. Overall, serum amino acid profile supports the enrichment protein

digestion and absorption in KEGG pathway of our transcriptome analysis, as the circulatory amino

acid abundancy drop.

Non-targeted inflammation-related metabolites were also significantly altered in the ZEA

group. We found that serum uric acid had the second highest fold change among the detected

untargeted metabolites when compared with the control group, suggesting the role of ZEA in

altering purine metabolism. High serum uric acid was indicated to promote inflammation and

contribute to the increased tumor cell proliferation [45], while the decline in glutamine was found

to be an indicator of systemic inflammation in CRC patients [40]. Apart from amino acids,

concentrations of lipid-related metabolites were also lower in the ZEA group (Fig. 3B) and

negatively correlated with tumor weight (Suppl. Table 3). We observed that serum

hydroxyeicosatetraenoic acids (HETE) level was lower in the ZEA group, potentially leading to

defects in PPARγ activation which is responsible for apoptosis in CRC [46]. Another lipid-related

metabolite, hydroxydocosahexaenoic acid (HDoHE), was also found to be negatively correlated

with tumor weight, in line with a previous study that revealed that a higher level of HDoHE was

associated with smaller CRC tumors in mice due to the anti-inflammatory property of HDoHE

[47]. These results suggested that ZEA could possibly hinder the production of anti-inflammatory

metabolites, leading to aggravated tumor growth.

It is well known that the gut microbiota plays an important role in regulating the host's immune

system. Alteration of the microbial community is known to influence the development of diseases,

such as irritable bowel disease, obesity, liver disease and cancer [48-50]. In our study, the

abundances of SCFA-producing bacteria, including *Rikenella* [51] and *Rikenellaceae RC9 gut*

group, and the anti-inflammatory bacteria *Candidatus Saccharimonas* [36] were lowered in the

ZEA-treated mice. The suppression of SCFA producing bacteria was confirmed with the decreased

caecal SCFAs concentration in our study. Our finding is consistent with a previous study that

revealed a significantly lower caecal SCFAs content in the ZEA-treated animal group [52]. Low

abundance of SCFAs, major metabolites from gut microbiota, and SCFAs-producing bacteria are

linked to colon cancer development [53].

We further conducted an integrative analysis of multi-omics data, including differentially

expressed genes, differentially abundant metabolites and differentially abundant genera. The

results revealed that the abundance of *Tuzzerella* is negatively correlated with serum uric acid level

which was previously reported to activate PI3K/AKT pathway [54]. Subsequently, the lower

abundance of *Tuzzerella* in the ZEA group could lead to tumor cell proliferation and PI3K/AKT

activation due to the increased level of uric acid. However, the role of *Tuzzerella* in tumor

pathogenesis is still unclear; thus, further study is needed to uncover its metabolic influence and

interactions with uric acid secretion. In addition, the abundance of *Tuzzerella* is negatively

correlated with the expression of BEST4 which could bind to PI3K and activate AKT. Further

studies are warranted to confirm and explain such correlations.

Apart from uric acid, 10 other differentially abundant serum metabolites, which were strongly

correlated with tumor weight, were also found to be strongly associated with the differentially

expressed genes or differentially abundant genera in our integrative analysis. Similar to uric acid,

p-cresol sulfate was found to be positively correlated with the DGKB gene, which was reported to

be associated with CRC malignant growth [29]. In this study, we observed a significantly higher

serum level of p-cresol sulfate in the ZEA group. P-cresol sulfate was reported to induce inflammation and tumor growth in colon cancer [55]. In addition, p-cresol sulfate is derived from microbial metabolism and conjugation in the colon from amino acids, phenylalanine and tyrosine [56]. This also explains the significantly lower phenylalanine levels in the ZEA group as phenylalanine could be metabolized or converted to tyrosine following microbial conversion. Moreover, among the identified serum metabolites, hippuric acid was significantly positively correlated with BEST4, JAK3, ROBO3 and CYP4F29P genes. In particular, BEST4, JAK3 and ROBO3 were reported to promote tumor cell proliferation through activating different signaling pathways, including the PI3K/AKT, JAK3/STAT3 and SLIT/ROBO pathways, respectively [30, 57, 58]. These results suggested that hippuric acid played a role in promoting tumor growth upon ZEA treatment through alteration of oncogenic gene expression. It is also worth noting that the serum level of hippuric was elevated upon ZEA treatment. The production of hippuric acid involves the conjugation of glycine to benzoic acid [59]. Alternatively, it may also be formed from the gut microbiota metabolism of phenylalanine [60]. These results possibly explain the significant reduction of the amino acid glycine and phenylalanine but elevated hippuric acid levels in the ZEA group. Furthermore, several indolic compounds were also significantly altered and were strongly associated with several differentially expressed genes in this study. Collectively, we showed that ZEA played an important part in altering gut microbiota, microbiota-derived metabolites and oncogenic gene expression during CRC progression.

By establishing a patient-derived xenograft CRC mouse model, we provided novel and comprehensive mechanistic insights into how ZEA promoted tumor development in CRC using integrative multi-omics approaches. However, there are several limitations in the current study. One major limitation of this study is that an animal model used here may not inform the same

human responses under ZEA intervention. Although human and mouse share similar physiology

and anatomy structure, only 15% of gut commensal bacteria are shared between mice and human.

Most human microbiome studies utilized faecal samples as human ferment indigestible food

components in the colon, while cecum is the site of fermentation in mouse [61]. This may lead to

discrepancies when translating to a clinical setting. Additionally, we were unable to distinguish

bacterial species and identify the metabolic features of the microbiome due to the limitation of the

16S rDNA sequencing. To more accurately estimate effect of ZEA on human subjects with CRC,

we suggested a humanized mouse model with shotgun metagenomic sequencing in the future.

Another limitation will be the small sample size with only nine mice per group, which could limit

the power of the study.

Conclusions

In summary, this study provided evidence that ZEA could promote tumor growth and

development in colon cancer. We demonstrated that an exposure to ZEA resulted in altered tumor

transcriptome, serum metabolome and gut microbiota. ZEA-induced tumor promotion is likely

through modulating the BEST4/AKT/ERK1/2 pathway, reducing circulatory amino acid, and

altering microbiota composition. Through integration and correlation analysis of multi-omics data,

we found that the genus *Tuzzerella* was significantly negatively correlated with the BEST4

expression level and the serum metabolite, uric acid, which was positively correlated with tumor

weight. The circulatory hippuric acid level was found to be significantly positively associated with

the tumor weight and the expression levels of oncogenic genes, such as BEST4, indicating its

contribution to the observed tumor growth. Since ZEA is commonly found in the human diet and

its impact can be persistent, further studies on the potential contribution of chronic exposure of

ZEA to colon cancer development are warranted.

Compliance with Ethics Requirement

586   All Institutional Guidelines for the care and use of animals were followed. Animal handling for the
587   experimentation was approved by the Committee on the Use of Live Animals in Teaching and Research (CULATR
588   No. 4785-18), the University of Hong Kong.

CRediT authorship contribution statement

591   Conceptualization, E.K.K.L. and H.E.; methodology, E.K.K.L., X.W.W, P.K.L., H.W.; investigation, E.K.K.L.,
592   X.W.W., P.K.L., H.C.W., D.Z., J.L. and H.E.; writing - original draft, E.K.K.L., X.W.W, writing - review and editing
593   E.K.K.L., X.W.W., P.K.L, J.C.Y.L, C.G.G., D.Z, J.L. and H.E.; funding acquisition, D.Z, J.L. and H.E. All authors
594   have read and agreed to the published version of the manuscript.

Data Availability Statement

597   Transcriptome and 16S rDNA data were archived in the National Center for Biotechnology Information (NCBI)
598   Short Read Archive (SRA): PRJNA875586.

Declaration of Competing Interest

601   The authors declare that they have no known competing financial interests or personal relationships that could
602   have appeared to influence the work reported in this paper.

Acknowledgments

605   The authors especially thanks Mr. Oscar Chow for providing technical assistance. The authors are grateful to Dr.
606   Sam Leung Kin Sum for creating the preliminary data on SCFAs. This study was supported by Shenzhen Basic
607   Research Program (JCYJ20190808182402941), Guangdong Basic and Applied Research Major Program
608   (2019B030302005), Collaborative Research Fund (C7013-19GF) in Hong Kong and City University of Hong Kong
609   internal grant (7005530, 7005756, 9678247, 9680310), and PolyU Start-up fund (1-BE3H).

Supplementary material

Supplementary data to this article can be found at https://doi.org/xxxx. Supplementary figure 1: Effect of
Zearalenone in body weight, feed consumption and vital organ weight of xenograft mice; Supplementary figure 2:
Heatmap of log-transformed expression levels (in the form of FPKM values) of all differentially expressed genes
identified by DESeq2 at padj <0.1; Supplementary figure 3: Correlations between ZEA treatment and potential
oncogenic signaling; Supplementary figure 4: Effect of ZEA treatment on metabolites in serum samples;
Supplementary figure 5: Heatmap of log-transformed relative abundance of significantly (FDR <0.05) different
genera between control and ZEA detected by ANCOM-BC; Supplementary figure 6: Effect on the SCFA levels in
the caecal contents upon treatment; Supplementary figure 7: qPCR analysis of the mRNA expression of BEST4;
Supplementary figure 8: Transcriptional changes in tumor cells in response to ZEA treatment; Supplementary Table

1: Amino acids that were altered by ZEA treatment compared to control group; Supplementary Table 2: List of metabolites detected in serum samples of control and experimental groups using Progenesis QI; Supplementary Table3: Metabolites strongly related to tumor weight with spearman's rank correlation coefficient; Supplementary Table4: Pearson correlation coefficient and significance level for DA metabolites, DE genes and DA genera; Supplementary methods: RNA extraction and Real time reverse transcription polymerase chain reaction (RT-qPCR).

References

1. Bray, F., et al., *Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries.* Ca-a Cancer Journal for Clinicians, 2018. **68**(6): p. 394-424.
2. Qie, Y., et al., *Environmental Estrogens and Their Biological Effects through GPER Mediated Signal Pathways*.* Environmental Pollution, 2021. **278**.
3. Belhassen, H., et al., *Zearalenone and its metabolites in urine and breast cancer risk: a case-control study in Tunisia.* Chemosphere, 2015. **128**: p. 1-6.
4. Yip, K.Y., et al., *Combined low-dose zearalenone and aflatoxin B1 on cell growth and cell-cycle progression in breast cancer MCF-7 cells.* Toxicology Letters, 2017. **281**: p. 139-151.
5. Kowalska, K., et al., *Estrogen receptor α is crucial in zearalenone-induced invasion and migration of prostate cancer cells.* Toxins, 2018. **10**(3): p. 98.
6. Abassi, H., et al., *The mycotoxin zearalenone enhances cell proliferation, colony formation and promotes cell migration in the human colon carcinoma cell line HCT116.* Toxicol Lett, 2016. **254**: p. 1-7.
7. Lo, E.K.K., et al., *Low dose of zearalenone elevated colon cancer cell growth through G protein-coupled estrogenic receptor.* Sci Rep, 2021. **11**(1): p. 7403.
8. Jackson, E.L. and H. Lu, *Three-dimensional models for studying development and disease: moving on from organisms to organs-on-a-chip and organoids.* Integr Biol (Camb), 2016. **8**(6): p. 672-83.
9. Ma, H., et al., *Correlation between microbes and colorectal cancer: tumor apoptosis is induced by sitosterols through promoting gut microbiota to produce short-chain fatty acids.* Apoptosis, 2019. **24**(1-2): p. 168-183.
10. Vipperla, K. and S.J. O'Keefe, *Diet, microbiota, and dysbiosis: a 'recipe' for colorectal cancer.* Food Funct, 2016. **7**(4): p. 1731-40.
11. Yoon, K. and N. Kim, *The Effect of Microbiota on Colon Carcinogenesis.* J Cancer Prev, 2018. **23**(3): p. 117-125.
12. Garrett, W.S., *The gut microbiota and colon cancer.* Science, 2019. **364**(6446): p. 1133-1135.
13. Sethi, V., et al., *Gut Microbiota Promotes Tumor Growth in Mice by Modulating Immune Response.* Gastroenterology, 2018. **155**(1): p. 33-37 e6.
14. Tan, S., et al., *Zearalenone-induced aberration in the composition of the gut microbiome and function impacts the ovary reserve.* Chemosphere, 2020. **244**: p. 125493.
15. Tan, S.J., et al., *Zearalenone-induced aberration in the composition of the gut microbiome and function impacts the ovary reserve.* Chemosphere, 2020. **244**.
16. Zhang, W., et al., *Changes in intestinal barrier functions and gut microbiota in rats exposed to zearalenone.* Ecotoxicology and Environmental Safety, 2020. **204**.
17. Zhang, W., et al., *Changes in intestinal barrier functions and gut microbiota in rats exposed to zearalenone.* Ecotoxicology and Environmental Safety, 2020. **204**: p. 111072.
18. Pfohl-Leszkowicz, A., L. Chekir-Ghedira, and H. Bacha, *Genotoxicity of zearalenone, an estrogenic mycotoxin: DNA adduct formation in female mouse tissues.* Carcinogenesis, 1995. **16**(10): p. 2315-20.
19. Nair, A.B. and S. Jacob, *A simple practice guide for dose conversion between animals and human.* J Basic Clin Pharm, 2016. **7**(2): p. 27-31.
20. Li, F.L., et al., *Acetylation accumulates PFKFB3 in cytoplasm to promote glycolysis and protects cells from cisplatin-induced apoptosis.* Nat Commun, 2018. **9**(1): p. 508.
21. Kopylova, E., L. Noé, and H. Touzet, *SortMeRNA: fast and accurate filtering of ribosomal RNAs in metatranscriptomic data.* Bioinformatics, 2012. **28**(24): p. 3211-3217.
22. Zheng, T., et al., *Mining, analyzing, and integrating viral signals from metagenomic data.* Microbiome, 2019. **7**(1): p. 1-15.
23. Fayzullina, S. and L.J. Martin, *Detection and Analysis of DNA Damage in Mouse Skeletal Muscle In Situ Using the TUNEL Method.* Jove-Journal of Visualized Experiments, 2014(94).
24. Crowe, A.R. and W. Yue, *Semi-quantitative Determination of Protein Expression UsingImmunohistochemistry Staining and Analysis: An Integrated Protocol.* Bio-protocol, 2019. **9**(24).

675   25.   Van den Boogaart, K.G. and R. Tolosana-Delgado, *"Compositions": a unified R package to analyze*
676   *compositional data.* Computers & Geosciences, 2008. **34**(4): p. 320-338.

677   26.   Lin, H. and S.D. Peddada, *Analysis of compositions of microbiomes with bias correction.* Nature
678   communications, 2020. **11**(1): p. 1-11.

679   27.   Zheng, X., et al., *A targeted metabolomic protocol for short-chain fatty acids and branched-chain amino*
680   *acids.* Metabolomics, 2013. **9**(4): p. 818-827.

681   28.   Cai, J., et al., *Orthogonal Comparison of GC-MS and (1)H NMR Spectroscopy for Short Chain Fatty Acid*
682   *Quantitation.* Anal Chem, 2017. **89**(15): p. 7900-7906.

683   29.   Luo, D.K., et al., *A Novel Seventeen-Gene Metabolic Signature for Predicting Prognosis in Colon Cancer.*
684   Biomed Research International, 2020. **2020**.

685   30.   He, X.S., et al., *Oncogenic potential of BEST4 in colorectal cancer via activation of PI3K/Akt signaling.*
686   Oncogene, 2022. **41**(8): p. 1166-1177.

687   31.   Mastrogamvraki, N. and A. Zaravinos, *Signatures of co-deregulated genes and their transcriptional*
688   *regulators in colorectal cancer (vol 6, 23, 2020).* Npj Systems Biology and Applications, 2020. **6**(1).

689   32.   Shen, A.L., et al., *EBF1-Mediated Upregulation of Ribosome Assembly Factor PNO1 Contributes to Cancer*
690   *Progression by Negatively Regulating the p53 Signaling Pathway.* Cancer Research, 2019. **79**(9): p. 2257-
691   2270.

692   33.   Liu, G., et al., *Biological system responses to zearalenone mycotoxin exposure by integrated metabolomic*
693   *studies.* J Agric Food Chem, 2013. **61**(46): p. 11212-21.

694   34.   Wang, H., et al., *Integrated Metabolomics and Transcriptomics Analyses Reveal Metabolic Mechanisms in*
695   *Porcine Intestinal Epithelial Cells under Zearalenone Stress.* Journal of Agricultural and Food Chemistry,
696   2022.

697   35.   Nittoli, A.C., et al., *Effects of α-zearalenol on the metabolome of two breast cancer cell lines by 1H-NMR*
698   *approach.* Metabolomics, 2018. **14**(3): p. 33.

699   36.   He, R., et al., *Cancer-Preventive Role of Bone Marrow-Derived Mesenchymal Stem Cells on Colitis-*
700   *Associated Colorectal Cancer: Roles of Gut Microbiota Involved.* Front Cell Dev Biol, 2021. **9**: p. 642948.

701   37.   Miyagi, Y., et al., *Plasma free amino acid profiling of five types of cancer patients and its application for*
702   *early detection.* PLoS One, 2011. **6**(9): p. e24143.

703   38.   Hashim, N.A.A., et al., *Global serum metabolomics profiling of colorectal cancer.* Mol Clin Oncol, 2019.
704   **11**(1): p. 3-14.

705   39.   Geijsen, A.J.M.R., et al., *Plasma metabolites associated with colorectal cancer stage: Findings from an*
706   *international consortium.* International Journal of Cancer, 2020. **146**(12): p. 3256-3266.

707   40.   Sirnio, P., et al., *Alterations in serum amino-acid profile in the progression of colorectal cancer: associations*
708   *with systemic inflammation, tumour stage and patient survival.* Br J Cancer, 2019. **120**(2): p. 238-246.

709   41.   Di Donato, S., et al., *A Serum Metabolomics Classifier Derived from Elderly Patients with Metastatic*
710   *Colorectal Cancer Predicts Relapse in the Adjuvant Setting.* Cancers (Basel), 2021. **13**(11).

711   42.   Zhu, J., et al., *Colorectal cancer detection using targeted serum metabolic profiling.* Journal of proteome
712   research, 2014. **13**(9): p. 4120-4130.

713   43.   Zuazo-Gaztelu, I. and O. Casanovas, *Unraveling the Role of Angiogenesis in Cancer Ecosystems.* Front
714   Oncol, 2018. **8**: p. 248.

715   44.   Leichtle, A.B., et al., *Serum amino acid profiles and their alterations in colorectal cancer.* Metabolomics,
716   2012. **8**(4): p. 643-653.

717   45.   Yan, J. and C. Zhu, *Hyperuricemia is a Adverse Prognostic Factor for Colon Cancer Patients.* Int J Gen
718   Med, 2021. **14**: p. 3001-3006.

719   46.   Pakiet, A., et al., *Changes in lipids composition and metabolism in colorectal cancer: a review.* Lipids Health
720   Dis, 2019. **18**(1): p. 29.

721   47.   Nakanishi, M., et al., *A novel bioactive derivative of eicosapentaenoic acid (EPA) suppresses intestinal tumor*
722   *development in ApcΔ14/+ mice.* Carcinogenesis, 2018. **39**(3): p. 429-438.

723   48.   Ohtani, N., S. Yoshimoto, and E. Hara, *Obesity and cancer: a gut microbial connection.* Cancer Res, 2014.
724   **74**(7): p. 1885-9.

725   49.   Qin, N., et al., *Alterations of the human gut microbiome in liver cirrhosis.* Nature, 2014. **513**(7516): p. 59-
726   64.

727   50.   Imhann, F., et al., *Interplay of host genetics and gut microbiota underlying the onset and clinical presentation*
728   *of inflammatory bowel disease.* Gut, 2018. **67**(1): p. 108-119.

729   51.   Shi, H., et al., *Dietary fucoidan of Acaudina molpadioides alters gut microbiota and mitigates intestinal*
730   *mucosal injury induced by cyclophosphamide.* Food Funct, 2017. **8**(9): p. 3383-3393.

731   52.   Cai, J., et al., *Bacillus velezensis A2 Inhibited the Cecal Inflammation Induced by Zearalenone by Regulating*
732   *Intestinal Flora and Short-Chain Fatty Acids.* Front Nutr, 2022. **9**: p. 806115.
733   53.   Hou, H., et al., *Gut microbiota-derived short-chain fatty acids and colorectal cancer: Ready for clinical*
734   *translation?* Cancer Letters, 2022. **526**: p. 225-235.
735   54.   Chen, M., et al., *Soluble uric acid increases PDZK1 and ABCG2 expression in human intestinal cell lines*
736   *via the TLR4-NLRP3 inflammasome and PI3K/Akt signaling pathway.* Arthritis Res Ther, 2018. **20**(1): p. 20.
737   55.   Rivas-Domínguez, A., et al., *The Role of DNA Damage Response in Dysbiosis-Induced Colorectal Cancer.*
738   Cells, 2021. **10**(8).
739   56.   Gryp, T., et al., *p-Cresyl Sulfate.* Toxins (Basel), 2017. **9**(2).
740   57.   Zhou, W.J., et al., *Slit-Robo signaling induces malignant transformation through Hakai-mediated E-cadherin*
741   *degradation during colorectal epithelial cell carcinogenesis.* Cell Res, 2011. **21**(4): p. 609-26.
742   58.   Lin, Q., et al., *Constitutive activation of JAK3/STAT3 in colon carcinoma tumors and cell lines: inhibition of*
743   *JAK3/STAT3 signaling induces apoptosis and cell cycle arrest of colon carcinoma cells.* Am J Pathol, 2005.
744   **167**(4): p. 969-80.
745   59.   Pallister, T., et al., *Hippurate as a metabolomic marker of gut microbiome diversity: Modulation by diet and*
746   *relationship to metabolic syndrome.* Sci Rep, 2017. **7**(1): p. 13670.
747   60.   Sarkissian, C.N., C.R. Scriver, and O.A. Mamer, *Quantitation of phenylalanine and its trans-cinnamic,*
748   *benzoic and hippuric acid metabolites in biological fluids in a single GC-MS analysis.* J Mass Spectrom,
749   2007. **42**(6): p. 811-7.
750   61.   Nguyen, T.L.A., et al., *How informative is the mouse for human gut microbiota research?* Disease models
751   & mechanisms, 2015. **8**(1): p. 1-16.