1   **From southern Africa and beyond: historical biogeography of the Ledebouriinae**

2   **(Scilloideae, Asparagaceae)**

Running title: Ledebouriinae (Scilloideae, Asparagaceae) historical biogeography

Cody Coyotee Howard[1*], Leevi Nanyeni[2], Neduvoto Mollel[3], David Chuba[4], Alexandre R.

Zuntini[5], Panagiota Malakasi[5], Timothy S. Harvey[6], Nico Cellinese[7]

[1] Department of Plant Biology, Ecology and Evolution, Oklahoma State University, Stillwater,

OK, USA

[2] National Botanical Research Institute, Windhoek, Namibia

[3] Tropical Pesticides Research Institute (TPRI), National Herbarium, Arusha, Tanzania

[4] Department of Biological Sciences, University of Zambia, Lusaka, Zambia

[5] Royal Botanic Gardens, Kew, Richmond, Surrey, TW9 3AE UK

[6] Plantae Novae, Thousand Oaks, CA, USA

[7] Florida Museum of Natural History, University of Florida, Gainesville, FL, USA

* corresponding author: cody.howard@okstate.edu

Acknowledgements

We are extremely grateful to Silke Rügheimer and Dr. Esmerialda Klaassen at the National

Botanical Research Institute in Windhoek, Namibia for their assistance with fieldwork and

specimen export while in Namibia. We would also like to thank Inge Pehlemann for being a

wonderful companion during excursions in search of Namibian *Ledebouria*. We are grateful to

the governments of Namibia, Zambia, and Tanzania for issuing collection and export permits.

Namibian collections were made under permit numbers 1784/2013, 1908/2014, 2056/2016, and

2185/2016. Collections from Zambia were made under permit number TJ/DNPW/101/13/18.

Tanzanian collections were made under permit number 2017-22-NA-2016-247. Plants were

imported under USDA permit numbers P37-09-00910, P37-16-00181, and P37-16-01462. Many

thanks to Dylan Hannon, Gottfriend Milkuhn, Tom Cole, and Tom McCoy for providing leaf

material from their collections. Lastly, we thank Killian Fleurial, Taylor La Val, and Emily B.

Sessa for assistance while in the field, and William Baker for assisting with obtaining sequence

data for outgroup taxa. Funds from the Huntington Botanical Gardens, the Cactus and Succulent

Society of America, the Pacific Bulb Society, the San Gabriel Cactus and Succulent Society, the

Florida Museum of Natural History, the American Society of Plant Taxonomists, the University

of Florida International Center, the University of Florida Department of Biology, the Botanical

Society of America, the Society of Systematic Biologists, Xeric Growers, and numerous private

donors helped to support this work.

Abstract

Aim

Within sub-Saharan Africa, plants inhabiting more seasonal and arid landscapes showcase

unique distributional patterns that hint at fascinating evolutionary histories. Research on plants in

these habitats have highlighted a complex interplay between climate and orogenic activities.

However, despite a steady improvement in our understanding of evolution within these diverse

and widespread habitats, many taxa within these regions remain understudied. The

Ledebouriinae (Scilloideae, Asparagaceae) are widespread throughout sub-Saharan Africa, and

can also be found in Madagascar, the Middle East, India, and Sri Lanka. Unfortunately, this

widespread distribution coupled with taxonomic uncertainties have made uncovering the

evolutionary history of the Ledebouriinae very difficult. Here, using the most comprehensive

sampling of the lineage to date, we investigate the timing and historical biogeography of these

bulbous monocots within and outside of Africa.

Location

Sub-Saharan Africa, Madagascar, Asia

Taxon

Ledebouriinae (Scilloideae, Asparagaceae)

Methods

We infer age estimates of major clades and subclades using penalized likelihood as implemented

in treePL. Capitalizing on our broad geographic sampling and using BioGeoBEARS, we then

reconstruct ancestral ranges across the phylogeny to investigate the role vicariance and dispersal

have played in the biogeographic history of the lineage.

Results

Our results suggest that the Ledebouriinae originated within the past ~30 myr in southeastern

sub-Saharan Africa, with the major subclades arising soon thereafter. Vicariance likely led to the

current distribution of *Ledebouria* in Asia, not long-distance dispersal as previously

hypothesized. The two *Ledebouria* overlap in eastern Africa, but have divergent biogeographical

histories, divided into mostly northern and southern clades, yet each has an independent dispersal

to Madagascar. A similar north-south split is seen in *Drimiopsis*. The predominantly sub-Saharan

African *Ledebouria* clade has a complex biogeographic history, with a rapid radiation estimated

~14 mya, likely driven by drastic climate change and mountain building in southern Africa.

Main conclusions

The expansion of seasonal and arid landscapes in sub-Saharan Africa coupled with mountain

building likely spurred the radiation of the Ledebouriinae as well as several subclades.

Fragmentation, due to Miocene-driven aridification, of a once widespread distribution led to the

current distribution in Asia.

Keywords Africa, *Drimiopsis*, geophytes, *Ledebouria*, *Resnova*, Scilloideae

1 INTRODUCTION

Modern-day Africa is dominated by arid and semi-arid landscapes that contain a diversity of habitats from deserts, woodlands and savannas, to name a few (Bobe, 2006; Linder, 2014). In sub-Saharan Africa, these drier ecosystems collectively form a fairly continuous sickle-shaped corridor that connects the floras of southwestern, northeastern and western Africa, and that skirts around the wet tropics of central and western Africa (Balinsky, 1962; Bellstedt et al., 2012; Jürgens, 1997) (Figure 1). The recent onset of aridity in Africa (i.e., approximately from the Mid-Eocene onwards) is hypothesized to have been caused by the synergistic activities of rapid global cooling, tectonic events (e.g., Eastern African rift) and oceanic upwelling (i.e., Benguela current) that altered precipitation patterns across the continent (Bobe, 2006; Couvreur et al., 2021; Hagen et al., 2021; Linder, 2017; Senut et al., 2009; Sepulchre et al., 2006). These changes had immense impacts on the evolutionary trajectories of countless lineages on the African continent. For example, in the early Eocene, equatorial Africa was largely covered by tropical forests, but aridification repeatedly fragmented and reduced these habitats over time (Couvreur, 2015; Couvreur et al., 2008; Hagen et al., 2021). Evidence from plant fossils and phylogenetic studies suggest that lineages with past widespread distributions across the continent, as well as between eastern Africa and southern Asia, are now disjunct as a result of recent aridification (Ali et al., 2013; Jacobs et al., 1999; Pokorny et al., 2015; Sanmartín et al., 2010; Zhou et al., 2011). Conversely, dry and more seasonal climates have been linked to the diversification and dispersal of lineages both in and out of Africa. Today, a diversity of taxa can be found in abundance within the more arid regions of Africa as well as surrounding areas with similar climates (e.g., Arabian Peninsula) (Bruyns et al., 2014; Coe & Skinner, 1993; Jürgens, 1997; Lorenzen et al., 2012; Nylinder et al., 2016). Therefore, it is imperative we gain a broad understanding of the

historical evolution of diverse lineages across different landscapes, habitats, and biomes in order

to continually refine our understanding of the past and future changes of biodiversity within

Africa.

Geophytes, herbaceous plants with renewal buds located belowground on structures such

as bulbs, corms, and stem tubers, are ubiquitous components of seasonal or disturbance-prone

habitats, and are phylogenetically diverse (Howard et al., 2019; Pausas et al., 2018; Tribble et al.,

2021). Within Africa, geophytes are major components of the Greater Cape Floristic Region and

the Mediterranean Basin (Buerki et al., 2012; Procheş et al., 2006). Although geophytes are

predominant elements of these two areas, the geophytic habit is widespread throughout sub-

Saharan Africa, particularly within seasonal or disturbance-prone (e.g., fire) habitats (Esler et al.,

1999; Kornas, 1985). Studies have reported that many African geophytes' origins coincide with

the onset of increased seasonality and/or aridity within the continent (i.e., since the Eocene), with

the majority of diversity evolving in response to the relatively more drastic climatic changes

since the Oligocene/Miocene (Ali et al., 2012, 2013; Buerki et al., 2012; del Hoyo et al., 2009;

Procheş et al., 2006). At a broad scale, therefore, geophytic lineages make excellent candidates

for understanding recent evolutionary and biogeographic dynamics within seasonal and arid

climates.

Scilloideae (Asparagaceae) are a bulbous geophyte lineage widespread both within and

outside of Africa (Speta, 1998a). This monocotyledonous clade consist of 1,000+ taxa found

throughout seasonal climates in Africa as well as Madagascar, Europe, the Middle East, and

Asia, with a single lineage found in South America (i.e., *Oziroe* Raf.; (Giranje & Nandikar,

2016; Speta, 1998b). Although only a handful of studies have investigated the historical

biogeography of the Scilloideae and its subclades, so far, all have pointed to sub-Saharan Africa

as the origin for the majority of the group (excluding *Oziroe* in South America), followed by a

complex history within and outside of the continent (Ali et al., 2012, 2013; Buerki et al., 2012;

Pfosser, 2012). However, most inferences have been made using a broad view of each groups'

distribution (i.e., in biogeographical studies sub-Saharan Africa has been subdivided into one to

three areas) (Ali et al., 2012, 2013; Buerki et al., 2012; Pfosser, 2012). Focusing on widespread

groups within the Scilloideae using a more detailed approach to the regionalization of Africa as

well as a greater taxon sampling, may provide refined insights into the biogeographical processes

that have impacted the dispersal of plants across and out of Africa.

The Ledebouriinae are an ideal group to study because they are widespread within sub-

Saharan Africa, with a handful of taxa found in Madagascar, Socotra, Yemen, India, and Sri

Lanka (Giranje & Nandikar, 2016; Venter, 1993) (Figure 1). This distribution is unique within

Scilloideae since many sympatric lineages with Ledebouriinae are also found in northern Africa

(Pfosser, 2012; Speta, 1998a), but the Ledebouriinae are absent from the Scilloideae-rich

Mediterranean Basin (Venter, 2008) (Figure 1). In sub-Saharan Africa, the Ledebouriinae are

predominantly found within more seasonal landscapes, with highest diversity in the Limpopo,

Mpumalanga, and KwaZulu-Natal regions of South Africa (Venter, 1993), yet some occurrences

are documented from more wet, tropical regions (Figure 1). Much diversity within the

Ledebouriinae, however, remains undescribed to science (Howard, 2014; Howard et al., 2022).

Additionally, an expanded phylogenomic analysis of the group suggests that a complex

biogeographical history awaits to be thoroughly examined (Howard et al., 2022). Previous dating

analyses have estimated the origin of the Ledebouriinae sometime within the last 25 myr in sub-

Saharan Africa (Ali et al., 2012; Buerki et al., 2012). However, this clade was not the focus of

study and therefore, taxon sampling was low. Additionally, results were unable to provide fine-

scale biogeographical patterns since sub-Saharan Africa was considered and analyzed as one

large area. A more detailed categorization of the various regions in the continent would provide a

better understanding of the biogeographical history of the group. For example, Ali et al. (2012)

found that Malagasy and Indian *Ledebouria* were sister to one another, which led the authors to

invoke long-distance dispersals from Madagascar to India. However, as stated by the authors,

Ledebouriinae samples from eastern, western, and northern sub-Saharan Africa were absent,

limiting confidence in these conclusions (Ali et al., 2012). Furthermore, the phylogenetic

reconstructions of Pfosser et al. (2012) suggested two independent dispersal events to

Madagascar, but low phylogenetic resolution diminished confidence in this hypothesis. A recent

phylogenomic analysis of the Ledebouriinae provided the framework to suggest (without testing)

multiple dispersals to Madagascar, and a potential migration out of Africa via the Arabian

Peninsula into India (Howard et al., 2022).

The widespread distribution of the Ledebouriinae provides us with the opportunity to

refine our understanding of evolution and biogeography within Africa, potentially during a time

of extensive climatic, geologic, and habitat change on the continent. Here, we investigate the

timing and historical dispersal of the Ledebouriinae both within and out of Africa. We ask: 1)

Which region within Africa acted as the putative origin of the group, and 2) What roles have

dispersal and vicariance played in the evolutionary history of the Ledebouriinae, if any? Our

specific focus on these questions and the use of an increased sampling of the Ledebouriinae

allow for a more thorough investigation into the evolutionary history of the group and improve

our knowledge on patterns and processes within Africa.

2 METHODS

2.1 Phylogenetic analysis

Ledebouriinae samples were obtained from the field, private collections, and herbarium vouchers (Howard et al., 2022). DNA extractions were performed using a modified CTAB protocol, followed by high throughput sequencing on an Illumina HiSeq using the Angiosperms353 universal probe set (Johnson et al., 2019). Raw reads were cleaned using SECAPR (Andermann et al., 2018), sequences were pulled using hybpiper (Johnson et al., 2016) and aligned using MAFFT v.7 (Katoh & Standley, 2013). See Howard et al. (2022) for more details on Ledebouriinae sequence acquisition and analysis as well as data used.

We incorporated outgroup taxa from the Plant and Fungal Tree of Life project (Baker et al., 2022) and the 1KP dataset (Matasci et al., 2014). Exons were only available for the outgroup taxa included to estimate divergence times using fossils and secondary calibration points. Phylogenetic reconstruction including the Ledebouriinae plus outgroups was performed on a concatenated, partitioned supermatrix of exons with 10% gappy/ambiguous sites removed using phyx (Brown et al., 2017). This matrix was analyzed using IQ-Tree v.2-rc1 (Nguyen et al., 2015) with 1000 ultrafast bootstraps (Nguyen et al., 2015) with 1000 ultrafast bootstraps, a best-fit partitioning scheme using the greedy algorithm of PartitionFinder (Lanfear et al., 2012), and a relaxed clustering percentage of 10 (Lanfear et al., 2014), followed by phylogenetic reconstruction (-m TESTMERGE). However, the exon-only dataset returned low support for many nodes within the Ledebouriinae. Therefore, we reran the IQ-Tree analysis with a topological constraint tree (-g) that was previously built from a supercontig (i.e., exons + introns) dataset of the Ledebouriinae (Howard et al., 2022).

2.2 Time calibration

We incorporated eight outgroup fossil calibration points, each with a minimum age

specified in Iles et al. (2015) (Table S1). A secondary calibration point at the crown node of

monocots was inferred between 131–135 mya based on previous analyses (Givnish et al., 2018;

Magallón et al., 2015).

Given the size of the dataset, we used penalized likelihood as implemented in treePL

(Smith & O'Meara, 2012) for time calibration. To incorporate uncertainty around age estimation,

we took a multi-tiered approach (see https://github.com/sunray1/treepl) similar to previous

studies estimating divergence times using large phylogenetic datasets (Emberts et al., 2020; Li et

al., 2019; Magallón et al., 2015). We generated 100 bootstrap replicates of our original exon

supermatrix alignment using RAxML v.8.2.0 (f -j option) (Stamatakis, 2014). A maximum

likelihood tree for each replicate with a corresponding partition file was then reconstructed using

a topological constraint (i.e., the phylogeny from the IQ-Tree analysis) to ensure consistent

calibration point placement, and a GTRGAMMA model of evolution. The resulting 100 "best

trees" were rooted on *Acorus gramineus* using phyx (Brown et al., 2017). In Step 1, each

replicate tree had a priming step completed with a random seed number and the thorough

command invoked. Step 2 was performed three independent times to assess convergence on the

best smoothing parameter for each individual tree. Step 2 also included the individual outputs

from each tree's previous priming step (e.g., optad, moredetail, etc.) as well as the cross-

validation (CV) steps, which were set to cvstart=10000, cvstop= .00000000000001,

cvmultstep=0.09. Lastly, Step 3 summarized each individual tree's CV output to determine and

scale each tree using the appropriate smoothing parameter. The 100 ultrametric phylogenies were

summarized using TreeAnnotator v.1.10.4 (Bouckaert et al., 2014) to obtain a maximum clade

credibility tree with 95% confidence intervals around each node and median node heights.

2.3 Biogeographical analysis

The biogeographical regionalization of Linder et al. (2012) was used for categorizing the

location of each sub-Saharan African Ledebouriinae accession. The exact distribution of

populations/taxa represented by many field-collected individuals (i.e., collections made by C.C.

Howard) remains to be fully assessed since they are undescribed species and/or are only known

from one locality. Additionally, many described species are currently known as occurring in

small geographic ranges or even single mountain tops (Lebatha, 2004; Venter, 2008).

Consequently, most samples were assigned only to known areas of occurrence. Accessions found

outside of Africa were coded according to their respective areas of occurrence: Madagascar,

Yemen, Socotra, or Sri Lanka). Another reason we coded each taxon using this approach is

exemplified by *Ledebouria revoluta*, reported from the Cape Region of South Africa to Sri

Lanka (Mwafongo et al., 2017; Venter, 1993). This widespread taxon likely represents a species

complex with a wide range of morphological variation (Brita Stedje, 1998), and therefore, often

this name is associated with individuals whose species identification is difficult (Mwafongo et

al., 2017). Consequently, and not surprisingly, *L. revoluta* is an ambiguous entity as evidenced

by its polyphyly within the Ledebouriinae (Howard et al., 2022). In total, 14 geographic regions

were used to categorize all terminals. To estimate ancestral ranges, we compared the maximum

likelihood implementations of three biogeographical models, DEC, DIVA-like, and BAYAREA-

like as implemented in BioGeoBEARS (Matzke, 2013a, 2013b, 2014) in R (R Core Team,

2016). We used log-likelihood and AICc values to determine the best model between the three.

Given the debate surrounding the use of models with the +j parameter (Matzke, 2022; Ree &

Sanmartín, 2018), we decided to not include them and, additionally, their implementation tends

to infer a high number of jump dispersals, which are unlikely biological scenarios for the

majority of the Ledebouriinae (see Discussion).

2.3.1 Biogeographical uncertainty

Within a subclade of *Ledebouria* Clade A, phylogenetic relationships are poorly

supported (Figure 2, node a) (Howard et al., 2022). However, given the potential rapid radiation

along the backbone of this group, we wanted to explore the biogeography of this clade since it

may contain an interesting history. To compensate for the uncertainty, we incorporated

topological and branch depth variability into our biogeographical analysis by using a random

sampling of trees from the posterior of a Bayesian analysis for the subclade of interest

(Ceccarelli et al., 2019; Magalhaes et al., 2021). To begin, we reduced our total alignment to the

most clock-like genes using SortaDate (Smith et al., 2018). This was done due to failure to reach

convergence when using the entire alignment (data not shown). We kept genes that were at least

10% concordant (bipartition >0.1), had a tree length greater than 7.24, and had root-to-tip

variation of less than 0.009. Cutoff values for the latter two were determined using the median

values for all gene trees, as previously reported (Pillon et al., 2021). Of the remaining

alignments, we included only taxa within the subclade of interest (Figure 2b) and removed

duplicate Ledebouriinae outgroup taxa as well as those putatively duplicate within the

Ledebouriinae ingroup (indicated by * in Figure 2). Each gene was then aligned using MAFFT

(Katoh & Standley, 2013) invoking the --auto option. Sites with 5% gaps/ambiguity were

removed using phyx (Brown et al., 2017). Due to failure to reach convergence using a

partitioned, concatenated alignment in BEAST v1.10.4 (Drummond & Rambaut, 2007), we

instead input an unpartitioned, concatenated alignment using a GTR+GAMMA model of

substitution, a speciation birth-death prior, and a lognormal distribution with an offset 17.1,

mean 1.0, stdev 1.0 (95% range 17.24–19.88) on the root node of the subclade. This initial value

was chosen based on the treePL divergence time estimates for this node. We ran an MCMC

chain of 300k generations sampling every 10k generations. Convergence was assessed using

Tracer v1.7.1 (Rambaut et al., 2018) to ensure all ESS values were above 200. We removed 50%

of trees as burnin and used the remaining trees in the analysis. We randomly sampled 100 trees

after burnin, and performed a DIVA-like analysis (i.e., the best model) on each tree followed by

stochastic mapping using BioGeoBEARS (Matzke, 2013a). Trees were summarized to obtain an

average number of biogeographical events within and between each region included in the

analysis. Scripts for running the analysis can be found on github

(https://github.com/ivanlfm/BGB_BSM_multiple_trees). The rationale for incorporating

phylogenetic uncertainty was developed by Ceccarelli et al. (2019) and Magalhaes et al. (2021).

3 RESULTS

3.1 Phylogenetic relationships and age estimates

We recovered a polyphyletic *Ledebouria*, and a monophyletic *Drimiopsis* and *Resnova*,

which is congruent with a recent study (Howard et al., 2022). However, shallow-level

relationships differed compared to Howard et al. (2022), particularly within the *Ledebouria*

clades, and especially within *Ledebouria* Clade A. This is due to using a topological constraint (-

g), which searches tree parameter space, over using a fixed topology (-te) which does not

perform a tree search (see http://www.iqtree.org/doc/Command-Reference). Given differences in

the inputs between Howard et al. (2022) and our exons-only datasets, we preferred a topological

constraint allowing for a tree search over forcing a fixed topology.

Within a penalized likelihood framework, the Scilloideae originated approximately 52.0

my (95% HPD 51.2 – 54.5) (Figure 2; Table 1). We recovered a median crown age estimate for

the Ledebouriinae of 28.0 myr (95% HPD 26.1–28.3), with age estimates of the four major

Ledebouriinae clades soon thereafter. The split between *Drimiopsis* and *Ledebouria* Clade B was

estimated at 27.1 mya (95% HPD 25.2–27.5). The crown of *Drimiopsis* was dated to 22.1 mya

(95% HPD 20.5–23.5), and *Ledebouria* Clade B was dated at 26.1 mya (95% HPD 24.0–26.5).

The split between *Resnova* and *Ledebouria* Clade A occurred at 26.3 mya (95% HPD 23.7–

25.2). The *Resnova* crown was dated at 17.9 mya (95% HPD 16.6–18.5), and *Ledebouria* Clade

A was estimated at 25.0 mya (95% HPD 23.0–25.4) (Table 1).

3.2 Biogeography

The DIVA-like model produced the most likely ancestral range estimates among the

models (LnL -220.54) (Table 2). The regions Natal+Zambezian were reconstructed as the

ancestral range for the Ledebouriinae ($p = .53$) and *Drimopsis* ($p = .84$), while Zambezian was

reconstructed for *Drimiopsis* + *Ledebouria* Clade B ($p = .95$) and *Ledebouria* Clade B ($p = .98$).

Within *Ledebouria* Clade B, the model favors a widespread distribution of Yemen+Zambezian

($p = .55$) followed by subsequent reconstructions containing various combinations of Zambezian

and other regions, or Zambezian alone (Figures 3 and S2). Within *Ledebouria* Clade B we also

recover one of the two dispersals of the Ledebouriinae to Madagascar (Figure 3).

The Kalahari+Natal regions were recovered as the ancestral range for *Resnova* +

Ledebouria Clade A ($p = .42$), whereas the Natal region was recovered for *Resnova* ($p = .99$),

and Kalahari for *Ledebouria* Clade A ($p = .51$) (Figures 3 and S2). Within *Ledebouria* Clade A,

we find an additional dispersal to Madagascar (Figure 3). Additionally, in summarizing the 10

biogeographic stochastic reconstructions (see section 2.3.1), we recovered a steady rise in the

number of lineages in the Zambezian and Kalahari regions from ~15 myr onwards in a subclade

of *Ledebouria* Clade A (Figure 3b). Additionally, despite some differences in topology between

these results and those of Howard et al. (2022), both studies overall share clades that largely

reflect geography. We recover two subclades of mostly Namibian taxa, a subclade containing

mostly South African taxa, and a subclade containing a mixture of Zambian, Tanzanian, and

Zimbabwean taxa (Figures 2 and 3).

4 DISCUSSION

4.1 Broad biogeographical patterns

Here, we present the best and most comprehensive sampling of the Ledebouriinae to date

(representing ~30% of described Ledebouriinae taxa (POWO, 2019) plus numerous undescribed

taxa) which has provided greater insights into the historical biogeography of this widespread,

bulbous lineage than previously uncovered. Our results suggest a rapid radiation along the

backbone of the Ledebouriinae, estimated between ~28–26 mya (Figure 2; Table 1) in

southeastern, sub-Saharan Africa (Figures 3, S1, S2). This region corresponds somewhat to the

current center of diversity of the Ledebouriinae, which is at the intersection of the Natal-

Kalahari-Zambezian biogeographical regions (Lebatha, 2004; Manning, 2020; Venter, 1993).

The four major subclades (*Ledebouria* Clade A, *Ledebouria* Clade B, *Resnova*, and *Drimiopsis*)

originated in neighboring regions to one another soon thereafter (Table 1; Figure 3). During this

timeframe in Africa, major shifts in climate and habitat composition have been inferred at both

local and continental scales, and at shallow and deep phylogenetic scales (Couvreur et al., 2021;

Gizaw et al., 2021; Hagen et al., 2021; Kandziora et al., 2022; Pokorny et al., 2015). From the

Eocene-Oligocene transition, global cooling promoted the expansion of seasonal and arid

climates as well as savanna, grassland, and fire-prone habitats in Africa, with greater increases

during the Miocene (Senut et al., 2009; Sepulchre et al., 2006). Furthermore, the Great

Escarpment underwent renewed uplift in the early Miocene, which increased habitat

heterogeneity that was coupled with an intensification of global cooling and aridification that

collectively led to lineage radiation and dispersal in many African taxa (Cowling et al., 2009;

Galley et al., 2007; Neumann & Bamford, 2015; Partridge & Maud, 1987). Being bulbous

geophytes, the Ledebouriinae may have been "pre-adapted" for these climatic and environmental

changes, and therefore diversified in response (Howard et al., 2020). Overall, our biogeographic

reconstructions and divergence time estimations lead us to hypothesize that the expansion of

seasonal habitats and extensive orogenic activity in southern, sub-Saharan Africa spurred the

radiation of the Ledebouriinae and promoted additional radiations and dispersals over time.

4.2 A tale of two *Ledebouria*

We estimate that the two *Ledebouria* lineages originated at slightly different times

(Figure 2; Table 1) in neighboring regions (i.e., Kalahari vs Zambezian), which lead to

subsequent divergent biogeographic histories (Figure 3). In general, the two *Ledebouria* overlap

in eastern Africa with a geographical divide between southern and northern sub-Saharan Africa

(Figure 2) (Howard et al., 2022). The divergent evolutionary histories and current overlapping

distribution of the two *Ledebouria* may have been influenced by the expansion and contraction

of wet, tropical, forested habitats in eastern Africa that repeatedly split a historically more

widespread distribution (Couvreur, 2015; Couvreur et al., 2021). These past habitat fluctuations

have created a mosaic of climates and vegetation types in eastern Africa, and have influenced the

evolutionary trajectory of countless lineages (Dagallier et al., 2020; Lorenzen et al., 2012). Other

lineages adapted to seasonal and arid habitats share similar distributions with the two *Ledebouria*

(Grace et al., 2015; Jürgens, 1997). For example, the two lineages overlap geographically with

that of stapeliads (Ceropegieae, Apocynaceae)—*Ledebouria* Clade B overlaps with the Northern

grade, whereas *Ledebouria* Clade A overlaps with the Pan-African clade (see Fig. 3 in Bruyns et

al. (2014)). We also find two dispersals of *Ledebouria* to Madagascar (Figure 3), which had been

previously hypothesized (Pfosser, 2012). Madagascar and mainland Africa have been separated

since at least the Paleocene (Couvreur et al., 2021), which suggests long distance dispersal from

Africa to Madagascar led to the origin of Ledebouriinae on the island.

4.2.1 The Voyage of *Ledebouria* Out of Africa, or *Ledebouria* Clade B

Previous studies invoked long distance dispersal from Madagascar to India to explain the

distribution of *Ledebouria* in Asia (Ali et al., 2012); however, we find little support for this

hypothesis. Reconstructions place the ancestral area of *Ledebouria* Clade B in the Zambezian

region followed by migration into other regions, such as Yemen and India, between ~20–15 mya

(Figures 2 and 3). During this time, the Arabian Peninsula and Socotra were connected or in

close proximity with mainland Africa (Edgell, 2006; Fleitmann et al., 2004; Jacobs, 2004; Rögl,

1999), which would have allowed overland dispersal to these regions. Pollen and wood fossils

indicate historical widespread distributions between eastern Africa and south Asia of other

closely related taxa (Bonnefille, 2010; Morgan et al., 1994). Additionally, fossils indicate that

forested habitats were present in the Arabian Peninsula during the Eocene and Oligocene, that

later gave way to more open, xeric, grassland habitats in the Miocene (Jacobs et al., 1999;

Whybrow & McClure, 1980). The modern-day equivalents of these conditions are commonly

inhabited by *Ledebouria* (Venter, 1993). Given the historical environments present in Africa and

south Asia during the Late Oligocene/Early Miocene, the ancestor of *Ledebouria yemenensis* +

L. nossibeensis likely had a widespread distribution that was fragmented as aridification

increased during the Miocene. Additionally, vicariance likely promoted allopatric speciation of

Socotran *Ledebouria* as the islands drifted away from mainland Africa over time. Vicariance has

been invoked for other lineages with hypothesized widespread historical ranges between Africa,

the Arabian Peninsula, Socotra, and/or southern Asia, including *Isodon* (Lamiaceae) (Yu et al.,

2014), *Aganope* (Fabaceae) (Sirichamorn et al., 2014), *Searsia* (Anacardiaceae) (Yang et al.,

2016), *Smilax* (Smilacaceae) (Chen et al., 2014), and several reptiles (Main et al., 2022; Smíd et

al., 2013; Tamar et al., 2016), many of which have similar age estimates and distributions as

Ledebouria Clade B. In other groups, such as the Urgineoideae (Asparagaceae) (Ali et al., 2013)

and *Uvaria* (Annonaceae) (Zhou et al., 2011), dispersals from Africa to south Asia by way of the

Arabian Peninsula have been hypothesized. Furthermore, *Ledebouria* seed has limited dispersal

capacity, which occurs primarily via sheet water flow (i.e., travels only short distances and is

rainfall dependent) (Venter, 1993). In years of limited rainfall and/or in flat terrains, seeds and

seedlings can be found surrounding the parent plant (CCH, pers. obs.), and the ephemeral nature

of the seeds in some taxa warrants germination soon after ripening (CCH and TSH, pers. obs.).

Thus, our age estimates and biogeographical reconstructions as well as the limited dispersal

ability of *Ledebouria* lead us to hypothesize that Clade B dispersed out of Africa to south Asia

via overland migration through the Arabian Peninsula. This historical, widespread distribution

was then fragmented by Miocene-driven aridification. Today, *L. yemenensis*, the only

Ledebouriinae currently recorded from the Arabian Peninsula, likely represents a relic of this

once widespread distribution as it is endemic to the cooler, higher elevation of the Yemeni

Highlands surrounded by arid lowlands.

4.2.2 Within Africa—*Ledebouria* Clade A

Our results suggest that *Ledebouria* Clade A originated slightly later and in a more

southerly region than that of *Ledebouria* Clade B (Table 1; Figures 2 and 3). Based on our

dataset, *Ledebouria* Clade A arose ~25–23 mya and remained predominantly within sub-

equatorial Africa (Figures 2 and 3). Despite the low resolution within some areas of Clade A

(Howard et al., 2022), we recovered multiple subclades that reflect geographic affinity (Figure

2). For example, we see a subclade of mixed geographic composition (i.e., *L. caesiomontana* + *L.*

sp. 12 CCH190) (node c, Figure 2) that suggests a widespread distribution between the Kalahari

and Zambezian regions that diverged and lead to *in situ* speciation in the Zambezian region

followed by a dispersal to Madagascar (Figure 3). We recovered an additional subclade of

predominantly Zambezian taxa (i.e., *L.* sp. 37 CCH169 + *L.* sp. 24 Uganda) (subclade in node a,

Figure 2) with evidence of a dispersal to the Sudanian region (i.e., *L.* sp. 24 Uganda).

Furthermore, two subclades of predominantly Namibian taxa (i.e., *L.* sp. 14 CCH109 + *L.* sp. 18

CCH218 (node b, Figure 2), and *L.* sp. 7 CCH066 + *L.* sp. 32 CCH145 (subclade in node a,

Figure 2)) suggest multiple dispersals to the region followed by *in situ* speciation (Figure 3).

Lastly, a subclade of mostly South African taxa (i.e., *L. galpinii* + *L. coriacea*) (subclade in node

a, Figure 2) was recovered (Figure 3). Overall, our results suggest that within a relatively short

time frame (~20–15 mya), a complex history of dispersals followed by *in situ* speciation events

occurred within *Ledebouria* Clade A in southern Africa (Figure 3; Table S2).

From the Miocene onwards, complex interactions between mountain uplift and climate

change have promoted diversification across lineages within southern Africa (García-Aloy et al.,

2017; Maswanganye et al., 2017; Neumann & Bamford, 2015; Nielsen et al., 2018). *Ledebouria*

Clade A may have originated in southeastern Africa then dispersed and diversified as the region

grew more heterogeneous and arid. For example, we found an instance of potentially rapid

radiation within *Ledebouria* Clade A between 20–17 mya (Figure 3b), which corresponds to a

clade with consistently poor phylogenetic resolution (Howard et al., 2022). This subclade

contains taxa from across southern sub-Saharan Africa with geographic signals within (i.e.,

clades of South African, Namibian, and Zambian taxa) (Figures 2 and 3). It is possible that the

Middle Miocene Climatic Optimum or Middle Miocene Climate Transition (Zachos et al., 2001)

coupled with renewed uplift of the Great Escarpment spurred this radiation, and we see a steady

rise in lineage accumulation from ~20–15 mya (Figure 3b). During the Middle Miocene, the

vegetation of South Africa was likely tropical, which gave way to more xerophytic shrubland by

Late Miocene due to a global cooling climate (Pound et al., 2012). This time period is associated

with diversification in the Crassulaceae (Lu et al., 2022) and also the expansion of C4

grasslands/savannas at a global scale (Bobe, 2006; Senut et al., 2009). Current biogeographical

patterns of *Aloe* suggest dispersal from southern Africa to East Africa as seasonality and

aridification increased across the continent within the past ~16 my (Grace et al., 2015).

Monsonia (Geraniaceae) share similar distributional patterns as *Ledebouria* Clade A with

multiple lineages comprised of mainly southwestern or southeastern taxa (i.e., southern African

split between eastern and western clades), which radiated within the past ~20–15 myr (García-

Aloy et al., 2017). Overall, we find a fascinating, albeit more complicated, evolutionary history

of *Ledebouria* Clade A within southern sub-Saharan Africa that remains to be fully resolved.

4.3 Limited, but intriguing insights into *Drimiopsis*

Based on our limited sampling of *Drimiopsis*, we find additional evidence for a southern

and northern sub-Saharan African divide (Figures 2 and 3), which is also seen in ungulates

(Lorenzen et al., 2012), *Lycaon pictus* (Canidae) (Marsden et al., 2012), *Encephalartos*

(Cycadaceae) (Mankga et al., 2020), and *Agama* (Reptilia) (Leaché et al., 2014). The

eastern/northern sub-Saharan African *Drimiopsis* arose ~19–17 mya (Figure 2), during a time of

increased mountain building in the region (~17 and ~13.5 mya; (Wichura et al., 2015) that was

followed by a shift towards a more arid climate (Bonnefille, 2010; Linder, 2017; Morgan et al.,

1994; Senut et al., 2009). Several East African *Drimiopsis* have overall more succulent leaves

compared to their southern African relatives (Lebatha, 2004), which may have been selected for

as eastern Africa experienced greater aridification compared to the southeast. Additionally, East

African *Drimiopsis* are known polyploids (Lebatha, 2004; B Stedje & Nordal, 1987), which may

have further spurred evolution in this region since polyploidy may provide an advantage in

particular environmental conditions (e.g., aridity) (Manzaneda et al., 2012; Sonnleitner et al.,

2010). In contrast, South African *Drimiopsis* occur in moist and shady habitats, and are not

found in the drier portions of southwestern Africa (Lebatha, 2004). Given the estimated age of

South African *Drimiopsis* (~13.6 mya) (Figure 2), uplift of the Great Escarpment may have

increased aridity in southwest Africa (Bobe, 2006; Partridge & Maud, 1987) that hindered

Drimiopsis establishment in this region, but at the same time promoted diversification in

southeastern Africa leading to its high diversity in the area (Lebatha, 2004). However, far greater

sampling of *Drimiopsis* from its entire distribution is needed to test hypotheses regarding their

evolutionary history. Ideally, an expanded phylogenetic framework would be coupled with

ploidal inferences, anatomical studies, and experimental manipulations to fully assess the

relationship between historical biogeography and morphological evolution.

4.4 Where to sample next?

Our taxon sampling heavily consisted of southern, sub-Saharan African taxa (Figure 1),

and largely excluded *Resnova*. Therefore, increased sampling of *Resnova* as well as *Ledebouria*

and *Drimiopsis* from western, central, and northern Africa will provide further insights into the

biogeographical history of the Ledebouriinae. For example, *Ledebouria* occurs in northern

Angola (Gregory Jongsma, pers. obs.) and Gabon (Figure 1). This leads us to wonder whether a

continuous distribution was once present along the central/western coast of Africa, or if the

Central African tropical forests have consistently obstructed Ledebouriinae dispersal within the

coastal, wet tropics. Furthermore, a *Ledebouria* accession from Uganda is nested within

Ledebouria Clade A, which suggests that by increasing sampling from northern/western Africa

we may uncover additional biogeographic events. Lastly, multiple, morphologically distinct

Ledebouria taxa are recorded from Socotra and surrounding islands (Miller & Alexander, 2010),

and India (Giranje & Nandikar, 2016). Incorporating these into future phylogenetic studies may

highlight further dispersals, or bolster support for allopatric speciation/vicariance as a major

evolutionary process leading to the current distribution of the group outside of Africa.

5 CONCLUSION

The Ledebouriinae are a widespread, bulbous lineage of monocots that remain poorly

understood taxonomically due to a high degree of phenotypic plasticity and significant under

collection of specimens. Our study provides a pivotal first step towards refining our

understanding of historical evolution and biodiversity within this fascinating lineage. The results

of our analyses highlight the complex biogeographic history of the Ledebouriinae within and

outside of sub-Saharan Africa. We find that the Ledebouriinae evolved within the past ~30 myr,

which suggests the group radiated in response to increasing climatic seasonality and orogenic

activity since. We find evidence that vicariance led to the current distribution of *Ledebouria* in

Asia and Socotra, with overland dispersal out of Africa into Asia likely occurring via the Arabian

Peninsula. We also find evidence of two independent dispersals to Madagascar. Within Africa,

we find evidence of a northern and southern sub-Saharan division between the two *Ledebouria*

lineages and within *Drimiopsis*. We also uncover a complex history of dispersals and *in situ*

speciation events within southern Africa in *Ledebouria* Clade A, a clade that warrants improved

phylogenetic resolution to further elucidate the processes affecting its current distribution. In

conclusion, our study shows the value of increasing research focus on understudied lineages

inhabiting seasonal and arid habitats.

Tables

Table 1. Comparison of median age estimates and associated uncertainty (95% highest
probability density (HPD)) for each major Ledebouriinae clade using penalized likelihood as
implemented in treePL.

Clade	median estimate	95% HPD
Scilloideae	52.2	51.2 – 54.5
Ledebouriinae	28.0	26.1 – 28.3
Ledebouria Clade A	25.0	23.0 – 25.4
Resnova	17.9	16.6 – 18.5
Clade A + *Resnova*	26.3	24.5 – 26.7
Drimiopsis	22.1	20.5 – 23.5
Ledebouria Clade B	26.1	24.0 – 26.5
Clade B + *Drimiopsis*	27.1	25.2 – 27.5

493   **Table 2**. Statistical outputs for each model from BioGeoBEARS. log-likelihood (LnL); rate of
494   dispersal (*d*); rate of extinction (*e*); Akaike's Information Criterion (AIC)

Model	LnL	Number of parameters	*d*	*e*	*j*	AIC
DEC	-267.1	2	0.01	0.01	0	538.4
DIVA-like	**-220.5**	**2**	**0.0027**	**0.009**	**0**	**445.2**
BAYAREA-like	-232.8	2	0.0036	0.04	0	469.8

 Figures

Figure 1. General distribution of the seasonal and arid ecosystems of sub-Saharan Africa (light
orange polygon) underlain by the distribution of the Ledebouriinae displayed using GBIF
specimen occurrence data (teal squres; GBIF.org (22 March 2022) GBIF Occurrence Download
https://doi.org/10.15468/dl.xmccae). Collection localities of samples used in this study are
denoted with maroon circles. Examples of habitats and associated Ledebouriinae taxa are
displayed in the images. a) *Ledebouria* sp. 8 CCH186, b) *Ledebouria* sp. 15 CCH027, c)
Drimiopsis botryoides subsp. *botryoides* CCH153. See Figure 2 for phylogenetic placement of
these example taxa. General distribution of seasonal landscapes adapted from Balinsky (1962),
Jürgens (1997), and Bellstedt et al. (2012).

Figure 2. Divergence times within the Ledebouriinae as estimated using penalized likelihood in treePL. For each clade, collection localities are overlaid on a map of Africa, Madagascar, and Asia to aid with discussions of geographical patterns. Clades labeled with a, b, and c in yellow are clades discussed within the manuscript. Numbers above blue bars denote median age estimates. Blue bars denote 95% highest posterior density (95% HPD). * indicate samples excluded from subsequent biogeographical analyses. Not all nodes are annotated with estimated divergences times for illustrative simplicity. For a fully annotated phylogeny, see Figure S1.

Figure 3. Biogeographical reconstruction using the DIVA-like model as implemented in
BioGeoBEARS and plotted using RevGadgets. a) Regionalization scheme used in the
classification of taxa for analysis. b) Summary of 10 stochastic mappings from a posterior
distribution of trees including only taxa from the indicated clade (solid black line). For pie chart
probabilities at each node, see Figure S2.

Data Availability

Raw reads from PAFTOL are available in ENA (accession PRJEB35285) and assembled reads

from www.treeoflife.kew.org. Raw reads of the Ledebouriinae are available from the SRA

(accession PRJNA721471). Alignments, species trees, biogeography inputs/outputs are all

available on Dryad (*doi: currently in process*) (but try https://doi.org/10.5061/dryad.r4gxd2gw).

Biosketch

Cody Coyotee Howard is an Assistant Professor at Oklahoma State University broadly interested

in geophyte evolution and biogeography, and he has a particular fascination with bulbous plants.

The Ledebouriinae represent one bulbous lineage that has captivated him for years, and these

plants will continue to do so for many more.

Author contributions: Conceptualization CCH, TSH, NC; Formal analysis, CCH and ARZ;

Investigation, CCH and ARZ; Resources, TSH, LN, DC, LN, PM, ARZ, NC; Writing —

Original Draft, CCH and NC; Writing — Review & Editing, all authors; Visualization, CCH;

Funding acquisition, CCH and NC.

Supporting information

Table S1. Fossil and secondary calibration points used in the penalized likelihood (treePL)

analysis. With the exception of monocots, all calibrations were set as minimum values in the

analysis.

node	minimum calibration	source
Zingiberaceae	72.1	Iles et al., 2015
Typhaceae	51.66	Iles, et al. 2015
Cyperaceae	47.0	Iles, et al. 2015
Amaryllidaceae	48.88	Pigg et al., 2018
Coryphoideae	83.6	Iles, et al. 2015
Agavoideae	14.5	Iles, et al. 2015
Xanthorrhoeaceae	38.0	Iles, et al. 2015
Goodyerinae	15.0	Iles, et al. 2015
Monocots	131 – 135	Magallón et al., 2015

539 **Table S2**. Top ten most common transition types found in a subclade of *Ledebouria* Clade A

540 (Figure 3b) using stochastic mapping of biogeographic events.

starting_range	ending_range	average_n_events	SD	transition_type
K	K	23.432	2.121	in situ speciation
Z	Z	19.252	1.297	in situ speciation
K	KN	3.393	0.517	dispersal
T	T	2.764	2.089	in situ speciation

C	C	1.29	1.912	in situ speciation
KZ	K	1.09	0.307	vicariance or subset sympatry or extinction
KZ	Z	1.09	0.307	vicariance or subset sympatry or extinction
K	KZ	1.029	0.215	dispersal
CK	C	0.912	0.703	vicariance or subset sympatry or extinction

Figure S1. Full dated phylogeny with 95% highest posterior density plots shown.

Figure S2. Biogeographical reconstruction using the DIVA-like model with nodes showing the probable range for each region at that node.

Bibliography

Ali, S. S., Pfosser, M., Wetschnig, W., Martínez-Azorín, M., Crespo, M. B., & Yu, Y. (2013). Out of Africa: Miocene dispersal, vicariance, and extinction within Hyacinthaceae subfamily Urgineoideae. *Journal of Integrative Plant Biology*, *55*(10), 950–964. https://doi.org/10.1111/jipb.12065

Ali, S. S., Yu, Y., Pfosser, M., & Wetschnig, W. (2012). Inferences of biogeographical histories within subfamily Hyacinthoideae using S-DIVA and Bayesian binary MCMC analysis implemented in RASP (Reconstruct Ancestral State in Phylogenies). *Annals of Botany*,

109(1), 95–107. https://doi.org/10.1093/aob/mcr274

Andermann, T., Cano, Á., Zizka, A., Bacon, C., & Antonelli, A. (2018). SECAPR-a

bioinformatics pipeline for the rapid and user-friendly processing of targeted enriched

Illumina sequences, from raw reads to alignments. *PeerJ*, *6*, e5175.

https://doi.org/10.7717/peerj.5175

Baker, W. J., Bailey, P., Barber, V., Barker, A., Bellot, S., Bishop, D., Botigué, L. R., Brewer,

G., Carruthers, T., Clarkson, J. J., Cook, J., Cowan, R. S., Dodsworth, S., Epitawalage, N.,

Françoso, E., Gallego, B., Johnson, M. G., Kim, J. T., Leempoel, K., … Forest, F. (2022). A

comprehensive phylogenomic platform for exploring the angiosperm tree of life. *Systematic*

Biology, *71*(2), 301–319. https://doi.org/10.1093/sysbio/syab035

Balinsky, B. I. (1962). Patterns of animal distribution on the African continent. *Ann. Cape Prov.*

Mus, *2*, 299–310.

Bellstedt, D. U., Galley, C., Pirie, M. D., & Linder, H. P. (2012). The migration of the

palaeotropical arid flora: zygophylloideae as an example. *Systematic Botany*, *37*(4), 951–

959. https://doi.org/10.1600/036364412X656608

Bobe, R. (2006). The evolution of arid ecosystems in eastern Africa. *Journal of Arid*

Environments, *66*(3), 564–584. https://doi.org/10.1016/j.jaridenv.2006.01.010

Bonnefille, R. (2010). Cenozoic vegetation, climate changes and hominid evolution in tropical

Africa. *Global and Planetary Change*, *72*(4), 390–411.

https://doi.org/10.1016/j.gloplacha.2010.01.015

Bouckaert, R., Heled, J., Kühnert, D., Vaughan, T., Wu, C.-H., Xie, D., Suchard, M. A.,

Rambaut, A., & Drummond, A. J. (2014). BEAST 2: a software platform for Bayesian

evolutionary analysis. *PLoS Computational Biology*, *10*(4), e1003537.

https://doi.org/10.1371/journal.pcbi.1003537

Brown, J. W., Walker, J. F., & Smith, S. A. (2017). Phyx: phylogenetic tools for unix.

Bioinformatics, *33*(12), 1886–1888. https://doi.org/10.1093/bioinformatics/btx063

Bruyns, P. V., Klak, C., & Hanáček, P. (2014). Evolution of the stapeliads (Apocynaceae-

Asclepiadoideae) - repeated major radiation across Africa in an Old World group. *Molecular*

Phylogenetics and Evolution, *77*, 251–263. https://doi.org/10.1016/j.ympev.2014.03.022

Buerki, S., Jose, S., Yadav, S. R., Goldblatt, P., Manning, J. C., & Forest, F. (2012). Contrasting

biogeographic and diversification patterns in two Mediterranean-type ecosystems. *Plos One*,

7(6), e39377. https://doi.org/10.1371/journal.pone.0039377

Ceccarelli, F. S., Koch, N. M., Soto, E. M., Barone, M. L., Arnedo, M. A., & Ramírez, M. J.

(2019). The grass was greener: repeated evolution of specialized morphologies and habitat

shifts in ghost spiders following grassland expansion in south america. *Systematic Biology*,

68(1), 63–77. https://doi.org/10.1093/sysbio/syy028

Chen, C., Qi, Z.-C., Xu, X.-H., Comes, H. P., Koch, M. A., Jin, X.-J., Fu, C.-X., & Qiu, Y.-X.

(2014). Understanding the formation of Mediterranean-African-Asian disjunctions: evidence

for Miocene climate-driven vicariance and recent long-distance dispersal in the Tertiary

relict Smilax aspera (Smilacaceae). *The New Phytologist*, *204*(1), 243–255.

https://doi.org/10.1111/nph.12910

Coe, M. J., & Skinner, J. D. (1993). Connections, disjunctions and endemism in the eastern and

southern african mammal faunas. *Transactions of the Royal Society of South Africa*, *48*(2),

233–255. https://doi.org/10.1080/00359199309520273

Couvreur, T. L. P., Chatrou, L. W., Sosef, M. S. M., & Richardson, J. E. (2008). Molecular

phylogenetics reveal multiple tertiary vicariance origins of the African rain forest trees.

BMC Biology, *6*, 54. https://doi.org/10.1186/1741-7007-6-54

Couvreur, T. L. P., Dauby, G., Blach-Overgaard, A., Deblauwe, V., Dessein, S., Droissart, V.,

Hardy, O. J., Harris, D. J., Janssens, S. B., Ley, A. C., Mackinder, B. A., Sonké, B., Sosef,

M. S. M., Stévart, T., Svenning, J.-C., Wieringa, J. J., Faye, A., Missoup, A. D., Tolley, K.

A., … Sepulchre, P. (2021). Tectonics, climate and the diversification of the tropical African

terrestrial flora and fauna. *Biological Reviews of the Cambridge Philosophical Society*,

96(1), 16–51. https://doi.org/10.1111/brv.12644

Couvreur, T. L. P. (2015). Odd man out: why are there fewer plant species in African rain

forests? *Plant Systematics and Evolution*, *301*(5), 1299–1313.

https://doi.org/10.1007/s00606-014-1180-z

Cowling, R. M., Procheş, S., & Partridge, T. C. (2009). Explaining the uniqueness of the Cape

flora: incorporating geomorphic evolution as a factor for explaining its diversification.

Molecular Phylogenetics and Evolution, *51*(1), 64–74.

https://doi.org/10.1016/j.ympev.2008.05.034

Dagallier, L.-P. M. J., Janssens, S. B., Dauby, G., Blach-Overgaard, A., Mackinder, B. A.,

Droissart, V., Svenning, J.-C., Sosef, M. S. M., Stévart, T., Harris, D. J., Sonké, B.,

Wieringa, J. J., Hardy, O. J., & Couvreur, T. L. P. (2020). Cradles and museums of generic

plant diversity across tropical Africa. *The New Phytologist*, *225*(5), 2196–2213.

https://doi.org/10.1111/nph.16293

del Hoyo, A., García-Marín, J. L., & Pedrola-Monfort, J. (2009). Temporal and spatial

diversification of the African disjunct genus Androcymbium (Colchicaceae). *Molecular*

Phylogenetics and Evolution, *53*(3), 848–861. https://doi.org/10.1016/j.ympev.2009.08.005

Drummond, A. J., & Rambaut, A. (2007). BEAST: Bayesian evolutionary analysis by sampling

trees. *BMC Evolutionary Biology*, *7*, 214. https://doi.org/10.1186/1471-2148-7-214

Edgell, H. S. (2006). *Arabian deserts: nature, origin and evolution*. Springer Science & Business Media.

Emberts, Z., St Mary, C. M., Howard, C. C., Forthman, M., Bateman, P. W., Somjee, U., Hwang, W. S., Li, D., Kimball, R. T., & Miller, C. W. (2020). The evolution of autotomy in leaf-footed bugs. *Evolution*, *74*(5), 897–910. https://doi.org/10.1111/evo.13948

Esler, K. J., Rundel, P. W., & Vorster, P. (1999). Biogeography of prostrate-leaved geophytes in semi-arid South Africa: hypotheses on functionality. *Plant Ecology*, *142*(1), 105–120.

Fleitmann, D., Matter, A., Burns, S. J., Al-Subbary, A., & Al-Aowah, M. A. (2004). Geology and Quaternary climate history of Socotra. *Fauna of Arabia*, *20*, 27–44.

Galley, C., Bytebier, B., Bellstedt, D. U., & Linder, H. P. (2007). The Cape element in the Afrotemperate flora: from Cape to Cairo? *Proceedings. Biological Sciences / the Royal Society*, *274*(1609), 535–543. https://doi.org/10.1098/rspb.2006.0046

García-Aloy, S., Sanmartín, I., Kadereit, G., Vitales, D., Millanes, A. M., Roquet, C., Vargas, P., Alarcón, M., & Aldasoro, J. J. (2017). Opposite trends in the genus Monsonia (Geraniaceae): specialization in the African deserts and range expansions throughout eastern Africa. *Scientific Reports*, *7*(1), 9872. https://doi.org/10.1038/s41598-017-09834-6

Giranje, P. T., & Nandikar, M. D. (2016). Synopsis of the genus *Ledebouria* Roth (Hyacinthaceae: Hyacinthoideae) in India. *Webbia*, *71*(2), 213–217. https://doi.org/10.1080/00837792.2016.1182324

Givnish, T. J., Zuluaga, A., Spalink, D., Soto Gomez, M., Lam, V. K. Y., Saarela, J. M., Sass, C., Iles, W. J. D., de Sousa, D. J. L., Leebens-Mack, J., Chris Pires, J., Zomlefer, W. B., Gandolfo, M. A., Davis, J. I., Stevenson, D. W., dePamphilis, C., Specht, C. D., Graham, S.

W., Barrett, C. F., & Ané, C. (2018). Monocot plastid phylogenomics, timeline, net rates of

species diversification, the power of multi-gene analyses, and a functional model for the

origin of monocots. *American Journal of Botany*, *105*(11), 1888–1910.

https://doi.org/10.1002/ajb2.1178

Gizaw, A., Gorospe, J. M., Kandziora, M., Chala, D., Gustafsson, L., Zinaw, A., Salomón, L.,

Eilu, G., Brochmann, C., Kolář, F., & Schmickl, R. (2021). Afro-alpine flagships revisited

II: elucidating the evolutionary relationships and species boundaries in the giant senecios

(Dendrosenecio, Asteraceae). *Alpine Botany*. https://doi.org/10.1007/s00035-021-00268-5

Grace, O. M., Buerki, S., Symonds, M. R., Forest, F., van Wyk, A. E., Smith, G. F., Klopper, R.

R., Bjorå, C. S., Neale, S., Demissew, S., Simmonds, M. S., & Rønsted, N. (2015).

Evolutionary history and leaf succulence as explanations for medicinal use in aloes and the

global popularity of Aloe vera. *BMC Evolutionary Biology*, *15*(1), 291.

https://doi.org/10.1186/s12862-015-0291-7

Hagen, O., Skeels, A., Onstein, R. E., Jetz, W., & Pellissier, L. (2021). Earth history events

shaped the evolution of uneven biodiversity across tropical moist forests. *Proceedings of the*

National Academy of Sciences of the United States of America, *118*(40).

https://doi.org/10.1073/pnas.2026347118

Howard, C. C., Crowl, A. A., Harvey, T. S., & Cellinese, N. (2022). Peeling back the layers:

First phylogenomic insights into the Ledebouriinae (Scilloideae, Asparagaceae). *Molecular*

Phylogenetics and Evolution, *169*, 107430. https://doi.org/10.1016/j.ympev.2022.107430

Howard, C. C., Folk, R. A., Beaulieu, J. M., & Cellinese, N. (2019). The monocotyledonous

underground: global climatic and phylogenetic patterns of geophyte diversity. *American*

Journal of Botany, *106*(6), 850–863. https://doi.org/10.1002/ajb2.1289

Howard, C. C., Landis, J. B., Beaulieu, J. M., & Cellinese, N. (2020). Geophytism in monocots

leads to higher rates of diversification. *The New Phytologist*, *225*(2), 1023–1032.

https://doi.org/10.1111/nph.16155

Howard, C. C. (2014). In the shade of the mopane: *Ledebouria* in Namibia. *Cactus and*

Succulent Journal, *86*(3), 92–98. https://doi.org/10.2985/015.086.0301

Iles, W. J., Smith, S. Y., Gandolfo, M. A., & Graham, S. W. (2015). Monocot fossils suitable for

molecular dating analyses. *Botanical Journal of the Linnean Society*, *178*(3), 346–374.

Jacobs, B. F., Kingston, J. D., & Jacobs, L. L. (1999). The Origin of Grass-Dominated

Ecosystems. *Annals of the Missouri Botanical Garden*, *86*(2), 590.

https://doi.org/10.2307/2666186

Jacobs, B. F. (2004). Palaeobotanical studies from tropical Africa: relevance to the evolution of

forest, woodland and savannah biomes. *Philosophical Transactions of the Royal Society of*

London. Series B, Biological Sciences, *359*(1450), 1573–1583.

https://doi.org/10.1098/rstb.2004.1533

Johnson, M. G., Gardner, E. M., Liu, Y., Medina, R., Goffinet, B., Shaw, A. J., Zerega, N. J. C.,

& Wickett, N. J. (2016). HybPiper: extracting coding sequence and introns for phylogenetics

from high-throughput sequencing reads using target enrichment. *Applications in Plant*

Sciences, *4*(7). https://doi.org/10.3732/apps.1600016

Johnson, M. G., Pokorny, L., Dodsworth, S., Botigué, L. R., Cowan, R. S., Devault, A.,

Eiserhardt, W. L., Epitawalage, N., Forest, F., Kim, J. T., Leebens-Mack, J. H., Leitch, I. J.,

Maurin, O., Soltis, D. E., Soltis, P. S., Wong, G. K.-S., Baker, W. J., & Wickett, N. J.

(2019). A universal probe set for targeted sequencing of 353 nuclear genes from any

flowering plant designed using k-medoids clustering. *Systematic Biology*, *68*(4), 594–606.

https://doi.org/10.1093/sysbio/syy086

Jürgens, N. (1997). Floristic biodiversity and history of African arid regions. *Biodiversity and*

Conservation, *6*, 495–514. https://doi.org/10.1023/a:1018325026863

Kandziora, M., Gehrke, B., Popp, M., Gizaw, A., Brochmann, C., & Pirie, M. D. (2022). The

enigmatic tropical alpine flora on the African sky islands is young, disturbed, and

unsaturated. *Proceedings of the National Academy of Sciences of the United States of*

America, *119*(22), e2112737119. https://doi.org/10.1073/pnas.2112737119

Katoh, K., & Standley, D. M. (2013). MAFFT multiple sequence alignment software version 7:

improvements in performance and usability. *Molecular Biology and Evolution*, *30*(4), 772–

780. https://doi.org/10.1093/molbev/mst010

Kornas, J. (1985). Adaptive strategies of African pteridophytes to extreme environments.

Proceedings of the Royal Society of Edinburgh, Section B: Biological Sciences, *86*, 391–

396.

Lanfear, R., Calcott, B., Ho, S. Y. W., & Guindon, S. (2012). Partitionfinder: combined selection

of partitioning schemes and substitution models for phylogenetic analyses. *Molecular*

Biology and Evolution, *29*(6), 1695–1701. https://doi.org/10.1093/molbev/mss020

Lanfear, R., Calcott, B., Kainer, D., Mayer, C., & Stamatakis, A. (2014). Selecting optimal

partitioning schemes for phylogenomic datasets. *BMC Evolutionary Biology*, *14*, 82.

https://doi.org/10.1186/1471-2148-14-82

Leaché, A. D., Wagner, P., Linkem, C. W., Böhme, W., Papenfuss, T. J., Chong, R. A., Lavin, B.

R., Bauer, A. M., Nielsen, S. V., Greenbaum, E., Rödel, M.-O., Schmitz, A., LeBreton, M.,

Ineich, I., Chirio, L., Ofori-Boateng, C., Eniang, E. A., Baha El Din, S., Lemmon, A. R., &

Burbrink, F. T. (2014). A hybrid phylogenetic-phylogenomic approach for species tree

715  estimation in African Agama lizards with applications to biogeography, character evolution,

716  and diversification. *Molecular Phylogenetics and Evolution*, *79*, 215–230.

717  https://doi.org/10.1016/j.ympev.2014.06.013

718 Lebatha, P. D. (2004). *A systematic revision of Drimiopsis Lindl. & Paxt. (Hyacinthaceae)*

719  [Doctoral dissertation].

720 Linder, H. P. (2014). The evolution of African plant diversity. *Frontiers in Ecology and*

721  *Evolution*, *2*, 1–14. https://doi.org/10.3389/fevo.2014.00038

722 Linder, H. P. (2017). East African Cenozoic vegetation history. *Evolutionary Anthropology*,

723  *26*(6), 300–312. https://doi.org/10.1002/evan.21570

724 Li, H.-T., Yi, T.-S., Gao, L.-M., Ma, P.-F., Zhang, T., Yang, J.-B., Gitzendanner, M. A., Fritsch,

725  P. W., Cai, J., Luo, Y., Wang, H., van der Bank, M., Zhang, S.-D., Wang, Q.-F., Wang, J.,

726  Zhang, Z.-R., Fu, C.-N., Yang, J., Hollingsworth, P. M., … Li, D.-Z. (2019). Origin of

727  angiosperms and the puzzle of the Jurassic gap. *Nature Plants*, *5*(5), 461–470.

728  https://doi.org/10.1038/s41477-019-0421-0

729 Linder, H. P., de Klerk, H. M., Born, J., Burgess, N. D., Fjeldså, J., & Rahbek, C. (2012). The

730  partitioning of Africa: statistically defined biogeographical regions in sub-Saharan Africa.

731  *Journal of Biogeography*, *39*(7), 1189–1205. https://doi.org/10.1111/j.1365-

732  2699.2012.02728.x

733 Lorenzen, E. D., Heller, R., & Siegismund, H. R. (2012). Comparative phylogeography of

734  African savannah ungulates. *Molecular Ecology*, *21*(15), 3656–3670.

735  https://doi.org/10.1111/j.1365-294X.2012.05650.x

736 Lu, M., Fradera-Soler, M., Forest, F., Barraclough, T. G., & Grace, O. M. (2022). Evidence

737  linking life-form to a major shift in diversification rate in Crassula. *American Journal of*

Botany, *109*(2), 272–290. https://doi.org/10.1002/ajb2.1797

Magalhaes, I. L. F., Santos, A. J., & Ramírez, M. J. (2021). Incorporating Topological and Age

Uncertainty into Event-Based Biogeography of Sand Spiders Supports Paleo-Islands in

Galapagos and Ancient Connections among Neotropical Dry Forests. *Diversity*, *13*(9), 418.

https://doi.org/10.3390/d13090418

Magallón, S., Gómez-Acevedo, S., Sánchez-Reyes, L. L., & Hernández-Hernández, T. (2015). A

metacalibrated time-tree documents the early rise of flowering plant phylogenetic diversity.

The New Phytologist, *207*(2), 437–453. https://doi.org/10.1111/nph.13264

Main, D. C., Jansen van Vuuren, B., Tilbury, C. R., & Tolley Conceptualisation, K. A. (2022).

Out of southern Africa: origins and cryptic speciation in Chamaeleo, the most widespread

chameleon genus. *Molecular Phylogenetics and Evolution*, 107578.

https://doi.org/10.1016/j.ympev.2022.107578

Mankga, L. T., Yessoufou, K., & Chitakira, M. (2020). On the origin and diversification history

of the African genus Encephalartos. *South African Journal of Botany*, *130*, 231–239.

https://doi.org/10.1016/j.sajb.2019.12.007

Manning, J. C. (2020). Systematics of *Ledebouria* sect. *Resnova* (Hyacinthaceae: Scilloideae:

Massonieae), with a new subtribal classification of Massonieae. *South African Journal of*

Botany, *133*, 98–110. https://doi.org/10.1016/j.sajb.2020.07.010

Manzaneda, A. J., Rey, P. J., Bastida, J. M., Weiss-Lehman, C., Raskin, E., & Mitchell-Olds, T.

(2012). Environmental aridity is associated with cytotype segregation and polyploidy

occurrence in *Brachypodium distachyon* (Poaceae). *The New Phytologist*, *193*(3), 797–805.

https://doi.org/10.1111/j.1469-8137.2011.03988.x

Marsden, C. D., Woodroffe, R., Mills, M. G. L., McNutt, J. W., Creel, S., Groom, R.,

Emmanuel, M., Cleaveland, S., Kat, P., Rasmussen, G. S. A., Ginsberg, J., Lines, R., André,
J.-M., Begg, C., Wayne, R. K., & Mable, B. K. (2012). Spatial and temporal patterns of
neutral and adaptive genetic variation in the endangered African wild dog (*Lycaon pictus*).
Molecular Ecology, *21*(6), 1379–1393. https://doi.org/10.1111/j.1365-294X.2012.05477.x

Maswanganye, K. A., Cunningham, M. J., Bennett, N. C., Chimimba, C. T., & Bloomer, P.
(2017). Life on the rocks: Multilocus phylogeography of rock hyrax (Procavia capensis)
from southern Africa. *Molecular Phylogenetics and Evolution*, *114*, 49–62.
https://doi.org/10.1016/j.ympev.2017.04.006

Matasci, N., Hung, L.-H., Yan, Z., Carpenter, E. J., Wickett, N. J., Mirarab, S., Nguyen, N.,
Warnow, T., Ayyampalayam, S., Barker, M., Burleigh, J. G., Gitzendanner, M. A., Wafula,
E., Der, J. P., dePamphilis, C. W., Roure, B., Philippe, H., Ruhfel, B. R., Miles, N. W., …
Wong, G. K.-S. (2014). Data access for the 1,000 Plants (1KP) project. *GigaScience*, *3*, 17.
https://doi.org/10.1186/2047-217X-3-17

Matzke, N. J. (2013a). BioGeoBEARS: BioGeography with Bayesian (and likelihood)
evolutionary analysis in R Scripts. *R Package, Version 0.2*, *1*.

Matzke, N. J. (2013b). Probabilistic historical biogeography: new models for founder-event
speciation, imperfect detection, and fossils allow improved accuracy and model-testing.
Frontiers of Biogeography, *5*(4). https://doi.org/10.21425/F5FBG19694

Matzke, N. J. (2014). Model selection in historical biogeography reveals that founder-event
speciation is a crucial process in island clades. *Systematic Biology*, *63*(6), 951–970.
https://doi.org/10.1093/sysbio/syu056

Matzke, N. J. (2022). Statistical comparison ofDEC andDEC +J is identical to comparison of
twoClaSSE submodels, and is therefore valid. *Journal of Biogeography*.

https://doi.org/10.1111/jbi.14346

Miller, A. G., & Alexander, D. (2010). Haemanthus grandifolius Balf.f.: an enigma resolved.

Edinburgh Journal of Botany, *53*(1), 43–47. https://doi.org/10.1017/S0960428600002717

Morgan, M. E., Kingston, J. D., & Marino, B. D. (1994). Carbon isotopic evidence for the

emergence of C4 plants in the Neogene from Pakistan and Kenya. *Nature*, *367*(6459), 162–

165. https://doi.org/10.1038/367162a0

Mwafongo, E., Vollsnes, A. V., Bjorå, C. S., Nordal, I., & Eriksen, A. B. (2017). Leaf

mottling/variegation and shape in the Ledebouria revoluta complex – Development, stability

and putative function. *Flora*, *236–237*, 33–43. https://doi.org/10.1016/j.flora.2017.09.009

Neumann, F. H., & Bamford, M. K. (2015). Shaping of modern southern African biomes:

Neogene vegetation and climate changes. *Transactions of the Royal Society of South Africa*,

70(3), 195–212. https://doi.org/10.1080/0035919X.2015.1072859

Nguyen, L.-T., Schmidt, H. A., von Haeseler, A., & Minh, B. Q. (2015). IQ-TREE: a fast and

effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Molecular*

Biology and Evolution, *32*(1), 268–274. https://doi.org/10.1093/molbev/msu300

Nielsen, S. V., Daniels, S. R., Conradie, W., Heinicke, M. P., & Noonan, B. P. (2018).

Multilocus phylogenetics in a widespread African anuran lineage (Brevicipitidae: *Breviceps*

) reveals patterns of diversity reflecting geoclimatic change. *Journal of Biogeography*,

45(9), 2067–2079. https://doi.org/10.1111/jbi.13394

Nylinder, S., Razafimandimbison, S. G., & Anderberg, A. A. (2016). From the Namib around the

world: biogeography of the Inuleae-Plucheinae (Asteraceae). *Journal of Biogeography*,

43(9), 1705–1716. https://doi.org/10.1111/jbi.12764

Partridge, T. C., & Maud, R. R. (1987). Geomorphic evolution of southern Africa since the

Mesozoic. *South African Journal of Geology*, *90*(2), 179–208.

Pausas, J. G., Lamont, B. B., Paula, S., Appezzato-da-Glória, B., & Fidelis, A. (2018).

Unearthing belowground bud banks in fire-prone ecosystems. *The New Phytologist*, *217*(4),

1435–1448. https://doi.org/10.1111/nph.14982

Pfosser, M. (2012). Phylogenetic relationships of Malagasy Hyacinthaceae. *Plant Ecology and*

Evolution, *145*(1), 65–72. https://doi.org/10.5091/plecevo.2012.590

Pigg, K. B., Bryan, F. A., & DeVore, M. L. (2018). *Paleoallium billgenseli* gen. et sp. nov.:

Fossil Monocot Remains from the Latest Early Eocene Republic Flora, Northeastern

Washington State, USA. *International Journal of Plant Sciences*, *179*(6), 477–486.

https://doi.org/10.1086/697898

Pillon, Y., Hopkins, H. C. F., Maurin, O., Epitawalage, N., Bradford, J., Rogers, Z. S., Baker, W.

J., & Forest, F. (2021). Phylogenomics and biogeography of Cunoniaceae (Oxalidales) with

complete generic sampling and taxonomic realignments. *American Journal of Botany*,

108(7), 1181–1200. https://doi.org/10.1002/ajb2.1688

Pokorny, L., Riina, R., Mairal, M., Meseguer, A. S., Culshaw, V., Cendoya, J., Serrano, M.,

Carbajal, R., Ortiz, S., Heuertz, M., & Sanmartín, I. (2015). Living on the edge: timing of

Rand Flora disjunctions congruent with ongoing aridification in Africa. *Frontiers in*

Genetics, *6*, 154. https://doi.org/10.3389/fgene.2015.00154

Pound, M. J., Haywood, A. M., Salzmann, U., & Riding, J. B. (2012). Global vegetation

dynamics and latitudinal temperature gradients during the Mid to Late Miocene (15.97–

5.33Ma). *Earth-Science Reviews*, *112*(1–2), 1–22.

https://doi.org/10.1016/j.earscirev.2012.02.005

POWO. (2019). *Plants of the world online. Facilitated by the Royal Botanic Gardens, Kew.*

http://www.plantsoftheworldonline.org/

Proches, Ş., Cowling, R. M., Goldblatt, P., Manning, J. C., & DA Snijman. (2006). An overview

of the Cape geophytes. *Biological Journal of the Linnean Society*, *87*(1), 27–43.

Rambaut, A., Drummond, A. J., Xie, D., Baele, G., & Suchard, M. A. (2018). Posterior

summarization in Bayesian phylogenetics using Tracer 1.7. *Systematic Biology*, *67*(5), 901–

904. https://doi.org/10.1093/sysbio/syy032

R Core Team. (2016). *R: A language and environment for statistical computing.* (3.3.2)

[Computer software]. R Foundation for Statistical Computing. https://www.R-project.org/

Ree, R. H., & Sanmartín, I. (2018). Conceptual and statistical problems with the DEC+J model

of founder-event speciation and its comparison with DEC via model selection. *Journal of*

Biogeography, *45*(4), 741–749. https://doi.org/10.1111/jbi.13173

Rögl, F. (1999). Mediterranean and Paratethys. Facts and hypotheses of an Oligocene to

Miocene paleogeography (short overview). *Geologica Carpathica*, *50*(4), 339–349.

Sanmartín, I., Anderson, C. L., Alarcon, M., Ronquist, F., & Aldasoro, J. J. (2010). Bayesian

island biogeography in a continental setting: the Rand Flora case. *Biology Letters*, *6*(5),

703–707. https://doi.org/10.1098/rsbl.2010.0095

Senut, B., Pickford, M., & Ségalen, L. (2009). Neogene desertification of Africa. *Comptes*

Rendus Geoscience, *341*(8), 591–602.

Sepulchre, P., Ramstein, G., Fluteau, F., Schuster, M., Tiercelin, J.-J., & Brunet, M. (2006).

Tectonic uplift and Eastern Africa aridification. *Science*, *313*(5792), 1419–1423.

https://doi.org/10.1126/science.1129158

Sirichamorn, Y., Thomas, D. C., Adema, F. A. C. B., & van Welzen, P. C. (2014). Historical

biogeography of *Aganope*, *Brachypterum* and *Derris* (Fabaceae, tribe Millettieae): insights

into the origins of Palaeotropical intercontinental disjunctions and general biogeographical

patterns in Southeast Asia. *Journal of Biogeography*, *41*(5), 882–893.

https://doi.org/10.1111/jbi.12262

Smíd, J., Carranza, S., Kratochvíl, L., Gvoždík, V., Nasher, A. K., & Moravec, J. (2013). Out of

Arabia: a complex biogeographic history of multiple vicariance and dispersal events in the

gecko genus Hemidactylus (Reptilia: Gekkonidae). *Plos One*, *8*(5), e64018.

https://doi.org/10.1371/journal.pone.0064018

Smith, S. A., Brown, J. W., & Walker, J. F. (2018). So many genes, so little time: A practical

approach to divergence-time estimation in the genomic era. *Plos One*, *13*(5), e0197433.

https://doi.org/10.1371/journal.pone.0197433

Smith, S. A., & O'Meara, B. C. (2012). treePL: divergence time estimation using penalized

likelihood for large phylogenies. *Bioinformatics*, *28*(20), 2689–2690.

https://doi.org/10.1093/bioinformatics/bts492

Sonnleitner, M., Flatscher, R., Escobar García, P., Rauchová, J., Suda, J., Schneeweiss, G. M.,

Hülber, K., & Schönswetter, P. (2010). Distribution and habitat segregation on different

spatial scales among diploid, tetraploid and hexaploid cytotypes of Senecio carniolicus

(Asteraceae) in the Eastern Alps. *Annals of Botany*, *106*(6), 967–977.

https://doi.org/10.1093/aob/mcq192

Speta, F. (1998a). Hyacinthaceae. *Flowering Plants: Monocotyledons*, 261–285.

Speta, F. (1998b). Systematische analyse der gattung Scilla L. sl (Hyacinthaceae). *Phyton*.

Stamatakis, A. (2014). RAxML version 8: a tool for phylogenetic analysis and post-analysis of

large phylogenies. *Bioinformatics*, *30*(9), 1312–1313.

https://doi.org/10.1093/bioinformatics/btu033

Stedje, B, & Nordal, I. (1987). Cytogeographical studies of Hyacinthaceae in Africa south of the

Sahara. *Nordic Journal of Botany*, *7*(1), 53–65.

Stedje, Brita. (1998). Phylogenetic relationships and generic delimitation of sub-SaharanScilla

(Hyacinthaceae) and allied African genera as inferred from morphological and DNA

sequence data. *Plant Systematics and Evolution*, *211*(1–2), 1–11.

https://doi.org/10.1007/BF00984908

Tamar, K., Carranza, S., Sindaco, R., Moravec, J., Trape, J.-F., & Meiri, S. (2016). Out of

Africa: Phylogeny and biogeography of the widespread genus Acanthodactylus (Reptilia:

Lacertidae). *Molecular Phylogenetics and Evolution*, *103*, 6–18.

https://doi.org/10.1016/j.ympev.2016.07.003

Tribble, C. M., Martínez-Gómez, J., Howard, C. C., Males, J., Sosa, V., Sessa, E. B., Cellinese,

N., & Specht, C. D. (2021). Get the shovel: morphological and evolutionary complexities of

belowground organs in geophytes. *American Journal of Botany*, *108*(3), 372–387.

https://doi.org/10.1002/ajb2.1623

Venter, S. (1993). *A revision of the genus Ledebouria Roth (Hyacinthaceae) in South Africa.*

[Doctoral dissertation].

Venter, S. (2008). Synopsis of the genus Ledebouria Roth (Hyacinthaceae) in South Africa.

Herbertia, *62*, 85–155.

Whybrow, P. J., & McClure, H. A. (1980). Fossil mangrove roots and palaeoenvironments of the

miocene of the eastern Arabian Peninsula. *Palaeogeography, Palaeoclimatology,*

Palaeoecology, *32*, 213–225. https://doi.org/10.1016/0031-0182(80)90041-3

Wichura, H., Jacobs, L. L., Lin, A., Polcyn, M. J., Manthi, F. K., Winkler, D. A., Strecker, M. R.,

& Clemens, M. (2015). A 17-My-old whale constrains onset of uplift and climate change in

east Africa. *Proceedings of the National Academy of Sciences of the United States of*

America, *112*(13), 3910–3915. https://doi.org/10.1073/pnas.1421502112

Yang, Y.-Y., Meng, Y., Wen, J., Sun, H., & Nie, Z.-L. (2016). Phylogenetic analyses of Searsia

(Anacardiaceae) from eastern Asia and its biogeographic disjunction with its African

relatives. *South African Journal of Botany*, *106*(1), 129–136.

https://doi.org/10.1016/j.sajb.2016.05.021

Yu, X.-Q., Maki, M., Drew, B. T., Paton, A. J., Li, H.-W., Zhao, J.-L., Conran, J. G., & Li, J.

(2014). Phylogeny and historical biogeography of *Isodon* (Lamiaceae): rapid radiation in

south-west China and Miocene overland dispersal into Africa. *Molecular Phylogenetics and*

Evolution, *77*, 183–194. https://doi.org/10.1016/j.ympev.2014.04.017

Zachos, J., Pagani, M., Sloan, L., Thomas, E., & Billups, K. (2001). Trends, rhythms, and

aberrations in global climate 65 Ma to present. *Science*, *292*(5517), 686–693.

https://doi.org/10.1126/science.1059412

Zhou, L., Su, Y. C. F., Thomas, D. C., & Saunders, R. M. K. (2011). 'Out-of-Africa' dispersal of

tropical floras during the Miocene climatic optimum: evidence from Uvaria (Annonaceae).

Journal of Biogeography, *39*(2), 322–335. https://doi.org/10.1111/j.1365-

2699.2011.02598.x