From southern Africa and beyond: historical biogeography of a monocotyledonous bulbous

geophyte

Running title: Ledebouriinae (Scilloideae, Asparagaceae) historical biogeography

Cody Coyotee Howard[1*], Leevi Nanyeni[2], Neduvoto Mollel[3], David Chuba[4], Alexandre R.

Zuntini[5], Panagiota Malakasi[5], Timothy S. Harvey[6], Nico Cellinese[7]

[1] Department of Plant Biology, Ecology, and Evolution, Oklahoma State University, Stillwater,

OK, USA

[2] National Botanical Research Institute, Windhoek, Namibia

[3] Tropical Pesticides Research Institute (TPRI), National Herbarium, Arusha, Tanzania

[4] Department of Biological Sciences, University of Zambia, Lusaka, Zambia

[5] Royal Botanic Gardens, Kew, Richmond, Surrey, TW9 3AE UK

[6] Plantae Novae, Thousand Oaks, CA, USA

[7] Florida Museum of Natural History, University of Florida, Gainesville, FL, USA

* corresponding author: cody.howard@okstate.edu

Acknowledgements

We are extremely grateful to Silke Rügheimer and Esmerialda Strauss at the National Botanical

Research Institute in Windhoek, Namibia for their assistance with fieldwork and specimen

export while in Namibia. We would also like to thank Inge Pehlemann for being a wonderful

companion during excursions in search of Namibian *Ledebouria*. We are grateful to the

governments of Namibia, Zambia, and Tanzania for issuing collection and export permits.

Namibian collections were made under permit numbers 1784/2013, 1908/2014, 2056/2016, and

2185/2016. Collections from Zambia were made under permit number TJ/DNPW/101/13/18.

Tanzanian collections were made under permit number 2017-22-NA-2016-247. Plants were

imported under USDA permit numbers P37-09-00910, P37-16-00181, and P37-16-01462. Many

thanks to Dylan Hannon, Gottfriend Milkuhn, Tom Cole, and Tom McCoy for providing leaf

material from their collections. Lastly, we thank Killian Fleurial, Taylor LaVal, and Emily B.

Sessa for assistance while in the field, and William Baker for assistance in obtaining sequence

data for outgroup taxa. Funds from the Huntington Botanical Gardens, the Cactus and Succulent

Society of America, the Pacific Bulb Society, the San Gabriel Cactus and Succulent Society, the

Florida Museum of Natural History, the American Society of Plant Taxonomists, the University

of Florida International Center, the University of Florida Department of Biology, the Botanical

Society of America, the Society of Systematic Biologists, Xeric Growers, and numerous private

donors helped to support this work.

Abstract

Aim

Within sub-Saharan Africa, plants inhabiting more seasonal and arid landscapes showcase

unique distributional patterns that hint at fascinating evolutionary histories. The Ledebouriinae

(Scilloideae, Asparagaceae) are widespread throughout such climates in sub-Saharan Africa, and

Madagascar, the Middle East, India, and Sri Lanka. Long-distance dispersal has been

hypothesized as leading to such a widespread distribution; however, low taxon sampling and

taxonomic uncertainties have made uncovering the history of the Ledebouriinae difficult. Here,

using the most comprehensive sampling of the lineage to date, we hypothesize that both

vicariance and dispersal events impacted the biogeographical history of these bulbous monocots

within and outside of Africa.

Location

Sub-Saharan Africa, Madagascar, Asia

Taxon

Ledebouriinae (Scilloideae, Asparagaceae)

Methods

We infer age estimates using penalized likelihood as implemented in treePL. Capitalizing on our

broad geographic sampling, we use 'BioGeoBEARS' to reconstruct ancestral ranges and

investigate the role of vicariance and dispersal.

Results

Our results suggest the Ledebouriinae originated within the past ~30 myr in southeastern sub-Saharan Africa, with the major subclades arising soon thereafter. Although long-distance dispersal cannot be fully ruled out, our results lead us to hypothesize vicariance was the major process responsible for the current distribution of *Ledebouria* in Eurasia. We recover two distinct *Ledebouria* groups that overlap in eastern Africa, but are divided into mostly northern and southern clades with divergent biogeographical histories, and each showing an independent dispersal to Madagascar. A similar north-south split is seen in *Drimiopsis*. Additionally, we recover a complex biogeographic history in the predominantly sub-Saharan African *Ledebouria* clade, with a rapid radiation estimated at ~14 mya.

Main conclusions

We recover evidence to suggest that the expansion of seasonal rainfall and aridity in sub-Saharan Africa, coupled with orogeny, may have fostered the diversification of the Ledebouriinae and many subclades. Miocene-driven aridification may have caused fragmentation of a once widespread distribution that led to their occurrence in Eurasia.

Keywords *Drimiopsis*, geophytes, *Ledebouria*, phylogenomics, *Resnova*, Scilloideae

1 INTRODUCTION

Modern-day Africa is dominated by arid and semi-arid landscapes that contain a diversity of habitats from deserts, woodlands and savannas, to name a few (Bobe, 2006; Linder, 2014). In sub-Saharan Africa, these drier ecosystems collectively form a fairly continuous sickle-shaped corridor that connects the floras of southwestern, northeastern and western Africa, and that skirts around the wet tropics of central and western Africa (Balinsky, 1962; Bellstedt et al., 2012; Jürgens, 1997) (Figure 1). The gradual increase in aridification in Africa is hypothesized to have been caused by the synergistic activities of rapid global cooling, tectonic events (e.g., Eastern African rift) and oceanic upwelling (i.e., Benguela current) that altered precipitation patterns across the continent (Bobe, 2006; Couvreur et al., 2021; Hagen et al., 2021; Linder, 2017; Senut et al., 2009; Sepulchre et al., 2006). These changes had immense impacts on the evolutionary trajectories of many lineages on the African continent. For example, in the early Eocene, equatorial Africa was largely covered by tropical forests, but aridification repeatedly fragmented and reduced these habitats over time (Couvreur, 2015; Couvreur et al., 2008; Hagen et al., 2021). Evidence from plant fossils and phylogenetic studies suggest that lineages with past widespread distributions across the continent, as well as between eastern Africa and southern Asia, are now disjunct as a result of recent aridification (Ali et al., 2013; Jacobs et al., 1999; Pokorny et al., 2015; Sanmartín et al., 2010; Zhou et al., 2011). Conversely, dry and more seasonal climates have been linked to an increase in the diversification and dispersal of lineages both in and out of Africa, and today, a rich diversity of taxa can be found within such climates in Africa as well as surrounding areas (e.g., Arabian Peninsula) (Bruyns et al., 2014; Coe & Skinner, 1993; Jürgens, 1997; Lorenzen et al., 2012; Nylinder et al., 2016). Therefore, it is imperative we gain a broad understanding of the historical evolution of diverse lineages across different landscapes, habitats,

and biomes, especially in fragmented regions, in order to continually refine our understanding of

the past and improve our ability to predict the ways in which African biodiversity may be shaped

in the future.

Geophytes, herbaceous plants with renewal buds located belowground on structures such

as bulbs, corms, and stem tubers, are ubiquitous components of seasonal or disturbance-prone

habitats, and are phylogenetically diverse (Howard et al., 2019; Pausas et al., 2018; Tribble et al.,

2021). Within Africa, geophytes are major components of the Greater Cape Floristic Region and

the Mediterranean Basin (Buerki et al., 2012; Proches et al., 2006). Although geophytes are

predominant elements of these two areas, the geophytic habit is widespread throughout sub-

Saharan Africa, particularly within seasonal or disturbance-prone (e.g., fire) habitats (Esler et al.,

1999; Kornas, 1985). Studies have reported that many African geophytes' origins coincide with

the onset of increased seasonality and/or aridity within the continent (i.e., since the Eocene), with

the majority of diversity evolving in response to the relatively more drastic climatic changes

since the Oligocene/Miocene (Ali et al., 2012, 2013; Buerki et al., 2012; del Hoyo et al., 2009;

Proches et al., 2006). At a broad scale, therefore, geophytic lineages make excellent candidates

for understanding relatively recent evolutionary and biogeographic dynamics within seasonal and

arid climates.

Scilloideae (Asparagaceae) are a bulbous geophyte lineage widespread both within and

outside of Africa (Speta, 1998a). This monocotyledonous clade consist of 1,000+ taxa found

throughout seasonal climates in Africa as well as Madagascar, Europe, the Middle East, and

Asia, with a single lineage found in South America (i.e., *Oziroe* Raf.; Giranje & Nandikar, 2016;

Speta, 1998a). Although only a handful of studies have investigated the historical biogeography

of the Scilloideae and its subclades, so far, all have pointed to sub-Saharan Africa as the origin

for the majority of the group (excluding *Oziroe* in South America), followed by a complex

history within and outside of the continent (Ali et al., 2012, 2013; Buerki et al., 2012; Pfosser,

2012). Age estimates for the Scilloideae suggest an origin within the past ~70–50 myr, and major

subclades within the past ~60–50 myr (Ali et al., 2013; Buerki et al., 2012); however, most

inferences have been made with low taxon sampling, and in biogeographical studies sub-Saharan

Africa has been broadly subdivided into one to three areas (Ali et al., 2012, 2013; Buerki et al.,

2012; Pfosser, 2012). Regardless, given the current age estimates, we hypothesize that the

majority of Scilloideae lineages diversified in response to an increase in climatic seasonality and

aridification that intensified from the Eocene onwards. Focusing on widespread groups within

the Scilloideae using a more detailed approach to the regionalization of Africa as well as a

greater taxon sampling may provide refined insights into the biogeographical processes that have

impacted the dispersal of the Scilloideae as well as other plant lineages found across and outside

of Africa.

The Ledebouriinae are an ideal group to study because they are widespread within sub-

Saharan Africa, with a handful of taxa found in Madagascar, Socotra, Yemen, India, and Sri

Lanka (Giranje & Nandikar, 2016; Venter, 1993) (Figure 1). This distribution is unique within

Scilloideae since many sympatric lineages with Ledebouriinae are also found in northern Africa

(Pfosser, 2012; Speta, 1998b), but the Ledebouriinae are absent from the Scilloideae-rich

Mediterranean Basin (Venter, 1993, 2008) (Figure 1). In sub-Saharan Africa, the Ledebouriinae

are predominantly found within regions that experience oscillating weather patterns of wet and

dry and/or cold and warm, with highest diversity in the Limpopo, Mpumalanga, and KwaZulu-

Natal regions of South Africa (Venter, 1993), yet some occurrences are documented from more

wet, tropical regions (e.g., Gabon) (Figure 1). Much diversity within the Ledebouriinae,

however, remains undescribed to science (Howard, 2014; Howard et al., 2022). Additionally, an

expanded phylogenomic analysis of the group suggests that a complex biogeographical history

awaits to be thoroughly examined (Howard et al., 2022). Previous dating analyses have estimated

the origin of the Ledebouriinae sometime within the last ~25 myr in sub-Saharan Africa (Ali et

al., 2012; Buerki et al., 2012). However, this clade was not the focus of study and therefore,

taxon sampling was low. Additionally, results were unable to provide fine-scale biogeographical

patterns since sub-Saharan Africa was considered and analyzed as one large area. Regardless,

given our current understanding of the Ledebouriinae distribution, their dispersal ability, and a

recent phylogenomic study on this group (Howard et al., 2022), we can generate several

hypotheses regarding major biogeographical events throughout their history.

Although Ledebouriinae diversity is highest within Africa, many taxa are also found

outside of the continent (Venter, 1993). However, this intriguing distribution has yet to be the

explicit focus of investigation. Previous phylogenetic studies have hypothesized that multiple

dispersals to Madagascar from mainland Africa have occurred within the Ledebouriinae (Pfosser,

2012). For example, within Madagascar, we find two endemics: *L. nossibeensis* and an

undescribed taxon with morphological affinities to *L. socialis* (Pfosser, 2012). In addition to their

morphological differences, these two Malagasy taxa are also found in distinct regions of the

island (northwest vs southwest; see Figure 1). Madagascar has also been hypothesized to have

played a role in the dispersal of *Ledebouria* to India. For example, the phylogenetic

reconstructions of Ali et al. (2012) recovered Malagasy and Indian *Ledebouria* as sisters, which

led the authors to invoke long-distance dispersal from Madagascar to India. However, as stated

by the authors, taxa from eastern, western, and northern sub-Saharan Africa were absent, limiting

confidence in these conclusions (Ali et al., 2012). Furthermore, low phylogenetic resolution from

Pfosser (2012) diminishes confidence in the hypothesized independent dispersal events to

Madagascar. Recently, a phylogenomic analysis of the Ledebouriinae provided the framework to

suggest (without testing) multiple dispersals to Madagascar, and a potential migration out of

Africa via the Arabian Peninsula into India (Howard et al., 2022). Therefore, we hypothesize that

at least two independent dispersals to Madagascar have occurred in the Ledebouriinae. This is

based on the distinct distribution and morphotypes of Malagasy taxa, as well as their independent

phylogenetic placement (Howard et al., 2022; Pfosser, 2012). Regarding dispersals out of Africa,

we hypothesize that a once widespread distribution between Africa and India by way of the

Middle East fragmented due to Miocene-driven aridification, isolating populations and

contracting the overall distribution ranges. This hypothesis is drawn from several lines of

evidence. First, *Ledebouria* seed has limited dispersal capacity, which occurs primarily via sheet

water flow (i.e., dispersal occurs only short distances and is rainfall dependent) (Venter, 1993).

In years of limited rainfall and/or in flat terrains, seeds and seedlings can be found surrounding

the parent plant (CCH, pers. obs.), and the ephemeral nature of the seeds in some taxa warrants

germination soon after fruit dehiscence (CCH and TSH, pers. obs.). Therefore, we hypothesize

that long distance dispersal from Africa or Madagascar to India is unlikely. Further evidence for

a once-widespread, now-fragmented distribution is found in the current distribution of the

Ledebouriinae in the Arabian Peninsula. Today, *L. yemenensis* is the only Ledebouriinae

currently recorded from the region, and it is endemic to the cooler, higher elevations of the

Yemeni Highlands that are surrounded by arid lowlands. We hypothesize this taxon represents a

relic of a once much larger range. Additionally, there are multiple, endemic taxa to the Socotra

archipelago (Miller & Morris, 2004). For example, *L. grandifolia* and a putative *L.* aff. *revoluta*

are found only on the island of Socotra, and *L. insularis* is endemic to the summit of Samha,

where low clouds provide a moist habitat relative to the surrounding desert landscape (Miller &

Morris, 2004). Therefore, given the multiple taxa located on different islands in the Socotra

archipelago, we hypothesize that vicariance led to allopatric speciation as the islands moved

farther away from the mainland. In summary, although the Ledebouriinae occur within

seasonally dry and arid regions, many taxa may not be adapted to intense aridity, such as that

found in much of the Middle East and Pakistan, that intensified during the Miocene.

The unique, widespread distribution of the Ledebouriinae provides us with the

opportunity to refine our understanding of evolution and biogeography within Africa, potentially

during a time of extensive climatic, geologic, and habitat change on the continent. Here, using a

detailed categorization of the African continent, we investigate the timing and historical dispersal

of the Ledebouriinae both within and out of Africa to test major biogeographical hypotheses: 1)

due to high diversity within southern Africa, this region may represent the putative origin of the

Ledebouriinae; 2) at least two independent dispersals occurred to Madagascar from mainland

Africa; 3) vicariance is the major biogeographical process leading to the widespread distribution

of *Ledebouria* in Eurasia. Our increased sampling of the Ledebouriinae allows for a more

thorough investigation into the evolutionary history of this group and will improve our

knowledge of biogeographical patterns and processes within Africa.

2 METHODS

2.1 Phylogenetic analysis

Ledebouriinae samples were obtained from the field, private collections, and herbarium

vouchers (Howard et al., 2022). DNA extractions were performed using a modified CTAB

protocol, followed by high throughput sequencing on an Illumina HiSeq using the

Angiosperms353 universal probe set (Johnson et al., 2019). Raw reads were cleaned using

SECAPR (Andermann et al., 2018), sequences were pulled using HYBPIPER (Johnson et al., 2016)

and aligned using MAFFT v.7 (Katoh & Standley, 2013). See Howard et al. (2022) for more

details on Ledebouriinae sequence acquisition and analysis as well as data used, which can all be

found on Dryad (https://doi.org/10.5061/dryad.nzs7h44q6).

We incorporated outgroup taxa from the Plant and Fungal Tree of Life project (Baker et

al., 2022) and the 1KP dataset (Matasci et al., 2014). Exons were only available for the outgroup

taxa included to estimate divergence times using fossils and secondary calibration points.

Phylogenetic reconstruction including the Ledebouriinae plus outgroups was performed on a

concatenated, partitioned supermatrix of exons with 10% gaps/ambiguous sites removed using

phyx (Brown et al., 2017). This matrix was analyzed using IQ-TREE v.2-rc1 (Nguyen et al., 2015)

with 1000 ultrafast bootstraps, a best-fit partitioning scheme using the greedy algorithm of

PARTITIONFINDER (Lanfear et al., 2012), and a relaxed clustering percentage of 10 (Lanfear et al.,

2014), followed by phylogenetic reconstruction (-m TESTMERGE). However, the exon-only

dataset returned low support for many nodes within the Ledebouriinae. Therefore, we reran the

IQ-TREE analysis with a topological constraint tree (-g) that was previously built from a

supercontig (i.e., exons + introns) dataset of the Ledebouriinae (Howard et al., 2022).

2.2 Time calibration

We incorporated eight outgroup fossil calibration points, each with a minimum age

specified in Iles et al. (2015) (Table S1). A secondary calibration point at the crown node of

monocots was inferred between 131–135 mya based on previous analyses (Givnish et al., 2018;

Magallón et al., 2015).

Given the size of the dataset, we used penalized likelihood as implemented in TREEPL

(Smith & O'Meara, 2012) for time calibration. To incorporate uncertainty around age estimation,

we took a multi-tiered approach (see https://github.com/sunray1/treepl) similar to previous

studies estimating divergence times using large phylogenetic datasets (Emberts et al., 2020; Li et

al., 2019; Magallón et al., 2015). We generated 100 bootstrap replicates of our original exon

supermatrix alignment using RAXML v.8.2.0 (f -j option) (Stamatakis, 2014). A maximum

likelihood tree for each replicate with a corresponding partition file was then reconstructed using

a topological constraint (i.e., the phylogeny from the IQ-TREE analysis) to ensure consistent

calibration point placement, and a GTRGAMMA model of evolution. The resulting 100 "best

trees" were rooted on *Acorus gramineus* using phyx (Brown et al., 2017). In Step 1, each

replicate tree had a priming step completed with a random seed number and the thorough

command invoked. Step 2 was performed three independent times to assess convergence on the

best smoothing parameter for each individual tree. Step 2 also included the individual outputs

from each tree's previous priming step (e.g., optad, moredetail, etc.) as well as the cross-

validation (CV) steps, which were set to cvstart=10000, cvstop= .00000000000001,

cvmultstep=0.09. Lastly, Step 3 summarized each individual tree's CV output to determine and

scale each tree using the appropriate smoothing parameter. The 100 ultrametric phylogenies were

summarized using TREEANNOTATOR v.1.10.4 (Bouckaert et al., 2014) to obtain a maximum clade

credibility tree with 95% confidence intervals around each node and median node heights.

2.3 Biogeographical analysis

The combined biogeographical regionalization of Linder et al. (2012) (see Figure 2 in

Linder et al., 2012) was used for categorizing the location of each sub-Saharan African

Ledebouriinae accession. The exact distribution of populations/taxa represented by many field-

collected individuals (i.e., collections made by C.C. Howard) remains to be fully assessed since

they are undescribed species and/or are only known from one locality. Additionally, many

described species are currently known as occurring in small geographic ranges or even single

mountain tops (Lebatha, 2004; Venter, 2008). Consequently, most samples were assigned only to

known areas of occurrence or areas they were collected. Likewise, accessions found outside of

Africa were coded according to their actual known provenance: Madagascar, Yemen, Socotra,

India, or Sri Lanka. Even when seemingly widespread, taxa were coded using their documented

occurrence, rather than their putative distribution inferred by ambiguous taxonomic knowledge.

For example, *Ledebouria revoluta* is reported from the Cape Region of South Africa to Sri

Lanka (Mwafongo et al., 2017; Venter, 1993), however, this widespread taxon is poorly

understood, and it may harbor undescribed diversity as evidenced by the wide range of

morphological variation across its entire distribution range (Stedje, 1998). Its diversity has not

been investigated and therefore, this name is often applied (or misapplied) to individuals whose

species identification is difficult (Mwafongo et al., 2017). Consequently, and not surprisingly, *L.*

revoluta has become a 'bucket' species, an ambiguous and complex entity as evidenced by its

polyphyly (Howard et al., 2022).

In total, 14 geographic regions were used to code taxa. To estimate ancestral ranges, we

compared the maximum likelihood implementations of three biogeographical models, DEC,

DIVA-like, and BAYAREA-like as implemented in 'BioGeoBEARS' (Matzke, 2013a, 2013b,

2014) in R (2016). We used log-likelihood and AICc values to determine the best model between

the three. Given the debate surrounding the use of models with the +j parameter (Matzke, 2022;

Ree & Sanmartín, 2018), we decided to not include them and, additionally, their implementation

tends to infer a high number of jump dispersals, which are unlikely biological scenarios for the

majority of the Ledebouriinae (see Introduction and Discussion).

2.3.1 Biogeographical uncertainty

Within a subclade of *Ledebouria* Clade A, phylogenetic relationships are poorly

supported (Figure 2, node a) (Howard et al., 2022). However, given the potential rapid radiation

along the backbone of this group, we wanted to explore the biogeography of this clade since it

may contain an interesting history. To compensate for the uncertainty, we incorporated

topological and branch depth variability into our biogeographical analysis by using a random

sampling of trees from the posterior of a Bayesian analysis for the subclade of interest

(Ceccarelli et al., 2019; Magalhaes et al., 2021). To begin, we reduced our total alignment to the

most clock-like genes using scripts available in SortaDate (Smith et al., 2018). This was done

due to failure to reach convergence when using the entire alignment (data not shown). We kept

genes that were at least 10% concordant (bipartition > 0.1), had a tree length greater than 7.24,

and had root-to-tip variation of less than 0.009. Cutoff values for the latter two were determined

using the median values for all gene trees, as previously reported (Pillon et al., 2021). Of the

remaining alignments, we included only taxa within the subclade of interest (Figure 2b) and

removed duplicate Ledebouriinae outgroup taxa as well as those putatively duplicate within the

Ledebouriinae ingroup (indicated by * in Figure 2). Each gene was then aligned using MAFFT

invoking the --auto option. Sites with >5% gaps/ambiguity were removed using phyx. Due to

failure to reach convergence using a partitioned, concatenated alignment in BEAST v1.10.4

(Drummond & Rambaut, 2007), we instead input an unpartitioned, concatenated alignment using

a GTR+GAMMA model of substitution, a speciation birth-death prior, and a lognormal

distribution with an offset 17.1, mean 1.0, stdev 1.0 (95% range 17.24–19.88) on the root node of

the subclade. This initial value was chosen based on the treePL divergence time estimates for this

node. We ran an MCMC chain of 300k generations sampling every 10k generations.

Convergence was assessed using TRACER v1.7.1 (Rambaut et al., 2018) to ensure all ESS values

were above 200. We removed 50% of trees as burnin and used the remaining trees in the

analysis. We randomly sampled 100 trees after burnin, and performed a DIVA-like analysis (i.e.,

the best model) on each tree followed by stochastic mapping using 'BioGeoBEARS'. Trees were

summarized to obtain an average number of biogeographical events within and between each

region included in the analysis. Scripts for running the analysis can be found on github

(https://github.com/ivanlfm/BGB_BSM_multiple_trees). The rationale for incorporating

phylogenetic uncertainty was developed by Ceccarelli et al. (2019) and Magalhaes et al. (2021).

3 RESULTS

3.1 Phylogenetic relationships and age estimates

We recovered a polyphyletic *Ledebouria*, and a monophyletic *Drimiopsis* and *Resnova*,

which is congruent with a recent study (Howard et al., 2022). However, shallow-level

relationships differed compared to Howard et al. (2022), particularly within the *Ledebouria*

clades, and especially within *Ledebouria* Clade A. This is due to using a topological constraint (-

g), which searches tree parameter space, over using a fixed topology (-te) which does not

perform a tree search (see http://www.iqtree.org/doc/Command-Reference). Given differences in

the inputs between Howard et al. (2022) and our exons-only datasets, we preferred a topological

constraint allowing for a tree search over forcing a fixed topology. One major difference between

our results and those of Howard et al. (2022) can be found in *Ledebouria* Clade B, specifically

the subclade containing mostly non-African taxa (i.e., *L. grandifolia* + *L. revoluta* India).

Howard et al. (2022) used supercontigs (exons + flanking intronic region) for phylogenetic

reconstruction, and they recovered *L. grandifolia* (Socotra) as sister to a clade containing *L.*

yemenensis (Yemen), which was sister to the remaining taxa. All branches within the clade were

recovered with high ultrafast bootstrap support (UF-BOOT). In this study (Figure 2), using only

exons, we recovered *L. yemenensis* as sister to the same clade with high support, not *L.*

grandifolia, which was instead recovered as sister to *L. revoluta* (Eritrea). Additionally, the node

proceeding *L. yemenensis* was recovered with low support (UF-BOOT of 76).

330   Within a penalized likelihood framework, the Scilloideae originated approximately 52.0

mya (95% CI 51.2–54.5) (Figure 2; Table 1). We recovered a median crown age estimate for the

Ledebouriinae of 28.0 myr (95% CI 26.1–28.3), with age estimates of the four major

Ledebouriinae clades soon thereafter. The split between *Drimiopsis* and *Ledebouria* Clade B was

estimated at 27.1 mya (95% CI 25.2–27.5). The crown of *Drimiopsis* was dated to 22.1 mya

(95% CI 20.5–23.5), and *Ledebouria* Clade B was dated at 26.1 mya (95% CI 24.0–26.5). The

split between *Resnova* and *Ledebouria* Clade A occurred at 26.3 mya (95% CI 23.7–25.2). The

Resnova crown was dated at 17.9 mya (95% CI 16.6–18.5), and *Ledebouria* Clade A was

estimated at 25.0 mya (95% CI 23.0–25.4) (Table 1).

3.2 Biogeography

340   The DIVA-like model produced the most likely ancestral range estimates among the

models (LnL -220.54) (Table 2). The regions Natal+Zambezian were reconstructed as the

ancestral range for the Ledebouriinae ($p = 0.53$) (the next best area being Kalahari + Zambezian;

$p = 0.29$) and *Drimopsis* ($p = 0.84$), while Zambezian was reconstructed for *Drimiopsis* +

Ledebouria Clade B ($p = 0.95$) and *Ledebouria* Clade B ($p = 0.98$). Within *Ledebouria* Clade B

(i.e., *L. yemenensis* + *L. revoluta* Somalia), the model strongly favors a widespread distribution

of Yemen+Zambezian ($p = 0.55$) (the next best area being Zambezian; $p = 0.38$) followed by

subsequent reconstructions containing various combinations of Zambezian and other regions, or

Zambezian alone (Figures 3 and S2). Within *Ledebouria* Clade B we recover one of two

dispersals to Madagascar (Figure 3).

The Kalahari+Natal ($p = 0.42$) or Natal ($p = 0.35$) regions were recovered as the ancestral

range for *Resnova* + *Ledebouria* Clade A, whereas the Natal region was recovered for *Resnova*

($p = 0.99$), and Kalahari for *Ledebouria* Clade A ($p = 0.51$) (the next best area being Natal; $p =$

0.18) (Figures 3 and S2). Within *Ledebouria* Clade A, we find an additional dispersal to

Madagascar (Figure 3). Additionally, in summarizing the 10 biogeographic stochastic

reconstructions (see section 2.3.1), we recovered a steady rise in the number of lineages in the

Zambezian and Kalahari regions from ~15 myr onwards in a subclade of *Ledebouria* Clade A

(Figure 3b). Additionally, despite some differences in topology between these results and those

of Howard et al. (2022), both studies overall share clades that largely reflect geography. We

recover two subclades of mostly Namibian taxa, a subclade containing mostly South African

taxa, and a subclade containing a mixture of Zambian, Tanzanian, and Zimbabwean taxa

(Figures 2 and 3).

4 DISCUSSION

4.1 Broad biogeographical patterns

Here, we present the best and most comprehensive sampling of the Ledebouriinae to date

(representing ~30% of described Ledebouriinae taxa (POWO, 2019) plus numerous undescribed

taxa), which has provided greater insights into the historical biogeography of this widespread,

bulbous lineage than previously uncovered. Our results suggest a rapid radiation along the

backbone of the Ledebouriinae, estimated between ~28–26 mya (Figure 2; Table 1) in

southeastern sub-Saharan Africa (Figures 3, S1). This region corresponds somewhat to the

current center of diversity of the Ledebouriinae, which is at the intersection of the Natal-

Kalahari-Zambezian biogeographical regions (Lebatha, 2004; Manning, 2020; Venter, 1993).

The four major subclades (*Ledebouria* Clade A, *Ledebouria* Clade B, *Resnova*, and *Drimiopsis*)

originated in neighboring regions to one another soon thereafter (Table 1; Figure 3). During this

timeframe in Africa, major shifts in climate and habitat composition have been inferred at both

local and continental scales, and at shallow and deep phylogenetic scales (Couvreur et al., 2021;

Gizaw et al., 2021; Hagen et al., 2021; Kandziora et al., 2022; Pokorny et al., 2015). From the

Eocene-Oligocene transition, global cooling promoted the expansion of seasonal and arid

climates as well as savanna, grassland, and fire-prone habitats in Africa, with greater increases

during the Miocene (Senut et al., 2009; Sepulchre et al., 2006). Furthermore, the Great

Escarpment underwent renewed and sporadic uplift during the Miocene, which increased habitat

heterogeneity that was coupled with an intensification of global cooling and aridification that

collectively led to lineage radiation and dispersal in many African taxa (Cowling et al., 2009;

Galley et al., 2007; Neumann & Bamford, 2015; Partridge & Maud, 1987). Being bulbous

geophytes, which overall are found in some of the driest and most seasonal climates (Howard et

al., 2019), the Ledebouriinae may have been "pre-adapted" for these climatic changes, and

therefore diversified in response (Howard et al., 2022). Overall, our biogeographic

reconstructions and divergence time estimations lead us to hypothesize that the expansion of

seasonal habitats coupled with orogenic activity in southern sub-Saharan Africa spurred the

radiation of the Ledebouriinae and promoted additional radiations and dispersals over time.

4.2 A tale of two *Ledebouria*

We estimate that the two *Ledebouria* lineages originated at slightly different times

(Figure 2; Table 1) in neighboring regions (i.e., Kalahari vs Zambezian), which led to subsequent

divergent biogeographic histories (Figure 3). In general, the two *Ledebouria* overlap in eastern

Africa with a geographical divide between southern and northern sub-Saharan Africa (Figure 2)

(Howard et al., 2022). The divergent evolutionary histories and current overlapping distribution

of the two *Ledebouria* may have been influenced by the expansion and contraction of wet,

tropical, forested habitats in eastern Africa that repeatedly split a historically more widespread

distribution (Couvreur, 2015; Couvreur et al., 2021). These past habitat fluctuations have created

a mosaic of climates and vegetation types in eastern Africa, and have influenced the evolutionary

trajectory of many lineages (Dagallier et al., 2020; Lorenzen et al., 2012). Other lineages adapted

to seasonal and arid habitats share similar distributions with the two *Ledebouria* (Grace et al.,

2015; Jürgens, 1997). For example, the two lineages overlap geographically with that of

stapeliads (Ceropegieae, Apocynaceae)—*Ledebouria* Clade B overlaps with the Northern grade,

whereas *Ledebouria* Clade A overlaps with the Pan-African clade (see Fig. 3 in Bruyns et al.,

(2014)). We also find two dispersals of *Ledebouria* to Madagascar (Figure 3), which was

previously hypothesized (Pfosser, 2012). Madagascar and mainland Africa have been separated

since at least the Paleocene (Couvreur et al., 2021), and our results therefore suggests long

distance dispersal from Africa to Madagascar led to the origin of Ledebouriinae on the island.

Interestingly, both independent dispersal events are estimated to have occurred at similar times

(i.e., the confidence intervals overlap) (Figures 2, 3, S1), but this timing does not match with the

direction of ocean current flows that would have facilitated transport via rafts from Africa to

Madagascar (J. R. Ali & Huber, 2010). However, dispersal rates for animals were estimated to

not have changed due to ocean current flow direction, but instead may have increased after the

current shifted to an East to West flow direction (Samonds et al., 2012). Although completely

speculative, but interesting to think about nonetheless, perhaps an environmental phenomenon,

such as a historical major cyclone (Samonds et al., 2012), facilitated overwater dispersal of

Ledebouriinae from Africa to Madagascar, leading to the similar age estimates we recovered.

Although our sampling is significantly improved relative to studies thus far, taxa from other

geographic areas remain unsampled, and therefore our age and biogeographic estimates may

change with an even more comprehensive dataset.

4.2.1 The Voyage of *Ledebouria* Out of Africa, or *Ledebouria* Clade B

Reconstructions place the ancestral area of *Ledebouria* Clade B in the Zambezian region

followed by migration into other regions, such as Yemen and India, between ~20–15 mya

(Figures 2 and 3). Although our age estimates do not allow us to unequivocally favor vicariance

over dispersal from Africa to Eurasia, the current distribution of the Ledebouriinae, coupled with

our knowledge of their biology and the overall results from this study, do allow us to form

hypotheses of how the group may have migrated out of Africa. The Arabian Peninsula and

Socotra were connected or in close proximity with mainland Africa from at least ~17 mya

(Bottenberg, 2012; Culek, 2013; Edgell, 2006; Fleitmann et al., 2004; Jacobs, 2004; Pirouz et al.,

2017; Rögl, 1999), which would have allowed overland dispersal between these regions.

Whether arrival into the Arabian Peninsula occurred before the full formation of the Red Sea or

through the persistent connection offered by the Sinai Peninsula, will always be hard to assess

with any confidence. Later, migration into Eurasia may have been facilitated by the

Gomphotherium land bridge, which was present between ~19–13 mya (Bialik et al., 2019; Okay

et al., 2010; Rögl, 1999; Sen, 2013). The node reconstructed as India + Zambezian is estimated

at 15.6 myr (95% CI 16.3–13.1), and therefore, we hypothesize the land bridge may have

allowed dispersal of Ledebouriinae into Eurasia, a hypothesis also inferred for several animal

lineages (Harzhauser et al., 2007; Sen, 2013). Additionally, as the Socotra archipelago drifted

away from the mainland, we also hypothesize that vicariance likely promoted allopatric

speciation in the region, which is evidenced by the different species found on different islands

(Miller & Morris, 2004). Our age estimates, however, cannot fully rule out multiple dispersals to

the archipelago, but the many species found within the island chain led us to favor a vicariant

origin for these taxa. The vicariance hypotheses put forth here are not unique and fit a pattern

recovered in multiple studies. In other groups with widespread distributions between Africa, the

Arabian Peninsula, Socotra, and/or southern Asia, such as *Isodon* (Lamiaceae) (Yu et al., 2014),

Aganope (Fabaceae) (Sirichamorn et al., 2014), *Searsia* (Anacardiaceae) (Yang et al., 2016),

Smilax (Smilacaceae) (Chen et al., 2014), and several reptiles (Smíd et al., 2013; Tamar et al.,

2016), vicariance has been hypothesized or alluded to for explaining distributions and

biogeography, and many of these lineages have similar age estimates and distributions as

Ledebouria Clade B. Lastly, pollen and wood fossils indicate historical widespread distributions

between eastern Africa and south Asia of closely related taxa (Bonnefille, 2010; Morgan et al.,

1994), and fossils indicate that forested habitats were present in the Arabian Peninsula during the

Eocene and Oligocene, that later gave way to more open, xeric, grassland habitats in the Miocene

(Jacobs et al., 1999; Whybrow & McClure, 1980). The modern-day equivalents of these

conditions are commonly inhabited by Ledebouriinae (Lebatha, 2004; Venter, 1993). However,

given phylogenetic and divergence times uncertainty and limited taxon sampling from this

region, we cannot yet completely rule out multiple dispersals to the Arabian Peninsula and/or the

Socotra Archipelago. Regardless, in summary, our age estimates and biogeographical

reconstructions as well as the limited dispersal ability of Ledebouriinae, lead us to hypothesize

that *Ledebouria* Clade B dispersed out of Africa to Eurasia via overland migration. The

historical, widespread distribution was subsequently fragmented by Miocene-driven aridification.

4.2.2 Within Africa—*Ledebouria* Clade A

Our results suggest that *Ledebouria* Clade A originated slightly later and in a more

southerly region than that of *Ledebouria* Clade B (Table 1; Figures 2 and 3). Based on our

dataset, *Ledebouria* Clade A arose ~25–23 mya and remained predominantly within sub-

equatorial Africa (Figures 2 and 3). Despite the low resolution within some areas of Clade A

(Howard et al., 2022), we recovered multiple subclades that reflect geographic affinity (Figure

2). For example, we see a subclade of mixed geographic composition (i.e., *L. caesiomontana* + *L.*

sp. 12 CCH190) (node c, Figure 2) that suggests a widespread distribution between the Kalahari

and Zambezian regions that diverged and lead to *in situ* speciation in the Zambezian region

followed by a dispersal to Madagascar (Figure 3). We recovered an additional subclade of

predominantly Zambezian taxa (i.e., *L.* sp. 37 CCH169 + *L.* sp. 24 Uganda) (subclade in node a,

Figure 2) with evidence of a dispersal to the Sudanian region (i.e., *L.* sp. 24 Uganda).

Furthermore, two subclades of predominantly Namibian taxa (i.e., *L.* sp. 14 CCH109 + *L.* sp. 18

CCH218 (node b, Figure 2), and *L.* sp. 7 CCH066 + *L.* sp. 32 CCH145 (subclade in node a,

Figure 2) suggest multiple dispersals to the region followed by *in situ* speciation (Figure 3).

Lastly, a subclade of mostly South African taxa (i.e., *L. galpinii* + *L. coriacea*) (subclade in node

a, Figure 2) was recovered (Figure 3). Overall, our results suggest that within a relatively short

time frame (between ~20–15 mya), a complex history of dispersals followed by *in situ* speciation

events occurred within *Ledebouria* Clade A in southern Africa (Figure 3; Table S2).

From the Miocene onwards, complex interactions between mountain uplift and climate

change promoted diversification across lineages within southern Africa (García-Aloy et al.,

2017; Maswanganye et al., 2017; Neumann & Bamford, 2015; Nielsen et al., 2018). *Ledebouria*

Clade A may have originated in southeastern Africa then dispersed and diversified as the region

grew more heterogeneous and arid. For example, we found an instance of potentially rapid

radiation within *Ledebouria* Clade A between ~20–17 mya (Figure 3b), which corresponds to a

clade with consistently poor phylogenetic resolution (Howard et al., 2022). This subclade

contains taxa from across southern sub-Saharan Africa with geographic signals within (i.e.,

clades of South African, Namibian, and Zambian taxa) (Figures 2 and 3). It is possible that the

Middle Miocene Climatic Optimum or Middle Miocene Climate Transition (Zachos et al., 2001)

coupled with renewed uplift of the Great Escarpment spurred this radiation, and we see a steady

rise in lineage accumulation from ~20–15 mya (Figure 3b). During the Middle Miocene, the

vegetation of South Africa was likely tropical, which gave way to more xerophytic shrubland by

Late Miocene due to a global cooling climate (Pound et al., 2012). This time period is associated

with diversification in the Crassulaceae (Lu et al., 2022) and also the expansion of C4

grasslands/savannas at a global scale (Bobe, 2006; Senut et al., 2009). Current biogeographical

patterns of *Aloe* suggest dispersal from southern Africa to East Africa as seasonality and

aridification increased across the continent within the past ~16 myr (Grace et al., 2015).

Monsonia (Geraniaceae) share similar distributional patterns as *Ledebouria* Clade A with

multiple lineages comprised of mainly southwestern or southeastern taxa (i.e., southern African

split between eastern and western clades), which radiated within the past ~20–15 myr (García-

Aloy et al., 2017). Overall, we find a fascinating, albeit more complicated, evolutionary history

of *Ledebouria* Clade A within southern sub-Saharan Africa that remains to be fully resolved.

4.3 Limited, but intriguing insights into *Drimiopsis*

Based on our limited sampling of *Drimiopsis*, we find additional evidence for a southern

and northern sub-Saharan African divide (Figures 2 and 3), which is also seen in ungulates

(Lorenzen et al., 2012), *Lycaon pictus* (Canidae) (Marsden et al., 2012), *Encephalartos*

(Cycadaceae) (Mankga et al., 2020), and *Agama* (Reptilia) (Leaché et al., 2014). The

eastern/northern sub-Saharan African *Drimiopsis* arose ~19–17 mya (Figure 2), during a time of

increased mountain building in the region (~17 and ~13.5 mya; Wichura et al., 2015) that was

followed by a shift towards a more arid climate (Bonnefille, 2010; Linder, 2017; Morgan et al.,

1994; Senut et al., 2009). Several East African *Drimiopsis* have overall more succulent leaves

compared to their southern African relatives (Lebatha, 2004), which may have been selected for

as eastern Africa experienced greater aridification compared to the southeast. Additionally, East

African *Drimiopsis* are known polyploids (Lebatha, 2004; Stedje & Nordal, 1987), which may

have further spurred evolution in this region since polyploidy may provide an advantage in

particular environmental conditions (e.g., aridity) (Manzaneda et al., 2012; Sonnleitner et al.,

2010). In contrast, South African *Drimiopsis* occur in moist and shady habitats, and are not

found in the drier portions of southwestern Africa (Lebatha, 2004). Given the estimated age of

South African *Drimiopsis* (~13.6 mya) (Figure 2), uplift of the Great Escarpment may have

increased aridity in southwest Africa (Bobe, 2006; Partridge & Maud, 1987) that hindered

Drimiopsis establishment in this region, but at the same time promoted diversification in

southeastern Africa leading to its high diversity in the area (Lebatha, 2004). However, far greater

sampling of *Drimiopsis* from its entire distribution is needed to test hypotheses regarding their

evolutionary history. Ideally, an expanded phylogenetic framework would be coupled with

ploidal inferences, anatomical studies, and experimental manipulations to fully assess the

relationship between historical biogeography and morphological evolution.

4.4 Where to sample next?

Our taxon sampling heavily consisted of southern sub-Saharan African taxa (Figure 1),

and largely excluded *Resnova*. Therefore, increased sampling of *Resnova* as well as *Ledebouria*

and *Drimiopsis* from western, central, and northern Africa will provide further insights into the

biogeographical history of the Ledebouriinae. For example, *Ledebouria* occurs in northern

Angola (Gregory Jongsma, pers. obs.) and Gabon (Figure 1). This leads us to wonder whether a

continuous distribution was once present along the central/western Atlantic coast of Africa, or if

the Central African tropical forests have consistently obstructed Ledebouriinae dispersal within

the coastal, wet tropics. Furthermore, a *Ledebouria* accession from Uganda is nested within

Ledebouria Clade A, which suggests that by increasing sampling from northern/western Africa

we may uncover additional biogeographic events in this clade. Lastly, multiple, morphologically

distinct *Ledebouria* taxa are recorded from Socotra and surrounding islands (Miller &

Alexander, 2010; Miller & Morris, 2004), and India (Giranje & Nandikar, 2016). Incorporating

these into future phylogenetic studies may highlight further dispersals, or bolster support for

allopatric speciation/vicariance as a major evolutionary process leading to the current

distribution of the group outside of Africa.

5 CONCLUSION

The Ledebouriinae are a widespread, bulbous lineage of monocots that remain poorly

understood taxonomically due to a high degree of phenotypic plasticity and significant under

collection of specimens. Our study provides a pivotal first step towards refining our

understanding of historical evolution and biodiversity within this fascinating lineage. The results

of our analyses highlight the complex biogeographic history of the Ledebouriinae within and

outside of sub-Saharan Africa. We find that the Ledebouriinae evolved within the past ~30 myr,

which suggests the group radiated in response to increasing climatic seasonality and orogenic

activity since. Overall, given the evidence, we hypothesize that vicariance led to the current

distribution of *Ledebouria* in Eurasia, with overland dispersal out of Africa potentially occurring

via the *Gomphotherium* land bridge. However, long distance dispersal to Socotra and southern

Asia cannot be fully ruled out at this time. We also find evidence of two independent dispersals

to Madagascar. Within Africa, we can infer a northern and southern sub-Saharan division

between the two *Ledebouria* lineages and within *Drimiopsis*. We also uncover a complex history

of dispersals and *in situ* speciation events within southern Africa in *Ledebouria* Clade A, a clade

that warrants improved phylogenetic resolution to further elucidate the processes affecting its

current distribution. In conclusion, our study shows the value of increasing research focus on

understudied lineages inhabiting seasonal and arid habitats.

Tables

Table 1. Comparison of median age estimates and associated uncertainty (95% confidence
intervals (CI)) for each major Ledebouriinae clade using penalized likelihood as implemented in
treePL.

Clade	median estimate	95% CI
Scilloideae	52.2	51.2–54.5
Ledebouriinae	28.0	26.1–28.3
Ledebouria Clade A	25.0	23.0–25.4
Resnova	17.9	16.6–18.5
Clade A + *Resnova*	26.3	24.5–26.7

Drimiopsis	22.1	20.5–23.5
Ledebouria Clade B	26.1	24.0–26.5
Clade B + *Drimiopsis*	27.1	25.2–27.5

Table 2. Statistical outputs for each model from BioGeoBEARS. log-likelihood (LnL); rate of dispersal (*d*); rate of extinction (*e*); Akaike's Information Criterion (AIC)

Model	LnL	Number of parameters	*d*	*e*	AIC
DEC	-267.1	2	0.01	0.01	538.4
DIVA-like	**-220.5**	**2**	**0.0027**	**0.009**	**445.2**
BAYAREA-like	-232.8	2	0.0036	0.04	469.8

Figures

[double column] Figure 1. General distribution of the seasonal and arid ecosystems of sub-Saharan Africa (light orange polygon) underlain by the general distribution of the Ledebouriinae, which is sufficiently captured by displaying GBIF specimen occurrence data (teal squares; GBIF.org (22 March 2022) GBIF Occurrence Download https://doi.org/10.15468/dl.xmccae) and the location of samples collected for this study (maroon circles). Do note though that sampling

gaps exist, for example in Angola, Nigeria, and South Sudan. Examples of habitats and
associated Ledebouriinae taxa are displayed in the images. a) *Ledebouria* sp. 8 CCH186, b)
Ledebouria sp. 15 CCH027, c) *Drimiopsis botryoides* subsp. *botryoides* CCH153. See Figure 2
for phylogenetic placement of these example taxa. General distribution of seasonal landscapes
adapted from Balinsky (1962), Jürgens (1997), and Bellstedt et al. (2012).

[double column] Figure 2. Divergence times within the Ledebouriinae as estimated using
penalized likelihood in treePL. For each clade, collection localities are overlaid on a map of
Africa, Madagascar, and Asia to aid with discussions of geographical patterns. Clades labeled
with a, b, and c in yellow are clades discussed within the manuscript. Black circles indicate
nodes with ultrafast bootstrap support values below 90 as recovered from the IQ-TREE analysis.
Numbers above blue bars denote median age estimates. Blue bars denote 95% confidence

intervals. * indicates samples excluded from subsequent biogeographical analyses. Not all nodes
are annotated with estimated divergences times for illustrative simplicity. For a fully annotated
phylogeny, see Figure S1.

[double column] Figure 3. Biogeographical reconstruction using the DIVA-like model as
implemented in 'BioGeoBEARS' and plotted using 'RevGadgets'. Colored boxes at nodes
display the region reconstructed with the highest probability; colors used at nodes match those in
the upper left corner map (i.e., Fig. 3b). a) Regionalization scheme from Linder et al. (2012).
used in the classification of taxa for biogeographical analysis. b) Summary of 10 stochastic
mappings from a posterior distribution of trees including only taxa from the indicated clade
(solid black line). For pie chart probabilities at each node, see Figure S2.

Data Availability

Raw reads from PAFTOL are available in ENA (accession PRJEB35285) and assembled reads

from www.treeoflife.kew.org. Raw reads of the Ledebouriinae are available from the SRA

(accession PRJNA721471). Alignments, species trees, biogeography inputs/outputs are all

available on Dryad (FOR REVIEW ONLY:

https://datadryad.org/stash/share/ApGYMXXwCCd1G873IOkciaD27YdQyNqg0lpvKT38Xic).

Biosketch

Cody Coyotee Howard is an Assistant Professor at Oklahoma State University broadly interested

in geophyte evolution and biogeography, and he has a particular fascination with bulbous plants.

The Ledebouriinae represent one bulbous lineage that has captivated him for years, and these

plants will continue to do so for many more.

Author contributions: Conceptualization CCH, TSH, NC; Formal analysis, CCH and ARZ;

Investigation, CCH and ARZ; Resources, TSH, LN, DC, LN, PM, ARZ, NC; Writing —

Original Draft, CCH and NC; Writing — Review & Editing, all authors; Visualization, CCH;

Funding acquisition, CCH and NC.

Supporting information

Table S1. Fossil and secondary calibration points used in the penalized likelihood (treePL)

analysis. With the exception of monocots, all calibrations were set as minimum values in the

analysis.

node	minimum calibration	source
Zingiberaceae	72.1	Iles et al. (2015)
Typhaceae	51.66	Iles et al. (2015)
Cyperaceae	47.0	Iles et al. (2015)
Amaryllidaceae	48.88	Pigg et al. (2018)
Coryphoideae	83.6	Iles et al. (2015)
Agavoideae	14.5	Iles et al. (2015)
Xanthorrhoeaceae	38.0	Iles et al. (2015)
Goodyerinae	15.0	Iles et al. (2015)
Monocots	131–135	Magallón et al. (2015)

619   **Table S2**. Top ten most common transition types found in a subclade of *Ledebouria* Clade A

620   (Figure 3b) using stochastic mapping of biogeographic events.

starting_range	ending_range	average_n_events	SD	transition_type
K	K	23.432	2.121	in situ speciation
Z	Z	19.252	1.297	in situ speciation
K	KN	3.393	0.517	dispersal
T	T	2.764	2.089	in situ speciation
C	C	1.29	1.912	in situ speciation
KZ	K	1.09	0.307	vicariance or subset sympatry or extinction
KZ	Z	1.09	0.307	vicariance or subset sympatry or extinction
K	KZ	1.029	0.215	dispersal
CK	C	0.912	0.703	vicariance or subset sympatry or extinction

Figure S1. Full dated phylogeny with 95% confidence internals shown.

Figure S2. Biogeographical reconstruction using the DIVA-like model with nodes showing the

probable range for each region at that node.

References

Ali, J. R., & Huber, M. (2010). Mammalian biodiversity on Madagascar controlled by ocean currents.
Nature, *463*(7281), 653–656.

Ali, S. S., Pfosser, M., Wetschnig, W., Martínez-Azorín, M., Crespo, M. B., & Yu, Y. (2013). Out of
Africa: Miocene dispersal, vicariance, and extinction within Hyacinthaceae subfamily Urgineoideae.
Journal of Integrative Plant Biology, *55*(10), 950–964. https://doi.org/10.1111/jipb.12065

Ali, S. S., Yu, Y., Pfosser, M., & Wetschnig, W. (2012). Inferences of biogeographical histories within
subfamily Hyacinthoideae using S-DIVA and Bayesian binary MCMC analysis implemented in RASP
(Reconstruct Ancestral State in Phylogenies). *Annals of Botany*, *109*(1), 95–107.
https://doi.org/10.1093/aob/mcr274

Andermann, T., Cano, Á., Zizka, A., Bacon, C., & Antonelli, A. (2018). SECAPR-a bioinformatics
pipeline for the rapid and user-friendly processing of targeted enriched Illumina sequences, from raw
reads to alignments. *PeerJ*, *6*, e5175. https://doi.org/10.7717/peerj.5175

Baker, W. J., Bailey, P., Barber, V., Barker, A., Bellot, S., Bishop, D., Botigué, L. R., Brewer, G.,
Carruthers, T., Clarkson, J. J., Cook, J., Cowan, R. S., Dodsworth, S., Epitawalage, N., Françoso, E.,
Gallego, B., Johnson, M. G., Kim, J. T., Leempoel, K., Maurin, O., Mcginnie, C., Pokorny, L., Roy, S.,
Stone, M., Toledo, E., Wickett, N. J., Zuntini, A. R., Eiserhardy, W. L., Kersey, P. J., Leitch, I. J., Forest,
F. (2022). A comprehensive phylogenomic platform for exploring the angiosperm tree of life. *Systematic*
Biology, *71*(2), 301–319. https://doi.org/10.1093/sysbio/syab035

Balinsky, B. (1962). Patterns of animal distribution on the African continent. *Ann. Cape Prov. Mus*, *2*,
299–310.

Bellstedt, D. U., Galley, C., Pirie, M. D., & Linder, H. P. (2012). The migration of the palaeotropical arid
flora: Zygophylloideae as an example. *Systematic Botany*, *37*(4), 951–959.
https://doi.org/10.1600/036364412X656608

Bialik, O. M., Frank, M., Betzler, C., Zammit, R., & Waldmann, N. D. (2019). Two-step closure of the
Miocene Indian Ocean Gateway to the Mediterranean. *Scientific Reports*, *9*(1), 8842.

Bobe, R. (2006). The evolution of arid ecosystems in eastern Africa. *Journal of Arid Environments*, *66*(3),
564–584. https://doi.org/10.1016/j.jaridenv.2006.01.010

Bonnefille, R. (2010). Cenozoic vegetation, climate changes and hominid evolution in tropical Africa.
Global and Planetary Change, *72*(4), 390–411. https://doi.org/10.1016/j.gloplacha.2010.01.015

Bottenberg, H. C. (2012). *Plate Kinematics of the Afro-Arabian Rift System with Emphasis on the Afar*
Depression, Ethiopia. Missouri University of Science and Technology.

Bouckaert, R., Heled, J., Kühnert, D., Vaughan, T., Wu, C.-H., Xie, D., Suchard, M. A., Rambaut, A., &
Drummond, A. J. (2014). BEAST 2: A software platform for Bayesian evolutionary analysis. *PLoS*
Computational Biology, *10*(4), e1003537. https://doi.org/10.1371/journal.pcbi.1003537

Brown, J. W., Walker, J. F., & Smith, S. A. (2017). Phyx: Phylogenetic tools for unix. *Bioinformatics*,
33(12), 1886–1888. https://doi.org/10.1093/bioinformatics/btx063

Bruyns, P. V., Klak, C., & Hanáček, P. (2014). Evolution of the stapeliads (Apocynaceae-
Asclepiadoideae)—Repeated major radiation across Africa in an Old World group. *Molecular*
Phylogenetics and Evolution, *77*, 251–263. https://doi.org/10.1016/j.ympev.2014.03.022

Buerki, S., Jose, S., Yadav, S. R., Goldblatt, P., Manning, J. C., & Forest, F. (2012). Contrasting
biogeographic and diversification patterns in two Mediterranean-type ecosystems. *Plos One*, *7*(6),
e39377. https://doi.org/10.1371/journal.pone.0039377

Ceccarelli, F. S., Koch, N. M., Soto, E. M., Barone, M. L., Arnedo, M. A., & Ramírez, M. J. (2019). The
grass was greener: Repeated evolution of specialized morphologies and habitat shifts in ghost spiders
following grassland expansion in south america. *Systematic Biology*, *68*(1), 63–77.
https://doi.org/10.1093/sysbio/syy028

Coe, M. J., & Skinner, J. D. (1993). Connections, disjunctions and endemism in the eastern and southern
african mammal faunas. *Transactions of the Royal Society of South Africa*, *48*(2), 233–255.
https://doi.org/10.1080/00359199309520273

Couvreur, T. L. P. (2015). Odd man out: Why are there fewer plant species in African rain forests? *Plant*
Systematics and Evolution, *301*(5), 1299–1313. https://doi.org/10.1007/s00606-014-1180-z

Couvreur, T. L. P., Chatrou, L. W., Sosef, M. S. M., & Richardson, J. E. (2008). Molecular phylogenetics
reveal multiple tertiary vicariance origins of the African rain forest trees. *BMC Biology*, *6*, 54.
https://doi.org/10.1186/1741-7007-6-54

Couvreur, T. L. P., Dauby, G., Blach-Overgaard, A., Deblauwe, V., Dessein, S., Droissart, V., Hardy, O.
J., Harris, D. J., Janssens, S. B., Ley, A. C., Mackinder, B. A., Sonké, B., Sosef, M. S. M., Stévart, T.,
Svenning, J.-C., Wieringa, J. J., Faye, A., Missoup, A. D., Tolley, K. A., Nicolas, V., Ntie, S., Frederic,
F., Robin, C., Guillocheau, F., Barboni, D.,Sepulchre, P. (2021). Tectonics, climate and the
diversification of the tropical African terrestrial flora and fauna. *Biological Reviews of the Cambridge*
Philosophical Society, *96*(1), 16–51. https://doi.org/10.1111/brv.12644

Cowling, R. M., Procheş, S., & Partridge, T. C. (2009). Explaining the uniqueness of the Cape flora:
Incorporating geomorphic evolution as a factor for explaining its diversification. *Molecular Phylogenetics*
and Evolution, *51*(1), 64–74. https://doi.org/10.1016/j.ympev.2008.05.034

Culek, M. (2013). Geological and morphological evolution of the Socotra Archipelago (Yemen) from the
biogeographical view. *Journal of Landscape Ecology*, *6*(3), 84–108.

Dagallier, L.-P. M. J., Janssens, S. B., Dauby, G., Blach-Overgaard, A., Mackinder, B. A., Droissart, V.,
Svenning, J.-C., Sosef, M. S. M., Stévart, T., Harris, D. J., Sonké, B., Wieringa, J. J., Hardy, O. J., &
Couvreur, T. L. P. (2020). Cradles and museums of generic plant diversity across tropical Africa. *The*
New Phytologist, *225*(5), 2196–2213. https://doi.org/10.1111/nph.16293

del Hoyo, A., García-Marín, J. L., & Pedrola-Monfort, J. (2009). Temporal and spatial diversification of
the African disjunct genus Androcymbium (Colchicaceae). *Molecular Phylogenetics and Evolution*,
53(3), 848–861. https://doi.org/10.1016/j.ympev.2009.08.005

Drummond, A. J., & Rambaut, A. (2007). BEAST: Bayesian evolutionary analysis by sampling trees.
BMC Evolutionary Biology, *7*, 214. https://doi.org/10.1186/1471-2148-7-214

Edgell, H. (2006). *Arabian deserts: Nature, origin and evolution*. Springer Science & Business Media.
https://books.google.com/books?hl=en&lr=&id=zMSTz_zvhi8C&oi=fnd&pg=PR17&dq=arabian+desert
s+nature+origin+and+evolution&ots=Unlin___dv&sig=IQW2MZBEYaku9W2lIkmzywZf38A

Emberts, Z., St Mary, C. M., Howard, C. C., Forthman, M., Bateman, P. W., Somjee, U., Hwang, W. S.,
Li, D., Kimball, R. T., & Miller, C. W. (2020). The evolution of autotomy in leaf-footed bugs. *Evolution*,
74(5), 897–910. https://doi.org/10.1111/evo.13948

Esler, K., Rundel, P., & Vorster, P. (1999). Biogeography of prostrate-leaved geophytes in semi-arid
South Africa: Hypotheses on functionality. *Plant Ecology*, *142*(1), 105–120.

Fleitmann, D., Matter, A., Burns, S., Al-Subbary, A., & Al-Aowah, M. (2004). Geology and Quaternary
climate history of Socotra. *Fauna of Arabia*, *20*, 27–44.

Galley, C., Bytebier, B., Bellstedt, D. U., & Linder, H. P. (2007). The Cape element in the Afrotemperate
flora: From Cape to Cairo? *Proceedings. Biological Sciences / the Royal Society*, *274*(1609), 535–543.
https://doi.org/10.1098/rspb.2006.0046

García-Aloy, S., Sanmartín, I., Kadereit, G., Vitales, D., Millanes, A. M., Roquet, C., Vargas, P., Alarcón,
M., & Aldasoro, J. J. (2017). Opposite trends in the genus Monsonia (Geraniaceae): Specialization in the
African deserts and range expansions throughout eastern Africa. *Scientific Reports*, *7*(1), 9872.
https://doi.org/10.1038/s41598-017-09834-6

Giranje, P. T., & Nandikar, M. D. (2016). Synopsis of the genus Ledebouria Roth (Hyacinthaceae:
Hyacinthoideae) in India. *Webbia*, *71*(2), 213–217. https://doi.org/10.1080/00837792.2016.1182324

Givnish, T. J., Zuluaga, A., Spalink, D., Soto Gomez, M., Lam, V. K. Y., Saarela, J. M., Sass, C., Iles, W.
J. D., de Sousa, D. J. L., Leebens-Mack, J., Chris Pires, J., Zomlefer, W. B., Gandolfo, M. A., Davis, J. I.,
Stevenson, D. W., dePamphilis, C., Specht, C. D., Graham, S. W., Barrett, C. F., & Ané, C. (2018).
Monocot plastid phylogenomics, timeline, net rates of species diversification, the power of multi-gene
analyses, and a functional model for the origin of monocots. *American Journal of Botany*, *105*(11), 1888–
1910. https://doi.org/10.1002/ajb2.1178

Gizaw, A., Gorospe, J. M., Kandziora, M., Chala, D., Gustafsson, L., Zinaw, A., Salomón, L., Eilu, G.,
Brochmann, C., Kolář, F., & Schmickl, R. (2021). Afro-alpine flagships revisited II: elucidating the
evolutionary relationships and species boundaries in the giant senecios (Dendrosenecio, Asteraceae).
Alpine Botany. https://doi.org/10.1007/s00035-021-00268-5

Grace, O. M., Buerki, S., Symonds, M. R., Forest, F., van Wyk, A. E., Smith, G. F., Klopper, R. R.,
Bjorå, C. S., Neale, S., Demissew, S., Simmonds, M. S., & Rønsted, N. (2015). Evolutionary history and
leaf succulence as explanations for medicinal use in aloes and the global popularity of Aloe vera. *BMC
Evolutionary Biology*, *15*(1), 291. https://doi.org/10.1186/s12862-015-0291-7

Hagen, O., Skeels, A., Onstein, R. E., Jetz, W., & Pellissier, L. (2021). Earth history events shaped the
evolution of uneven biodiversity across tropical moist forests. *Proceedings of the National Academy of
Sciences of the United States of America*, *118*(40). https://doi.org/10.1073/pnas.2026347118

Harzhauser, M., Kroh, A., Mandic, O., Piller, W. E., Göhlich, U., Reuter, M., & Berning, B. (2007).
Biogeographic responses to geodynamics: A key study all around the Oligo–Miocene Tethyan Seaway.
Zoologischer Anzeiger-A Journal of Comparative Zoology, *246*(4), 241–256.

Howard, C. C. (2014). In the shade of the mopane: Ledebouria in Namibia. *Cactus and Succulent
Journal*, *86*(3), 92–98. https://doi.org/10.2985/015.086.0301

Howard, C. C., Crowl, A. A., Harvey, T. S., & Cellinese, N. (2022). Peeling back the layers: First
phylogenomic insights into the Ledebouriinae (Scilloideae, Asparagaceae). *Molecular Phylogenetics and
Evolution*, *169*, 107430. https://doi.org/10.1016/j.ympev.2022.107430

Howard, C. C., Folk, R. A., Beaulieu, J. M., & Cellinese, N. (2019). The monocotyledonous underground: Global climatic and phylogenetic patterns of geophyte diversity. *American Journal of Botany*, *106*(6), 850–863. https://doi.org/10.1002/ajb2.1289

Iles, W., Smith, S., Gandolfo, M., & Graham, S. (2015). Monocot fossils suitable for molecular dating analyses. *Botanical Journal of the Linnean Society*, *178*(3), 346–374.

Jacobs, B. F. (2004). Palaeobotanical studies from tropical Africa: Relevance to the evolution of forest, woodland and savannah biomes. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, *359*(1450), 1573–1583. https://doi.org/10.1098/rstb.2004.1533

Jacobs, B. F., Kingston, J. D., & Jacobs, L. L. (1999). The Origin of Grass-Dominated Ecosystems. *Annals of the Missouri Botanical Garden*, *86*(2), 590. https://doi.org/10.2307/2666186

Johnson, M. G., Gardner, E. M., Liu, Y., Medina, R., Goffinet, B., Shaw, A. J., Zerega, N. J. C., & Wickett, N. J. (2016). HybPiper: Extracting coding sequence and introns for phylogenetics from high-throughput sequencing reads using target enrichment. *Applications in Plant Sciences*, *4*(7). https://doi.org/10.3732/apps.1600016

Johnson, M. G., Pokorny, L., Dodsworth, S., Botigué, L. R., Cowan, R. S., Devault, A., Eiserhardt, W. L., Epitawalage, N., Forest, F., Kim, J. T., Leebens-Mack, J. H., Leitch, I. J., Maurin, O., Soltis, D. E., Soltis, P. S., Wong, G. K.-S., Baker, W. J., & Wickett, N. J. (2019). A universal probe set for targeted sequencing of 353 nuclear genes from any flowering plant designed using k-medoids clustering. *Systematic Biology*, *68*(4), 594–606. https://doi.org/10.1093/sysbio/syy086

Jürgens, N. (1997). Floristic biodiversity and history of African arid regions. *Biodiversity and Conservation*, *6*, 495–514. https://doi.org/10.1023/a:1018325026863

Kandziora, M., Gehrke, B., Popp, M., Gizaw, A., Brochmann, C., & Pirie, M. D. (2022). The enigmatic tropical alpine flora on the African sky islands is young, disturbed, and unsaturated. *Proceedings of the National Academy of Sciences of the United States of America*, *119*(22), e2112737119. https://doi.org/10.1073/pnas.2112737119

Katoh, K., & Standley, D. M. (2013). MAFFT multiple sequence alignment software version 7: Improvements in performance and usability. *Molecular Biology and Evolution*, *30*(4), 772–780. https://doi.org/10.1093/molbev/mst010

Kornas, J. (1985). Adaptive strategies of African pteridophytes to extreme environments. *Proceedings of the Royal Society of Edinburgh, Section B: Biological Sciences*, *86*, 391–396.

Lanfear, R., Calcott, B., Ho, S. Y. W., & Guindon, S. (2012). Partitionfinder: Combined selection of partitioning schemes and substitution models for phylogenetic analyses. *Molecular Biology and Evolution*, *29*(6), 1695–1701. https://doi.org/10.1093/molbev/mss020

Lanfear, R., Calcott, B., Kainer, D., Mayer, C., & Stamatakis, A. (2014). Selecting optimal partitioning schemes for phylogenomic datasets. *BMC Evolutionary Biology*, *14*, 82. https://doi.org/10.1186/1471-2148-14-82

Leaché, A. D., Wagner, P., Linkem, C. W., Böhme, W., Papenfuss, T. J., Chong, R. A., Lavin, B. R., Bauer, A. M., Nielsen, S. V., Greenbaum, E., Rödel, M.-O., Schmitz, A., LeBreton, M., Ineich, I., Chirio, L., Ofori-Boateng, C., Eniang, E. A., Baha El Din, S., Lemmon, A. R., & Burbrink, F. T. (2014). A hybrid phylogenetic-phylogenomic approach for species tree estimation in African Agama lizards with applications to biogeography, character evolution, and diversification. *Molecular Phylogenetics and Evolution*, *79*, 215–230. https://doi.org/10.1016/j.ympev.2014.06.013

Lebatha, P. (2004). *A systematic revision of Drimiopsis Lindl. & Paxt. (Hyacinthaceae)* [THESIS.DOCTORAL]. https://repository.nwu.ac.za/handle/10394/913

Li, H.-T., Yi, T.-S., Gao, L.-M., Ma, P.-F., Zhang, T., Yang, J.-B., Gitzendanner, M. A., Fritsch, P. W., Cai, J., Luo, Y., Wang, H., van der Bank, M., Zhang, S.-D., Wang, Q.-F., Wang, J., Zhang, Z.-R., Fu, C.-N., Yang, J., Hollingsworth, P. M., … Li, D.-Z. (2019). Origin of angiosperms and the puzzle of the Jurassic gap. *Nature Plants*, *5*(5), 461–470. https://doi.org/10.1038/s41477-019-0421-0

Linder, H. P. (2014). The evolution of African plant diversity. *Frontiers in Ecology and Evolution*, *2*, 1–14. https://doi.org/10.3389/fevo.2014.00038

Linder, H. P. (2017). East African Cenozoic vegetation history. *Evolutionary Anthropology*, *26*(6), 300–312. https://doi.org/10.1002/evan.21570

Linder, H. P., de Klerk, H. M., Born, J., Burgess, N. D., Fjeldså, J., & Rahbek, C. (2012). The partitioning of Africa: Statistically defined biogeographical regions in sub-Saharan Africa. *Journal of Biogeography*, *39*(7), 1189–1205. https://doi.org/10.1111/j.1365-2699.2012.02728.x

Lorenzen, E. D., Heller, R., & Siegismund, H. R. (2012). Comparative phylogeography of African savannah ungulates. *Molecular Ecology*, *21*(15), 3656–3670. https://doi.org/10.1111/j.1365-294X.2012.05650.x

Lu, M., Fradera-Soler, M., Forest, F., Barraclough, T. G., & Grace, O. M. (2022). Evidence linking life-form to a major shift in diversification rate in Crassula. *American Journal of Botany*, *109*(2), 272–290. https://doi.org/10.1002/ajb2.1797

Magalhaes, I. L. F., Santos, A. J., & Ramírez, M. J. (2021). Incorporating Topological and Age Uncertainty into Event-Based Biogeography of Sand Spiders Supports Paleo-Islands in Galapagos and Ancient Connections among Neotropical Dry Forests. *Diversity*, *13*(9), 418. https://doi.org/10.3390/d13090418

Magallón, S., Gómez-Acevedo, S., Sánchez-Reyes, L. L., & Hernández-Hernández, T. (2015). A metacalibrated time-tree documents the early rise of flowering plant phylogenetic diversity. *The New Phytologist*, *207*(2), 437–453. https://doi.org/10.1111/nph.13264

Mankga, L. T., Yessoufou, K., & Chitakira, M. (2020). On the origin and diversification history of the African genus Encephalartos. *South African Journal of Botany*, *130*, 231–239. https://doi.org/10.1016/j.sajb.2019.12.007

Manning, J. C. (2020). Systematics of Ledebouria sect. Resnova (Hyacinthaceae: Scilloideae: Massonieae), with a new subtribal classification of Massonieae. *South African Journal of Botany*, *133*, 98–110. https://doi.org/10.1016/j.sajb.2020.07.010

Manzaneda, A. J., Rey, P. J., Bastida, J. M., Weiss-Lehman, C., Raskin, E., & Mitchell-Olds, T. (2012). Environmental aridity is associated with cytotype segregation and polyploidy occurrence in Brachypodium distachyon (Poaceae). *The New Phytologist*, *193*(3), 797–805. https://doi.org/10.1111/j.1469-8137.2011.03988.x

Marsden, C. D., Woodroffe, R., Mills, M. G. L., McNutt, J. W., Creel, S., Groom, R., Emmanuel, M., Cleaveland, S., Kat, P., Rasmussen, G. S. A., Ginsberg, J., Lines, R., André, J.-M., Begg, C., Wayne, R. K., & Mable, B. K. (2012). Spatial and temporal patterns of neutral and adaptive genetic variation in the endangered African wild dog (Lycaon pictus). *Molecular Ecology*, *21*(6), 1379–1393. https://doi.org/10.1111/j.1365-294X.2012.05477.x

Maswanganye, K. A., Cunningham, M. J., Bennett, N. C., Chimimba, C. T., & Bloomer, P. (2017). Life
on the rocks: Multilocus phylogeography of rock hyrax (Procavia capensis) from southern Africa.
Molecular Phylogenetics and Evolution, *114*, 49–62. https://doi.org/10.1016/j.ympev.2017.04.006

Matasci, N., Hung, L.-H., Yan, Z., Carpenter, E. J., Wickett, N. J., Mirarab, S., Nguyen, N., Warnow, T.,
Ayyampalayam, S., Barker, M., Burleigh, J. G., Gitzendanner, M. A., Wafula, E., Der, J. P., dePamphilis,
C. W., Roure, B., Philippe, H., Ruhfel, B. R., Miles, N. W., Graham, S. W., Mathews, S., Surek B.,
Melkonian, M., Soltis, D. E., Soltis, P. S., Rothfels, C., Pokorny, L., Shaw, J. A., DeGironimo, L.,
Stevenson, D. W., Villarreal, J. C., Chen, T., Kutchan, T. M., Roly, M., Baucom, R. S., Deyholos, M. K.,
Samudrala, R., Tian, Z., Wu, X., Sun, X., Zhang, Y., Wang, J., Leebens-Mack, J., Wong, G. K.-S. (2014).
Data access for the 1,000 Plants (1KP) project. *GigaScience*, *3*, 17. https://doi.org/10.1186/2047-217X-3-
17

Matzke, N. J. (2013a). BioGeoBEARS: BioGeography with Bayesian (and likelihood) evolutionary
analysis in R Scripts. *R Package, Version 0.2*, *1*.

Matzke, N. J. (2013b). Probabilistic historical biogeography: New models for founder-event speciation,
imperfect detection, and fossils allow improved accuracy and model-testing. *Frontiers of Biogeography*,
5(4). https://doi.org/10.21425/F5FBG19694

Matzke, N. J. (2014). Model selection in historical biogeography reveals that founder-event speciation is a
crucial process in island clades. *Systematic Biology*, *63*(6), 951–970.
https://doi.org/10.1093/sysbio/syu056

Matzke, N. J. (2022). Statistical comparison ofDEC andDEC +J is identical to comparison of twoClaSSE
submodels, and is therefore valid. *Journal of Biogeography*. https://doi.org/10.1111/jbi.14346

Miller, A. G., & Alexander, D. (2010). Haemanthus grandifolius Balf.f.: An enigma resolved. *Edinburgh
Journal of Botany*, *53*(1), 43–47. https://doi.org/10.1017/S0960428600002717

Miller, A. G., & Morris, M. (2004). *Ethnoflora of the Soqotra Archipelago.* Royal Botanic Garden
Edinburgh.

Morgan, M. E., Kingston, J. D., & Marino, B. D. (1994). Carbon isotopic evidence for the emergence of
C4 plants in the Neogene from Pakistan and Kenya. *Nature*, *367*(6459), 162–165.
https://doi.org/10.1038/367162a0

Mwafongo, E., Vollsnes, A. V., Bjorå, C. S., Nordal, I., & Eriksen, A. B. (2017). Leaf
mottling/variegation and shape in the Ledebouria revoluta complex – Development, stability and putative
function. *Flora*, *236–237*, 33–43. https://doi.org/10.1016/j.flora.2017.09.009

Neumann, F. H., & Bamford, M. K. (2015). Shaping of modern southern African biomes: Neogene
vegetation and climate changes. *Transactions of the Royal Society of South Africa*, *70*(3), 195–212.
https://doi.org/10.1080/0035919X.2015.1072859

Nguyen, L.-T., Schmidt, H. A., von Haeseler, A., & Minh, B. Q. (2015). IQ-TREE: a fast and effective
stochastic algorithm for estimating maximum-likelihood phylogenies. *Molecular Biology and Evolution*,
32(1), 268–274. https://doi.org/10.1093/molbev/msu300

Nielsen, S. V., Daniels, S. R., Conradie, W., Heinicke, M. P., & Noonan, B. P. (2018). Multilocus
phylogenetics in a widespread African anuran lineage (Brevicipitidae: Breviceps) reveals patterns of
diversity reflecting geoclimatic change. *Journal of Biogeography*, *45*(9), 2067–2079.
https://doi.org/10.1111/jbi.13394

Nylinder, S., Razafimandimbison, S. G., & Anderberg, A. A. (2016). From the Namib around the world:
Biogeography of the Inuleae-Plucheinae (Asteraceae). *Journal of Biogeography*, *43*(9), 1705–1716.
https://doi.org/10.1111/jbi.12764

Okay, A. I., Zattin, M., & Cavazza, W. (2010). Apatite fission-track data for the Miocene Arabia-Eurasia
collision. *Geology*, *38*(1), 35–38.

Partridge, T., & Maud, R. (1987). Geomorphic evolution of southern Africa since the Mesozoic. *South*
African Journal of Geology, *90*(2), 179–208.

Pausas, J. G., Lamont, B. B., Paula, S., Appezzato-da-Glória, B., & Fidelis, A. (2018). Unearthing
belowground bud banks in fire-prone ecosystems. *The New Phytologist*, *217*(4), 1435–1448.
https://doi.org/10.1111/nph.14982

Pfosser, M. (2012). Phylogenetic relationships of Malagasy Hyacinthaceae. *Plant Ecology and Evolution*,
145(1), 65–72. https://doi.org/10.5091/plecevo.2012.590

Pillon, Y., Hopkins, H. C. F., Maurin, O., Epitawalage, N., Bradford, J., Rogers, Z. S., Baker, W. J., &
Forest, F. (2021). Phylogenomics and biogeography of Cunoniaceae (Oxalidales) with complete generic
sampling and taxonomic realignments. *American Journal of Botany*, *108*(7), 1181–1200.
https://doi.org/10.1002/ajb2.1688

Pirouz, M., Avouac, J.-P., Hassanzadeh, J., Kirschvink, J. L., & Bahroudi, A. (2017). Early Neogene
foreland of the Zagros, implications for the initial closure of the Neo-Tethys and kinematics of crustal
shortening. *Earth and Planetary Science Letters*, *477*, 168–182.

Pokorny, L., Riina, R., Mairal, M., Meseguer, A. S., Culshaw, V., Cendoya, J., Serrano, M., Carbajal, R.,
Ortiz, S., Heuertz, M., & Sanmartín, I. (2015). Living on the edge: Timing of Rand Flora disjunctions
congruent with ongoing aridification in Africa. *Frontiers in Genetics*, *6*, 154.
https://doi.org/10.3389/fgene.2015.00154

Pound, M. J., Haywood, A. M., Salzmann, U., & Riding, J. B. (2012). Global vegetation dynamics and
latitudinal temperature gradients during the Mid to Late Miocene (15.97–5.33Ma). *Earth-Science*
Reviews, *112*(1–2), 1–22. https://doi.org/10.1016/j.earscirev.2012.02.005

POWO. (2019). *Plants of the world online. Facilitated by the Royal Botanic Gardens, Kew.*
http://www.plantsoftheworldonline.org/

Procheş, Ş., Cowling, R., Goldblatt, P., Manning, J., & DA Snijman. (2006). An overview of the Cape
geophytes. *Biological Journal of the Linnean Society*, *87*(1), 27–43.

R Core Team. (2016). *R: A language and environment for statistical computing.* (3.3.2). R Foundation for
Statistical Computing. https://www.R-project.org/

Rambaut, A., Drummond, A. J., Xie, D., Baele, G., & Suchard, M. A. (2018). Posterior summarization in
Bayesian phylogenetics using Tracer 1.7. *Systematic Biology*, *67*(5), 901–904.
https://doi.org/10.1093/sysbio/syy032

Ree, R. H., & Sanmartín, I. (2018). Conceptual and statistical problems with the DEC+J model of
founder-event speciation and its comparison with DEC via model selection. *Journal of Biogeography*,
45(4), 741–749. https://doi.org/10.1111/jbi.13173

Rögl, F. (1999). Mediterranean and Paratethys. Facts and hypotheses of an Oligocene to Miocene
paleogeography (short overview). *Geologica Carpathica*, *50*(4), 339–349.

Samonds, K. E., Godfrey, L. R., Ali, J. R., Goodman, S. M., Vences, M., Sutherland, M. R., Irwin, M. T.,
& Krause, D. W. (2012). Spatial and temporal arrival patterns of Madagascar's vertebrate fauna explained
by distance, ocean currents, and ancestor type. *Proceedings of the National Academy of Sciences*,
109(14), 5352–5357.

Sanmartín, I., Anderson, C. L., Alarcon, M., Ronquist, F., & Aldasoro, J. J. (2010). Bayesian island
biogeography in a continental setting: The Rand Flora case. *Biology Letters*, *6*(5), 703–707.
https://doi.org/10.1098/rsbl.2010.0095

Sen, S. (2013). Dispersal of African mammals in Eurasia during the Cenozoic: Ways and whys. *Geobios*,
46(1–2), 159–172.

Senut, B., Pickford, M., & Ségalen, L. (2009). Neogene desertification of Africa. *Comptes Rendus*
Geoscience, *341*(8), 591–602.

Sepulchre, P., Ramstein, G., Fluteau, F., Schuster, M., Tiercelin, J.-J., & Brunet, M. (2006). Tectonic
uplift and Eastern Africa aridification. *Science*, *313*(5792), 1419–1423.
https://doi.org/10.1126/science.1129158

Smith, S. A., Brown, J. W., & Walker, J. F. (2018). So many genes, so little time: A practical approach to
divergence-time estimation in the genomic era. *Plos One*, *13*(5), e0197433.
https://doi.org/10.1371/journal.pone.0197433

Smith, S. A., & O'Meara, B. C. (2012). TreePL: divergence time estimation using penalized likelihood
for large phylogenies. *Bioinformatics*, *28*(20), 2689–2690. https://doi.org/10.1093/bioinformatics/bts492

Sonnleitner, M., Flatscher, R., Escobar García, P., Rauchová, J., Suda, J., Schneeweiss, G. M., Hülber, K.,
& Schönswetter, P. (2010). Distribution and habitat segregation on different spatial scales among diploid,
tetraploid and hexaploid cytotypes of Senecio carniolicus (Asteraceae) in the Eastern Alps. *Annals of*
Botany, *106*(6), 967–977. https://doi.org/10.1093/aob/mcq192

Speta, F. (1998a). Hyacinthaceae. *Flowering Plants: Monocotyledons*, 261–285.

Speta, F. (1998b). Systematische analyse der gattung Scilla L. sl (Hyacinthaceae). *Phyton*.
http://agris.fao.org/agris-search/search.do?recordID=US201301793397

Stamatakis, A. (2014). RAxML version 8: A tool for phylogenetic analysis and post-analysis of large
phylogenies. *Bioinformatics*, *30*(9), 1312–1313. https://doi.org/10.1093/bioinformatics/btu033

Stedje, B. (1998). Phylogenetic relationships and generic delimitation of sub-SaharanScilla
(Hyacinthaceae) and allied African genera as inferred from morphological and DNA sequence data. *Plant*
Systematics and Evolution, *211*(1–2), 1–11. https://doi.org/10.1007/BF00984908

Stedje, B., & Nordal, I. (1987). Cytogeographical studies of Hyacinthaceae in Africa south of the Sahara.
Nordic Journal of Botany, *7*(1), 53–65.

Tribble, C. M., Martínez-Gómez, J., Howard, C. C., Males, J., Sosa, V., Sessa, E. B., Cellinese, N., &
Specht, C. D. (2021). Get the shovel: Morphological and evolutionary complexities of belowground
organs in geophytes. *American Journal of Botany*, *108*(3), 372–387. https://doi.org/10.1002/ajb2.1623

Venter, S. (1993). *A revision of the genus Ledebouria Roth (Hyacinthaceae) in South Africa.*
[THESIS.DOCTORAL]. http://nrfnexus.nrf.ac.za/handle/20.500.11892/82782

Venter, S. (2008). Synopsis of the genus Ledebouria Roth (Hyacinthaceae) in South Africa. *Herbertia*,
62, 85–155.

Whybrow, P. J., & McClure, H. A. (1980). Fossil mangrove roots and palaeoenvironments of the miocene
of the eastern Arabian Peninsula. *Palaeogeography, Palaeoclimatology, Palaeoecology*, *32*, 213–225.
https://doi.org/10.1016/0031-0182(80)90041-3

Wichura, H., Jacobs, L. L., Lin, A., Polcyn, M. J., Manthi, F. K., Winkler, D. A., Strecker, M. R., &
Clemens, M. (2015). A 17-My-old whale constrains onset of uplift and climate change in east Africa.
Proceedings of the National Academy of Sciences of the United States of America, *112*(13), 3910–3915.
https://doi.org/10.1073/pnas.1421502112

Zachos, J., Pagani, M., Sloan, L., Thomas, E., & Billups, K. (2001). Trends, rhythms, and aberrations in
global climate 65 Ma to present. *Science*, *292*(5517), 686–693. https://doi.org/10.1126/science.1059412

Zhou, L., Su, Y. C. F., Thomas, D. C., & Saunders, R. M. K. (2011). 'Out-of-Africa' dispersal of tropical
floras during the Miocene climatic optimum: Evidence from Uvaria (Annonaceae). *Journal of*
Biogeography, *39*(2), 322–335. https://doi.org/10.1111/j.1365-2699.2011.02598.x