Title: Mouse Memory CD8 T cell subsets defined by Tissue Resident Memory (T_{RM}) Integrin

Expression Exhibit Distinct Metabolic Profiles

Running title: CD8 T cell subsets exhibit distinct metabolic profiles

Mike Sportiello[1,2], Alexis Poindexter[1], Emma C. Reilly[1], Adam Geber[1,2], Kris Lambert Emo[1],

Taylor N. Jones[1], David J. Topham[1]

[1]Center for Vaccine Biology and Immunology, University of Rochester Medical Center, Rochester,

NY, United States

[2] Medical Scientist Training Program, University of Rochester Medical Center, Rochester, NY

14642, USA

* Correspondence:

David J. Topham, PhD

David_Topham@urmc.rochester.edu

Phone: (585) 273-1403

Fax: (585) 273-2452

Abstract

Tissue-resident memory CD8 T cells (T_{RM}) principally reside in peripheral non-lymphoid tissues

such as lung and skin and confer protection against a variety of illnesses ranging from infections to

cancers. The functions of different memory CD8 T cell subsets have been linked with distinct

metabolic pathways and differ from other CD8 T cell subsets. For example, skin-derived memory T

cells undergo fatty acid oxidation and oxidative phosphorylation to a greater degree than circulating

memory and naïve cells. Lung T_{RM} cells defined by the cell surface expression of integrins exist as

distinct subsets that differ in gene expression and function. We hypothesize that T_{RM} subsets with

different integrin profiles will utilize unique metabolic programs. To test this, differential expression

and pathway analysis were conducted on RNAseq datasets from mouse lung T_{RM} yielding significant

differences related to metabolism. Next, metabolic models were constructed and the predictions were

interrogated using functional metabolite uptake assays. The levels of oxidative phosphorylation,

mitochondrial mass, and neutral lipids were measured. Furthermore, to investigate the potential

relationships to T_{RM} development, T-cell differentiation studies were conducted *in vitro* with varying

concentrations of metabolites. These demonstrated that lipid conditions impact T cell survival, and

that glucose concentration impacts the expression of canonical T_{RM} marker CD49a, with no effect on

central memory-like T-cell marker CCR7. In summary, it is demonstrated that mouse resident

memory T cells subsets defined by integrin-expression in the lung have unique metabolic profiles and

that nutrient abundance can alter differentiation.

1   Introduction

Tissue-resident memory T cells (T_{RM}) and tumor-infiltrating lymphocytes (TIL) with a T_{RM}-like phenotype (CD103+ CD49a+) confer protection against a variety of illnesses ranging from infectious diseases to cancers. As demonstrated in the skin, transfer of T_{RM} from immune to naïve hosts is sufficient to prevent herpes simplex virus type 1 (HSV-1)-associated pathology and, comparably, the presence of antigen-specific CD49a+ CD103+ TIL in a mouse model of melanoma can limit tumor growth (1, 2). In the lungs, T_{RM} are critical for protection against SARS-CoV-2 (COVID-19), influenza, and respiratory syncytial virus, even in the absence of antibodies specific for those viruses (3, 4, 5, 6). T_{RM} have been shown to act as both antigen-specific sentinels linking the innate and adaptive immune systems as well as cytotoxic mediators (7). This functional capacity makes them important players in immunity across multiple organ systems, particularly barrier tissues where nutrient substrate availability varies, and disease states.

T_{RM} are heterogeneous in terms of integrin expression, migration, effector potential, and gene expression (7). Furthermore, the functions of different memory CD8 T cell subsets have been linked with distinct metabolic pathways. For example, skin T_{RM} undergo fatty acid oxidation to a greater degree than other circulating memory and naive cells which tend to have higher glycolytic fluxes (8, 9). However, it is unclear if this programming is universal among all memory T-cell subsets and whether it holds true in other tissues such as the lung. It is also not yet known whether T_{RM} CD8 T cell subsets that express different combinations of surface integrins CD49a and CD103 share metabolic profiles. RNAseq on these populations revealed distinct metabolic transcriptomes (7), suggesting the overall metabolic profiles may differ, which was supported by ex vivo assays. *In vitro* studies to drive differentiation into T_{RM}-like phenotypes were used to test whether manipulating nutrient availability affected integrin-based subset development.

2 Materials and Methods

2.1 RNA sequencing analysis

RNA sequencing data and lists of differentially expressed genes were obtained from previously published work from our lab (GSE179653). Full code that generated those differentially expressed genes as well as further analyses and figures is publicly available at https://github.com/sportiellomike/Lung-CD8-T-cell-Immunometabolism .

2.2 Metabolic modelling

Mouse metabolic model iMM1865 was used as a starting point, onto which our publicly accessible transcriptomic data for CD49a^{pos}CD103^{pos} (Double Positive, DP), CD49a^{pos}CD103^{neg} (CD49a Single Positive, CD49a SP), or CD49a^{neg}CD103^{neg} (Double Negative, DN) was applied using the Gene Inactivity Moderated by Metabolism and Expression (GIMME algorithm) (10). A threshold value of the median transcript per kilobase million (TPM) for the 1,865 genes in the metabolic model was used. Reaction constraints were added according to previously published research (11). All code used to generate input data, figures, and the models themselves is available at https://github.com/sportiellomike/Lung-CD8-T-cell-Immunometabolism.

2.3 Mice

Mice were housed in approved microisolator cages, within a pathogen-free vivarium facility that is accredited by the Association for Assessment and Accreditation of Laboratory Animal Care. The vivarium is staffed with a number of personnel including those with expertise in husbandry, technical skills, and veterinary personnel. C57BL/6J mice (Jackson Laboratories) used for experiments were infected 8-10 weeks after birth. This study was carried out in strict accordance with the recommendations in the *Guide for the Care and Use of Laboratory Animals* as defined by the NIH.

81   The Institutional Animal Care and Use Committee of the University of Rochester approved all

82   protocols.

2.4   Virus and infection

84   Mice were anesthetized with 3,3,3-tribromoethanol. Upon verification of sedation, mice were

85   infected intranasally with 10^5 EID_{50} of HKx31 human influenza A virus in 30uL volume. After

86   observation to ensure they recovered from anesthesia, mice were monitored daily for weight and

87   overall morbidity to comply with all regulations.

2.5   Lung and Bronchoalveolar lavage processing

Bronchoalveolar lavage (BAL) was collected by inserting a flexible Teflon catheter into a slit in the

trachea in proximity to the larynx. A 1mL syringe was connected to the catheter and the lungs were

flushed 3 times with 1x PBS (volumes of 1mL, 900uL, and 800uL). BAL was spun down at 424xg

for 10 minutes. Red blood cells were eliminated through ACK mediated lysis. Cells were washed in

1x PBS containing 1% FBS (PBS+serum) and resuspended in PBS+serum.

The lungs were harvested and separated into right and left lobes prior to collection into RPMI + 8%

FBS (Corning #10040CV). Lungs were transferred into Miltenyi Gentle MACS tubes containing 2

mL of 2mg/mL collagenase II (Worthington #LS004177) and mechanically ground up by a Gentle

MACS machine. 3 additional mL of 2 mg/mL collagenase II solution was added before incubation

for 30 min at 37° C. Further mechanical digestion by Gentle MACS was done at this point. Tubes

were spun at 233xg at room temperature for 2 minutes.

Pellets were then dissolved in existing supernatant and transferred through 100 µm strainers.

Contents were quantitatively transferred using 5 mL of RPMI + 8% FBS. Tubes were spun at 424xg

for 6 minutes at room temperature. The supernatant was discarded, and dissolved in 4 mL 40%

Percoll solution, which was then layered on 3 mL 75% Percoll solution. These tubes were then spun

at 931xg for 20 minutes at room temperature with no centrifuge break so as not to disturb the layers

of the gradient.

The top layer of lipid-rich liquid was aspirated off, and the interface (approximately 2 mL) of the two

Percoll densities was collected where the cells were. Tubes were filled with RPMI + 8% FBS and

spun down at 424xg for 6 minutes at room temperature.

2.6   Spleen processing to achieve single-cell suspension

C57BL/6J mouse spleens were obtained from sacrificed mice and mechanically digested by grinding

between two frosted microscope slides without exogenous enzyme added. After grinding, Single-cell

suspensions were passed through a plastic mesh to capture large non-single cellular material and

centrifuged in conical vials at 424xg at 4°C for 6 minutes. Cells were resuspended in 3 mL of ACK

lysis buffer (Cold Spring Harbor Protocol: doi:10.1101/pdb.rec083295) for 5 minutes at room

temperature. Volume was brought up to 15 mL in PBS + serum followed by centrifuging at 424xg at

4°C for 6 minutes.

2.7 T_{RM}-like Media preparation

The following were added to 500 mL DMEM (Corning Catalog No. 10-017-CV): 50 mL fetal bovine

serum, 5 mL L-glutamine (Fisher Scientific Catalog No. 25030081), 12.5 mL HEPES buffer (Fisher

Scientific Catalog No. 15630080), 6.5 MEM Non-Essential Amino Acids Solution (Fisher Scientific

11140050), 0.6 mL of 1:300 dilution of beta-mercaptoethanol in PBS, 5 mL penicillin-streptomycin

(Fisher Scientific Catalog No. 15140122), 0.6 mL gentamycin (Fisher Scientific Catalog No.

15710064). Contents were sterile-filtered.

2.8   Fatty Acid Uptake

Fatty Acid Uptake Media (FUM) was created by dissolving 2 µL stock solution made according to

package insert per 1 mL of Assay Buffer (AAT Bioquest Screen Quest Fluorimetric Fatty Acid

Uptake Assay Kit #36385). Once a single cell suspension was obtained from the harvesting protocol,

cells were plated in 100 µL of FUM and 100 µL Leibowitz media for a final volume of 200 µL.

Vehicle control used equivalent volume of dimethyl sulfoxide (DMSO) in Assay Buffer. Cells were

incubated for 30 minutes at 37° C, then washed 1.5 times with PBS + 1% serum before staining.

2.9   Neutral Lipid Staining

Bodipy 505/515 (Fisher Scientific Catalog No. D3921) was dissolved in DMSO to yield a stock of 1

mg/µL concentration. Once a single cell suspension was obtained from the harvesting protocol, each

sample of cells in 1 mL of PBS were inoculated with Bodipy to yield a final concentration of 2

mg/mL, or DMSO for the vehicle control. Samples were incubated for 30 minutes at 37° C, then

washed 1.5 times with PBS + 1% serum before staining.

2.10   2-(N-(7-Nitrobenz-2-oxa-1,3-diazol-4-yl)Amino)-2-Deoxyglucose Uptake

2-(N-(7-Nitrobenz-2-oxa-1,3-diazol-4-yl)Amino)-2-Deoxyglucose (2-NBDG) (Abcam Catalog No.

ab146200) was dissolved in DMSO to yield a 20 mM stock solution. After a single cell suspension

was obtained from the harvesting protocol, cells were resuspended in 500 µL non-supplemented

Leibowitz media and incubated at 37° C with a final 2-NBDG concentration of 80 µM or DMSO for

the vehicle control for 35 minutes at 37° C. Samples were washed 1.5 times with PBS + 1% serum

before staining.

2.11   Mitochondrial Mass Quantification

Mitotracker Green FM (Mitotracker) (Fisher Scientific Catalog No. M7514) was dissolved in DMSO

to obtain a 1 mM stock solution. After a single cell suspension was obtained from the harvesting

protocol, each sample was dissolved in 500 µL PBS + 1% serum. Samples were inoculated with

Mitotracker for a final concentration of 50 nM or equivalent volume of DMSO for the vehicle control

and incubated for 45 minutes at 37° C.

2.12 Oxidative Phosphorylation Quantification

Tetramethylrhodamine, Ethyl Ester, Perchlorate (TMRE) (Cayman Chemical Catalog No. 701310)

was prepared in a 0.5 mM working stock by dissolving in DMSO, according to package insert. After

a single cell suspension was obtained from the harvesting protocol, TMRE was added to two separate

aliquots of 5 mL RPMI + 8% FBS (TMRE Media) for a final concentration of 250 nM. To one of

these aliquots, 2-[2-[4-(trifluoromethoxy)phenyl]hydrazinylidene]-propanedinitrile (FCCP) was

added (FCCP-TMRE Media) for a final concentration of 1 µM. Samples were split into two and

dissolved into 500 µL of either TMRE Media or FCCP-TMRE Media. Samples were incubated at

37° C for 30 minutes. For vehicle control, DMSO was added in place of TMRE. Cells were washed

1.5 times with PBS + 1% FBS prior to staining.

2.13 T$_{RM}$-like cell culture conditions

24 well plates were coated with anti-CD3/28 stimulating antibodies (Biolegend Catalog Nos. 100302

and 102102) at final concentration of 5 µg/mL each overnight at 4° C. The next day, 2 million single

cell splenocytes were plated in 1 mL T$_{RM}$-like culture media onto each well with 7.5 ng/mL human

IL-2 (Obtained from the NIH). Two days later, cells were collected, centrifuged at 424xg at 4° C for

6 minutes, and plated onto non-coated 24 well plates in 1 mL TRM-like culture media with 7.5

ng/mL IL-2 for three days. After that, cells were collected, centrifuged at 424xg at 4° C for 6

minutes, and plated in TRM-like culture media with 10 ng/mL mouse IL-7 (Biolegend Catalog No.

577804) and 10 ng/mL human TGFβ1 (Fisher Scientific PHG9214) for 6 days.

Additional glucose (Fisher Scientific Catalog No. A2494001) and lipid supplement (Millipore Sigma

Catalog No. L5146) were added to TRM-like cell culture media starting from the day of spleen

harvest (day 0) throughout the end of the experiment (day 11, day 6 post TGFβ1 addition). Soluble

lipid stores were created in Kolliphor P 188 (Millipore Sigma K4894) according to lipid supplement

instructions.

2.14 Staining for Flow Cytometry

Cells were suspended in 100 µL staining buffer for 30 minutes at 4° C. Staining buffer was created

by adding staining antibodies to PBS below for a final concentration of 1 µg/mL each, or 0.2 µL of

live-dead stock solution/100 µL staining buffer. Live-dead stock solution was prepared according to

package insert available on the vendor website. αCD16/32 was added for a final concentration of 2.5

µg/mL. αCD45 was previously injected at time of organ harvest intravenously at a concentration of

0.2 µg stock antibody/100 µL PBS in a volume of 100 µL. See table S1 for more information. All

antibodies used Bangs Laboratories compensation beads (Bangs #556) which were stained an

equivalent time. Live-dead stain used Life Technologies ArC reactive beads (Invitrogen A10346).

Because the vast majority of cells at day 14 post infection will have been antigen experienced, CD44

was not used as a marker on day 14 experiments. Experiments were run on an 18-color BD LSR II

flow cytometer. Gating strategies are displayed in in the supplementary file.

2.15 MFI Determination and Normalization

The four subsets of interest were manually gated after selecting for lymphocytes (FSC-A and SSC-

A), singlets (FSC-A and FSC-H), live cells, IV-CD45^{neg} cells, TCRβ^{high} and CD8α^{high} cells. DN

(Double Negative) was defined as CD103^{low} and CD49a^{low}, DP (Double Positive) as CD103^{high} and

CD49a^{high}, CD103 SP (CD103 Single Positive) as CD103^{high} CD49a^{low}, and CD49a SP (CD49a

Single Positive) as CD49a^{high} and CD103^{low}.

Median fluorescence intensities (MFIs) were calculated using Flowjo (version 10.6.2). Data was

normalized by dividing the MFI of each subset by the sum of all four subsets' MFIs for each mouse

and multiplying by 4. For the assays using TMRE, FCCP was subtracted from the TMRE MFI for

each subset within each mouse. That number was divided by the sum of MFIs of the uncoupled

control (FCCP+TMRE) subsets for each mouse. Normalized data can be found at

https://github.com/sportiellomike.

2.16 Statistical Tests

After repeated measures ANOVA, Multiple pairwise comparisons were evaluated using the Student's

T-test with Benjamini-Hochberg correction. If $P_{adjusted}<0.05$, results were deemed significant. R

Packages ggplot2 (version 3.3.2) and rstatix (0.7.0) were used for data plotting and statistical testing

respectively. The code used to perform statistical tests and generate figures can be found here:

https://github.com/sportiellomike.

2.17 Ethics statement

This study was carried out in strict accordance with the recommendations in the Guide for the Care

and Use of Laboratory Animals as defined by the NIH. Animal protocols were reviewed and

approved by the Institutional Animal Care and Use Committee of the University of Rochester in

writing. All animals were housed in a centralized and Association for Assessment and Accreditation

of Laboratory Animal Care accredited research animal facility that is fully staffed with trained

husbandry, technical, and veterinary personnel. Our UCAR number is 2006-029.

Euthanasia was performed by animal welfare committee reviewed approved methods.

3 **Results**

3.1 *Pathway analysis for metabolic analyses.*

T_{RM} can be defined by their surface protein expression, with multiple markers having been used

including CD69, CD49a, and CD103. In the mouse lung, CD44 expression identifies antigen-

experienced memory T cells while CD49a marks T_{RM}, and when used in combination with CD103,

yields two T_{RM} subsets: CD49a^{pos}CD103^{pos} (Double Positive, DP) and CD49a^{pos}CD103^{neg} (CD49a

Single Positive, CD49a SP) (7). At later memory time points, very few CD49a^{neg}CD103^{pos} (CD103

Single Positive, CD103 SP) cells remain, suggesting they have either died, left the tissue, or altered

their integrin expression. CD49a^{neg}CD103^{neg} (Double Negative DN) cells represent mixed

phenotypes, though more than half are CD44^{high} CD62L^{low} T cells consistent with an Effector

Memory profile (T_{EM}) and may be circulating (7). To more fully describe the metabolic profiles of

T_{RM} based on integrin expression, we utilized the publicly accessible RNA sequencing data set we

previously reported (7). This data set was generated from CD44^{high} mouse lung tissue CD8 T cells 21

days post-infection (dpi) that were sorted by CD49a and CD103 expression.

Using P_{adj} <0.05 as a cutoff, the data were analyzed to identify genes with a minimum of $\text{Log}_2(x) =$

0.5 Fold Change (LFC) to determine up- or downregulated genes, respectively. Per pathway analysis

utilizing both the Reactome and KEGG databases, there were no significant differences when

comparing DP to CD49a SP. However, several metabolic pathways were upregulated in DP vs DN,

including *Metabolism of lipids and lipoproteins, Fatty acid, triacylglycerol, and ketone body*

metabolism, and *Cholesterol biosynthesis.* A selection of pathways is plotted in **Figure 1A-B**. The

complete set of significantly enriched pathways can be found in the Supplementary file

"Significantly Enriched Pathways.xlsx".

Interestingly, CD49a SP and DP did not share any strictly metabolic pathways from the Reactome

when compared to DN yet were indistinguishable from one another at the global transcriptomic level

(7) (**Figure 1B**). They also did not share any pathways with each other in the KEGG database beyond

Steroid biosynthesis (**Figure 1B**). Metabolic differences between DP and DN but not CD49a SP and

DN could imply metabolic differences between CD49a SP and DP that these pathway analysis

methods are not sensitive enough to distinguish. In summary, T_{RM} subsets defined by integrin

expression were metabolically distinguishable from DN cells based on gene expression.

3.2   *Metabolic modeling predicts lipid-centric metabolism in T_{RM}*

Having performed basic pathway analyses and discovering a possible enrichment for lipid-centric

programming in T_{RM}, we developed an approach for modeling the metabolome of lung CD8 T cells

using the entirety of the transcriptomic data and used this more intensive approach to verify the

results of the basic analysis.

Metabolic models are networks of metabolites and the reactions that utilize and produce them.

Generally, these reactions are catalyzed using genome-encoded enzymes. Three metabolic

computational models were constructed: one for lung CD8 T cells expressing both CD49a and

CD103 (*tissuemodelDP*), one for the single positive subset (*tissuemodelCD49aSP*), and one for the

subset expressing neither integrin (*tissuemodelDN*). With these models, we used the expression of

enzymes present in a system to predict the amount of each metabolite present. These models together

incorporate nearly all antigen-experienced CD8 T cells in mouse lungs, encompassing both T_{RM} and

non-T_{RM} memory T cells.

Briefly, the published mouse metabolic model iMM1865 was used as a starting point (12). Then, the

Gene Inactivity Moderated by Metabolism and Expression (GIMME) algorithm was implemented in

the COBRA Toolbox (13) to conduct a flux balance analysis using our RNAseq transcript level data

to inform the level of flux for each reaction (10). GIMME alters the level of flux through a reaction

to minimize the total inconsistency score. Assuming a larger quantity of enzyme should produce a

larger flux through that reaction, GIMME increases (or decreases) flux through different reactions

depending on the level of transcript for that enzyme. When building metabolic models, an objective

function must be selected for which flux is to be maximized given the constraints. In line with most

other metabolic models, "BIOMASS" was selected, which itself is a combination of several reactions

related to cell maintenance that must have positive flux for cells to be viable. Given our constraints, a

single, nonzero optimal solution was predicted for all three of our models.

Our model predicted that our DP subset had the greatest flux through its *Beta Oxidation of Fatty*

Acids pathway, consistent with our initial pathway analysis (**Figure 1C**). In addition, the CD49a SP

subset differed from the DP subset with less flux through that pathway than both DP and DN, again

consistent with the initial pathway analysis. These results, in addition to our basic pathway analysis,

imply differences in T_{RM} metabolism between DP and CD49a SP subsets, as well as between

integrin-expressing and DN subsets.

3.3 T_{RM} CD8 T cell subsets have unique metabolic profiles directly *ex vivo*

Thus far, our transcriptomic analyses were largely consistent with that demonstrated for bulk T_{RM} in

the skin. Nonetheless, transcriptomic data and metabolic modeling do not clearly demonstrate that

functional differences exist between cell subsets whether circulating or tissue memory. The Seahorse

XF Mito Fuel Flex assay which (ideally) requires large cell numbers is considered a "gold standard"

for measuring a number of metabolic metrics such as oxygen consumption and acidification rate.

However, due to limiting cell numbers and our interest in the resting (not restimulated) metabolic

state of memory cells, the signal was not high and reproducible enough to draw conclusions,

demonstrating this approach is not feasible with unstimulated primary memory T cells. Instead, to

capture single-cell data, a flow-cytometric approach was performed. T cells with T_{RM}-like integrin expression are evident in mouse lung T cells 14 days post-infection (dpi) with HKx31 influenza A virus and may include cells with memory potential, while by 60 dpi memory is well established. To determine if the T cell subsets display any differences in their mitochondria, the mitochondrial mass of cells of different integrin profiles was evaluated by Mitotracker Green FM (Mitotracker) staining. Mitotracker preferentially accumulates in mitochondria by covalently binding free thiol groups independent of mitochondrial membrane potential (14). Both airway cells harvested through bronchoalveolar lavage (BAL) and cells remaining in the tissue after lavage (likely lung parenchymal cells) were examined for Mitotracker staining. Tissue vasculature-associated cells were excluded using a standard intravenous (IV) labeling approach (7).

We found no differences in Mitotracker staining in the lung or BAL at day 14 (**Figure 2A and S2**). However, at day 60 dpi, CD49a SP had significantly lower levels of mitochondrial mass than DP and DN counterparts in the lung, while DP had the highest staining (**Figure 2B**). This suggests that as memory T cell subsets develop, mitochondrial mass changes.

The question remained of whether the cell populations displayed differential mitochondrial functions, not just different mitochondrial masses. Oxidative phosphorylation is a primary role of the mitochondrion, as it is the most efficient means by which a cell can generate ATP. Cells with equivalent mitochondrial mass could potentially undergo differential rates of oxidative phosphorylation and therefore ATP production. Oxidative phosphorylation is linked to mitochondrial membrane polarization. TMRE, a fluorescent dye that preferentially stains polarized mitochondrial membranes was used as a proxy for processes that polarize the mitochondrion, though TMRE is not a direct readout of oxidative phosphorylation. To remove background staining, an equivalent sample was treated with both TMRE and the decoupling agent FCCP, whose signal was subtracted from the treatment group's signal.

Cells in the lung expressing CD103 had increased TMRE staining implying a greater degree of

membrane polarization and therefore oxidative phosphorylation 14 dpi (**Figure 2C**). Later, at 60+

dpi, no subset was significantly different (**Figure 2D**). At 60+ dpi, DP and DN cells have more

mitochondrial mass than CD49a SP cells (**Figure 2B**), while having the same membrane potential

(**Figure 2D**). Though not the only possibility, this is consistent with cells that have a higher spare

respiratory capacity (SRC), i.e., they may have the capability to use their higher amounts of

mitochondria to produce more ATP after reactivation.

Based on the observed differences, we wanted to investigate whether nutrient uptake was altered in a

similar way. Glucose is a metabolically essential molecule in a variety of cellular processes as both a

source of cell energy in the form of ATP as well as a carbon source in anabolism. To evaluate the

extent to which glucose was being taken up by the CD8 T cells, we utilized the glucose analog 2-

NBDG. This fluorescent analog undergoes facilitated diffusion through glucose transporters GLUT1-

4 as well as through active transport through SGLT1 and 2, recapitulating the kinetics of glucose

transport (15, 16, 17, 18, 19, 20). It is phosphorylated by hexokinase after entering the cell, which

traps the fluorescent molecule in the cell in a form that cannot proceed further in glycolysis.

Lung DP and CD49a SP displayed similar levels of 2-NBDG staining 14 dpi, however, these

populations displayed lower levels of staining compared with DN and CD103 SP cells, suggesting

that CD49a expressing subsets may be less glucose-dependent than CD49a negative (non-T_{RM})

counterparts (**Figure 2E**). A similar pattern was observed in BAL samples, with CD49a[pos] subsets

taking up lower levels of the dye (**Figure S2**). This pattern persisted at later memory time points 60+

dpi (**Figure 2F**) and is consistent with studies done on T_{RM} in the skin, which noted an increased

dependence on free fatty acids (FFAs) as opposed to glucose (9).

Having demonstrated significant differences in the uptake of the glucose analog 2-NBDG, we next

investigated the ability of T cells to take up free fatty acids with the hypothesis that lung T_{RM} utilize a

more lipid-centric metabolism, similar to skin (9). A fluorescently conjugated fatty acid (FFA) was

utilized to measure uptake. In the lung parenchyma, CD49a^{pos} subsets (CD49a SP and DP) took up

the highest amount of FFA. (**Figure 2G**). Similar results were found among the BAL subsets. These

findings were similar but distinct at 60+ dpi: DP and DN took up more FFAs than CD49a SP (**Figure**

2H). Other studies that have reported a dependence of T_{RM} on FFAs have not used CD49a as a

marker (9) and therefore may have excluded the CD49a^{pos} CD103^{neg} subset while possibly including

the CD103 SP subset. We expected our results to demonstrate that both DP and CD49a SP (the latter

not studied by other metabolic studies) were equally dependent on these FFAs in the lung, but we

found this only to be true at 14 dpi. At 60+ dpi, DP were much more like DN than their CD49a SP

T_{RM} counterparts. One interpretation consistent with the data is that CD49a SP utilizes a less lipid-

centric metabolic strategy than their DP counterparts.

We next wanted to determine if there were further differences in metabolic lipids. To address this,

Bodipy, a molecule that is taken up passively and stains cellular stores of neutral lipids, was used to

quantify the amount of neutral lipid present in lung T cells (21). Neutral lipids serve as a store of

potential energy as well as a carbon source, a valuable resource if the cells were to expand and

undergo effector functions in a nutrient-deficient environment.

Nearly identical patterns for BAL and lung were observed at 14 dpi (**Figure 2I, Figure S2**). DP cells

displayed the highest levels of neutral lipid storage, followed by CD49a SP then DN, with the lowest

levels in CD103 SP. While all six comparisons demonstrate that the four subsets are significantly

different from one another, it is remarkable that the subsets expressing CD49a (CD49a SP and DP)

stain brightly for neutral lipids while those not expressing CD49a (CD103 SP and DN) display low

levels. Having quick access to neutral lipids for both energy and carbon needs may prove important

for their protective function and/or long-term memory formation given these differences observed at

14 dpi were not seen at 60+ dpi (**Figure 2J**).

3.4   Glucose and lipid metabolites alter the phenotype of T$_{RM}$-like cells in vitro.

Noting the myriad of differences between T$_{RM}$ and non-T$_{RM}$, as well as among T$_{RM}$ subsets, we

hypothesized that the metabolic microenvironment is deterministic in their differentiation. Given that

tissue-specific metabolic substrates are impossible to control, an *in vitro* approach was employed.

T$_{RM}$ differentiation in vivo requires TGFβ (22). Similarly, T cells with T$_{RM}$-like integrin profiles can

be generated by stimulating naïve mouse splenocytes under defined conditions that include TGFβ

(23). Briefly, T cells were initially activated by culture on αCD3/CD28 coated plates for two days in

the presence of IL-2, followed by three days in non-coated plates supplemented with IL-2. T$_{RM}$-like

differentiation was achieved by placing these cells in IL-7 and TGFβ for six more days to elicit a

T$_{RM}$-like surface phenotype (23). To test the effects of glucose or lipids within as well as across

levels, both the concentration of glucose as well as lipids were independently varied (**Figure S3A**).

Glucose concentrations began with the concentration of glucose in standard cell culture media (4.5

mg/ml). This concentration has been demonstrated to ensure optimal T cell viability and activation,

especially over the course of several days in culture since glucose is rapidly utilized by cells, though

it is higher than that found in the blood or most other tissue microenvironments. To assess the impact

of lipids on T-cell differentiation *in vitro*, a broad range of exogenous lipid concentrations, which

includes both fatty acids and cholesterol (Millipore Sigma #L5146), was used. Lipid supplementation

varied from no exogenous lipid to 8x the concentration often used for cell culture.

First, the effect on CD49a expression was examined. Without exogenous lipid added, there was no

effect of glucose concentration on frequency, but with 2x or 4x lipid supplementation, generally, as

the concentration of glucose increased, the cell number (normalized per volume) and proportion of

CD49a positive cells both decreased (**Figures 3A-3B, S3, Supplementary File "2D combined stats

sheet.xlsx"**). The relationship between the absolute number of CD49a positive cells per volume and

glucose concentration was linear. 8x lipid supplementation was toxic as the number of live CD8 T

cells for the group was effectively zero. Overall, this paints a picture of increasing glucose

concentration causing a decrease in both the frequency and the absolute number of CD49a-positive

cells. This contrasts with the frequency of central memory marker CCR7 positivity (**Figures 4A, 4B,

and S3, Supplementary File "2D combined stats sheet.xlsx"**), which did not vary with lipid

supplementation, though increasing concentrations of glucose did seem to decrease the frequency of

CCR7 positive cells.

Looking at the effect of lipid supplementation at constant glucose levels on the differentiation of

CD8 T cells *in vitro*, a contrasting pattern emerged: within each concentration of glucose a 2x lipid

supplement increased the frequency of CD49a positive cells at the lower concentrations of glucose

tested (4.5 and 9.5 mg/mL) (**Figure 3B, Supplementary File "2D combined stats sheet.xlsx"**).

Further addition of lipid to 4x did not increase the frequency of CD49a positive cells. The absolute

number of CD49a positive cells per volume peaked at 2x and decreased upon further addition of lipid

for all concentrations of glucose tested. A similar but less drastic finding occurred for CCR7 positive

cells, but not for the frequency of positive cells (**Figure 4B, Supplementary File "2D combined

stats sheet.xlsx"**). Patterns for CD49a^{pos}CD103^{pos} cells were similar to those of CD49a SP (**Figure

S5**).

Taken together, these results imply that increasing glucose availability antagonizes the generation or

maintenance of memory CD8 T cells independent of lipid supplementation and affects the expression

of CD49a, at least in this assay. Furthermore, lipid supplementation is beneficial for CD49a

expression up to a point but can be toxic at too high a level.

4 Discussion

Using a genomic approach, we found memory T cell subsets in the lung exhibit distinct metabolic profiles. Furthermore, *in vitro* experiments conducted to drive T cells towards T_{RM}-like surface phenotypes in the presence of varying metabolic substrates suggest metabolite availability can influence memory T cell differentiation and/or survival. *In vivo* metabolite availability may be due to differences in local concentrations, the cell's ability to transport metabolic substrates into the cell, or the expression of receptor(s) needed to take up these metabolites from outside the cell.

Differential expression, pathway, and metabolic modeling of our RNAseq data led to the hypothesis that lung T_{RM} cells have distinct metabolic programming compared with circulating cells. Using both functional and descriptive methods, we produced a dataset for a variety of cell functions including mitochondrial profiling and nutrient uptake at both 14 and 60+ dpi. Integrating these results, what seems clear is that DP and CD49a SP, though different from each other in a few assays, share a similar pattern. This pattern is distinct from DN T cells, a mixed population consisting mostly of T cells with the phenotypes of effector memory and some central memory (7). The methods employed suggest DP and CD49a SP both take up more fatty acids, store more neutral lipids, and take up less glucose than DN cells at 14 dpi. At later memory time points, DP and CD49a SP diverge: DP cells have more mitochondrial mass and take up less glucose and more fatty acid than CD49a SP. Generally, memory cells have a higher SRC than effector cells, suggesting one possible advantage is to convey spare respiratory capacity (24). An increased spare respiratory capacity in DP but not CD49a SP could imply different roles in memory and effector responses. This should be cautiously interpreted since this SRC was not directly measured here, meriting further studies to validate this conclusion.

These findings are consistent with the interpretation that DP cells exhibit a more typical "resting"

memory phenotype and that CD49a SP could represent a more "activated" cell type. We have

previously reported that the CD49a SP subset has a number of more effector-like features such as

having a higher frequency of Granzyme B and perforin double positive cells, making them poised for

cell killing (7).

The metabolic profile of an individual T cell as well as a T cell subset is important functionally in

differentiation as well as effector function. Indeed, this is a leading hypothesis as to why some T cells

become T_{RM} and some do not (23). The functions of these subsets during a subsequent infection may

in part be achieved through the availability of different nutrients: if T_{RM} depend mostly on the uptake

of free fatty acids whereas naive and effector T cells depend more on glucose uptake, both may be

able to perform their functions without competing for energy resources. Furthermore, these profiles

can directly impact the cell signaling the cells are receiving. It is known that altering the ability of

naïve CD8 T cells to take up glucose directly impacts their likelihood of becoming memory cells

such that the most glucose-starved T cells become memory cells (25). Having a metabolic profile

that, at baseline, is not dependent on glucose may ensure that the internal cell signaling conferring

this memory phenotype continues to do so in a positive feedback manner. TCR stimulation is directly

related to the specificity of binding and the abundance of a TCR's peptide:MHC complex. Others

have shown that T cells take up more glucose when more activated (26). Having an abundance of

high-affinity interactions with peptide:MHC complexes may cause a hyper-local effect of glucose

starvation. The same group demonstrated that decreased availability of glucose led to increased

memory CD8 T cell populations. A lipid-centric metabolism by memory T-cells may assist naïve T

cells to respond to infection because they don't compete with other memory cells for glucose making

it less likely that memory T cells will starve out the naïve responses. This may support eliciting a

more diverse group of memory T cells at the end of an infection, which would increase the likelihood

of a productive immune response upon subsequent heterologous infection. An important caveat to

our interpretation is that NBDG doesn't always correlate with glucose uptake, even though it is a

glucose analog itself. As a result, more research must be done to confirm these conclusions.

The causal effect of metabolic factors on CD8 T cell differentiation has been studied but never in the

context of T_{RM} differentiation or in consideration of different T_{RM} subsets. In short, we found glucose

to be detrimental to the *in vitro* driven expression of T_{RM} marker CD49a in a linear manner, whereas

we found increasing lipid beneficial, to a point. Conversely, glucose did not affect the expression of

central memory marker CCR7, whereas the effect of lipids was similar for both CD49a and CCR7.

Optimal concentrations may exist that increase or decrease the expression of these markers. We

believe these findings have direct implications for the creation of therapies like CAR-T cells which

necessitate *ex vivo* culture conditions. For example, conditions could be optimized for the creation of

T-cells with a T_{RM}-like phenotype as observed among tumor-infiltrating lymphocytes associated with

positive outcomes (27, 28, 29, 30). Future studies should assess this possibility.

5   Figure Legends

Figure 1. Transcriptomic analysis of lung CD8 T cells 21 days post infection

Reactome and KEGG databases were queried with the lists of differentially expressed genes between

(A) DP and DN,or (B) CD49a SP and DN. Criteria for inclusion as differentially expressed genes

include $P_{adj} < 0.05$ and Log_2(Fold Change) > 0.5 or Log_2(Fold Change) < -0.5. Adjusted P value

cutoff for enrichment is $P_{adj}<0.05$. Only select pathways are shown here. A complete list can be

found in the supplemental. Datasets were generated using 10 total mice per N with N=3 per subset.

Metabolic models were created using the GIMME algorithm and the level of flux through Beta

Oxidation was plotted (C). DP (Double Positive, CD49a^{pos}CD103^{pos}); DN (Double Negative,

CD49a^{neg}CD103^{neg}); CD49a SP (CD49a Single Positive, CD49a^{pos}CD103^{neg}).

Figure 2. Normalized mouse IV-label-negative lung CD8 T cell MFIs from flow cytometric

assays.

Normalized MFIs of IV-label-negative lung CD8 T cells plotted for each assay at 14 and 60+ days

post infection derived from flow cytometric analysis. $*P < 0.05$, $**P < 0.01$, $***P < 0.001$, $****P <$

0.0001. DP (<u>D</u>ouble <u>P</u>ositive, CD49a^{pos}CD103^{pos}); DN (<u>D</u>ouble <u>N</u>egative, CD49a^{neg}CD103^{neg});

CD49a SP (<u>CD49a</u> <u>S</u>ingle <u>P</u>ositive, CD49a^{pos}CD103^{neg}) ; CD103 SP (<u>CD103</u> <u>S</u>ingle <u>P</u>ositive,

CD49a^{neg}CD103^{pos}). Mitochondrial mass was assessed with Mitotracker Green FM (A and B).

Mitochondrial membrane potential was assessed with TMRE (C and D). Glucose uptake was

assessed with glucose analogue 2-NBDG (E and F). Fatty acid uptake was assessed with

fluorescently conjugated fatty acids (G and H). Neutral lipid stores were assessed with Bodipy (I and

J). Data generated from two independent experiments of N= 4-10 mice per experiment.

Figure 3. Metabolite concentration impact on T$_{RM}$-like cell differentation and maintenance *in*

***vitro*.**

CD49a positive cells per µL (A) and percent CD49a positive (B) were assessed in metabolite

alteration assays where both lipid (0x-4x) and glucose (4.5-24.5 mg/mL) were altered after activation

in the presence of IL-2 and cultured with IL-7 and TGFβ. $*P < 0.05$, $**P < 0.01$, $***P < 0.001$,

$****P < 0.0001$. Small circles represent individual data points. Data generated from two independent

experiments of N=5 mice per experiment.

Figure 4. Metabolite concentration impact on T$_{CM}$-like cell differentation and maintenance *in*

***vitro*.**

CCR7 positive cells per µL (A) and percent CD49a positive (B) were assessed in metabolite

alteration assays where both lipid (0x-8x) and glucose (4.5-24.5 mg/mL) were altered after activation

in the presence of IL-2 and cultured with IL-7 and TGFβ. *P < 0.05, **P < 0.01, ***P < 0.001,

****P < 0.0001. Small circles represent individual data points. Data generated from two independent

experiments of N=5 mice per experiment.

Supplementary Figure 1: Gating scheme and representative fluorescent dye staining for flow

cytometry assays pictured in Figure 2.

Supplementary Figure 2: Normalized mouse bronchoalveolar lavage non-intravenously labelled

CD44^{pos} CD8 T cell MFIs from flow cytometric assays. *P < 0.05, **P < 0.01, ***P < 0.001, ****P

< 0.0001. DP (<u>D</u>ouble <u>P</u>ositive, CD49a^{pos}CD103^{pos}); CD49aSP (<u>CD49a</u> <u>S</u>ingle <u>P</u>ositive,

CD49a^{pos}CD103^{neg}); CD103SP (<u>CD103</u> <u>S</u>ingle <u>P</u>ositive, CD49a^{neg}CD103^{pos}); DN (<u>D</u>ouble <u>N</u>egative,

CD49a^{neg}CD103^{neg}). Data were generated from two independent experiments of 4-10 mice each.

Supplementary Figure 3. Assay layout for T cell differentiation studies: 16 combinations (4

glucose treatment groups, 4 lipid treatment groups permuted) were assayed. Glucose groups included

4.5, 9.5, 14.5, and 24.5 mg/mL. Lipid groups included 0x, 2x, 4x, and 8x recommended

concentrations (A). Gating scheme for differentiation studies (B).

Supplementary Figure 4. Metabolite concentration impact on T cell viability *in vitro*: Percent

alive in metabolite alteration assays where both lipid and glucose were altered. Data were generated

from two independent experiments of 5 mice each.

Supplementary Figure 5. Metabolite concentration impact on CD49a^{pos}CD103^{pos} T$_{RM}$-like cell

differentiation and maintenance *in vitro*: Percent CD49a^{pos}CD103^{pos} (B) and cell number per μL

(A) in metabolite alteration assays where both lipid (0x-4x) and glucose (4.5-24.5 mg/mL) were

altered after activation in the presence of IL-2 and culture with IL-7 and TGFβ. *P < 0.05, **P <

0.01, ***P < 0.001, ****P < 0.0001. Dots represent individual mice. Data were generated from two

independent experiments of 5 mice each.

Supplementary Table 1: Antibody information and catalog numbers. Additional information to

ensure reproducibility in methods.

Significantly Enriched Pathways.xlsx: The complete set of significantly enriched pathways from

which a subset are plotted in Figure 1 is contained in this table.

2D combined stats sheet.xlsx: The statistical comparisons for figures 3, 4, and Supplementary figure

5 are contained in this table.

6   Author Contributions

MS designed and performed experiments, analyzed and interpreted data, and wrote the manuscript.

AP assisted in conducting experiments and assay optimization. AG, KL, ECR, and TJ edited the

manuscript. ECR and KL provided technical support. DJT provided overall direction, procured

funding, interpreted data, and edited the manuscript.

7   Acknowledgments

We acknowledge the important contributions of the University of Rochester Medical Center

Genomic Research Center, the Flow Cytometry Core, and Vivarium. In addition, we are gracious for

the reagents provided by Drs. Andrea Amitrano and Minsoo Kim.

8   Data Availability Statement

The datasets for this study can be found on GitHub (https://github.com/sportiellomike/Lung-CD8-T-

cell-Immunometabolism). Previously published RNAseq data sets were used and are available here:

Geo with accession GSE179653 (https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE179653).

9   References

1.   Park SL, Buzzai A, Rautela J, Hor JL, Hochheiser K, Effern M, et al. Tissue-resident memory CD8(+) T cells promote melanoma-immune equilibrium in skin. Nature. 2019;565(7739):366-71.

2.   Park SL, Zaid A, Hor JL, Christo SN, Prier JE, Davies B, et al. Local proliferation maintains a stable pool of tissue-resident memory T cells after antiviral recall responses. Nat Immunol. 2018;19(2):183-91.

3.   Ray SJ, Franki SN, Pierce RH, Dimitrova S, Koteliansky V, Sprague AG, et al. The collagen binding alpha1beta1 integrin VLA-1 regulates CD8 T cell-mediated immune protection against heterologous influenza infection. Immunity. 2004;20(2):167-79.

4.   Jozwik A, Habibi MS, Paras A, Zhu J, Guvenel A, Dhariwal J, et al. RSV-specific airway resident memory CD8+ T cells and differential disease severity after experimental human infection. Nat Commun. 2015;6:10224.

5.   Poon MML, Rybkina K, Kato Y, Kubota M, Matsumoto R, Bloom NI, et al. SARS-CoV-2 infection generates tissue-localized immunological memory in humans. Sci Immunol. 2021;6(65):eabl9105.

6.   Carbone FR. Unique properties of tissue-resident memory T cells in the lungs: implications for COVID-19 and other respiratory diseases. Nat Rev Immunol. 2023;23(5):329-35.

7.   Reilly EC, Sportiello M, Emo KL, Amitrano AM, Jha R, Kumar ABR, et al. CD49a Identifies Polyfunctional Memory CD8 T Cell Subsets that Persist in the Lungs After Influenza Infection. Front Immunol. 2021;12:728669.

8.   Pan Y, Kupper TS. Metabolic Reprogramming and Longevity of Tissue-Resident Memory T Cells. Front Immunol. 2018;9:1347.

9.   Pan Y, Tian T, Park CO, Lofftus SY, Mei S, Liu X, et al. Survival of tissue-resident memory T cells requires exogenous lipid uptake and metabolism. Nature. 2017;543(7644):252-6.

10.   Becker SA, Palsson BO. Context-specific metabolic networks are consistent with experiments. PLoS Comput Biol. 2008;4(5):e1000082.

11.   Han F, Li G, Dai S, Huang J. Genome-wide metabolic model to improve understanding of CD4(+) T cell metabolism, immunometabolism and application in drug design. Mol Biosyst. 2016;12(2):431-43.

12.   Khodaee S, Asgari Y, Totonchi M, Karimi-Jafari MH. iMM1865: A New Reconstruction of Mouse Genome-Scale Metabolic Model. Sci Rep. 2020;10(1):6177.

13.   Heirendt L, Arreckx S, Pfau T, Mendoza SN, Richelle A, Heinken A, et al. Creation and analysis of biochemical constraint-based models using the COBRA Toolbox v.3.0. Nat Protoc. 2019;14(3):639-702.

14.   Presley AD, Fuller KM, Arriaga EA. MitoTracker Green labeling of mitochondrial proteins and their subsequent analysis by capillary electrophoresis with laser-induced fluorescence detection. J Chromatogr B Analyt Technol Biomed Life Sci. 2003;793(1):141-50.

15.   Lu Y-T, Ma X-L, Xu Y-H, Hu J, Wang F, Qin W-Y, et al. A Fluorescent Glucose Transport Assay for Screening SGLT2 Inhibitors in Endogenous SGLT2-Expressing HK-2 Cells. Nat Prod Bioprospect. 2019;9(1):13-21.

16.   Fang P, Yu M, Min W, Wan D, Han S, Shan Y, et al. Effect of baicalin on GLUT4 expression
and glucose uptake in myotubes of rats. Life Sciences. 2018;196:156-61.

17.   Ferreira JM, Burnett AL, Rameau GA. Activity-dependent regulation of surface glucose
transporter-3. J Neurosci. 2011;31(6):1991-9.

18.   Marín-Juez R, Rovira M, Crespo D, van der Vaart M, Spaink HP, Planas JV. GLUT2-
mediated glucose uptake and availability are required for embryonic brain development in zebrafish.
J Cereb Blood Flow Metab. 2015;35(1):74-85.

19.   Meireles P, Sales-Dias J, Andrade CM, Mello-Vieira J, Mancio-Silva L, Simas JP, et al.
GLUT1-mediated glucose uptake plays a crucial role during Plasmodium hepatic infection. Cell
Microbiol. 2017;19(2).

20.   Blodgett AB, Kothinti RK, Kamyshko I, Petering DH, Kumar S, Tabatabai NM. A
fluorescence method for measurement of glucose transport in kidney cells. Diabetes Technol Ther.
2011;13(7):743-51.

21.   Rumin J, Bonnefond H, Saint-Jean B, Rouxel C, Sciandra A, Bernard O, et al. The use of
fluorescent Nile red and BODIPY for lipid measurement in microalgae. Biotechnology for Biofuels.
2015;8(1):42.

22.   Casey KA, Fraser KA, Schenkel JM, Moran A, Abt MC, Beura LK, et al. Antigen-
independent differentiation and maintenance of effector-like resident memory T cells in tissues. J
Immunol. 2012;188(10):4866-75.

23.   Bromley SK, Akbaba H, Mani V, Mora-Buch R, Chasse AY, Sama A, et al. CD49a Regulates
Cutaneous Resident Memory CD8(+) T Cell Persistence and Response. Cell Rep.
2020;32(9):108085.

24.   van der Windt Gerritje JW, Everts B, Chang C-H, Curtis Jonathan D, Freitas Tori C, Amiel E,
et al. Mitochondrial Respiratory Capacity Is a Critical Regulator of CD8+ T Cell Memory
Development. Immunity. 2012;36(1):68-78.

25.   Sukumar M, Liu J, Ji Y, Subramanian M, Crompton JG, Yu Z, et al. Inhibiting glycolytic
metabolism enhances CD8+ T cell memory and antitumor function. J Clin Invest.
2013;123(10):4479-88.

26.   Jacobs SR, Herman CE, Maciver NJ, Wofford JA, Wieman HL, Hammen JJ, et al. Glucose
uptake is limiting in T cell activation and requires CD28-mediated Akt-dependent and independent
pathways. J Immunol. 2008;180(7):4476-86.

27.   Loi S, Sirtaine N, Piette F, Salgado R, Viale G, Van Eenoo F, et al. Prognostic and predictive
value of tumor-infiltrating lymphocytes in a phase III randomized adjuvant breast cancer trial in
node-positive breast cancer comparing the addition of docetaxel to doxorubicin with doxorubicin-
based chemotherapy: BIG 02-98. J Clin Oncol. 2013;31(7):860-7.

28.   Denkert C, Loibl S, Noske A, Roller M, Muller BM, Komor M, et al. Tumor-associated
lymphocytes as an independent predictor of response to neoadjuvant chemotherapy in breast cancer. J
Clin Oncol. 2010;28(1):105-13.

29.   Wang R, Song Y, Hu T, Wang X, Jiang Y, Zhang D, et al. Decreased CD8(+) Lymphocytic
Infiltration in Multifocal and Multicentric Glioblastomas. Front Oncol. 2021;11:748277.

30.   Smazynski J, Webb JR. Resident Memory-Like Tumor-Infiltrating Lymphocytes (TILRM):
Latest Players in the Immuno-Oncology Repertoire. Front Immunol. 2018;9:1741.

A

DP vs DN

Steroid biosynthesis
Cholesterol biosynthesis Homo sapiens R-HSA-191273
Regulation of cholesterol biosynthesis by SREBP (SREBF) Homo sapiens R-HSA-1655829
Glycosphingolipid biosynthesis
Sphingolipid signaling pathway
Fatty acid, triacylglycerol, and ketone body metabolism Homo sapiens R-HSA-535734
Metabolism of lipids and lipoproteins Homo sapiens R-HSA-556833
Metabolism Homo sapiens R-HSA-1430728
Metabolism of proteins Homo sapiens R-HSA-392499
Purine metabolism Homo sapiens R-HSA-73847
Metabolism of amino acids and derivatives Homo sapiens R-HSA-71291
Selenoamino acid metabolism Homo sapiens R-HSA-2408522
Selenocysteine synthesis Homo sapiens R-HSA-2408557

-10000 -5000 -1000 -500 -100 -50 -10 0 10 50 100 500

B

CD49a vs DN

Steroid biosynthesis
Cholesterol biosynthesis Homo sapiens R-HSA-191273
Regulation of cholesterol biosynthesis by SREBP (SREBF) Homo sapiens R-HSA-1655829
Glycosphingolipid biosynthesis
Sphingolipid signaling pathway
Fatty acid, triacylglycerol, and ketone body metabolism Homo sapiens R-HSA-535734
Metabolism of lipids and lipoproteins Homo sapiens R-HSA-556833
Metabolism Homo sapiens R-HSA-1430728
Metabolism of proteins Homo sapiens R-HSA-392499
Purine metabolism Homo sapiens R-HSA-73847
Metabolism of amino acids and derivatives Homo sapiens R-HSA-71291
Selenoamino acid metabolism Homo sapiens R-HSA-2408522
Selenocysteine synthesis Homo sapiens R-HSA-2408557

-100 -50 -10 0 10 50 100

Combined Score

P_{adj}

0.01 0.02 0.03

C

Net flux of Beta Oxidation by Subset

Net Flux (mmol g^{-1} DW^{-1} h^{-1})

DP   CD49a SP   DN