1   **Mouse lung Tissue Resident Memory (TRM) CD8 T cell subsets have distinct metabolic**

2   **profiles from each other and from non-TRM**

Mike Sportiello[1,2], Alexis Poindexter[1], Emma C. Reilly[1], Adam Geber[1,2], Kris Lambert Emo[1],

Taylor N. Jones[1], David J. Topham[1]

[1]Center for Vaccine Biology and Immunology, University of Rochester Medical Center, Rochester,

NY, United States

[2] Medical Scientist Training Program, University of Rochester Medical Center, Rochester, NY 14642,

USA

* Correspondence:

David J. Topham, PhD

David_Topham@urmc.rochester.edu

Abstract

Tissue resident memory CD8 T cells (T_{RM}) principally reside in peripheral non-lymphoid tissues and

confer protection against a variety of illnesses ranging from infections to cancers. The functions of

different CD8 T cell subsets in the skin have been linked with distinct metabolic pathways and differ

from other CD8 T cell subsets. For example, they undergo fatty acid oxidation and oxidative

phosphorylation to a greater degree than circulating memory and naïve cells. However, it is unclear

if this programming is universal. Our work demonstrates T_{RM} are not homogenous and, based on

integrin expression, exist as subsets with distinct gene expression and function.   We hypothesize

that lung T_{RM} with different integrin profiles will utilize unique metabolic programs. First, differential

expression and pathway analysis was conducted on RNAseq datasets yielding important differences

between subsets. Next, metabolic models were constructed and interrogated. Functional metabolite

uptake assays were conducted, and the levels of oxidative phosphorylation, mitochondrial mass,

and neutral lipids were measured. Furthermore, T cell differentiation studies were conducted varying

concentrations of metabolites to investigate the causal relationship in T_{RM} development,

demonstrating that, *in vitro*, lipid conditions impact memory cell maintenance generally, and that

glucose concentration impacts T_{RM}-like differentiation specifically, with no effect on central memory-

like T cell differentiation. In summary, a diverse set of analyses were utilized to demonstrate lung

T_{RM} have unique metabolic profiles and that metabolic microenvironments alter differentiation *in*

vitro.

1   Introduction

Tissue resident memory T cells (T_{RM}) confer protection against a variety of illnesses ranging from

infectious diseases to cancers. As demonstrated in the skin, transfer of T_{RM} from immune to naïve

hosts is sufficient to prevent herpes simplex virus type 1 (HSV-1)-associated pathology and,

comparably, the presence of antigen-specific T_{RM} in a mouse model of melanoma can limit tumor

growth (1, 2). In the lungs, T_{RM} are critical for protection against influenza and respiratory syncytial

virus, even in the absence of antibodies specific for those viruses (3, 4). T_{RM} have been shown to act

as both antigen-specific sentinels linking the innate and adaptive immune systems as well as cytotoxic

mediators themselves (5). This functional capacity makes them important players in immunity across

multiple organ systems and disease states.

TRM are heterogenous in terms of integrin expression, migration, effector potential, and gene

expression (5). Furthermore, the functions of different CD8 T cell subsets have been linked with

distinct metabolic pathways. For example, skin T_{RM} display a unique metabolic profile relative to other

CD8 memory cells subsets. They undergo fatty acid oxidation to a greater degree than other

circulating memory cells and naïve cells which tend to have higher glycolytic fluxes (6, 7). However, it

is unclear if this programming is universal among all T_{RM} subsets and whether it holds true in other

tissues such as the lung. It is also not yet know whether memory CD8 T cell subsets that express

different combinations of surface integrins CD49a and CD103 in addition to S1PR1 antagonist CD69

share metabolic profiles. Performing RNAseq on these populations yielded distinct transcriptomes (5),

suggesting the overall metabolic profiles may differ, perhaps to assist in their survival and function.

Furthermore, these unique integrin expression profiles could be linked to different tissue localization

and may require differential metabolic cell programming.

2 Materials and Methods

2.1 RNA sequencing analysis

RNA sequencing data and lists of differentially expressed genes were obtained from previously published work from our lab (GSE179653). Full code that generated those differentially expressed genes as well as further analyses and figures is publicly available at https://github.com/sportiellomike.

2.2 Metabolic modelling

Mouse metabolic model iMM1865 was used as a starting point, onto which our publicly accessible transcriptomic data for CD49a^{pos}CD103^{pos} (Double Positive, DP), CD49a^{pos}CD103^{neg} (CD49a Single Positive, CD49a SP), or CD49a^{neg}CD103^{neg} (Double Negative, DN) was applied using the Gene Inactivity Moderated by Metabolism and Expression (GIMME algorithm) (8). A threshold value of the median transcript per kilobase million (TPM) for the 1,865 genes in the metabolic model was used. Reaction constraints were added according to previously published research (9). All code used to generate input data, figures, and the models themselves is available at https://github.com/sportiellomike.

2.3 Mice

Mice were housed in approved microisolator cages, within a pathogen-free facility that is accredited by the Association for Assessment and Accreditation of Laboratory Animal Care. It is staffed with a number of personnel including those with expertise in husbandry, technical skills, and veterinary personnel. C57BL/6J mice (Jackson Laboratories) used for experiments were infected 8-10 weeks after birth. This study was carried out in strict accordance with the recommendations in the *Guide for the Care and Use of Laboratory Animals* as defined by the NIH. The Institutional Animal Care and Use Committee of the University of Rochester approved all protocols.

2.4 Virus and infection

Mice were anesthetized with 3,3,3-tribromoethanol. Upon verification of sedation, mice were infected

intranasally with 10^5 EID_{50} of HKx31 human influenza A virus in 30uL volume. After observation to

ensure they recovered from anesthesia, mice were monitored daily for weight and overall morbidity to

comply with all regulations.

2.5   Lung and Bronchoalveolar lavage processing

Bronchoalveolar lavage (BAL) was collected by inserting a flexible Teflon catheter into a slit in the

trachea in proximity to the larynx. A 1mL syringe was connected to the catheter and the lungs were

flushed 3 times with 1x PBS (volumes of 1mL, 900uL, and 800uL). BAL was spun down at 424xg for

10 minutes. Red blood cells were eliminated through ACK mediated lysis. Cells were washed in 1x

PBS containing 1% FBS (PBS+serum) and resuspended in PBS+serum.

The lungs were harvested and separated into right and left lobes prior to collection into RPMI + 8%

FBS (Corning #10040CV). Lungs were transferred into Miltenyi Gentle MACS tubes containing 2 mL

of 2mg/mL collagenase II (Worthington #LS004177) and mechanically ground up by a Gentle MACS

machine. 3 additional mL of 2 mg/mL collagenase II solution was added before incubation for 30 min

at 37° C. Further mechanical digestion by Gentle MACS was done at this point. Tubes were spun at

233xg at room temperature for 2 minutes.

Pellets were then dissolved in existing supernatant and transferred through 100 μm strainers.

Contents were quantitatively transferred using 5 mL of RPMI + 8% FBS. Tubes were spun at 424xg

for 6 minutes at room temperature. The supernatant was discarded, and dissolved in 4 mL 40% Percoll

solution, which was then layered on 3 mL 75% Percoll solution. These tubes were then spun at 931xg

for 20 minutes at room temperature with no centrifuge break so as not to disturb the layers of the

gradient.

The top layer of lipid-rich liquid was aspirated off, and the interface (approximately 2 mL) of the two

Percoll densities was collected where the cells were. Tubes were filled with RPMI + 8% FBS and spun

down at 424xg for 6 minutes at room temperature.

2.6 Spleen processing to achieve single cell suspension

C57BL/6J mouse spleens were obtained from sacrificed mice and mechanically digested by grinding between two frosted microscope slides without exogenous enzyme added. After grinding, Single cell suspensions were passed through plastic mesh to capture large non-single cellular material and centrifuged in conical vials at 424xg at 4°C for 6 minutes. Cells were resuspended in 3 mL of ACK lysis buffer (Cold Spring Harbor Protocol: doi:10.1101/pdb.rec083295) for 5 minutes at room temperature. Volume was brought up to 15 mL in PBS + serum followed by centrifuging at 424xg at 4°C for 6 minutes.

2.7 T_{RM}-like Media preparation

The following were added to 500 mL DMEM (Corning Catalog No. 10-017-CV): 50 mL fetal bovine serum, 5 mL L-glutamine (Fisher Scientific Catalog No. 25030081), 12.5 mL HEPES buffer (Fisher Scientific Catalog No. 15630080), 6.5 MEM Non-Essential Amino Acids Solution (Fisher Scientific 11140050), 0.6 mL of 1:300 dilution of beta mercaptoethanol in PBS, 5 mL penicillin streptomycin (Fisher Scientific Catalog No. 15140122), 0.6 mL gentamycin (Fisher Scientific Catalog No. 15710064). Contents were sterile filtered.

2.8 Fatty Acid Uptake

Fatty Acid Uptake Media (FUM) was created by dissolving 2 µL stock solution made according to package insert per 1 mL of Assay Buffer (AAT Bioquest Screen Quest Fluorimetric Fatty Acid Uptake Assay Kit #36385). Once a single cell suspension was obtained from the harvesting protocol, cells were plated in 100 µL of FUM and 100 µL Leibowitz media for a final volume of 200 µL. Vehicle control used equivalent volume of dimethyl sulfoxide (DMSO) in Assay Buffer. Cells were incubated for 30 minutes at 37° C, then washed 1.5 times with PBS + 1% serum before staining.

2.9 Neutral Lipid Staining

Bodipy 505/515 (Fisher Scientific Catalog No. D3921) was dissolved in DMSO to yield a stock of 1

mg/µL concentration. Once a single cell suspension was obtained from the harvesting protocol, each

sample of cells in 1 mL of PBS were inoculated with Bodipy to yield a final concentration of 2 mg/mL,

or DMSO for the vehicle control. Samples were incubated for 30 minutes at 37° C, then washed 1.5

times with PBS + 1% serum before staining.

2.10 2-(N-(7-Nitrobenz-2-oxa-1,3-diazol-4-yl)Amino)-2-Deoxyglucose Uptake

2-(N-(7-Nitrobenz-2-oxa-1,3-diazol-4-yl)Amino)-2-Deoxyglucose (2-NBDG) (Abcam Catalog No.

ab146200) was dissolved in DMSO to yield a 20 mM stock solution. After a single cell suspension was

obtained from the harvesting protocol, cells were resuspended in 500 µL non-supplemented Leibowitz

media and incubated at 37° C with a final 2-NBDG concentration of 80 µM or DMSO for the vehicle

control for 35 minutes at 37° C. Samples were washed 1.5 times with PBS + 1% serum before staining.

2.11 Mitochondrial Mass Quantification

Mitotracker Green FM (Mitotracker) (Fisher Scientific Catalog No. M7514) was dissolved in DMSO to

obtain a 1 mM stock solution. After a single cell suspension was obtained from the harvesting protocol,

each sample was dissolved in 500 µL PBS + 1% serum. Samples were inoculated with Mitotracker

for a final concentration of 50 nM or equivalent volume of DMSO for the vehicle control and incubated

for 45 minutes at 37° C.

2.12 Oxidative Phosphorylation Quantification

Tetramethylrhodamine, Ethyl Ester, Perchlorate (TMRE) (Cayman Chemical Catalog No. 701310)

was prepared in a 0.5 mM working stock by dissolving in DMSO, according to package insert. After a

single cell suspension was obtained from the harvesting protocol, TMRE was added to two separate

aliquots of 5 mL RPMI + 8% FBS (TMRE Media) for a final concentration of 250 nM. To one of these

aliquots, 2-[2-[4-(trifluoromethoxy)phenyl]hydrazinylidene]-propanedinitrile (FCCP) was added

(FCCP-TMRE Media) for a final concentration of 1 µM. Samples were split into two and dissolved into

500 μL of either TMRE Media or FCCP-TMRE Media. Samples were incubated at 37° C for 30

minutes. For vehicle control, DMSO was added in place of TMRE. Cells were washed 1.5 times with

PBS + 1% FBS prior to staining.

2.13 T_{RM}-like cell culture conditions

24 well plates were coated with anti-CD3/28 stimulating antibodies (Biolegend Catalog Nos. 100302

and 102102) at final concentration of 5 μg/mL each overnight at 4° C. The next day, 2 million single

cell splenocytes were plated in 1 mL T_{RM}-like culture media onto each well with 7.5 ng/mL human IL-

2 (Obtained from the NIH). Two days later, cells were collected, centrifuged at 424xg at 4° C for 6

minutes, and plated onto non-coated 24 well plates in 1 mL TRM-like culture media with 7.5 ng/mL

IL-2 for three days. After that, cells were collected, centrifuged at 424xg at 4° C for 6 minutes, and

plated in TRM-like culture media with 10 ng/mL mouse IL-7 (Biolegend Catalog No. 577804) and 10

ng/mL human TGFβ1 (Fisher Scientific PHG9214) for 6 days.

Additional glucose (Fisher Scientific Catalog No. A2494001) and lipid supplement (Millipore Sigma

Catalog No. L5146) were added to TRM-like cell culture media starting from the day of spleen

harvest (day 0) throughout the end of the experiment (day 11, day 6 post TGFβ1 addition). Soluble

lipid stores were created in Kolliphor P 188 (Millipore Sigma K4894) according to lipid supplement

instructions.

2.14 Staining for Flow Cytometry

Cells were suspended in 100 μL staining buffer for 30 minutes at 4° C. Staining buffer was created by

adding staining antibodies to PBS below for a final concentration of 1 μg/mL each, or 0.2 μL of live-

dead stock solution/100 μL staining buffer. Live-dead stock solution was prepared according to

package insert available on the vendor website. αCD16/32 was added for a final concentration of 2.5

μg/mL. αCD45 was previously injected at time of organ harvest intravenously at a concentration of 0.2

μg stock antibody/100 μL PBS in a volume of 100 μL. See table S1 for more information. All antibodies

used Bangs Laboratories compensation beads (Bangs #556) which were stained an equivalent time.

Live-dead stain used Life Technologies ArC reactive beads (Invitrogen A10346). Because the vast

majority of cells at day 14 post infection will have been antigen experienced, CD44 was not used as

a marker on day 14 experiments. Experiments were run on an 18-color BD LSR II flow cytometer.

Gating strategy is displayed in Figure S1.

2.15   MFI Determination and Normalization

The four subsets of interest were manually gated after selecting for lymphocytes (FSC-A and SSC-

A), singlets (FSC-A and FSC-H), live cells, IV-CD45^{neg} cells, TCRβ^{high} and CD8α^{high} cells. DN (Double

Negative) was defined as CD103^{low} and CD49a^{low}, DP (Double Positive) as CD103^{high} and CD49a^{high},

CD103 SP (CD103 Single Positive) as CD103^{high} CD49a^{low}, and CD49a SP (CD49a Single Positive)

as CD49a^{high} and CD103^{low}.

Median fluorescence intensities (MFIs) were calculated using Flowjo (version 10.6.2). Data was

normalized by dividing the MFI of each subset by the sum of all four subsets' MFIs for each mouse

and multiplying by 4. For the assays using TMRE, FCCP was subtracted from the TMRE MFI for each

subset within each mouse. That number was divided by the sum of MFIs of the uncoupled control

(FCCP+TMRE)   subsets   for   each   mouse.   Normalized   data   can   be   found   at

https://github.com/sportiellomike.

2.16   Statistical Tests

After repeated measures ANOVA, Multiple pairwise comparisons were evaluated using the Student's

T test with Benjamini-Hochberg correction. If $P_{adjusted}<0.05$, results were deemed significant. For the

linear regression analysis, Pearson's method was used. R Packages ggplot2 (version 3.3.2) and

rstatix (0.7.0) were used for data plotting and statistical testing respectively. The code used to perform

statistical tests and generate figures can be found here: https://github.com/sportiellomike.

3   Results

3.1   Pathway analysis for metabolic analyses.

T_{RM} can be defined by their surface protein expression and multiple markers have been used to

including CD69, CD49a, and CD103. In the lung, CD44 expression identifies antigen-experienced

memory T cells while CD49a^{pos} marks T_{RM}, and when used in combination with CD103, yields two T_{RM}

subsets: CD49a^{pos}CD103^{pos} (Double Positive, DP) and CD49a^{pos}CD103^{neg} (CD49a Single Positive,

CD49a SP) (5). At later memory time points, very few CD49a^{neg}CD103^{pos} (CD103 Single Positive,

CD103 SP) remain, suggesting they have either died, left the tissue, or altered their integrin

expression. CD49a^{neg}CD103^{neg} cells that remain at this time point resemble T cells with an Effector

Memory phenotype (T_{EM}) and may be circulating (5). To more fully describe the metabolic profiles of

T_{RM} based on integrin expression, we utilized the publicly accessible RNA sequencing data set we

previously reported (5). This data set was generated from antigen-experienced mouse lung tissue

CD8 T cells 21 days post infection (dpi) that were sorted by CD49a and CD103 expression and

submitted for poly-A enriched mRNA sequencing. Double Negative cells (DN) were defined as

CD103^{neg} and CD49a^{neg}, Double Positive (DP) as CD103^{pos} and CD49a^{pos}, CD103 Single Positive

(CD103 SP) as CD103^{pos} CD49a^{neg}, and CD49a Single Positive (CD49a SP) as CD49a^{pos} and

CD103^{neg}.

Using P_{adj} <0.05 as a cutoff, the data were analyzed using Log_2(Fold Change) (LFC) cutoffs of >0.5

or <-0.5 to determine up- or downregulated genes, respectively. Per pathway analysis, there were no

enriched pathways in either the up- or downregulated genes when comparing DP to CD49a SP. Seven

pathways from the Reactome database were upregulated in DP vs DN, including *Metabolism of lipids*

and lipoproteins, *Fatty acid, triacylglycerol, and ketone body metabolism*, and *Cholesterol*

biosynthesis (**Figure 1A**), with a large number of downregulated pathways, largely related to RNA

translation and decay. Equivalent analysis using the KEGG database yielded eight upregulated

pathways including *Steroid biosynthesis*, with just one downregulated pathway *Ribosome* (**Figure**

1B). Many pathways including *Fatty acid biosynthesis*, *Fatty acid degradation*, and *Fatty acid*

elongation were not significantly different in the pathway analysis using KEGG.

Interestingly, CD49a SP, though indistinguishable from DP at the global transcriptomic level (5), did

not share any strictly metabolic pathways from Reactome with DP (**Figure 1C**). They also did not

share any with each other in the KEGG database beyond *Steroid biosynthesis* (**Figure 1D**).

Specifically, CD49a SP did not share *Cholesterol biosynthesis*, *Regulation of cholesterol biosynthesis*

by SREBP (SREBF), *Metabolism of lipids and lipoproteins*, or *Fatty acid, triacylglycerol, and ketone*

body metabolism which were all upregulated in DP vs DN but not CD49a SP vs DN. Metabolic

differences between DP and DN but not CD49a SP and DN could imply metabolic differences between

CD49a SP and DP that these pathway analysis methods are not sensitive enough to distinguish.

3.2   *Metabolic modelling predicts lipid-centric metabolism in T$_{RM}$*

Having performed basic pathway analyses and discovering a possible enrichment for lipid-centric

programming in T$_{RM}$, we were interested in modelling the metabolome of lung CD8 T cells using the

entirety of the transcriptomic data to see if this more intensive approach yielded similar results.

Therefore, three metabolic computational models were constructed: one for lung CD8 T cells

expressing both CD49a and CD103 (*tissuemodelDP*), one for the single positive subset

(*tissuemodelCD49aSP*) and one for the subset expressing neither integrin (*tissuemodelDN*).

Because so few CD103 SP cells exist at memory timepoints, a metabolic model was not created for

this subset. These novel models together represent nearly all antigen experienced CD8 T cells in

mouse lung, encompassing both T$_{RM}$ and non-T$_{RM}$ cells that happen to be trafficking through the

lung.

Briefly, the published mouse metabolic model iMM1865 was used as a starting point (10). Then, the

Gene Inactivity Moderated by Metabolism and Expression (GIMME) algorithm was implemented in

COBRA Toolbox (11) to conduct a flux balance analysis using our RNAseq transcript level data to

inform the level of flux for each reaction (8). When building metabolic models, an objective function

must be selected for which flux is to be maximized given the constraints. In line with the majority of

other metabolic models, "BIOMASS" was selected, which itself is a combination of a number of

reactions related to cell maintenance that must have positive flux for cells to be viable. Given our

constraints, a single, nonzero optimal solution was predicted for all three of our models.

Our model predicted that our DP subset had the greatest flux through its Beta Oxidation of Fatty

Acids pathway, consistent with our initial pathway analysis (Figure 2). In addition, our CD49aSP

subset differed from our DP subset with less flux through that pathway than both DP and DN, again

consistent with initial pathway analysis. These results, in addition to our initial pathway analysis,

imply differences in T_{RM} metabolism between DP and CD49a SP subsets.

3.3 Tissue resident memory CD8 T cell subsets have unique metabolic profiles directly ex vivo

Thus far, our transcriptomic analyses were largely consistent with that demonstrated for T_{RM} in skin.

Nonetheless, transcriptomic data and metabolic modelling do not clearly demonstrate that functional

differences exist between T_{RM} subsets or between circulating cells and T_{RM}. To further understand

these T_{RM}, a Seahorse XF Mito Fuel Flex Test assay was performed. However, even after pooling the

T cells subsetted from 40 mice, the signal was not high and reproducible enough to draw conclusions

(data not shown), suggesting this approach is not feasible in this context. Instead a flow-cytometric

approach was performed. To determine if the T_{RM} subsets display any differences in their

mitochondria, the mitochondrial mass of cells of different integrin profiles at 14 days post infection

(dpi) were evaluated by Mitotracker Green FM (Mitotracker) staining. Mitotracker preferentially

accumulates in mitochondria by covalently binding free thiol groups independent of mitochondrial

membrane potential (12). Mice were infected with HKx31 influenza virus and cells were collected at

14 dpi, a time when CD8 T cells with a T_{RM} phenotype are numerous. Both airway cells harvested

through bronchoalveolar lavage (BAL) and cells remaining in the tissue after lavage (likely lung

parenchymal cells) were examined for Mitotracker staining. To ensure that cells in the lung tissue

were not associated with or within the vasculature, a standard intravenous (IV) labeling exclusion

approach was utilized (5).

We found that both DP and CD103 SP contain higher mitochondrial mass than the DN subset (**Figure**

3A). Similar results were found in BAL, though the comparison between CD103 SP and DN did not

achieve significance (**Figure S2**). Mitochondrial mass was also assessed at 62 dpi. CD103 SP cells

were not included in this analysis because so few cells could be detected. At this later timepoint,

CD49a SP had significantly lower levels of mitochondrial mass than its DP and DN counterparts in

the lung (**Figure 3B**). Day 62 may be considered by some to be a more classical "memory" time point,

suggesting mitochondrial mass changes in some subsets as memory develops.

The question of whether the cell populations displayed differential mitochondrial functions, not just

different mitochondrial masses, remained. Oxidative phosphorylation is a primary role of the

mitochondrion, as it is the most efficient means by which a cell can generate ATP. Cells with equivalent

mitochondrial mass could potentially undergo differential rates of oxidative phosphorylation and

therefore ATP production. To evaluate this level of oxidative phosphorylation, TMRE, a fluorescent

dye preferentially taken up by active mitochondria with membrane potentials, was used. To remove

background staining, an equivalent sample was treated with both TMRE and the decoupling agent

FCCP, whose MFI was subtracted from the treatment group's MFI.

Cells in the lung expressing CD103 (DP and CD103 SP) had increased TMRE staining implying a

greater degree of oxidative phosphorylation (**Figure 3C**). Later, at 62 dpi, no comparison was

significantly different (**Figure 3D**). At 62 dpi, DP and DN cells have more mitochondrial mass than the

CD49 SP subset (**Figure 3B**), while undergoing the same amount of oxidative phosphorylation

(**Figure 3D**). This implies that at later memory timepoints, DP and DN cells have a higher spare

respiratory capacity (SRC): they may have the capability to use their higher amounts of mitochondria

to produce more ATP after reactivation. This is a well-documented phenomenon when comparing

memory T cells to effector T cells or naïve T cells, but never before have different T_{RM} subsets been

compared this to each other (13). Generally, memory cells have a higher SRC than effector cells. This

increased spare respiratory capacity in DP but not CD49a SP could imply different roles in memory

and effector responses in the lung post infection.

Based on the observed differences, we wanted to investigate whether nutrient uptake was altered in

a similar way. Glucose is a metabolically essential molecule in a variety of cellular processes as both

an eventual source of cell energy in the form of ATP as well as a carbon source in anabolism. To

evaluate the extent to which glucose was being taken up by the CD8 T cells, we utilized the glucose

analogue 2-NBDG. This fluorescent analogue undergoes facilitated diffusion through glucose

transporters GLUT1-4 as well as through active transport through SGLT1 and 2, recapitulating the

kinetics of glucose transport (14-19). It is phosphorylated by hexokinase after entering the cell, which

traps the fluorescent molecule in the cell, but cannot proceed further in glycolysis.

DP and CD49a SP displayed similar levels of glucose uptake 14 dpi, however, these populations

displayed lower levels of 2-NBDG staining compared with DN and CD103 SP cells, suggesting that

CD49a expressing subsets may be less glucose-dependent than CD49a negative counterparts

(**Figure 3E**). A similar pattern was observed in BAL samples, with CD49a^{pos} subsets taking up lower

levels of the dye. This pattern was conserved through later memory timepoints 60+ dpi (**Figure 3F**)

and are consistent with studies done on T$_{RM}$ in the skin, which noted an increased dependence on

free fatty acids (FFAs) as opposed to glucose (7) in T$_{RM}$.

Based on these findings, we hypothesized that lung T$_{RM}$ utilize a more lipid-centric metabolism. This

was originally hypothesized in skin due to its lipid-rich environment, though the lung is significantly

less lipid-rich (7). Having demonstrated significant differences in glucose uptake, we next investigated

the ability of T cells to take up free fatty acids. A fluorescently conjugated fatty acid (FFA) was utilized.

In the lung parenchyma, CD49a^{pos} subsets (CD49a SP and DP) took up the highest amount of FFA.

(**Figure 3G**). Similar results were found in BAL. These findings were similar but importantly distinct at

60+ dpi: DP took up more FFAs than CD49a SP which in turn took up equivalent FFAs as DN (**Figure**

3H). Other studies that have reported a dependence of T$_{RM}$ on FFAs have never used CD49a as a

marker (7). We expected our results to demonstrate that both DP and CD49a SP (the latter not studied

by other metabolic studies) were dependent on these FFAs in the lung, but we found this only to be

true at 14 dpi. At 60+ dpi, The CD49a SP were much more similar to DN, indicating an important

functional difference. From this data, we hypothesize that CD49a SP utilizes a less lipid centric

metabolic strategy to maintain residency than their DP counterparts.

We next wanted to determine if there were further differences in lipid metabolism. To address this,

Bodipy, a molecule that is taken up passively and stains cellular stores of neutral lipids, was used to

quantify the amount of neutral lipids present in lung T cells (20). Neutral lipids serve as a store of

potential energy as well as a carbon source, a valuable resource if the cells were to expand and

undergo effector functions in a nutrient deficient environment.

Nearly identical patterns for BAL and lung were observed 14 dpi (**Figure 3I, J, Figure S2**). DP cells

displayed the highest levels of neutral lipid storage, followed by CD49a SP, DN, with the lowest levels

in CD103 SP. While all six comparisons demonstrate that the four subsets are significantly different

from one another, it is remarkable that the subsets expressing CD49a (CD49a SP and DP) stain

brightly for neutral lipids while those not expressing CD49a (CD103 SP and DN) display low levels.

Having quick access to neutral lipids for both energy and carbon needs may prove important for their

function. Interestingly, the differences observed at 14 dpi were not seen at 60+ dpi.

3.4   Glucose-rich metabolic microenvironments alter differentiation of T_{RM}-like cells in vitro

Noting the myriad of differences between T_{RM} and non-T_{RM}, as well as among T_{RM} subsets, we

hypothesized that these effects may not only be phenotypic differences among these CD8 T cell

subsets, but that the metabolic microenvironment is deterministic in their differentiation. To test this,

T_{RM}-like T cells were generated *in vitro* so that that the metabolic microenvironment could be closely

controlled. T_{RM}-like cells can be generated *in vitro* stimulating naïve mouse splenocytes on anti-

CD3/CD28 coated plates for two days in the presence of IL-2 followed by three days in non-coated

plates supplemented with IL-2, and finally in IL-7 and TGF-β for six days to illicit a T_{RM}-like

phenotype, as previously described (21). Both the concentration of glucose as well as lipids were

varied (**Figure 4**) to test the effects given levels of glucose or lipids have as well as across levels.

First, the effect of glucose was examined within each lipid concentration, ranging from 0x to 8x

recommended supplementation (Millipore Sigma #L5146). Without exogenous lipid added, there

was no effect of glucose concentration on frequency, but with 2x or 4x lipid supplementation,

generally, as the concentration of glucose increased, the cell number normalized per volume and

frequency of CD49a positive cells both decreased (**Figures 5A and 5B**). The number of live CD8 T

cells for the 8x lipid supplementation group was effectively zero by this timepoint, and for the other

three groups the linear relationship between absolute number of CD49a positive cells per volume

and glucose concentration was significant for the 0x, 2x, and 4x lipid supplementation groups

(P=2.9e-6, 1.4e-9, 8.2e-6, R=-0.66, -0.79, and -0.64 respectively) (**Figure 6A**). In line with that

finding, the relationship between the frequency of CD49a positive cells and glucose concentration

was significant for both the 2x and 4x lipid supplementation groups (P=4.5e-13 and 1.3e-4, R= -0.87

and -0.57 respectively) (**Figure 6B**). Overall, this paints a picture of increasing glucose

concentration causing a decrease in both the frequency and absolute number of CD49a positive

cells. This is in contrast to the frequency of central memory marker CCR7 positivity (**Figures 6C and**

6D, 7A-7D), which was not significant for any lipid supplementation concentration, though increasing

concentrations of glucose did seem to decrease the number of CCR7 positive cells per microliter at

lipid concentrations 0x, 2x, and 4x.

When analyzing the effect of lipid on the differentiation of CD8 T cells *in vitro*, a contrasting pattern

emerged: within each concentration of glucose, increasing lipid concentration from 0x to 2x

increased the frequency of CD49a positive cells at the lowest and second lowest concentrations of

glucose tested (4.5 and 9.5 mg/mL) (**Figure 5C**), but further addition of lipid to 4x did not increase

the frequency of CD49a positive cells. The absolute number of CD49a positive cells per volume

peaked at 2x and decreased upon further addition of lipid for all concentrations of glucose tested. A

similar finding occurred for CCR7 positivity (**Figure 5D**).

Taken together, these results imply that high glucose concentration was detrimental to the long-term

viability or maintenance of CD8 T cells independent of lipid supplementation, and affects the

differentiation of CD49a positive cells, not CCR7 positive cells, at least in this assay. Furthermore,

lipid supplementation is beneficial up to a point, and then may become toxic as measured by the

number of live CD8 T cells in culture. It is beneficial for both CD49a and CCR7 positivity.

4   Discussion

We found memory T cell subsets in the lung exhibit distinct metabolic profiles. Furthermore,

experiments conducted to differentiate T_{RM} in the presence of varying metabolic substrates suggest

metabolite availability can influence memory T cell differentiation and survival. This availability may

be due local presence or lack of presence, the cell's ability to transport said metabolite into the cell,

or expression of receptor(s) to interact with these metabolites from outside the cell.

Differential expression, pathway, and metabolic modeling analyses of our RNAseq experiment led to

the hypotheses that lung T_{RM} have distinct metabolic programming compared with circulating cells.

Using both functional and descriptive methods, we produced a dataset for a variety of cell functions

including mitochondrial profiling, nutrient uptake, and more at both 14 and 60+ dpi. Synthesizing these

data together, what seems clear is that DP and CD49a SP, though different from each other in a few

assays, share a similar pattern. This pattern is distinct from DN, the subset presumably consisting of

T cells mostly with the phenotypes of effector memory and some central memory (5). DP and CD49a

SP both take up more fatty acids, store more neutral lipids, and take up less glucose than DN cells at

14 dpi. At later memory time points, DP and CD49a SP diverge: DP have more mitochondrial mass,

uptake less glucose, and uptake more fatty acid than CD49a SP. These changes from 14 to 60 dpi

could be a result from a change in the actual metabolic profile of these cells, a change in integrin

expression, cells dying, and/or cells leaving the tissue. These differences make the CD49a SP

population, well validated to be resident in the lung, an interesting subset of T_{RM} with respect to their

metabolism. These findings all correlate with a more "resting" memory phenotype for DP versus the

perhaps more "activated" CD49a SP. We have previously reported that the CD49a SP subset has a

number of features that correlate with a more effector-like phenotype such as having a higher

frequency of Granzyme B and perforin double positive cells, poised for cell killing (5).

The metabolic profile of an individual T cell as well as a T cell subset population are important

functionally, as we know many cell signaling molecules are limiting. Indeed, this is a leading

hypothesis as to why some T cells become T_{RM} and some do not (21). The functions of these subsets

during a subsequent infection may in part be achieved through the availability of different nutrients: if

T_{RM} depend mostly on uptake of free fatty acids whereas naive and effector T cells depend on glucose

uptake, both may be able to perform their functions more optimally with less competition for energy

resources existing between them. Furthermore, these profiles directly impact the cell signaling the

cells may be receiving. It is known that altering the ability for naïve CD8 T cells to take up glucose

directly impacts its future likelihood of becoming a memory cell (22). Having a metabolic profile that,

at baseline, is not dependent on glucose may ensure that the internal cell signaling conferring this

memory phenotype continues to do so in a positive feedback manner. Furthermore, upon infection,

the most glucose-starved T cells become memory cells (22). TCR stimulation is directly related to

specificity of binding and the abundance of a TCR's peptide:MHC complex. Others have shown that

T cells take up more glucose when more activated (23). Having an abundance of high-affinity

interactions with peptide:MHC complexes may cause a hyper local effect of glucose starvation. The

same group demonstrated that decreased availability of glucose led to increased memory CD8 T cell

populations. The fact that these T_{RM} and other memory cells seem to demonstrate lipid-centric

metabolism may assist naïve T cells to respond to infection because they don't compete with other

memory cells for glucose: if memory cells utilize one energy source (lipids), but naïve are more

dependent on another (glucose), it is less likely that memory T cells will starve out the naïve

responses, eliciting a more diverse group of memory T cells at the end of that infection, which would

increase the likelihood of a productive immune response upon a subsequent heterologous infection

since there is a more diverse and potentially cross-reactive set of T cells.

The causal effect of metabolic factors on CD8 T cell differentiation has been studied but never in the

context of T_{RM} differentiation. In short, we found glucose to be detrimental for the expression of T_{RM}

marker CD49a in a linear manner, whereas we found increasing lipid beneficial, to a point. Conversely,

glucose did not affect the expression of central memory marker CCR7, whereas the effect of lipids

was similar for both CD49a and CCR7: there exists an optimal concentration above and below which

decreases the expression of these markers. We believe these findings have direct implications for the

creation of therapies like CAR-T cells which necessitate *ex vivo* culture conditions: these conditions

should be optimized for the most efficacious T cell response which could include the optimization of

these culture conditions for the creation of T_{RM}, since tumor infiltrating lymphocytes have a T_{RM}-like

phenotype and are associated with positive outcomes (24-27). Future studies should assess this

possibility.

5   Figure Legends

Figure 1. Significantly up and downregulated Reactome and KEGG pathways in mouse lung

memory CD8 T cells 21 days post infection

(A) Reactome and KEGG databases were queried with the lists of differentially expressed genes

between DP and DN. (B) Reactome and KEGG databases were queried with the lists of differentially

expressed genes between CD49a SP and DN. Criteria for inclusion as differentially expressed genes

include P_{adj} < 0.05 and Log_2(Fold Change) > 0.5 or Log_2(Fold Change) < -0.5. DP. Adjusted P value

cutoff for enrichment is P_{adj}<0.05. (<u>D</u>ouble <u>P</u>ositive, CD49a^{pos}CD103^{pos}); DN (<u>D</u>ouble <u>N</u>egative,

CD49a^{neg}CD103^{neg}); CD49a SP (<u>CD49a</u> <u>S</u>ingle <u>P</u>ositive, CD49a^{pos}CD103^{neg}).

Figure 2. Fatty acid oxidation net fluxes for DP, CD49a SP, and DN models.

Metabolic models were created and beta oxidation of fatty acid net flux (mmols per gram dryweight

hours) levels were reported for each memory CD8 T cell subset. DP (<u>D</u>ouble <u>P</u>ositive,

CD49a^{pos}CD103^{pos}); DN (<u>D</u>ouble <u>N</u>egative, CD49a^{neg}CD103^{neg}); CD49a SP (<u>CD49a</u> <u>S</u>ingle <u>P</u>ositive,

CD49a^{pos}CD103^{neg}).

Figure 3. Normalized mouse IV-label-negative lung CD44^{pos} CD8 T cell MFIs from flow

cytometric assays.

451 Normalized MFIs plotted for each assay at 14 and 60+ days post infection. *P < 0.05, **P < 0.01, ***P

452 < 0.001, ****P < 0.0001. DP (<u>D</u>ouble <u>P</u>ositive, CD49a^{pos}CD103^{pos}); DN (<u>D</u>ouble <u>N</u>egative,

453 CD49a^{neg}CD103^{neg}); CD49a SP (<u>CD49a</u> <u>S</u>ingle <u>P</u>ositive, CD49a^{pos}CD103^{neg}) ; CD103 SP (<u>CD103</u>

454 <u>S</u>ingle <u>P</u>ositive, CD49a^{neg}CD103^{pos}).

Figure 4. Assay layout for T cell differentiation studies.

456 16 combinations (4 glucose treatment groups, 4 lipid treatment groups permuted) were assayed.

Figure 5. Metabolite concentration impact on T$_{RM}$-like cell differentation and maintenance *in*

***vitro*.**

459 Percent CD49a positive (A and C) and cell number per µL (B and D) in metabolite alteration assays

460 where both lipid (0x-8x) and glucose (4.5-24.5 mg/mL) were altered. *P < 0.05, **P < 0.01, ***P <

461 0.001, ****P < 0.0001. Small circles represent individual data points.

Figure 6. Simple linear regression modeling the effect of glucose concentration on CD49a and

CCR7 positivity.

464 Simple linear regression of (A) CD49a positive cells per µL ~ glucose concentration, (B) percent

465 CD49a positive ~ glucose concentration, (C) CCR7 positive cells per µL ~ glucose concentration, and

466 (D) percent CCR7 positive ~ glucose concentration. Small circles represent individual data points.

467 Large circles represent the mean of those for each glucose cocentration. Background represents 95%

468 confidence interval of the best fit line.

Figure 7. Metabolite concentration impact on T$_{CM}$-like cell differentation and maintenance *in vitro*.

Percent CCR7 positive (A and C) and cell number per μL (B and D) in metabolite alteration assays where both lipid (0x-8x) and glucose (4.5-24.5 mg/mL) were altered. *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, ****$P < 0.0001$. Small circles represent individual data points.

6   Author Contributions

MS designed and performed experiments, analyzed and interpreted data, and wrote the manuscript. AP assisted in conducting experiments and assay optimization. AG, KL, ECR, and TJ edited the manuscript. ECR and KL provided technical support. DJT provided overall direction, procured funding, interpreted data, and edited the manuscript.

7   Acknowledgments

We acknowledge the important contribution of the University of Rochester Medical Center Genomic Research Center, Flow Core, and Vivarium staff. In addition, we are gracious and acknowledge the supplying of reagents by Andrea Amitrano, PhD and Minsoo Kim, PhD.

8   Data Availability Statement

The datasets for this study can be found on GitHub (https://github.com/sportiellomike). Previously published RNAseq data sets were used and are available here: Geo with accession GSE179653 (https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE179653).

9   References

1.   Park SL, Buzzai A, Rautela J, Hor JL, Hochheiser K, Effern M, et al. Tissue-resident memory CD8(+) T cells promote melanoma-immune equilibrium in skin. Nature. 2019;565(7739):366-71.

2.   Park SL, Zaid A, Hor JL, Christo SN, Prier JE, Davies B, et al. Local proliferation maintains a stable pool of tissue-resident memory T cells after antiviral recall responses. Nat Immunol. 2018;19(2):183-91.

3.   Ray SJ, Franki SN, Pierce RH, Dimitrova S, Koteliansky V, Sprague AG, et al. The collagen
binding alpha1beta1 integrin VLA-1 regulates CD8 T cell-mediated immune protection against
heterologous influenza infection. Immunity. 2004;20(2):167-79.

4.   Jozwik A, Habibi MS, Paras A, Zhu J, Guvenel A, Dhariwal J, et al. RSV-specific airway
resident memory CD8+ T cells and differential disease severity after experimental human infection.
Nat Commun. 2015;6:10224.

5.   Reilly EC, Sportiello M, Emo KL, Amitrano AM, Jha R, Kumar ABR, et al. CD49a Identifies
Polyfunctional Memory CD8 T Cell Subsets that Persist in the Lungs After Influenza Infection. Front
Immunol. 2021;12:728669.

6.   Pan Y, Kupper TS. Metabolic Reprogramming and Longevity of Tissue-Resident Memory T
Cells. Front Immunol. 2018;9:1347.

7.   Pan Y, Tian T, Park CO, Lofftus SY, Mei S, Liu X, et al. Survival of tissue-resident memory
T cells requires exogenous lipid uptake and metabolism. Nature. 2017;543(7644):252-6.

8.   Becker SA, Palsson BO. Context-specific metabolic networks are consistent with
experiments. PLoS Comput Biol. 2008;4(5):e1000082.

9.   Han F, Li G, Dai S, Huang J. Genome-wide metabolic model to improve understanding of
CD4(+) T cell metabolism, immunometabolism and application in drug design. Mol Biosyst.
2016;12(2):431-43.

10.   Khodaee S, Asgari Y, Totonchi M, Karimi-Jafari MH. iMM1865: A New Reconstruction of
Mouse Genome-Scale Metabolic Model. Sci Rep. 2020;10(1):6177.

11.   Heirendt L, Arreckx S, Pfau T, Mendoza SN, Richelle A, Heinken A, et al. Creation and
analysis of biochemical constraint-based models using the COBRA Toolbox v.3.0. Nat Protoc.
2019;14(3):639-702.

12.   Presley AD, Fuller KM, Arriaga EA. MitoTracker Green labeling of mitochondrial proteins
and their subsequent analysis by capillary electrophoresis with laser-induced fluorescence detection.
J Chromatogr B Analyt Technol Biomed Life Sci. 2003;793(1):141-50.

13.   van der Windt Gerritje JW, Everts B, Chang C-H, Curtis Jonathan D, Freitas Tori C, Amiel E,
et al. Mitochondrial Respiratory Capacity Is a Critical Regulator of CD8+ T Cell Memory
Development. Immunity. 2012;36(1):68-78.

14.   Lu Y-T, Ma X-L, Xu Y-H, Hu J, Wang F, Qin W-Y, et al. A Fluorescent Glucose Transport
Assay for Screening SGLT2 Inhibitors in Endogenous SGLT2-Expressing HK-2 Cells. Nat Prod
Bioprospect. 2019;9(1):13-21.

15.   Fang P, Yu M, Min W, Wan D, Han S, Shan Y, et al. Effect of baicalin on GLUT4 expression
and glucose uptake in myotubes of rats. Life Sciences. 2018;196:156-61.

16.   Ferreira JM, Burnett AL, Rameau GA. Activity-dependent regulation of surface glucose
transporter-3. J Neurosci. 2011;31(6):1991-9.

17.   Marín-Juez R, Rovira M, Crespo D, van der Vaart M, Spaink HP, Planas JV. GLUT2-
mediated glucose uptake and availability are required for embryonic brain development in zebrafish.
J Cereb Blood Flow Metab. 2015;35(1):74-85.

18.   Meireles P, Sales-Dias J, Andrade CM, Mello-Vieira J, Mancio-Silva L, Simas JP, et al.
GLUT1-mediated glucose uptake plays a crucial role during Plasmodium hepatic infection. Cell
Microbiol. 2017;19(2).

19.   Blodgett AB, Kothinti RK, Kamyshko I, Petering DH, Kumar S, Tabatabai NM. A
fluorescence method for measurement of glucose transport in kidney cells. Diabetes Technol Ther.
2011;13(7):743-51.

20.   Rumin J, Bonnefond H, Saint-Jean B, Rouxel C, Sciandra A, Bernard O, et al. The use of
fluorescent Nile red and BODIPY for lipid measurement in microalgae. Biotechnology for Biofuels.
2015;8(1):42.

21.   Bromley SK, Akbaba H, Mani V, Mora-Buch R, Chasse AY, Sama A, et al. CD49a Regulates
Cutaneous Resident Memory CD8(+) T Cell Persistence and Response. Cell Rep.
2020;32(9):108085.

22.   Sukumar M, Liu J, Ji Y, Subramanian M, Crompton JG, Yu Z, et al. Inhibiting glycolytic
metabolism enhances CD8+ T cell memory and antitumor function. J Clin Invest.
2013;123(10):4479-88.

23.   Jacobs SR, Herman CE, Maciver NJ, Wofford JA, Wieman HL, Hammen JJ, et al. Glucose
uptake is limiting in T cell activation and requires CD28-mediated Akt-dependent and independent
pathways. J Immunol. 2008;180(7):4476-86.

24.   Loi S, Sirtaine N, Piette F, Salgado R, Viale G, Van Eenoo F, et al. Prognostic and predictive
value of tumor-infiltrating lymphocytes in a phase III randomized adjuvant breast cancer trial in
node-positive breast cancer comparing the addition of docetaxel to doxorubicin with doxorubicin-
based chemotherapy: BIG 02-98. J Clin Oncol. 2013;31(7):860-7.

25.   Denkert C, Loibl S, Noske A, Roller M, Muller BM, Komor M, et al. Tumor-associated
lymphocytes as an independent predictor of response to neoadjuvant chemotherapy in breast cancer. J
Clin Oncol. 2010;28(1):105-13.

26.   Wang R, Song Y, Hu T, Wang X, Jiang Y, Zhang D, et al. Decreased CD8(+) Lymphocytic
Infiltration in Multifocal and Multicentric Glioblastomas. Front Oncol. 2021;11:748277.

27.   Smazynski J, Webb JR. Resident Memory-Like Tumor-Infiltrating Lymphocytes (TILRM):
Latest Players in the Immuno-Oncology Repertoire. Front Immunol. 2018;9:1741.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7