1

OCA-B/Pou2af1 is sufficient to promote CD4$^+$ T cell memory and prospectively identifies memory precursors

Wenxiang Sun[1,2,†,¶], Erik P. Hughes[1,2,†], Heejoo Kim[1,2,†,Ψ], Jelena Perovanovic[1,2,§], Krystal R. Charley[1,2], Junhong Du[1,2], Andrea Ibarra[1,2], J. Scott Hale[1], Matthew A. Williams[1,2], Dean Tantin[1,2,3,*]

[1]Department of Pathology, University of Utah School of Medicine, Salt Lake City, UT 84112, U.S.A.
[2]Huntsman Cancer Institute, University of Utah School of Medicine, Salt Lake City, UT 84112, U.S.A.
[3]Lead contact

[*]Correspondence author: Dean Tantin
Email: dean.tantin@path.utah.edu

Author contributions:
M.A.W., D.T. designed research; W.S., E.P.H., H.K., K.R.C., J.D., A.I. performed research; J.S.H., M.A.W. contributed research tools; W.S., H.K., J.P., K.R.C. analyzed data; W.S., E.P.H., H.K., J.P., K.R.C., J.D., A.I., J.S.H., M.A.W., D.T. wrote the manuscript.

Competing interest statement: The authors declare that they have no competing interests.

Classification: Immunology

Keywords: Immunological memory ｜ CD4 T cells ｜ Pou2af1/OCA-B

[†]Equal contributors
[¶]Current address: Sumitomo Pharma Oncology, 3900 N. Traverse Mountain Blvd., Suite 100, Lehi, Utah 84043
[Ψ]Current address: Janssen Research & Development, LLC., Neuroimmunology Drug Discovery, 3210 Merryfield Row, San Diego, CA 92121.
[§]Current address: Recursion Pharmaceuticals, 41 400 W, Salt Lake City, UT 84101

Words in abstract: 162
Words in MS (excluding references): 6,471

Abstract

The molecular mechanisms leading to the establishment of immunological memory are inadequately understood, limiting the development of effective vaccines and durable anti-tumor immune therapies. Here we show that ectopic OCA-B expression is sufficient to drive T cells towards a memory fate, while having minimal effects on primary antiviral effector response. At primary effector response, short lived effector T cell compartments are expanded but show increased expression of *Gadd45b* and *Socs2*, while memory precursor effector cell clusters show increased expression of *Bcl2*, *Il7r* and *Tcf7* on a per cell basis, consistent with increased memory potential. Using an OCA-B mCherry reporter mouse line, we observe high OCA-B expression in CD4$^+$ T$_{CM}$ cells. We show that during viral infection, endogenously elevated OCA-B expression prospectively identifies memory precursor cells with increased survival capability and memory recall potential. Cumulatively, the results demonstrate that OCA-B is both necessary and sufficient to promote CD4 T cell memory in vivo and can be used to prospectively identify memory precursor cells.

Significance

CD4$^+$ T cell memory is incompletely defined and memory progenitors are difficult to identify. Here, we show that expression of the OCA-B transcription coactivator in CD4$^+$ T cells is necessary and sufficient to drive productive memory. Using a novel mCherry reporter mouse strain, we show that OCA-B expression can be used to enrich responding effector cells with elevated potential to form memory. The results show that OCA-B expression in T cells is sufficient to promote CD4$^+$ memory formation and marks memory precursor cells.

Introduction

The ability to respond robustly to secondary antigen exposures (immune memory) is a defining

feature of the adaptive immune system and forms the physiologic basis for vaccination (1). Immune

memory can also drive autoimmune and antitumor responses (2-5). Strategies that promote memory

have the potential to improve both vaccine efficacy and tumor immunotherapy. Central memory T

cells (T_{CM}), which home to secondary lymphoid organs, are particularly long lived and demonstrate

high proliferative capacity upon stimulation.

Multiple transcription factors and epigenetic regulators are known to modulate formation or

function of T_{CM}, including Blimp1 (6-8), Suv39h1 (9), T-bet (10, 11), Foxo1 (12, 13), Bcl6 (14, 15),

Eomes (16) and TCF1 (17). Many of these factors regulate memory less clearly in CD4$^+$ compared to

CD8$^+$ T cells. In CD4$^+$ T cells, the lymphocyte-restricted transcriptional coregulator OCA-B (also

known as OBF-1 and Pou2af1) is induced upon T cell antigen stimulation (18, 19). Germline OCA-B

deletion results in normal primary T cell responses to LCMV, but attenuates CD4$^+$ memory T cell

formation and recall response. Elevated *Ocab* (*Pou2af1*) mRNA levels are seen in CD4$^+$CD25^{lo} early

effector cells that are enriched for memory potential (20). OCA-B is also associated with autoimmunity

(21-24).

Here, we show that OCA-B deletion in T cells selectively reduces memory responses. We

show that ectopic OCA-B expression does not alter effector cell numbers at peak T cell response to

LCMV, but is sufficient to expand the memory recall effector pool at the expense of empty vector

(EV)-transduced controls. The effect becomes progressively stronger at later timepoints. mRNA

expression profiling of effector cells ectopically expressing OCA-B reveals gene expression changes

associated with effector and memory properties, signaling and homing. These include global

increases in *Socs2* and *Gadd45b*, expansion of short-lived terminal effector T cell populations, and

increases in *Tcf7* and *Il7r* in specific memory precursor populations. Using an OCA-B reporter mouse,

we show that elevated levels of endogenous OCA-B in effector T cells during the primary response

prospectively identifies CD4$^+$ T effector cells with enhanced survival and expansion capability

following viral rechallenge.

Results and Discussion

OCA-B in T cells is required for robust CD4$^+$ T cells memory recall responses

We used a conditional *Ocab* (*Pou2af1*) mouse allele crossed to a CD4-Cre driver, which efficiently

deletes OCA-B in T cells (24), to study effects of T cell-specific OCA-B loss. We infected *Ocab*$^{fl/fl}$;CD4-

Cre and control *Ocab*$^{fl/fl}$ mice with LCMV^{Arm} and measured specific T cell responses using I-A^b/gp$_{66-77}$

tetramer staining. No significant differences in the frequency of positive cells were observed at 8 days

post-primary infection (dpi, Fig. S1*A*). Rechallenging these mice with *Listeria monocytogenes*

expressing the LCMV glycoprotein epitope (Lm-gp61) (25) elicited a trending but nonsignificant

decrease in the frequency of CD4$^+$ I-A^b/gp$_{66-77}$$^+$ CD4$^+$ T cells undergoing memory recall response in

Ocab$^{fl/fl}$;CD4-Cre mice (Fig. S1*A*, D40+7). The responding cells showed little change in IL-2/IFNγ

double-producing CD4$^+$ T cells, but a significant decrease in IFNγ single-producers (Fig. S1*B*, *C*). The

same cells showed minimal changes in ICOS, PD-1, Ly6C or CD62L expression (Fig. S1*D*). These

results indicate that infection of T cell-conditional OCA-B knockout mice with LCMV results in

deficiencies in memory recall response IFNγ expression.

The relatively small reductions in frequencies of tetramer$^+$ CD4$^+$ T cells at recall response may

reflect selection for a small percentage of cells escaping OCA-B deletion. This selection can be

circumvented using bone marrow chimeras in which conditional deficient or control cells are competed

against wild-type T cells in the same mice. We generated bone marrow radiation chimeras in which

wild-type donors were co-engrafted with congenically-marked *Ocab$^{fl/fl}$*;CD4-Cre or control *Ocab$^{fl/fl}$* cells

in the same recipients. After 8 weeks to allow engraftment, recipient mice were infected with LCMV.

The ratio of antigen-reactive *Ocab$^{fl/fl}$* control effector CD4$^+$ T cells to wild-type competitor was close to

1:1 at 8 dpi. OCA-B deficient (*Ocab$^{fl/fl}$*;CD4-Cre) cells were slightly reduced relative to wild-type

competitor (Fig. S1*E*). To investigate the effects of T cell-specific OCA-B loss on memory recall

responses, we rechallenged mice with Lm-gp61. Following rechallenge, the control to wild-type cell

ratio remained close to 1:1 at both primary and memory recall response, while wild-type competitor

cells were greatly increased at the expense of OCA-B deficient cells specifically at recall. (>8-fold, Fig.

S1*E*, *F*). These findings indicate that, similar to T cell-conditional Oct1 deficiency and global OCA-B

deficiency (26), T cell-conditional OCA-B deficiency in bone marrow chimeras results in severe and

selective deficiencies in CD4$^+$ T cell memory recall responses.

Ectopic OCA-B expression is sufficient to enhance CD4$^+$ T cell memory responses in vivo

We transduced Thy1.1$^+$ SMARTA TCR transgenic donor CD4$^+$ T cells with a retroviral vector

expressing mouse OCA-B. In parallel, Ly5.1$^+$ (CD45.1$^+$) control SMARTA cells were transduced with

EV. SMARTA mice express a transgenic TCR recognizing an immunodominant LCMV glycoprotein

(GP$_{61-80}$) CD4$^+$ epitope (27). GFP$^+$ Thy1.1$^+$ SMARTA expressing OCA-B and GFP$^+$ Ly5.1 SMARTA

(EV) cells were sorted, mixed 1:1 and adoptively co-transferred into Thy1.2$^+$Ly5.2$^+$ recipients (Fig.

1*A*). Efficient OCA-B expression was confirmed using fibroblasts (Fig. 1*B*). After transfer, recipient

mice were infected with LCMV and the cell populations were detected by flow cytometry using GFP

expressed from MSCV and congenic markers from the donor T cells. The host T cells mounted

normal responses to LCMV, allowing SMARTA cells to be monitored in the context of a normal host

response. At 8 dpi, effector EV and OCA-B transduced cells expanded equivalently, maintaining the

same ratio in both spleen and peripheral blood (Fig. 1*C* and D). Another group of mice were infected

with LCMV and allowed to clear the virus and form memory. Recall responses were induced by re-

challenge using Lm-gp61. At peak T cell response to rechallenge (52+7 and 110+7 dpi), ~50-fold

more OCA-B overexpressing SMARTA cells were present (Fig. 1*C* and *D*). Ratios of percentages

from multiple mice are quantified in Fig. 1*E*. These results indicate that OCA-B expression enhances

the ability of CD4$^+$ T cells to induce memory responses.

130   At 8 dpi, GFP$^+$ SMARTA effector T cells were uniformly CD44^{hi}CD62L^{lo} with few changes

induced by ectopic OCA-B expression (Fig. 1*F*). ICOS and PD-1 levels were slightly diminished by

OCA-B overexpression. Ly6C was not significantly altered (Fig. 1*F* and *G*). To investigate if ectopic

OCA-B expression induces gene expression changes in effector cells that promote later memory

formation, we profiled gene expression using splenic SMARTA T cells at 8 dpi. CD4$^+$ cells were first

isolated by magnetic isolation, then Ly5.1$^+$/5.2$^+$ and GFP$^+$ cells were further isolated by FACS. We

identified ~600 down-regulated genes and ~200 up-regulated genes (Table S1). Unsupervised

hierarchical clustering identified differentially expressed gene clusters (Fig. 2*A*). Down-regulated

genes included *Sell*, which encodes CD62L, and genes specific to alternative CD4$^+$ lineages, such as

Bcl6 and *Foxp3*. The top up-regulated gene, *Pou2af1*, encodes OCA-B, providing a quality control.

Other up-regulated genes were associated with effector activity (*Tbx21*, *Zeb2*), exhaustion

(*Ikzf2*/*Helios*) and attenuation of T cell signaling and proliferation (*Socs2*, *Gadd45b*). *Gadd45b*

promotes autoimmune disease and anti-tumor immune responses (28-31). Example genome tracks

are shown in Fig. 2*B*. Gene ontology (GO) analysis of the up-regulated genes identified terms

associated with H3K9me3 regulation. Down-regulated genes by contrast were strongly associated

with H3K27me3 in CD4$^+$ and CD8$^+$ memory T cells (Fig. 2*C*). ChIP-X Enrichment Analysis (ChEA)

(32) identifies Tp53 and Smc3 as factors that also regulate the gene set induced by OCA-B ectopic

expression (Fig. 2*D*). p53 and SMC3 may therefore also control expression of these genes alongside

OCA-B, or could themselves be controlled by OCA-B to mediate gene induction via an indirect

mechanism. Sets of significant Molecular Function, Epigenomic Roadmap and ChEA GO terms are

shown in Table S2. These changes may be sufficient to confer potential increased memory formation,

as small advantages compounded over time can result in the large observed effects.

152   Two other genes associated with T cell survival and memory formation, *Il7r* and *Tcf7*, were

decreased in bulk RNA-seq. *Il7r* encodes a cytokine receptor critical for T cell survival, while Tcf7

encodes TCF1 which promotes naïve, Tfh and memory T cell transcriptional programs (11, 17).

Effector CD4$^+$ T cell populations are heterogeneous, with memory progenitor cells comprising a small

proportion of the effector T cell population (11, 20, 33). In bulk RNA-seq, icreased expression of

genes that promote memory in rare memory progenitors may therefore be obscured by decreases in

terminal effector cells. To test this, we performed single-cell RNA-sequencing (scRNA-seq) at D8

post-LCMV infection. Uniform manifold approximation and projection (UMAP) using EV-transduced

cells revealed a variety of populations (Fig. 3*A*, Table S3). Clusters 1 and 4 were enriched for genes

such as *Ccr7*, *Bcl2*, *Slamf6* and *Tcf7* (both clusters), *Tox* (cluster 1), and *Cxcr5*, *Il7r*, *Icos* and *Cd69*

(cluster 4, Table S3). These clusters likely contain memory precursors. Clusters 0, 2 and 6 by contrast

were enriched in *Ccl5*, *Ifng* and *Tbx21*, respectively (Table S3), and likely comprise terminal effectors.

Cluster 5 was enriched for genes encoding histones, cyclin-dependent kinases and proliferating cell

nuclear antigen (PCNA) and likely represent proliferating effectors.

OCA-B-transduction resulted in more uniform clustering with fewer and more tightly spaced

clusters identified using the same parameters (Fig. 3*B*). Cluster 0 effector cells were marked by *Zeb2*,

Klrg1 and *Tbx21* (Table S3). Clusters 1 and 2 were also associated with effector activity. OCA-B

transduced cluster 3 was associated with effector/memory progenitor activity similar to EV-transduced

clusters 1 and 4 (Fig. 3*C*). Although there were proportionally fewer of these cells compared to the

EV-transduced clusters 1 and 4, multiple memory-associated genes were increased on a per cell

basis. *Tcf7* was increased by >4-fold in OCA-B cluster 3 relative to other clusters, but only 2-fold in EV

clusters 1 and 4 (Table S3). *Slamf6* was increased by >3-fold in OCA-B cluster 3, but <2.5-fold in EV

clusters 1 and 4. *Bcl2* and *Il7r* were similarly more strongly expressed relative to the other clusters in

OCA-B- compared to EV-transduced cells. Additionally, *Foxo1* was not over-represented in either

memory-associated EV cluster but was enriched in OCA-B cluster 3, while *Ikzf2*, which has been

associated with CD4[+] T cell exhaustion (34), was not enriched in cluster 3 but was enriched in the

equivalent EV-transduced clusters (Table S3). Simultaneously, OCA-B-transduced clusters showed

broad increases in the expression of *Socs2*, which encodes an attenuator of T cell signal transduction,

and *Gadd45b*, which promotes growth arrest (Fig. 3*C*). These findings suggest that OCA-B

transduced cells in cluster 3 are qualitatively superior to their EV-transduced counterparts at forming

memory.

Clustering EV- and OCA-B-transduced cells together allows the relative proportions of cells

with common gene expression features to be determined. Again, specific clusters were enriched for

effector cells (Fig. 3*D*, clusters 0, 1, 2, 4, 6), proliferating effector cells (5) and effector cells with

memory progenitor potential (3). Feature plots for key genes are shown in Fig. 3*E*. In the OCA-B-

transduced condition, cluster 3 contained proportionally fewer cells but showed elevated expression

of genes such as *Il7r* and *Gadd45b* on a per cell basis (Fig. 3*F*). These findings identify changes in

gene expression and effector T cell populations consistent with increased memory potential.

OCA-B knock-in reporter mice label CD4[+] memory T cell populations

To monitor OCA-B expression in viable cells, we knocked 3 mCherry reporter cassettes preceded by

P2A elements into the mouse *Pou2af1* (*Ocab*) locus. The resulting reporter allele contained the

mCherry cassettes inserted immediately before the stop codon and the 3' UTR (Fig. 4*A*). A successful

hemizygous knock-in founder and a homozygote generated by intercrossing the progeny were

confirmed by PCR (Fig. S2*A*). Hemizygous and homozygous reporter mice retained normal OCA-B

protein expression (Fig. S2*B*). We used flow cytometry to identify peripheral blood mCherry-

expressing cells from hemizygous and homozygous reporter mice and wild-type controls. OCA-B was

readily detectable in B cells (Fig. S2*C, D*). Peripheral T cell expression was weaker, forming a

shoulder in hemizygous reporters, but a discrete peak in homozygotes (Fig. S2*C, D*). We therefore

used heterozygous mice to characterize B cell populations, and homozygous mice to study T cells.

Robust mCherry expression could be detected in the bone marrow, peritoneal cavity, spleen

and lymph nodes of specific pathogen-free hemizygous reporter mice. Total thymocytes by contrast

expressed little OCA-B (Fig. S2*E*). We used IgM and B220 to discriminate developing, immature and

recirculating bone marrow B cells. OCA-B expression was readily detected in developing, immature

and recirculating B cell populations compared to wild-type controls (Fig. S2*F*). Expression was nearly

uniform in the different populations, except for a small population of immature B cells that lacked

OCA-B expression. Peritoneal B-1 and B-2 B cells were also homogeneous, with stronger expression

in B-2 relative to B-1 cells (Fig. S2*G*). Splenic B cells can be partitioned into transitional and mature

populations based on CD93 (AA4.1) staining. T1/2/3 subsets stratified by CD23 and IgM uniformly

expressed OCA-B (Fig. S2*H*). CD21 and CD23 expression divides mature splenic B cells into

marginal zone (MZ), follicular and non-follicular populations, and all three populations expressed

OCA-B strongly (Fig. S2*I*). Expression was largely uniform, but there was a significant population of

newly formed $CD21^{lo}CD23^{lo}$ B cells expressing no OCA-B. The nature and significance of these cells

is unknown. These results document strong reporter activity in B cell populations, with the highest

level of activity in MZ and B-1 B cells. MZ B cells are largely absent in OCA-B knockout mice (35).

These results indicate that the OCA-B reporter robustly labels B cells.

Gating out B cells, approximately 5% of thymocytes from unchallenged, specific pathogen-free

homozygous reporter mice expressed OCA-B (Fig. S2*J*). mCherry expression could be identified in

CD4/CD8 double-negative (DN), DP and SP compartments, with the highest amounts in DN and CD8

SP cells (Fig. S2*K*). In the spleen, the relative abundance of $CD4^+$ and $CD8^+$ T cells was unaltered by

the homozygous reporter allele (Fig. 4*B*). Splenic $CD4^+$ T cells express mCherry at ~50-fold lower

levels compared to B cells and with a wider range of expression (Fig. 4*C*). Peripheral blood T cell

mCherry levels varied over a similar range (not shown).

We used CD62L and CD44 to stratify resting splenic $CD4^+$ and $CD8^+$ cells into naïve,

activated/effector and central memory compartments. Reporter activity could be detected in

approximately 7% of naïve ($CD62L^{hi}CD44^{lo}$) cells (Fig. 4*D*). The nature of these cells (e.g., Tregs,

Tscm, recent thymic emigrants) is unknown. Progressively larger fractions of early activated

($CD62L^{lo}CD44^{lo}$), activated ($CD62L^{lo}CD44^{hi}$) and T_{CM} ($CD62L^{hi}CD44^{hi}$) cells expressed mCherry (Fig.

4*D*). Naïve $CD8^+$ T cells showed comparatively higher expression levels, but decreased expression in

the activated state and similar expression in T_{CM} cells (Fig. 4*E*). Therefore, within the $CD4^+$ T cell

compartment, increased OCA-B expression as measured by mCherry fluorescence progressively

labels greater fractions of naïve, activated and central memory $CD4^+$ cells.

Elevated OCA-B expression prospectively enriches memory precursor CD4[+] T cells

To assess OCA-B responses during infection, memory formation and rechallenge, we infected reporter mice with LCMV and used MHC tetramers to identify splenic T cells recognizing immunodominant LCMV epitopes. Prior to infection, >90% of cells were naïve phenotype, with 13.7% of these mCherry[+]. ~50% of activated CD44[hi] cells expressed mCherry (Fig. S3*A*, D0). Similarly, 50% of 8 dpi effector cells expressed mCherry regardless of tetramer status (Fig. S3*B*, D8). After LCMV clearance, memory timepoint mice showed higher percentages of mCherry (D43, 61.6%), though with lower expression per cell. Rechallenge with Lm-gp61 to generate CD4[+] T cell antigen-specific recall responses reduced mCherry positivity compared to resting memory, and high-affinity tetramer[+] cells declining from >60% to <10% mCherry (Fig. S3*A*, D42+7). In contrast, a higher percentage of CD8[+] reporter T cells expressed mCherry prior to infection. Higher percentages of tetramer[+] mCherry[+] were also present at 5 dpi, but by 8 dpi (peak T cell response) mCherry was reduced (Fig. S3*B*). Antigen specific memory CD8[+] T cells also showed reduced mCherry levels relative to CD4[+] (Fig. S3*B*). We did not collect data on CD8[+] rechallenge because Lm-gp61 does not express the CD8-dominant epitope recognized by H2-D[b] LCMV tetramers.

To determine if OCA-B expression in responding CD4[+] T cells is associated with increased memory potential, we crossed the reporter allele to a SMARTA TCR transgenic background to fix the TCR specificity and to the Ly5.1 congenic marker to track engrafted cells. We transferred cells into recipients, infected with LCMV, isolated mCherry[hi] and mCherry[lo] cells at 8 dpi, transferred large numbers of cells (8×10^5) into naïve secondary recipient mice, and isolated splenic CD4[+] T cells to assess contraction (Fig. 5*A*). More cells were recovered from mice engrafted with mCherry[hi] cells, indicating that these cells contracted less than mCherry[lo] controls (Fig. 5*B*, *C*). Interestingly the cells that did survive showed similar mCherry fluorescence (Fig. 5*D*). These findings indicate that effector CD4[+] cells expressing higher OCA-B undergo less cell death when transferred into naïve animals.

To test the ability of the cells expressing OCA-B to mount antigen recall responses, we similarly transferred congenically marked SMARTA cells into recipient mice and infected with LCMV, but isolated mCherry[hi] and mCherry[lo] cells at 43 dpi. 4×10^4 of memory SMARTA cells were transferred into naïve hosts, which were challenged with LCMV and analyzed at 7 dpi (Fig. 5*E*). mCherry[hi] SMARTA cells mounted better recall responses compared to mCherry[lo] (Fig. 5*F*, *G*) though again with little difference in mCherry fluorescence intensity (Fig. 5*H*). This result indicates a plasticity in OCA-B expression whereby expression normalizes following antigen re-encounter.

In mice, OCA-B expression can be detected, varying over a 10-fold range, in responding cells as early as one day after infection with lymphocytic choriomeningitis virus (LCMV) (26). Memory progenitor cells characterized by low CD25 expression can be identified as early as early as 3 dpi (14,

20). We transferred SMARTA homozygous OCA-B mCherry reporter T cells into congenic recipients,

infected with LCMV, and collected spleens at 3 dpi to study mCherry and CD25 levels (Fig. 5*I*). A

large fraction of responding CD25^{lo} SMARTA cells expressed mCherry (Fig. 5*J*). SMARTA cells from

the same mouse expressing low levels of CD25 also expressed high levels of TCF1, as expected

(20). mCherry and TCF1 were not compared as cell fixation quenches mCherry, however we were

able to directly compare CD25 and OCA-B. CD25^{hi} cells expressed a range of OCA-B, while CD25^{lo}

cells were highly enriched for mCherry^{hi} cells (Fig. 5*K*, *L*). These findings show that high OCA-B

expression can be used as a method to prospectively mark live CD4^{+} memory precursor effector T

cells with augmented capacity to form central memory. Identifying transcriptional regulators that define

the memory precursor effector CD4^{+} T cells will promote advances in understanding the formation of

CD4 T cell memory. Cumulatively, this study indicates that OCA-B expression within CD4^{+} T cells is

both necessary for, and sufficient to promote, the emergence of T$_{CM}$.

Materials and methods

Mice

All mice used in this study were on the C57BL6/J strain background. All mouse experiments were approved by the University of Utah Institutional Animal Care and Use Committee (protocol 00001553). *Pou2af1* (*Ocab*) conditional (floxed) mice were described previously (24). *Pou2af1*-3×mCherry reporter knock-in mice were generated on a C57BL/6N background (Biocytogen), but were backcrossed >5 times to C57BL/6J prior to generation of data herein.

LCMV and *Listeria* infection

LCMV Armstrong 53b (LCMV[Arm]) (36) was grown in BHK cells and titered using Vero cells (27). For primary infection, 8-to 12-week-old *Ocab*[fl/fl] or *Ocab*[fl/fl];CD4-Cre mice were inoculated i.p. with 2×10^5 plaque-forming units (PFU) of LCMV intraperitoneally.

For heterologous rechallenge with *Listeria monocytogenes* expressing LCMV glycoprotein 61-80 (Lm-gp61) (25), bacteria were grown in log phase in BHI media, and concentrations were calculated using OD at 600 nm (OD of $1=1 \times 10^9$ CFU/mL). Mice were rechallenged intravenously with 2×10^5 colony forming units (CFU) of Lm-gp61 the indicated number of days after primary infection as published (37).

Bone marrow radiation chimeras

Radiation chimeras were generated as published (38). Briefly, male Ly5.1 C57BL/6J mice were used as recipients. Mice received a split dose of 900 Rad (2×450 RAD, spaced 1 hr apart), and were engrafted one day later with 1 million mixed donor bone marrow cells. Competitor bone marrow cells from Ly5.2/Thy1.1 but otherwise wild-type mice, and experimental cells from Ly5.2/Thy1.2 *Ocab*[fl/fl] or littermate *Ocab*[fl/fl];CD4-Cre mice, were isolated from femurs and tibias of donor mice. After red blood cell lysis, CD3[+] cells were depleted using biotinylated anti-CD3 antibodies (eBioScience) and anti-biotin magnetic MicroBeads (Miltentyi Biotech) according to the manufacturer's instructions. CD3-depeleted bone marrow cells were mixed 1:1 and injected retro-orbitally. The chimeras were rested for 8 weeks to allow engraftment, after which mice were infected as above. Virus-reactive cells were identified by LCMVgp$_{66-77}$:I-Ab tetramer staining.

pMSCV-IRES-GFP-OCA-B vector generation

Wild-type *Ocab* cDNA was amplified from the plasmid pcDNA3.1-OCA-B (39) (a gift from Robert Roeder, Rockefeller University) by PCR and cloned into the pMSCV-IRES-GFP (pMIGR1) retroviral vector (a gift from Ryan O'Connell, University of Utah). Primers used for amplification were mOcab-F-XhoI, 5'CTCGAGCTGTCTGCTTCAAAGAGAAAAGGCAAC; mOcab-R-*Eco*RI, 5'

GAATTCCTAAAAGCCCTCCACGGAGAGGGT. The PCR product was gel purified, digested with

*Eco*RI and *Xho*I, and inserted into a similarly digested pMIGR1 backbone using T4 DNA ligase to

generate pMIGR1-OCA-B. All constructs were validated by resequencing.

CD4$^+$ T cell isolation and stimulation

Naïve splenic CD4$^+$ T cells were isolated and simulated as described previously (40). Briefly, spleens

were dissociated by grinding and passing through a 70 μm nylon strainer. Red blood cells were lysed

by ACK lysis buffer (150 mM NH$_4$Cl, 10 mM KHCO$_3$, 0.1 mM EDTA). Cells were isolated from mice

using a naïve CD4$^+$ T cell isolation kit (Miltenyi), and stimulated in culture using 10 µg/mL plate-bound

anti-CD3ε and 2 µg/mL anti-CD28 antibodies (eBioscience).

Ectopic OCA-B expression in CD4$^+$ T cells

T cells from Ly5.1 SMARTA and Thy1.1 SMARTA mice were purified using a CD4$^+$ T cell isolation kit

(Miltenyi). Cells were placed into RPMI 1640 medium supplemented with 10% fetal bovine serum, 50

U/mL penicillin, 50 μg/mL streptomycin, 2 mM L-glutamine, 1 mM sodium pyruvate, 1× MEM

nonessential amino acids, 55 μM 2-mercaptoethanol, and 20 IU/mL recombinant human IL-2. All

components were from ThermoFisher with the exception of IL-2, which was supplied by R&D

Systems. Cells were simultaneously transduced with pMIGR retroviruses packaged in 239T cells, with

or without expressed mouse OCA-B cDNA, using spin infection. For spin infection, cells were

centrifuged with 293T retroviral supernatant at 1000×g for 2 hr at 37°C in the presence of 4 µg/mL

polybrene (Sigma). On the following day, transduced GFP$^+$ cells were isolated by fluorescent-

activated cell sorting. 2×10^4 pMIGR1- and pMIGR1-OCA-B-transduced cells were combined and co-

transferred into wild-type (Ly5.2/Thy1.2) C57BL6/J mice. Chimeric mice were infected with LCMV and

re-challenged with Lm-gp61 as described above.

Flow cytometry

Spleens were dissociated and passed through a 70 μm nylon strainer. Red blood cells were lysed by

ACK lysis buffer (150 mM NH$_4$Cl, 10 mM KHCO$_3$, 0.1 mM EDTA). For intracellular staining, cell

suspensions in RPMI supplemented with 10% fetal bovine serum were re-stimulated for 4 hr with 1

µM LCMV GP$_{61-80}$ or 0.1 µM GP$_{33-41}$ peptide along with Brefeldin A (GolgiPlug Becton-Dickinson, 1

µl/mL) as published (41, 42). Cells were subsequently fixed by cell fixation/permeabilization solution

(Cytofix/Cytoperm, Becton-Dickinson) according to the manufacturer's protocol. For tetramer staining,

APC-conjugated gp$_{66-77}$:I-A^b and gp$_{33-41}$:H-2D^b tetramers were provided by the NIH Tetramer Core

Facility (Emory Vaccine Center). Cell suspensions were incubated at 37°C for 3 hr in RPMI with the

tetramer followed by cell-surface staining. Tetramer fluorescence was normalized with isotype

tetramer (hCLIP:I-A^b) staining. Antibodies used for flow cytometry are presented in the Key Resources

Table.

Bulk RNAseq

CD4$^+$ T cells were purified from female Ly5.1 SMARTA and Ly5.1/5.2 SMARTA mice, and separately

transduced with pMIGR1-OCA-B or pMIGR1 (EV), respectively. On the following day, transduced

GFP$^+$ cells were sorted, 20,000 of each combined 1:1, and injected intravenously into 24 male

C57BL/6J male recipients. 24h later, each recipient mouse was injected intraperitoneally with 2x10^5

pfu LCMV. After eight days, the 24 mice were divided into six groups of four mice by combining four

spleens together. CD4$^+$ T cells were purified using a Miltenyi CD4 T cell isolation kit. Cells then were

stained with PerCP-Cy5.5 Ly5.1 and AF700 Ly5.2 antibodies. GFP$^+$Ly5.1$^+$Ly5.2$^+$ (EV) and

GFP$^+$Ly5.1$^+$ (OCA-B-expressing) cells were sorted by fluorescence-activated cell sorting (FACS) and

used for RNA purification (RNeasy Mini Kit, QIAGEN). RNA concentrations were determined using a

Quant-iT RNA assay kit and a Qubit fluorometer (ThermoFisher). Because limited RNA from EV-

transduced cells was obtained, the six EV samples were combined into three samples. Five OCA-B-

transduced samples (each comprising four mice) were submitted for RNAseq analysis. These showed

a high degree of concordance (not shown). Total RNA samples (200-500 ng) were hybridized with

Ribo-Zero Plus (Illumina) to deplete cytoplasmic and mitochondrial and ribosomal RNA from samples.

RNA sequencing libraries were prepared as described using the Stranded Total RNA Prep, Ligation

with Ribo-Zero Plus kit (Illumina) and IDT for Illumina RNA UD Indexes Set A, Ligation (Illumina).

Purified libraries were qualified on an Agilent Technologies 2200 TapeStation using a D1000

ScreenTape assay. The molarity of adapter-modified molecules was defined by qPCR using the Kapa

Biosystems Kapa Library Quant Kit. Individual libraries were normalized to 0.65 nM in preparation for

Illumina sequence analysis. Sequencing libraries were chemically denatured and applied to an

Illumina NovaSeq flow cell using the NovaSeq XP workflow (20043131). Following transfer of the flow

cell to an Illumina NovaSeq 6000 instrument, a 150 x 150 cycle paired-end sequence run was

performed using a NovaSeq 6000 S4 reagent Kit v1.5 (Illlumina).

Bulk RNA-seq analysis

Bulk RNA-seq analysis was performed as previously described (43). Briefly, Reads were aligned to

Mm10 using STAR (v2.7.3a) and checked for quality using multiqc (v1.10). Between 12 and 14 million

pair-end reads were generated for each sample, with >98% of aligned reads mapping to the correct

strand and >93% of the reads uniquely aligned to the gene. Differentially expressed genes were

identified using DESeq2 version 1.24.0 (44) with a 5% FDR cutoff. Features with zero counts and 5 or

fewer reads in every sample were removed from the analysis. Genes increased by 1.4-fold or more or

decreased by 2-fold or more, and with adjusted *p*<0.05 were selected as differentially expressed (0.5

< log2FC < -1; padj ≤ 0.05). Figures were generated in R version 4.0.0 using functions from ggplots

libraries and pheatmap.

Single-cell RNA-seq

To profile gene expression at the single-cell level, splenic GFP+Ly5.1+Ly5.2+ (EV-transduced) and

GFP+Ly5.1+ (OCA-B-transduced) SMARTA cells from four pooled mice per condition were isolated,

resuspended in PBS with 0.04% bovine serum albumin (ThermoFisher), and filtered through 40 μm

strainers. Viability and cell count were assessed using a Countess II (ThermoFisher). Equilibrium to

targeted cell recovery of 6,000 cells along with Gel Beads and reverse transcription reagents were

loaded to Chromium Single Cell A to form Gel-bead-in Emulsions (GEMs). Within individual GEMs,

cDNA generated from captured and barcoded mRNA was synthesized by reverse transcription at

53°C for 45 min. Samples were then heated to 85°C for 5 min. Single cell transcriptomes were

assessed using a 10X Genomics Chromium Single Cell Gene Expression instrument. Individual cells

were tagged with 16 bp barcodes, and specific transcripts with 10 bp Unique Molecular Identifiers

(UMIs) according to manufacturer instructions.

Single-cell RNA-seq analysis

Single cell transcriptome data were analyzed and clustered as described previously (43). Sequences

from the Chromium platform were de-multiplexed and aligned using CellRanger ver. 3.1.0 (10X

Genomics) with default parameters mm10-3.0.0. Clustering, filtering, variable gene selection and

dimensionality reduction were performed using Seurat ver.4.0.4 (45) according to the following

workflow: 1, Cells with <200 detected genes were excluded further analysis. 2, Cells with <5% UMIs

mapping to mitochondrial genes and *Cd44* expression >0 were retained for downstream analysis, as

SMARTA cells are uniformly responding to LCMV glycoprotein antigen at this time. 3, The UMI counts

per ten thousand were log-normalized for each cell using the natural logarithm. 4, Variable genes

(2000 features) were selected using the FindVariableFeatures function. 5, Common anchors between

the conditions were identified using FindIntegrationAnchors function that were further used to

integrate these sets. 6, Gene expression levels in the integrated set were scaled along each gene and

linear dimensional reduction was performed. The number of principal components was decided

through the assessment of statistical plots (JackStrawPlot and ElbowPlot). 7, Cells were clustered

using a shared nearest neighbor (SNN) modularity optimization-based clustering algorithm and

visualized using two-dimensional uniform manifold approximation and projection (UMAP). 8, One

cluster in each condition marked predominantly by mitochondrial genes (indicative of dying cells) was

excluded from the analysis. This resulted in 12,886 EV-transduced cells with 45,748 mean reads per

cell, and 2,595 median genes per cell. The total number of reads was 589,513,069, with 59.6%

mapping to exonic regions. Similarly, there were 11,709 OCA-B-transduced cells with 68,931 mean

reads per cell, and 2,668 median genes per cell. Total number of reads was 807,111,486, with 57.4%

mapping to exonic regions. Clusters were identified using manual interrogation of gene expression

enrichment for each population (Table S3), the ImmGene cluster identity predictor (Table S4, S5), and

PanglaoDB annotation terms. Clusters comprised of contaminating macrophages and neutrophils

were manually deleted.

Memory progenitor isolation, normalization and transfer

To assess contraction of the effector cells with different OCA-B levels, splenic CD4$^+$ T cells were

purified from 8 week-old female Ly5.1/5.2 homozygous knock-in reporter, SMARTA TCR transgenic

mice. 2×10^4 SMARTA cells were intravenously injected into 15 male Ly5.2/5.2 C57BL/6J recipients,

which were infected intraperitoneally 1 day later with 2×10^5 plaque-forming units of LCMV in 250 μL

phosphate-buffered saline. After 8 days, responding Ly5.1$^+$5.2$^+$ SMARTA T cells were sorted into

mCherry^{hi} and mCherry^{lo} populations. Cells were pooled and 8×10^5 mCherry^{hi} and mCherry^{lo} cells

were separately injected intravenously into 7 naïve age-matched 9 week-old male C57BL/6J

secondary recipients in the case of mCherry^{lo} and 5 secondary recipients in the case of mCherry^{hi}.

After 8 additional days, mice were sacrificed and splenic Ly5.1$^+$5.2$^+$ SMARTA cells were counted by

flow cytometry. To assess proliferative responses to rechallenge, 15 similar virus-naïve secondary

recipients were rested for 43 days post-infection prior to isolation of mCherry^{hi} and mCherry^{lo}

populations. 1.3×10^4 SMARTA cells were transferred into naïve recipients, which were infected with

LCMV one day later. 7 days post-infection, splenic CD4$^+$Ly5.1$^+$5.2$^+$ SMARTA cells were evaluated by

flow cytometry. Mice with poor infection/response (total splenocyte count $<6\times10^7$) were excluded from

the analysis.

Quantification and statistical analysis

Excel (Microsoft) and Prism (GraphPad) were used for statistics and graphing. Two-tailed Student T-

tests were used to ascribe statistical significance unless otherwise indicated. For all figures, *=*p*-

value≤0.05; **=*p*≤0.005; ***=p<0.001. All error bars denote ± standard error of the mean.

450 **Acknowledgements:** We thank M. Chandrasekharan for help with graphics. We thank J. Marvin and

451 the University of Utah Health Sciences Center Flow Cytometry Core facility. We thank O. Allen, B.

452 Dalley and the High-Throughput Genomics Core. MHC tetramers were provided by the NIH Tetramer

453 Core Facility (Atlanta, GA). We thank R. Roeder for the mouse OCA-B cDNA and R. M. O'Connell for

454 the pMIGR1 plasmid.

456 **Funding:** This work was supported by NIH grant R01AI100873 to DT.

460 **Data and materials availability:** All data are available in the main text or the supplementary

461 materials. Materials used in the study are available to any researcher for purposes of reproducing or

462 extending the findings. Bulk and single-cell RNA-seq data have been deposited at the Gene

463 Expression Omnibus GEO) website (https://www.ncbi.nlm.nih.gov/geo/, GSE214310).

 Figure legends

Fig. 1. Ectopic OCA-B expression enhances CD4+ T cell memory recall responses in vivo. (*A*)

Experimental schematic for OCA-B transduction and T cell mixed adoptive transfer. Ly5.1$^+$ SMARTA

cells were transduced with pMSCV-IRES-GFP (pMIGR1, EV) and Thy1.1$^+$ SMARTA cells were

transduced with pMIGR1 expressing mouse OCA-B (pMIGR1-OCA-B). The donors were age- and

sex-matched littermates. The next day following transduction, GFP$^+$Ly5.1$^+$ SMARTA (EV) cells and

GFP$^+$Thy1.1$^+$ SMARTA expressing OCA-B were sorted, combined 1:1, co-transferred into

Ly5.2$^+$/Thy1.2$^+$ C57BL6/J recipient mice, and infected with LCMV. Recipients were rechallenged with

Lm-gp61 on either day 52 or 110 post-LCMV infection, and in each case analyzed after 7 days. (*B*)

Lysates from NIH-3T3 cells transduced with either empty pMIGR1 (EV) or pMIGR1-OCA-B were

immunoblotted using antibodies against OCA-B. α-tubulin is shown as an internal loading standard.

(*C*) Flow cytometric analysis of Ly5.1$^+$ pMIGR1 EV-transduced and Thy1.1$^+$ pMIGR1-OCA-B-

transduced (OCA-B) SMARTA cells in the spleen and blood of a representative recipient mouse at

peak effector response (D8), peak memory recall response (52+7) and long-term memory recall

(D110+7). Cells were first gated based on CD44, CD4 and GFP positivity. (*D*) Mean relative

percentages of splenic and blood total EV- or pMIGR1-OCA-B-transduced (GFP$^+$) SMARTA cells

were plotted at D8, D52+7 and 110+7. Each datapoint represents an individual mouse harboring both

EV- and pMIGR1-OCA-B-transduced cells. 5, 7 and 10 mice were sacrificed at each timepoint to

obtain measurements from spleen. All mice were bled at D8, and remaining mice were bled at

subsequent timepoints. (*E*) Mean ratios of pMIGR1-OCA-B-transduced relative to EV-transduced

SMARTA cells (OCA-B/EV) are plotted. Splenic (top) and peripheral blood (bottom) SMARTA T cells

are shown at day 8 post-infection, or 7 days post-rechallenge with Lm-gp61 (52+7 or 110+7). (*F*) Flow

cytometry showing relative percentages of splenic GFP$^+$CD44^{hi} EV-transduced (blue) or pMIGR1-

OCAB-transduced (red) SMARTA cells expressing ICOS, PD-1, Ly6C or CD62L at day 8 post-

infection. Representative data showing the two populations of cells within a single mouse are shown

at day 8 post-infection. (*G*) Percent expression for ICOS, PD-1, CD62L and Ly6C was quantified from

four mice using splenic EV- and pMIGR1-OCAB-transduced GFP$^+$CD44^{hi} SMARTA cells at 8 days

post-infection. Four replicates were performed. For CD26L, the four datapoints are superimposed,

and for Ly6C two of the datapoints are superimposed. Individual mice are connected by lines.

Fig. 2. Gene expression changes associated with ectopic OCA-B expression in SMARTA transgenic

primary effector T cells. (*A*) Heatmap showing top up- and down-regulated genes. Genes with log2

fold-change >0.5 or <-1.0, and padj $\leq$ 0.05 are shown. Example genes are shown at right. Control

and OCA-B-transduced cells were purified from the same mice using magnetic isolation and FACS.

(*B*) Example genome tracks showing two up-regulated genes, *Socs2* and *Gadd45b*. *Pou2af1* (*Ocab*)

is shown as a positive control. (*C*) Top Epigenomic roadmap and Molecular function GO terms for the

set of OCA-B up- and down-regulated genes. (*D*) Top ChIP-X Enrichment Analysis (ChEA 2016)

factors potentially regulating the same set of genes up- and down-regulated by OCA-B ectopic

expression.

Fig. 3. Single-cell RNA-seq analysis of OCA-B-transduced vs. control SMARTA cells at 8 days post

LCMV infection. (*A*) UMAP projections of independently clustered EV-transduced (GFP$^+$) SMARTA

cells purified similar to Figs. 2 and 3, and subjected to scRNA-seq. Cluster identities were annotated

using a combination of gene expression enrichment (Table S3), cluster identity predictor ImmGen

identity scores (Table S4) and PanglaoDB annotation terms. Clusters 1 and 4 were associated with

memory formation. Top PanglaoDB annotation terms for cluster 4 are shown below. (*B*) Similar UMAP

projection using OCA-B-transduced cells purified from the same mice as in (A). Cluster identities were

annotated similarly to (A). Cluster 3 in OCA-B-transduced cells was most strongly associated with

memory formation. Top PanglaoDB annotation terms for cluster 3 are shown below.

(*C*) Feature plots highlighting expression of representative genes for the two UMAP projections in (A)

and (B). (*D*) UMAP projections of the the EV- and OCA-B-transduced single-cell RNA-seq datasets

clustered together. For each projection, the percentage of each cluster relative to the total number

cells in condition is shown. Cluster identities were annotated using a combination of gene expression

enrichment (Table S3), cluster identity predictor ImmGen identity scores (Table S5) and PanglaoDB

annotation terms. (*E*) Feature plots highlighting expression of representative genes for the combined

projections. (*F*) Violin plots depicting expression of six genes (*Il7r*, *Zeb2*, *Gadd45b*, *Foxo1*, *Tcf7* and

Tox) across each of the 10 clusters in EV-transduced cells (blue, left) and OCA-B-transduced cells

(red, right).

Fig. 4. mCherry expression in splenic T cell populations in specific pathogen-free homozygous OCA-

B-3×mCherry mice. (*A*) Mouse *Pou2af1* locus, targeting vector and the targeted *Pou2af1* locus. (*B*)

CD4 and CD8 expression are shown for an allelic series of example wild-type (+/+) and homozygous

(KI/KI) OCA-B knock-in reporter mice. (*C*) OCA-B homozygous reporter expression in splenic CD4$^+$ T

cell populations. (*D*) The same CD4$^+$ cells as in (C) were further stratified by CD62L and CD44 into

naïve, early activated, activated and central memory (Tcm) populations. CD62L^{lo}CD44$_{lo}$ cells are likely

recently activated cells that have down-regulated CD62L but not yet up-regulated CD44. (*E*) Similar

analysis as in (D) except for CD8$^+$ cells.

Fig. 5. CD4$^+$ T cells expressing high levels of OCA-B reporter activity preferentially form central

memory cells. (*A*) Schematic for assessing contraction of mCherry^{hi} vs. mCherry^{lo} SMARTA T cells

following LCMV infection. Congenically marked SMARTA T cells were isolated from donor mice,

transferred into naïve secondary recipients and infected with LCMV. At 8 days post-infection,

mCherry^{hi} and mCherry^{lo} populations were isolated by FACS. 8×10^5 SMARTA cells were transferred

into naïve recipients. After 8 days, splenic CD4$^+$Ly5.1$^+$5.2$^+$ SMARTA cells were evaluated by flow

cytometry. (*B*) Flow cytometric analysis of SMARTA donor T cells engrafted into naïve mice to monitor

rates of decline. Representative mice engrafted with mCherry^{lo} (left) or mCherry^{hi} (right) cells are

shown. (*C*) Quantification of averaged donor T cell numbers from mice engrafted with mCherry^{lo} or

mCherry^{hi} cells. N=7 for the mCherry^{lo} group and N=5 for the mCherry^{hi} group. (*D*) Similar analysis to

(C) except quantifying mean mCherry fluorescence intensity. (*E*) Schematic for assessing recall

responses of mCherry^{hi} vs. mCherry^{lo} SMARTA T cells following LCMV infection. Congenically

marked SMARTA T cells were isolated from donor mice, transferred into naïve secondary recipients

and infected with LCMV. However, mice were allowed to clear LCMV and form memory. After 43

days, mCherry^{hi} and mCherry^{lo} memory T cell populations were isolated by FACS. 1.3×10^4 SMARTA

cells were transferred into naïve recipients, which were infected with LCMV one day later. 7 days

post-infection, splenic CD4$^+$Ly5.1$^+$5.2$^+$ SMARTA cells were evaluated by flow cytometry. (*F*) Flow

cytometric analysis of SMARTA donor T cells 7 days post-rechallenge to monitor recall responses.

Representative mice engrafted with mCherry^{lo} (left) or mCherry^{hi} (right) cells are shown. (*G*)

Quantification of averaged donor T cell percentages and numbers from mice engrafted with mCherry^{lo}

or mCherry^{hi} cells. N=7 for the mCherry^{lo} group and N=5 for the mCherry^{hi} group. (*H*) Similar analysis

to (G) except quantifying mean mCherry fluorescence intensity. (*I*) Schematic for assessing mCherry

levels in SMARTA T at early timepoints following LCMV infection. 2×10^5 congenically marked

(CD45.1$^+$) SMARTA T cells were isolated from donor mice and transferred into naïve secondary

recipients. One day later mice were intraperitoneally infected with 2×10^5 PFU LCMV. Splenic T cells

were collected at 3 dpi. (*J*) Gated unfixed splenic CD4$^+$ T cells were assessed for the SMARTA

congenic marker Ly5.1 and mCherry (left), while fixed Ly5.1$^+$ (SMARTA) cells from the same mice

were used to assess CD25 and TCF1 (right). (*K*) CD25 and mCherry expression were assessed from

a representative animal. Left: distribution of SMARTA cell CD25 and mCherry expression in SMARTA

mice. Right: CD25 levels of gated mCherry^{hi} and mCherry^{lo} SMARTA cells displayed as a

concatenated histogram. (*L*) Quantification of averaged mCherry MFI in CD25^{hi} and CD25^{lo} cells (left),

and frequencies (center) and numbers (right) of CD25^{lo} cells in mCherry^{lo} or mCherry^{hi} cells SMARTA

T cells. N=6 independent recipient mice.

References and Notes

1. F. Sallusto, A. Lanzavecchia, K. Araki, R. Ahmed, From vaccines to memory and back. *Immunity* **33**, 451-463 (2010).

2. S. V. Gearty *et al.*, An autoimmune stem-like CD8 T cell population drives type 1 diabetes. *Nature* **602**, 156-161 (2022).

3. A. Schnell *et al.*, Stem-like intestinal Th17 cells give rise to pathogenic effector T cells during autoimmunity. *Cell* **184**, 6281-6298 e6223 (2021).

4. P. C. Rosato *et al.*, Virus-specific memory T cells populate tumors and can be repurposed for tumor immunotherapy. *Nat Commun* **10**, 567 (2019).

5. G. Cafri *et al.*, Memory T cells targeting oncogenic mutations detected in peripheral blood of epithelial cancer patients. *Nat Commun* **10**, 449 (2019).

6. R. L. Rutishauser *et al.*, Transcriptional repressor Blimp-1 promotes CD8(+) T cell terminal differentiation and represses the acquisition of central memory T cell properties. *Immunity* **31**, 296-308 (2009).

7. A. Kallies, A. Xin, G. T. Belz, S. L. Nutt, Blimp-1 transcription factor is required for the differentiation of effector CD8(+) T cells and memory responses. *Immunity* **31**, 283-295 (2009).

8. H. M. Shin *et al.*, Epigenetic modifications induced by Blimp-1 Regulate CD8(+) T cell memory progression during acute virus infection. *Immunity* **39**, 661-675 (2013).

9. L. Pace *et al.*, The epigenetic control of stemness in CD8(+) T cell fate commitment. *Science* **359**, 177-186 (2018).

10. H. D. Marshall *et al.*, Differential expression of Ly6C and T-bet distinguish effector and memory Th1 CD4(+) cell properties during viral infection. *Immunity* **35**, 633-646 (2011).

11. N. S. Joshi *et al.*, Inflammation directs memory precursor and short-lived effector CD8(+) T cell fates via the graded expression of T-bet transcription factor. *Immunity* **27**, 281-295 (2007).

12. R. Hess Michelini, A. L. Doedens, A. W. Goldrath, S. M. Hedrick, Differentiation of CD8 memory T cells depends on Foxo1. *J Exp Med* **210**, 1189-1200 (2013).

13. R. R. Rao, Q. Li, M. R. Gubbels Bupp, P. A. Shrikant, Transcription factor Foxo1 represses T-bet-mediated effector functions and promotes memory CD8(+) T cell differentiation. *Immunity* **36**, 374-387 (2012).

14. M. Pepper, A. J. Pagan, B. Z. Igyarto, J. J. Taylor, M. K. Jenkins, Opposing signals from the Bcl6 transcription factor and the interleukin-2 receptor generate T helper 1 central and effector memory cells. *Immunity* **35**, 583-595 (2011).

15. H. Ichii *et al.*, Bcl6 is essential for the generation of long-term memory CD4+ T cells. *Int Immunol* **19**, 427-433 (2007).

16. A. Banerjee *et al.*, Cutting edge: The transcription factor eomesodermin enables CD8+ T cells to compete for the memory cell niche. *J Immunol* **185**, 4988-4992 (2010).

17. D. T. Utzschneider *et al.*, T Cell Factor 1-Expressing Memory-like CD8(+) T Cells Sustain the Immune Response to Chronic Viral Infections. *Immunity* **45**, 415-427 (2016).

18. P. Sauter, P. Matthias, The B cell-specific coactivator OBF-1 (OCA-B, Bob-1) is inducible in T cells and its expression is dispensable for IL-2 gene induction. *Immunobiology* **198**, 207-216 (1997).

19. S. Zwilling, A. Dieckmann, P. Pfisterer, P. Angel, T. Wirth, Inducible expression and phosphorylation of coactivator BOB.1/OBF.1 in T cells. *Science* **277**, 221-225. (1997).

20. J. P. Snook, C. Kim, M. A. Williams, TCR signal strength controls the differentiation of CD4(+) effector and memory T cells. *Sci Immunol* **3**, eaas9103 (2018).

21. S. Chevrier, T. Kratina, D. Emslie, A. Karnowski, L. M. Corcoran, Germinal center-independent, IgM-mediated autoimmunity in sanroque mice lacking Obf1. *Immunol Cell Biol* **92**, 12-19 (2014).

22. M. Nakamura *et al.*, Genome-wide association study identifies TNFSF15 and POU2AF1 as susceptibility loci for primary biliary cirrhosis in the Japanese population. *Am J Hum Genet* **91**, 721-728 (2012).

23. M. Ban *et al.*, Linkage disequilibrium screening for multiple sclerosis implicates JAG1 and POU2AF1 as susceptibility genes in Europeans. *J Neuroimmunol* **179**, 108-116 (2006).

24. H. Kim *et al.*, Targeting OCA-B/Pou2af1 blocks activated, autoreactive T cells in the pancreas
and type-1 diabetes. *J Exp Med* **218**, e20200533 (2021).
25. H. Shen *et al.*, Recombinant Listeria monocytogenes as a live vaccine vehicle for the induction
of protective anti-viral cell-mediated immunity. *Proc Natl Acad Sci U S A* **92**, 3987-3991
(1995).
26. A. Shakya *et al.*, Oct1 and OCA-B are selectively required for CD4 memory T cell function. *J*
Exp Med **212**, 2115-2131 (2015).
27. A. Oxenius, M. F. Bachmann, R. M. Zinkernagel, H. Hengartner, Virus-specific MHC-class II-
restricted TCR-transgenic mice: effects on humoral and cellular immune responses after viral
infection. *Eur J Immunol* **28**, 390-400 (1998).
28. L. Liu *et al.*, Gadd45 beta and Gadd45 gamma are critical for regulating autoimmunity. *J Exp*
Med **202**, 1341-1347 (2005).
29. P. Rodriguez-Jimenez *et al.*, Growth arrest and DNA damage-inducible proteins (GADD45) in
psoriasis. *Sci Rep* **11**, 14579 (2021).
30. Y. Luo *et al.*, Suppression of collagen-induced arthritis in growth arrest and DNA damage-
inducible protein 45beta-deficient mice. *Arthritis Rheum* **63**, 2949-2955 (2011).
31. S. Ju *et al.*, Gadd45b and Gadd45g are important for anti-tumor immune responses. *Eur J*
Immunol **39**, 3010-3018 (2009).
32. A. Lachmann *et al.*, ChEA: transcription factor regulation inferred from integrating genome-
wide ChIP-X experiments. *Bioinformatics* **26**, 2438-2444 (2010).
33. J. N. Blattman *et al.*, Estimating the precursor frequency of naive antigen-specific CD8 T cells.
J Exp Med **195**, 657-664 (2002).
34. A. Crawford *et al.*, Molecular and transcriptional basis of CD4(+) T cell dysfunction during
chronic infection. *Immunity* **40**, 289-302 (2014).
35. T. Samardzic, D. Marinkovic, P. J. Nielsen, L. Nitschke, T. Wirth, BOB.1/OBF.1 deficiency
affects marginal-zone B-cell compartment. *Mol Cell Biol* **22**, 8320-8331 (2002).
36. R. Ahmed, A. Salmi, L. D. Butler, J. M. Chiller, M. B. Oldstone, Selection of genetic variants of
lymphocytic choriomeningitis virus in spleens of persistently infected mice. Role in suppression
of cytotoxic T lymphocyte response and viral persistence. *J Exp Med* **160**, 521-540 (1984).
37. C. Kim, D. C. Jay, M. A. Williams, Stability and function of secondary Th1 memory cells are
dependent on the nature of the secondary stimulus. *J Immunol* **189**, 2348-2355 (2012).
38. D. M. Mitchell, E. V. Ravkov, M. A. Williams, Distinct roles for IL-2 and IL-15 in the
differentiation and survival of CD8+ effector and memory T cells. *J Immunol* **184**, 6719-6730
(2010).
39. Y. Luo, R. G. Roeder, Cloning, functional characterization, and mechanism of action of the B-
cell-specific transcriptional coactivator OCA-B. *Mol. Cell. Biol.* **15**, 4115-4124 (1995).
40. A. Shakya, J. Kang, J. Chumley, M. A. Williams, D. Tantin, Oct1 is a switchable, bipotential
stabilizer of repressed and inducible transcriptional states. *J Biol Chem* **286**, 450-459 (2011).
41. K. Murali-Krishna *et al.*, Counting antigen-specific CD8 T cells: a reevaluation of bystander
activation during viral infection. *Immunity* **8**, 177-187 (1998).
42. M. A. Williams, E. V. Ravkov, M. J. Bevan, Rapid culling of the CD4+ T cell repertoire in the
transition from effector to memory. *Immunity* **28**, 533-545 (2008).
43. J. Perovanovic *et al.*, Oct1 cooperates with the Smad family of transcription factors to promote
mesodermal lineage specification. *Sci. Signal.* **16**, eadd5750 (2023).
44. M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion for RNA-
seq data with DESeq2. *Genome Biol* **15**, 550 (2014).
45. T. Stuart *et al.*, Comprehensive Integration of Single-Cell Data. *Cell* **177**, 1888-1902 e1821
(2019).

Sun et al. Figure 1

A SMARTA LCMV D8

EV OCA-B

Foxp3
Sell
Ccr1
Zbtb16
Bcl6
Ifngr2
Il7r
Pou2af1 (Ocab)
Smad3
Tbx21 (Tbet)
Notch3
Zeb2
Ikzf2 (Helios)
Socs2
Gadd45b

B *Pou2af1*

Socs2

Gadd45b

C

OCA-B upregulated genes

Epigenomic roadmap HM	*p*-value
H3K9me3 Breast vHMEC	0.048
Molecular function	*p*-value
Toll-like receptor binding	0.006
Aspartic-type peptidase activity	0.007

OCA-B downregulated genes

Epigenomic roadmap HM	*p*-value
H3K27me3 CD4 Memory Primary Cells	1.45E-04
H3K27me3 CD8 Memory Primary Cells	9.30E-04
Molecular function	*p*-value
Superoxide-generating NADPH oxidase activator activity	1.29E-09
Cytokine receptor activity	1.24E-06

D

OCA-B upregulated genes

ChEA 2016	*p*-value
Tp53	0.01936
Smc3	0.02544

OCA-B downregulated genes

ChEA 2016	*p*-value
IRF8	1.24E-20
SMRT	1.14E-15
MYB	1.73E-09
Nerf2	4.79E-09
NCOR	1.52E-08