1

Transcription coactivator OCA-B/Pou2af1 is sufficient to promote T cell-intrinsic

CD4 memory and prospectively identifies memory precursors

Authors: Wenxiang Sun[1,2,†], Erik P. Hughes[1,2,†], Heejoo Kim[1,2,†], Jelena Perovanovic[1,2], Krystal R. Charley[1,2], Junhong Du[1,2], Andrea Ibarra[1,2], J. Scott Hale[1], Matthew A. Williams[1,2], Dean Tantin[1,2,3,*]

[1]Department of Pathology, University of Utah School of Medicine, Salt Lake City, UT 84112, U.S.A.

[2]Huntsman Cancer Institute, University of Utah School of Medicine, Salt Lake City, UT 84112, U.S.A.

[3]Lead contact

[*]Correspondence to Dean Tantin: dean.tantin@path.utah.edu

[†]Equal contributors

W. Sun's current affiliation is Sumitomo Pharma Oncology, 3900 N. Traverse Mountain Blvd., Suite 100, Lehi, Utah 84043

H. Kim's current affiliation is Janssen Research & Development, LLC., Neuroimmunology Drug Discovery, 3210 Merryfield Row, San Diego, CA 92121.

J. Perovanovic's current affiliation is Recursion Pharmaceuticals, 41 400 W, Salt Lake City, UT 84101

Characters in title (including spaces): 128

Words in abstract: 142

Characters in manuscript (no spaces, excluding legends, methods, references): 19,921

Total main figures and tables: 5

The molecular mechanisms leading to the establishment of immunological memory are inadequately understood, limiting the development of effective vaccines and durable anti-tumor immune therapies. Here we show that ectopic OCA-B expression is sufficient to drive T cells towards a memory fate, while having minimal effects on primary antiviral effector response. At primary effector response, short lived effector T cell compartments are expanded but show increased expression of *Gadd45b* and *Socs2*, while memory precursor effector cell clusters show increased expression of *Bcl2*, *Il7r*, *Tcf7* and *Slamf6*, consistent with increased memory potential. Using an OCA-B mCherry reporter mouse allele, we observe high reporter expression in CD4$^+$ T$_{CM}$ cells. We show that endogenously elevated OCA-B expression prospectively identifies memory precursor cells with increased survival capability and memory recall potential. Cumulatively, the results demonstrate that OCA-B is necessary and sufficient to promote CD4 T cell memory in vivo.

Introduction

The ability to respond robustly to secondary antigen exposures (immune memory) is a defining feature of the adaptive immune system and forms the physiologic basis for vaccination (Sallusto et al., 2010). Immune memory can also drive autoimmune and antitumor responses (Cafri et al., 2019; Gearty et al., 2022; Rosato et al., 2019; Schnell et al., 2021). Strategies that promote memory have the potential to improve both vaccine efficacy and tumor immunotherapy. Central memory T cells (T_{CM}), which home to secondary lymphoid organs, are particularly long lived and demonstrate high proliferative capacity upon stimulation.

Multiple transcription factors and epigenetic regulators are known to modulate formation or function of T_{CM}, including Blimp1 (Kallies et al., 2009; Rutishauser et al., 2009; Shin et al., 2013), Suv39h1 (Pace et al., 2018), T-bet (Joshi et al., 2007; Marshall et al., 2011), Foxo1 (Hess Michelini et al., 2013; Rao et al., 2012), Bcl6 (Ichii et al., 2007; Pepper et al., 2011), Eomes (Banerjee et al., 2010) and TCF1 (Utzschneider et al., 2016). Many of these factors manifest pro-memory functions more clearly in $CD8^+$ compared to $CD4^+$ T cells, where the molecular basis of memory is more poorly defined. In $CD4^+$ T cells, the lymphocyte-restricted transcriptional coregulator OCA-B (also known as OBF-1 and Pou2af1) is induced upon T cell antigen stimulation (Sauter and Matthias, 1997; Zwilling et al., 1997). Germline OCA-B deletion results in normal primary T cell responses to LCMV, but attenuates $CD4^+$ memory T cell formation and recall response. Elevated *Ocab* (*Pou2af1*) mRNA levels are seen in $CD4^+CD25^{lo}$ early effector cells that are enriched for memory potential (Snook et al., 2018). OCA-B is also associated with autoimmunity (Ban et al., 2006; Chevrier et al., 2014; Kim et al., 2021; Nakamura et al., 2012).

Here, we show that OCA-B deletion in T cells selectively reduces memory responses. We show that ectopic OCA-B expression does not alter effector cell numbers at peak T cell response to LCMV, but is sufficient to expand the memory recall effector pool at the expense of empty vector-transduced controls. The effect becomes progressively stronger at later timepoints. mRNA expression profiling of effector cells ectopically expressing OCA-B reveals gene expression changes associated with effector and memory properties, signaling and homing. These include global increases in *Socs2* and *Gadd45b*, expansion of short-lived terminal effector T cell populations, and increases in *Tcf7* and *Il7r* in specific memory precursor populations. Using an OCA-B reporter mouse, we show that elevated levels of endogenous OCA-B in effector T cells during the primary response prospectively identifies $CD4^+$ T effector cells with enhanced survival and expansion capability following viral rechallenge.

Results and Discussion

OCA-B in T cells is required for robust CD4$^+$ T cells memory recall responses

We used a conditional *Ocab* (*Pou2af1*) mouse allele crossed to a CD4-Cre driver, which efficiently deletes OCA-B in T cells (Kim et al., 2021), to study effects of T cell-specific OCA-B loss. We infected *Ocab$^{fl/fl}$*;CD4-Cre and control *Ocab$^{fl/fl}$* mice with LCMV^{Arm} and measured specific T cell responses using I-A^b/gp$_{66-77}$ tetramer staining. No significant differences in the frequency of positive cells were observed at 8 days post-primary infection (dpi, Fig. S1 A). Rechallenging these mice with *Listeria monocytogenes* expressing the LCMV glycoprotein epitope (Lm-gp61) (Shen et al., 1995) elicited a trending but nonsignificant decrease in the frequency of CD4$^+$ I-A^b/gp$_{66-77}^+$ CD4$^+$ T cells undergoing memory recall response in *Ocab$^{fl/fl}$*;CD4-Cre mice (Fig. 1A, D40+7). The responding cells showed little change in IL-2/IFNγ double-producing CD4$^+$ T cells, but a significant decrease in IFNγ single-producers (Fig. S1 B, C). The same cells showed minimal changes in ICOS, PD-1, Ly6C or CD62L expression (Fig. S1 D). These results indicate that infection of T cell-conditional OCA-B knockout mice with LCMV results in deficiencies in memory recall response IFNγ expression.

The relatively small reductions in frequencies of tetramer$^+$ CD4$^+$ T cells at recall response may reflect selection for a small percentage of cells escaping OCA-B deletion. This selection can be circumvented using bone marrow chimeras in which conditional deficient or control cells are competed against wild-type T cells in the same mice. We generated bone marrow radiation chimeras in which wild-type donors were co-engrafted with congenically-marked *Ocab$^{fl/fl}$*;CD4-Cre or control *Ocab$^{fl/fl}$* cells in the same recipients. After 8 weeks to allow engraftment, recipient mice were infected with LCMV. The ratio of antigen-reactive *Ocab$^{fl/fl}$* control effector CD4$^+$ T cells to wild-type competitor was close to 1:1 at 8 dpi. OCA-B deficient (*Ocab$^{fl/fl}$*;CD4-Cre) cells were slightly reduced relative to wild-type competitor (Fig. S1 E). To investigate the effects of T cell-specific OCA-B loss on memory recall responses, we rechallenged mice with Lm-gp61. Following rechallenge, the control to wild-type cell ratio remained close to 1:1 at both primary and memory recall response, while wild-type competitor cells were greatly increased at the expense of OCA-B deficient cells specifically at recall. (>8-fold, Fig. S1 E, F). These findings indicate that, similar to T cell-conditional Oct1 deficiency and global OCA-B deficiency (Shakya et al., 2015), T cell-conditional OCA-B deficiency in bone marrow chimeras results in severe and selective deficiencies in CD4$^+$ T cell memory recall responses.

Ectopic OCA-B expression is sufficient to enhance CD4$^+$ T cell memory responses in vivo

We transduced Thy1.1$^+$ SMARTA TCR transgenic donor T cells with an MSCV vector expressing mouse OCA-B. In parallel, Ly5.1$^+$ (CD45.1$^+$) control SMARTA cells were transduced with empty

vector (EV). SMARTA mice express a transgenic TCR recognizing an immunodominant LCMV

glycoprotein (GP$_{61-80}$) CD4$^+$ epitope (Oxenius et al., 1998). The cells were mixed 1:1 and adoptively

co-transferred into Thy1.2$^+$Ly5.2$^+$ recipients as depicted in Fig. 1 A. Fibroblasts, which express no

endogenous OCA-B, efficiently express OCA-B protein when transduced with pMIG-OCA-B (Fig. 1

B). After transfer, recipient mice were infected with LCMV and the cell populations were detected by

flow cytometry using GFP expressed from MSCV and congenic markers from the donor T cells. The

host T cells mounted normal responses to LCMV, allowing SMARTA cells to be monitored in the

context of a normal host response. At 8 dpi, effector EV and OCA-B transduced cells expanded

equivalently, maintaining the same ratio in both spleen and peripheral blood (Fig. 1 C and D). Another

group of mice were infected with LCMV and allowed to clear the virus and form memory. Recall

responses were induced by re-challenge using Lm-gp61. At peak T cell response to rechallenge (52+7

and 110+7 dpi), ~50-fold more OCA-B overexpressing SMARTA cells were present (Fig. 1 C and D).

Ratios of percentages from multiple mice are quantified in Fig. 1 E. These results indicate that OCA-B

expression enhances the ability of CD4$^+$ T cells to induce memory responses.

At 8 dpi, GFP$^+$ SMARTA effector T cells were uniformly CD44^{hi}CD62L^{lo} with few changes

induced by ectopic OCA-B expression (Fig. 1 F). ICOS and PD-1 levels were slightly diminished by

OCA-B overexpression. Ly6C was not significantly altered (Fig. 1 F and G). To investigate if ectopic

OCA-B expression induces gene expression changes in effector cells that promote later memory

formation, we profiled gene expression using splenic SMARTA T cells at 8 dpi. CD4$^+$ cells were first

isolated by magnetic isolation, then Ly5.1+/5.2$^+$ and GFP$^+$ cells were further isolated by FACS. We

identified ~600 down-regulated genes and ~200 up-regulated genes (Table S1). Unsupervised

hierarchical clustering identified differentially expressed gene clusters (Fig. 2 A). Down-regulated

genes included *Sell*, which encodes CD62L, and genes specific to alternative CD4$^+$ lineages, such as

Bcl6 and *Foxp3*. The top up-regulated gene, *Pou2af1*, encodes OCA-B, providing a quality control.

Other up-regulated genes were associated with effector activity (*Tbx21*, *Zeb2*), exhaustion

(*Ikzf2*/*Helios*) and attenuation of T cell signaling and proliferation (*Socs2*, *Gadd45b*). *Gadd45b*

promotes autoimmune disease and anti-tumor immune responses (Ju et al., 2009; Liu et al., 2005; Luo

et al., 2011; Rodriguez-Jimenez et al., 2021). Gene ontology (GO) analysis of the up-regulated genes

identified terms associated with H3K9me3 regulation. Down-regulated genes by contrast were strongly

associated with H3K27me3 in CD4$^+$ and CD8$^+$ memory T cells (Fig. 2 C). ChIP-X Enrichment

Analysis (ChEA) (Lachmann et al., 2010) identifies Tp53 and Smc3 as factors that also regulate the

gene set induced by OCA-B ectopic expression (Fig. 2 D). p53 and SMC3 may therefore also control

expression of these genes alongside OCA-B, or could themselves be controlled by OCA-B to mediate

gene induction via an indirect mechanism. Sets of significant Molecular Function, Epigenomic
Roadmap and ChEA GO terms are shown in Table S2. These changes may be sufficient to confer
potential increased memory formation, as small advantages compounded over time can result in the
large observed effects.

Two other genes associated with T cell survival and memory formation, *Il7r* and *Tcf7* (Joshi et
al., 2007; Utzschneider et al., 2016), were counterintuitively decreased in bulk RNA-seq. *Il7r* for
example encodes a cytokine receptor important for T cell survival. Effector CD4[+] T cell populations
are heterogeneous, with memory progenitor cells comprising a small proportion of the effector T cell
population (Blattman et al., 2002; Joshi et al., 2007; Snook et al., 2018). Increased *Il7r* expression in
memory progenitors may be obscured in bulk RNA-seq by decreases in the larger populations of
terminal effector cells, which may lead to more culling to create greater capacity for memory. We
therefore also performed single-cell RNA-sequencing (scRNA-seq) at D8 post-LCMV infection.
Uniform manifold approximation and projection (UMAP) using EV-transduced cells revealed a variety
of populations (Fig. 3 A, Table S3). Clusters 1 and 4 were enriched for genes such as *Ccr7*, *Bcl2*,
Slamf6 and *Tcf7* (both clusters), *Tox* (cluster 1), and *Cxcr5*, *Il7r*, *Icos* and *Cd69* (cluster 4, Table S3).
These clusters likely contain memory precursors. "T Memory Cells" was identified as the top cluster 4
term (Fig. 3 A). Clusters 0, 2 and 6 by contrast were enriched in *Ccl5*, *Ifng* and *Tbx21*, respectively
(Table S3), and likely comprise terminal effectors. Cluster 5 was enriched for genes encoding histones,
cyclin-dependent kinases and proliferating cell nuclear antigen (PCNA) and likely represent
proliferating effectors. OCA-B-transduction resulted in more uniform clustering with fewer and more
tightly spaced clusters identified using the same parameters (Fig. 3 B). Cluster 0 contained the largest
number of cells and was marked by *Zeb2*, *Klrg1* and *Tbx21* (Table S3). Clusters 1 and 2 were also
associated with effector activity. One difference between the empty vector- vs. OCA-B-transduced
effector cell clusters was the selective expression of *Socs2*, which encodes an attenuator of T cell
signal transduction, more robustly in the OCA-B-transduced condition (Fig. 3 C). These results suggest
an expansion of short-lived effector cells in the OCA-B-transduced condition.

OCA-B transduced cluster 3 was associated with effector/memory progenitor activity (Fig. 3).
Interestingly, *Ikzf2*, which has been associated with CD4[+] T cell exhaustion (Crawford et al., 2014),
was not enriched in the memory-associated cluster 3 but was enriched in the equivalent cluster 4 in
EV-transduced cells (Table S3). Several memory-associated genes in the memory progenitor OCA-B-
expressing cluster 3 were increased compared to equivalent memory progenitor EV-transduced
clusters. *Tcf7* was increased by >4-fold in OCA-B cluster 3, but only 2-fold in empty vector clusters 1
and 4 (Table S3). *Slamf6* was increased by >3-fold in OCA-B cluster 3, but <2.5-fold in empty vector

clusters 1 and 4. *Bcl2* and *Il7r* were similarly more strongly expressed relative to the other clusters in OCA-B- compared to empty vector-transduced cells. These results are consistent with the bulk RNA-seq in that at single-cell resolution we identify memory progenitor clusters with increased *Tcf7* and *Il7r* in the background of global decreases. *Foxo1* was not over-represented in either memory-associated empty vector cluster but was enriched in OCA-B cluster 3. Feature plots showing representative genes in the two sets of clusters are shown in Fig. 3 C. These findings suggest that OCA-B transduced cells in cluster 3 are qualitatively superior to their empty vector-transduced counterparts at forming memory.

Clustering empty vector- and OCA-B-transduced cells together allows the relative proportions of cells with common gene expression features to be determined. Again, specific clusters were enriched for effector cells (Fig. 3 D, clusters 0, 1, 4, 6), proliferating effector cells (5) and effector cells with memory progenitor potential (2 and 3). Feature plots for key genes are shown in Fig. 3 E. Cluster 2, which was enriched for genes associated with both effector and memory activity (e.g. *Ifng*, *Zeb2*) harbored >3-fold more cells in the OCA-B expressing condition compared to controls (33% vs. 10%), however cluster 3 was also associated with memory (Fig. 3 E) but contained fewer than half as many cells. Although there were fewer cells in cluster 3, analysis of gene expression levels (Fig. 3 F) showed that *Il7r*, *Zeb2* and *Gadd45b* gene expression were elevated in OCA-B-transduced cells. *Foxo1*, *Tcf7* and *Tox* by contrast were expressed similarly. Together these findings identify changes in gene expression and effector T cell populations that predict increased memory potential.

OCA-B knock-in reporter mice label CD4$^+$ memory T cell populations

To monitor OCA-B expression in viable cells, we knocked 3 mCherry reporter cassettes preceded by P2A elements into the mouse *Pou2af1* (*Ocab*) locus. The resulting reporter allele contained the mCherry cassettes inserted immediately before the stop codon and the 3' UTR (Fig. S2 A). A successful hemizygous knock-in founder and a homozygote generated by intercrossing the progeny were confirmed by PCR (Fig. S2 B). Hemizygous and homozygous reporter mnice retained normal OCA-B protein expression (Fig. S2 C). We used flow cytometry to identify peripheral blood mCherry-expressing cells from hemizygous and homozygous reporter mice and wild-type controls. OCA-B was readily detectable in B cells (Fig. S2 D, E). Peripheral T cell expression was weaker, forming a shoulder in hemizygous reporters, but a discrete peak in homozygotes (Fig. S2 D, E). We therefore used heterozygous mice to characterize B cell populations, and homozygous mice to study T cells.

Robust mCherry expression could be detected in the bone marrow, peritoneal cavity, spleen and lymph nodes of specific pathogen-free hemizygous reporter mice. Total thymocytes by contrast

expressed little OCA-B (Fig. S2 F). We used IgM and B220 to discriminate developing, immature and recirculating bone marrow B cells. OCA-B expression was readily detected in developing, immature and recirculating B cell populations compared to wild-type controls (Fig. S2 G). Expression was nearly uniform in the different populations, except for a small population of immature B cells that lacked OCA-B expression. Peritoneal B-1 and B-2 B cells were also homogeneous, with stronger expression in B-2 relative to B-1 cells (Fig. S6 H). Splenic B cells can be partitioned into transitional and mature populations based on CD93 (AA4.1) staining. T1/2/3 subsets stratified by CD23 and IgM uniformly expressed OCA-B (Fig. S6 I). CD21 and CD23 expression divides mature splenic B cells into marginal zone (MZ), follicular and non-follicular populations, and all three populations expressed OCA-B strongly (Fig. S6 J). Expression was largely uniform, but there was a significant population of newly formed $CD21^{lo}CD23^{lo}$ B cells expressing no OCA-B. The nature and significance of these cells is unknown. These results document strong reporter activity in B cell populations, with the highest level of activity in MZ and B-1 B cells. MZ B cells are largely absent in OCA-B knockout mice (Samardzic et al., 2002). These results indicate that the OCA-B reporter robustly labels B cells.

Gating out B cells, approximately 5% of thymocytes from unchallenged, specific pathogen-free homozygous reporter mice expressed OCA-B (Fig. S2 K). mCherry expression could be identified in CD4/CD8 double-negative (DN), DP and SP compartments, with the highest amounts in DN and CD8 SP cells (Fig. S2 L). In the spleen, the relative abundance of $CD4^+$ and $CD8^+$ T cells was unaltered by the homozygous reporter allele (Fig. 4 A). Splenic $CD4^+$ T cells express mCherry at ~50-fold lower levels compared to B cells and with a wider range of expression (Fig. 4 B). Peripheral blood T cell mCherry levels varied over a similar range (not shown).

We used CD62L and CD44 to stratify resting splenic $CD4^+$ and $CD8^+$ cells into naïve, activated/effector and central memory compartments. Reporter activity could be detected in approximately 7% of naïve ($CD62L^{hi}CD44^{lo}$) cells (Fig. 4 C). The nature of these cells (e.g., Tregs, Tscm, recent thymic emigrants) is unknown. Progressively larger fractions of early activated ($CD62L^{lo}CD44^{lo}$), late activated ($CD62L^{lo}CD44^{hi}$) and T_{CM} ($CD62L^{hi}CD44^{hi}$) cells expressed mCherry (Fig. 4 C). Naïve $CD8^+$ T cells showed comparatively higher expression levels, but decreased expression in the activated state and similar expression in T_{CM} cells (Fig. 4 D). Therefore, within the $CD4^+$ T cell compartment, increased OCA-B expression as measured by mCherry fluorescence progressively labels greater fractions of naïve, activated and central memory $CD4^+$ cells.

Elevated OCA-B expression prospectively enriches memory precursor $CD4^+$ T cells

To assess OCA-B responses during infection, memory formation and rechallenge, we infected reporter

mice with LCMV and used MHC tetramers to identify splenic T cells recognizing immunodominant

LCMV epitopes. Prior to infection, >90% of cells were naïve phenotype, with 13.7% of these

mCherry$^+$. ~50% of activated CD44^{hi} cells expressed mCherry (Fig. S3 A, D0). Similarly, 50% of 8

dpi effector cells expressed mCherry regardless of tetramer status (Fig. S3 B, D8). After LCMV

clearance, memory timepoint mice showed higher percentages of mCherry (D43, 61.6%), though with

lower expression per cell. Rechallenge with Lm-gp61 to generate CD4$^+$ T cell antigen-specific recall

responses reduced mCherry positivity compared to resting memory, and high-affinity tetramer$^+$ cells

declining from >60% to <10% mCherry (Fig. S3 A, D42+7). In contrast, a higher percentage of CD8$^+$

reporter T cells expressed mCherry prior to infection. Higher percentages of tetramer$^+$ mCherry$^+$ were

also present at 5 dpi, but by 8 dpi (peak T cell response) mCherry was reduced (Fig. S3 B). Antigen

specific memory CD8$^+$ T cells also showed reduced mCherry levels relative to CD4$^+$ (Fig. S3 B). We

did not collect data on CD8$^+$ rechallenge because Lm-gp61 does not express the CD8-dominant

epitope recognized by H2-D^b LCMV tetramers.

To determine if OCA-B expression in responding CD4$^+$ T cells is associated with increased

memory potential, we crossed the reporter allele to a SMARTA TCR transgenic background to fix the

TCR specificity and to the Ly5.1 congenic marker to track engrafted cells. We transferred cells into

recipients, infected with LCMV, isolated mCherry^{hi} and mCherry^{lo} cells at 8 dpi, transferred large

numbers of cells ($8{\times}10^5$) into naïve secondary recipient mice, and isolated splenic CD4$^+$ T cells to

assess contraction (Fig. 5 A). More cells were recovered from mice engrafted with mCherry^{hi} cells,

indicating that these cells contracted less than mCherry^{lo} controls (Fig. 5 B, C). Interestingly the degree

of mCherry fluorescence in the cells that did survive was the same in both groups, though with

considerably more variance associated with mCherry^{lo} (Fig. 5 D) These findings suggest that effector

CD4$^+$ cells expressing higher OCA-B undergo less cell death.

To test the ability of the cells expressing OCA-B to mount antigen recall responses, we

similarly transferred congenically marked SMARTA cells into recipient mice and infected with

LCMV, but isolated mCherry^{hi} and mCherry^{lo} cells at 43 dpi. $4{\times}10^4$ of memory SMARTA cells were

transferred into naïve hosts, which were challenged with LCMV and analyzed at 7 dpi (Fig. 5 E).

mCherry^{hi} SMARTA cells mounted better recall responses compared to mCherry^{lo} (Fig. 5 F, G) though

again with little difference in mCherry fluorescence (Fig. 5 H). This result indicates a plasticity in

OCA-B expression whereby expression normalizes following antigen re-encounter.

In mice, OCA-B expression can be detected, varying over a 10-fold range, in responding cells as early as one day after infection with lymphocytic choriomeningitis virus (LCMV) (Shakya et al., 2015). Memory progenitor cells characterized by low CD25 expression can be identified as early as early as 3 dpi (Snook et al., 2018). We transferred SMARTA homozygous OCA-B mCherry reporter T cells into congenic recipients, infected with LCMV, and collected spleens at 3 dpi to study mCherry and CD25 levels (Fig. 5I). A large fraction of responding CD25^{lo} SMARTA cells expressed mCherry (Fig. 5J). SMARTA cells from the same mouse expressing low levels of CD25 also expressed high levels of TCF1, as expected (Snook et al., 2018). mCherry and TCF1 were not compared as cell fixation quenches mCherry, however we were able to directly compare CD25 and OCA-B. CD25^{hi} cells expressed a range of OCA-B, while CD25^{lo} cells were highly enriched for mCherry^{hi} cells (Fig. 5K, L). These findings show that high OCA-B expression can be used as a method to prospectively mark live CD4$^+$ memory precursor effector T cells with augmented capacity to form central memory. Identifying transcriptional regulators that define the memory precursor effector CD4$^+$ T cells will promote advances in understanding the formation of CD4 T cell memory. Cumulatively, this study indicates that OCA-B expression within CD4$^+$ T cells is both necessary for, and sufficient to promote, the emergence of T$_{CM}$.

Materials and methods

Mice

All mice used in this study were on the C57BL6/J strain background. All mouse experiments were approved by the University of Utah Institutional Animal Care and Use Committee (protocol 00001553). *Pou2af1* (*Ocab*) conditional (floxed) mice were described previously (Kim et al., 2021). *Pou2af1*-3×mCherry reporter knock-in mice were generated on a C57BL/6N background (Biocytogen), but were backcrossed >5 times to C57BL/6J prior to generation of data herein.

LCMV and *Listeria* infection

LCMV Armstrong 53b (LCMV^{Arm}) (Ahmed et al., 1984) was grown in BHK cells and titered using Vero cells (Oxenius et al., 1998). For primary infection, 8-to 12-week-old *Ocab$^{fl/fl}$* or *Ocab$^{fl/fl}$*;CD4-Cre mice were inoculated i.p. with 2×10^5 plaque-forming units (PFU) of LCMV intraperitoneally. For heterologous rechallenge with *Listeria monocytogenes* expressing LCMV glycoprotein 61-80 (Lm-gp61) (Shen et al., 1995), bacteria were grown in log phase in BHI media, and concentrations were calculated using OD at 600 nm (OD of $1=1\times10^9$ CFU/mL). Mice were rechallenged intravenously with 2×10^5 colony forming units (CFU) of Lm-gp61 the indicated number of days after primary infection as published (Kim et al., 2012).

Bone marrow radiation chimeras

Radiation chimeras were generated as published (Mitchell et al., 2010). Briefly, male Ly5.1 C57BL/6J mice were used as recipients. Mice received a split dose of 900 Rad (2×450 RAD, spaced 1 hr apart), and were engrafted one day later with 1 million mixed donor bone marrow cells. Competitor bone marrow cells from Ly5.2/Thy1.1 but otherwise wild-type mice, and experimental cells from Ly5.2/Thy1.2 *Ocab$^{fl/fl}$* or littermate *Ocab$^{fl/fl}$*;CD4-Cre mice, were isolated from femurs and tibias of donor mice. After red blood cell lysis, CD3$^+$ cells were depleted using biotinylated anti-CD3 antibodies (eBioScience) and anti-biotin magnetic MicroBeads (Miltentyi Biotech) according to the manufacturer's instructions. CD3-depeleted bone marrow cells were mixed 1:1 and injected retro-orbitally. The chimeras were rested for 8 weeks to allow engraftment, after which mice were infected as above. Virus-reactive cells were identified by LCMVgp$_{66-77}$:I-Ab tetramer staining.

pMSCV-IRES-GFP-OCA-B vector generation

Wild-type *Ocab* cDNA was amplified from the plasmid pcDNA3.1-OCA-B (Luo and Roeder, 1995) (a gift from Robert Roeder, Rockefeller University) by PCR and cloned into the pMSCV-IRES-GFP (pMIGR1) retroviral vector (a gift from Ryan O'Connell, University of Utah). Primers used for amplification were mOcab-F-XhoI, 5'CTCGAGCTGTCTGCTTCAAAGAGAAAAGGCAAC; mOcab-R-*Eco*RI, 5' GAATTCCTAAAAGCCCTCCACGGAGAGGGT. The PCR product was gel purified, digested with *Eco*RI and *Xho*I, and inserted into a similarly digested pMIGR1 backbone using T4 DNA ligase to generate pMIG-OCA-B. All constructs were validated by resequencing.

CD4+ T cell isolation and stimulation

Naïve splenic CD4+ T cells were isolated and simulated as described previously (Shakya et al., 2011). Briefly, spleens were dissociated by grinding and passing through a 70 μm nylon strainer. Red blood cells were lysed by ACK lysis buffer (150 mM NH_4Cl, 10 mM $KHCO_3$, 0.1 mM EDTA). Cells were isolated from mice using a naïve CD4+ T cell isolation kit (Miltenyi), and stimulated in culture using 10 μg/mL plate-bound anti-CD3ε and 2 μg/mL anti-CD28 antibodies (eBioscience).

Ectopic OCA-B expression in CD4+ T cells

T cells from Ly5.1 SMARTA and Thy1.1 SMARTA mice were purified using a CD4+ T cell isolation kit (Miltenyi). Cells were placed into RPMI 1640 medium supplemented with 10% fetal bovine serum, 50 U/mL penicillin, 50 μg/mL streptomycin, 2 mM L-glutamine, 1 mM sodium pyruvate, 1× MEM nonessential amino acids, 55 μM 2-mercaptoethanol, and 20 IU/mL recombinant human IL-2. All components were from ThermoFisher with the exception of IL-2, which was supplied by R&D Systems. Cells were simultaneously transduced with pMIGR retroviruses packaged in 239T cells, with or without expressed mouse OCA-B cDNA, using spin infection. For spin infection, cells were centrifuged with 293T retroviral supernatant at 1000×*g* for 2 hr at 37°C in the presence of 4 μg/mL polybrene (Sigma). On the following day, transduced GFP+ cells were isolated by fluorescent-activated cell sorting. 2×10^4 pMIGR1- and pMIGR1-OCA-B-transduced cells were combined and co-transferred into wild-type (Ly5.2/Thy1.2) C57BL6/J mice. Chimeric mice were infected with LCMV and re-challenged with Lm-gp61 as described above.

Flow cytometry

Spleens were dissociated and passed through a 70 μm nylon strainer. Red blood cells were lysed by ACK lysis buffer (150 mM NH_4Cl, 10 mM $KHCO_3$, 0.1 mM EDTA). For intracellular staining, cell

suspensions in RPMI supplemented with 10% fetal bovine serum were re-stimulated for 4 hr with 1

μM LCMV GP_{61-80} or 0.1 μM GP_{33-41} peptide along with Brefeldin A (GolgiPlug Becton-Dickinson, 1

μl/mL) as published (Murali-Krishna et al., 1998; Williams et al., 2008). Cells were subsequently fixed

by cell fixation/permeabilization solution (Cytofix/Cytoperm, Becton-Dickinson) according to the

manufacturer's protocol. For tetramer staining, APC-conjugated gp_{66-77}:I-A^b and gp_{33-41}:H-2D^b

tetramers were provided by the NIH Tetramer Core Facility (Emory Vaccine Center). Cell suspensions

were incubated at 37°C for 3 hr in RPMI with the tetramer followed by cell-surface staining. Tetramer

fluorescence was normalized with isotype tetramer (hCLIP:I-A^b) staining. Antibodies used for flow

cytometry are presented in the Key Resources Table.

Bulk RNAseq

$CD4^+$ T cells were purified from female Ly5.1 SMARTA and Ly5.1/5.2 SMARTA mice, and

separately transduced with pMIG-OCA-B or empty vector (EV), respectively. On the following day,

transduced GFP^+ cells were sorted, 20,000 of each combined 1:1, and injected intravenously into 24

male C57BL/6J male recipients. 24h later, each recipient mouse was injected intraperitoneally with

$2x10^5$ pfu LCMV. After eight days, the 24 mice were divided into six groups of four mice by

combining four spleens together. $CD4^+$ T cells were purified using a Miltenyi CD4 T cell isolation kit.

Cells then were stained with PerCP-Cy5.5 Ly5.1 and AF700 Ly5.2 antibodies. $GFP^+Ly5.1^+Ly5.2^+$

(EV) and $GFP^+Ly5.1^+$ (OCA-B-expressing) cells were sorted by fluorescence-activated cell sorting

(FACS) and used for RNA purification (RNeasy Mini Kit, QIAGEN). RNA concentrations were

determined using a Quant-iT RNA assay kit and a Qubit fluorometer (ThermoFisher). Because limited

RNA from EV-transduced cells was obtained, the six EV samples were combined into three samples.

Five OCA-B-transduced samples (each comprising four mice) were submitted for RNAseq analysis.

These showed a high degree of concordance (not shown). Total RNA samples (200-500 ng) were

hybridized with Ribo-Zero Plus (Illumina) to deplete cytoplasmic and mitochondrial and ribosomal

RNA from samples. RNA sequencing libraries were prepared as described using the Stranded Total

RNA Prep, Ligation with Ribo-Zero Plus kit (Illumina) and IDT for Illumina RNA UD Indexes Set A,

Ligation (Illumina). Purified libraries were qualified on an Agilent Technologies 2200 TapeStation

using a D1000 ScreenTape assay. The molarity of adapter-modified molecules was defined by qPCR

using the Kapa Biosystems Kapa Library Quant Kit. Individual libraries were normalized to 0.65 nM

in preparation for Illumina sequence analysis. Sequencing libraries were chemically denatured and

applied to an Illumina NovaSeq flow cell using the NovaSeq XP workflow (20043131). Following

transfer of the flow cell to an Illumina NovaSeq 6000 instrument, a 150 x 150 cycle paired-end

sequence run was performed using a NovaSeq 6000 S4 reagent Kit v1.5 (Illlumina).

Bulk RNA-seq analysis

Bulk RNA-seq analysis was performed as previously described (Perovanovic et al.). Briefly, Reads

were aligned to *Mm10* using STAR (v2.7.3a) and checked for quality using multiqc (v1.10). Between

12 and 14 million pair-end reads were generated for each sample, with >98% of aligned reads mapping

to the correct strand and >93% of the reads uniquely aligned to the gene. Differentially expressed

genes were identified using DESeq2 version 1.24.0 (Love et al., 2014) with a 5% FDR cutoff. Features

with zero counts and 5 or fewer reads in every sample were removed from the analysis. Genes

increased by 1.4-fold or more or decreased by 2-fold or more, and with adjusted $p<0.05$ were selected

as differentially expressed ($0.5 < log2FC < -1$; padj ≤ 0.05). Figures were generated in R version 4.0.0

using functions from ggplots libraries and pheatmap.

Single-cell RNA-seq

To profile gene expression at the single-cell level, splenic GFP$^+$Ly5.1$^+$Ly5.2$^+$ (EV-transduced) and

GFP$^+$Ly5.1$^+$ (OCA-B-transduced) SMARTA cells from four pooled mice per condition were isolated,

resuspended in PBS with 0.04% bovine serum albumin (ThermoFisher), and filtered through 40 μm

strainers. Viability and cell count were assessed using a Countess II (ThermoFisher). Equilibrium to

targeted cell recovery of 6,000 cells along with Gel Beads and reverse transcription reagents were

loaded to Chromium Single Cell A to form Gel-bead-in Emulsions (GEMs). Within individual GEMs,

cDNA generated from captured and barcoded mRNA was synthesized by reverse transcription at 53°C

for 45 min. Samples were then heated to 85°C for 5 min. Single cell transcriptomes were assessed

using a 10X Genomics Chromium Single Cell Gene Expression instrument. Individual cells were

tagged with 16 bp barcodes, and specific transcripts with 10 bp Unique Molecular Identifiers (UMIs)

according to manufacturer instructions.

Single-cell RNA-seq analysis

Single cell transcriptome data were analyzed and clustered as described previously (Perovanovic et

al.). Sequences from the Chromium platform were de-multiplexed and aligned using CellRanger ver.

3.1.0 (10X Genomics) with default parameters mm10-3.0.0. Clustering, filtering, variable gene

selection and dimensionality reduction were performed using Seurat ver.4.0.4 (Stuart et al., 2019)

according to the following workflow: 1, Cells with <200 detected genes were excluded further

analysis. 2, Cells with <5% UMIs mapping to mitochondrial genes and *Cd44* expression >0 were

retained for downstream analysis, , as SMARTA cells are uniformly responding to LCMV

glycoprotein antigen at this time. 3, The UMI counts per ten thousand were log-normalized for each

cell using the natural logarithm. 4, Variable genes (2000 features) were selected using the

FindVariableFeatures function. 5, Common anchors between the conditions were identified using

FindIntegrationAnchors function that were further used to integrate these sets. 6, Gene expression

levels in the integrated set were scaled along each gene and linear dimensional reduction was

performed. The number of principal components was decided through the assessment of statistical

plots (JackStrawPlot and ElbowPlot). 7, Cells were clustered using a shared nearest neighbor (SNN)

modularity optimization-based clustering algorithm and visualized using two-dimensional uniform

manifold approximation and projection (UMAP). 8, One cluster in each condition marked

predominantly by mitochondrial genes (indicative of dying cells) was excluded from the analysis. This

resulted in 12,886 empty vector-transduced cells with 45,748 mean reads per cell, and 2,595 median

genes per cell. The total number of reads was 589,513,069, with 59.6% mapping to exonic regions.

Similarly, there were 11,709 OCA-B-transduced cells with 68,931 mean reads per cell, and 2,668

median genes per cell. Total number of reads was 807,111,486, with 57.4% mapping to exonic regions.

Clusters were identified using manual interrogation of gene expression enrichment for each population

(Table S3), the ImmGene cluster identity predictor (Table S4, S5), and PanglaoDB annotation terms.

Memory progenitor isolation, normalization and transfer

To assess contraction of the effector cells with different OCA-B levels, splenic $CD4^+$ T cells were

purified from 8 week-old female Ly5.1/5.2 homozygous knock-in reporter, SMARTA TCR transgenic

mice. 2×10^4 SMARTA cells were intravenously injected into 15 male Ly5.2/5.2 C57BL/6J recipients,

which were infected intraperitoneally 1 day later with 2×10^5 plaque-forming units of LCMV in 250 μL

phosphate-buffered saline. After 8 days, responding $Ly5.1^+5.2^+$ SMARTA T cells were sorted into

$mCherry^{hi}$ and $mCherry^{lo}$ populations. Cells were pooled and 8×10^5 $mCherry^{hi}$ and $mCherry^{lo}$ cells

were separately injected intravenously into 7 naïve age-matched 9 week-old male C57BL/6J secondary

recipients in the case of $mCherry^{lo}$ and 5 secondary recipients in the case of $mCherry^{hi}$. After 8

additional days, mice were sacrificed and splenic $Ly5.1^+5.2^+$ SMARTA cells were counted by flow

cytometry. To assess proliferative responses to rechallenge, 15 similar virus-naïve secondary recipients

were rested for 43 days post-infection prior to isolation of $mCherry^{hi}$ and $mCherry^{lo}$ populations.

1.3×10^4 SMARTA cells were transferred into naïve recipients, which were infected with LCMV one

day later. 7 days post-infection, splenic $CD4^+Ly5.1^+5.2^+$ SMARTA cells were evaluated by flow

cytometry. Mice with poor infection/response (total splenocyte count $<6 \times 10^7$) were excluded from the

analysis.

Quantification and statistical analysis

Excel (Microsoft) and Prism (GraphPad) were used for statistics and graphing. Two-tailed Student T-

tests were used to ascribe statistical significance unless otherwise indicated. For all figures, *=*p*-

value≤0.05; **=*p*≤0.005; ***=p<0.001. All error bars denote ± standard error of the mean.

Acknowledgements: We thank M. Chandrasekharan for help with graphics. We thank J. Marvin and

the University of Utah Health Sciences Center Flow Cytometry Core facility. We thank O. Allen, B.

Dalley and the High-Throughput Genomics Core. MHC tetramers were provided by the NIH Tetramer

Core Facility (Atlanta, GA). We thank R. Roeder for the mouse OCA-B cDNA and R. M. O'Connell

for the pMIGR1 plasmid.

Funding: This work was supported by NIH grant R01AI100873 to DT.

Author's contributions:

Conceptualization: DT

Methodology: WS, HK, JP, MAW

Investigation: WS, HK, EH, JD, AI

Visualization: WS, HK, JP

Funding acquisition: DT

Project administration: DT

Supervision: JSH, MAW, DT

Writing – original draft: WS, HK, JP, EH, JD, AI, JSH, MAW, DT

Competing interests: The authors declare that they have no competing interests.

Data and materials availability: All data are available in the main text or the supplementary

materials. Materials used in the study are available to any researcher for purposes of reproducing or

extending the findings. Bulk and single-cell RNA-seq data have been deposited at the Gene Expression

Omnibus GEO) website (https://www.ncbi.nlm.nih.gov/geo/, GSE214310).

Figures legends

Figure 1. **Ectopic OCA-B expression enhances CD4$^+$ T cell memory recall responses in vivo.**
(A) Experimental schematic for OCA-B transduction and T cell mixed adoptive transfer. Ly5.1$^+$ SMARTA cells were transduced with pMSCV-IRES-GFP (pMIG) empty vector (EV) and Thy1.1$^+$ SMARTA cells were transduced with pMIG expressing mouse OCA-B (pMIG-OCA-B). The donors were age- and sex-matched littermates. Following transduction, the two populations of SMARTA cells were combined 1:1, co-transferred into Ly5.2$^+$/Thy1.2$^+$ C57BL6/J recipient mice, and infected with LCMV. Recipients were rechallenged with Lm-gp61 on either day 52 or 110 post-LCMV infection, and in each case analyzed after 7 days. **(B)** Lysates from 3T3 cells transduced with either empty pMIG empty vector (EV) or pMIG-OCA-B were immunoblotted using antibodies against OCA-B. α-tubulin is shown as an internal loading standard. **(C)** Flow cytometric analysis of Ly5.1$^+$ pMIG empty vector-transduced (EV) and Thy1.1$^+$ pMIG-OCA-B-transduced (OCA-B) SMARTA cells in the spleen and blood of a representative recipient mouse at peak effector response (D8), peak memory recall response (52+7) and long-term memory recall (D110+7). Cells were first gated based on CD44, CD4 and GFP positivity. **(D)** Mean relative percentages of splenic and blood total EV- or pMIG-OCA-B-transduced (GFP$^+$) SMARTA cells were plotted at D8, D52+7 and 110+7. Each datapoint represents an individual mouse harboring both EV- or pMIG-OCA-B-transduced cells. 5, 7 and 10 mice were sacrificed at each timepoint to obtain measurements from spleen. All mice were bled at D8, and remaining mice were bled at subsequent timepoints. **(E)** Mean ratios of pMIG-OCA-B-transduced relative to empty vector-transduced SMARTA cells (OCA-B/EV) are plotted. Splenic (top) and peripheral blood (bottom) SMARTA T cells are shown at day 8 post-infection, or 7 days post-rechallenge with Lm-gp61 (52+7 or 110+7). **(F)** Flow cytometry showing relative percentages of splenic GFP$^+$CD44^{hi} EV-transduced (blue) or pMIG-OCAB-transduced (red) SMARTA cells expressing ICOS, PD-1, Ly6C or CD62L at day 8 post-infection. Representative data showing the two populations of cells within a single mouse are shown at day 8 post-infection. **(G)** Percent expression for ICOS, PD-1, CD62L and Ly6C was quantified from four mice using splenic EV- and pMIG-OCAB-transduced GFP$^+$CD44^{hi} SMARTA cells at 8 days post-infection. Four replicates were performed. For CD26L, the four datapoints are superimposed, and for Ly6C two of the datapoints are superimposed. Individual mice are connected by lines.

Figure 2. **Gene expression changes associated with ectopic OCA-B expression in SMARTA transgenic primary effector T cells. (A)** Heatmap showing top up- and down-regulated genes. Genes

with log2 fold-change >0.5 or <-1.0, and padj ≤ 0.05 are shown. Example genes are shown at right.

Control and OCA-B-transduced cells were purified from the same mice using magnetic isolation and

FACS. **(B)** Example genome tracks showing two up-regulated genes, *Socs2* and *Gadd45b*. *Pou2af1*

(*Ocab*) is shown as a positive control. **(C)** Top Epigenomic roadmap and Molecular function GO terms

for the set of OCA-B up- and down-regulated genes. **(D)** Top ChIP-X Enrichment Analysis (ChEA

2016) factors potentially regulating the same set of genes up- and down-regulated by OCA-B ectopic

expression.

Figure 3. **Single-cell RNA-seq analysis of SMARTA cells transduced with OCA-B or empty**

vector. (A) UMAP projections of independently clustered empty vector-transduced (GFP$^+$) SMARTA

cells purified similar to Figs. 2 and 3, and subjected to scRNA-seq. Cluster identities were annotated

using a combination of gene expression enrichment (Table S3), cluster identity predictor ImmGen

identity scores (Table S4) and PanglaoDB annotation terms. Clusters 1 and 4 were associated with

memory formation. Top PanglaoDB annotation terms for cluster 4 are shown below. **(B)** Similar

UMAP projection using OCA-B-transduced cells purified from the same mice as in (A). Cluster

identities were annotated similarly to (A). Cluster 3 in OCA-B-transduced cells was most strongly

associated with memory formation. Top PanglaoDB annotation terms for cluster 3 are shown below.

(C) Feature plots highlighting expression of representative genes for the two UMAP projections in (A)

and (B). **(D)** UMAP projections of the the EV- and OCA-B-transduced single-cell RNA-seq datasets

clustered together. For each projection, the percentage of each cluster relative to the total number cells

in condition is shown. Cluster identities were annotated using a combination of gene expression

enrichment (Table S3), cluster identity predictor ImmGen identity scores (Table S5) and PanglaoDB

annotation terms. **(E)** Feature plots highlighting expression of representative genes for the combined

projections.

(F) Violin plots depicting expression of six genes (*Il7r*, *Zeb2*, *Gadd45b*, *Foxo1*, *Tcf7* and *Tox*) across

each of the 10 clusters in EV-transduced cells (blue, left) and OCA-B-transduced cells (red, right).

Figure 4. **mCherry expression in splenic T cell populations in specific pathogen-free homozygous**

OCA-B-3×mCherry mice. (A) CD4 and CD8 expression are shown for an allelic series of example

wild-type (+/+) and homozygous (KI/KI) OCA-B knock-in reporter mice. **(B)** OCA-B homozygous

reporter expression in splenic CD4$^+$ T cell populations. **(C)** The same CD4$^+$ cells as in (B) were further

stratified by CD62L and CD44 into naïve effector (Teff) and central memory (Tcm) populations.

CD62L^{lo}CD44$_{lo}$ cells are likely recently activated cells that have down-regulated CD62L but not yet

up-regulated CD44. **(D)** Similar analysis as in (C) except for CD8$^+$ cells.

Figure 5. **CD4$^+$ T cells expressing high levels of OCA-B reporter activity preferentially form**

central memory cells. (A) Schematic for assessing contraction of mCherry^{hi} vs. mCherry^{lo} SMARTA

T cells following LCMV infection. Congenically marked SMARTA T cells were isolated from donor

mice, transferred into naïve secondary recipients and infected with LCMV. At 8 days post-infection,

mCherry^{hi} and mCherry^{lo} populations were isolated by FACS. 8×10^5 SMARTA cells were transferred

into naïve recipients. After 8 days, splenic CD4$^+$Ly5.1$^+$5.2$^+$ SMARTA cells were evaluated by flow

cytometry. **(B)** Flow cytometric analysis of SMARTA donor T cells engrafted into naïve mice to

monitor rates of decline. Representative mice engrafted with mCherry^{lo} (left) or mCherry^{hi} (right) cells

are shown. **(C)** Quantification of averaged donor T cell numbers from mice engrafted with mCherry^{lo}

or mCherry^{hi} cells. N=7 for the mCherry^{lo} group and N=5 for the mCherry^{hi} group. **(D)** Similar analysis

to (C) except quantifying mean mCherry fluorescence intensity. **(E)** Schematic for assessing recall

responses of mCherry^{hi} vs. mCherry^{lo} SMARTA T cells following LCMV infection. Congenically

marked SMARTA T cells were isolated from donor mice, transferred into naïve secondary recipients

and infected with LCMV. However, mice were allowed to clear LCMV and form memory. After 43

days, mCherry^{hi} and mCherry^{lo} memory T cell populations were isolated by FACS. 1.3×10^4 SMARTA

cells were transferred into naïve recipients, which were infected with LCMV one day later. 7 days

post-infection, splenic CD4$^+$Ly5.1$^+$5.2$^+$ SMARTA cells were evaluated by flow cytometry. **(F)** Flow

cytometric analysis of SMARTA donor T cells 7 days post-rechallenge to monitor recall responses.

Representative mice engrafted with mCherry^{lo} (left) or mCherry^{hi} (right) cells are shown. **(G)**

Quantification of averaged donor T cell percentages and numbers from mice engrafted with mCherry^{lo}

or mCherry^{hi} cells. N=7 for the mCherry^{lo} group and N=5 for the mCherry^{hi} group. **(H)** Similar

analysis to (G) except quantifying mean mCherry fluorescence intensity. **(I)** Schematic for assessing

mCherry levels in SMARTA T at early timepoints following LCMV infection. 2×10^5 congenically

marked (CD45.1$^+$) SMARTA T cells were isolated from donor mice and transferred into naïve

secondary recipients. One day later mice were intraperitoneally infected with 2×10^5 PFU LCMV.

Splenic T cells were collected at 3 dpi. **(J)** Gated unfixed splenic CD4$^+$ T cells were assessed for the

SMARTA congenic marker Ly5.1 and mCherry (left), while fixed Ly5.1$^+$ (SMARTA) cells from the

same mice were used to assess CD25 and TCF1 (right). **(K)** CD25 and mCherry expression were

assessed from a representative animal. Left: distribution of SMARTA cell CD25 and mCherry

expression in SMARTA mice. Right: CD25 levels of gated mCherry^{hi} and mCherry^{lo} SMARTA cells

displayed as a concatenated histogram. (**L**) Quantification of averaged mCherry MFI in CD25^{hi} and

CD25^{lo} cells (left), and frequencies (center) and numbers (right) of CD25^{lo} cells in mCherry^{lo} or

mCherry^{hi} cells SMARTA T cells. N=6 independent recipient mice.

Supplementary Materials

Figure S1. **OCA-B in T cells is required for robust polyclonal memory CD4+ T cell formation and**

cytokine expression following LCMV infection. (A) *Ocab$^{fl/fl}$*;CD4-Cre and control *Ocab$^{fl/fl}$* mice

were infected with LCMV and rechallenged with Lm-gp61 40 days after LCMV infection. Splenic I-

A^{b}/gp$_{66-77}$ tetramer-positive, CD4^{+}CD44^{+} T cells were analyzed by flow cytometry 8 days after

primary infection and 7 days after heterologous rechallenge. n=5 for each group. n.s.=not significant.

(B) Splenocytes from LCMV-infected *Ocab$^{fl/fl}$*;CD4-Cre or control *Ocab$^{fl/fl}$* mice were restimulated in

vitro with gp$_{61-80}$ peptide for 4 hours, and IFNγ and IL-2 expression in CD4^{+} T cells examined by flow

cytometry. Cells were gated based on I-A^{b}/gp$_{66-77}$ tetramer, CD4 and CD44 positivity. Data from

representative mice are shown. **(C)** Mean T cell frequencies of CD4^{+}CD44^{+}IFNγ^{+}IL-2^{-} T cells are

shown. n=5 for each group. **(D)** Surface marker expression in polyclonal OCA-B T cell-deficient and

control splenic CD4^{+} effector T cells at 8 days post-infection or 7 days post-rechallenge. Example flow

cytometry panels of ICOS, Ly6C, CD62L. PD-1 was additionally performed for memory recall

responses. Cells were gated for I-A^{b}-LCMV$^{gp66-77}$ positivity. **(E)** Flow cytometry plot showing

Thy1.1^{+}Ly5.2^{+} but otherwise genetically wild-type (WT) donor bone marrow cells engrafted 50:50

with Ly5.2^{+} control cells lacking the Cre driver (*fl/fl*) or experimental T cell-conditional *Ocab$^{fl/fl}$*;CD4-

Cre donor cells (*fl/fl*;Cre). Recipient hosts were Ly5.1^{+}. Splenic effector cells were recovered at

primary response (D8) or seven days post-rechallenge (D42+7). Cells were gated based on I-A^{b}/gp$_{66-77}$

tetramer positivity. Example mice are shown. **(F)** Mean wild-type competitor to control of OCA-B

deficient T cell ratios are shown. n=3 for each group.

Figure S2. **OCA-B-3×mCherry knock-in reporter mouse characterization in B and developing T**

cell populations. (A) Mouse *Pou2af1* locus, targeting vector and the targeted *Pou2af1* locus. **(B)** PCR

genotyping OCA-B-3✕mCherry reporter mice. PCR yields a 494 bp product for the wild-type *Pou2af1*

allele and a 373 bp product for the targeted *Pou2af1* allele. +=wild-type allele. KI=knock-in allele. **(C)**

OCA-B immunoblot using splenocytes from wild-type (+/+) and heterozygous (+/KI) or homozygous

knock-in OCA-B-3✕mCherry knock-in mice. β-actin was used as a loading control. **(D)** Flow

cytometry showing mCherry expression in peripheral blood cells of wild-type mice (+/+), or either heterozygous (+/KI) or homozygous (KI/KI) OCAB-3×mCherry knock-in reporter mice. **(E)** Similar flow cytometry plots as in (D) except gated on the indicated populations. **(F)** mCherry (OCA-B) expression in a 4.5 week-old heterozygous female OCA-B-3×mCherry reporter mouse. Cells were harvested from bone marrow (BM), peritoneal cavity (PC), spleen (SP), lymph node (LN) or thymus. After removal of red blood cells, mCherry expression was measured in total cell populations. **(G)** Wild-type (+/+) and heterozygous knock-in (+/KI) bone marrow cells were further stratified into B1 and B2 populations based on B220 and IgM expression Top plots show relative proportions of B1 ($IgM^{hi}B220^{lo}$) B2 ($IgM^{lo}B220^{hi}$) cells in wild-type and two example heterozygous reporter mice. Bottom plots show mCherry expression in gated populations. **(H)** Analysis of peritoneal cavity B cell populations from +/+ and +/KI reporter mice. Cells were stratified into pro/pre, immature and recirculating B cell compartments based on B220 and IgM expression. Top plots show relative proportions of the cells in mice of the indicated genotypes. Bottom plots show mCherry levels in the gated populations. **(I)** Splenic transitional B cells were stratified using CD93 (AA4.1), CD23 and IgM antibodies. One wild-type and two example heterozygous reporter mice are shown. **(J)** Similar analysis to (I) except using CD21 and CD23 antibodies to visualize follicular and MZ B cell populations. **(K)** mCherry expression in total B220-negative thymocytes from wild-type (+/+) or homozygous knock-in reporter mice was analyzed by flow cytometry. Specific pathogen-free mice were used. **(L)** Thymocytes were stained with CD4 and CD8 antibodies to visualize double-negative, double-positive and single-positive T cell developmental stages in homozygous knock-in reporter mice. Right: mCherry expression in developing reporter T cells was subdivided by quadrant.

Fig. S3. **OCA-B expression in antigen-responsive T cells following LCMV infection. (A)** Flow cytometry images are shown of homozygous OCA-B-3×mCherry knock-in reporter mice at different timepoints following LCMV infection (D0-43) and following rechallenge with Lm-gp61. Mice were challenged on different days and spleens were harvested and processed on the same day. Top panels: percentages of cells expressing CD44 and class II LCMV tetramer (I-A^b LCMV$^{gp66-77}$) are shown. Bottom panels: for the same timepoints cells gated from each quadrant in the top panels are plotted. Each flow panel is representative of biological duplicate mice sacrificed at each timepoint. **(B)** Cells from the same mice as above were gated for CD8 expression and stained with class I LCMV tetramers (H2-D^b LCMV$^{gp33-41}$). Analysis of recall response timepoints was not possible for antigen-specific

CD8 cells because mice were rechallenged with Lm-gp61, which lacks CD8 immunodominant

epitopes.

Table S1. **Differentially expressed bulk RNAseq genes comparing GFP+ SMARTA T cells**

expressing empty vector or OCA-B at day 8 post-LCMV infection (cutoffs: -1>log2FC>0.5,

***p*<0.05)**

Table S2. **Gene Ontology analysis using differentially expressed bulk RNA-seq transcripts**

comparing GFP+ SMARTA T cells transduced with empty vector or OCA-B at day 8 post-

LCMV infection (cutoffs: -1>log2FC>0.5, *p*<0.05)

Table S3. **Single-cell RNA-seq gene enrichments by cluster in GFP+ SMARTA T cells transduced**

with empty vector or OCA-B at day 8 post-LCMV infection.

Table S4. **Top 5 single-cell RNA-seq ImmGen identity scores per cluster using GFP+ SMARTA T**

cells transduced with empty vector or OCA-B at day 8 post-LCMV infection.

Table S5. **Integrated gene enrichment analysis of single-cell RNA-seq gene enrichments by**

cluster in GFP+ SMARTA T cells transduced with empty vector or OCA-B at day 8 post-LCMV

infection.

References and Notes

Ahmed, R., A. Salmi, L.D. Butler, J.M. Chiller, and M.B. Oldstone. 1984. Selection of genetic variants of lymphocytic choriomeningitis virus in spleens of persistently infected mice. Role in suppression of cytotoxic T lymphocyte response and viral persistence. *J Exp Med* 160:521-540.

Ban, M., D. Booth, R. Heard, G. Stewart, A. Goris, K. Vandenbroeck, B. Dubois, M. Laaksonen, J. Ilonen, M. Alizadeh, G. Edan, M.C. Babron, D. Brassat, M. Clanet, I. Cournu-Rebeix, B. Fontaine, G. Semana, R. Goedde, J. Epplen, A. Weber, C. Infante-Duarte, F. Zipp, C. Rajda, K. Bencsik, L. Vecsei, S. Heggarty, C. Graham, S. Hawkins, M. Liguori, P. Momigliano-Richiardi, D. Caputo, L.M. Grimaldi, M. Leone, L. Massacesi, C. Milanese, M. Salvetti, G. Savettieri, M. Trojano, B. Bielecki, M.P. Mycko, K. Selmaj, M. Santos, P. Maciel, C. Pereira, A. Silva, B.M. Silva, F. Coraddu, M.G. Marrosu, E. Akesson, J. Hillert, P. Datta, A. Oturai, H.F. Harbo, A. Spurkland, R. Goertsches, P. Villoslada, M. Eraksoy, A. Hensiek, A. Compston, E. Setakis, J. Gray, T.W. Yeo, and S. Sawcer. 2006. Linkage disequilibrium screening for multiple sclerosis implicates JAG1 and POU2AF1 as susceptibility genes in Europeans. *J Neuroimmunol* 179:108-116.

Banerjee, A., S.M. Gordon, A.M. Intlekofer, M.A. Paley, E.C. Mooney, T. Lindsten, E.J. Wherry, and S.L. Reiner. 2010. Cutting edge: The transcription factor eomesodermin enables CD8+ T cells to compete for the memory cell niche. *J Immunol* 185:4988-4992.

Blattman, J.N., R. Antia, D.J. Sourdive, X. Wang, S.M. Kaech, K. Murali-Krishna, J.D. Altman, and R. Ahmed. 2002. Estimating the precursor frequency of naive antigen-specific CD8 T cells. *J Exp Med* 195:657-664.

Cafri, G., R. Yossef, A. Pasetto, D.C. Deniger, Y.C. Lu, M. Parkhurst, J.J. Gartner, L. Jia, S. Ray, L.T. Ngo, M. Jafferji, A. Sachs, T. Prickett, P.F. Robbins, and S.A. Rosenberg. 2019. Memory T cells targeting oncogenic mutations detected in peripheral blood of epithelial cancer patients. *Nat Commun* 10:449.

Chevrier, S., T. Kratina, D. Emslie, A. Karnowski, and L.M. Corcoran. 2014. Germinal center-independent, IgM-mediated autoimmunity in sanroque mice lacking Obf1. *Immunol Cell Biol* 92:12-19.

Crawford, A., J.M. Angelosanto, C. Kao, T.A. Doering, P.M. Odorizzi, B.E. Barnett, and E.J. Wherry. 2014. Molecular and transcriptional basis of CD4(+) T cell dysfunction during chronic infection. *Immunity* 40:289-302.

Gearty, S.V., F. Dundar, P. Zumbo, G. Espinosa-Carrasco, M. Shakiba, F.J. Sanchez-Rivera, N.D. Socci, P. Trivedi, S.W. Lowe, P. Lauer, N. Mohibullah, A. Viale, T.P. DiLorenzo, D. Betel, and A. Schietinger. 2022. An autoimmune stem-like CD8 T cell population drives type 1 diabetes. *Nature* 602:156-161.

Hess Michelini, R., A.L. Doedens, A.W. Goldrath, and S.M. Hedrick. 2013. Differentiation of CD8 memory T cells depends on Foxo1. *J Exp Med* 210:1189-1200.

Ichii, H., A. Sakamoto, M. Arima, M. Hatano, Y. Kuroda, and T. Tokuhisa. 2007. Bcl6 is essential for the generation of long-term memory CD4+ T cells. *Int Immunol* 19:427-433.

Joshi, N.S., W. Cui, A. Chandele, H.K. Lee, D.R. Urso, J. Hagman, L. Gapin, and S.M. Kaech. 2007. Inflammation directs memory precursor and short-lived effector CD8(+) T cell fates via the graded expression of T-bet transcription factor. *Immunity* 27:281-295.

Ju, S., Y. Zhu, L. Liu, S. Dai, C. Li, E. Chen, Y. He, X. Zhang, and B. Lu. 2009. Gadd45b and Gadd45g are important for anti-tumor immune responses. *Eur J Immunol* 39:3010-3018.

Kallies, A., A. Xin, G.T. Belz, and S.L. Nutt. 2009. Blimp-1 transcription factor is required for the differentiation of effector CD8(+) T cells and memory responses. *Immunity* 31:283-295.

Kim, C., D.C. Jay, and M.A. Williams. 2012. Stability and function of secondary Th1 memory cells are dependent on the nature of the secondary stimulus. *J Immunol* 189:2348-2355.

Kim, H., J. Perovanovic, A. Shakya, Z. Shen, C.N. German, I. A., J.L. Jafek, N.-P. Lin, B.D. Evavold, D.H. Chou, P.E. Jensen, X. He, and D. Tantin. 2021. Targeting OCA-B/Pou2af1 blocks activated, autoreactive T cells in the pancreas and type-1 diabetes. *J Exp Med* 218:e20200533.

Lachmann, A., H. Xu, J. Krishnan, S.I. Berger, A.R. Mazloom, and A. Ma'ayan. 2010. ChEA: transcription factor regulation inferred from integrating genome-wide ChIP-X experiments. *Bioinformatics* 26:2438-2444.

Liu, L., E. Tran, Y. Zhao, Y. Huang, R. Flavell, and B. Lu. 2005. Gadd45 beta and Gadd45 gamma are critical for regulating autoimmunity. *J Exp Med* 202:1341-1347.

Love, M.I., W. Huber, and S. Anders. 2014. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol* 15:550.

Luo, Y., D.L. Boyle, D. Hammaker, M. Edgar, G. Franzoso, and G.S. Firestein. 2011. Suppression of collagen-induced arthritis in growth arrest and DNA damage-inducible protein 45beta-deficient mice. *Arthritis Rheum* 63:2949-2955.

Luo, Y., and R.G. Roeder. 1995. Cloning, functional characterization, and mechanism of action of the B-cell-specific transcriptional coactivator OCA-B. *Mol. Cell. Biol.* 15:4115-4124.

Marshall, H.D., A. Chandele, Y.W. Jung, H. Meng, A.C. Poholek, I.A. Parish, R. Rutishauser, W. Cui, S.H. Kleinstein, J. Craft, and S.M. Kaech. 2011. Differential expression of Ly6C and T-bet distinguish effector and memory Th1 CD4(+) cell properties during viral infection. *Immunity* 35:633-646.

Mitchell, D.M., E.V. Ravkov, and M.A. Williams. 2010. Distinct roles for IL-2 and IL-15 in the differentiation and survival of CD8+ effector and memory T cells. *J Immunol* 184:6719-6730.

Murali-Krishna, K., J.D. Altman, M. Suresh, D.J. Sourdive, A.J. Zajac, J.D. Miller, J. Slansky, and R. Ahmed. 1998. Counting antigen-specific CD8 T cells: a reevaluation of bystander activation during viral infection. *Immunity* 8:177-187.

Nakamura, M., N. Nishida, M. Kawashima, Y. Aiba, A. Tanaka, M. Yasunami, H. Nakamura, A. Komori, M. Nakamuta, M. Zeniya, E. Hashimoto, H. Ohira, K. Yamamoto, M. Onji, S. Kaneko, M. Honda, S. Yamagiwa, K. Nakao, T. Ichida, H. Takikawa, M. Seike, T. Umemura, Y. Ueno, S. Sakisaka, K. Kikuchi, H. Ebinuma, N. Yamashiki, S. Tamura, Y. Sugawara, A. Mori, S. Yagi, K. Shirabe, A. Taketomi, K. Arai, K. Monoe, T. Ichikawa, M. Taniai, Y. Miyake, T. Kumagi, M. Abe, K. Yoshizawa, S. Joshita, S. Shimoda, K. Honda, H. Takahashi, K. Hirano, Y. Takeyama, K. Harada, K. Migita, M. Ito, H. Yatsuhashi, N. Fukushima, H. Ota, T. Komatsu, T. Saoshiro, J. Ishida, H. Kouno, H. Kouno, M. Yagura, M. Kobayashi, T. Muro, N. Masaki, K. Hirata, Y. Watanabe, Y. Nakamura, M. Shimada, N. Hirashima, T. Komeda, K. Sugi, M. Koga, K. Ario, E. Takesaki, Y. Maehara, S. Uemoto, N. Kokudo, H. Tsubouchi, M. Mizokami, Y. Nakanuma, K. Tokunaga, and H. Ishibashi. 2012. Genome-wide association study identifies TNFSF15 and POU2AF1 as susceptibility loci for primary biliary cirrhosis in the Japanese population. *Am J Hum Genet* 91:721-728.

Oxenius, A., M.F. Bachmann, R.M. Zinkernagel, and H. Hengartner. 1998. Virus-specific MHC-class II-restricted TCR-transgenic mice: effects on humoral and cellular immune responses after viral infection. *Eur J Immunol* 28:390-400.

Pace, L., C. Goudot, E. Zueva, P. Gueguen, N. Burgdorf, J.J. Waterfall, J.P. Quivy, G. Almouzni, and S. Amigorena. 2018. The epigenetic control of stemness in CD8(+) T cell fate commitment. *Science* 359:177-186.

Pepper, M., A.J. Pagan, B.Z. Igyarto, J.J. Taylor, and M.K. Jenkins. 2011. Opposing signals from the Bcl6 transcription factor and the interleukin-2 receptor generate T helper 1 central and effector memory cells. *Immunity* 35:583-595.

Perovanovic, J., Y. Wu, H. Abewe, Z. Shen, E.P. Hughes, J. Gertz, M.B. Chandrasekharan, and D. Tantin. 2023. Oct1 cooperates with the Smad family of transcription factors to promote mesodermal lineage specification. *Sci. Signal.* 16:eadd5750.

Rao, R.R., Q. Li, M.R. Gubbels Bupp, and P.A. Shrikant. 2012. Transcription factor Foxo1 represses T-bet-mediated effector functions and promotes memory CD8(+) T cell differentiation. *Immunity* 36:374-387.

Rodriguez-Jimenez, P., L. Fernandez-Messina, M.C. Ovejero-Benito, P. Chicharro, P. Vera-Tome, A. Vara, D. Cibrian, P. Martinez-Fleta, M. Jimenez-Fernandez, I. Sanchez-Garcia, M. Llamas-Velasco, F. Abad-Santos, F. Sanchez-Madrid, E. Dauden, and H. de la Fuente. 2021. Growth arrest and DNA damage-inducible proteins (GADD45) in psoriasis. *Sci Rep* 11:14579.

Rosato, P.C., S. Wijeyesinghe, J.M. Stolley, C.E. Nelson, R.L. Davis, L.S. Manlove, C.A. Pennell, B.R. Blazar, C.C. Chen, M.A. Geller, V. Vezys, and D. Masopust. 2019. Virus-specific memory T cells populate tumors and can be repurposed for tumor immunotherapy. *Nat Commun* 10:567.

Rutishauser, R.L., G.A. Martins, S. Kalachikov, A. Chandele, I.A. Parish, E. Meffre, J. Jacob, K. Calame, and S.M. Kaech. 2009. Transcriptional repressor Blimp-1 promotes CD8(+) T cell terminal differentiation and represses the acquisition of central memory T cell properties. *Immunity* 31:296-308.

Sallusto, F., A. Lanzavecchia, K. Araki, and R. Ahmed. 2010. From vaccines to memory and back. *Immunity* 33:451-463.

Samardzic, T., D. Marinkovic, P.J. Nielsen, L. Nitschke, and T. Wirth. 2002. BOB.1/OBF.1 deficiency affects marginal-zone B-cell compartment. *Mol Cell Biol* 22:8320-8331.

Sauter, P., and P. Matthias. 1997. The B cell-specific coactivator OBF-1 (OCA-B, Bob-1) is inducible in T cells and its expression is dispensable for IL-2 gene induction. *Immunobiology* 198:207-216.

Schnell, A., L. Huang, M. Singer, A. Singaraju, R.M. Barilla, B.M.L. Regan, A. Bollhagen, P.I. Thakore, D. Dionne, T.M. Delorey, M. Pawlak, G. Meyer Zu Horste, O. Rozenblatt-Rosen, R.A. Irizarry, A. Regev, and V.K. Kuchroo. 2021. Stem-like intestinal Th17 cells give rise to pathogenic effector T cells during autoimmunity. *Cell* 184:6281-6298 e6223.

Shakya, A., A. Goren, A. Shalek, C.N. German, J. Snook, V.K. Kuchroo, N. Yosef, R.C. Chan, A. Regev, M.A. Williams, and D. Tantin. 2015. Oct1 and OCA-B are selectively required for CD4 memory T cell function. *J Exp Med* 212:2115-2131.

Shakya, A., J. Kang, J. Chumley, M.A. Williams, and D. Tantin. 2011. Oct1 is a switchable, bipotential stabilizer of repressed and inducible transcriptional states. *J Biol Chem* 286:450-459.

Shen, H., M.K. Slifka, M. Matloubian, E.R. Jensen, R. Ahmed, and J.F. Miller. 1995. Recombinant Listeria monocytogenes as a live vaccine vehicle for the induction of protective anti-viral cell-mediated immunity. *Proc Natl Acad Sci U S A* 92:3987-3991.

Shin, H.M., V. Kapoor, T. Guan, S.M. Kaech, R.M. Welsh, and L.J. Berg. 2013. Epigenetic modifications induced by Blimp-1 Regulate CD8(+) T cell memory progression during acute virus infection. *Immunity* 39:661-675.

Snook, J.P., C. Kim, and M.A. Williams. 2018. TCR signal strength controls the differentiation of CD4(+) effector and memory T cells. *Sci Immunol* 3:eaas9103.

Stuart, T., A. Butler, P. Hoffman, C. Hafemeister, E. Papalexi, W.M. Mauck, 3rd, Y. Hao, M. Stoeckius, P. Smibert, and R. Satija. 2019. Comprehensive Integration of Single-Cell Data. *Cell* 177:1888-1902 e1821.

Utzschneider, D.T., M. Charmoy, V. Chennupati, L. Pousse, D.P. Ferreira, S. Calderon-Copete, M. Danilo, F. Alfei, M. Hofmann, D. Wieland, S. Pradervand, R. Thimme, D. Zehn, and W. Held. 2016. T Cell Factor 1-Expressing Memory-like CD8(+) T Cells Sustain the Immune Response to Chronic Viral Infections. *Immunity* 45:415-427.

Williams, M.A., E.V. Ravkov, and M.J. Bevan. 2008. Rapid culling of the CD4+ T cell repertoire in the transition from effector to memory. *Immunity* 28:533-545.

798   Zwilling, S., A. Dieckmann, P. Pfisterer, P. Angel, and T. Wirth. 1997. Inducible expression and
799   phosphorylation of coactivator BOB.1/OBF.1 in T cells. *Science* 277:221-225.

C

OCA-B upregulated genes

Epigenomic roadmap HM	*p*-value
H3K9me3 Breast vHMEC	0.048
Molecular function	*p*-value
Toll-like receptor binding	0.006
Aspartic-type peptidase activity	0.007

OCA-B downregulated genes

Epigenomic roadmap HM	*p*-value
H3K27me3 CD4 Memory Primary Cells	1.45E-04
H3K27me3 CD8 Memory Primary Cells	9.30E-04
Molecular function	*p*-value
Superoxide-generating NADPH oxidase activator activity	1.29E-09
Cytokine receptor activity	1.24E-06

D

OCA-B upregulated genes

ChEA 2016	*p*-value
Tp53	0.01936
Smc3	0.02544

OCA-B downregulated genes

ChEA 2016	*p*-value
IRF8	1.24E-20
SMRT	1.14E-15
MYB	1.73E-09
Nerf2	4.79E-09
NCOR	1.52E-08

Sun et al. Figure 3

Sun et al. Figure 5