Transcription coactivator OCA-B/Pou2af1 is necessary and sufficient to promote T cell-intrinsic CD4 memory

Wenxiang Sun[1,2,¶,$], Heejoo Kim[1,2,Ψ,$], Jelena Perovanovic[1,2,§], Erik Hughes[1,2], Junhong Du[1,2], Andrea Ibarra[1,2], J. Scott Hale[1], Matthew A. Williams[1,2], Dean Tantin[1,2,3,*]

[1]Department of Pathology, University of Utah School of Medicine, Salt Lake City, UT 84112, U.S.A.

[2]Huntsman Cancer Institute, University of Utah School of Medicine, Salt Lake City, UT 84112, U.S.A.

[3]Lead contact

[*]Correspondence: dean.tantin@path.utah.edu

[$]Equal contributors

[¶]Current address: Sumitomo Dainippon Pharma Oncology, 3900 N. Traverse Mountain Blvd., Suite 100, Lehi, Utah 84043

[Ψ]Current address: Janssen Research & Development, LLC., Neuroimmunology Drug Discovery, 3210 Merryfield Row, San Diego, CA 92121.

[§]Current address: Recursion Pharmaceuticals, 41 400 W, Salt Lake City, UT 84101

Characters in title (including spaces): 128

Words in abstract: 218

Character count (including spaces, excluding references): 63,307

Total main figures and tables: 6

In brief eTOC blurb
Sun et al. use T cell conditional knockout and ectopic expression to show that OCA-B is required for, and sufficient to promote, CD4[+] memory T cell formation in vivo. Using fluorescent reporter mice, they show that OCA-B expression can be used to prospectively identify effector T cells with enhanced memory potential.

Highlights
- **Ectopic OCA-B expression is sufficient to promote Th1 memory**
- **OCA-B controls the expression of genes in responding CD4[+] T cells that regulate effector response and memory**
- **OCA-B expression can be used to prospectively identify CD4[+] memory precursor cells**
- **OCA-B expression is reduced after memory cells are formed**

SUMMARY

37
The molecular mechanisms leading to the establishment of durable immunological memory
are inadequately understood, limiting the development of effective vaccines and durable anti-
tumor immune therapies. Using a T cell-conditional knockout mouse model and a viral
pathogen, we show that expression of the transcription cofactor OCA-B (Pou2af1/Bob.1/OBF-
1) within T cells is necessary for proper CD4$^+$ memory T cell formation. We also show that
ectopic OCA-B expression is sufficient to drive T cells towards a memory fate, while having
minimal effects on primary antiviral effector response. Bulk and single-cell gene expression
profiling comparing cells transduced with OCA-B and empty vector at primary effector
response identifies changes in gene expression consistent with later memory formation,
including genes increased (*Tbx21*, *Il7r*, *Gadd45b*, *Socs2*) in specific subpopulations by ectopic
OCA-B expression. Short-lived effector T cell compartments are expanded but show increased
expression of *Gadd45b* and *Socs2*, while clusters of effector cells with memory potential show
increased expression of *Bcl2*, *Il7r*, *Tcf7* and *Slamf6*. We also describe an OCA-B-mCherry
reporter mouse allele that selectively labels B and T lymphocytes, and shows high reporter
expression in CD4$^+$ T$_{CM}$ cells. We show that elevated OCA-B expression prospectively
identifies cells with increased survival capability and memory recall potential. Cumulatively,
the results demonstrate that OCA-B is necessary and sufficient to promote CD4 T cell memory
in vivo.

Introduction

The ability to respond robustly to secondary antigen exposures ("immunological memory") is a defining feature of the adaptive immune system and forms the physiologic basis for vaccination(Sallusto et al., 2010). Immunological memory not only provides critical protection against pathogens, but can also drive autoimmune and anti-tumor responses (Cafri et al., 2019; Chee et al., 2014; Elyaman et al., 2008; Hurton et al., 2016; Kawakami et al., 2005; Raphael et al., 2020; Rosato et al., 2019; Yeo et al., 2018). Central memory T cells (T_{CM}), which home to secondary lymphoid organs, are particularly long-lived and demonstrate high proliferative capacity upon stimulation. In CD4$^+$ T cells, where the molecular basis of memory remains incompletely defined, the transcriptional co-regulator OCA-B (also known as Bob.1, OBF-1 and Pou2af1) (Luo and Roeder, 1995; Strubin et al., 1995) is mostly silent in naïve T cells, but becomes induced upon stimulation (Sauter and Matthias, 1997; Zwilling et al., 1997). OCA-B docks with its cognate transcription factor Oct1/Pou2f1 to regulate T cell target genes (Shakya et al., 2015). In vitro, OCA-B's effects on target gene expression are maximal after withdrawing the stimulus, then resting and re-stimulating T cells (Shakya *et al.*, 2015). OCA-B mediates these effects at least in part by recruiting histone lysine demethylases such as Jmjd1a (Kdm3a, Jhdm2a) to remove inhibitory histone methylation modifications at silent but previously activated target loci (Shakya *et al.*, 2015).

In mice, OCA-B expression can be detected, varying over a 10-fold range, in responding cells one day after infection with lymphocytic choriomeningitis virus (LCMV) (Shakya *et al.*, 2015). Germline OCA-B deletion results in normal primary T cell responses to LCMV, but severely attenuates CD4$^+$ memory T cell formation and recall response. Elevated levels of *Ocab* (*Pou2af1*) mRNA are seen in CD4$^+$CD25^{lo} early effector cells that are enriched for the ability to form memory after viral clearance (Snook et al., 2018). T cell-specific OCA-B knockout using a conditional mouse allele strongly attenuates spontaneous autoimmune diabetes, with autoreactive TCR specificities in pancreas either deleted or associated with anergy (Kim et al., 2021). Cumulatively, these findings demonstrate potent roles for OCA-B in antigen recall response, immune memory and autoimmunity.

Here, we use a conditional *Pou2af1* (*Ocab*) mouse allele to test T cell-intrinsic effects of OCA-B deletion on CD4$^+$ central memory formation and recall responses. We find that OCA-B deletion in T cells reduces memory responses while leaving primary timepoints unaffected. OCA-B deficient cells that do respond to secondary infection show defects in IFNγ expression, reverting to levels observed in the initial infection. Using retroviral transduction of primary T cells, we show that ectopic OCA-B expression does not alter effector cell numbers at 8 days post-infection (peak T cell response to LCMV), but is sufficient to expand the memory recall effector pool at the expense of empty vector-transduced controls. The effect becomes progressively stronger at later timepoints. mRNA expression profiling of effector cells, using both bulk and single-cell RNA-seq approaches, reveals changes in

genes associated with effector and memory properties, signaling and homing in cells ectopically

expressing OCA-B. These include global increases in *Socs2* and *Gadd45b*, expansion of short-lived

terminal effector T cell populations, and increases in *Tcf7* and *Il7r* in specific OCA-B-transduced

memory precursor populations. Finally, we generated an OCA-B-3×mCherry reporter knock-in mouse

allele. This allele efficiently labels B cell populations in different compartments such as bone marrow,

spleen and peritoneal cavity. Although they have 10-100-fold less fluorescence relative to B cells, in

the homozygous condition this allele efficiently labels T cells. Unexpectedly, reporter mice mark rare

populations of T cells in the thymus, and naïve CD4$^+$ T cells with some degree of OCA-B expression.

We find that CD62L^{hi}CD44^{hi} T$_{CM}$ cells have higher frequencies and levels of OCA-B expression

compared to naïve cells. Using this reporter to chart OCA-B expression in T cells in response to

LCMV^{Arm}, we show that OCA-B expression is maximal in antigen-specific memory cells and

unexpectedly declines upon rechallenge. Finally, we show that elevated endogenous OCA-B

expression prospectively identifies CD4$^+$ T effector cells responding to LCMV with enhanced survival

capability and expansion potential following viral rechallenge.

Star Methods

"Key resources" table [uploaded separately]

Resource availability

Lead contact

Further information and requests for resources and reagents should be directed to and will be fulfilled by the lead contact, Dean Tantin (dean.tantin@path.utah.edu).

Materials availability

The MSCV-GFP-OCA-B plasmid and OCA-B-3×mCherry reporter mice are available upon request from the Lead Contact with a completed Materials Transfer Agreement.

Data and code availability

This manuscript does not report original code.

Experimental model and subject details

Laboratory mice

All mice used in this study were on the C57BL6/J strain background. All animal experiments were approved by the University of Utah Institutional Animal Care and Use Committee (17-05008). *Pou2af1* (*Ocab*) conditional (floxed) mice were described previously (Kim *et al.*, 2021). *Pou2af1*-3×mCherry reporter knock-in mice were generated on a C57BL/6N background (Biocytogen), but all subsequent mouse crosses (>5 prior to generation of data herein) were to C57BL/6J.

Method details

LCMV and *Listeria* infection

LCMV Armstrong 53b (LCMV^{Arm}) (Ahmed et al., 1984) was grown in BHK cells and titered using Vero cells (Oxenius et al., 1998). For primary infection, 8-to 12-week-old $Ocab^{fl/fl}$ or $Ocab^{fl/fl}$;CD4-Cre mice were inoculated i.p. with 2×10^5 plaque-forming units (PFU) of LCMV^{Arm}. For heterologous rechallenge with *Listeria monocytogenes* expressing LCMV glycoprotein 61-80 (Lm-gp61) (Shen et al., 1995), bacteria were grown in log phase in BHI media, and concentrations were calculated using OD at 600 nm (OD of 1=1×10^9 CFU/mL). Mice were rechallenged intravenously with 2×10^5 colony forming units (CFU) of Lm-gp61 the indicated number of days after primary infection as published (Kim et al., 2012).

Bone marrow radiation chimeras

Radiation chimeras were generated as published (Mitchell et al., 2010). Briefly, male Ly5.1 C57BL/6J

mice were used as recipients. Mice received a split dose of 900 Rad (2×450 RAD, spaced 1 hr apart),

and were engrafted one day later with 1 million mixed donor bone marrow cells. Competitor bone

marrow cells from Ly5.2/Thy1.1 but otherwise wild-type mice, and experimental cells from

Ly5.2/Thy1.2 *Ocab^{fl/fl}* or littermate *Ocab^{fl/fl}*;CD4-Cre mice, were isolated from femurs and tibias of

donor mice. After red blood cell lysis, CD3$^+$ cells were depleted using biotinylated anti-CD3 antibodies

(eBioScience) and anti-biotin magnetic MicroBeads (Miltentyi Biotech) according to the

manufacturer's instructions. CD3-depeleted bone marrow cells were mixed 1:1 and injected retro-

orbitally. The chimeras were rested for 8 weeks to allow engraftment, after which mice were infected

as above. Virus-reactive cells were identified by LCMVgp$_{66-77}$:I-Ab tetramer staining.

pMSCV-IRES-GFP-OCA-B vector generation

Wild-type *Ocab* cDNA was amplified from the plasmid pcDNA3.1-OCA-B (Luo and Roeder, 1995) (a

gift from Robert Roeder, Rockefeller University) by PCR and cloned into the pMSCV-IRES-GFP

(pMIGR1) retroviral vector (a gift from Ryan O'Connell, University of Utah). Primers used for

amplification were mOcab-F-XhoI, 5'CTCGAGCTGTCTGCTTCAAAGAGAAAAGGCAAC; mOcab-R-

*Eco*RI, 5' GAATTCCTAAAAGCCCTCCACGGAGAGGGT. The PCR product was gel purified,

digested with *Eco*RI and *Xho*I, and inserted into a similarly digested pMIGR1 backbone using T4 DNA

ligase to generate pMIG-OCA-B. All constructs were validated by resequencing.

CD4$^+$ T cell isolation and stimulation

Naïve splenic CD4$^+$ T cells were isolated and simulated as described previously (Shakya et al., 2011).

Briefly, spleens were dissociated by grinding and passing through a 70 μm nylon strainer. Red blood

cells were lysed by ACK lysis buffer (150 mM NH$_4$Cl, 10 mM KHCO$_3$, 0.1 mM EDTA). Cells were

isolated from mice using a naïve CD4$^+$ T cell isolation kit (Miltenyi), and stimulated in culture using 10

μg/mL plate-bound anti-CD3ϵ and 2 μg/mL anti-CD28 antibodies (eBioscience).

Ectopic OCA-B expression in CD4$^+$ T cells

T cells from Ly5.1 SMARTA and Thy1.1 SMARTA mice were purified using a CD4$^+$ T cell isolation kit

(Miltenyi). Cells were placed into RPMI 1640 medium supplemented with 10% fetal bovine serum, 50

U/mL penicillin, 50 μg/mL streptomycin, 2 mM L-glutamine, 1 mM sodium pyruvate, 1× MEM

nonessential amino acids, 55 μM 2-mercaptoethanol, and 20 IU/mL recombinant human IL-2. All

components were from ThermoFisher with the exception of IL-2, which was supplied by R&D

Systems. Cells were simultaneously transduced with pMIGR retroviruses packaged in 239T cells, with

or without expressed mouse OCA-B cDNA, using spin infection. For spin infection, cells were

centrifuged with 293T retroviral supernatant at $1000 \times g$ for 2 hr at 37°C in the presence of 4 µg/mL

polybrene (Sigma). On the following day, transduced GFP$^+$ cells were isolated by fluorescent-

activated cell sorting. 2×10^4 pMIGR1- and pMIGR1-OCA-B-transduced cells were combined and co-

transferred into wild-type (Ly5.2/Thy1.2) C57BL6/J mice. Chimeric mice were infected with LCMV^{Arm}

and re-challenged with Lm-gp61 as described above.

Flow cytometry

Spleens were dissociated and passed through a 70 µm nylon strainer. Red blood cells were lysed by

ACK lysis buffer (150 mM NH$_4$Cl, 10 mM KHCO$_3$, 0.1 mM EDTA). For intracellular staining, cell

suspensions in RPMI supplemented with 10% fetal bovine serum were re-stimulated for 4 hr with 1

µM LCMV GP$_{61-80}$ or 0.1 µM GP$_{33-41}$ peptide along with Brefeldin A (GolgiPlug Becton-Dickinson, 1

µl/mL) as published (Murali-Krishna et al., 1998; Williams et al., 2008). Cells were subsequently fixed

by cell fixation/permeabilization solution (Cytofix/Cytoperm, Becton-Dickinson) according to the

manufacturer's protocol. For tetramer staining, APC-conjugated gp$_{66-77}$:I-A^b and gp$_{33-41}$:H-2D^b

tetramers were provided by the NIH Tetramer Core Facility (Emory Vaccine Center). Cell suspensions

were incubated at 37°C for 3 hr in RPMI with the tetramer followed by cell-surface staining. Tetramer

fluorescence was normalized with isotype tetramer (hCLIP:I-A^b) staining. Antibodies used for flow

cytometry are presented in the Key Resources Table.

Bulk RNAseq

CD4$^+$ T cells were purified from female Ly5.1 SMARTA and Ly5.1/5.2 SMARTA mice, and separately

transduced with pMIG-OCA-B or empty vector (EV), respectively. On the following day, transduced

GFP$^+$ cells were sorted, 20,000 of each combined 1:1, and injected intravenously into 24 male

C57BL/6J male recipients. 24h later, each recipient mouse was injected intraperitoneally with 2×10^5

pfu LCMV^{Arm}. After eight days, the 24 mice were divided into six groups of four mice by combining four

spleens together. CD4$^+$ T cells were purified using a Miltenyi CD4 T cell isolation kit. Cells then were

stained with PerCP-Cy5.5 Ly5.1 and AF700 Ly5.2 antibodies. GFP$^+$Ly5.1$^+$Ly5.2$^+$ (EV) and

GFP$^+$Ly5.1$^+$ (OCA-B-expressing) cells were sorted by fluorescence-activated cell sorting (FACS) and

used for RNA purification (RNeasy Mini Kit, QIAGEN). RNA concentrations were determined using a

Quant-iT RNA assay kit and a Qubit fluorometer (ThermoFisher). Because limited RNA from EV-

transduced cells was obtained, the six EV samples were combined into three samples. One of the six

OCAB OE sample failed. Therefore, three EV-transduced and five OCA-B-transduced samples were

eventually submitted for RNAseq analysis. Total RNA samples (200-500 ng) were hybridized with

Ribo-Zero Plus (Illumina) to deplete cytoplasmic and mitochondrial and ribosomal RNA from samples.

RNA sequencing libraries were prepared as described using the Stranded Total RNA Prep, Ligation

with Ribo-Zero Plus kit (Illumina) and IDT for Illumina RNA UD Indexes Set A, Ligation (Illumina).

Purified libraries were qualified on an Agilent Technologies 2200 TapeStation using a D1000

ScreenTape assay. The molarity of adapter-modified molecules was defined by qPCR using the Kapa

Biosystems Kapa Library Quant Kit. Individual libraries were normalized to 0.65 nM in preparation for

Illumina sequence analysis. Sequencing libraries were chemically denatured and applied to an

Illumina NovaSeq flow cell using the NovaSeq XP workflow (20043131). Following transfer of the flow

cell to an Illumina NovaSeq 6000 instrument, a 150 x 150 cycle paired-end sequence run was

performed using a NovaSeq 6000 S4 reagent Kit v1.5 (IIllumina).

Bulk RNA-seq analysis

Bulk RNA-seq analysis was performed as previously described (Perovanovic et al.). Briefly, Reads

were aligned to *Mm10* using STAR (v2.7.3a) and checked for quality using multiqc (v1.10). Between

12 and 14 million pair-end reads were generated for each sample, with >98% of aligned reads

mapping to the correct strand and >93% of the reads uniquely aligned to the gene. Differentially

expressed genes were identified using DESeq2 version 1.24.0 (Love et al., 2014) with a 5% FDR

cutoff. Features with zero counts and 5 or fewer reads in every sample were removed from the

analysis. Genes increased by 1.4-fold or more or decreased by 2-fold or more, and with adjusted

$p<0.05$ were selected as differentially expressed ($0.5 < \log2FC < -1$; padj ≤ 0.05). Figures were

generated in R version 4.0.0 using functions from ggplots libraries and pheatmap.

Single-cell RNA-seq

To profile gene expression at the single-cell level, splenic GFP[+]Ly5.1[+]Ly5.2[+] (EV-transduced) and

GFP[+]Ly5.1[+] (OCA-B-transduced) SMARTA cells were collected similarly as for bulk RNA-seq. Cells

were resuspended in PBS with 0.04% bovine serum albumin (ThermoFisher), and filtered through 40

μm strainers. Viability and cell count were assessed using a Countess II (ThermoFisher). Equilibrium

to targeted cell recovery of 6,000 cells along with Gel Beads and reverse transcription reagents were

loaded to Chromium Single Cell A to form Gel-bead-in Emulsions (GEMs). Within individual GEMs,

cDNA generated from captured and barcoded mRNA was synthesized by reverse transcription at

53°C for 45 min. Samples were then heated to 85°C for 5 min. Single cell transcriptomes were

assessed using a 10X Genomics Chromium Single Cell Gene Expression instrument. Individual cells

were tagged with 16 bp barcodes, and specific transcripts with 10 bp Unique Molecular Identifiers

(UMIs) according to manufacturer instructions.

Single-cell RNA-seq analysis

Single cell transcriptome data were analyzed and clustered as described previously (Perovanovic *et*

al.). Sequences from the Chromium platform were de-multiplexed and aligned using CellRanger ver.

3.1.0 (10X Genomics) with default parameters mm10-3.0.0. Clustering, filtering, variable gene

selection and dimensionality reduction were performed using Seurat ver.4.0.4 (Stuart et al., 2019)

according to the following workflow: 1, Cells with <200 detected genes were excluded further analysis.

2, Cells with <5% UMIs mapping to mitochondrial genes and *Cd44* expression >0 were retained for

downstream analysis. 3, The UMI counts per ten thousand were log-normalized for each cell using the

natural logarithm. 4, Variable genes (2000 features) were selected using the FindVariableFeatures

function. 5, Common anchors between the conditions were identified using FindIntegrationAnchors

function that were further used to integrate these sets. 6, Gene expression levels in the integrated set

were scaled along each gene and linear dimensional reduction was performed. The number of

principal components was decided through the assessment of statistical plots (JackStrawPlot and

ElbowPlot). 7, Cells were clustered using a shared nearest neighbor (SNN) modularity optimization-

based clustering algorithm and visualized using two-dimensional uniform manifold approximation and

projection (UMAP). 8, One cluster in each condition marked predominantly by mitochondrial genes

(indicative of dying cells) was excluded from the analysis. This resulted in 12,886 empty vector-

transduced cells with 45,748 mean reads per cell, and 2,595 median genes per cell. The total number

of reads was 589,513,069, with 59.6% mapping to exonic regions. Similarly, there were 11,709 OCA-

B-transduced cells with 68,931 mean reads per cell, and 2,668 median genes per cell. Total number

of reads was 807,111,486, with 57.4% mapping to exonic regions.

Memory progenitor isolation, normalization and transfer

To assess contraction of the effector cells with different OCA-B levels, splenic CD4[+] T cells were

purified from 8 week-old female Ly5.1/5.2 homozygous knock-in reporter, SMARTA TCR transgenic

mice. 2×10^4 SMARTA cells were intravenously injected into 15 male Ly5.2/5.2 C57BL/6J recipients,

which were infected intraperitoneally 1 day later with 2×10^5 plaque-forming units of LCMV[Arm] in 250 μL

phosphate-buffered saline. After 8 days, responding Ly5.1[+]5.2[+] SMARTA T cells were sorted into

mCherry[hi] and mCherry[lo] populations. Cells were pooled and 8×10^5 mCherry[hi] and mCherry[lo] cells

were separately injected intravenously into 7 naïve age-matched 9 week-old male C57BL/6J

secondary recipients in the case of mCherry[lo] and 5 secondary recipients in the case of mCherry[hi].

After 8 additional days, mice were sacrificed and splenic Ly5.1[+]5.2[+] SMARTA cells were counted by

flow cytometry. To assess proliferative responses to rechallenge, 15 similar virus-naïve secondary

recipients were rested for 43 days post-infection prior to isolation of mCherry[hi] and mCherry[lo]

populations. 1.3×10^4 SMARTA cells were transferred into naïve recipients, which were infected with

LCMV one day later. 7 days post-infection, splenic CD4[+]Ly5.1[+]5.2[+] SMARTA cells were evaluated by

flow cytometry. Mice with poor infection/response (total splenocyte count $<6\times10^7$) were excluded from
the analysis.
281
Quantification and statistical analysis
283 Excel (Microsoft) and Prism (GraphPad) were used for statistics and graphing. Two-tailed Student T-
284 tests were used to ascribe statistical significance unless otherwise indicated. For all figures, *=p-
285 value≤0.05; **=p≤0.005; ***=p<0.001. All error bars denote ± standard error of the mean.

Results

OCA-B is required for robust CD4$^+$ T cells memory recall responses

We previously used OCA-B whole-body knockout T cells, SMARTA TCR transgenic mice, and mixed adoptive transfer assays to demonstrate that OCA-B is required for robust monoclonal CD4$^+$ T cell memory responses. SMARTA mice express a transgenic TCR specific for an immunodominant CD4$^+$ epitope of the LCMV glycoprotein (GP$_{61-80}$) (Oxenius *et al.*, 1998). While *Ocab$^{-/-}$* SMARTA CD4$^+$ effector T cell expansion was equivalent to wild-type, ~50-fold fewer *Ocab$^{-/-}$* cells populated the memory compartment after re-challenge (Shakya *et al.*, 2015). These experiments were limited by the fact that 1) T cells were subject to effects of development in an OCA-B deficient environment and 2) monoclonal responses were studied. We used a conditional *Ocab* (*Pou2af1*) mouse allele crossed onto a CD4-Cre driver, which efficiently deletes OCA-B in T cells (Kim *et al.*, 2021), to study effects of OCA-B loss during polyclonal LCMV responses. We infected age- and sex-matched *Ocab$^{fl/fl}$*;CD4-Cre and littermate control *Ocab$^{fl/fl}$* mice with 2×10^5 pfu LCMV^{Arm} intraperitoneally, and measured specific T cell responses using I-A^b/gp$_{66-77}$ tetramer staining. No significant differences in the frequency of positive cells were observed at 8 days post-primary infection (Figure 1A). Rechallenging these mice with *Listeria monocytogenes* expressing the LCMV glycoprotein epitope (Lm-gp61) (Shen *et al.*, 1995) elicited a trending decrease in the frequency of CD4$^+$ I-A^b/gp$_{66-77}$-positive CD4$^+$ T cells undergoing memory recall response in *Ocab$^{fl/fl}$*;CD4-Cre mice (Figure 1A, D40+7). The responding cells showed no change in percentages of IL-2/IFNγ double-producing CD4$^+$ T cells, but a significant decrease in IFNγ single-producers (Figure 1B,C). The same cells showed minimal changes in Inducible T cell co-stimulator (ICOS), PD-1, Ly6C or CD62L expression (Figure S1).

The relatively small reductions in frequencies of tetramer-positive CD4$^+$ T cells at recall response may reflect selection for a small percentage of cells escaping OCA-B deletion. Infection of mice with T cell conditional deletion of *Oct1*/*Pou2f1* (which encodes the transcription factor with which OCA-B docks), indicated a strong selection for cells that escape Cre-mediated deletion in antigen-specific CD4$^+$ memory T cells (Shakya *et al.*, 2015). This selection can be circumvented by performing LCMV infections using bone marrow chimeras in which conditional-deficient or control cells are competed against wild-type T cells in the same animals. We generated bone marrow radiation chimeras in which wild-type donors were co-engrafted 1:1 with congenically-marked *Ocab$^{fl/fl}$*;CD4-Cre or control *Ocab$^{fl/fl}$* cells into the same recipient animals. In this case, a potential competitive disadvantage conferred by OCA-B deficiency can be identified relative to co-transferred wild-type competitors, and relative to control OCA-B sufficient cells carrying the same congenic markers. Bone marrow was depleted of T cells prior to engraftment (see Methods). After 8 weeks to allow engraftment, recipient mice were infected with LCMV^{Arm} (Figure 1D). The ratio of antigen-reactive

Ocab^{fl/fl} control effector CD4$^+$ T cells to wild-type competitor was close to 1:1 at 8 day post-infection.

OCA-B deficient (*Ocab^{fl/fl}*;CD4-Cre) cells were slightly reduced relative to wild-type competitor

(maximum difference 1.6 fold, Figure 1E). To investigate the effects of T cell-specific OCA-B loss on

memory recall responses, we rechallenged mice with Lm-gp61. Following rechallenge, the control to

wild-type cell ratio remained close to 1:1 at both primary and memory recall response, while wild-type

competitor cells were greatly increased at the expense of OCA-B deficient cells specifically at recall.

(>8-fold, Figure 1E,F). Therefore, much like T cell-conditional Oct1 deficiency (Shakya *et al.*, 2015), T

cell-conditional OCA-B deficiency in bone marrow chimeras results in competitive defects selectively

in CD4$^+$ T cell memory recall responses.

Ectopic OCA-B expression is sufficient to enhance CD4$^+$ T cell memory responses in vivo.

To test if ectopic expression of OCA-B is sufficient to induce more robust memory responses in CD4$^+$

T cells, we generated an MSCV-IRES-GFP (pMIGR1) vector expressing mouse OCA-B (pMIG-OCA-

B). Using this construct, we transduced Thy1.1$^+$ SMARTA TCR transgenic donor T cells. In parallel,

Ly5.1$^+$ (CD45.1$^+$) control SMARTA cells were transduced with empty vector (EV). The cells were

mixed 1:1 and adoptively co-transferred into Thy1.2$^+$Ly5.2$^+$ recipients as depicted in Figure 2A. NIH-

3T3 cells, which express no endogenous OCA-B, efficiently express OCA-B protein when transduced

with pMIG-OCA-B (Figure 2B). After transfer, recipient mice were infected with LCMV^{Arm} and the cell

populations were detected by flow cytometry using GFP expressed from MSCV and congenic markers

from the donor T cells. The host T cells mounted normal responses to LCMV, allowing SMARTA cells

to be monitored in the context of a normal host response. At 8 days post-infection, effector EV- and

OCA-B-transduced cells expanded equivalently, maintaining the same ratio in both spleen and

peripheral blood (Figure 2C and D, D8). Another group of mice were infected with LCMV^{Arm} and

allowed to clear the virus and form memory. Recall responses were induced by re-challenge 52 days

post-infection using Lm-gp61. At peak T cell response to rechallenge (day 52+7), ~50-fold more OCA-

B overexpressing SMARTA cells were present (Figure 2C and D, D52+8). We also tested the effect of

OCA-B overexpression on long-term memory by secondary challenge at day 110. In the case of long-

term memory recall responses, responding SMARTA T cells were even more strongly skewed

towards cells ectopically expressing OCA-B (Figure 2C and D, D110+7), indicating a progressive

increase in memory T cell potential with time. Ratios of percentages from multiple mice are quantified

in Figure 2E. These results indicate that the OCA-B expression enhances the ability of CD4$^+$ T cells to

induce memory responses.

Effector T cells that are capable of forming memory can be partly distinguished by the

expression of specific surface proteins, though the markers are more robust in CD8$^+$ than CD4$^+$ cells

(Laidlaw et al., 2016; Marshall et al., 2011). To investigate how OCA-B overexpression induces more

robust memory responses, we characterized the expression of ICOS, programmed cell death protein-

1 (PD-1) (PD-1), CD62L (L-selectin) and Ly6C in OCA-B-overexpressing and empty vector control

SMARTA cells at 8 days post-infection. GFP$^+$ SMARTA effector T cells were uniformly CD44^{hi} and

unchanged by OCA-B expression (Figure 2F). ICOS and PD-1 levels were slightly decreased by

OCA-B overexpression. All GFP$^+$ SMARTA T cells were CD62L^{lo}, however expression was slightly

decreased in MIGR1-OCA-B-transduced cells compared to EV (Figure 2F and G). Ly6C by contrast

was not significantly altered (Figure 2F and G).

Ectopic OCA-B expression results in gene expression changes in effector T cells that precede

and predict enhanced memory.

Although EV- and OCA-B-transduced SMARTA transgenic CD4$^+$ T cells expand equivalently in

response to primary LCMV infection, we hypothesized that ectopic OCA-B expression induces gene

expression changes in effector cells that promote later memory formation. Because we did not

observe large changes in tested surface markers, we profiled gene expression using splenic SMARTA

T cells from recipient mice processed and isolated similarly to the cells in Figure 2C. CD4$^+$ cells were

first isolated by magnetic isolation, then Ly5.1+/5.2$^+$ and GFP$^+$ cells were further isolated by FACS

(see methods). Three EV groups (each comprising 8 mice) and five OCA-B ectopic expression groups

(each comprising four mice) were used for bulk RNA-seq, which showed a high degree of

concordance (Figure S2A). Using log2 fold-change cutoffs of <-1 (down-regulated) and >0.5 (up-

regulated) and adjusted $p<0.05$, we identified a large set of ~600 down-regulated genes and a smaller

group of ~200 up-regulated genes (Table S1).

Unsupervised hierarchical clustering identified groups of genes consistently differentially

expressed between the two conditions (Figure 3A). Down-regulated genes included *Sell*, which

encodes CD62L and which was also down-regulated at the protein level (Figure 2F-G). Down-

regulated genes also surprisingly included a number of genes specific to alternative lymphocyte

lineages, such as *Bcl6* and *Foxp3*, which encode master transcription regulators of germinal center B

cells and Tfh cells and of regulatory T cells respectively. The top up-regulated gene (3.4-fold) was

Ocab (*Pou2af1*) itself, providing a quality control (Figure 3A,B). Other up-regulated genes were

associated with effector T cell activity (*Tbx21*/*Tbet*, *Zeb2*), CD4$^+$ T cell exhaustion (*Ikzf2*/*Helios*),

attenuation of T cell signaling and proliferation (*Socs2*, *Gadd45b*) and fatty acid metabolism (*Acacb*).

Example tracks for up- and down-regulated genes are shown in Figure 3B and Figure S2B. Gene

ontology (GO) analysis of the up-regulated genes identified terms associated with H3K9me3

regulation. Down-regulated genes by contrast were strongly associated with H3K27me3 in CD4$^+$ and

CD8$^+$ memory T cells (Figure 3C). Because H3K27me3 is a repressive histone modification, the

further down-regulation of genes associated with this modification is consistent with promotion of

memory phenotypes. ChIP-X Enrichment Analysis (ChEA) (Lachmann et al., 2010) identifies Tp53

and Smc3 as factors that also regulate the gene set induced by OCA-B ectopic expression (Figure

3D). p53 and SMC3 may therefore also control expression of these genes alongside OCA-B, or could

themselves be controlled by OCA-B to mediate gene induction via an indirect mechanism. The

complete sets of significant (*p*<0.05) Molecular Function, Epigenomic Roadmap and ChEA GO terms

are shown in Table S2.

We intersected the differential gene expression list with prior bulk RNA-seq experiments using

isolated naïve T OCA-B germline knockouts stimulated, rested and restimulated in vitro, and with

ChIP-seq experiments identifying targets of Oct1 and OCA-B in these cells (Shakya *et al.*, 2015).

Although the formats for these experiments were different, we reasoned that the intersection of these

gene sets would represent robust regulated genes and direct targets. Only 1.6% of genes (10 of 620)

down-regulated by OCA-B ectopic expression relative to empty vector were up-regulated by OCA-B

knockout in vitro in the prior study (Table S3). In contrast, a larger percentage of genes increased by

OCA-B ectopic expression (10 of 215, 5%) were down-regulated by OCA-B loss in vitro. These were

Tbx21 (*Tbet*), *Zbtb32*, *Gadd45b*, *Socs2*, *Il1rl1*, *Zeb2*, *Tmem159*, *Tnfsf9*, *Sytl3* and *Plekhf1*. For ChIP-

seq, we compared the set of 2680 targets co-bound by Oct1 and OCA-B (Shakya *et al.*, 2015) to the

620 genes down-regulated by ectopic OCA-B expression. Of these, 43 were identified as direct

Oct1/OCA-B targets. Down-regulated direct target genes include *Bcl6*, *Id3*, *Il7r* and *Sell* (*Cd62l*). Of

the 215 up-regulated genes, 25 were identified including *Gadd45b*, *Smad3* and *Tbx21* (*Tbet*) and

Zbtb32 (Table S4). Overall, these findings reveal that OCA-B induces complex changes in the

expression of key genes in T cells, many of which are direct targets. Some of these are consistent

with attenuation of signaling, growth arrest and later memory formation, while others are more

consistent with increased effector function and reduced survival. These findings can be explained by

the fact that CD4[+] T cell effector responses are heterogenous.

Single-cell resolution RNA-sequencing identifies putative memory progenitor cells ectopically

expressing OCA-B.

Tcf7 is a transcription factor that has been associated with CD4[+] T cell memory (Utzschneider et al.,

2016). It has been identified as an OCA-B target gene based on prior ChIPseq (Shakya *et al.*, 2015),

but was not significantly differentially expressed in bulk RNA-seq. Similarly, *Il7r* encodes a component

of IL-7 receptor, a common gamma chain cytokine receptor important for T cell survival that has also

been implicated in memory formation (Joshi et al., 2007). This gene is also a direct OCA-B target

gene, but counter-intuitively is down-regulated in bulk RNA-seq (Figure 3A). *Zeb2* is associated with

effector function in CD8[+] T cells (Guan et al., 2018; Omilusik et al., 2015) and was unexpectedly

induced (Figure 3A, Table S1). Memory progenitor cells comprise a small proportion of the effector T

cell population (Blattman et al., 2002; Joshi *et al.*, 2007; Snook *et al.*, 2018). Because bulk RNA-seq

approaches lack the granularity to identify changes in relative sub-populations or in the genes they

express, we also performed single-cell RNA-sequencing (scRNA-seq) at D8 post-LCMV^{Arm} infection.

Splenic SMARTA TCR transgenic CD4$^+$ and GFP$^+$ T cells isolated from four pooled mice per condition

were isolated as in the bulk RNA-seq, combined and subjected to scRNA-seq. Similar to a published

scRNA-seq study using effector SMARTA cells (Ciucci et al., 2019), we filtered the data by requiring

Cd44 gene expression, as SMARTA cells should be uniformly responding to LCMV at this time.

Approximately 13,000 and 12,000 empty vector and OCA-B-expressing SMARTA T cells were profiled

with an average depth of ~46,000 and ~69,000 reads/cell, respectively (see Methods).

Uniform manifold approximation and projection (UMAP) using empty vector-transduced cells

revealed a variety of populations (Figure 4A) almost uniformly expressing CD3 cluster genes such as

Cd3e as well as *Cd4* (Figure S3). Two small clusters (10 and 11) lacked CD3 gene expression and

were likely contaminants. Cluster 8 lacked *Cd4*. We used the cluster identity predictor with the

ImmGen database, gene expression enrichment for each population relative to the others (Table S5)

and PanglaoDB annotation terms to help define the nature of the clusters. Cluster 8 for example was

enriched for *Cd8a*, *Prf1*, *Gzmb* and other hallmark cytotoxic T lymphocyte genes, marking these cells

as rare CD8$^+$ cells. Contaminating clusters 10 and 11 were myeloid in nature.

Of the remaining clusters, 1 and 4 for were enriched for genes such as *Ccr7*, *Bcl2*, *Slamf6* and

Tcf7 (both clusters), *Tox* (cluster 1), and *Cxcr5*, *Il7r*, *Icos* and *Cd69* (cluster 4, Table S5). These

clusters likely contain memory precursors, with cluster 4 more memory-like than cluster 1. Cluster

annotation terms associated with clusters 4 are shown at the bottom of Figure 4A. Cluster 4 identifies

"T Memory Cells" as the first term (adj. *p*=3.4e-9). Cluster 1 was also associated with memory based

on annotation terms, albeit more weakly (Table S5 and not shown). Clusters 0, 2 and 6 by contrast

were enriched in *Ccl5* (*Rantes*), *Ifng* and *Tbx21*, respectively (Table S5), and likely comprise terminal

effectors. Cluster 5 was enriched for genes encoding histones, cyclin-dependent kinases and

proliferating cell nuclear antigen (PCNA) and likely represent proliferating effector T cells. Other

clusters also contained gene expression hallmarks of effector T cells (Figure 4A, Table S5).

UMAP projections of OCA-B-transduced cells resulted in more uniform clustering (Figure 4B).

Fewer, more tightly spaced clusters were identified using the same parameters. Like empty vector-

transduced SMARTA cells, almost all the cells expressed *Cd3e* and *Cd4* (Figure S3). Only one

contaminating myeloid population (Figure 4B, cluster 6) was identified, and unlike empty vector-

transduced cells there was no CD8-expressing cluster. The relative lack of rare CD8-expressing cells

likely explains the decrease in *Cd8a* and *Cd8b1* gene expression in bulk RNA-seq (Figure 2A).

Cluster 0 contained the largest number of cells and was marked by *Zeb2* expression, as well as other

short-lived effector cell markers such as *Klrg1* and *Tbx21* (Table S5). In cells transduced with empty

vector, *Zeb2* was enriched in rarer effector populations (Figure 4A, cluster 6) and in CD8$^+$ cells

(cluster 8), explaining the enrichment for this gene in bulk RNA-seq. One difference between the

empty vector- vs. OCA-B-transduced effector cell clusters was the selective co-expression of *Socs2*,

which encodes an attenuator of T cell signal transduction, only in the OCA-B-transduced condition
(Figure 4C). These results suggest an expansion of short-lived (terminal) effector cells in the OCA-B-
transduced condition. Similar to cluster 5 of EV-transduced cells, cluster 5 cells in the OCA-B-
transduced condition strongly expressed proliferative markers like histones and *Pcna*. Cluster 4 was
most equivalent to cluster 7 in empty vector-transduced cells.

The remaining cells were more cohesively clustered compared to their empty vector-
transduced counterparts and associated with either effector activity (cluster 0, 1, 2) or
effector/memory progenitor activity (cluster 3). The top GO term associated with cluster 3 was also "T
memory cells" (Figure 4B, bottom). Interestingly, *Ikzf2*, which has been associated with CD4[+] T cell
exhaustion (Crawford et al., 2014), was not enriched in the memory-associated cluster 3, but was
enriched in the equivalent cluster 4 in EV-transduced cells (Table S5). Furthermore, a number of
memory-associated genes in the memory progenitor OCA-B-expressing cluster 3 were more strongly
expressed compared to equivalent memory progenitor EV-transduced clusters. Genes expressed
more frequently in cluster 3 compared to the surrounding clusters included *Tcf7* (99.3% of cells), *Bcl2*,
Il7r and *Foxo1* (Table S5). *Tcf7* for example was increased by greater than 4-fold in OCA-B cluster 3,
but only slightly more than 2-fold in empty vector clusters 1 and 4 (Table S5). *Slamf6* was increased
by greater than 3-fold in OCA-B cluster 3, but by less than 2.5-fold in empty vector clusters 1 and 4.
Bcl2 and *Il7r* were similarly more strongly expressed relative to the other clusters in OCA-B-
compared to empty vector-transduced cells. *Foxo1* was not over-represented in either of the
equivalent empty vector clusters but was enriched in OCA-B cluster 3. Feature plots showing
representative genes in the two sets of clusters are shown in Figure 4C. These findings suggest that
OCA-B transduced cells in cluster 3, although not increased in numbers, are qualitatively superior to
their empty vector-transduced counterparts at forming memory.

Clustering empty vector- and OCA-B-transduced cells together allows the relative proportions
of cells with common gene expression features to be determined. Clustering together also generated
rare clusters of contaminating CD8 T cell and myeloid cells (Figure 4D clusters 7, 8 and 9). Other
clusters were enriched for likely effector cells (clusters 0, 1, 4 and 6), proliferating effector cells
(cluster 5) and effector cells with memory progenitor potential (clusters 2 and 3). Feature plots for key
genes are shown in Figure 4E. Cluster 2, which was associated with expression of genes associated
with both effector and memory activity (e.g. *Ifng*, *Zeb2*) harbored greater that 3-fold more cells in the
OCA-B expressing condition (33% vs. 10%), however cluster 3 was also associated with memory
(e.g., expressing *Tcf7* and *Il7r* at high levels, Figure 4E) yet contained proportionally fewer than half
as many cells as controls. Although there were fewer cells in this cluster, analysis of gene expression
levels (Figure 4F) showed that *Il7r*, *Zeb2* and *Gadd45b* gene expression were all elevated in this
cluster in OCA-B-transduced comparted to empty vector-transduced cells. *Foxo1*, *Tcf7* and *Tox* by

contrast were expressed at similar levels. Together these findings identify quantitative and qualitative

changes in gene expression and effector T cell populations that predict increased memory potential.

OCA-B knock-in reporter mice fluorescently label CD4$^+$ memory T cell populations

To monitor OCA-B expression in viable cells, we used CRISPR/Cas9 to knock 3 mCherry reporter

cassettes into the mouse *Pou2af1* (*Ocab*) locus upstream of the 3' UTR. The mCherry cassettes were

preceded by P2A elements, allowing for independent translation. *Pou2af1* is compact, spanning just

30 kb. The 3' UTR is ~1kb and lacks significant conservation, making it ideal for reporter cassette

insertion. The resulting reporter allele contained the mCherry cassettes inserted between the stop

codon and the 3' UTR (Figure S4A). A successful heterozygous knock-in founder and a homozygote

generated by intercrossing the progeny were confirmed by PCR genotyping (Figure S4B). Both

animals retained normal OCA-B protein expression, indicating that reporter knock-in homozygosity

does not interfere with endogenous OCA-B protein expression (Figure S4C). We used flow cytometry

to identify peripheral blood mCherry-expressing cells from wild-type, heterozygous and homozygous

reporter mice. The expression and function of OCA-B are best characterized in B lymphocytes

(Betzler et al., 2022; Casellas et al., 2002). OCA-B expression in B cells was readily detectable

(Figure S4D,E). T cells formed a slight shoulder in the heterozygous condition, but were also

detectable in homozygous mice (Figure S4D,E). We therefore used homozygous mice exclusively to

study T cells, and heterozygous mice to perform a brief characterization of B cell populations.

We first profiled OCA-B expression in different B cell populations in unchallenged, specific

pathogen-free heterozygous reporter mice. Robust mCherry expression could be detected in organs

harboring large numbers of developing and mature B cells – bone marrow, peritoneal cavity, spleen

and lymph node (Figure S5A). There was relatively little expression by contrast in total thymocytes. In

the bone marrow, we used antibodies against IgM and B220 to discriminate developing (Pro/Pre),

immature and recirculating B cells. The presence of the reporter allele conferred no major changes in

the relative populations (Figure S5B). The largest change was a 30% drop in the relative percentage

of developing B cells. This difference was unlikely to be due to decreased OCA-B function, as

heterozygous reporter mice express normal levels of OCA-B (Figure S4C) and as whole body OCA-B

knockouts do not show major deficits in developing (immature) B cell numbers and at most 2-fold

differences in mature B cells (Kim et al., 1996; Nielsen et al., 1996; Schubart et al., 1996). OCA-B

expression was readily detected in the three populations (Figure S5B, bottom). Expression was nearly

uniform compared to control wild-type mice lacking the reporter allele, except for a small population of

immature B cells that lack OCA-B expression. In the peritoneal cavity, we used anti-IgM and anti-

B220 and visualize B-1 and B-2 B cells. Again, only small differences in percentages were present

(Figure S5C, top). OCA-B expression was largely uniform, and more robust OCA-B expression was

detected in B-2 relative to B-1 cells (bottom).

Splenic B cells can be partitioned into transitional (immature) and mature populations based

on CD93 (AA4.1) staining. We observed no major changes in total transitional splenic B cells, or in

T1/2/3 subsets, in heterozygous reporter mice compared to wild-type controls (Figure S5D). The

populations expressed OCA-B uniformly and equivalently. Mature splenic B cell populations can be

divided into marginal zone, follicular and non-follicular cells based on CD21 and CD23 expression.

The relative populations again showed few changes in the presence of the reporter allele, and all

three populations expressed OCA-B strongly (Figure S5E). Expression was largely uniform, though

there was an unexpected significant population of newly-formed CD21^{lo}CD23^{lo} B cells expressing no

OCA-B. These results document strong reporter activity in B cell populations, with the highest level of

activity in the more innate-like marginal zone and B-1 B cells. These results indicate that the OCA-B

reporter robustly labels B cells, and has negligible effects on their relative abundances.

We then characterized OCA-B expression in T cells from unchallenged, specific pathogen-free

homozygous reporter mice. Gating out B cells, approximately 5% of thymocytes expressed some level

of mCherry (Figure S6A). Nevertheless, reporter thymi show normal distributions of double-positive

and CD4/CD8 single-positive cells (Figure S6B), consistent with findings that OCA-B deletion does

not affect T cell development (Shakya *et al.*, 2015). mCherry expression could be identified in all

CD4/CD8 double-negative, double-positive and single positive compartments, with the highest

amounts seen in double-negative and CD8 single-positive cells (Figure S6B). In the spleen, we tested

an allelic series of heterozygous and homozygous reporter alleles. As expected, the relative

abundance of CD4^{+} and CD8^{+} T cells was unaltered (Figure 5A). As expected, splenic B cells robustly

and nearly uniformly express mCherry (Figure 5B). CD4^{+} T cells express detectable amounts of

mCherry but at 10- 100-fold lower levels compared to B cells (Figure 5C). Interestingly, a greater

proportion of total CD8^{+} T cells in specific pathogen-free mice expressed OCA-B (Figure S7A).

Additionally, the range of mCherry expression was wider in T cells compared to B cells. While B cell

intensity varied over a 6-fold range (Figure 5B), T cells in homozygous knock-in mice varied over 50-

fold (Figure 5C, right panel and Figure S7A). Peripheral blood T cells varied over a similar range (not

shown).

We used antibodies against CD62L and CD44 to stratify resting splenic CD4^{+} and CD8^{+} cells

into naïve, activated/effector and central memory compartments. Unexpectedly, reporter activity could

be detected in approximately 7% of naïve (CD62L^{hi}CD44^{lo}) cells (Figure 5D). The nature of these cells

(e.g., Tregs, recent thymic emigrants) is unknown. A larger fraction of activated CD62L^{lo}CD44^{hi} cells

(37%) showed reporter activity, while CD62L^{lo} cells that had not yet up-regulated CD44 showed levels

intermediate between naïve and activated. CD62L^{hi}CD44^{hi} T$_{CM}$ cells showed the highest percentage

of reporter activity, with half the cells expressing mCherry (Figure 5D). These results indicate that

some but not all T$_{CM}$ cells continue to express OCA-B. In addition, mCherry mean fluorescence

intensity (MFI) increased as cells progressed from naïve to activated to T$_{CM}$ (Figure S7B). CD8^{+} T

cells showed comparatively higher expression levels in naïve cells, but decreased expression in the

activated state and similar expression in T_{CM} cells (Figure 5E).

Elevated OCA-B expression prospectively identifies memory precursor CD4$^+$ T cells

We used the reporter mice together with LCMV infection and MHC class I and class II tetramers

recognizing immunodominant LCMV epitopes to assess OCA-B responses during infection,

subsequent memory formation and rechallenge. Mice were sacrificed at each timepoint to assess

mCherry expression splenic T cells. Prior to infection, >90% of cells were naïve CD44^{lo} phenotype,

with 13.7% of these cells and 49.5% of the remaining CD44^{hi} cells expressing mCherry (Figure 6A,

D0). These results are comparable to Figure 5D. At 8 days post-infection, significant numbers of

tetramer-positive effector cells accumulated, and of these 50% were mCherry-positive, similar to

tetramer-negative CD44^{hi} cells (D8). Memory timepoint mice showed even higher percentages of

mCherry$^+$ cells (D43, 61.6%), though with a lower degree of expression per cell. We also rechallenged

these mice with Lm-gp61 to assess antigen recall responses. Interestingly, responding CD44^{hi} CD4$^+$ T

cells reduced mCherry positivity compared to resting memory, and high-affinity tetramer-positive cells

showed significant mCherry diminution, declining from >60% to <10% positivity (Figure S8A, D42+7).

587   In contrast to CD4$^+$ mCherry reporter T cells, CD8$^+$ reporter T cells showed significant

differences. A higher percentage of D0 CD44^{lo} and CD44^{hi} cells express mCherry, and there were

detectably higher percentages of tetramer-positive fluorescent cells at day 5, but by day 8 (the peak of

the primary T cell LCMV response) mCherry levels were greatly reduced. Memory timepoints also

showed reduced levels of mCherry fluorescence in antigen-specific cells relative to CD4$^+$ T cells

(Figure S8B). We did not collect data on CD8$^+$ rechallenge because Lm-gp61 does not express the

CD8-dominant epitope recognized by H2-D^b LCMV tetramers.

594   To determine if elevated OCA-B levels in T cells responding to LCMV are associated with

increased memory potential, we crossed the reporter allele onto a SMARTA TCR transgenic

background to fix the TCR specificity and onto Ly5.1 as a means of tracking cells. We then adoptively

transferred congenically marked SMARTA cells into recipient animals, infected with LCMV, isolated

mCherry^{hi} and mCherry^{lo} cells at day 8 post-infection, and transferred large numbers of cells (8×10^5)

into naïve host animals. After 8 days, we isolated splenic CD4+ T cells to assess donor cell

frequencies (Figure 6A). Significantly more cells were recovered from recipient animals engrafted with

mCherry^{hi} cells, indicating that the rate of loss of these cells is lower than for control mCherry^{lo} cells

(Figure 6B,C). Averaged percentages of splenic SMARTA cells in this experiment were trending

towards statistical significance but did not reach $p<0.05$ (not shown). Interestingly, the degree of

mCherry fluorescence in the cells that did survive was the same in both groups, though with

considerably more variance associated with mCherry^{lo} donor cells (Figure 6D) These findings strongly

suggest that effector CD4^{+} cells expressing higher levels of OCA-B undergo less cell death.

To test the ability of the cells expressing OCA-B to mount superior antigen recall responses,

we similarly transferred congenically marked SMARTA cells into recipient animals, infected with

LCMV, but isolated mCherry^{hi} and mCherry^{lo} cells at a memory timepoint after viral clearance (day

43). Smaller numbers (4×10^{4}) of memory SMARTA cells were then transferred into naïve hosts which

were challenged with LCMV and analyzed at day 7 (Figure 6E). Recipient mice engrafted with

mCherry^{hi} SMARTA cells and infected with LCMV mounted better recall responses compared to

mCherry^{lo} (Figure 6F,G). Again, there was little difference in mCherry fluorescence in the responding

cells (Figure 6H), indicating a plasticity in OCA-B expression whereby expression normalizes following

antigen re-encounter. Cumulatively, these results indicate that OCA-B expression labels the central

memory compartment and can be used to enrich memory progenitor cells prospectively.

Discussion

Immunological memory provides long-lasting protection against chronic and opportunistic infection. Strategies that promote memory have the potential to improve both vaccine efficacy and tumor immunotherapy. Multiple transcription factors and epigenetic regulators have been reported to modulate formation or function of CD4$^+$ and CD8$^+$ T$_{CM}$ cells, including Oct1 and OCA-B (Shakya *et al.*, 2015), Blimp1 (Kallies et al., 2009; Rutishauser et al., 2009; Shin et al., 2013), Suv39h1 (Pace et al., 2018), graded levels of T-bet (Joshi *et al.*, 2007; Marshall *et al.*, 2011), co-expressed Tox and TCF1 (Utzschneider *et al.*, 2016), Foxo1 (Hess Michelini et al., 2013; Rao et al., 2012), Bcl6 (Ichii et al., 2007; Pepper et al., 2011), Eomes (Banerjee et al., 2010) and ThPOK (Ciucci *et al.*, 2019). Many of these factors appear to function more precisely in CD8$^+$ compared to CD4$^+$ T cells, and additionally none have been shown to be individually sufficient to selectively promote T cell memory formation or function in vivo. Here we show that loss of the lymphocyte-restricted transcription coactivator OCA-B in T cells confers selective defects in CD4 memory recall response and cytokine production, and that ectopic OCA-B expression does not skew primary effector CD4$^+$ T cell numbers, but is sufficient to promote subsequent memory T cell recall responses. Using an OCA-B reporter mouse strain, we show that elevated levels of endogenous OCA-B in effector T cells during the primary response can be used to prospectively identify memory progenitor CD4$^+$ T cells.

OCA-B is a transcriptional coregulatory protein named for its strong expression in the B cell lineage. OCA-B binding to immunoglobulin variable region promoters can regulate Ig repertoires (Betzler *et al.*, 2022; Casellas *et al.*, 2002). After B cell activation, OCA-B is essential for the generation of germinal centers and the generation of high-affinity antibodies (Kim *et al.*, 1996; Schubart *et al.*, 1996). The requirement for germinal center formation has been shown to be B cell-intrinsic (Qin et al., 1998). Unlike B cells, developing and naïve CD4$^+$ T cells express negligible amounts of OCA-B. Instead, OCA-B becomes expressed following T cell activation (Shakya *et al.*, 2015). Naïve CD8$^+$ cells express OCA-B more strongly compared to CD4$^+$ cells, and expression is further increased upon activation (Kim *et al.*, 2021). OCA-B has been associated with autoimmunity (Ban et al., 2006; Chevrier et al., 2014; Kim *et al.*, 2021; Nakamura et al., 2012).

Prior work used T cell-conditional mice deficient in Oct1, the transcription factor with which OCA-B docks, and competitive mixed bone marrow chimeras to identify decreases in memory potential in T cell lacking Oct1 (Shakya *et al.*, 2015). Oct1 conditional mice directly infected with LCMV showed few differences. This was likely due to selection for cells that fail to delete the conditional allele, as significant bias against recombination could be detected in antigen-specific CD4$^+$ memory T cells at day 8 post-infection compared to antigen-nonspecific T cells in the same animal. The positive results that were obtained with bone marrow chimeras required *fl/Δ* mice with one allele germline-deleted (Shakya *et al.*, 2015). In contrast to the findings with Oct1, T cell-conditional loss of

OCA-B in polyclonal mice directly infected with LCMV[Arm] results in detectable deficiencies in memory

recall responses and IFNγ expression. The finding that direct infection of OCA-B T cell conditional-

deficient mice results in stronger phenotypes vis-à-vis Oct1 suggests a more complete deletion of the

Ocab allele, reduced selection for retention of the undeleted (floxed) allele, and/or stronger biological

effects of OCA-B deletion. The latter may include possible Oct1-independent effects of OCA-B. In

competitive mixed bone marrow chimera experiments, T cell-conditional OCA-B loss showed severe

deficiencies in memory recall responses compared to wild-type competitor bone marrow and

compared to littermate *Ocab* conditional bone marrow lacking the Cre driver.

Using LCMV-specific SMARTA TCR transgenic mice, we also show that ectopic OCA-B

expression promotes CD4$^+$ memory T cell recall responses in vivo. In these experiments, OCA-B is

ectopically expressed prior to pathogen exposure, and cells accumulate at the expense of empty

vector-transduced control cells selectively at memory recall but not primary response timepoints.

Although memory is selectively affected, OCA-B mediates changes in responding CD44^{hi} effector

cells that are predictive of later memory formation. ICOS, PD-1 and CD62L (L-selectin) were slightly

reduced in OCA-B-transduced SMARTA T cells relative to empty vector-transduced cells isolated

from the same animals. *Sell*, which encodes CD62L, is a direct OCA-B target gene, as is *Icos*

(Shakya *et al.*, 2015). We also performed bulk and single-cell RNA-seq to delineate gene expression

changes that may be predictive of later enhanced T$_{CM}$ formation. Bulk RNA-seq identifies a number of

genes whose expression was increased in OCA-B- compared to empty vector-transduced

populations, including *Gadd45b*, *Socs2* and *Zeb2*. These changes may be sufficient to confer

potential increased memory formation, as small advantages compounded over time can result in the

large observed effects. *Gadd45b* promotes autoimmune diseases including lupus, psoriasis and

arthritis(Liu et al., 2005; Luo et al., 2011; Rodriguez-Jimenez et al., 2021), as well as for strong anti-

tumor immune responses(Ju et al., 2009). Also up-regulated is *Acacb*, which is important for fatty acid

synthesis. Memory T cells are known to rely on oxidation of fatty acid stores (Michalek et al., 2011;

Pearce et al., 2009). ChIP-X Enrichment Analysis identifies Tp53 and Smc3, proteins associated with

growth arrest and topological DNA-DNA interactions, as potential regulators of the up-regulated gene

set. Other genes were decreased in bulk RNA-seq, including genes associated with H3K27me3 and

repressed in memory cells. The further repression of these genes is consistent with the memory

advantage conferred by OCA-B expression.

A subset of differentially expressed genes correspond to previously identified OCA-B targets

based on ChIP-seq results using in vitro stimulation of T cells lacking OCA-B (Shakya *et al.*, 2015).

For this analysis, genes down-regulated by OCA-B loss were compared to genes induced by ectopic

expression, and the converse. Using this method, better overlap was identified with genes whose

expression is reduced by OCA-B loss and increased by ectopic expression compared to the converse.

Many of the gene repression events could therefore be indirect, however there are repressed direct

OCA-B targets, e.g. *Sell*, which encodes CD62L. Recent findings that OCA-B can also mediate

repression as well as activation (Lu et al., 2019) are consistent with this possibility. If the activity of

OCA-B switches between activation and repression comparing cells in vitro vs. in vivo, many targets

from this analysis would be missed. More work is needed to uncover mechanisms that dictate OCA-B

transcription output in T cells.

Two other genes associated with T cell survival and memory formation, *Il7r* and *Tcf7* (Joshi *et*

al., 2007; Utzschneider *et al.*, 2016), are counterintuitively decreased in bulk RNA-seq. *Il7r* for

example encodes a cytokine receptor important for T cell survival. Effector CD4$^+$ T cell populations

can manifest heterogeneity, with sub-populations skewed towards later memory formation (Joshi *et*

al., 2007; Snook *et al.*, 2018). Increased *Il7r* expression in memory progenitors for example may be

obscured in bulk RNA-seq by decreases in the larger populations of terminal effector cells, which may

lead to more culling to create greater capacity for memory formation. Consistently, at single-cell

resolution, we identified memory-like cell clusters with increased *Il7r* in the background of global

down-regulation. The memory advantages conferred by OCA-B cannot be solely mediated by

negative factors that cull effector cells, because both T cells ectopically expressing OCA-B and

competing cells expressing empty vector would benefit and show equal representation in recall

responses. Therefore, other effects of OCA-B must reinforce memory formation in a memory cell-

intrinsic manner. Like *Il7r, Tcf7* expression is decreased in bulk RNA-seq but increased in likely

memory progenitor populations to a higher degree than corresponding populations in cells transduced

with empty vector.

We generated an OCA-B knock-in mouse reporter allele, and confirmed robust expression in

different B cell populations in unchallenged mice. Most B cells were almost uniformly labeled,

however there were sub-populations of immature bone marrow and splenic B cells that completely

lack OCA-B expression. The nature and significance of these cells is unknown. Expression was

highest in marginal zone B cells, which are largely absent in OCA-B knockout mice (Samardzic et al.,

2002). We confirm that expression is 50-100-fold lower in T cells overall compared to B cells. Within

the CD4$^+$ T cell compartment, increased OCA-B expression as measured by mCherry fluorescence

progressively labels greater fractions of naïve, activated and central memory CD4$^+$ cells. Upon LCMV

infection, homozygous reporter mice display robust expression of OCA-B, with half of all tetramer-

positive CD4$^+$ T cells showing robust mCherry expression. An even greater fraction of splenic

tetramer-positive cells show reporter expression at memory timepoints, though fluorescence intensity

is lower. This pattern was not recapitulated in CD8$^+$ cells, which showed earlier expression of

mCherry and poor induction in both primary effector, memory and memory recall timepoints. This

pattern is consistent with prior findings that naïve CD8$^+$ T cells express OCA-B at higher levels

compared to naïve CD4$^+$ cells (Kim et al., 2019).

These studies have identified a method of prospectively marking viable CD4$^+$ memory

precursor effector T cells. Such methods are largely lacking compared to CD8$^+$ T cells, where markers

such as IL7R and KLRG1 can be used to identify memory precursor cells (Joshi *et al.*, 2007).

Identifying similar markers and transcriptional regulators that define the memory precursor effector

CD4$^+$ T cells will promote major advances in understanding the formation of CD4 T cell memory. We

isolated OCA-B^{hi} vs OCA-B^{lo} SMARTA transgenic effector CD4$^+$ T cells at 8 days post-infection, which

results in roughly equal partitioning of cells into each group. Populations of effector cells with elevated

OCA-B expression contracted less in naïve animals, and were better able to mount recall responses

to antigenic rechallenge compared to cells with low or absent OCA-B expression. These results

indicate that OCA-B reporter activity can be used to prospectively identify live CD4$^+$ effector cells with

augmented capacity to form central memory cells. Cumulatively, the results indicate that OCA-B

expression within CD4$^+$ T cells is both necessary for, and sufficient to promote, the emergence of T$_{CM}$.

Abbreviations

ChEA, ChIP-X Enrichment Analysis; EV, empty vector; FACS, fluorescence-activated cell sorting; GEMs, Gel-bead-in EMulsions; ICOS, Inducible T cell co-stimulator; LCMV, lymphocytic choriomeningitis virus; Lm-gp61, *Listeria monocytogenes* expressing LCMV gp61-80; PCNA, proliferating cell nuclear antigen; pMIG, pMSCV-IRES-GFP; scRNA-seq, single-cell RNA-sequencing; T_{CM}, central memory T cell; UMAP, uniform manifold approximation and projection; UMIs, Unique Molecular Identifiers

Declarations

Consent for publication

Not applicable

Availability of supporting data

Please contact author for data requests.

Competing interests

The sponsors had no role in study design, collection, analysis and interpretation of data, in the writing of the report or in the decision to submit the article for publication.

Author's contributions

DT conceived the study and designed experiments, supervised the study, and provided administrative, technical and material support. WS, HK and AI acquired and interpreted the data. JP performed bioinformatic analyses. JSH and MAW provided material and intellectual support. All authors were involved in writing, reviewing and revising the manuscript.

Acknowledgements

We thank M. Chandrasekharan for help with graphics. We thank J. Marvin and the University of Utah Health Sciences Center Flow Cytometry Core facility. We thank O. Allen, B. Dalley and the High-Throughput Genomics Core. MHC tetramers were provided by the NIH Tetramer Core Facility (Atlanta, GA). We thank R. Roeder for the mouse OCA-B cDNA and R. M. O'Connell for the pMIGR1 plasmid.

Funding

This work was supported by NIH grant AI100873 to DT.

 Figure legends

Figure 1. OCA-B in T cells is required for robust polyclonal memory CD4$^+$ T cell formation and
cytokine expression following LCMV infection.

A) *Ocab$^{fl/fl}$*;CD4-Cre and control *Ocab$^{fl/fl}$* mice were infected with LCMV^{Arm}, and rechallenged with Lm-
gp61 40 days after LCMV infection. Splenic I-A^b/gp$_{66-77}$ tetramer-positive, CD4$^+$CD44$^+$ T cells were
analyzed by flow cytometry 8 days after primary infection and 7 days after heterologous rechallenge.
n=5 for each group. n.s.=not significant.

B) Splenocytes from LCMV-infected *Ocab$^{fl/fl}$*;CD4-Cre or control *Ocab$^{fl/fl}$* mice were restimulated in
vitro with gp$_{61-80}$ peptide for 4 hours, and IFNγ and IL-2 expression in CD4$^+$ T cells examined by flow
cytometry. Cells were gated based on I-A^b/gp$_{66-77}$ tetramer, CD4 and CD44 positivity. Data from
representative animals are shown.

C) Mean T cell frequencies of CD4$^+$CD44$^+$IFNγ^+IL-2$^-$ T cells are shown. n=5 for each group.

D) Experimental schematic for mixed bone marrow reconstitution and LCMV challenge. Bone marrow
cells were depleted of mature T cells prior to engraftment (see methods).

E) Flow cytometry plot showing Thy1.1$^+$Ly5.2$^+$ but otherwise genetically wild-type (WT) donor bone
marrow cells engrafted 50:50 with Ly5.2$^+$ control cells lacking the Cre driver (*fl/fl*) or experimental T
cell-conditional *Ocab$^{fl/fl}$*;CD4-Cre donor cells (*fl/fl*;Cre). Recipient hosts were Ly5.1$^+$. Splenic effector
cells were recovered at primary response (D8) or seven days post-rechallenge (D42+7). Cells were
gated based on I-A^b/gp$_{66-77}$ tetramer positivity. Example animals are shown.

F) Mean wild-type competitor to control of OCA-B deficient T cell ratios are shown. n=3 for each
group.

Figure 2. Ectopic OCA-B expression enhances CD4$^+$ T cell memory recall responses in vivo.

A) Experimental schematic for OCA-B transduction and T cell mixed adoptive transfer. Ly5.1$^+$
SMARTA cells were transduced with pMSCV-IRES-GFP (pMIG) empty vector (EV) and Thy1.1$^+$
SMARTA cells were transduced with pMIG expressing mouse OCA-B (pMIG-OCA-B). Following
transduction, the two populations of SMARTA cells were combined 1:1, co-transferred into
Ly5.2$^+$/Thy1.2$^+$ C57BL6/J recipient mice, and infected with LCMV^{Arm}. Recipients were rechallenged
with Lm-gp61 on either day 52 or 110 post-LCMV infection, and in each case analyzed after 7 days.

B) Lysates from 3T3 cells transduced with either empty pMIG empty vector (EV) or pMIG-OCA-B
were immunoblotted using antibodies against OCA-B. α-tubulin is shown as an internal loading
standard.

C) Flow cytometric analysis of Ly5.1$^+$ pMIG empty vector-transduced (EV) and Thy1.1$^+$ pMIG-OCA-B-
transduced (OCA-B) SMARTA cells in the spleen and blood of a representative recipient mouse at

peak effector response (D8), peak memory recall response (52+7) and long-term memory recall

(D110+7). Cells were first gated based on CD44, CD4 and GFP positivity.

D) Mean relative percentages of splenic and blood total EV- or pMIG-OCA-B-transduced (GFP$^+$)

SMARTA cells were plotted at D8, D52+7 and 110+7. Each datapoint represents an individual mouse

harboring both EV- or pMIG-OCA-B-transduced cells. 5, 7 and 10 mice were sacrificed at each

timepoint to obtain measurements from spleen. All mice were bled at D8, and remaining mice were

bled at subsequent timepoints.

E) Mean ratios of pMIG-OCA-B-transduced relative to empty vector-transduced SMARTA cells (OCA-

B/EV) are plotted. Splenic (top) and peripheral blood (bottom) SMARTA T cells are shown at day 8

post-infection, or 7 days post-rechallenge with Lm-gp61 (52+7 or 110+7).

F) Flow cytometry showing relative percentages of splenic GFP$^+$CD44^{hi} EV-transduced (blue) or

pMIG-OCAB-transduced (red) SMARTA cells expressing ICOS, PD-1, Ly6C or CD62L at day 8 post-

infection. Representative data showing the two populations of cells within a single mouse are shown

at day 8 post-infection.

G) Percent expression for ICOS, PD-1, CD62L and Ly6C was quantified from four mice using splenic

EV- and pMIG-OCAB-transduced GFP$^+$CD44^{hi} SMARTA cells at 8 days post-infection. Four replicates

were performed. For CD26L, the four datapoints are superimposed, and for Ly6C two of the

datapoints are superimposed. Individual mice are connected by lines.

Figure 3. Gene expression changes associated with ectopic OCA-B expression in SMARTA

transgenic primary effector T cells.

A) Heatmap showing top up- and down-regulated genes. Genes with log2 fold-change >0.5 or <-1.0,

and padj ≤ 0.05 are shown. Example genes are shown at right. Control and OCA-B-transduced cells

were purified from the same mice using magnetic isolation and FACS.

B) Example genome tracks showing two up-regulated genes, *Socs2* and *Gadd45b*. *Pou2af1* (*Ocab*) is

shown as a positive control.

C) Top Epigenomic roadmap and Molecular function GO terms for the set of OCA-B up- and down-

regulated genes.

D) Top ChIP-X Enrichment Analysis (ChEA 2016) factors potentially regulating the same set of genes

up- and down-regulated by OCA-B ectopic expression.

Figure 4. Single-cell RNA-seq analysis of SMARTA cells transduced with OCA-B or empty

vector.

A) UMAP projections of independently clustered empty vector-transduced (GFP$^+$) SMARTA cells

purified similar to Figures 2 and 3, and subjected to scRNA-seq. Cluster identities were annotated

using a combination of gene expression enrichment (Table S5), cluster identity predictor ImmGen

identity scores (Table S6) and PanglaoDB annotation terms. Clusters 1 and 4 were associated with

memory formation. Top PanglaoDB annotation terms for cluster 4 are shown below.

B) Similar UMAP projection using OCA-B-transduced cells purified from the same mice as in (A).

Cluster identities were annotated similarly to (A). Cluster 3 in OCA-B-transduced cells was most

strongly associated with memory formation. Top PanglaoDB annotation terms for cluster 3 are shown

below.

C) Feature plots highlighting expression of representative genes for the two UMAP projections in (A)

and (B).

D) UMAP projections of the the EV- and OCA-B-transduced single-cell RNA-seq datasets clustered

together. For each projection, the percentage of each cluster relative to the total number cells in

condition is shown. Cluster identities were annotated using a combination of gene expression

enrichment (Table S7), cluster identity predictor ImmGen identity scores and PanglaoDB annotation

terms.

E) Feature plots highlighting expression of representative genes for the combined projections.

F) Violin plots depicting expression of six genes (*Il7r*, *Zeb2*, *Gadd45b*, *Foxo1*, *Tcf7* and *Tox*) across

each of the 10 clusters in EV-transduced cells (blue, left) and OCA-B-transduced cells (red, right).

Figure 5. mCherry expression in splenic T cell populations in specific pathogen-free

homozygous OCA-B-3×mCherry mice.

A) CD4 and CD8 expression are shown for an allelic series of example wild-type (+/+), heterozygous

(+/KI) and homozygous OCA-B knock-in reporter mice.

B) OCA-B reporter expression in splenic B cells using the same allelic series of mice as in (A). A more

complete analysis of B cell populations is shown in Figure S5.

C) OCA-B reporter expression in splenic CD4$^+$ T cell populations from the same allelic series as in

(A).

D) The same CD4$^+$ cells as in (C) were further stratified by CD62L and CD44 into naïve effector (Teff)

and central memory (Tcm) populations. CD62L^{lo}CD44$_{lo}$ cells are likely recently activated cells that

have down-regulated CD62L but not yet up-regulated CD44.

E) Similar analysis as in (D) except for CD8$^+$ cells.

Figure 6. CD4$^+$ T cells expressing high levels of OCA-B reporter activity preferentially form

central memory cells.

A) Schematic for assessing contraction of mCherry^{hi} vs. mCherry^{lo} SMARTA T cells following LCMV

infection. Congenically marked SMARTA T cells were isolated from donor mice, transferred into naïve

secondary recipients and infected with LCMV. At 8 days post-infection, mCherry[hi] and mCherry[lo]

populations were isolated by FACS. 8×10^5 SMARTA cells were transferred into naïve recipients. After

8 days, splenic CD4[+]Ly5.1[+]5.2[+] SMARTA cells were evaluated by flow cytometry.

B) Flow cytometric analysis of SMARTA donor T cells after 8 days in a naïve animal to monitor rates

of decline. Representative animals engrafted with mCherry[lo] (left) or mCherry[hi] (right) cells are shown.

C) Quantification of averaged donor T cell numbers from mice engrafted with mCherry[lo] or mCherry[hi]

cells. N=7 for the mCherry[lo] group and N=5 for the mCherry[hi] group.

D) Similar analysis to (C) except quantifying mean mCherry fluorescence intensity.

E) Schematic for assessing recall responses of mCherry[hi] vs. mCherry[lo] SMARTA T cells following

LCMV infection. Congenically marked SMARTA T cells were isolated from donor mice, transferred

into naïve secondary recipients and infected with LCMV as in (A). However, mice were allowed to

clear LCMV and form memory. After 43 days, mCherry[hi] and mCherry[lo] memory T cell populations

were isolated by FACS. 1.3×10^4 SMARTA cells were transferred into naïve recipients, which were

infected with LCMV one day later. 7 days post-infection, splenic CD4[+]Ly5.1[+]5.2[+] SMARTA cells were

evaluated by flow cytometry.

F) Flow cytometric analysis of SMARTA donor T cells 7 days post-rechallenge to monitor recall

responses. Representative animals engrafted with mCherry[lo] (left) or mCherry[hi] (right) cells are

shown.

G) Quantification of averaged donor T cell percentages and numbers from mice engrafted with

mCherry[lo] or mCherry[hi] cells. N=7 for the mCherry[lo] group and N=5 for the mCherry[hi] group.

H) Similar analysis to (G) except quantifying mean mCherry fluorescence intensity.

Supplemental Figures

Figure S1, related to Figure 1. Surface marker expression in polyclonal OCA-B T cell-deficient

and control splenic CD4[+] effector T cells at 8 days post-infection or 7 days post-rechallenge.

Example flow cytometry panels of ICOS, Ly6C, CD62L. PD-1 was additionally performed for memory

recall responses. Cells were gated for I-A[b]-LCMV[gp66-77] positivity.

Figure S2, related to Figure 2. Bulk RNA-seq replicate correlations and example genome

tracks.

Euclidian distance clustering of bulk RNA-seq replicates to show divergence between replicates and

conditions.

Figure S3, related to Figure 4. Quality control *Cd3e* and *Cd4* mRNA expression in separately

clustered single-cell RNA-seq.

A) Feature plots showing gene expression intensities for *Cd3e* (top) and *Cd4* (bottom) in EV-transduced (left) and OCA-B-transduced (right) SMARTA T cells at day 8 post-infection with LCMV.

B) Violin plots showing *Cd4* gene expression by cluster for each condition.

Figure S4, related to Figure 5. OCA-B-3×mCherry knock-in reporter mice.

A) Mouse *Pou2af*1 locus, targeting vector and the targeted *Pou2af1* locus.

B) PCR genotyping OCAB-3×mCherry reporter mice. PCR yields a 494 bp product for the wild-type *Pou2af1* allele and a 373 bp product for targeted *Pou2af1* allele. +=wild-type allele. KI=knock-in allele.

C) OCA-B immunoblot using splenocytes from wild-type (+/+) and heterozygous (+/KI) or homozygous knock-in OCAB-3×mCherry knock-in mice. β-actin was used as a loading control.

D) Flow cytometry showing mCherry expression in peripheral blood cells of wild-type mice (+/+), or either heterozygous (+/KI) or homozygous (KI/KI) OCAB-3×mCherry knock-in reporter mice.

E) Similar flow cytometry plots as in (D) except gated on the indicated populations.

Figure S5, related to Figure 5. Characterization OCA-B expression in different B cell populations using heterozygous reporter mice.

A) mCherry (OCA-B) expression in a 4.5 week-old heterozygous female OCA-B-3×mCherry reporter mouse. Cells were harvested from bone marrow (BM), peritoneal cavity (PC), spleen (SP), lymph node (LN) or thymus. After removal of red blood cells, mCherry expression was measured in total cell populations.

B) Wild-type (+/+) and heterozygous knock-in (+/KI) bone marrow cells were further stratified into B1 and B2 populations based on B220 and IgM expression Top plots show relative proportions of B1 (IgM^{hi}B220^{lo}) B2 (IgM^{lo}B220^{hi}) cells in wild-type and two example heterozygous reporter mice. Bottom plots show mCherry expression in gated populations.

C) Analysis of peritoneal cavity B cell populations from +/+ and +/KI reporter mice. Cells were stratified into pro/pre, immature and recirculating B cell compartments based on B220 and IgM expression. Top plots show relative proportions of the cells in animals of the indicated genotypes. Bottom plots show mCherry levels in the gated populations.

D) Splenic transitional B cells were stratified using CD93 (AA4.1), CD23 and IgM antibodies. One wild-type and two example heterozygous reporter mice are shown.

E) Similar analysis to (D) except using CD21 and CD23 antibodies to visualize follicular and marginal zone B cell populations.

Figure S6, related to Figure 5. OCA-B reporter expression in thymocytes.

A) mCherry expression in total B220-negative thymocytes from wild-type (+/+) or homozygous knock-in reporter mice was analyzed by flow cytometry. Specific pathogen-free mice were used.

B) Thymocytes were stained with CD4 and CD8 antibodies to visualize double-negative, double-positive and single-positive T cell developmental stages in homozygous knock-in reporter mice. Right: mCherry expression in developing reporter T cells was subdivided by quadrant.

Figure S7, related to Figure 5. OCA-B reporter expression in total splenic CD4$^+$ and CD8$^+$ T cells.

A) mCherry expression was tested in spleens from 9 week-old male homozygous knock-in reporter mice. Specific pathogen-free mice were used.

B) mCherry mean fluorescence intensity was calculated from four 9 week-old male homozygous knock-in reporter mice using flow cytometry and gating identical to Figure 5D and E.

Figure S8, related to Figure 6. OCA-B reporter expression in antigen-responsive T cells following LCMV infection.

A) Flow cytometry images are shown of homozygous OCA-B-3×mCherry knock-in reporter mice at different timepoints following LCMV infection (D0-43) and following rechallenge with Lm-gp61. Mice were challenged on different days and spleens were harvested and processed on the same day. Top panels: percentages of cells expressing CD44 and class II LCMV tetramer (I-A^b LCMV$^{gp66-77}$) are shown. Bottom panels: for the same timepoints cells gated from each quadrant in the top panels are plotted. Each flow panel is representative of biological duplicate mice sacrificed at each timepoint.

B) Cells from the same mice as above were gated for CD8 expression and stained with class I LCMV tetramers (H2-D^b LCMV$^{gp33-41}$). Analysis of recall response timepoints was not possible for antigen-specific CD8 cells because animals were rechallenged with Lm-gp61, which lacks CD8 immunodominant epitopes.

Supplemental Tables

Table S1, related to Figure 3. Differentially expressed bulk RNAseq genes comparing GFP$^+$ SMARTA T cells expressing empty vector or OCA-B at day 8 post-LCMV infection (cutoffs: -1>log2FC>0.5, *p*<0.05).

Table S2, related to Figure 3. Gene Ontology analysis using differentially expressed bulk RNA-seq transcripts comparing GFP$^+$ SMARTA T cells transduced with empty vector or OCA-B at day 8 post-LCMV infection (cutoffs: -1>log2FC>0.5, *p*<0.05).

Table S3, related to Figure 3. Intersection of differentially expressed bulk RNAseq genes comparing GFP⁺ SMARTA T cells transduced with OCA-B relative to empty vector at 8 days post-LCMV infection (cutoffs: -1>log2FC>0.5, _p_<0.05) with differentially expressed genes in restimulated T cells lacking OCA-B, published previously (Shakya et al. 2015).

Table S4, related to Figure 3. Intersection between previously published Oct1/OCA-B targets (Shakya et al. 2015, 3173 targets*) and bulk RNA-seq comparing GFP⁺ SMARTA T cells transduced with empty vector or OCA-B at day 8 post post-LCMV infection (cutoffs: - 1>log2FC>0.5, _p_<0.05).

Table S5, related to Figure 4. Single-cell RNA-seq gene enrichments by cluster in GFP⁺ SMARTA T cells transduced with empty vector or OCA-B at day 8 post-LCMV infection.

Table S6, related to Figure 4. Top 5 single-cell RNA-seq ImmGen identity scores per cluster using GFP⁺ SMARTA T cells transduced with empty vector or OCA-B at day 8 post-LCMV infection.

Table S7, related to Figure 4. Integrated gene enrichment analysis of single-cell RNA-seq gene enrichments by cluster in GFP⁺ SMARTA T cells transduced with empty vector or OCA-B at day 8 post-LCMV infection.

References

Ahmed, R., Salmi, A., Butler, L.D., Chiller, J.M., and Oldstone, M.B. (1984). Selection of genetic variants of lymphocytic choriomeningitis virus in spleens of persistently infected mice. Role in suppression of cytotoxic T lymphocyte response and viral persistence. J Exp Med *160*, 521-540. 10.1084/jem.160.2.521.

Ban, M., Booth, D., Heard, R., Stewart, G., Goris, A., Vandenbroeck, K., Dubois, B., Laaksonen, M., Ilonen, J., Alizadeh, M., et al. (2006). Linkage disequilibrium screening for multiple sclerosis implicates JAG1 and POU2AF1 as susceptibility genes in Europeans. J Neuroimmunol *179*, 108-116.

Banerjee, A., Gordon, S.M., Intlekofer, A.M., Paley, M.A., Mooney, E.C., Lindsten, T., Wherry, E.J., and Reiner, S.L. (2010). Cutting edge: The transcription factor eomesodermin enables CD8+ T cells to compete for the memory cell niche. J Immunol *185*, 4988-4992. jimmunol.1002042 [pii] 10.4049/jimmunol.1002042 [doi].

Betzler, A.C., Fiedler, K., Hoffmann, T.K., Fehling, H.J., Wirth, T., and Brunner, C. (2022). BOB.1/OBF.1 is required during B-cell ontogeny for B-cell differentiation and germinal center function. Eur J Immunol *52*, 404-417. 10.1002/eji.202149333.

Blattman, J.N., Antia, R., Sourdive, D.J., Wang, X., Kaech, S.M., Murali-Krishna, K., Altman, J.D., and Ahmed, R. (2002). Estimating the precursor frequency of naive antigen-specific CD8 T cells. J Exp Med *195*, 657-664. 10.1084/jem.20001021.

Cafri, G., Yossef, R., Pasetto, A., Deniger, D.C., Lu, Y.C., Parkhurst, M., Gartner, J.J., Jia, L., Ray, S., Ngo, L.T., et al. (2019). Memory T cells targeting oncogenic mutations detected in peripheral blood of epithelial cancer patients. Nat Commun *10*, 449. 10.1038/s41467-019-08304-z.

Casellas, R., Jankovic, M., Meyer, G., Gazumyan, A., Luo, Y., Roeder, R., and Nussenzweig, M. (2002). OcaB Is Required for Normal Transcription and V(D)J Recombination of a Subset of Immunoglobulin kappa Genes. Cell *110*, 575.

Chee, J., Ko, H.J., Skowera, A., Jhala, G., Catterall, T., Graham, K.L., Sutherland, R.M., Thomas, H.E., Lew, A.M., Peakman, M., et al. (2014). Effector-memory T cells develop in islets and report islet pathology in type 1 diabetes. J Immunol *192*, 572-580.

Chevrier, S., Kratina, T., Emslie, D., Karnowski, A., and Corcoran, L.M. (2014). Germinal center-independent, IgM-mediated autoimmunity in sanroque mice lacking Obf1. Immunol Cell Biol *92*, 12-19.

Ciucci, T., Vacchio, M.S., Gao, Y., Tomassoni Ardori, F., Candia, J., Mehta, M., Zhao, Y., Tran, B., Pepper, M., Tessarollo, L., et al. (2019). The Emergence and Functional Fitness of Memory CD4(+) T Cells Require the Transcription Factor Thpok. Immunity *50*, 91-105 e104. 10.1016/j.immuni.2018.12.019.

Crawford, A., Angelosanto, J.M., Kao, C., Doering, T.A., Odorizzi, P.M., Barnett, B.E., and Wherry, E.J. (2014). Molecular and transcriptional basis of CD4(+) T cell dysfunction during chronic infection. Immunity *40*, 289-302. 10.1016/j.immuni.2014.01.005.

Elyaman, W., Kivisakk, P., Reddy, J., Chitnis, T., Raddassi, K., Imitola, J., Bradshaw, E., Kuchroo, V.K., Yagita, H., Sayegh, M.H., and Khoury, S.J. (2008). Distinct functions of autoreactive memory and effector CD4+ T cells in experimental autoimmune encephalomyelitis. Am J Pathol *173*, 411-422. 10.2353/ajpath.2008.080142.

Guan, T., Dominguez, C.X., Amezquita, R.A., Laidlaw, B.J., Cheng, J., Henao-Mejia, J., Williams, A., Flavell, R.A., Lu, J., and Kaech, S.M. (2018). ZEB1, ZEB2, and the miR-200 family form a counterregulatory network to regulate CD8(+) T cell fates. J Exp Med *215*, 1153-1168. 10.1084/jem.20171352.

Hess Michelini, R., Doedens, A.L., Goldrath, A.W., and Hedrick, S.M. (2013). Differentiation of CD8 memory T cells depends on Foxo1. J Exp Med *210*, 1189-1200. jem.20130392 [pii] 10.1084/jem.20130392 [doi].

Hurton, L.V., Singh, H., Najjar, A.M., Switzer, K.C., Mi, T., Maiti, S., Olivares, S., Rabinovich, B., Huls, H., Forget, M.A., et al. (2016). Tethered IL-15 augments antitumor activity and promotes a stem-cell

memory subset in tumor-specific T cells. Proc Natl Acad Sci U S A *113*, E7788-e7797.
10.1073/pnas.1610544113.
Ichii, H., Sakamoto, A., Arima, M., Hatano, M., Kuroda, Y., and Tokuhisa, T. (2007). Bcl6 is essential
for the generation of long-term memory CD4+ T cells. Int Immunol *19*, 427-433. dxm007 [pii]
10.1093/intimm/dxm007 [doi].

Joshi, N.S., Cui, W., Chandele, A., Lee, H.K., Urso, D.R., Hagman, J., Gapin, L., and Kaech, S.M.
(2007). Inflammation directs memory precursor and short-lived effector CD8(+) T cell fates via the
graded expression of T-bet transcription factor. Immunity *27*, 281-295.
Ju, S., Zhu, Y., Liu, L., Dai, S., Li, C., Chen, E., He, Y., Zhang, X., and Lu, B. (2009). Gadd45b and
Gadd45g are important for anti-tumor immune responses. Eur J Immunol *39*, 3010-3018.
10.1002/eji.200839154.
Kallies, A., Xin, A., Belz, G.T., and Nutt, S.L. (2009). Blimp-1 transcription factor is required for the
differentiation of effector CD8(+) T cells and memory responses. Immunity *31*, 283-295.
Kawakami, N., Odoardi, F., Ziemssen, T., Bradl, M., Ritter, T., Neuhaus, O., Lassmann, H., Wekerle,
H., and Flugel, A. (2005). Autoimmune CD4+ T cell memory: lifelong persistence of encephalitogenic
T cell clones in healthy immune repertoires. J Immunol *175*, 69-81.
Kim, C., Jay, D.C., and Williams, M.A. (2012). Stability and function of secondary Th1 memory cells
are dependent on the nature of the secondary stimulus. J Immunol *189*, 2348-2355.
10.4049/jimmunol.1200244.
Kim, H., Dickey, L., Stone, C., Jafek, J.L., Lane, T.E., and Tantin, D. (2019). T cell-selective deletion
of Oct1 protects animals from autoimmune neuroinflammation while maintaining neurotropic pathogen
response. J. Neuroinflamm. *16*. ARTN 133 10.1186/s12974-019-1523-3.
Kim, H., Perovanovic, J., Shakya, A., Shen, Z., German, C.N., A., I., Jafek, J.L., Lin, N.-P., Evavold,
B.D., Chou, D.H., et al. (2021). Targeting OCA-B/Pou2af1 blocks activated, autoreactive T cells in the
pancreas and type-1 diabetes. J Exp Med *218*, e20200533. 10.1084/jem.20200533.
Kim, U., Qin, X.F., Gong, S., Stevens, S., Luo, Y., Nussensweig, M., and Roeder, R.G. (1996). The B-
cell-specific transcription coactivator OCA-B/OBF-1/Bob-1 is essential for normal production of
immunoglobulin isotypes. Nature *383*, 542-547.
Lachmann, A., Xu, H., Krishnan, J., Berger, S.I., Mazloom, A.R., and Ma'ayan, A. (2010). ChEA:
transcription factor regulation inferred from integrating genome-wide ChIP-X experiments.
Bioinformatics *26*, 2438-2444. 10.1093/bioinformatics/btq466.
Laidlaw, B.J., Craft, J.E., and Kaech, S.M. (2016). The multifaceted role of CD4(+) T cells in CD8(+) T
cell memory. Nat Rev Immunol *16*, 102-111. 10.1038/nri.2015.10.
Liu, L., Tran, E., Zhao, Y., Huang, Y., Flavell, R., and Lu, B. (2005). Gadd45 beta and Gadd45
gamma are critical for regulating autoimmunity. J Exp Med *202*, 1341-1347. 10.1084/jem.20051359.
Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. Genome Biol *15*, 550.
Lu, X., Chu, C.S., Fang, T., Rayon-Estrada, V., Fang, F., Patke, A., Qian, Y., Clarke, S.H., Melnick,
A.M., Zhang, Y., et al. (2019). MTA2/NuRD Regulates B Cell Development and Cooperates with
OCA-B in Controlling the Pre-B to Immature B Cell Transition. Cell Rep *28*, 472-485 e475.
10.1016/j.celrep.2019.06.029.
Luo, Y., Boyle, D.L., Hammaker, D., Edgar, M., Franzoso, G., and Firestein, G.S. (2011). Suppression
of collagen-induced arthritis in growth arrest and DNA damage-inducible protein 45beta-deficient
mice. Arthritis Rheum *63*, 2949-2955. 10.1002/art.30497.
Luo, Y., and Roeder, R.G. (1995). Cloning, functional characterization, and mechanism of action of
the B-cell-specific transcriptional coactivator OCA-B. Mol. Cell. Biol. *15*, 4115-4124.
Marshall, H.D., Chandele, A., Jung, Y.W., Meng, H., Poholek, A.C., Parish, I.A., Rutishauser, R., Cui,
W., Kleinstein, S.H., Craft, J., and Kaech, S.M. (2011). Differential expression of Ly6C and T-bet
distinguish effector and memory Th1 CD4(+) cell properties during viral infection. Immunity *35*, 633-
646.
Michalek, R.D., Gerriets, V.A., Jacobs, S.R., Macintyre, A.N., MacIver, N.J., Mason, E.F., Sullivan,
S.A., Nichols, A.G., and Rathmell, J.C. (2011). Cutting edge: distinct glycolytic and lipid oxidative

metabolic programs are essential for effector and regulatory CD4+ T cell subsets. J Immunol *186*,
3299-3303. 10.4049/jimmunol.1003613.
Mitchell, D.M., Ravkov, E.V., and Williams, M.A. (2010). Distinct roles for IL-2 and IL-15 in the
differentiation and survival of CD8+ effector and memory T cells. J Immunol *184*, 6719-6730.
jimmunol.0904089 [pii] 10.4049/jimmunol.0904089 [doi].
Murali-Krishna, K., Altman, J.D., Suresh, M., Sourdive, D.J., Zajac, A.J., Miller, J.D., Slansky, J., and
Ahmed, R. (1998). Counting antigen-specific CD8 T cells: a reevaluation of bystander activation
during viral infection. Immunity *8*, 177-187. 10.1016/s1074-7613(00)80470-7.
Nakamura, M., Nishida, N., Kawashima, M., Aiba, Y., Tanaka, A., Yasunami, M., Nakamura, H.,
Komori, A., Nakamuta, M., Zeniya, M., et al. (2012). Genome-wide association study identifies
TNFSF15 and POU2AF1 as susceptibility loci for primary biliary cirrhosis in the Japanese population.
Am J Hum Genet *91*, 721-728.
Nielsen, P.J., Georgiev, O., Lorentz, B., and Schaffner, W. (1996). B lymphocytes are impaired in
mice lacking the transcriptional co-activator Bob1/OCA-B/OBF1. Eur J Immunol *26*, 3214-3218.
Omilusik, K.D., Best, J.A., Yu, B., Goossens, S., Weidemann, A., Nguyen, J.V., Seuntjens, E.,
Stryjewska, A., Zweier, C., Roychoudhuri, R., et al. (2015). Transcriptional repressor ZEB2 promotes
terminal differentiation of CD8+ effector and memory T cell populations during infection. J Exp Med
212, 2027-2039. 10.1084/jem.20150194.
Oxenius, A., Bachmann, M.F., Zinkernagel, R.M., and Hengartner, H. (1998). Virus-specific MHC-
class II-restricted TCR-transgenic mice: effects on humoral and cellular immune responses after viral
infection. Eur J Immunol *28*, 390-400.
Pace, L., Goudot, C., Zueva, E., Gueguen, P., Burgdorf, N., Waterfall, J.J., Quivy, J.P., Almouzni, G.,
and Amigorena, S. (2018). The epigenetic control of stemness in CD8(+) T cell fate commitment.
Science *359*, 177-186. 10.1126/science.aah6499.
Pearce, E.L., Walsh, M.C., Cejas, P.J., Harms, G.M., Shen, H., Wang, L.S., Jones, R.G., and Choi, Y.
(2009). Enhancing CD8 T-cell memory by modulating fatty acid metabolism. Nature *460*, 103-107.
nature08097 [pii] 10.1038/nature08097 [doi].
Pepper, M., Pagan, A.J., Igyarto, B.Z., Taylor, J.J., and Jenkins, M.K. (2011). Opposing signals from
the Bcl6 transcription factor and the interleukin-2 receptor generate T helper 1 central and effector
memory cells. Immunity *35*, 583-595.
Perovanovic, J., Wu, Y., Abewe, H., Shen, Z., Hughes, E., Gertz, J., Chandrasekharan, M.B., and
Tantin, D. Oct1 recruits the histone lysine demethylase Utx to canalize lineage specification.
submitted https://www.biorxiv.org/content/10.1101/2020.12.01.406488v2.
Qin, X.F., Reichlin, A., Luo, Y., Roeder, R.G., and Nussenzweig, M.C. (1998). OCA-B integrates B cell
antigen receptor-, CD40L- and IL 4-mediated signals for the germinal center pathway of B cell
development. EMBO J *17*, 5066-5075.
Rao, R.R., Li, Q., Gubbels Bupp, M.R., and Shrikant, P.A. (2012). Transcription factor Foxo1
represses T-bet-mediated effector functions and promotes memory CD8(+) T cell differentiation.
Immunity *36*, 374-387. S1074-7613(12)00088-X [pii] 10.1016/j.immuni.2012.01.015 [doi].
Raphael, I., Joern, R.R., and Forsthuber, T.G. (2020). Memory CD4(+) T Cells in Immunity and
Autoimmune Diseases. Cells *9*, 531. 10.3390/cells9030531.
Rodriguez-Jimenez, P., Fernandez-Messina, L., Ovejero-Benito, M.C., Chicharro, P., Vera-Tome, P.,
Vara, A., Cibrian, D., Martinez-Fleta, P., Jimenez-Fernandez, M., Sanchez-Garcia, I., et al. (2021).
Growth arrest and DNA damage-inducible proteins (GADD45) in psoriasis. Sci Rep *11*, 14579.
10.1038/s41598-021-93780-x.
Rosato, P.C., Wijeyesinghe, S., Stolley, J.M., Nelson, C.E., Davis, R.L., Manlove, L.S., Pennell, C.A.,
Blazar, B.R., Chen, C.C., Geller, M.A., et al. (2019). Virus-specific memory T cells populate tumors
and can be repurposed for tumor immunotherapy. Nat Commun *10*, 567. 10.1038/s41467-019-08534-
1.
Rutishauser, R.L., Martins, G.A., Kalachikov, S., Chandele, A., Parish, I.A., Meffre, E., Jacob, J.,
Calame, K., and Kaech, S.M. (2009). Transcriptional repressor Blimp-1 promotes CD8(+) T cell
terminal differentiation and represses the acquisition of central memory T cell properties. Immunity *31*,
296-308.

Sallusto, F., Lanzavecchia, A., Araki, K., and Ahmed, R. (2010). From vaccines to memory and back.
Immunity *33*, 451-463. 10.1016/j.immuni.2010.10.008.
Samardzic, T., Marinkovic, D., Nielsen, P.J., Nitschke, L., and Wirth, T. (2002). BOB.1/OBF.1
deficiency affects marginal-zone B-cell compartment. Mol Cell Biol *22*, 8320-8331.
10.1128/MCB.22.23.8320-8331.2002.
Sauter, P., and Matthias, P. (1997). The B cell-specific coactivator OBF-1 (OCA-B, Bob-1) is inducible
in T cells and its expression is dispensable for IL-2 gene induction. Immunobiology *198*, 207-216.
S0171-2985(97)80041-1 [pii] 10.1016/S0171-2985(97)80041-1 [doi].
Schubart, D.B., Rolink, A., Kosco-Vilbois, R.G., Botteri, F., and Matthias, P. (1996). B-cell-specific
coactivator OBF-1/OCA-B/Bob1 required for immune response and germinal centre formation. Nature
383, 538-542.
Shakya, A., Goren, A., Shalek, A., German, C.N., Snook, J., Kuchroo, V.K., Yosef, N., Chan, R.C.,
Regev, A., Williams, M.A., and Tantin, D. (2015). Oct1 and OCA-B are selectively required for CD4
memory T cell function. J Exp Med *212*, 2115-2131. 10.1084/jem.20150363.
Shakya, A., Kang, J., Chumley, J., Williams, M.A., and Tantin, D. (2011). Oct1 is a switchable,
bipotential stabilizer of repressed and inducible transcriptional states. J Biol Chem *286*, 450-459.
10.1074/jbc.M110.174045.
Shen, H., Slifka, M.K., Matloubian, M., Jensen, E.R., Ahmed, R., and Miller, J.F. (1995). Recombinant
Listeria monocytogenes as a live vaccine vehicle for the induction of protective anti-viral cell-mediated
immunity. Proc Natl Acad Sci U S A *92*, 3987-3991. 10.1073/pnas.92.9.3987.
Shin, H.M., Kapoor, V., Guan, T., Kaech, S.M., Welsh, R.M., and Berg, L.J. (2013). Epigenetic
modifications induced by Blimp-1 Regulate CD8(+) T cell memory progression during acute virus
infection. Immunity *39*, 661-675. 10.1016/j.immuni.2013.08.032.
Snook, J.P., Kim, C., and Williams, M.A. (2018). TCR signal strength controls the differentiation of
CD4(+) effector and memory T cells. Sci Immunol *3*, eaas9103. 10.1126/sciimmunol.aas9103.
Strubin, M., Newell, J.W., and Matthias, P. (1995). OBF-1, a novel B cell-specific coactivator that
stimulates immunoglobulin promoter activity through association with octamer-binding proteins. Cell
80, 497-506.
Stuart, T., Butler, A., Hoffman, P., Hafemeister, C., Papalexi, E., Mauck, W.M., 3rd, Hao, Y.,
Stoeckius, M., Smibert, P., and Satija, R. (2019). Comprehensive Integration of Single-Cell Data. Cell
177, 1888-1902 e1821. 10.1016/j.cell.2019.05.031.
Utzschneider, D.T., Charmoy, M., Chennupati, V., Pousse, L., Ferreira, D.P., Calderon-Copete, S.,
Danilo, M., Alfei, F., Hofmann, M., Wieland, D., et al. (2016). T Cell Factor 1-Expressing Memory-like
CD8(+) T Cells Sustain the Immune Response to Chronic Viral Infections. Immunity *45*, 415-427.
10.1016/j.immuni.2016.07.021.
Williams, M.A., Ravkov, E.V., and Bevan, M.J. (2008). Rapid culling of the CD4+ T cell repertoire in
the transition from effector to memory. Immunity *28*, 533-545.
Yeo, L., Woodwyk, A., Sood, S., Lorenc, A., Eichmann, M., Pujol-Autonell, I., Melchiotti, R., Skowera,
A., Fidanis, E., Dolton, G.M., et al. (2018). Autoreactive T effector memory differentiation mirrors beta
cell function in type 1 diabetes. J Clin Invest *128*, 3460-3474.
Zwilling, S., Dieckmann, A., Pfisterer, P., Angel, P., and Wirth, T. (1997). Inducible expression and
phosphorylation of coactivator BOB.1/OBF.1 in T cells. Science *277*, 221-225.

A I-A^b/gp$_{66-77}$ Tetramer$^+$

n.s.   *p*=0.07

% CD4$^+$CD44$^+$ cells

D8   D40+7

● *Ocab*$^{fl/fl}$
● *Ocab*$^{fl/fl}$;CD4-Cre

B *Ocab*$^{fl/fl}$ *Ocab*$^{fl/fl}$;CD4-Cre

D8   4.97   1.03   5.62   1.69

D40+7   23.3   5.04   3.87   4.98

IFNγ

IL-2

C IFNγ$^+$IL-2$^-$

*   n.s.

p=0.06

% CD4$^+$ T cells

● D8
● D40+7

Ocab$^{fl/fl}$ *Ocab*$^{fl/fl}$;CD4-Cre

D 1:1   BM cells

Thy1.1 WT+Thy1.2 *Ocab*$^{fl/fl}$ or
Thy1.1 WT+Thy1.2 *Ocab*$^{fl/fl}$;CD4-Cre

Irradiated C57BL/6J
Ly5.1 recipients

reconstitute   LCMV^{Arm} Lm-gp61

8 weeks

D8   D42   D42+7

E I-A^b/gp$_{66-77}$ Tetramer$^+$

D8   D42+7

Ocab$^{fl/fl}$ *f/f*   WT   *f/f*   WT

48.2   51.8   48.4   51.6

Ocab$^{fl/fl}$;CD4-Cre   *f/f*;Cre   WT   *f/f*;Cre   WT

37.9   62.1   10.5   89.3

Ly5.2

Thy1.1

F **

Ratio WT/control or WT/OCA-B deficient

D8   D42+7

● *Ocab*$^{fl/fl}$
● *Ocab*$^{fl/fl}$;CD4-Cre

Sun, Kim et al. Figure 2

A

SMARTA LCMV D8

EV   OCA-B

Foxp3
Sell

Ccr1
Zbtb16
Bcl6

Ifngr2

Il7r

Pou2af1 (Ocab)
Smad3
Tbx21 (Tbet)
Notch3
Zeb2
Ikzf2 (Helios)
Socs2
Gadd45b

B

EV [0-80]   20 kb
OCA-B [0-80]
Pou2af1

EV [0-3.6]   5 kb
OCA-B [0-3.6]
Socs2

EV [0-5.4]   5 kb
OCA-B [0-5.4]
Gadd45b

C

OCA-B upregulated genes

Epigenomic roadmap HM	*p*-value
H3K9me3 Breast vHMEC	0.048
Molecular function	***p*-value**
Toll-like receptor binding	0.006
Aspartic-type peptidase activity	0.007

OCA-B downregulated genes

Epigenomic roadmap HM	*p*-value
H3K27me3 CD4 Memory Primary Cells	1.45E-04
H3K27me3 CD8 Memory Primary Cells	9.30E-04
Molecular function	***p*-value**
Superoxide-generating NADPH oxidase activator activity	1.29E-09
Cytokine receptor activity	1.24E-06

D

OCA-B upregulated genes

ChEA 2016	*p*-value
Tp53	0.01936
Smc3	0.02544

OCA-B downregulated genes

ChEA 2016	*p*-value
IRF8	1.24E-20
SMRT	1.14E-15
MYB	1.73E-09
Nerf2	4.79E-09
NCOR	1.52E-08

Sun, Kim et al. Figure 5