Brain mapping and spatial protein profiling reveal functional connectivity deficits and

molecular changes following repetitive mild traumatic brain injury in wild-type mice.

Sakthivel Ravi[1,2,3], Marangelie Criado-Marrero[1,2,3], Daylin Barroso[1,2,3], Isadora M Braga[3],

Mackenzie Bolen[2,3], Uriel Rubinovich[1,3], Gabriela P. Hery[1,5], Matteo M Grudny[4], John Koren

III[1,2,3], Stefan Prokop[1,3,5,6], Marcelo Febo[3,4], Jose Francisco Abisambra[1,2,3,7, *]

[1]Center for Translational Research in Neurodegenerative Disease, [2]Department of Neuroscience,

[3]McKnight Brain Institute, [4]Department of Psychiatry, [5]Department of Pathology, [6]Fixel Institute

for Neurological Diseases, [7]Brain Injury Rehabilitation and Neuroresilience (BRAIN) Center

University of Florida, Gainesville, FL 32610, USA

16 *Correspondence: j.abisambra@ufl.edu

Abstract

Background: Repetitive mild traumatic brain injury (rmTBI) is a leading and severe threat to cognition that often goes undiagnosed. A major challenge in developing diagnostics and treatments for the consequences of rmTBI is the fundamental knowledge gaps that explain how TBI promotes brain dysfunction. It is both critical and urgent to understand the neuropathological and functional consequences of rmTBI to develop effective therapeutic strategies. In this study, we defined the extent of altered brain functional connectivity (FC) and expression of neuropathological markers after rmTBI.

Methods: We performed two rmTBI (2x 0.6 J impacts 24 h apart) in male and female C57BL/6J wild-type (WT) (~2.5-3mo) mice using CHIMERA or sham procedures. At 5-6 days post-injury (dpi), we measured changes in brain volume and FC using T2-weighted images, resting-state functional MRI (rsfMRI) and graph theory analyses. We used diffusion tensor imaging (DTI) to assess microstructural changes in white matter tracts. In addition, at 7dpi, we measured changes in Iba1 and GFAP in specific brain regions to determine the extent of gliosis. The expression of disease-associated protein markers in grey and white matter (WM) regions were evaluated using the NanoString-GeoMx digital spatial protein profiling spatial profiling (DSP) platform.

Results: The rsfMRI data revealed aberrant changes in connectivity such as node clustering coefficient, global and local efficiency, participation coefficient, eigenvector centrality, and betweenness centrality in thalamus and other key brain regions that process visual, auditory, and somatosensory information. Using DTI, we found that fractional anisotropy (FA) and axial diffusivity in the optic tract was significantly decreased. Also, mean, radial, and axial diffusivity (L1) were significantly increased in the hippocampus. DSP revealed that phospho-serine 199 tau (pS199) as well as glial markers such as GFAP, cathepsin-D, and Iba1 were significantly increased in the optic tract. In thalamic nuclei, the neuroinflammatory marker GPNMB was increased significantly, and the cell proliferation marker Ki-67 was significantly decreased in the rmTBI group. Our data suggest that rmTBI significantly alters brain functional connectivity and causes a profound inflammatory response in gray matter regions, beyond chronic white matter damage.

Keywords: CHIMERA, repetitive mild TBI, diffusion tensor imaging, microglia, resting state fMRI, optic tract, thalamus.

Introduction

Repetitive mild traumatic brain injuries (rmTBI) are the most frequently diagnosed form of head injury in the United States [1–4]. These repetitive brain injuries result from multiple and subsequent impacting blows to the head, and they are common in subjects engaging in contact sports as well as by active military personnel. Moreover, evidence suggests that TBIs, and in particular rmTBIs, significantly increase risk for numerous neurodegenerative disorders, including chronic traumatic encephalopathy (CTE) and Alzheimer's disease (AD), among other AD-related disorders (ADRD) [5–9]

The mechanisms linking rmTBIs and the pathogenesis of neurodegenerative disorders remain unknown. However, advances in monitoring both the anatomical and molecular changes that occur after rmTBI suggest links between TBI-induced brain alterations are detectable. For example, diffuse tensor imaging (DTI) in MRI identifies shears in white matter tracts that result in diffuse axonal injury (DAI), which is a typical sign of rmTBI [10]. In addition, increased expression of the microglial marker, Iba1 (ionized calcium-binding adapter molecule 1) and astrocytic marker, GFAP (glial fibrillary acidic protein) reflect gliosis response after injury. Though these network and molecular changes reveal considerable mechanistic information and can be found in TBIs of different severity, distinct markers of rmTBI are not clearly established.

In this study, we used functional brain imaging and digital spatial protein profiling to identify markers of altered neural function following rmTBI. Given that rmTBI causes rotational damage that ruptures white matter axonal tracts, we examined the effects of rmTBI on regions that possess high and low amount of white matter (optic tract and commissures and thalamic

nuclei, respectively). Additionally, due to the purported link between TBI and cognitive damage, we examined brain structures highly involved in ADRDs (hippocampus and cortex). We identified region-specific abnormalities resulting from rmTBI. These distinct imaging and protein surrogates of brain damage strongly suggest that, though the global consequences of rmTBIs are unpredictable, the local region-specific effects are consistent and are reflected at the gross structural and fine molecular levels. This information reveals novel imaging and protein biomarkers that could help determine outcomes and inform mechanistic knowledge gaps.

Materials and Methods

Animals

Male and female C57BL/6J (WT) mice were purchased from the Jackson Laboratory (Bar Harbor, ME, USA). Two to five mice of same-sex groups were housed in standard room conditions with 12 h light and dark cycle and allowed to access free food and water ad libitum. All the animal procedures were approved by the Institutional Animal Care and Use Committee (IACUC), University of Florida.

Repetitive mild traumatic brain injury (rmTBI) by CHIMERA

The rmTBI procedure was followed as suggested by Namjoshi et al. 2014[11]. 2.5-3 months old male and female WT mice were subjected to two mild closed head injuries (0.6 J) at 24h intervals using the Closed-Head Impact Model of Engineered Rotational Acceleration (CHIMERA) impactor. Briefly, mice were anesthetized with isoflurane (induction 3-4% and maintenance 1-2.5%), and lubricating eye ointment was applied to prevent corneal drying. Meloxicam (10 mg/kg) was administered subcutaneously right before injury for two days to alleviate pain. Animals were positioned supine in the holding bed of the impactor. The head was aligned in a flat position over a hole in the head plate to receive piston strikes closer to the

bregma. After impact, animals were transferred immediately to the recovery chamber pre-

warmed at 38°C and monitored until fully ambulatory. Sham mice were exposed to all these

procedures, except for the impact. At 7dpi, transcardial perfusions were performed with sterile

ice-cold 0.9 % saline and the brain tissues were harvested. One half of the hemibrain was micro-

dissected and stored at -80°C ultra-deep freezer for further analysis. The other half of the

hemibrain was fixed in 10% neutral buffered formalin for further histological analysis.

T2 and resting-state functional Magnetic Resonance Imaging (rsfMRI) data acquisition

Anatomical T2-weighted, diffusion weighted and rsfMRI scans were collected between

5-6 dpi. Sham and rmTBI mice were anesthetized with a mixture of isoflurane (Induction 4%,

maintenance: 1%) and medical grade air (0.1-0.15L/min) and positioned prone on a custom-built

holder with bed, ports for warm water flow, a bite bar and a nose port for gas anesthetic delivery.

A respiratory pad was placed underneath the abdomen to monitor respiration throughout the

entire imaging session. The core body temperature of the mice was maintained at 36-37°C by

placing the circulating warm water setup connected to the water bath. Lubricating eye ointment

was applied to prevent corneal drying, and a radiofrequency (RF) coil was placed on top of the

head. The images were acquired using an 11.1T MRI scanner (Magnex Scientific Ltd, Oxford,

UK) with high power gradient sets (RRI BFG-240/120-S7; maximum gradient strength of 1000

mT/m at 325 Amps and a 200 ms rise time; RRI, Billerica, MA). The MRI system was controlled

by a Bruker Paravision 6.01 console (Bruker BioSpin, Billerica, MA). A custom-made 2.0 cm x

2.5 cm quadrature RF surface transmit/receive coil tuned to 470.7 MHz (1H resonance) was used

for B1 excitation and signal detection (RF engineering lab, Advanced Magnetic Resonance

Imaging and Spectroscopy Facility, Gainesville, FL). A T2- weighted Turbo Rapid Acquisition

with Refocused Echoes (TurboRARE) sequence was acquired with the following parameters:

effective echo time (TE) = 41.42 ms, repetition time (TR) = 4s, RARE factor =16, number of

averages = 12, the field of view of 15 mm x 15 mm and 0.9 mm thick slice, a data matrix of 256

x 256 and 14 interleaved ascending coronal (axial) slices covering the entire brain from the

rostral-most extent of the anterior frontal cortical surface, caudally towards the upper brainstem

and cerebellum. The resting-state functional images were collected using a single-shot spin-echo

echo planar imaging sequence with the following parameters: TE = 15 ms, TR = 2 s, 600

repetitions, the field of view = 15 x 15 mm and 0.9 mm thick slice, and a data matrix of 64 x 48

with 14 interleaved ascending coronal slices in the same space as the T2 anatomical. Two

additional single repetition SE-EPI scans were collected, one with phase encode gradient lobes

('PE blips') collected along the positive gradient direction and the other collected with PE blips

reversed to the negative PE direction (Bruker Paravision binary method for distortion correction

is kindly provided by Dr. Matthew Budde, Medical College of Wisconsin).

rsfMRI image pre-processing

Functional connectivity of rmTBI mice brain was analyzed by graph theoretical analysis, an

approach highly used to assess novel biomarkers of brain function in neurological diseases and

aging [12]. SE-EPI distortions were corrected using TOPUP in FSL [13]. The pair of opposite

phase-encode blip scans were used to estimate the susceptibility-induced off-resonance field,

which is used to unwarp subsequent EPI volumes [13]. Image pre-processing and analysis was

carried via in-house UNIX terminal integrating the following tools: Analysis of Functional

NeuroImages (AFNI)[14], FMRIB Software Library (FSL)[15–17] and Advanced Normalization

Tools (ANTs) [18,19]. Anatomical and distortion-corrected functional scan masks outlining

mouse brain boundaries were generated in MATLAB using Three-Dimensional Pulsed Coupled

Neural Networks (PCNN3D) [20] and then manually edited. Then, 3dDespike in AFNI was used

to remove time series spikes and 3dvolreg for image volume alignment. Preprocessed scans were

cropped and a high-pass temporal filter (<0.009Hz) was used (3dTproject) to remove slow

variations (temporal drift) in the fMRI signal. Independent component analysis (ICA)

decomposition was then applied using Multivariate Exploratory Optimized Decomposition into

Independent Components (FSL MELODIC version 3.0) to preprocessed scans to assess noise

components in each subjects' native space prior to spatial smoothing and registration. The

resulting components were assessed, and in most cases all components contained noise-related

signal along brain edges, in ventricular voxels, and large vessel regions. These components were

suppressed using a soft ('non-aggressive') regression approach, as implemented in FMRIB

Software Library (FSL 6.0.3) using fsl_regfilt [15]. A low-pass filter (>0.12Hz) and spatial

smoothing (0.4mm FWHM) was then applied to the fMRI scans.

Preprocessed anatomical and fMRI scans were aligned to a parcellated mouse common

coordinate framework (version 3, or CCFv3) template [21]. Bilateral region of interest (ROI)-

based nodes (64 total) was created with the guidance of the annotated CCFv3 parcellation and

using tools in ITKSNAP and FSL. In ITKSNAP, the template along with the overlaid

parcellation were used to find the left hemisphere voxel coordinates for each of the nodes

included in this study, which were distributed as evenly as possibly without overlap across the

mouse brain template. The node coordinates were positioned in subregions of all areas for

example the optic tract and thalamus. We created 0.6 mm diameter spheric nodes in the template

space (resolution: $0.05mm^3$) centered on the voxel coordinates.   The right hemispheric

representations of the same nodes were then created to complete left and right representations for

each node.   For subject-to-atlas registration, fMRI scans are upsampled from a native space

0.234x0.3125 x0.9 mm resolution (spatially smoothed at 0.4 mm FWHM) to a downsampled

template resolution of $0.1mm^3$. Thus, the spheric nodes, in effect, were designed to sample fMRI

signals at a native space slightly over single-voxel resolution.

Anatomical scans were cropped and N4 bias field correction [22] applied to T2 images to

correct intensity variations due to RF field inhomogeneities [23]. The extracted brain maps were

linearly registered to the mouse template using FSL linear registration tool (FLIRT)[15], using a

correlation ratio search cost, full 180-degree search terms, 12 degrees of freedom and trilinear

interpolation. The linear registration output was then nonlinearly warped to the template space

using ANTs (antsIntroduction.sh script). Anatomical-to-atlas linear and nonlinear transformation

matrices were applied to fMRI scans at a later stage. Brain extraction using a mask (see above)

was first applied to fMRI scans and the cropped scans were then aligned to their respective

higher resolution anatomical scans. Timeseries functional images were split into 600 individual

volumes and the first in the series was linearly aligned to the anatomical scan using FLIRT (same

parameters as above, except 6 degrees of freedom was used in this step). ANTs

(antsRegistrationSyNQuick.sh script) was used to warp the lower resolution functional images to

their structural (using a single stage step deformable b-spline syn with a 26-step b-spline

distance). Linear and nonlinear warping matrices for fMRI-to-anatomical alignment were applied

to individual scans in the time series, then the merged 4-D functional timeseries were moved to

the atlas space using the prior anatomical-to-template transformation matrices.

A total of 64 ROI masks, divided into 32 left and 32 right ROI's, were included in our

analyses. Center voxel coordinates (see above) were used for 3D network visualizations in

BrainNet viewer in MATLAB [24]. Signal timeseries were extracted from preprocessed fMRI

scans with the assistance of ROI mask overlays. This generated 64 individual ROI text files per

subject that contained L2-normalized resting state signals as a vector of 600 data points. The

timeseries files were used in cross-correlations and in calculations of Pearson r coefficients for

every pairwise combinations of ROIs (1dCorrelate in AFNI). The resulting number of pairwise

correlations was 1,952 per subject (after removing 64 self-correlations). Correlation coefficients

were imported to MATLAB and Fisher's transform applied to ensure a normal distribution of z

values prior to analyses.

Functional Network Analysis

Functional network analysis was completed as previously reported [25]. Briefly, the

Brain Connectivity Toolbox [26] and MATLAB were used to determine weighted matrices.

Edge densities thresholds were set in a range from 2 to 40% to calculate the following global

network metrics: Clustering Coefficient (tendency of nodes to cluster and connect within the

network), Characteristic Path Length (average of shortest path length between all pairs of nodes

in the network), Transitivity (probability of neighboring nodes to be interconnected within the

network), Global Efficiency (efficiency to communicate across distant brain regions), Louvain

Modularity (density of connections within a neural cluster), and Small World Index (high local

clustering with shortest path length). Unless otherwise indicated in the figures, a 10% threshold

was used for all the node-specific measures such as Node Strength (how strongly a node directly

connects to other nodes), Node degree (number of connections with other nodes), Eigenvector

Centrality (node's influence in a network), Participation Coefficient (node interaction within

single or multiple communities), Local Efficiency (capacity to integrate information between

neighboring nodes), and Betweenness Centrality (node's influences over the flow of information

in the network). Significant alterations in brain connectivity between sham and rmTBI were

determined by using an unpaired T-test.

Brain volume and diffusion tensor imaging (DTI): Image Processing and Analysis

Anatomical T2-weighted images of sham and rmTBI mice were used to measure the

brain volume using ITK-SNAP software. Diffusion weighted scans were acquired at 5-7 dpi

using a 4-shot, 2-shell spin echo planar diffusion imaging (DTI EPI) sequence in Bruker

Paravision, with TR = 4 seconds, TE = 19 ms, number of averages = 4, gradient duration δ = 3

ms, diffusion time Δ = 8 ms, 54 images with 3 different diffusion weightings, two b=0, 6

directions with b=600 s/mm^2, and 46 directions with b=2000 s/mm^2. A navigator signal was used

by the Bruker reconstruction software to improve signal stability in the 4-shot EPI. Image

saturation bands were placed on either side and below the brain to suppress non-brain signal

during image acquisition. Diffusion images had the same FOV and slice thickness as the

anatomical scan but with a lower resolution data matrix size of 128 x 96 and 17 slices

(resolution: 0.117 mm x 0.117 mm x 0.7 mm) in the same space as anatomical scans. This

allowed careful manual outlining of regions to be analyzed by using T2 scans and diffusion maps

(see below).

Diffusion MRI scans were processed using tools available on FMRIB software library -

FSL [27] and DSI Studio [28] as previously reported for mouse diffusion scans obtained at

11.1Tesla [29]. Individual diffusion images were examined carefully and removed from the

image series due to low signal to noise or the presence of motion artifact. The gradient b-vector

files were corrected by removing gradient directions for the removed images. Eddy correction

was used for adjusting slight movements during image acquisition and gradient files rotated

according to the motion correction vectors. After eddy correction, tensor element reconstruction

and estimates of 1st, 2nd and 3rd eigenvectors and eigenvalues (λ1, λ2, λ3, where λ1 is axial

diffusivity and the average of λ2 and λ3 provides radial diffusivity values) was performed using

weighted least squares regression on DTIFIT in FSL [30]. This last step generated independent

images of FA, mean, axial and radial diffusivities (MD, AD, and RD, respectively).

Regions of interest were manually selected in ITK-SNAP to estimate their mean intensity

and volumes. These included thalamus, optic tract, fimbria, corpus callosum, hippocampus, and

amygdala. Significant alterations in brain microstructure were determined by using an unpaired

T-test.

Immunohistochemistry (IHC)

Formalin-fixed, Paraffin-embedded coronal brain sections (5 µm) were used for the IHC

study. Both sham and rmTBI slides were processed simultaneously. The slides were

deparaffinized in xylene twice for 5 min each and rehydrated sequentially in the gradient of

ethanol followed by water for 3 min each. The antigen retrieval was performed by immersing

slides in 10mM citrate with 0.5% tween20 (pH 6.0) and 30 min incubation in the steamer. The

endogenous peroxidases were removed by incubating the slides with the mixture of 0.3%

H_2O_2/PBS and 10% Triton X-100 (100µl for 200 ml of 0.3% H_2O_2 solution) for 20 min. Blocking

was performed with 10% normal goat serum/PBS-T (0.05% tween) for 30 min at room

temperature (RT). After blocking, the brain sections were incubated overnight with the following

antibodies: Iba1 (1:1000, PA5-27436, Invitrogen) and GFAP (GA5) (1:1000, CS3670S, Cell

Signaling) at 4°C for microglia and astrocytes, respectively. After incubation, the slides were

washed with PBS-T and incubated for 30 min with corresponding biotinylated secondary

antibody [Goat Anti-Rabbit IgG Antibody (H+L), Biotinylated-(BA-1000); Goat Anti-Mouse

IgG Antibody, Biotinylated, R.T.U. BP-9200, Vector laboratories). Further, the sections were

incubated with avidin-biotin complex (ABC) reagent for 20 min (VECTASTAIN Elite ABC-

HRP Kit, Peroxidase Standard), PK-6100) following washes with PBS-T and developing with 3,

3'-diaminobenzidine (DAB) kit [KPL DAB Reagent Set, SeraCare (5510-0031)]. Then, the

slides were washed twice in PBS, 5 min each, and counter-stained with hematoxylin for 1-2 min.

The slides were washed briefly in water and dehydrated with the gradient of alcohol (70%

ethanol, 90% ethanol and 100% ethanol) for 3 min each and incubated in xylene twice for 5 min

each. The sections were mounted with Cytoseal Mountant 60 (Epredia 83104) and air dried. The

slides were scanned using an Aperio image scanner (20X objective lens, Scan Scope™ XT,

Aperio Technologies, Inc. Vista, CA, USA) and quantified using Image Scope software

(v12.4.3.5008) positive pixel count program. Three sections per mouse at 55 µm apart were

averaged to calculate the expression of Iba1 and GFAP.

Spatial proteomics analysis using the NanoString GeoMx Digital Spatial protein profiling

platform

Formalin-fixed paraffin-embedded brain tissue was sectioned coronal at 5 µm thickness.

DSP staining and assay procedures were conducted according to the manufacturer-directed

protocol. Briefly, to identify regions of interest (ROI), nuclei were fluorescently labeled with

SYTO 13 (GeoMx Nuclear Stain Morphology kit, Item no. 121300303, Nanostring, Seattle,

WA), and microglia cells were labeled with an Iba1 (GeoMx Alzheimer's Morphology Kit, Item

no. 121300306, NanoString, Seattle, WA, United States) antibody. The regions of interest (ROIs)

were selected based on visual identification of anatomic regions using the nuclear stain, aided by

increased Iba1 expression in white matter regions of injured animals. The following ROI were

selected: primary somatic sensory area, dentate gyrus polymorph layer, thalamus, and the optic

tract. The following protein cores and modules were used for multiplexed protein quantification:

Neural Cell Profiling (Item no. 121300120), Parkinson's disease (PD) pathology (Item no.

121300122), Alzheimer's Disease (AD) pathology (Item no. 121300121), AD pathology

extended (Item no. 121300123), Glial cell subtyping (Item no. 121300125), and autophagy (Item

no. 121300124) markers. Samples were processed on the NanoString GeoMx system and the

nCounter max system following manufacturer's instructions. Data were analyzed using GeoMx

DSP analysis suit Version 2.4.0.421 as suggested by manufacturer instructions. Briefly, the Field

of view (FOV) QC was adjusted to 45% to check the low count score. All the QC passed

segments were normalized to housekeepers such as GAPDH and Histone H3. Further

background corrections were performed with the negative background targets such as Rb IgG, Rt

IgG2a, and Rt IgG2b. background corrected files were changed to tab-delimited files, and

custom scripts with slight modifications were used to generate volcano plots.

Statistical analyses

GraphPad Prism 9 software was used to perform all the statistical analysis. An unpaired

T-test was used to determine significant differences in nodal properties between control and

injured mice. Multiple t-tests were performed to measure the difference between density

thresholds in rsfMRI data. An unpaired T-test was performed within the GeoMx DSP analysis

suit to analyze the NanoString GeoMx Spatial Profiling data. All the results were represented as

mean $\pm$ SD; $p<0.05$ is considered statistically significant.

Results

We used the Closed Head Injury Model of Engineered Rotational Acceleration

(CHIMERA) that reproducibly causes DAI [11,31,32]. We injured 2.5-3mo C57Bl/6 mice, twice

within a 24hr interval, and we measured outcomes after the primary injury phase (Fig. 1). As

expected from clinical correlations with T2-weighted images, we found no overt structural or

volumetric changes in the injured mice compared to controls (Fig. 2A-C).

Given that changes in brain network organization strongly associate with cognitive

dysfunction [33], we then coupled resting state functional MRI (rsfMRI) and graph theory

network analyses to identify spatial and functional information following TBI [34]. We

performed focal measurements in a rsfMRI modality such as node clustering coefficient, global

and local efficiency, participation coefficient, eigenvector centrality, and betweenness centrality

in specific brain regions that process visual, auditory, somatosensory information, fractional

anisotropy (FA) and axial diffusivity in the optic tract; all in the absence of differences in brain

volume. We developed graphs of functional brain networks by defining nodes (macroscopic

regions of the brain) and edges (links). We selected 64 nodes (Supplementary table 1) that are

closely associated with white matter tracts (Fig. 2D & E) for functional connectivity analysis.

We calculated the node clustering coefficient, which is the index of the number of connected

neighboring nodes [35]. We found that the clustering coefficient in specific thalamic and visual

nodes were the most pronounced and significantly different changes identified (Fig. 2F-2I).

We then measured global efficiency and found that the edge density at 4% and 6%

threshold was significantly decreased (p=0.024 and p=0.008, respectively) in the rmTBI group

(Fig. 3A). Focusing on these densities, we found that local efficiency at the 4% (Fig. 3B-3C) and

6% (Fig. 3D-3F) density thresholds were significantly decreased in specific thalamic and visual

nuclei. No changes were observed in the net clustering coefficient and small world index

between sham and rmTBI groups (Fig. 3G-3H). In contrast, the net characteristic pathlength,

revealing about functional network integration, was significantly increased (p=0.0367) in the

rmTBI group at 4% edge density (Fig. 3I), but no changes were observed at the edge density

ranges between 6-40%.

The greatest differences following rmTBI were detected in the strength of a node's connections within its module (participation coefficient) and the influence of distinct nodes on the network (eigenvector centrality). We found that the participation coefficient in the left ventral posteromedial thalamic nuclei was significantly decreased ($p=0.0026$) in the rmTBI group (Fig. 4A). Conversely, participation coefficient was significantly increased in the right splenium of corpus callosum ($p=0.0028$), right central lateral parafascicular thalamic region ($p=0.0137$), right visual anteromedial area ($p=0.0373$), and right visual postrhinal area ($p=0.0375$) of the rmTBI group (Fig. 4B-4E). Eigenvector centrality in specific brain regions such as genu of corpus callosum ($p=0.0166$), the body ($p=0.0329$) and splenium of corpus callosum ($p=0.0134$), and the thalamic paracentral nucleus ($p=0.0392$) were significantly increased in the rmTBI group compared to sham (Fig. 4F-4I). The node betweenness centrality in the thalamic dorsal lateral geniculate nucleus was significantly increased in the rmTBI group (Fig. 4J). These highly unique characteristics reveal alterations to unique brain regions that define rmTBI node specificity. Meanwhile, other network features such as global network strength, Louvain modularity, assortativity, and Gamma values showed modest or undetectable differences between the sham and rmTBI groups. Only, the net lambda values, which indicate the degree of network randomness within neighboring nodes, was significantly decreased at wide edge densities from 8% to 40% ($p<0.05$) in rmTBI group (Fig. S1).

Given FC differences in specific nodes, we examined cerebral microstructure by measuring the diffusion of water molecules in the thalamus, optic tract, and corpus callosum. We also focused on the fimbria, hippocampus, and amygdala because of their important role in learning and memory (Fig. 5A). We observed that the fractional anisotropy (FA), which measures white matter integrity, was significantly decreased ($p=0.0240$) in the optic tract of the

rmTBI mice (Fig. 5B). The mean diffusivity (MD) (p=0.0259) in the hippocampus was

significantly increased after rmTBI. A higher diffusion rate (increased water content) suggests

the presence of edema or inflammation in this region (Fig. 5C). Two other measures of axonal

integrity, radial diffusivity (RD; p=0.0303) and axial diffusivity (L1; p=0.0413) were also

increased in the hippocampus of the rmTBI group, suggesting greater diffusion rate running

parallel and perpendicular to white matter fiber tracts (Fig. 5D & 5E). As expected, due to the

limited white matter content, we did not detect volumetric changes in the thalamus, fimbria, and

amygdala between the sham and rmTBI groups.

We measured white matter gliosis by measuring localized changes in the levels of Iba1

and GFAP using immunohistochemistry. We detected a robust glial response in the optic tract

(Fig. 6A-B and 6D-E) and corpus callosum (Fig. 6G-H) of injured mice. As established in

previous reports, Iba1-positive microglia were significantly increased (p<0.0001) in the optic

tract of injured mice (Fig. 6C). Similarly, GFAP-positive signal was significantly increased in

both the optic tract (p<0.0001; Fig 6F) and corpus callosum (p<0.0320; Fig. 6I) of the rmTBI.

The rsfMRI and DTI data identified four brain regions that were significantly altered by

the injuries: the primary somatosensory area, hippocampus, thalamus, and optic tract. To

determine whether the injuries impacted the levels of specific proteins in these areas, we

performed digital spatial protein profiling using the NanoString GeoMx DSP platform. We

measured changes in the levels of 72 proteins altered in ADRDs int the following ROI's primary

somatosensory area, dentate gyrus polymorph area, thalamus, and optic tract.

Levels of distinct proteins were significantly increased or decreased in the three areas of

interest (Fig. 7A-B and Table 1). We observed no changes in the dentate gyrus polymorph layer

(Data not shown). Myelin basic protein was significantly downregulated in the primary

somatosensory area, which reflects white matter damage (Fig. 7N&O). Interestingly, both total

tau and phospho-tau S396 (pS396) levels were significantly decreased but levels of one phospho-

tau species, pS199 tau, were increased in the rmTBI group (Fig. 7C-7F). Various

neuroinflammatory markers such as Iba1, GFAP, Cathepsin-D (Fig. 7H-7J) were significantly

increased in the optic tract of the rmTBI group. The expression of neuroinflammatory marker

GPNMB was also significantly increased, and cell proliferation marker Ki-67 was significantly

decreased in the thalamic region (Fig. 7K-7M).

These data align with both expected and previously reported alterations to molecular

processes following head injury, including neuroinflammation (e.g. Iba1), white matter damage

(e.g. MBP), and microtubule disruption (e.g. tau). Unexpectedly, changes in the levels of these

proteins were unique to specific brain regions, suggesting that the injury-induced alterations

unleash distinct consequences. Importantly, these distinct changes could be the root of the

imaging alterations identified in the rsfMRI, and they may serve as biomarkers that distinguish

clinical outcomes.

Discussion

The long-term goal of this study was to establish novel mechanistic information that

connects head injuries to neurodegenerative disorders. Here, we identified rmTBI-induced

imaging abnormalities and aberrant accumulation of unique proteins. We determined that injuries

impact brain regions in distinct manners, and these distinctions may explain why rmTBI

manifests differently between subjects. We also identified proteins that are increased or

decreased depending on the brain region (Table 1). Our data strongly suggest that the molecular

pathways leading to accumulation or depletion of these proteins contributes to the imaging

abnormalities we identified.

Investigating brain networks using rsfMRI and graph theory offers powerful functional

measures, all of which can be translated to the clinic. Network alterations, such as the ones

described herein, have strong diagnostic potential for TBI among other brain disorders [36–38].

Recent work using rsfMRI identified broad functional connectivity deficits in human subjects;

these defects correlated with diminished choice-reaction scores, showing high sensitivity. Our

work improves upon pre-clinical and clinical studies because we dissected more granular aspects

of functional connectivity in a mouse model of rmTBI and at ultra-high field [25,39–41].

Functional graph metrics such as clustering coefficient, characteristic path length, centrality, and

efficiency are quantifiable features that reveal how information flows throughout the network

[42], and they were impaired following rmTBI (Fig. 2-4). Our findings are concordant with other

pre-clinical models of mild head injury showing local alterations in betweenness centrality,

clustering coefficient, and local efficiency, while global changes were intact [43], showing that

these robust measures could be applicable to milder and [44,45] repetitive TBI.

Although we did not detect global changes (as reported in Boroda et al., 2021[46]) in

clustering coefficient, the local clustering coefficient at 10% edge density was significantly

reduced in the visual anteromedial area and various sub-regions of the thalamus in the rmTBI

group. Further, we observed a significant decrease in net global and local efficiency in the

rmTBI group at thresholds below 10% densities which contain true connections; this measure is

commonly used in clinical practice [44,45]. These results positively reveal that local integration,

efficiency, and integrative information processing across distal brain regions were significantly

altered following rmTBI. Consistent with previous reports, we did not observe changes in the

small-world index and net clustering coefficient after mTBI [25].

We also calculated eigenvector and betweenness centrality to assess the highly influential

nodes and nodes at the intersection between the shortest path, respectively [25]. We observed

high betweenness centrality scores in thalamus dorsal lateral geniculate and high eigenvector

centrality scores in modules with nodes located in sub-regions of corpus callosum and thalamus.

Our results indicate that the injury increased the distance between connections of the thalamus

and corpus callosum thereby impacting the connectivity between clusters in these networks. We

note that these nodal changes are not observed in all TBI models. Yang et al., 2021[25] reported

no differences in betweenness centrality in Controlled cortical impact (CCI) rats at 30 dpi. Given

that nodes in eigenvector centrality scores in the contralateral cortex can be restored from 2 to 30

days after injury, it will be important to assess these metrics at later post-injury periods to

confirm the long-term repercussions of rmTBI. Confirming these changes at different timepoints

will provide valuable information about compensatory mechanisms and reorganization of

damaged networks after TBI.

The thalamus is an important deep gray matter region of the brain composed a bundle of

white matter tracts and groups of nuclei, which plays a vital role in global multifunctional

pathways. It is highly susceptible to damage due to sudden acceleration or deceleration

movements during head trauma [38]. Consistent with previous reports, our results show that

rmTBI induced by CHIMERA damages thalamic nuclei. Given its numerous roles in integrating

brain signaling, damage to the thalamus may result in varied manifestations after TBI; this may

underlie why patients present broad symptomatology.

In addition to the network-associated changes, aberrant signaling at the molecular level is

a probable pathogenic factor in the transition from rmTBI to chronic neurodegeneration. We

observed significant changes in distinct proteins that aligned with unique brain regions. For

example, significant accumulation of Iba1 and GFAP in the optic tract, which was consistent

with the immunohistochemistry results and previous studies using CHIMERA [11], indicate that

rmTBI induced gliosis. In addition, glycoprotein non-metastatic melanoma protein B (GPNMB)

was significantly increased in the thalamus of the rmTBI group. GPNMB is highly expressed in

glia, and is linked with risk for Parkinson's disease [47,48].

Given their limited and unclear molecular links with TBI, follow-up experiments

evaluating the prognostic value of GPNMB, Ki67, and cathepsin-D after rmTBI could unveil

new biomarkers. Interestingly, total and pS396 tau levels were decreased in the optic tract were

decreased, while pS199 tau was increased. We hypothesize that this could be the result of tau

cleavage resulting from head injury; importantly, it suggests that distinct post-translationally

modified tau species could extravasate and reflect the state of injuries thereby serving as more

reliable biomarkers [7]

The outcomes of the present study provide important information about the effect of

rmTBI and its primary molecular mechanism on the underlying pathology at 5-7dpi; this period

corresponds to secondary injury, where complex molecular pathways are unleashed after injury.

Importantly, changes in this time point could precede the long-term consequences of rmTBI and

connect injuries with the pathophysiology of TBI-associated neurodegenerative diseases. Our

future work on the assessment of the post-TBI effect together with behavioral outcomes will help

to understand the molecular mechanisms of TBI and their long-term consequences on cognition.

Conclusion

Using highly translational imaging modalities and digital spatial protein profiling we

report abnormalities following CHIMERA-induced rmTBI 5-7dpi. These measures may serve as

potential biomarkers that can diagnose rmTBI and predict clinical outcomes. Moreover, protein

changes could be responsible for the molecular pathways involved in the acute and chronic

consequences of rmTBI. Studies to identify the longevity of these abnormalities and their

association with cognitive dysfunction are currently underway.

Acknowledgements

This work was primarily supported by grant 1 R01 AG074584-01 from NIH/NIA. A

portion of this work was performed in the McKnight Brain Institute at the National High

Magnetic Field Laboratory's Advanced Magnetic Resonance Imaging and Spectroscopy

(AMRIS) Facility, which is supported by National Science Foundation Cooperative Agreement

DMR-1644779 and the State of Florida. This work was supported in part by an NIH award, S10

RR025671, for MRI/S instrumentation.

Conflict of interest:

The   authors   declare   no   conflict   of   interest.

References:

1. Cassidy JD, Carroll LJ, Peloso PM, Borg J, von Holst H, Holm L, et al. Incidence, risk factors and prevention of mild traumatic brain injury: Results of the WHO Collaborating Centre Task Force on Mild Traumatic Brain Injury. J Rehabil Med Suppl. 2004;28–60.

2. Schwab K, Terrio HP, Brenner LA, Pazdan RM, McMillan HP, MacDonald M, et al. Epidemiology and prognosis of mild traumatic brain injury in returning soldiers. Neurology. 2017;88:1571–9.

3. Nolan A, Hennessy E, Krukowski K, Guglielmetti C, Chaumeil MM, Sohal VS, et al. Repeated Mild Head Injury Leads to Wide-Ranging Deficits in Higher-Order Cognitive Functions Associated with the Prefrontal Cortex. J Neurotrauma. 2018;35:2425–34.

4. Pavlov V, Thompson-Leduc P, Zimmer L, Wen J, Shea J, Beyhaghi H, et al. Mild traumatic brain injury in the United States: demographics, brain imaging procedures, health-care utilization and costs. Brain Inj. Taylor & Francis; 2019;33:1151–7.

5. McKee AC, Cantu RC, Nowinski CJ, Hedley-Whyte ET, Gavett BE, Budson AE, et al. Chronic traumatic encephalopathy in athletes: Progressive tauopathy after repetitive head injury. J Neuropathol Exp Neurol. 2009;68:709–35.

6. Giunta B, Obregon D, Velisetty R, Sanberg PR, Borlongan C V., Tan J. The immunology of traumatic brain injury: a prime target for Alzheimer's disease prevention. J Neuroinflammation. ???; 2012;9:1.

7. Abisambra JF, Scheff S. Brain Injury in the Context of Tauopathies. J Alzheimer's Dis. 2014;40:495–518.

8. Jassam YN, Izzy S, Whalen M, McGavern DB, El Khoury J. Neuroimmunology of Traumatic Brain Injury: Time for a Paradigm Shift. Neuron. Elsevier Inc.; 2017;95:1246–65.

9. Walker A, Chapin B, Abisambra J, DeKosky ST. Association between single moderate to

severe traumatic brain injury and long-term tauopathy in humans and preclinical animal models:

a systematic narrative review of the literature. Acta Neuropathol Commun. BioMed Central;

2022;10:1–20.

10. Arfanakis K, Haughton VM, Carew JD, Rogers BP, Dempsey RJ, Meyerand ME. Diffusion

tensor MR imaging in diffuse axonal injury. AJNR Am J Neuroradiol. 2002;23:794–802.

11. Namjoshi DR, Cheng WH, McInnes KA, Martens KM, Carr M, Wilkinson A, et al. Merging

pathology with biomechanics using CHIMERA (Closed-Head Impact Model of Engineered

Rotational Acceleration): a novel, surgery-free model of traumatic brain injury. Mol

Neurodegener. 2014;9:55.

12. Pompilus M, Colon-Perez LM, Grudny MM, Febo M. Contextual experience modifies

functional connectome indices of topological strength and efficiency. Sci Rep. Nature Publishing

Group UK; 2020;10:1–15.

13. Andersson JLR, Skare S, Ashburner J. How to correct susceptibility distortions in spin-echo

echo-planar images: application to diffusion tensor imaging. Neuroimage. 2003;20:870–88.

14. Cox RW. AFNI: Software for Analysis and Visualization of Functional Magnetic Resonance

Neuroimages. Comput Biomed Res. 1996;29:162–73.

15. Jenkinson M, Bannister P, Brady M, Smith S. Improved Optimization for the Robust and

Accurate Linear Registration and Motion Correction of Brain Images. Neuroimage.

2002;17:825–41.

16. Smith SM. Fast robust automated brain extraction. Hum Brain Mapp. 2002;17:143–55.

17. Smith SM, Jenkinson M, Johansen-Berg H, Rueckert D, Nichols TE, Mackay CE, et al.

Tract-based spatial statistics: Voxelwise analysis of multi-subject diffusion data. Neuroimage.

2006;31:1487–505.

18. Avants BB, Tustison NJ, Song G, Cook PA, Klein A, Gee JC. A reproducible evaluation of

ANTs similarity metric performance in brain image registration. Neuroimage. 2011;54:2033–44.

19. Grandjean J, Zerbi V, Balsters JH, Wenderoth N, Rudin M. Structural Basis of Large-Scale

Functional Connectivity in the Mouse. J Neurosci. 2017;37:8092–101.

20. Chou N, Jiarong Wu, Bai Bingren J, Anqi Qiu, Kai-Hsiang Chuang. Robust Automatic

Rodent Brain Extraction Using 3-D Pulse-Coupled Neural Networks (PCNN). IEEE Trans Image

Process. 2011;20:2554–64.

21. Lein ES, Hawrylycz MJ, Ao N, Ayres M, Bensinger A, Bernard A, et al. Genome-wide atlas

of gene expression in the adult mouse brain. Nature. 2007;445:168–76.

22. Tustison NJ, Avants BB, Cook PA, Yuanjie Zheng, Egan A, Yushkevich PA, et al. N4ITK:

Improved N3 Bias Correction. IEEE Trans Med Imaging. 2010;29:1310–20.

23. Kenkel WM, Yee JR, Moore K, Madularu D, Kulkarni P, Gamber K, et al. Functional

magnetic resonance imaging in awake transgenic fragile X rats: evidence of dysregulation in

reward processing in the mesolimbic/habenular neural circuit. Transl Psychiatry. 2016;6:e763–

e763.

24. Xia M, Wang J, He Y. BrainNet Viewer: A Network Visualization Tool for Human Brain

Connectomics. Csermely P, editor. PLoS One. 2013;8:e68910.

25. Yang Z, Zhu T, Pompilus M, Fu Y, Zhu J, Arjona K, et al. Compensatory functional

connectome changes in a rat model of traumatic brain injury. Brain Commun. 2021;3:1–17.

26. Rubinov M, Sporns O. Complex network measures of brain connectivity: Uses and

interpretations. Neuroimage. Elsevier Inc.; 2010;52:1059–69.

27. Smith SM, Jenkinson M, Woolrich MW, Beckmann CF, Behrens TEJ, Johansen-Berg H, et

al. Advances in functional and structural MR image analysis and implementation as FSL.

Neuroimage. 2004;23:S208–19.

28. Yeh F-C, Verstynen TD, Wang Y, Fernández-Miranda JC, Tseng W-YI. Deterministic

Diffusion Fiber Tracking Improved by Quantitative Anisotropy. Zhan W, editor. PLoS One.

2013;8:e80713.

29. Colon-Perez LM, Ibanez KR, Suarez M, Torroella K, Acuna K, Ofori E, et al. Neurite

orientation dispersion and density imaging reveals white matter and hippocampal microstructure

changes produced by Interleukin-6 in the TgCRND8 mouse model of amyloidosis. Neuroimage.

2019;202:116138.

30. Behrens TEJ, Woolrich MW, Jenkinson M, Johansen-Berg H, Nunes RG, Clare S, et al.

Characterization and propagation of uncertainty in diffusion-weighted MR imaging. Magn Reson

Med. 2003;50:1077–88.

31. McNamara EH, Grillakis AA, Tucker LB, McCabe JT. The closed-head impact model of

engineered rotational acceleration (CHIMERA) as an application for traumatic brain injury pre-

clinical research: A status report. Exp Neurol. Elsevier; 2020;333:113409.

32. Cheng WH, Martens KM, Bashir A, Cheung H, Stukas S, Gibbs E, et al. CHIMERA

repetitive mild traumatic brain injury induces chronic behavioural and neuropathological

phenotypes in wild-type and APP/PS1 mice. Alzheimers Res Ther. 2019;11:6.

33. Pandit AS, Expert P, Lambiotte R, Bonnelle V, Leech R, Turkheimer FE, et al. Traumatic

brain injury impairs small-world topology. Neurology. 2013;80:1826–33.

34. Ham TE, Sharp DJ. How can investigation of network function inform rehabilitation after

traumatic brain injury? Curr Opin Neurol. 2012;25:662–9.

35. Yang Z, Zhu T, Pompilus M, Fu Y, Zhu J, Arjona K, et al. Compensatory functional

connectome changes in a rat model of traumatic brain injury. Brain Commun. 2021;3:1–17.

36. Khazaee A, Ebrahimzadeh A, Babajani-Feremi A. Identifying patients with Alzheimer's

disease using resting-state fMRI and graph theory. Clin Neurophysiol. International Federation

of Clinical Neurophysiology; 2015;126:2132–41.

37. Liu J, Li M, Pan Y, Lan W, Zheng R, Wu FX, et al. Complex Brain Network Analysis and Its

Applications to Brain Disorders: A Survey. Complexity. 2017;2017.

38. Tang L, Ge Y, Sodickson DK, Miles L, Zhou Y, Reaume J, et al. Thalamic resting-state

functional networks: Disruption in patients with mild traumatic brain injury. Radiology.

2011;260:831–40.

39. Sharp DJ, Beckmann CF, Greenwood R, Kinnunen KM, Bonnelle V, De Boissezon X, et al.

Default mode network functional and structural connectivity after traumatic brain injury. Brain.

2011;134:2233–47.

40. Hillary FG, Slocomb J, Hills EC, Fitzpatrick NM, Medaglia JD, Wang J, et al. Changes in

resting connectivity during recovery from severe traumatic brain injury. Int J Psychophysiol.

Elsevier B.V.; 2011;82:115–23.

41. Li F, Liu Y, Lu L, Shang S, Chen H, Haidari NA, et al. Rich-club reorganization of

functional brain networks in acute mild traumatic brain injury with cognitive impairment. Quant

Imaging Med Surg. 2022;12:3932–46.

42. Ajilore O, Lamar M, Leow A, Zhang A, Yang S, Kumar A, et al. Depression. Am J Geriatr

Psychiatry. 2015;22:195–206.

43. Kim E, Seo HG, Seong MY, Kang MG, Kim H, Lee MY, et al. An exploratory study on

functional connectivity after mild traumatic brain injury: Preserved global but altered local

organization. Brain Behav. 2022;1–13.

44. Hagmann P, Cammoun L, Gigandet X, Meuli R, Honey CJ, Wedeen VJ, et al. Mapping the

Structural Core of Human Cerebral Cortex. Friston KJ, editor. PLoS Biol. 2008;6:e159.

45. Perry A, Wen W, Lord A, Thalamuthu A, Roberts G, Mitchell PB, et al. The organisation of

the elderly connectome. Neuroimage. 2015;114:414–26.

46. Boroda E, Armstrong M, Gilmore CS, Gentz C, Fenske A, Fiecas M, et al. Network topology

changes in chronic mild traumatic brain injury (mTBI). NeuroImage Clin. Elsevier Inc.;

2021;31:102691.

47. Murthy MN, Blauwendraat C, Guelfi S, Hardy J, Lewis PA, Trabzuni D. Increased brain

expression of GPNMB is associated with genome wide significant risk for Parkinson's disease

on chromosome 7p15.3. Neurogenetics. neurogenetics; 2017;18:121–33.

48. Kia DA, Zhang D, Guelfi S, Manzoni C, Hubbard L, Reynolds RH, et al. Identification of

Candidate Parkinson Disease Genes by Integrating Genome-Wide Association Study,

Expression, and Epigenetic Data Sets. JAMA Neurol. 2021;78:464–72.

Figure legends:

Fig. 1. CHIMERA validation. The schematic representation and experimental design showing

the number of rmTBIs, rsfMRI and tissue collection. Brains were collected at 7 dpi for

Immunohistochemistry (IHC) and NanoString GeoMx spatial protein profiling analyses.

Fig. 2. Clustering coefficient analysis reveals that the network integration and efficiency in

specific brain regions are disrupted following rmTBI. Representative T2 weighted anatomical

brain images of sham and rmTBI mice **(A&B)** were used for brain volume analysis.

Quantification of brain volume revealed no changes between sham and rmTBI groups **(C)**. 3D

functional connectome maps of sham and rmTBI mice brain shows the position of nodes and

edges in the region of interest (ROIs) [total ROI = 64] **(D&E)**. Clustering coefficient in various

regions of thalamus and visual area **(F-I)** were significantly altered in rmTBI group compared to

sham. Unpaired t-test, mean ± SD, Sham (N= 8), rmTBI (N=10).

Fig. 3. Global and Local efficiency analysis reveals that efficiency in specific brain regions

are disrupted following rmTBI. Global efficiency **(A)** was significantly decreased at the

density threshold of 4% and 6% in rmTBI group compared to Sham. Local efficiency at 4%

threshold **(B & C)** was significantly decreased in thalamus anterior pretectal nucleus (L-Left)

and thalamus mediodorsal (R-Right) regions of Injured mice. Similarly, Local efficiency at 6%

threshold **(D-F)** was significantly decreased in thalamus anterior pretectal nucleus (L), thalamus

dorsal lateral geniculate (L) and visual anterior area (R) of injured mice compared to Sham. No

changes in global clustering coefficient **(G)** and small world index **(H)** were observed. The

global characteristic pathlength **(I)** at 4% threshold was significantly increased in injured mice

group compared to sham. The global metrices were analyzed by Multiple Unpaired t-test, mean ±

standard error at edge densities ranges 2 to 40%. The groups comparison was analyzed by

Unpaired t-test, mean ± SD, Sham (N= 8), rmTBI (N=10).

Fig. 4. rmTBI disrupted connectivity between modules, increased eigenvector centrality

and node betweenness centrality. Nodes in left (L) thalamus ventral posteromedial **(A)** region

had low participation coefficient, but it is significantly increased in the right (R) splenium of

corpus callosum, thalamus (R) and visual regions (R) of rmTBI group relative to control **(B-E).**

Increased eigenvector centrality shows increased high influence of specific nodes on connected

networks following rmTBI in thalamic regions and corpus callosum **(F-I).** The node

betweenness centrality shows the nodes at the intersection between shortest path in the specific

thalamus region **(J)** was significantly altered. Density threshold at which the participation

coefficient and node eigenvector centrality were calculated. Unpaired t-test, mean ± SD.

Fig. 5. rmTBI altered tissue microstructure in specific brain regions. Representative DTI brain image shows **(A)** ROIs selected for microstructure pattern analysis. rmTBI increased fractional anisotropy (FA) in the optic tract **(B)**. Mean diffusivity (MD) and radial diffusivity (RD) were significantly increased in hippocampus of the injured mice **(C&D)**. Axial diffusivity (L1) was significantly increased in the optic tract and hippocampus of the injured mice **(E)** compared to sham. Unpaired t-test, mean $\pm$ SD.

Fig. 6. rmTBI induced gliosis in white matter tracts. Iba1 expression was significantly increased in the optic tract of injured mice compared to sham **(A-C)**. The red arrowhead indicates microglia. Brain coronal view: 1 mm; Scale bar for inset (enlarged view of the optic tract) = 100 µm. ****$p<0.0001$, Unpaired t-test; Sham (N= 15), rmTBI (N=20). GFAP expression was significantly increased in the optic tract **(D-F)** and corpus callosum **(G-I)** of injured mice compared to sham. The blue arrowhead indicates astrocytes. Brain coronal view: 1 mm; Scale bar for inset (enlarged view of the optic tract and corpus callosum) = 100 µm. (****$p<0.0001$; *$p= 0.0320$, Unpaired t-test), Sham (N= 15), rmTBI (N=19).

Fig. 7. rmTBI increased the expression of disease associated proteins in white matter and grey matter regions. Coronal sections of sham and rmTBI mice shows the region of interest (primary somatic sensory area, dentate gyrus polymorph layer, thalamus, and the optic tract) selected for NanoString GeoMx spatial protein profiling analysis **(A&B)**. Volcano plots **(C)** and bar graphs **(D-J)** show significant alteration in the expression of Alzheimer's disease (AD) pathological markers, glial and neural cell markers in the optic tract of the injured mice. Volcano plot **(K)** and bar graphs show **(L&M)** significant changes in the expression of cell proliferation and AD associated, neuroinflammatory markers in the thalamus of the injured mice. Myelin basic protein was significant decreased in the primary somatosensory area following rmTBI

(N&O). Protein levels were normalized to the expression of housekeeping proteins. Unpaired t-test, mean ± SD, N=4-8.

Supplementary information Legend:

Fig. S1. Network disruption by rmTBI is not widespread. No differences in the Network strength **(A)**, Louvain modularity **(B)**, Assortativity **(C)**, Transitivity **(D)** and Gamma **(E)**. Net lambda was significantly decreased in rmTBI group at various density threshold levels compared to sham **(F)**. Multiple unpaired t-test, mean error ± SD at edge densities ranges 2 to 40%.

Table Legend:

Table. 1. Summary of the expression of various neuropathological markers following rmTBI identified through NanoString GeoMx spatial protein profiling analysis.

Supplementary table Legend:

Supplementary table. 1. The complete list of 64 ROIs (left and right) analyzed through rsfMRI and graph theory.

688   **Table. 1**

ROIs	Protein Biomarker	Expression	Full target name	Cell profiling panel	p-value
Primary somatosensory area	Myelin basic protein	Down	Myelin basic protein	Neural Cell Profiling Core	0.0059
Thalamus	GPNMB	Up	glycoprotein NMB	Glial cell subtyping module	0.0157
	Ki-67	Down	marker of proliferation Ki-67	Neural Cell Profiling Core	0.0466
Optic tract	CTSD	Up	cathepsin D	Glial cell subtyping module	0.0014
	Iba1	Up	allograft inflammatory factor 1	Neural Cell Profiling Core	0.0024
	pTau(S199)	Up	microtubule-associated protein tau	AD pathology extended	0.0067
	GFAP	Up	glial fibrillary acidic protein	Neural Cell Profiling Core	0.0110
	Tau	Down	microtubule-associated protein tau	AD pathology module	0.0092
	MERTK	Down	MER proto-oncogene, tyrosine kinase	Glial cell subtyping module	0.0273
	pTau(S396)	Down	microtubule-associated protein tau	AD pathology extended	0.0491

 Supplementary table. 1

ROI #	Abbreviation	Region
1	L VISAA	Visual Anterior area
2	L VISRLA	Visual Rostrolateral area
3	L VISAMA	Visual Anteromedial area
4	L VISLA	Visual Lateral area
5	L VISPV1	Visual primary visual
6	L VISPV2	Visual primary visual
7	L VISPL	Visual Posterolateral
8	L VISPM	Visual Posteromedial
9	L VISPR	Visual postrhinal
10	L THA MED	Thalamus mediodorsal
11	L THA L	Thalamus anterolateral complex of ventral thalamus
12	L THA AC	Thalamus laterodorsal
13	L THA PC	Thalamus posterior complex
14	L THA VP	Thalamus ventral posteromedial
15	L THA para	Thalamus paracentral nucleus
16	L THA PL	Thalamus posterior lateral
17	L THA CL	Thalamus central lateral – parafascicular
18	L THA AP	Thalamus anterior pretectal nucleus
19	L THA DL	Thalamus dorsal lateral geniculate
20	L THA V	Thalamus ventromedial
21	L TEMP	Temporal association areas
22	L DEN	dentate nuclei of cerebellum
23	L CERE	interposed complex of cerebellum
24	L CP G	genu of corpus callosum
25	L CP1	Body of corpus callosum 1
26	L CP2	Body of corpus callosum 2
27	L SCP1	splenium of corpus callosum 1
28	L SCP2	splenium of corpus callosum 2
29	L SCP3	splenium of corpus callosum 3
30	L OT1	Optic tract 1
31	L OT2	Optic tract 2
32	L OT3	Optic tract 3
33	R VISAA	Visual Anterior area
34	R VISRLA	Visual Rostrolateral area
35	R VISAMA	Visual Anteromedial area
36	R VISLA	Visual Lateral area

37	R VISPV1	Visual primary visual
38	R VISPV2	Visual primary visual
39	R VISPL	Visual Posterolateral
40	R VISPM	Visual Posteromedial
41	R VISPR	Visual postrhinal
42	R THA MED	Thalamus mediodorsal
43	R THA L	Thalamus anterolateral complex of ventral thalamus
44	R THA AC	Thalamus laterodorsal
45	R THA PC	Thalamus posterior complex
46	R THA VP	Thalamus ventral posteromedial
47	R THA para	Thalamus paracentral nucleus
48	R THA PL	Thalamus posterior lateral
49	R THA CL	Thalamus central lateral – parafascicular
50	R THA AP	Thalamus anterior pretectal nucleus
51	R THA DL	Thalamus dorsal lateral geniculate
52	R THA V	Thalamus ventromedial
53	R TEMP	Temporal association areas
54	R DEN	dentate nuceli of cerebellum
55	R CERE	interposed complex of cerebellum
56	R CP G	genu of corpus callosum
57	R CP1	Body of corpus callosum 1
58	R CP2	Body of corpus callosum 2
59	R SCP1	splenium of corpus callosum 1
60	R SCP2	splenium of corpus callosum 2
61	R SCP3	splenium of corpus callosum 3
62	R OT1	Optic tract 1
63	R OT2	Optic tract 2
64	R OT3	Optic tract 3

Fig. 1. Study design. Schematic representation showing the number of rmTBIs, rsfMRI and tissue collection. Brains were collected at 7 dpi for Immunohistochemistry (IHC) and nanostring GeoMx spatial protein profiling analyses.

Fig. 2. Clustering coefficient analysis reveals that the network integration and efficiency in specific brain regions are disrupted following rmTBI. Representative T2-weighted anatomical brain images of sham and rmTBI mice **(A&B)** were used for brain volume analysis. Quantification of brain volume revealed no changes between sham and rmTBI groups **(C)**.
3D functional connectome maps of sham and rmTBI mice brain shows the position of nodes and edges in the region of interest (ROIs) [total ROI = 64] **(D & E)**. Clustering coefficient in various regions of thalamus and visual area **(F-I)** were significantly altered in rmTBI group compared to sham. Unpaired t-test, mean ± SD, Sham (N= 8), rmTBI (N=10).

Fig. 3. Global and Local efficiency analysis reveals that efficiency in specific brain regions are disrupted following rmTBI.
Global efficiency **(A)** was significantly decreased at the density threshold of 4% and 6% in rmTBI group compared to Sham. Local efficiency at 4% threshold **(B & C)** was significantly decreased in thalamus anterior pretectal nucleus (L) and thalamus mediodorsal (R) regions of Injured mice. Similarly, Local efficiency at 6% threshold **(D-F)** was significantly decreased in thalamus anterior pretectal nucleus (L), thalamus dorsal lateral geniculate (L) and visual anterior area (R) of injured mice compared to Sham. No changes in global clustering coefficient **(G)** and small world index **(H)** were observed. The global characteristic pathlength **(I)** at 4% threshold was significantly increased in injured mice group compared to sham. The global metrices were analyzed by Multiple Unpaired t-test, mean ± standard error at edge densities ranges 2 to 40%. The groups comparison was analyzed by Unpaired t-test, mean ± SD, Sham (N= 8), rmTBI (N=10).

Fig. 4. rmTBI disrupted connectivity between modules, increased eigenvector centrality and node betweenness centrality. Nodes in left (L) thalamus ventral posteromedial **(A)** region had low participation coefficient, but it is significantly increased in the right (R) splenium of corpus callosum, thalamus (R) and visual regions (R) of rmTBI group relative to control **(B-E)**. Increased eigenvector centrality shows increased high influence of specific nodes on connected networks following rmTBI in thalamic regions and corpus callosum **(F-I)**. The node betweenness centrality shows the nodes at the intersection between shortest path in the specific thalamus region **(J)** was significantly altered. Density threshold at which the participation coefficient and node eigenvector centrality were calculated Unpaired t-test, mean ± SD.

Fig. 5. rmTBI altered tissue microstructure in specific brain regions. Representative DTI brain image shows **(A)** ROIs selected for microstructure pattern analysis. rmTBI increased fractional anisotropy (FA) in the optic tract **(B)**. Mean diffusivity (MD) and radial diffusivity (RD) were significantly increased in hippocampus of the injured mice **(C&D)**. Axial diffusivity (L1) was significantly increased in the optic tract and hippocampus of the injured mice **(E)** compared to sham. Unpaired t-test, mean ± SD.

Fig. 7. rmTBI increased the expression of disease associated proteins in white matter and grey matter regions. Coronal sections of sham and rmTBI mice shows the region of interest (primary somatic sensory area, dentate gyrus polymorph layer, thalamus, and the optic tract) selected for NanoString GeoMx spatial protein profiling analysis **(A&B)**. Volcano plots **(C)** and bar graphs **(D-J)** show significant alteration in the expression of Alzheimer's disease (AD) pathological markers, glial and neural cell markers in the optic tract of the injured mice. Volcano plot **(K)** and bar graphs show **(L&M)** significant changes in the expression of cell proliferation and AD associated, neuroinflammatory markers in the thalamus of the injured mice.

Myelin basic protein was significant decreased in the primary somatosensory area following rmTBI **(N&O)**.

Protein levels were normalized to the expression of house keeping proteins. Unpaired t-test, mean ± SD, N=4-8.

Fig. 6. rmTBI induced gliosis in white matter tracts. Iba1 expression was significantly increased in the optic tract of injured mice compared to sham **(A-C)**. The red arrowhead indicates microglia. Brain coronal view: 1 mm; Scale bar for inset (enlarged view of the optic tract) = 100 µm. ****p<0.0001, Unpaired t-test; Sham (N= 15), rmTBI (N=20). GFAP expression was significantly increased in the optic tract **(D-F)** and corpus callosum **(G-I)** of injured mice compared to sham. The blue arrowhead indicates astrocytes. Brain coronal view: 1 mm; Scale bar for inset (enlarged view of the optic tract and corpus callosum) = 100 µm. (****p<0.0001; *p= 0.0320, Unpaired t-test), Sham (N= 15), rmTBI (N=19).