Title: Dynamic refinement of behavioral structure mediates dopamine-dependent credit

assignment

Dopamine initially reinforces spatially similar and temporally proximal actions to actions that

trigger dopamine release, and drives a gradual refinement of the entire behavioral repertoire to

home-in on reward-producing actions.

Authors: Jonathan C.Y. Tang[1], Vitor Paixao[2,3], Filipe Carvalho[2,4], Artur Silva[2], Andreas Klaus[2],

Joaquim Alves da Silva[2], Rui M. Costa[1,2,5]*

Affiliations:

[1]Department of Neuroscience, Zuckerman Mind Brain Behavior Institute, Columbia University,

New York, NY 10027, USA

[2]Champalimaud Neuroscience Programme, Champalimaud Research, Champalimaud Foundation

Lisbon, Portugal

[3]Kinetikos, Coimbra, Portugal

[4]Open Ephys Production Site, Lisbon, Portugal

[5]Allen Institute, Seattle, WA 98109, USA

*Correspondence: rc3031@columbia.edu

Abstract

Animals exhibit a diverse behavioral repertoire when exploring new environments and can learn which actions or action sequences produce positive outcomes. Dopamine release upon encountering reward is critical for reinforcing reward-producing actions[1-3]. However, it has been challenging to understand how credit is assigned to the exact action that produced dopamine release during continuous behavior. We investigated this problem with a novel self-stimulation paradigm in which specific spontaneous movements triggered optogenetic stimulation of dopaminergic neurons. We uncovered that dopamine self-stimulation rapidly and dynamically changes the structure of the entire behavioral repertoire. Initial stimulations reinforced not only the stimulation-producing target action, but also actions similar to the target and actions that occurred a few seconds before stimulation. Repeated pairings led to gradual refinement of the behavioral repertoire leading animals to home in on the target action. Reinforcement of action sequences revealed further temporal dependencies of behavioral refinement. Action pairs that tend to be spontaneously separated by long time intervals promoted a stepwise credit assignment, with early refinement of actions most proximal to stimulation and subsequent refinement of more distal actions. Thus, a retrospective reinforcement mechanism promotes gradual refinement of the entire behavioral repertoire to assign credit to specific actions and action sequences that lead to dopamine release.

Main Text

Background

Animals spontaneously transition amongst a repertoire of movements when exploring new

environments. Movements or movement sequences that produce positive outcomes are

reinforced and increase in frequency to maximize the obtainment of those outcomes[4,5].

However, it is still not completely clear how animals assign credit to the exact action that

produce reward in the context of a continuous behavioral space. This credit assignment

problem[2,6–9] during spontaneous behavior poses at least two main challenges. First, it is unclear

how animals come to preferentially perform a specific reward-producing action or action

sequence above other possibilities in the behavioral repertoire. Second, it is unclear how animals

derive contingency between a reward-producing action and reward if there can be variable delays

between action performance and reward delivery.

Dopamine (DA) has been proposed to mediate credit assignment[6,10]. At the cellular level, DA

can facilitate synaptic plasticity in corticostriatal synapses[11] within a critical time window that is

behaviorally relevant[12–14]. Still, it is unknown how DA changes the dynamics of spontaneous

behavior to mediate credit assignment. We therefore developed a paradigm to investigate how

DA shapes the evolution of continuous behavior during action learning to gain insights into the

process of credit assignment.

Conventional operant conditioning paradigms[5,15–19] have helped derive principles of behavioral

reinforcement, but they are not ideal for studying action credit assignment. In general, such

paradigms do not permit the clean isolation of actions as the trigger for reward versus particular

locations or objects. In such paradigms, animals are also required to perform a series of

consummatory actions, such as approaching and interacting with reward-delivering devices to

retrieve reward. These requirements make it difficult to investigate how credit is assigned to a

specific action or action sequence in the behavioral repertoire during continuous behavior.

Until recently, technological and conceptual limits have made it difficult to study how the entire

structure of continuous behavior evolves as naive animals come to associate specific action or

action sequences with reward. To address previous limitations, we developed a new approach to

study action credit assignment. This approach directly reinforces specific spontaneous action(s)

by triggering dopaminergic neuron (DA neuron) excitation and DA release upon action

performance. It combines wireless inertial sensors, unsupervised clustering of continuous

behavior[20,21] and optogenetics[22] into a closed-loop system linking specific action performance to

immediate phasic DA release (Methods; Fig. 1a-f). This paradigm permits action detection and

reinforcement without requiring an animal to approach or interact with a place/object/cue, or to

perform consummatory behavior. These combined features overcome the aforementioned

caveats associated with conventional paradigms.

Rapid reinforcement of actions via closed-loop dopamine stimulation

To implement the action detection component of the closed loop system, we first classified the

entire behavioral repertoire of individual mice[23] mice in a grey-walled open field using inertial

sensors and unsupervised affinity propagation clustering[20,21] (Fig. 1d). Self-paced behavior was

monitored using a novel, wireless inertial sensor system (WEAR; Methods) that allows minimal

movement restraints, high resolution behavior monitoring and fast data transmission to open-

source hardware and software for online experimentation (Fig. 1b, Extended Data Fig. 1a).

Affinity propagation clustering is particularly well suited to cluster an unknown number of

clusters[20], is computationally efficient[24], and easily outputs similarity between clusters.

Clustering begins by processing accelerometer and gyroscope data to extract 4 features

discriminating postural changes, movement momentum, head and head-body rotations, and total

body accelerations. Feature values from 300 ms long segments of behavior were discretized into

histograms, upon which pairwise similarity comparisons could be made using a Earth-Mover's

Distance (EMD)[25] metric. The similarity matrix of all possible pairwise comparisons were fed

into an unsupervised affinity propagation clustering algorithm[20] (Methods), identifying naturally

occurring repertoire of 300 ms long behavioral clusters[21], or "actions" (Fig. 1c,Extended Data

Fig. 1b). The choice of 300 ms long movements was informed by previous studies[21,26]. Using

these parameters, we identified over 30 clusters of spontaneous behavior per individual (34.3 +/-

2.1 and 35.6 +/- 2.5 total actions per ChR2-YFP and YFP mice, respectively; mean +/- standard

deviation,15 ChR2-YFP and 10 YFP mice). We chose particular clusters of actions to be

reinforced (hereby named target action A).

To implement closed-loop reinforcement, we used Cre-dependent AAV viruses (EF1a-DIO-

expression cassette) to express channelrhodopsin ChR2-YFP[22] or the control protein YFP

bilaterally in DA neurons of the ventral tegmental area (VTA) [27,28] of DAT-Cre mice (Fig.

1a,Extended Data Fig. 2a-c). Using the wireless inertial sensor, we tracked behavior

continuously in a white open field and used the similarity metric to match ongoing 300 ms

behavioral segments to exemplars representing each mouse's repertoire of actions (Fig. 1d-e).

Upon a match to a defined target action (target action A), a 25 hz, 600 ms long train of

optogenetic stimulation was delivered to DA neurons of the VTA parabrachial pigmented area

(PBP) (30-60 ms delay, Fig. 1e). These target action As were different for different animals, and

were dispersed across a behavioral space (Fig. 1g).To evaluate whether stimulation parameters

triggered DA release similar in magnitude to that triggered by sucrose reward in food restricted

mice, we delivered random optogenetic stimulations to ChR2-YFP- or YFP-expressing VTA DA

neurons while monitoring DA release with the GRAB rDA1m sensor [29] in both ventral and

dorsal striatum (Fig. 1f). We also measured DA release in the same animals upon delivery of

sucrose while they were food deprived. Sucrose presentation led to a sharp increase in DA

release in both ventral and dorsal striatum (Fig. 1f). Interestingly, optogenetic stimulation of DA

neurons in VTA with the parameters described above, resulted in a similar phasic increase in DA

not only in ventral striatum but also in dorsal striatum (Fig. 1f). This is consistent with emerging

evidence showing the existence of dorsal striatum-projecting VTA neurons[30,31]. Thus, our

optogenetic stimulation triggered DA release similar in decay and spatial localization to that

triggered by sucrose reward in food restricted mice (Fig. 1f), offering us a suitable approach to

interrogate how pairing DA release with specific action performance leads to credit assignment.

Closed loop reinforcement for a specific action occurred over a 3-day, 60-90 minute/session

protocol designed to probe both intra- and inter-session changes in behavior (Fig. 1h-m,

Extended Data Fig. 3). Optogenetic stimulation of VTA DA neurons upon execution of a

particular target action (action A) resulted in significant increase in the frequency of action A for

ChR2-YFP, but not YFP mice (Fig. 1h, Extended Data Fig.3b). Increased action A in ChR2-YFP

animals depends on optogenetic stimulation, as removal of closed-loop stimulations resulted in

progressive extinction of action A (Fig.3h, Extended Data Fig.3d). Resuming paired stimulation

led to rapid re-instatement of action A (Fig. 1h, Extended Data Fig.3c,e). Interestingly, during

extinction, ChR2-YFP animals kept performing exploratory unrewarded bursts of action A,

which could explain rapid reinstatement (Extended Data Fig. 3e,f). This paradigm revealed that

just a few pairings with DA leads to rapid reinforcement, as changes in multiple parameters

including decreased trigger latency, increased action A frequency and increased average

behavioral similarity towards action A become significant following 10-15 stimulations (Fig. 1i,

Extended Data Fig. 4a-b).

We next examined if only action A changed in frequency or if other non-stimulated actions also

changed with closed-loop reinforcement of action A. We calculated baseline-normalized

frequency of all actions in the repertoire and ordered them as a function of similarity to the target

action (Fig. 1j). Earth-Mover's Distance (EMD)[21,25] was used to measure each action exemplar's

similarity to the target exemplar (Methods), with lower EMD value indicating increased

similarity. Surprisingly, we observed that optogenetic stimulation resulted in a dramatic change

in the entire behavioral repertoire. We observed that early in training actions most similar to

target tended to also increase in frequency (Fig. 1j-l, Extended Data Fig. 4c) whereas actions

most dissimilar to target tended to decrease in frequency. Repeated pairing led to refinement of

the actions that were performed at high frequency, and by late stages action A became the

predominant action being performed, with a sharp drop-off of non-target action frequencies as

similarity to target decreased (Fig. 1k-l). Such effects were not observed in YFP controls

(Extended Data Fig. 4d-e). These data suggested that early reinforcement results in rapid

reshaping of the entire behavioral repertoire, biasing animals towards actions similar to the target

action, and continued pairing resulted in gradual refinement and assignment of credit to the

specific target action.

Dynamics of behavioral refinement during reinforcement

To better describe individual action dynamics during reinforcement, we categorized actions (511

actions, *n*=15 ChR2-YFP animals) by the trajectories of their changes in frequency throughout

learning (Methods). Three meaningful types of trajectories were categorized, comprising over

94% of all actions. These types were characterized by either initial increase that remained stable

(Sustained Increase), initial increases that decreased over time (Transient Increase) and initial

decreases that remained stable (Decreased) (Fig 1m, Extended Data Fig. 5-6). We again

confirmed that the frequency dynamics type of each particular action was related to its similarity

to target, regardless of whether actions were sorted based on their raw or percentile similarity

scores (Extended Data Fig. 6b-c). Actions most similar to target were predominately Sustained

Increase types, while moderately similar actions mostly comprised of Sustained Increase or

Transient Increase types and more dissimilar actions are more of the Decreased type (Extended

Data Fig. 6b-c). Taken together, these finer resolution analyses indicate again that the dynamics

of action frequency are related in great part to the similarity to target action.

Reinforcement and refinement after reversal of action-reward contingencies

Next, we asked if animals could follow changes in contingency between action and closed-loop

DA stimulation. We therefore chose a different action, action B, which is clearly distinct from

the action A for each animal (Methods, Fig. 2a, Extended Data Fig. 1c) and started delivering

DA neuron optogenetic stimulation after action B. Chosen action A/B pairs were relatively

dissimilar in the context of entire action similarity distributions (Fig. 2b). Upon reinforcement,

previously trained ChR2-YFP, but not YFP animals showed increased action B performance

over time (Fig. 2c-e, Extended Data Fig. 7). In contrast, action A frequency changes clearly

moved in the opposite direction from that of action B over time (Fig. 2c). Maintenance of action

B performance depended on continual reinforcement (Fig. 2c, Extended Data Fig. 7d-e). Similar

to action A, action B credit assignment unfolds by initially biasing the entire repertoire, i.e.,

increasing the frequency of similar actions and reducing the frequency of dissimilar actions. This

was again followed by gradually refining for action B relative to similar actions as pairing

progressed (Fig. 2d-e, Extended Data Fig. 7f). To confirm that action learning is contingent on

action B appearing before reinforcement, we subjected trained animals to a contingency

degradation protocol in which we delivered a similar number of random stimulations uncoupled

to action B performance. Action B performance decreased following contingency degradation

and could be re-instated upon resuming the action B-stimulation contingency (Fig. 2f, Extended

Data Fig. 7g). These experiments indicate that animals can follow changes in the contingency

between actions and DA release and assign credit to a new action through a similar process of

behavioral repertoire refinement.

Although animals show similar patterns of behavioral refinement for actions A and B, animals

that previously credited an action (action A) for DA release did initially respond to reinforcement

of a new action (action B) differently from naïve animals (Fig. 2g-j). Whereas naïve animals

responded to initial reinforcements for target action A by significantly increasing action A

performance relative to the non-target action B (Fig. 2g,i,left graph), animals with a history of

reinforcement on action A animals responded to initial reinforcements of action B by increasing

non-target action A performance (Fig. 2g,i,right graph). This trend reverses later such that target

action B becomes significantly increased over the non-target action A (Fig. 2g,i,right graph).

YFP control animals showed no such trends (Fig. 2h,j). Thus, DA reinforcement does not simply

reinforce the recently performed, temporally contiguous action, but trigger previously credited

actions in the face of a new action-reward contingency that is not yet learned. This suggest again

that animals learned the contingency between action performance and DA release.

Temporal constraints of DA-dependent reinforcement

The contingency degradation results above indicate that the temporal relation between target

action and DA phasic activity is important for reinforcement (Fig. 2e). Reinforcement is thought

to occur on behavior that precedes reward in time[10,12,14,19], and while temporal contiguity

between action and reinforcement has long been recognized[32–34], it is not clear how the position

of an action relative to the time of DA phasic activity influences its subsequent frequency. We

investigated if in addition to behavioral similarity, the temporal relationship between action and

stimulation influenced the dynamics of behavioral repertoire evolution during reinforcement and

credit assignment.

We observed that the median inter-target action interval decreased with stimulation in ChR2-

YFP mice (Fig. 3a,b). We therefore examined the distribution of the action dynamic types

categorized above (Sustained Increase, Transient Increase, Decreased) according to both an

action's similarity to target and the median time of that action's performance leading into target

during baseline, before reinforcement protocol began (Fig. 3c-e). Action dynamic types showed

distinct distribution patterns for these two dependent variables (similarity and time). Further,

these two dependent variables were not significantly collinear (Methods). Thus, action similarity

to target as well as baseline temporal proximity to target should together predict action dynamic

type upon reinforcement better than either factor alone. To test this idea, we performed

multinomial logistic regression to assess whether 1- or 2-factor models best fit the observed

dynamics pattern that an action would follow upon reinforcement (Fig. 3f,g). The two-factor

model outperformed either one-factor models, and prediction of action dynamics type with this

model was significantly above chance as assessed by precision-recall curves, which is suitable

for evaluating datasets with imbalanced categories[35] (Fig. 3g). The beta coefficients indicated

that increased similarity to target and decreased median time to target increases prediction of

Sustained Increase and Transient Increase dynamic types relative to Decreased types

(Supplementary Table). These results suggest that DA may reshape behavioral repertoire by

reinforcing not only actions similar to the target action but also actions that happen to be

performed temporally close to the reinforcer, as suggested before[10,12,14,19].

To more rigorously test whether DA reinforcement acts in a retrospective or prospective manner,

we increased the resolution of analysis by examining 1st order action transitions leading into and

out of stimulation (Fig. 3h-j). By focusing analysis on action transitions enriched within specific

1.2 second moving windows, one could distinguish more clearly behavior that occurred leading

up to, during, and after DA stimulation. Our analyses showed that action transitions enriched in

windows up to 1.2 seconds prior to stimulation onset, as well as during stimulation, are

reinforced early on (Fig. 3i). However, this did not occur to action transitions following

stimulation, suggesting an asymmetric process. Indeed, action transitions enriched in windows

leading into stimulation were also preferentially reinforced relative to those enriched in windows

after stimulation (Fig. 3j). Thus, DA stimulation promotes reinforcement of behaviors occurring

during stimulation and a few seconds before stimulation.

Credit assignment for action sequences

In the real world, when animals are spontaneously shifting between actions in their repertoire,

outcomes are often not the result of a single action but rather of a sequence of actions performed

at variable intervals. We therefore investigated the dynamics of reinforcement when the release

of DA is contingent upon the performance of a sequence of 2 actions (target action 1 and 2, T1

and T2). We applied closed loop optogenetics to ask whether naïve animals can learn a T1$\rightarrow$T2

reinforcement rule, where the delays between T1 and T2 are governed by the spontaneous

behavior of the animals and not experimentally controlled (n=15 ChR2-YFP and 10 YFP mice,

Fig. 4a, Extended Data Fig. 2a,d-e, Extended Data Fig. 8-10). Various T1/T2 pairs were

sampled, with focus on sequences sharing general commonalities in movement order across

animals (Extended Data Fig. 1d,f-g). Overall, mice learned to increase the performance of a

sequence of two actions to obtain DA stimulation. Some animals showed a ChR2-dependent

increase in reinforcement within 5 sessions, but others experienced a lag in learning (Fig. 4b).

We hypothesized that this could relate to the initial time distance between T2 trigger and the

closest distal T1 (T1$\rightarrow$T2 interval). Indeed, animals reinforced for action pairs with initially long

interval values tended to show slower learning curves (Fig. 4c-d). To capture a learning time

point whereby individuals reach similar rising phase in their respective learning curves, a

criterion frequency was set (Methods). 14 of 15 trained animals eventually reached criterion

(Fig. 4e; Extended Data Fig. 8a-c). Sequence performance depended on continuing DA

reinforcement (Fig. 4f,g). Learning was also revealed by decreases in the median T1$\rightarrow$ T2 time

intervals (Fig.4h-i) and convergence of T1-to-T2 frequency ratio towards 1 (Fig. 4j). To quantify

the specific credit assignment of T1 and T2 we used a refinement index that compares the

median frequency of actions uniquely similar to T1 with those uniquely similar to T2, with the

frequencies normalized by either that of T1 or T2 (Methods). Values lower than 1 indicate that

the target actions are being performed even more frequently than similar actions, and thus

indicate greater refinement (Methods). By the end of learning, T1 and T2 became credited as the

reward-producing actions relative to their similar counterparts (Fig. 4k). YFP controls did not

show any of these trends (Fig.4c-d,4g-h). Thus, closed loop reinforcement promoted learning of

a two-action sequence rule in freely moving mice starting from a naïve state.

Importantly, the initial median T1$\rightarrow$T2 interval performed by ChR2-YFP animals was inversely

related to the eventual number of sessions required for each animal to reach criterion frequency

(Fig. 4l). A sigmoidal curve was fit to the data, showing that animals with longer open field

T1$\rightarrow$T2 intervals beyond the sigmoidal midpoint tended to face sudden increase in sessions to

reach criterion frequency (Fig. 4l). ChR2-YFP animals were divided according to the half-

maximum point of the sigmoidal curve into 'Fast Learners' and 'Slow Learners'. Fast Learners

quickly reached criterion frequency and low T1$\rightarrow$T2 time intervals, whereas Slow Learners

experienced a time lag in reaching criterion frequency and low T1$\rightarrow$T2 intervals. Slow Learners

tended to suddenly increase the frequency of sequence performance in sessions that showed a

drop in the median T1$\rightarrow$T2 interval to below 2-4 seconds (Fig.4d,h). In contrast, there was no

stable sigmoidal relationship between T1-T2 action similarities and sessions to criterion

frequency (Extended Data Fig. 8d). Thus, the initial median time distances between distal action

T1 and proximal action T2(which produced DA stimulation) modulated how fast animals learned

to effectively perform the reinforced action sequence.

If DA is acting retrospectively to reinforce actions performed earlier in time, we hypothesized

that the action most proximal to reinforcement, T2, should experience earlier refinement relative

to the distal action, T1. We again used the median target normalized frequencies of actions

uniquely related to T1 or T2 as refinement indices (Methods). Proximal T2 clearly refines

towards its most refined level earlier than the distal T1, at least in some animals (Fig. 5a). By

subtracting the area under the refinement curve for T1 from the curve for T2, one could calculate

differential refinement between the two actions. Positive values indicate refinement

preferentially favoring T2, and vice versa. A linear relationship was found between open field

median T1→T2 interval and differential refinement between T1 and T2 (Fig. 5b). This suggests

for longer T1→T2 median intervals, the proximal action T2 spends more sessions being more

refined than the distal action T1. In contrast, there was no significant linear relationship between

the initial intervals between the execution of the proximal action that led to reward and the next

initiation of the sequence (T2→T1) or of the similarity between T1 and T2 actions, and the

dynamics of differential refinement between T1 and T2 (Fig. 5b, right graph, Extended Data Fig.

9a).

We next investigated if the differential refinement between T1 and T2 was different for slow and

fast learners. We analyzed changes in T1-T2 refinement curves relative to 'Starting Points' at

which the refinement indices of T1 and T2 are most similar or are biased towards the distal T1

rather than the proximal T2 action (Methods). All Slow Learners showed a pattern where they

initially refine the repertoire of T2 from these Starting Points, and after reaching a maximum

Turning Point, they start showing a bias towards greater T1 refinement (Fig. 5c). Notably, by

these Turning Points the median intervals of T1→T2, but not T2→T1 events had decreased

significantly relative to initial values (Fig. 5d, Extended Data Fig. 9b). Therefore, the median

T1→T2 interval decrease occurred before a decrease in the interval to perform the next sequence

(T2→T1), which started decreasing after the Turning Point (Fig. 5e). Using these learning

landmarks, we asked more rigorously how animals homed in on T1 vs T2 over time (Fig. 5f,

Extended Data Fig.10a). We found that animals initially refined the action proximal to DA

stimulation (T2, between Starting Point and Turning Point), whereas T1 refinement occurred

several sessions later, after the Turning point (Fig. 5f, Extended Data Fig.10a). Indeed, the

Turning Point coincided with an increased probability of the T1 being found within 3.6 secs

before T2 and reinforcement (Fig. 5g-h). These results indicate that animals can assign credit to

sequences of actions that lead to reinforcement, following similar retrospective dynamics that

were observed for single actions, whereby the actions most proximal to reinforcement are refined

earlier and the actions more distal to reinforcement refined later, when they probabilistically start

to occur within a few seconds of DA release.

Discussion

Our results demonstrate that DA reinforcement promotes single action credit assignment from a

naïve state through a dynamic process whereby the entire behavioral repertoire is restructured.

During the initial stages of reinforcement both actions similar to the target action and actions that

were performed in close temporal proximity of the target action increase in frequency, while

very dissimilar actions decrease in frequency. With repeated reinforcement there is a process of

gradual refinement that homes in on the action that produces DA release. In the case of action

sequences, we observe a similar gradual refinement process whereby credit assignment for the

action sequence is accomplished by early refinement for the actions most temporally proximal to

reinforcement, followed by later refinement for the more temporally distal actions.

Previous synaptic and cellular studies[36,37] proposed that DA reinforcement may act

retrospectively to reinforce behavior. By utilizing the closed loop system, we rigorously tested

this prediction. Since retrospective reinforcement of behavior is not confined to the target action

alone, it facilitates credit assignment to a stimulation-producing action even when reinforcement

is delayed; stimulation-producing action pairs that tend to be performed closed together in time

were learned much faster than pairs that tended to be performed far apart in time. Intriguingly,

animals eventually learned to assign credit to distal stimulation-producing actions even in the

latter scenario. This is characterized by a gradual process whereby early on, the median time

interval between distal and proximal target actions decreased and the repertoire proximal to

reinforcement was preferentially refined to favor the performance of the proximal target action.

As the distal target action became significantly more likely to occur within second timescale

distance prior to reinforcement, retrospective reinforcement of the correct stimulation-producing

sequences became increasingly likely, resulting in whole behavioral refinement for the distal

target as well, hence increasing sequence performance (Fig. 5g).

It has been suggested that retrospective reinforcement of behavior is mediated by DA modulation

of an eligibility trace left by action potential-triggered synaptic plasticity[10]. Studies of DA action

at the striatal synaptic level[36,37] indicate that the timescale within which retrospective

reinforcement may occur is on the order of a few seconds, but the behavioral consequences have

remained elusive until now. Our behavioral findings are consistent with cellular studies in that

behavior occurring within a few seconds leading into DA stimulation are reinforced. It is also

noteworthy that distal T1 refinement in two action reinforcement occurs after the closest T1 to

DA stimulation has become more probable within a few seconds of stimulation. The cutoff of

retrospective reinforcement by phasic DA activities within a few seconds could explain the

sudden increase in sessions required to reach criterion frequency amongst animals that were

reinforced for action pairs with initially longer median time separations. Retrospective

behavioral reinforcement may be mediated by DA modulation of Ca2+ influx left by earlier

spiking activities. Ca2+ influx triggered by NMDA receptors would increase adenosine 3',5'-

cyclic monophosphate at thin distal dendrites of medium spiny neurons, leading to transient and

localized protein kinase A activity specifically within the retrospective time window, as

regulated by high phosphodiesterase activity[14]. Similar actions have more similar and

overlapping striatal neural ensemble activities[21]. Arrival of DA upon activation of action-specific

ensembles may reinforce not only a specific action, but also similar actions. As striatal

ensembles specific to actions are activated and a trial of eligibility traces is left temporally, DA

arrival could set the stage for retrospective reinforcement of a spatially graded repertoire of

actions within a few seconds, resulting in the observed behavioral learning patterns. Future

studies testing these ideas would clarify how synaptic plasticity and cellular ensemble activities

integrate to produce a dynamic refinement process, resulting in the behavioral principles for

credit assignment revealed here.

END OF MAIN TEXT

Methods

Animals: All experiments were approved by the Portuguese DGAV and Champalimaud Centre for the Unknown Ethical Committee and performed in accordance with European guidelines. They were also performed according to National Institutes of Health (NIH) guidelines and approved by the Institutional Animal Care and Use Committee of Columbia University. 3-5 months old DAT-Cre male mice in the C57/BL6J background[23] were used.

Sample Sizes, randomization, and blinding. For sample size, we applied a power of 0.8, significance of $p<0.05$, and standard variation of 20% of the mean. We determined sample sizes of 4-8 mice per group for different mean-based tests (matched pairs, 2 groups). No formal method of randomization was used; littermates were equally divided among the groups being compared. The experimenter was not blinded of the experimental groups. Optogenetic manipulations were performed automatically via a computer algorithm and not manually by the experimenter.

Recombinant adeno-associated viral vectors, stereotaxic injections, and implants. 750 nl of rAAV.EF1a.DIO.hChR2(H134R).eYFP or rAAV.EF1a.DIO.eYFP ($3-4 \times 10^{12}$ vg/ml, AAV5, University of North Carolina Vector Core; $1-2 \times 10^{13}$ vg/ml, AAV1, Addgene, 27056-AAV1 and 20298-AAV1) were injected into each hemisphere of the VTA of 3-4 month old DAT-Cre mice. For viral injections, the coordinates are AP - 3.52 mm, ML - +/- 0.35 mm, DV – 4.3 mm. Injections were made at 0.2 Hz pulses. Each pulse injects 4.6 nl volume. Injected needles were kept in place in the injection site for ~15 minutes before withdrawal. For each mouse, a dual optic fiber cannula (200/240 μm diameter, 6 mm length, 0.7 mm center-to-center FLT, 0.22 NA;

Doric, DFC_200/240-0.22_6mm_DF1.0_FLT) was placed 200 μm above the injection site and

fixed to the skull. Next, a 4-position receptacle connector (Harwin Inc., M52-5000445) was fixed

anteriorly to the dual optic fiber cannula, with its posterior edge set at -0.6 mm. Skull implants

are then fixed with dental cement. A 4-position connector (Harwin Inc., M52-040023V0445)

with pins removed from one end was used to cap the receptacle connector.

For photometry experiments, 3-5 month old DAT-Cre males were used. The conditions used for

VTA injections and implants were as above. Additionally, 1 μl and 500 nl of AAV9-hSyn-

GRAB-rDA1m (2 x 10^13 vg/ml; Addgene, 140556-AAV9) were injected into the dorsal

striatum (AP 0.5 mm, ML +2.1 (right), DV 2.3 (from brain surface)) and ventral striatum (AP

1.15mm, ML +1.65 (right), DV 4.2 (from Bregma)) , respectively. For photometry fiber

implants, mono fiberoptic cannula were used (400/430 μm diameter, 4 mm length (dorsal

striatum) and 6 mm length (ventral striatum), 0.37 NA, 1.25 diameter ferrule, flat; Doric,

MFC_400/430-0.37_6mm_MF1.25_FLT (ventral striatum) and MFC_400/430-

0.37_4mm_MF1.25_FLT (dorsal striatum)). Implants were inserted at a 22 degrees angle. For

dorsal striatum implantation, the cannula entered the skull at AP 0.5 mm and ML 3.03 mm at 22-

degree angle. The angled implant penetrated the brain from its surface for 1.92 mm. For ventral

striatum implantation, the cannula entered the skull at AP 2.85 mm at 22 degrees angle, ML 1.65

mm. The angled implant penetrated the brain from its surface for 4.25 mm.

WEAR motion sensor system. The WEAR motion sensor family was developed by the

Champalimaud Hardware platform and Costa lab as a wired or wireless solution to obtain self-

centered 9-axis motion data based on 3-axis accelerometer, gyroscope, and magnetometer

(https://www.cf-hw.org/harp/wear). The wired version is a very small and extremely lightweight

device (200mg) that can sample motion data up to 500 Hz and at the same time provide current

up to 500mA that can be used to power LEDs for optogenetic experiments or stimulating

electrodes. The wireless version is small and lightweight (~1.8g) and can sample motion data up

to 200 Hz while having the ability to provide up to 50 mA that can be used to power LEDs for

optogenetic experiments or stimulating electrodes. The battery of the wireless WEAR allows

recordings up to 4 h at 200 Hz sampling rate and even more at lower sampling rates. These

devices communicate with the computers through a base station based on the HARP design

developed by the Champalimaud Hardware Platform, which can be accessed through a software

GUI to easily change sensor parameters to best fit the experimental needs. The base stations have

several important hardware features such as 2 digital inputs and outputs, an analog input, 2

outputs for camera triggering, and a clock sync input and output that provides hardware-based

synchronization. The sensor can be started or stopped by software or pin. The WEAR motion

sensor family and base station are all open source (repository

at https://bitbucket.org/fchampalimaud/workspace/projects/HP). Moreover, the WEAR devices

are compatible with the Bonsai visual reactive programming software (https://bonsai-rx.org/),

also open source, and allow the integration and synchronization of the streams of data being

collected using the WEAR sensor with other data sources such as cameras.

Taking these specs and features together, the WEAR allows researchers to acquire high-

resolution motion data wirelessly and for long periods of time, without being computationally

very demanding. The 9-dimensional motion data acquired through WEAR is simple to process,

easy to connect to analysis software, which allowed the fast online behavior classification that

was fundamental for the experiments described in this paper.

Open field experiment. One-month post-surgery, mice were habituated to head-mounted

equipment over 2 days. On day 1, an actual or mock wireless inertial sensor (~2.5 cm H x 1 cm L

x 0.5 cm W with ~ 2.5-3.0 cm antennae, ~1.8 g weight) glued to the 4-position connector

(Harwin Inc., M52-040023V0445) was attached to the implanted receptacle connector on the

skull cap. Individual mice roamed freely in the home cage for 1 hour. On day 2, an actual

wireless inertial sensor and mono fiberoptic patchcord (200/220 μm diameter, 0.22 NA; Doric

DFP_200/220/900-0.22_2m_DF1.0-2FC) was attached to the skull cap via a mating sleeve.

Patchcords were attached to 1x2 fiber-optic rotary joint (intensity division, 0.22 NA; Doric,

FRJ_1x2i_FC-2FC) and mice roam freely in home cage for 1 hour. On open field recording day,

sensor/patchcord habituated mice were anesthetized by isoflurane, attached to equipment,

subjected to calibration protocol described below, and individually placed in an open field box

inside a sound insulated chamber. The open field box is made of 410 x 400 mm grey opaque

acrylic walls and a 410 x 400 mm white matte acrylic base. Individual mice were allowed to

behave freely inside the box for 75 minutes. The wireless inertial sensor (~1.8 g in weight,

WEAR wireless sensor v1.1; Champalimaud Scientific Hardware Platform) conveys motion

information sampled at 200 hz (set on WEAR v1.3.2 software; Champalimaud Scientific

Hardware Platform) to a receiver base-station (Harp basestation v1.1 or v. 1.2, Assembly v0,

Harp v1.4, Firmware v1.5; Champalimaud Scientific Hardware Platform), which conveys the

information to the experimental computer running a Bonsai script (Bonsai[38] editor v2.3.1) to

capture and record motion data and video information. Video was captured with a camera (Flea3

FL3-U3-I3Y3M(17450451), Point Grey Research) coupled to a 1/2" format lens (NMV-6WA,

Navitar).

Calibration. To ensure sensor stability within sessions, several approaches were employed. First, a coated mating sleeve was attached to the dual optic fiber cannula that sits immediately posterior to the sensor. The sleeve was thickened with black tape to a desired outer diameter such that it stabilized the sensor in the anterior-posterior direction. Second, the metal pins in the 4-position connector glued to the sensor were thickened with solder to stabilize their fit inside the receptacle connector in the skull cap. This protects against displacement in all directions. Third, stretchable black tape was wound around the base of the attached sensor and sleeve-covered cannula, further protecting against shifts in sensor positioning.

To control for possible variation in sensor positioning across sessions, a calibration approach was developed. Wireless inertial sensor was attached to individual isoflurane-anesthetized mice and the sensor was secured with the above strategies. Next, individual mice was placed in a custom-made calibration rig. The essential element of the rig is a vertical stainless-steel pole suspended above a stably secured table. In the setup used, the vertical pole was fixed to the horizontal edge of a vertically reversed "L" shape, stainless steel post assembly mounted on a breadboard (Thorlabs). The space between the lower end of the vertical pole and the table is enough for an individual mouse to slide underneath. The lower end of the vertical pole is fixed to a custom-made connector that resembles the connecting end of the fiberoptic patchcord. To perform calibration, individual isoflurane-anesthetized mice was securely attached to the vertical pole via a mating sleeve bridging the connection to the mouse's cannula implant. Next, replicate readings of the immobilized inertial sensor were made on Bonsai. Next, mice were attached to the experimental patchcord and allowed to recover in home cage for 20 minutes or until individual

mice are clearly recovered and behaviorally active. Individual mice were then placed in open-

field box for experimentation.

Calibration involves rotating all accelerometer and gyroscope readings from the inertial sensor

by a rotation matrix such that the final gravitational field vector of the stationary sensor, when

mounted on the mouse and fixed to the calibration rig, is in a universal frame of reference

whereby there is zero vertical tilt. In other words, the only non-zero acceleration is on the

universal z-axis (pointing down). To accomplish this, the accelerometer pitch and roll orientation

angles of the fixed stationary accelerometer were determined and then applied to calculate the

rotation matrix. The rotation matrix is multiplied by the sensor accelerometer and gyroscope

readings to remove the stationary vertical tilt from the sensor. To account for possible drift in

gyroscope baseline over time, a daily reading of stationary gyroscope baseline was made with a

mock cement skull cap attached to the sensor just before the start of each experimental day. The

baseline gyroscope readings were subtracted from all gyroscope values before the rotation matrix

is applied to sensor data.

Action Selection. After open field run in the grey-walled box, off-line behavioral clustering was

performed on calibrated sensor data. To identify the natural action repertoire of individual mice,

we quantified behavior using acceleration and gyroscope time series features in a similar fashion

as described previously[21]. For the ground truth analysis, we used: 1.) Gravitational acceleration

(GA) along the anterior-posterior (A-P) axis for the discrimination of postural changes - GAap.

2.) Raw sensor acceleration along the dorsal-ventral (D-V) axis to quantify movement

momentum – ACCdv. 3.) D-V axis of gyroscope to extract head head-body rotational

information – GYRdv. 4.) Total body acceleration to differentiate resting state from movement.

Total body acceleration (TotBA) was defined as:

$TotBA = sqrt(BAap^2 + BAml^2 + BAdv^2),$

where BAap, ml and dv represent the body acceleration of the anterior-posterior, medio-lateral

and dorsal-ventral axis, respectively. We calculated each individual BA component by median-

filtering the raw acceleration signals followed by a fourth-order Butterworth high-pass (0.5Hz)

filter. For the gravitational acceleration (GA) axis, the BA components were subtracted from the

median filtered raw signal axis.

All four time series features were binned into non overlapping 300 ms long window segments[26].

The values of each bin and per feature were then discretized, using fixed thresholds, producing a

summary distribution of each segment. For GAap and ACCdv we used 10 equal size threshold

values, plus two added bins between the limits and infinity to capture an approximated

distribution of values within each window bin. For GYRdv we used 5 thresholds (0, ±50, ±100)

to discriminate left and right turns. For TotBA, a single threshold was used to separate moving

from resting. The threshold was kept constant for all experiments and was set to the average

value separating the bimodal distribution of logTotBA (natural logarithm of TotBA feature). For

each 300-ms window segment we get four resulting histograms, one for each feature. The feature

histograms were individually normalized to obtain probability distributions and used to calculate

the pairwise similarities between segments.

We used the "earth mover's" (EM) distance as a measure of similarity[25]:

S = -(dEM/4)^2

where dEM is the sum of the normalized EM distances for the 4 features (GAap, ACCdv,

GYRdv and TotBA) defined above. The bin normalizations constrain S values within the range

[-1,0], specifically, -1 and 0 define the maximum dissimilarity and identity between the two

probability distributions, respectively. Finally, to produce a continuous unbiased classification of

behavioral states, the similarity measures were clustered using affinity propagation[20], with the

preference parameter set to the minimal value of the similarity matrix; this particular value was

used for its stable number of behavioral clusters within its range.

Using the behavioral clusters identified by affinity propagation clustering of the grey open field

behavior[13] as a ground truth for the true identity of each 300 ms histogram, we were able to

simulate and evaluate the precision with which the Earth Mover's Distance (EMD) metric[21,25]

could be applied for cluster matching online. Notable difference between the EMD metric used

here is the use of the 4 features mentioned above rather than the 3 features used previously[21], as

well as the multiplication of the similarity score by -1 such that the range of possible scores from

maximal identity to dissimilarity is 0 to 1, respectively. Although the EMD cluster matching

outcome correlates strongly with affinity propagation clustering, some false positive and false

negatives may occur. Several filters were set to optimize cluster selection for reinforcement: 1.)

We selected for clusters that show low false positive rate (<5.5%) and below the 60^{th} percentile

false positive rate amongst all clusters per animal. 2.) We selected against clusters with high

false negative rates (> 90^{th} percentile of clusters per animal). 3.) We selected against clusters that

tend to be performed serially within a short time interval. We calculated the probability that a

target cluster or its top 5 most similar clusters (determined by EMD score) would reappear 3-18

seconds after the first occurrence of the target cluster. Clusters that tend to be repeated either by

itself or have a high probability of having similar clusters appear within this 15 second window

(> 90^{th} percentile for median and range of probabilities of cluster appearing in window) were

removed from selection pool. 4.) We filtered against clusters whose matching by EMD would be

more sensitive to anterior-posterior shifts of the inertial sensor (although we already protected

against this possibility with the safeguards above) (> 90^{th} percentile for percent deviation from

original cluster matching after shifts of accelerometer reading in the anterior or posterior

direction). For each cluster, percent deviation is calculated first by summing up the total absolute

cluster matching changes from original cluster matching data in the anterior and posterior shifted

datasets. Next, the sum of deviation in the two altered datasets is divided by two and then

divided by the total of cluster calls from the original dataset, and multiplied by 100 to get percent

deviation from original cluster matching result. 5.) We selected for clusters that show fully

accelerating movement (cluster exemplar value of less than the maximum value of 1 in the body

acceleration feature bin of histogram). To choose dissimilar clusters per animal, an algorithm

was written filtering clusters of each animal's repertoire based on the feature histogram values of

each cluster's representative, or exemplar. Thresholds were set along the GAap and GYRdv

features to divide cluster exemplars based on the distribution of values within these feature

histograms. For each repertoire, all histogram values from all cluster exemplars are pooled to

create a pooled histogram. The range of bins with non-zero values for each feature are identified.

The algorithm then filters cluster exemplars in the repertoire for non-zero values in the high,

medium, low, or high+low value bins. For example, action A identification occurs by selecting

for a cluster exemplar with median counts falling in the high GAap and GYRdy value bins.

action B would then be selected by filtering for an exemplar with median counts falling in the

low GAap and GYRdy value bins. This results in actions that are highly dissimilar. For example,

EMD similarity scores comparing action A to action B almost always, except for 1 ChR2-YFP

animal, fall in the more dissimilar end of a distribution of scores created by comparing action A

to all actions in each animal. Hereafter, clusters will be referred to as actions.

Closed-Loop Optogenetics. For close loop optogenetics, a computer running a Bonsai script

captured and recorded wireless sensor motion data and video information as described above in

grey-walled open-field experiment. Here, data is also streamed to a custom MATLAB code

which analyzes action composition changes over the course of action reinforcement, we used the

EMD metric[21] to label individual 300 ms motion histograms with an action ID. For each arriving

300-ms segment we calculate the EMD distance between each cluster exemplar (or

representative) of the ground truth cluster library from the grey open field behavior recording.

The motion features histogram is assigned to the action for which comparison with the exemplar

gave the lowest EMD score (most similar to target) amongst all comparisons. Decision making

for stimulation has a range of 35-55 ms time gap between action performance and sent decision

for stimulation. To trigger optogenetics, a Multi-Pulse Width Modulation (PWM) generator

(Harp Multi-PWM Generator hardware v1.1, Assembly v1, Harp v1.4, Firmware v1.1; Harp

Multi-PWM Generator software v2.1.0; Champalimaud Scientific Platform) converts each

decision to trigger laser into electrical signals for 15 light pulses of 10 ms pulse duration at 25

Hz, with each train of pulses occurring over 600 ms and at 25% duty cycle. The multi-PWM

signal is passed through a 12 V, 7.2 W amplifier (Champalimaud Scientific Platform) and fixed

frequency driver (Opto-electronic, MODA110-D4-30 (2001.320220)) to control the activities of

a 473 nm, blue low noise laser (Shanghai Dream Lasers Technology, Co, Ltd. SDL-473-200T),

which was sent through an acousto-optic modulator (Opto-electronic, MTS110-A3-V1S (1001 /

330433)). The laser component that is modulated is then reflected by a mirror and funneled to a

mono fiberoptic patchcord, which is then coupled to a commutator. The output laser is then

passed through a dual-optic fiber patchord and connected to the implant cannula. Power

adjustment out of the tip of patchcord was made so that ~5mW was emitted from each end of the

dual optic fiber cannula. To ensure common time stamps from different channels, a clock

synchronization device (Harp Clock Sync v1.0; Champalimaud Scientific Platform) was

performed between the basestation and multi-PWM device.

Single action sequence selection. Mice were placed in a white open field box for closed loop

reinforcement protocol. Individual mice were subjected to a single session of protocol each day,

with sessions following each other on consecutive days. The white open field box is made of

410 x 400 mm white matte acrylic walls and a 410 x 400 mm white matte acrylic base. To

acquire baseline behavior, individual mice were allowed to behave freely inside the box for 30

minutes on the first action A selection session. Closed loop reinforcement by blue laser

stimulation of VTA DA neurons were made available for 60 minutes. 90 minutes of closed loop

reinforcement were made available for individual mice during sessions 2 and 3. For session 4, an

extinction protocol was carried out comprising of 20-minute maintenance of reinforced behavior

with laser availability, followed by 60 minutes of extinction of reinforced behavior without laser

availability, followed by 20-minute re-acquisition of reinforced behavior with laser availability.

To select for action B, a repeat of the protocol described above for action A was performed

starting on the day following extinction protocol of action A. Upon completion of the

reinforcement and extinction protocols for action B, a contingency degradation protocol was

performed comprising of 20-minute maintenance of action B with laser availability, followed by

60 minutes of contingency degradation of reinforced behavior by triggering laser randomly,

followed by 40-minute re-acquisition of reinforced behavior with laser availability for action B

performance.

Photometry experiment. One-month post-surgery, mice were habituated to head-mounted

equipment for 2 days. On day 1, habituation was made to wireless inertial sensor as described

above. On day 2, a multi-fiber bundled patch cord (3 fiber bundle, 400/440 μm diameter for a

maximum of inner diameter at 900 μm, 0.37 NA, 3.5 m long, 1.25 mm fiber tip diameter, low-

autofluorescence; Doric, BBP(3)_400/440/900-0.37_3.5m_FCM-3xMF1.25_LAF) was attached

to individual mice in addition to the wireless sensor and optogenetic patchcord. Individual mice

were allowed to habituate to the equipment for 1 hour in its home cage. On photometry recording

day, mice were subjected to 30 frames per second photometry recording (Neurophotometrics),

with 75-150 μW 560 nm LED illuminating rDA1m, and equivalent closed loop optogenetic

parameters described above were used. To test for DA release in the context of closed loop

optogenetic setup, an average of 30 hits of blue light were delivered randomly within the span of

30 minutes. To evaluate DA release in the context of food reward, mice were placed on food

deprivation protocol and kept within 85% of original weight. Mice were placed in an operant

chamber with a nosepoke linked to a lick detector (PyControl). Each lick detection triggers

dispensing 2 μl 10% sucrose. Since animals tend to accidentally trigger lick detector at the

beginning of sessions, between 40-50 sucrose dispensing events were gathered per animal and

rDA1m activities associated with the last 35 rewards of the session were used for analysis.

Two action sequence selection. Two action sequence selection occurs as follows: after

sensor/patchcord habituation and grey open field behavior recording, offline behavioral

clustering and action filtering were performed as for single action selection. For each animal,

median time intervals between all possible pairs of actions during open field were calculated as

described above. Across animals, T1/T2 pairs with median T1→T2 interval values varying

between 2 and 10 seconds, and with the feature of going from a head down(T1) to a head up(T2)

movement, were chosen for reinforcement.

On the first reinforcement session, a 30-minute baseline was taken when laser stimulation was

not available for reinforcement. Laser became available for reinforcement in all subsequent

sessions until extinction experiment. During reinforcement periods, when closed-loop system

detects performance of the proximal action (T1) of interest, the algorithm enters a state where

laser is triggered upon performance of the distal action (T2), regardless of the amount of time

that has elapsed between the latest T1 and T2. On Session 1, 60 minutes of laser availability was

given while in all subsequent reinforcement sessions, 90 minutes of laser availability was given.

Histology and Immunohistochemistry. After behavioral sessions were completed, mice were

deeply anesthetized with isoflurane and perfused transcardially in PBS and then 4% PFA/PBS.

Dissected brains with skulls attached were perfused in 4% PFA in PBS at 4 degrees Celsius

overnight. The next day, brains were rinsed 3 times in PBS. Next, brain regions including VTA

and implants were sectioned by vibratome into 50 or 100 μm slices. Slices are then subjected to

immunohistochemistry using the reagents below. Standard immunohistochemistry protocols

were applied to stain for the following reagents - Rabbit anti-GFP 488 conjugate (1:1000;

Molecular Probes A21311). Mouse Anti-TH (1:5000; Immunostar Th 22941) with Goat Anti-

Mouse - IgG (H+L) Highly cross-adsorbed secondary antibody - Alexa Fluor647 (1:1000;

ThermoFisher, A-21236), DAPI (1:1000 of 20 mg/mL stock; Sigma, D9542).

Imaging. Zeiss Axio Imager M2 microscope was used to acquire brain section pictures. 10x tiled

images were taken through the relevant fluorescent channels. The M2 is equipped with a fast

Colibri.7 LED illumination for excitation of fluorophores. Images are captured with a high-

sensitivity monochromatic sCMOS camera (Hamamatsu Orca Flash 4.0 v2). The objective used

for the images is a ZEISS Plan-ApoChromat 10x/0.45, which allows to resolve up to 577 nm

when using a wavelength of observation of 520nm and it is fully corrected for chromatic and

spherical aberrations. Implant locations were determined using standard mouse atlas[39].

Single action selection analyses. For target action frequency analysis, we analyzed frequencies

within 25-minute windows at 4 time points: Baseline (before first reinforcement trigger), Early

(after first reinforcement trigger in Session 1 (action A) or 5 (action B)), Mid (after 2-minute

mark in Session 2 (action A) or 6 (action B)), Late (after 2-minute mark in Session 3 (action A)

or 7 (action B)). For 3D action repertoire plots, baseline normalized frequencies were plotted and

actions whose time series include NaN or Infinity values were discarded from the plot. (Plotted

actions: 509 of 514 actions, 15 ChR2YFP animals (action A); 427 of 443 actions, 13 ChR2YFP

animals (action B); 355 of 356 actions, 10 YFP animals (action A); 341 of 356 actions, 10 YFP

animals (action B)).

Three parameters were assessed for rapid behavioral adaptation following cumulative closed

loop reinforcements: latency between Target A triggered reinforcements, Target A frequency and

average behavioral similarity to Target A. To calculate the latency parameter, the average

latency between 10 consecutive Target A triggered reinforcements following a specified number

of cumulated reinforcements were taken and then normalized by the average latency taken over

the final 10 baseline Target A instances that in simulations would have triggered reinforcement.

To calculate the frequency parameter, the frequency of Target A triggered reinforcements over

the course of 1 minute following a specified number of cumulated reinforcements were taken and

then normalized by frequency of the final 10 baseline Target A instances that in simulations

would have triggered reinforcement. To calculate the behavioral similarity parameter, the

average behavioral similarity (EMD score) to Target A between 10 consecutive Target A

triggered reinforcement events following a specified number of cumulated reinforcements were

taken and then normalized by the corresponding value taken over the final 10 baseline Target A

instances that in simulations would have triggered reinforcement.

rDA1m Fiber Photometry Analyses. To evaluate DA release in the context of food reward, the

delta F/Fo signal was plotted for rDA1m signal aligned to lick detection/reward trigger. The

baseline Fo value was taken as the median rDA1m raw fluorescence signal of the 10 time points

(333.33 milliseconds) preceding the trigger event. To test whether DA release is triggered in the

context of the closed loop system, the activity of the rDA1m sensor was quantified. Delta F/Fo

was calculated by subtracting baseline value from each fluorescent rDA1m value of a

smoothened time series (smooth function, default moving average filter, MATLAB), and then

dividing the outcome by the baseline value. To account for control ChR2-independent effects,

the average delta F/Fo trace of ChR2-YFP animals were subtracted from the corresponding

average trace of YFP animals, giving the differential delta F/Fo used for the plots. The standard

deviation of ChR2-YFP minus YFP curves were obtained by taking the square root of the sum of

squared variances of ChR2-YFP and YFP delta F/Fo curves.

Categorizing behavioral actions by temporal dynamics. To categorize behavioral actions by

temporal dynamics, moving mean of action counts was used as input. Various window sizes

were examined; 2.5-minute windows moving at 300 ms steps were found suitable for analyses.

The baseline frequency (f0) was the average of 5 minutes of moving mean data preceding the

first reinforcement event. Early frequency rate (f1) was the average of 30 minutes moving means

immediately following the first reinforcement event. Mid- and Late frequency rates were taken

from Day 2 (f2) and Day3 (f3), respectively. f2 and f3 rates were calculated from the beginning

30 minutes period after moving windows has accumulated enough bins (2.5 minutes) following

the start of the session. Significant positive modulation above baseline was judged if in 500

consecutive moving windows (2.5 minutes period) in Early/Mid or Late stages the frequency rate

of all bins were greater than the 99th percentile bin of baseline frequency. Significant negative

modulation below baseline was judged if in 500 consecutive moving windows (2.5-minute

period) in Early/Mid or Late stages the frequency rate of all bins were less than or equal to the 5th

percentile bin of baseline frequency. Actions that showed both significantly positive and

negative modulation at Early/Mid or Late stages when compared to baseline were delegated to

positive modulation group. For figure plotting, time-course median frequencies of action

dynamic types were downsampled 10-fold. To investigate the relationship between target

similarity and frequency, two approaches were taken. To perform multiple comparison statistics,

actions were binned by their percentile ranking in terms of similarity to target (EMD). This is

because action distribution based on raw EMD binning was not even. Percentile binning allowed

for even distribution of actions amongst the groups. To examine the distribution of action

dynamic type frequencies in terms of target similarity, a binning by raw EMD score (0.5 score

binwidth) was used because this allowed for clear visualization of the relationship between target

similarity and frequency. Alternatively, percentile binning of EMD score was also used and gave

similar trends.

Criterion for action dynamic types. Action dynamics were grouped according as follows: 1.)

Increasing actions showed significant increase in f0 to f1/2 and f1 to f2/3 comparisons and

showed either significant increase or unchanged frequency in f1/2 to f3 comparisons. 2.)

Sustained actions showed significant increase in f0 to f1/2 comparisons, and unchanged

frequency in f1 to f2/3 and f1/2 to f3 comparisons. 3.) Transient actions showed significant

increase in f0 to f1/2 comparisons, and significant decrease in f1/2 to f3 comparisons. 4.)

Decreasing actions showed significant decrease in f0 to f1/2 and f0 to f3 comparisons. 5.) Other

actions were all remaining actions that did not fall in the above groups. In the main figure only

dynamic subtypes with more than 10 members are shown.

Extinction analyses. 10 minutes portions from different time windows along the extinction

protocols (Session 4 for action A and Session 8 for action B) were chosen. Early maintenance

(M^1) starts from the first instance of target action performance in the session. Late maintenance

(M^2) is the portion preceding the first performance of target upon extinction. Early extinction

(E^1) begins at the first instance of target performance upon extinction. Late extinction (E^3) is the

portion preceding the first performance of target upon re-acquisition. Mid extinction (E^2) begins

at the midpoint between the starts of E^1 and E^3. Early re-acquisition (R^1) starts at the first

performance of target upon re-acquisition condition. Late re-acquisition (R^2) is the final portion

of the extinction protocol.

Action burstiness analysis. To evaluate action burstiness, or dispersion, we used Fano factor

(variance/mean) as a measure. A survey of moving mean frequencies of reinforced actions across

animals suggest that actions are more dispersed during the extinction phase, but the timescale

with which this may occur is variable. To identify a suitable timescale to detect dispersion across

reinforced actions, we screened a range of window sizes (600 ms to 5 minutes windows in 600

ms steps) with which to calculate moving window frequencies, and then calculate Fano factor in

varying time segments. We chose a moving window of 15 seconds (50 x 300 ms action units) to

construct moving mean frequencies. This window size consistently gave decreased Fano factor

in baseline vs. maintenance session across animal, a result that would be expected as

reinforcement led to stable target action performance.

Single action reinforcement, inter-target, and inter-action interval analyses. To quantity inter-target action intervals, the median amount of time that transpired between the start of successive target actions over the course of a time window was calculated. The time periods analyzed were: 1.) Baseline from the start of Day 1 (Sessions 1 and 5 for action A and B, respectively) until the first reinforcement event. 2-4.) Days 1 to 3 reinforcement. For reinforcement periods, behavior from the start of the first reinforcement event of that session until the end of session were analyzed. We considered the possibility that including the time interval between consecutive repeating of target actions (resulting in an inter-target action interval of 300 ms) would greatly affect the result. To test this, we removed values collected from consecutively repeating target actions. However, this did not affect result interpretations. Thus, we included intervals from consecutively repeating target actions in the presented analyses. For single action reinforcement, the median amount of time between the closest occurring action of interest and target action was calculated for both pre-target and post-target intervals.

Multinomial logistic regression predicting action dynamic types. To test whether intrinsic and baseline action properties are predictive of classifiable action dynamics during single action reinforcement from naïve state, two factors were considered. The factors are Earth Mover's Distance (EMD) similarity of action to target and median time interval of closest action of interest prior to target appearance at baseline condition.

To perform multinomial logistic regression, data from both dependent variables were log-transformed after addition of a constant value of 1. Transformed data were tested for collinearity

by examining scatter plots, Pearson's correlation coefficients, Variance Inflation Factors (VIF)

and condition indices. The two variables showed some correlation, but the coefficient value was

not above typical thresholds[40,41] and direct collinearity diagnostics did not show significant

collinearity (Pearson's correlation: $0.67 < 0.8$[40], VIFs: $1.82 < 5$-10[42], condition indices: $6.6 < 10$-

30[43]). Multinomial logistic regression was performed using MATLAB functions mnrfit and

mnrval. Non-Target A actions from all animals from reinforcement of action A were included

except those whose reinforcement dynamics were previously classified as "Other" types (n = 30

actions from a total of 514 actions, 15 ChR2-YFP animals). Decreasing dynamics type actions

were used as the reference group. Model accuracies were assessed using a 20-repeat, 10-fold

cross-validation approach for a total of 200 unique models for Real data, and 10,000 unique

models from 50 shuffled datasets.

To evaluate multinomial logistic regression, the deviance measure was used to judge model

fitting. Model performances were judged by area under precision-recall curve as this criterion is

suitable for imbalanced categories in the data[35]. A model containing both dependent variables

was found to outperform that of any single variable, even after consideration for penalties for an

extra factor (Akaike Information Criterion). The lack of significant collinearity between

dependent variables was supported by the stability of two relevant parameters, beta-coefficient

directions and significant p-values, across 200 cross-validation models and single- and double-

factor regression conditions (See Supplementary Information for tables).

Dopamine retrospective window analysis. To analyze whether DA reinforces actions proximal

to target, baseline rates of action transitions occurring close to reinforced action were examined.

First, a matrix tabulating 300 ms action counts from 2.4 seconds before to 2.4 seconds after each

theoretical target-triggered laser stimulation (600 ms in length) during baseline condition was

constructed. Next, all possible 600 ms action transitions (ex. X→Y) for each animal were then

counted using the above matrix, resulting in an action transition type (row) vs. time bin (column)

matrix where the counts of each action transition type occurring in specific 600 ms transition

windows (ex. X→Y) were recorded (sum across rows). This will be called the count matrix.

Next, the relative enrichment of each action transition type in a specific transition window

against all transition windows was calculated by dividing the action transition count matrix by

the total number of action transitions per type (probability across rows). Next, action transition

probability within a sliding 1.2 second transition window (containing a total of three action

transitions) relative to surrounding temporal environment (3.6 seconds) was derived by

subtracting the total number of action transitions per type within the surrounding 3.6 second

window from the total number of action transitions per type within the 1.2 second sliding

window of interest. This will be called the differential probability matrix. Next, action transition

types that showed greater than a threshold of 0.001 relative probability within sliding 1.2 second

windows of interest over the corresponding surrounding windows were filtered and kept for the

next step. Next, for each sliding 1.2 second window, the count matrix from above was analyzed

to select for action transition types that occurred between 2 to 6 times during the 30 minutes

baseline period (0.067 to 0.2 action transitions per minute). The count range was chosen to filter

out single events while selecting for action transitions with low initial frequencies over the

baseline period and analysis time range. Since the range of probabilities of specific action

transition types could vary greatly between different sliding 1.2 second windows, filtering as

above also balances the distribution of action transition probabilities amongst all action transition

types analyzed across sliding 1.2 second transition windows. The above process results in a list

of action transition types enriched for each sliding 1.2 second transition window, and baseline

normalized frequencies of these action transition types upon reinforcement in subsequent

sessions were calculated. Note that baseline normalized frequencies were calculated from all

occurrences of specific action transition types, regardless of their time distance in relationship to

target occurrence. Baseline normalized frequencies of individual action transition types were

averaged within animals and the means between animals are averaged to produce animal-

balanced results. Identical data trends and conclusions could be reached even if baseline

normalized frequencies of all action transitions were used for analyses.

Two action sequence experiment analyses. Two action sequence frequency was quantified in

terms of laser triggers per minutes. To assess learning across animals, the baseline frequency was

subtracted from frequencies of all reinforcement sessions. A criterion baseline subtracted

frequency of 3.2 triggers per minute was set after considering the range of baseline subtracted

frequencies observed in the open field and reinforcement sessions all animals. The criterion is set

such that it is > 20 % above the highest baseline-subtracted frequency value seen at open field

condition. The criterion point consistently falls above the open field frequencies of all animals

and marks the rising phase of all reinforcement frequency curves.

T1→T2 intervals were quantified as the time distance between the end of the latest distal action

(T1) and the end of the proximal action (T2) that triggers laser. T2→T1 intervals were quantified

as the time distance between the end of T2 that triggers laser and the end of the next closest T1.

To produce equivalent measures in open field and baseline conditions, laser trigger events were

simulated by scanning across the data as if reinforcement was available.

Significance testing was performed on 14 of 15 ChR2-YFP animals that reached criterion

frequency (ChR2-YFP Criterion). The lone animal that did not reach criterion frequency was

removed because the T1→T2 median interval was still very high after session 10. This animal

was subsequently subjected to single action reinforcement protocol to assess its ability to learn

T1 and subsequently T2. Next, the animal was again subjected to T1→T2 reinforcement

protocol. These results indicate that this animal was capable of action learning for both T1 and

T2 separately, and for T1→T2 sequence after learning of each individual action.

Reinforcement sessions for the 14 ChR2-YFP animals that reached beyond criterion frequency

continued until the T1→T2 interval has been decreased to below at least a median of 2 seconds.

As YFP animals do not decrease the T1→T2 median interval over sessions, we stopped

reinforcement at session 20.

Two action sequence extinction. Extinction session begins with a 25-minute maintenance

period for two action-sequence reinforcement, followed by a 40-minute extinction period when

laser was inactive, followed by a 25-minute re-acquisition period whereby reinforcement was

made available again. To quantify performance for plotting, frequency was calculated over 5

minutes bins and then normalized to the last 5 minutes bin of the maintenance condition. For

significant testing, raw frequencies were analyzed at the last 5 minutes of maintenance,

extinction, and re-acquisition conditions.

Two action sequence refinement. To measure refinement for T1 and T2 in the two-action

sequence, actions that were uniquely related to one but not the other were identified. Actions

performed by each animal in their open field repertoires were ranked by their EMD similarity

scores to T1 or T2. The top-12 actions (within action repertoires ranging between 30-40 actions)

most similar to either T1 or T2 were identified. Actions common to both T1 and T2 in these lists

were removed, leaving actions uniquely similar to T1 or T2. We required at least 3 non-target

actions to be uniquely related to each of T1 and T2. One of the animals did not meet this

requirement, because less than 3 actions were uniquely similar to each of T1 and T2 when

considering the top-12 actions related to T1 or T2. For this animal, we relaxed the stringency by

considering actions that uniquely belong as the top-9 actions most similar to either T1 or T2. We

took the median target-normalized frequency of these uniquely similar actions to T1 or T2 as the

refinement index. A refinement index of above or around 1 indicates little to no refinement of

uniquely related actions to target. Refinement index below 1 indicates refinement relative to

target; the lower the score the more refinement. Refinement curves were smoothened using the

Savitzky-Golay filter to improve visualization of trends. To better compare the progress of

refinement between T1- and T2-related actions, refinement indices were scaled such that the

minimum value amongst all sessions for individual animals would be zero and target-normalized

median frequency of 1 would remain at a scaled value of 1.

Relationship between T1→T2 interval and sessions to criterion frequency. To describe the

trend in a T1→T2 interval vs. sessions to criterion frequency scatter plot, non-linear sigmoidal fit

was tested against a 4th order polynomial fit. A linear fit was also tested. Sigmoidal fitting gave

the best result. The same fitting was tested for T2 → T1 interval vs. sessions to criterion

frequency, but the fit was poor and midpoint was unstable. For the T1→T2 sigmoidal curve,

half-maximum was 2.59 sessions to criterion frequency and midpoint was 4.69 seconds of open

field median interval. The half-maximum value was used to divide ChR2-YFP animals into slow

(above half-max) and fast (below half-max) learners.

Differential refinement analyses. The difference in area between T1 and T2 scaled refinement

curves over sessions was used to assess the relative refinement status between T1 and T2 over

sequence learning. The difference in areas were summed up using the trapezoid method across

sessions until the session when both T1 and T2 has or had reached minimal scaled refinement.

Next, the relationship between open field median interval and average difference in area under

T1 – T2 refinement curves per session was tested. Linear regression proved most suitable for

fitting (Goodness-of-fit: $R^2 = 0.66$). The fit for T1→T2 linear line was $y = 0.1893x – 0.7050$.

Slope was significantly non-zero ($p = 0.0004$). The same fitting was tested for T2 → T1 interval

vs. difference in area under T1 – T2 refinement curves per session ($y = 0.00736x + 0.1356$), but

the fit was poor, and goodness of fit was low (Goodness-of-fit: $R^2 = 0.07$). The slope was not

significantly non-zero ($p = 0.7063$).

Starting Point identification for evaluating progression of differential T1/T2 refinement. To

more precisely examine whether proximal action (T2) refinement precedes that of distal action

(T1) in Slow Learners, it was important to consider refinement progression of T1 relative to T2.

To rule out any bias towards proximal refinement because of initial bias towards proximal T2

refinement, a specific session was chosen as a Starting Point for analysis for each animal. This

Starting Point is defined by an early session in which T1 and T2 were relatively similar in

refinement levels or when the distal action T1 was more refined than proximal T2. To identify

these Starting Points, a scan was made retrospective from the session for which the T1→T2 time

interval is close to final value (less than or equal to a median of 3 seconds). Using this approach,

we identified earlier sessions in which distal T1 refinement was equal to or greater than proximal

T2 (T2 – T1 refinement curve area less than or equal to 0). The latest such session was set as the

Starting Point for analysis. If at no point early in learning did an animal have a session where

proximal (T1) action is most refined relative to distal (T2) action, an early session of closest T1

and T2 refinement was used as the Starting Point. The initial T2-T1 refinement curve area

difference calculated from the Starting Point to next session was subtracted from all T2-T1 area

differences calculated in subsequent sessions. This value is called the Starting Point subtracted

refinement difference. This made it possible to clearly track the change in relative refinement of

distal(T1) vs. proximal(T2) actions over time (Values above zero indicate T2>T1 refinement,

and values below zero indicate T1>T2 refinement). To identify the Turning Points for each

animal, sessions carrying the local maximum value of the Starting Point subtracted refinement

difference were identified for each animal. To calculate Starting Point subtracted refinement,

scaled refinement values from sessions of interest were subtracted from that of the Starting Point

session defined above.

Odds ratio analysis. For odds ratio calculation, the total amount of open field → Turning Point

session (second of two consecutive sessions used to calculate the refinement difference at

Turning Point as mentioned above) and Turning Point → session of criterion frequency median

interval changes were summed up for T1→T2 and T2→T1 intervals, respectively. Next, the

proportion of total interval change stemming from the open field condition→Turning Point

period, and from Turning Point→session reaching criterion frequency period, were calculated.

Next, the proportion of open field→Turning Point interval change was divided by the proportion

of Turning Point → session reaching criterion frequency period interval change for T1→T2 and

T2→T1 interval types, respectively. This gives the odds ratio.

T1 probability rank and refinement change across time bins from T2 trigger. For every

actual or simulated trigger for T1→T2 performance, the first occurrences of every action before

or after T2 triggers were counted at specific 300 ms time bins for up to 6 seconds before and

after T2 trigger. This was done for the specific conditions of baseline, Starting Point, Turning

Point, session passing criterion frequency, and last session. The probability of an action

occurring at a specific 300 ms time bin was calculated for all actions in the repertoire, and the

values were used to determine probability rank in terms of percentiles (100 percentile is most

probable action relative to all actions at a specific 300 ms time bin). To assess total T1

probability rank change within 0.3-1.8 or 2.1-3.6 second time bins, the area under the curve was

determined and values were normalized by subtraction from each animal's corresponding

baseline values. Refinement change was calculated by first taking the median probability rank of

actions most uniquely related to T1 at varying time distances before or after T2 trigger. This

value is then normalized by T1 probability rank to arrive at a refinement index. The area under

the curve was determined and values were normalized by subtraction from each animal's

corresponding baseline values. Decreasing values from Starting Point indicate increasing

refinement.

1005

Statistical Analysis:

Standard statistical analyses were performed on Prism (GraphPad Software, Inc.) and

permutation/bootstrap analyses were performed on MATLAB (MathWorks Inc.). To determine

appropriate tests for comparisons, datasets were assessed for normality using Anderson-Darling,

D'Agostino & Pearson, Shapiro-Wilk and/or Kolmogorov-Smirnov tests whenever applicable.

Datasets were also visualized for normality using QQ plots and assessed for equal variance by

examining the Residual plot (Residuals vs. Predicted Y). Parametric or non-parametric tests were

chosen based on the combination of these analyses. Data were transformed logarithmically (with

or without addition of a constant prior to transformation) whenever it was appropriate to promote

normality and equal variance. Unless specified, sphericity was not assumed, and Geisser-

Greenhouse correction was applied in all ANOVA tests. The appropriate post hoc multiple

comparisons tests were applied to compare between the means of specific conditions wherever

applicable. Significance was set at alpha $= 0.05$. For bootstrap analysis, significance was

determined by asking whether the original target action mean Fano factor was greater or less

than the 95% confidence interval of the bootstrap distribution. Permutation test was applied in

the comparisons between regression models because of the large sample size discrepancy

between groups. Bonferroni p adjustment was used to account for multiple comparisons in this

case. For detailed description of statistical procedures please refer to Supplementary Information.

1024

1025

1026

1027

1028

Acknowledgements:

We thank V.Athalye for helpful discussions and manuscript feedback, A. Vaz and C. Carvalho for mouse colony management, members of the Costa laboratory for comments, and the help from the Scientific Hardware Platform, Histopathology Platform, Scientific Software Platform and Advanced Bioimaging & BioOptics Experimental Platform (member of the Portuguese Platform of Bioimaging (PPBI-POCI-01-0145-FEDER-022122) of the Champalimaud Institute, S. Mutlu, D. Bento, P. Carriço., P. Silva, J. Araujo for hardware/software assistance, I. Marcelo for assistance with multinomial logistic regression, N. Loureiro for assistance with inertial sensor calculations , M. Mendoça and C. Alcacer for help with open field constructions. M. Carey lab for sharing apparatus. S. Fusi for project feedback. This work was supported by Life Sciences Research Fellowship and NINDS K99/R00 Award (1K99NS112575) granted to J.C.Y.T and National Institute of Health funding (5U19NS104649) to R.M.C.

Author Contributions:

J.C.Y.T and R.M.C. designed the study, interpreted results and wrote the paper. J.C.Y.T. performed and analyzed experiments. J.C.Y.T, V.P., F.C. designed close loop optogenetic system. J.C.Y.T., F.C. and A.S. executed assembly of the closed loop optogenetic system. F.C. and A.S. designed and assembled software and hardware. A.S. designed and assembled wireless inertial sensor and hardware. A.K. contributed Earth Mover's Distance code and was involved in early conceptions of the closed loop system. J.A.d.S., F.C. and A.S. designed and assembled the WEAR system. R.M.C. supervised the project. All authors edited the paper.

Competing Interests: F.C. is the Director of Open Ephys Production Site.

1052

Additional Information: Supplementary Information is available for this paper.

Code availability. MATLAB (MathWorks) codes used for data analysis are available from the corresponding author.

Data availability. Source Data are available from the corresponding author upon reasonable request.

Correspondence and requests for materials should be addressed to rc3031@columbia.edu

References

1. Schultz, W. Predictive reward signal of dopamine neurons. *J. Neurophysiol.* **80**, 1–27 (1998).

2. Schultz, W., Dayan, P. & Montague, P. R. A neural substrate of prediction and reward. *Science* **275**, 1593–1599 (1997).

3. Glimcher, P. W. Understanding dopamine and reinforcement learning: the dopamine reward prediction error hypothesis. *Proc. Natl. Acad. Sci. U. S. A.* **108 Suppl 3**, 15647–15654 (2011).

4. Thorndike, E. L. *Animal intelligence: Experimental studies.* viii, 297 (Macmillan Press, 1911). doi:10.5962/bhl.title.55072.

5. Skinner, B. F. *The behavior of organisms: an experimental analysis.* 457 (Appleton-Century, 1938).

6. Redgrave, P. & Gurney, K. The short-latency dopamine signal: a role in discovering novel actions? *Nat. Rev. Neurosci.* **7**, 967–975 (2006).

7.   Minsky, M. Steps toward Artificial Intelligence. *Proc. IRE* **49**, 8–30 (1961).

8.   Hull, C. L. *Principles of behavior: an introduction to behavior theory.* x, 422 (Appleton-

Century, 1943).

9.   Sutton, R. S. *Reinforcement learning an introduction /. Adaptive computation and machine*

learning (MIT Press, c1998.).

10.   Izhikevich, E. M. Solving the Distal Reward Problem through Linkage of STDP and

Dopamine Signaling. *Cereb. Cortex* **17**, 2443–2452 (2007).

11.   Reynolds, J. N. J., Hyland, B. I. & Wickens, J. R. A cellular mechanism of reward-related

learning. *Nature* **413**, 67–70 (2001).

12.   Shindou, T., Shindou, M., Watanabe, S. & Wickens, J. A silent eligibility trace enables

dopamine-dependent synaptic plasticity for reinforcement learning in the mouse striatum.

Eur. J. Neurosci. **49**, 726–736 (2019).

13.   Fisher, S. D. *et al.* Reinforcement determines the timing dependence of corticostriatal

synaptic plasticity in vivo. *Nat. Commun.* **8**, 334 (2017).

14.   Yagishita, S. *et al.* A critical time window for dopamine actions on the structural plasticity of

dendritic spines. *Science* **345**, 1616–1620 (2014).

15.   Jin, X., Tecuapetla, F. & Costa, R. M. Basal ganglia subcircuits distinctively encode the

parsing and concatenation of action sequences. *Nat. Neurosci.* **17**, 423–430 (2014).

16.   Cui, G. *et al.* Concurrent activation of striatal direct and indirect pathways during action

initiation. *Nature* **494**, 238–242 (2013).

17.   Jin, X. & Costa, R. M. Start/stop signals emerge in nigrostriatal circuits during sequence

learning. *Nature* **466**, 457–462 (2010).

18. Tervo, D. G. R. *et al.* Behavioral Variability through Stochastic Choice and Its Gating by

Anterior Cingulate Cortex. *Cell* **159**, 21–32 (2014).

19. Skinner, B. F. 'Superstition' in the pigeon. *J. Exp. Psychol.* **38**, 168–172 (1948).

20. Frey, B. J. & Dueck, D. Clustering by Passing Messages Between Data Points. *Science* **315**,

972–976 (2007).

21. Klaus, A. *et al.* The Spatiotemporal Organization of the Striatum Encodes Action Space.

Neuron **95**, 1171-1180.e7 (2017).

22. Boyden, E. S., Zhang, F., Bamberg, E., Nagel, G. & Deisseroth, K. Millisecond-timescale,

genetically targeted optical control of neural activity. *Nat. Neurosci.* **8**, 1263–1268 (2005).

23. Lammel, S. *et al.* Diversity of transgenic mouse models for selective targeting of midbrain

dopamine neurons. *Neuron* **85**, 429–438 (2015).

24. Dueck, D. Affinity Propagation: Clustering Data by Passing Messages. in (2009).

25. Rubner, Y., Tomasi, C. & Guibas, L. J. The Earth Mover's Distance as a Metric for Image

Retrieval. *Int. J. Comput. Vis.* **40**, 99–121 (2000).

26. Wiltschko, A. B. *et al.* Mapping Sub-Second Structure in Mouse Behavior. *Neuron* **88**,

1121–1135 (2015).

27. Phillips, A. G. & Fibiger, H. C. The role of dopamine in maintaining intracranial self-

stimulation in the ventral tegmentum, nucleus accumbens, and medial prefrontal cortex. *Can.*

J. Psychol. Can. Psychol. **32**, 58–66 (1978).

28. Corbett, D. & Wise, R. A. Intracranial self-stimulation in relation to the ascending

dopaminergic systems of the midbrain: a moveable electrode mapping study. *Brain Res.* **185**,

1–15 (1980).

29. Sun, F. *et al.* Next-generation GRAB sensors for monitoring dopaminergic activity in vivo.

Nat. Methods **17**, 1156–1166 (2020).

30. Beier, K. T. *et al.* Circuit Architecture of VTA Dopamine Neurons Revealed by Systematic

Input-Output Mapping. *Cell* **162**, 622–634 (2015).

31. Howe, M. W. & Dombeck, D. A. Rapid signalling in distinct dopaminergic axons during

locomotion and reward. *Nature* **535**, 505–510 (2016).

32. Schultz, W. Behavioral Theories and the Neurophysiology of Reward. *Annu. Rev. Psychol.*

57, 87–115 (2006).

33. Dickinson, A. The 28th Bartlett Memorial Lecture Causal learning: An associative analysis.

Q. J. Exp. Psychol. Sect. B **54**, 3–25 (2001).

34. Elsner, B. & Hommel, B. Contiguity and contingency in action-effect learning. *Psychol. Res.*

68, 138–154 (2004).

35. Saito, T. & Rehmsmeier, M. The precision-recall plot is more informative than the ROC plot

when evaluating binary classifiers on imbalanced datasets. *PloS One* **10**, e0118432 (2015).

36. Yagishita, S. *et al.* A critical time window for dopamine actions on the structural plasticity of

dendritic spines. *Science* **345**, 1616–1620 (2014).

37. Shindou, T., Shindou, M., Watanabe, S. & Wickens, J. A silent eligibility trace enables

dopamine-dependent synaptic plasticity for reinforcement learning in the mouse striatum.

Eur. J. Neurosci. **49**, 726–736 (2019).

38. Lopes, G. *et al.* Bonsai: an event-based framework for processing and controlling data

streams. *Front. Neuroinformatics* **9**, 7 (2015).

39. Paxinos, George & Franklin, K. B. J. The mouse brain in stereotaxic coordinates / George

Paxinos, Keith B.J. Franklin. (2001).

40. Berry, W. D., Feldman, S. & Stanley Feldman, D. *Multiple regression in practice*. (Sage,

1985).

41. Kim, J. H. Multicollinearity and misleading statistical results. *Korean J. Anesthesiol.* **72**,

558–569 (2019).

42. Neter, J., Kutner, M. H., Nachtsheim, C. J., Wasserman, W., & others. Applied linear

statistical models. (1996).

43. Belsley, D. A., Kuh, Edwin. & Welsch, R. E. *Regression diagnostics identifying influential*

data and sources of collinearity. (Wiley, 2004).

1150

Figures and Figure Legends:

Fig. 1. **Learning of a single action from the naïve state as mediated by closed loop**

optogenetics. a, Injection scheme. **b**, Wireless inertial sensor. **c**, Sensor data processing. **d**, Open

field behavioral clustering and action reinforcement. **e,** Closed loop schematic. **f,** Dopamine release

in dorsal and ventral striatum (n = 70 sucrose rewards, 2 ChR2-YFP mice; n = 66 and 65 random

stimulations, 2 ChR2-YFP and 2 YFP animals, respectively). Plots were mean, S.E.M. **g.** Action A

exemplar locations in behavioral space. **h-m,** ChR2-dependent reinforcement of Action A ($n = 15$

ChR2-YFP animals (green). $n = 10$ YFP animals (grey)). Plots were mean, S.E.M. **h,** Left: Head-

mount setup. Right: Light green/grey lines represent individual ChR2-YFP/YFP animals,

respectively. **i,** Rapid increase in target action performance in response to close-loop

reinforcements. Significant Time x Group Interactions (Supplementary Information). Plots were

mean, S.E.M. **j,** Evolution of pooled behavior repertoire (n = 509 actions, ChR2-YFP mice) across

learning. **k,** Early/Late cross-sectional views of (**j**) (Early: baseline normalized frequency >1, green

circles, < 1, magenta triangles). Blue dashed lines - single phase log decay fits. Bottom inset graph

shows Early/Late fitted lines normalized to 1 at EMD=0. **l,** Raw frequencies across learning and

target similarity percentile groups. Plots were mean, S.E.M. Two-way mixed effects statistics in

Supplementary Information. **m,** Pie chart summarizing distribution of actions according to their

dynamics within reinforced Action A (left) or other actions (right). Asterisks: **** $p < 0.0001$. ***

$p < 0.001$. ** $p < 0.01$. * $p < 0.05$. n.s. – not significant. See Supplementary Information for

statistical/sample details.

Figure 2. Transitioning from learned action to reinforcing new action. a-j, Animals reinforcing

for Action A (*n* = 15 ChR2-YFP) to Action B (*n* = 13 of 15 ChR2-YFP). *n* = 10 YFP animals. **a.**

Action A and B exemplar locations in behavioral space. **b,** Action similarity comparisons (A vs. B;

n = 15/10, ChR2-YFP/YFP; All vs. A; *n* = 514/356, ChR2-YFP/YFP) or Action B (All vs. B; *n* =

443/356, ChR2YFP/YFP). Plot indicates median/interquartile range. **c,** Reinforcement for Action A

and B in ChR2-YFP animals. Plot indicates mean/S.E.M. **d,** Evolution of pooled action repertoire (*n*

= 427 ChR2-YFP actions) reinforced for Action B. **e,** Early/Late cross-sectional views of (**d**). Blue

dashed lines indicate fitted decay curve. Bottom inset graph shows normalized Early/Late fitted

curves. **f,** Contingency degradation of Action B. Target random laser triggers frequencies (bottom) is based on initial Action B performance prior to contingency degradation. Plots indicate mean/S.E.M. **g-j,** Action A (blue) induced by reinforcement for Action B in experienced ChR2-YFP animals. **g-h,** Moving mean frequencies over reinforcement for Action A or B. Dashed, vertical line mark first reinforcement. Plots are mean/S.E.M (colored fill). Bin1/Bin2 are time bins for (**i-j**). **i-j,** Frequency measures within time bins noted in **(g,h).** Repeated measures two-way ANOVA reveal significant difference across time and actions A/B frequencies (not shown). Šidák's post hoc comparisons. Asterisks except in (**h**): **** $p < 0.0001$. ** $p < 0.01$. * $p < 0.05$. n.s. – not significant. See Supplementary Information for statistical/sample details.

Figure 3. Dopamine mediates retrospective reinforcement of freely moving behavior. a-b,

ChR2-dependent reinforcement decrease inter-action intervals for Action A (*n* = 15 ChR2-YFP) and

B (*n* = 13 of 15 ChR2-YFP). *n* = 10 YFP animals. Plots are mean/S.E.M (**a-b**). Significant

difference across time and ChR2-YFP/YFP (Mixed Effect Model. Action A: $F_{(3,69)}$ = 72.26, p <

0.0001. Action B: $F_{(3,62)}$ = 33.78, p < 0.0001.) **b**, Post-hoc Tukey's multiple comparisons of (**a**). **c-**

d, Distribution of action dynamic types (*n* = 464 actions, 15 ChR2-YFP animals) according to target

similarity (**c,d**), median time to target (**c,e**). **d-e**, Violin plots show median/quartiles. Two-tailed

permutation tests with Bonferroni-adjusted p-values. **f-g**, Multinomial logistic regression of all

factor combinations in Real data (200 models) versus Shuffled data (10,000 models). **f.** Groups

differ across combinations (repeated measures, two-way ANOVA. $F_{(2,30594)}$ = 518.2, p <

0.0001.). Post-hoc Dunnett multiple comparisons. Plots are mean/std. **g**, Performance of double-factor regression model measured with area under the precision-recall curves (AUPRC). Two-tailed permutation test with Bonferroni-adjusted p-value. Plots are mean/S.E.M. **h**, Identifying moving window-enriched action transitions. **i.** ChR2-dependent reinforcement for Action A increases action transitions prior to and within stimulation window. Plots indicate mean/S.E.M. **j,** Quantification of (**i**). Significant difference across time and Retrospective/Forward reinforcement directions (Mixed Effect Modeling. ChR2-YFP Session1: $F(6,168) = 114.8$, $p < 0.0001$. ChR2-YFP Session 3: $F(6,168) = 46.62$, $p < 0.0001$, YFP Session1: $F(6,108) = 10.52$, $p < 0.0001$. YFP Session 3: $F(6,168) = 0.8992$, $p = 0.4984$). Post-hoc Šidák multiple comparisons. **** $p < 0.0001$, *** $p < 0.001$, ** $p < 0.01$, * $p < 0.05$, n.s. – not significant. See Supplementary Information for statistical/sample details.

Figure 4. Relationship between pre-reinforcement inter-action intervals and learning of a two-action sequence. **a**, Schema. **b-l,** $n = 15$ (**b,d,h**) or 14 (**e-g,i-l**) ChR2-YFP, 6 YFP animals. Repeated measures one-way ANOVA, post hoc Šidák tests applied in (**e,g,i,k**). Plots of individuals in (**d-e**). **b**, ChR2-dependent increase in T1→T2 triggers (no laser during open field / baseline). **c**, Open field inter-action intervals of T1/T2 pairs chosen. Same color codes in (**d,h**). **d**, Individual learning curves labeled by color codes in (**c**). **e**, Frequency changes over conditions ($F_{(1.911,24.85)}=51.02$, $p<0.0001$). **f-g**, Extinction of T1→T2 sequence (ChR2-YFP). **f**, Plot shows mean(black)/S.E.M.(orange fill)/individuals(grey). **g,** Frequency changes over extinction conditions ($F_{(1.073, 12.87)} = 52.96$, $p<0.0001$). **h-i**, ChR2-dependent decrease in T1→T2 intervals. ($F_{(1.377, 17.90)} = 35.95$, $p<0.0001$) (**i**). **j**, T2:T1 frequency ratios (ChR2-YFP) **k**, Target refinement shown by median target normalized frequencies of related actions. (T1: $F_{(1.237, 16.08)} = 43.38$. T2:

F(1.171, 15.22) = 48.74. Both p<0.0001). Individual color code as in (c,g). **l**, Sigmoidal relationship between open field T1→T2 interval and sessions to criterion frequency.

Figure 5. Behavioral process underlying learning of a two action sequence. $n = 14$ ChR2-YFP (7 Slow Learners). **a**, T1/T2 refinements in two ChR2-YFP individuals. **b**, Linear relationship between initial T1→T2 interval and differential T1-T2 refinement. Non-zero slope significance: T1→T2, p = 0.0004, T2→T1, p = 0.7063. **c**, Progression of differential T1-T2 refinement from Starting Point in Individual Slow Learners. **d,** T1→T2 interval significantly decreased by Turning Point in Slow

Learners. Repeated-measures 2-way ANOVA. Post hoc Tukey's test. **e,** Odds ratio of T1→T2 / T2→T1 interval changes. Paired Wilcoxon test (p = 0.0312, n = 7 animals). **f,** Preferential refinement of T2 relative to T1 by Turning Point in Slow Learners. Raw scaled refinement indices. Repeated measures, mixed effects model. Significant main effects. Time (F $(2.184, 26.20)$ = 54.21, $p < 0.0001$). Post-hoc Šidák test. **g,** First occurrences of T1 before (left) and after (right) T2 triggers across learning stages. **h,** Quantification of pooled time bins from (**g**). Repeated measures, 2-way ANOVA for learning stage vs. rank change. First T1 Before and After T2 Trigger groups differ across learning stage and total T1 rank change. (Proximal bins $(0.3\text{-}1.8$ sec): $F(3,36)$ = 3.126. p=0.0376. Distal bins $(2.1$ to 3.6 sec): $F(3,36)$ = 7.701. p<0.001). Post-hoc Šidák relative to Starting Point values. **g,** Model for learning initially distantly separated T1→T2 sequences. Time not drawn to scale. **** $p < 0.0001$. *** $p < 0.001$. ** $p < 0.01$. * $p < 0.05$. n.s. – not significant. All bar plots indicate mean +/- S.E.M. See Supplementary Information for statistical/sample details.