A formula for calculating 30-item Geriatric Depression Scale (GDS-30) scores from the 15-item version (GDS-15)

Yuge Zhang[1], Marco Hoozemans[1], Mirjam Pijnappels[1], Sjoerd M. Bruijn[1,2,3*]

[1]Department of Human Movement Sciences, Faculty of Behavioural and Movement Sciences, Vrije Universiteit Amsterdam, Amsterdam, The Netherlands

[2]Institute of Brain and Behavior Amsterdam, Amsterdam, The Netherlands

[3]Biomechanics Laboratory, Fujian Medical University, Quanzhou, Fujian, PR China

* **Correspondence to:**
Sjoerd M. Bruijn
Van der Boechorststraat 9, 1081 BT Amsterdam, The Netherlands
Email address: s.m.bruijn@vu.nl

Abstract

The Geriatric Depression Scale with 30 items (GDS-30) and with 15 items (GDS-15) are both valid tools for assessing depression in older adults, but their absolute values are not directly comparable. Here, we used a dataset (n=431) with GDS-30 scores from a project concerning fall-risk assessment in older adults (FARAO) to develop and validate a formula which can be used to convert GDS-15 scores into GDS-30 scores. We found that the GDS-15 score cannot simply be multiplied by 2 to obtain the GDS-30 scores and that estimations of GDS-30 from GDS-15 are not affected by age, sex and MMSE. Therefore, the optimal formula to estimate the GDS-30 score from the GDS-15 score was: GDS-30_estimated = 1.57 + 1.95 × GDS-15. This formula yielded an estimate of GDS-30 with an explained variance of 79%, compared to 63% when GDS-15 was simply multiplied by 2. Researchers that have used the GDS-15 and want to compare their outcomes to other studies that reported only the GDS-30 are advised to use this formula.

Keywords: depression; GDS; older adults

1. Introduction

Our society is ageing, and it is well known that ageing comes with physical and mental problems. For instance, the prevalence of depression in older adults is around 28.4% worldwide (Hu and others 2022), with ranges of 10% to 30% in hospitals and nursing homes (Mitchell and others 2010b). The Geriatric Depression Scale with 30 items (GDS-30), designed by Yesavage and others (1983), has been shown to be a valid and reliable tool for assessing both major and minor depression in nursing home patients (Jongenelis and others

2005), primary care (Mitchell and others 2010a) and medical settings (Mitchell and others (2010b). The GDS-30 is a questionnaire, consisting of 30 yes/no items, which asks about depressive feelings on the day of administration. For most of these items, a "yes" answer indicates higher depressive symptoms. A sum-score of 10 is the cut-off between normal and depression and a sum-score of 20 is the dividing line between mild and severe depression. In 1986, a shortened version of the GDS-30 was proposed by Sheikh and Yesavage (1986) with 15 selected items (from the GDS-30) that had the highest correlation with depressive symptoms. The GDS-15 was also found to have good sensitivity and specificity, among older adult psychiatric inpatients (Mitchell and others 2010a; Mitchell and others 2010b), so it is a good substitute for the GDS-30 (Lesher and Berryhill 1994).

In a recent study, we came across a situation where we had administered the GDS-15, whereas the literature we were comparing our results to mostly reported results in terms of the GDS-30. The simplest solution to being able to compare our GDS-15 results to those who used the GDS-30 would have been multiplying the GDS-15 sum scores by 2. However, it was unclear if these scales are directly comparable as, to the best of our knowledge, there is no literature which provides a valid formula to 'convert' GDS-15 scores to equivalent GDS-30 scores that goes beyond the 'multiply by 2' method. Considering all items in GDS-15 come from the more extensive GDS-30, GDS-15 scores can be obtained from completely administered GDS-30 scores. Thus, here, we used a previously collected GDS-30 dataset from a community living sample to compare the validity of multiplying the GDS-15 sum scores by 2 to get GDS-30 sum scores with the validity of a linear conversion formula from GDS-15 to GDS-30 scores.

2. Methods

2.1 Data collection

The dataset we used was from a project concerning fall-risk assessment in older adults (FARAO) performed at the Vrije Universiteit, Amsterdam (van Schooten and others 2015). Participants were independently living individuals living in Amsterdam and surroundings and were included if their ages were between 65 and 99 years, and their mini mental state exam score (MMSE) was between 19 and 30. All items of the GDS-15 were contained in GDS-30, which participants filled in. The medical ethical committee of the Vrije Universiteit medical center approved the protocol (#2010/290) and all participants provided informed consent. From the individual scores on the GDS-30 items, the GDS-30 and GDS-15 scores were calculated by summing the scores of the items (after reversing the scores of the items for which this was required) according to Yesavage and others (1983) and Kurlowicz and Greenberg (2007), respectively.

2.2 Statistical analysis

Statistical analysis was performed in MATLAB R2021a (MathWorks, U.S.). We used linear regression to determine the relationship between GDS-30 sum score (outcome variable) and

GDS-15 sum score (predictor variable). To test whether the obtained model was better than
simply multiplying the GDS-15 scores by 2, we checked the goodness of fit, which was
evaluated by the coefficient of determination (R^2) and measured the error of the model in
predicting the GDS-30 by root mean square error (RMSE, with lower RMSE values indicating
smaller errors)(Hyndman and Koehler 2006). Moreover, we evaluated whether the intercept
of the obtained model was significantly different from 0 (i.e., 95% confidence interval (CI) did
not include zero), and whether the regression coefficient of the GDS-15 score as predictor
variable was significantly different from 2 (i.e., the 95% confidence interval did not include 2).
Age, sex and MMSE score were explored for interaction, as it might be expected that these
variables affect the relationship between GDS-15 scores and GDS-30 scores, i.e. that this
relationship is different for different levels of the potential effect modifiers.

3. Results

3.1 Demographic characteristics

From the FARAO cohort, 431 participants satisfied the inclusion criteria and filled in all items of
GDS-30 questionnaires (see table 1). The mean age was 76.4 (SD 7.5) years, the mean GDS-30
sum score was 5.4 (4.7), and the mean GDS-15 sum score was 2.0 (2.1). This latter result already
suggests that GDS-15 scores cannot simply be multiplied by 2 to get GDS-30 scores.

89
Table 1. Demographic Characteristics

Characteristics	Mean (SD)
Age (years)	76.4(7.5)
Sex (%, male)	41.5%
MMSE	27.4(2.4)
GDS-30 scores	5.4 (4.7)
GDS-15 scores	2.0 (2.1)

3.2 Prediction of GDS-30 scores from GDS-15 scores

Age, sex and MMSE score showed no significant interaction with GDS-15 scores in the prediction
of GDS-30 scores (see supplementary table 1). Therefore, only GDS-15 was considered as the
single predictor variable of GDS-30 in the regression model. The linear regression model to
estimate GDS-30 from GDS-15 was GDS-30_estimated = 1.57 + 1.95 × GDS-15. The R^2 of the linear
model predicting GDS-30 from GDS-15 was 0.79, and RMSE was 2.13 (see figure 1; marker size
represents the number of subjects for that data point). When simply multiplying GDS-15 scores
by 2, the R^2 was 0.63, and RMSE was 2.59, indicating that our linear model was better than
simply multiplying GDS-15 scores by 2. The intercept was significantly different from zero, (95% CI
[1.29,1.84]), but the regression coefficient of GDS-15 was not significantly different from 2 (95%
CI [1.86,2.05]).

104
105 **Figure 1. Linear regression relation between GDS-15 & GDS-30 score;** marker size represents the
106 number of subjects for that data point

4. Discussion

We set out to find a formula to estimate equivalent GDS-30 scores from GDS-15. In our
cohort of over 400 older adults, we found the formula GDS-30=1.57 + 1.95 × GDS-15, which
yielded better performance than simply multiplying GDS-15 scores by 2, as shown by the
higher explained variance (R^2) and lower error (RMSE). Although the slope was not
significantly different from 2, the intercept was statistically higher than zero. This indicates
that GDS-30 scores calculated using the GDS-15 times 2 method systematically
underestimate the better fitting linear model by 1.6 points, which is also clearly visible in
figure 1. Our analyses indicated that GDS-30 scores calculated from GDS-15 scores by using
the suggested formula have a better goodness of fit and lower error than simply multiplying
a GDS-15 score by two. In addition, the formula appeared to be robust as the relationship
was not affected by age, sex and MMSE score. The formula can be useful when comparing
the geriatric depression scale among different studies.
Our study has some limitations. First, as the data were collected in a community-dwelling
sample, the validity of our formula in a more severely depressed sample remains to be
tested. Second, although using non-integer estimates of the GDS-30 to compare with other
studies is not a problem, for clinical interpretation, integer values are probably preferred.
Lastly, our linear regression leads to a subject with score 0 on the GDS-15 having a score of
1.57 for the estimated GDS-30. While this may seem odd, it should be kept in mind that this
is still well below the cut-off for depression (which is 10).
127

5. Conclusion

In conclusion, the linear regression formula developed in our study can be used effectively

for estimating GDS-30 scores from GDS-15 assessments.

Acknowledgements: Many thanks to Kimberley van Schooten of Neuroscience Research
Australia, University of New South Wales, and Roel Weijer of the Department of Neurology,
Leiden University Medical Center, who collected the data. YZ was funded by the China
Scholarship Council (CSC) (202009110145). MP and SMB were funded by a grant from the
Dutch Organization for Scientific Research (NWO) (no. 91714344 and 016.Vidi.178.014,
respectively).

Author contribution: YZ contributed to formal analysis, software, funding acquisition, writing
the original draft, reviewing and editing of the draft. MH contributed to formal analysis,
methodology, reviewing and editing of the draft. MP contributed to investigation, data
curation, funding acquisition, resources, and supervision. SMB contributed to
conceptualization, formal analysis, funding acquisition, methodology, supervision, reviewing
and editing of the draft. All authors gave final approval of the version to be submitted.

Conflicts of Interest: The authors declare no conflict of interest.

Correspondence and requests for materials should be addressed to SMB.
145
146

147   **Reference**

148   Hu, T.; Zhao, X.; Wu, M.; Li, Z.; Luo, L.; Yang, C.; Yang, F. Prevalence of depression in older adults: A
149   systematic review and meta-analysis. Psychiatry Research. 311:114511; 2022
150   Hyndman, R.J.; Koehler, A.B. Another look at measures of forecast accuracy. International Journal of
151   Forecasting. 22:679-688; 2006
152   Jongenelis, K.; Eisses, A.M.H.; Gerritsen, D.L.; Derksen, M.; Beekman, A.T.F.; Kluiter, H.; Ribbe, M.W.
153   Diagnostic accuracy of the original 30-item and shortened versions of the Geriatric
154   Depression Scale in nursing home patients. International Journal of Geriatric Psychiatry.
155   20:1067-1074; 2005
156   Kurlowicz, L.; Greenberg, S.A. The Geriatric Depression Scale (GDS). AJN The American Journal of
157   Nursing. 107:67-68; 2007
158   Lesher, E.L.; Berryhill, J.S. VALIDATION OF THE GERIATRIC DEPRESSION SCALE - SHORT-FORM AMONG
159   INPATIENTS. Journal of Clinical Psychology. 50:256-260; 1994
160   Mitchell, A.J.; Bird, V.; Rizzo, M.; Meader, N. Diagnostic validity and added value of the geriatric
161   depression scale for depression in primary care: A meta-analysis of GDS(30) and GDS(15).
162   Journal of Affective Disorders. 125:10-17; 2010a
163   Mitchell, A.J.; Bird, V.; Rizzo, M.; Meader, N. Which Version of the Geriatric Depression Scale is Most
164   Useful in Medical Settings and Nursing Homes? Diagnostic Validity Meta-Analysis. American
165   Journal of Geriatric Psychiatry. 18:1066-1077; 2010b
166   van Schooten, K.S.; Pijnappels, M.; Rispens, S.M.; Elders, P.J.M.; Lips, P.; van Dieën, J.H. Ambulatory
167   fall-risk assessment: Amount and quality of daily-life gait predict falls in older adults. The
168   Journals of Gerontology: Series A. 70:608-615; 2015
169   Yesavage, J.A.; Brink, T.L.; Rose, T.L.; Lum, O.; Huang, V.; Adey, M.; Leirer, V.O. Development and
170   validation of a geriatric depression screening scale: A preliminary report. Journal of
171   Psychiatric Research. 17:37-49; 1983

172   Yesavage, J.A.; Sheikh, J.I. 9/Geriatric depression scale (GDS) recent evidence and development of a
173   shorter version. Clinical gerontologist. 5:165-173; 1986
174