A formula for calculating 30-item Geriatric Depression Scale (GDS-30) scores from the 15-item version (GDS-15)

Yuge Zhang[1], Marco Hoozemans[1], Mirjam Pijnappels[1], Sjoerd M. Bruijn[1,2,3*]

[1]Department of Human Movement Sciences, Faculty of Behavioural and Movement Sciences, Vrije Universiteit Amsterdam, Amsterdam, The Netherlands

[2]Institute of Brain and Behavior Amsterdam, Amsterdam, The Netherlands

[3]Biomechanics Laboratory, Fujian Medical University, Quanzhou, Fujian, PR China

* Corresponding:
Sjoerd M. Bruijn
Van der Boechorststraat 9, 1081 BT Amsterdam, The Netherlands
Email address: s.m.bruijn@vu.nl

Abstract

The Geriatric Depression Scale with 30 items (GDS-30) and with 15 items (GDS-15) are both valid tools for assessing depression in older adults, but their absolute values are not directly comparable. Here, we used a dataset (n=431) with GDS-30 scores from a project concerning fall-risk assessment in older adults (FARAO) to develop and validate a formula which can be used to convert GDS-15 scores into GDS-30 scores. We found that the GDS-15 score cannot simply be multiplied by 2 to obtain the GDS-30 scores. Instead, the optimal formula to estimate the GDS-30 score from the GDS-15 score was: GDS-30_estimated = 1.57 + 1.95 × GDS-15. This formula yielded an estimate of GDS-30 with an explained variance of 79%, compared to 63% when GDS-15 was simply multiplied by 2. Researchers that have used the GDS-15 and want to compare their outcomes to other studies that reported only the GDS-30 are advised to use this formula.

Keywords: depression; GDS; older adults

1. Introduction

Our society is ageing, and it is well known that ageing comes with physical and mental problems. For instance, the prevalence of depression in older adults is around 28.4% worldwide (Hu and others 2022), with ranges of 10% to 30% in hospitals and nursing homes (Mitchell and others 2010b). The Geriatric Depression Scale with 30 items (GDS-30), designed by Yesavage and others (1983), has been shown to be a valid and reliable tool for assessing both major and minor depression in nursing home patients (Jongenelis and others 2005), primary care (Mitchell and others 2010a) and medical settings (Mitchell and others (2010b).

In 1986, a shortened version of the GDS-30 was proposed by Sheikh and Yesavage (1986) with 15 selected items (from the GDS-30) that had the highest correlation with depressive symptoms. The GDS-15 was also found to be diagnostically valid and reliable, so it is an adequate substitute for the GDS-30 (Lesher and Berryhill 1994; Mitchell and others 2010b).

In a recent study, we came across a situation where we had administered the GDS-15, whereas the literature we were comparing our results to mostly reported results in terms of the GDS-30. The simplest solution to be able to compare our GDS-15 results to those who used the GDS-30 would have been multiplying the GDS-15 sum scores by 2. However, it was unclear if these scales are directly comparable as, to the best of our knowledge, there is no literature which provides a valid formula to 'convert' GDS-15 scores to equivalent GDS-30 scores that goes beyond the 'multiply by 2' method. Considering all items in GDS-15 come from the more extensive GDS-30, GDS-15 scores can be obtained from completely administered GDS-30 scores. Thus, here, we used a previously collected GDS-30 dataset to compare the validity of multiplying the GDS-15 sum scores by 2 to get GDS-30 sum scores with the validity of a linear conversion formula from GDS-15 to GDS-30 scores.

2. Methods

2.1 Data collection

The dataset we used was from a project concerning fall-risk assessment in older adults (FARAO) performed at the Vrije Universiteit, Amsterdam (van Schooten and others 2015). Participants were recruited in Amsterdam and surroundings and were included if their ages were between 65 and 99 years, and their mini mental state exam score (MMSE) was between 19 and 30. Participants filled in all items of the GDS-30, which includes all items of the GDS-15. The medical ethical committee of the Vrije Universiteit medical center approved the protocol (#2010/290) and all participants provided informed consent. From the individual scores on GDS-30 items, the GDS-30 and GDS-15 scores were calculated according to Yesavage and others (1983) and Kurlowicz and Greenberg (2007), respectively.

2.2 Statistical analysis

Statistical analysis was performed in MATLAB R2021a (MathWorks, U.S.). We used a linear regression to determine the relationship between GDS-30 sum score (outcome variable) and GDS-15 sum score (predictor variable). To test whether the obtained model was better than simply multiplying the GDS-15 scores by 2, we checked the goodness of fit, which was evaluated by the coefficient of determination (R^2). Moreover, we evaluated whether the intercept of the obtained model was significantly different from 0 (i.e., 95% confidence interval (CI) did not include zero), and whether the regression coefficient of the GDS-15 score as predictor variable was significantly different from 2 (i.e., the 95% confidence interval did not include 2).

3. Results

3.1 Demographic characteristics

From the FARAO cohort, 431 participants satisfied the inclusion criteria and filled in all items of GDS-30 questionnaires (see table 1). The mean age was 76.4 (SD 7.5) years, the mean GDS-30 sum score was 5.4 (4.7), and the mean GDS-15 sum score was 2.0 (2.1). This latter result already indicates that GDS-15 scores cannot simply be multiplied by 2 to get GDS-30 scores.

Table 1. Demographic Characteristics

Characteristics	Mean (SD)
Age (years)	76.4(7.5)
Gender (%, male)	41.5%
MMSE	27.4(2.4)
GDS-30 scores	5.4 (4.7)
GDS-15 scores	2.0 (2.1)

3.2 Prediction of GDS-30 scores from GDS-15 scores

The R^2 of a linear model predicting GDS-30 from GDS-15 was 0.79, and it's RMSE was 2.13 (see figure 1). When simply multiplying GDS-15 scores by 2, the R^2 was 0.63, and RMSE was 2.59, indicating that our linear model was better than simply multiplying GDS-15 scores by 2. The linear regression model to estimate GDS-30 from GDS-15 was GDS-30_estimated = 1.57 + 1.95 × GDS-15. The intercept was significantly different from zero, (95% CI [1.29,1.84]), but the regression coefficient of GDS-15 was not significantly different from 2 (95% CI [1.86,2.05]).

Figure 1. Linear regression relation between GDS-15 & GDS-30 score

4. Discussion

We set out to find a formula to estimate equivalent GDS-30 scores from GDS-15. In our cohort of over 400 older adults, we found the formula GDS-30=1.57 + 1.95 × GDS-15, which yielded better performance than simply multiplying GDS-15 scores by 2, as shown by the higher explained variance (R^2) and lower error (RMSE). Although the slope was not significantly different from 2, the intercept was statistically higher than zero. This indicates that GDS-30 scores calculated using the GDS-15 times 2 method systematically underestimate the better fitting linear model by 1.6 points, which is also clearly visible in figure 1. In conclusion, our analyses suggest that 1) GDS-30 scores simply calculated by multiplying a GDS-15 score by 2 results in lower goodness of fit than using the suggested formula, 2) GDS-30 scores can be validly estimated from GDS-15 scores using the suggested formula. This formula can be useful when comparing the geriatric depression scale among different studies.

5. Conclusion

In conclusion, the linear regression formula developed in our study can be used validly for estimating GDS-30 scores from GDS-15 assessments.

Acknowledgements: Many thanks to Kimberley van Schooten of Neuroscience Research Australia, University of New South Wales, and Roel Weijer of the Department of Neurology, Leiden University Medical Center, who collected the data. YZ was funded by the China Scholarship Council (CSC) (202009110145). MP and SMB were funded by a grant from the Dutch Organization for Scientific Research (NWO) (no. 91714344 and 016.Vidi.178.014, respectively).

Author contribution: YZ contributed to formal analysis, software, funding acquisition, writing the original draft, reviewing and editing of the draft. MH contributed to formal analysis, methodology, reviewing and editing of the draft. MP contributed to investigation, data curation, funding acquisition, resources, and supervision. SMB contributed to conceptualization, formal analysis, funding acquisition, methodology, supervision, reviewing and editing of the draft. All authors gave final approval of the version to be submitted.

Conflicts of Interest: The authors declare no conflict of interest.

Correspondence and requests for materials should be addressed to SMB.

Reference

Hu, T.; Zhao, X.; Wu, M.; Li, Z.; Luo, L.; Yang, C.; Yang, F. Prevalence of depression in older adults: A systematic review and meta-analysis. Psychiatry Research. 311:114511; 2022

Jongenelis, K.; Eisses, A.M.H.; Gerritsen, D.L.; Derksen, M.; Beekman, A.T.F.; Kluiter, H.; Ribbe, M.W. Diagnostic accuracy of the original 30-item and shortened versions of the Geriatric Depression Scale in nursing home patients. International Journal of Geriatric Psychiatry. 20:1067-1074;

129  2005

130 Kurlowicz, L.; Greenberg, S.A. The Geriatric Depression Scale (GDS). AJN The American Journal of
131   Nursing. 107:67-68; 2007

132 Lesher, E.L.; Berryhill, J.S. VALIDATION OF THE GERIATRIC DEPRESSION SCALE - SHORT-FORM AMONG
133   INPATIENTS. Journal of Clinical Psychology. 50:256-260; 1994

134 Mitchell, A.J.; Bird, V.; Rizzo, M.; Meader, N. Diagnostic validity and added value of the geriatric
135   depression scale for depression in primary care: A meta-analysis of GDS(30) and GDS(15).
136   Journal of Affective Disorders. 125:10-17; 2010a

137 Mitchell, A.J.; Bird, V.; Rizzo, M.; Meader, N. Which Version of the Geriatric Depression Scale is Most
138   Useful in Medical Settings and Nursing Homes? Diagnostic Validity Meta-Analysis. American
139   Journal of Geriatric Psychiatry. 18:1066-1077; 2010b

140 van Schooten, K.S.; Pijnappels, M.; Rispens, S.M.; Elders, P.J.M.; Lips, P.; van Dieën, J.H. Ambulatory fall-
141   risk assessment: Amount and quality of daily-life gait predict falls in older adults. The Journals
142   of Gerontology: Series A. 70:608-615; 2015

143 Yesavage, J.A.; Brink, T.L.; Rose, T.L.; Lum, O.; Huang, V.; Adey, M.; Leirer, V.O. Development and
144   validation of a geriatric depression screening scale: A preliminary report. Journal of Psychiatric
145   Research. 17:37-49; 1983

146 Yesavage, J.A.; Sheikh, J.I. 9/Geriatric depression scale (GDS) recent evidence and development of a
147   shorter version. Clinical gerontologist. 5:165-173; 1986

148