1   No evidence of sexual conflict for a novel sperm-derived seminal fluid protein in *Caenorhabditis*

2   nematodes

4   Short Title: NSPF proteins promote fitness without sexual conflict

7   Katja R. Kasimatis[1]*, Christine Rehaluk[1], Locke Rowe[1], and Asher D. Cutter[1]*

9   [1]Department of Ecology and Evolutionary Biology, University of Toronto, Toronto, Ontario,

10   Canada

12   *Corresponding authors: k.kasimatis@utoronto.ca (KRK) and asher.cutter@utoronto.ca (ADC)

ABSTRACT

Post-insemination molecular interactions involving sperm are an ideal system for linking genotype, phenotype, and fitness. Nematode sperm contain subcellular vesicles known as membranous organelles (MOs) that fuse with the cell membrane upon sperm activation to release their soluble contents into the extracellular space. The second most abundant proteins in the MOs is the conserved Nematode-Specific Peptide family, group F (NSPF) gene family. We hypothesize that these proteins contribute to seminal fluid upon MO fusion and are part of post-insemination reproductive tract dynamics. We used experimental evolution to compete the wildtype allele against a deletion allele in 10 replicate populations. We calculated a significant mean selective disadvantage of 0.1% for the deletion allele, which indicated that the NSPF genes are an evolutionarily important component of male reproductive success. This conclusion was reinforced by qualitative trends from lower powered single-generation male fertility assays. We then characterized the functional location of NSPF proteins during fertilization using whole-worm immunostaining of a His-tagged *nspf-1* transgene. Throughout male development, we found that NSPF presence and abundance was correlated with reproductive maturity in males, localizing to the seminal vesicle during L4 and to the spicules throughout adulthood. We confirmed that NSPF proteins are transferred to females during mating and hypothesize that proteins adhere to the spicules during transfer. NSPF proteins localize to the vulva and uterus lumen when transferred to mated females and in unmated adult hermaphrodites. These results suggest that the uterine localization of the NSPF proteins is likely a functional property of both male-derived sperm and self-sperm and not simply incidental to the point of transfer during mating. Our study demonstrates that nematodes use a novel mechanism for generating seminal fluid proteins and shows that the highly abundant NSPF proteins likely have a female-beneficial fitness effect.

Keywords: seminal fluid proteins, reproductive fitness, experimental evolution, *Caenorhabditis elegans*

INTRODUCTION

To understand the extent of adaptive evolution, we must link a phenotype with its underlying genetic basis and effect on fitness (Barrett & Hoekstra, 2011). By connecting the layers of the genotype-phenotype-fitness map we gain insight on the mechanisms that drive adaptation (Lewontin, 1974). Understanding these linkages will often demand an integrated, interdisciplinary experimental approach, which is necessarily challenging. Post-insemination reproductive interactions, however, offer an ideal stage of the lifecycle for mapping these relationships because their impact on reproductive fitness is direct, such that an adaptive advantage of one variant over another should lead to an increase in reproductive success. Additionally, the interactions involving cells, proteins, and small molecules simplifies the mapping of their genetic basis (McDonough *et al.*, 2016). Despite these direct and simplified linkages, connecting reproductive genes with a clear, consistent phenotype and a reproducible, quantifiable effect on fitness has proved challenging.

Post-insemination reproduction studies have largely focused on interactions mediated by seminal fluid. Seminal fluid is a complex mixture of cells, proteins, and other small molecules that are transferred to the female along with sperm (Poiani, 2006; Perry *et al.*, 2013). As a whole, seminal fluid is critical for male fertility – sperm alone is not enough to ensure fertilization – and is implicated in complex behaviors, such as sperm competition, male-female signaling and conflict (Poiani, 2006). Most functional work on these seminal fluid proteins comes from the study of dozens of *Drosophila* accessory gland proteins (Acps) (Chapman, 2011). The genes underlying many of these Acps are characterized by their elevated evolutionary rate relative to non-reproductive proteins, indicating adaptive evolution (Begun *et al.*, 2000; Swanson & Vacquier, 2002; Clark *et al.*, 2006; Wilburn & Swanson, 2016). The complexity of seminal fluid proteins coupled with their rapid evolution and some evidence of harmful effects on females have led to the paradigm that these proteins are beneficial to male fitness but detrimental female fitness (Sirot *et al.*, 2015). Perhaps the best studied Acp, Sex Peptide, has become a textbook example of post-insemination sexual conflict (Arnqvist & Rowe, 2005; Hopkins & Perry, 2022). However, this protein does not always fit the conflict paradigm it has come to represent (Hopkins & Perry, 2022). In particular, molecular evolution analysis of Sex Peptide indicates molecular constraint, which is not consistent with evolution driven by sexual conflict (Cirera & Aguade, 1997). Additionally, when individual components of fitness have been quantified, environmental

factors appear to impact when Sex Peptide is detrimental or even beneficial to female fitness

(Hopkins & Perry, 2022). These deeper analyses along with similar studies on other Acps (Wong

& Rundle, 2013) suggest that seminal fluid evolution is more nuanced than the sexual conflict

narrative alone and indicate that employing new approaches and new taxa may provide a more

precise and complete understanding of the complex roles and fitness effects of seminal fluid

proteins.

Caenorhabditis nematodes are powerful model system for integrating across the layers of

the genotype-phenotype-fitness landscape and are ideal for expanding our view on the extent of

evolution by sexual conflict and discovering new roles for seminal fluid proteins. First, mating

system differs across the genus, with *C. elegans* representing a lineage transition from obligate

outcrossing to hermaphroditic self-fertilization (Baldi *et al.*, 2009). Using the genetic tools

developed in *C. elegans*, researchers can manipulate hermaphrodites to develop as pseudo-

females that rely on males to reproduce, which allows direct comparison of the genetics and

fitness effects between self-sperm and male-derived sperm to disentangle conflict and

cooperation. Additionally, broader contrasts can be made among closely related species with

different mating systems and therefore varying opportunities for sexual conflict (Ting *et al.*,

2014; Yin *et al.*, 2018). Second, the small size and rapid generation time of nematodes make

them amenable to fitness assays within and across multiple generations. Multi-generation assays

allow for detection and quantification of even small fitness effects. Finally, nematodes have

unique amoeboid-like crawling sperm which can be visualized through the transparent cuticle,

allowing for spectacular live visualization of sperm and seminal fluid dynamics inside of females

or hermaphrodites (Smith & Stanfield, 2011; Ting *et al.*, 2014; Hansen *et al.*, 2015; Yin *et al.*,

2018). Very few seminal fluid proteins, however, have been identified and functionally

examined in *Caenorhabditis* (Smith & Stanfield, 2011; Yin *et al.*, 2018). Thus, this system

provides an underexploited opportunity for discovering new features of reproductive evolution

that can be evaluated and, perhaps, challenge existing paradigms.

An intriguing component of nematode sperm cells are subcellular vesicles known as

membranous organelles (MOs). In inactive spermatids, MOs are located around the periphery of

the cell (Ellis & Stanfield, 2014). Upon ejaculation, spermatids are released from the seminal

vesicle and undergo activation as they pass through the vas deferens. During activation, MOs

fuse with the cell membrane and create cup-like structures as well as releasing their soluble

protein contents into the extracellular space (Ellis & Stanfield, 2014). No accessory gland-like

tissues have been identified in *Caenorhabditis* and the source – as well as contents – of seminal

fluid are unknown. Consequently, we hypothesize that these MO fusions provide a novel

contribution to the seminal fluid proteome. Characterization of the MO proteome identified the

Nematode-Specific Peptide family, group F (NSPF) as the second most abundant set of proteins

in the MO (Kasimatis *et al.*, 2018). Kasimatis *et al.* (2018) additionally showed that the NSPF

genes are conserved in both sequence identity and synteny across the genus. Transcriptomic data

support the molecular evolution analyses by capturing NSPF expression in *C. elegans* males

along with NSPF ortholog expression in *C. remanei* males (Albritton *et al.*, 2014). Interestingly,

transcriptomic data also identify expression in *C. elegans* hermaphrodites but not *C. briggsae*

hermaphrodites, suggesting that this gene family has been conserved in at least some self-sperm

pathways (Albritton *et al.*, 2014).

Here we investigate the function of the NSPF gene family in fertilization by

interconnecting the layers of the genotype-phenotype-fitness landscape. We estimate the

contribution of the NSPF genes to female reproductive fitness using a combination of within

generation fertility assays and 20 generations of experimental evolution. We then use

immunostaining to identify the functional regions that NSPF proteins are associated with in

males and females/hermaphrodites. Our results indicate that the NSPF proteins are a component

of seminal fluid with a net fitness benefit and imply that even a highly abundant seminal fluid

protein can have a subtle function.

123   **MATERIALS AND METHODS**

Molecular biology

To create a tagged *nspf-1* gene, guides targeting the sequence in the same intergenic region

utilized by the ttTi5650 MosSCI site have previously been described (Dickinson *et al.*, 2015) and

inserted into pDD162 (Addgene #47549) to make plasmid pMS4. The repair template plasmid

(pKK4) was assembled using the NEBuild HiFi Kit (NEB) from a combination of a restriction

digest fragment and a synthetic gBlock Gene Fragment (IDT). The gBlock included the 647 base

pairs upstream of *nspf-1* as the promoter region, the *nspf-1* coding sequence with six histidines

added before the stop codon, and the *tbb-2* 3' UTR. The plasmid was purified using the GeneJET

Plasmid Miniprep Kit (Thermo Scientific) and assembly junctions were verified by Sanger

sequencing. A complete list of primers can be found in Table S1.

Strain generation

All strains used in this study are listed in Table S2. The NSPF genes were previously deleted

using CRISPR/Cas9 to create strain PX814 (fxDf1 (II: 2,484,339 - 2,487,244)) (Kasimatis *et al.*,

2018). To generate an obligate outcrossing deletion strain, JK574 (*fog-2*(q71) V) was crossed

with PX814 and then backcrossed 5 times with JK574 to create the obligate outcrossing, deletion

strain VX194. The deletion region was verified by Sanger sequencing.

Insertion of the His-tagged *nspf-1* transgene was done by CRISPR/Cas9 using standard

methods (Dickinson *et al.*, 2015). Briefly, a mixture of 10ng/µL repair template plasmid,

50ng/µL plasmid encoding CAS9 and the guide RNA, and 2.5ng/µL of the co-injection plasmid

pZCS16 was injected into the gonad of young adult hermaphrodites from PX814. Hygromycin B

(BioShop) was added to the injection plates at a final concentration of 250µg/mL three days after

injection and hygromycin resistance (HygR) was used as a selectable event. Worms were kept at

25ºC for the first five days immediately following injections and then transferred for 20ºC for

subsequent generations. Successful insertion was confirmed by PCR (Table S1) and Sanger

sequencing. Worms were then backcrossed five times to PX814 to create the final strain VX300

(yf*Si1*[*nspf-1*p::*nspf-1*::6xHis::*tbb-2* 3' UTR + loxP, II:8420157]). We tagged the C-terminal of

nspf-1 since these proteins are predicted signaling peptides (Kasimatis *et al.*, 2018) and, thus, the

C-terminal is likely included in the mature peptide.

Fertility Assays

We assayed the overall reproductive success of VX194 deletion males late in adulthood. We

isolated virgin deletion males on day 1 of adulthood. On day 5 adulthood, single males were

mated with three wildtype, virgin females (strain JK574) for 24 hours. As a control, wildtype

males (strain JK574) were mated to wildtype females following the same timing and male to

female ratio. Matings were conducted on small NGM-agar plates (35 mm diameter) seeded with

10µL OP50 *Escherichia coli*. After 24 hours, each male was removed and the females were

transferred to a new small plate to continue laying eggs. Females were transferred to new plates

every 24 hours until progeny production ceased. The total number of progeny was counted as a

measure of each male's reproductive success (File S1). Fertility data were analyzed using R (R

Core Team, 2020), with the significance of non-competitive reproductive success evaluated

using Welch's Two Sample t-test and an analysis of the power of the comparison computed.

Previous fecundity data were replotted using the data available in Kasimatis et al. (2018)

Additional File 9.

Experimental evolution design and worm culture

To create a base population, reciprocal crosses of virgin females and males from strains JK574

and VX194 were performed. The F1 heterozygous progeny were divided into 10 replicate

populations of 500 worms with equal numbers of each sex to start the first generation. After

generation one, populations were expanded to a census size of N = 1,000 worms per replicate

and evolved for 20 generations. To start each generation, age synchronized larval stage 1 (L1)

worms were plated onto four 10 cm NGM-agar plates seeded with OP50 *E. coli* at 20ºC with a

density of 250 worms per plate (Brenner, 1974; Kenyon *et al.*, 1993). Worms were left to mate

for the first five days of adulthood before eggs were once again collected on day six to

synchronize for the next generation. During this mating time, worms within each replicate were

pooled and redistributed to fresh NGM-agar plates twice to ensure they never starved. A subset

of worms from all replicates was frozen every five generations for future experiments. At

generation eight all worms were frozen and the experiment paused for five months due to

COVID-19 lockdown restrictions.

Amplicon sequencing and read calling

We performed amplicon sequencing on pooled samples of 500-800 L1 worms from experimental

evolution generations 11 and 20. Two control samples were generated, each with approximately

700 worms: 1) F1 heterozygous progeny generated by crossing strains JK574 and VX194, and 2)

pooled worms generated by mixing an equal number of JK574 and VX194 worms and sampling

from this pool. The wildtype allele was amplified using forward primer (5' to 3')

TGCAATGGATTTCTGGTGATAAC and reverse primer (5' to 3')

TCTGGCAGTAGGAAAGTGAG, yielding a product of size 230 base pairs. The deletion allele

was amplified using forward primer (5' to 3') AGGATAAGAGAGACCCAGAC and reverse

primer (5' to 3') TCTGGCAGTAGGAAAGTGAG, yielding a product of size 275 base pairs.

PCR products were generated using the OneTaq Quick-Load 2x PCR Master Mix with Standard

Buffer (NEB) in accordance with manufacturer instructions with *C. elegans* genomic DNA. PCR

products were purified using QIAquick PCR Purification kit (Qiagen) and normalized to

20ng/uL. Products were sequenced using GENEWIZ Amplicon-EZ services.

Reads for each sample were filtered using bioawk (https://github.com/lh3/bioawk) to

exclude reads <50 base pairs in length and with a mean quality score <30. Reads in each sample

>230 base pairs in length were then blasted to the deletion reference sequence and retained if the

blast score was significant. All remaining reads were then blasted to the unique reference

sequence for both the wildtype (bases 1-102) and deletion (bases 1-147) alleles. A read was

called as a wildtype or deletion allele based on the best E-value and bit score (File S2). Final

counts ranged from 89,721 to 132,685 reads per sample.

Estimating selection

Read data were analyzed using the R statistical language v4.0.0 (R Core Team, 2020). Read

counts across all time points were fit using a generalized linear mixed model with a binomial

logistic distribution: $glm((WT_{count}, DEL_{count}) \sim time)$. We fit two different linear models to

account for biases during PCR amplification as suggested by the frequency of each allele in the

control samples (i.e., 73% wildtype and 27% deletion). Model 1 assumed the frequency of each

allele to be 50% (i.e., 500 copies of the wildtype allele and 500 copies of the deletion allele for a

heterozygous population of $N = 500$), since all worms were heterozygous at the start of the

experiment. Model 2 used only the allele frequencies estimated from the amplicon sequencing at

generations 11 and 20. While we report the results of both models, Model 2 provides a better fit

($AIC_{Model\ 1} = 146,842$; $AIC_{Model\ 2} = 146,670$) and represents a more conservative estimate of

selection.

The neutral expectation was determined by simulating in R allele frequency dynamics at

a single locus in a population subject to genetic drift and random mating. We simulated 100

populations of 1,000 individuals. In the first generation, all individuals were heterozygotes. The

genotype frequencies of females and males were multiplied to give a matrix of the relative

frequencies of the possible random mating pairs. Mating pairs were then randomly sampled from

this distribution to generate offspring to start the next generation.

Whole-worm immunohistochemistry

Worms were formaldehyde-fixed and stained following a modified version of the Ruvkun and

Finney protocol (WormAtlas). Briefly, worms were cultured at 20ºC on 10cm NGM-agar plates

containing 100ng/mL carbenicillin and seeded with formaldehyde killed *E. coli* OP50 (following

Rogers *et al.*, 2006). At each developmental stage used, populations of at least 1,000 worms were

fixed in a 1% formaldehyde solution and flash frozen in dry ice with ethanol. Incorrect antibody

binding was blocked using solution of: 0.3M glycine, 10mM imidazole, 1% BSA, 0.5% skim

milk, and 0.25% Triton-X-100 in PBS. Worms were stained for 1 hour at room temperature

using a 1/4000 dilution of Recombinant Alex Fluor 488 Anti-6x His Tag antibody (Abcam

#ab237336) and co-stained with DAPI. The complete method along with buffer recipes can be

found in File S3.

To examine the localization of *nspf-1* in males and hermaphrodites across development,

VX300 male-rich populations (~40% males) were age synchronized by bleaching

{Kenyon:1988uc} and then formaldehyde-fixed during: L3 (28 hours post-L1), L4 (40 hours

post-L1), day 1 of adulthood (46 hours post-L1), day 2 of adulthood (70 hours post- L1), and day

8 of adulthood (190 hours post-L1). As a control, JK574 males and pseudo-females were

concurrently cultured and fixed at the same developmental stages. In our hands, worms

developed more slowly and less synchronously on plates with dead OP50 than on live OP50.

Thus, the exact time in development may be asynchronous between worms, but the

developmental stages were accurate. At least 100 stained worms from each strain × sex ×

developmental stage were scored for the presence and location of fluorescence (File S4). The

localization of fluorescence was characterized as occurring in one or more of the following

regions: head, distal gonad, proximal gonad (including the spermathecae or seminal vesicle),

uterus/vas deferens, vulva/cloaca, or tail. Worms were categorized as "not stained" if no

fluorescence was observed.

To determine whether or not males transfer NSPF-1 to females during mating, we crossed

JK574 pseudo-females with VX300 males. Both strains were age-synchronized by bleaching

followed by isolation of virgin pseudo-females (n = 400) and males (n = 150) during late L4 to

10cm NGM-agar plates containing 100ng/mL carbenicillin and seeded with formaldehyde-killed

E. coli OP50 (n = 3 replicate plates). Worms were mated overnight and then formaldehyde-fixed

and stained. One hundred pseudo-females were scored for the presence and location of

fluorescence.

Count data for the presence of fluorescence were analyzed using a χ^2 test based on a

contingency table of strain × presence of fluorescence for each sex and developmental stage

separately. Specifically, we tested the null hypothesis of no difference between counts of stained

individuals and counts of unstained individuals between the control (JK574) and His-tagged

(VX300) strains. All analyses were conducted in R (Team'', 2020).

Dissected gonad immunohistochemistry

Testes were methanol-fixed and stained following a modified version of the Shakes *et al.* (2009)

protocol. Briefly, worms were cultured at 20ºC on 10cm NGM-agar plates and seeded with *E.*

coli OP50. Young adult males were decapitated using a 30 gauge needle to release the gonad and

then transferred to ColorFrost Plus slides (Fisher Scientific). Gonads were fixed in methanol

followed by acetone. Incorrect antibody binding was blocked using solution of: 0.3M glycine,

10mM imidazole, 1% BSA, and 0.25% Triton-X-100 in PBS. Worms were stained for 1 hour at

room temperature using a 1/4000 dilution of Recombinant Alex Fluor 488 Anti-6x His Tag

antibody (Abcam #ab237336) and co-stained with DAPI. The complete method along with

buffer recipes can be found in File S3. Image stacks were processed using ImageJ.

RESULTS

Deletion of the NSPF genes decreases male fitness

Previous experiments on young adult male fertility showed no significant difference in total

fecundity between wildtype males and males with all three copies of the NSPF genes deleted

(i.e., deletion allele) (Fig. S1A) (Kasimatis *et al.*, 2018). Using old males (day 5 of adulthood),

we repeated these non-competitive, total fecundity assays to determine whether NSPF proteins

improved male sperm fertilization ability in a phase of the life cycle when male fertility is

declining overall. Although the mean fecundity of old males with the deletion allele trended

lower than wildtype males (x_{del} = 459; x_{wt} = 481.9), the effect was not significant (p = 0.85; Fig.

S1A). This nominal decrease in fecundity of males with the deletion allele was similar to what

was previously seen with young adult males under a competitive male setting, suggesting that

NSPF function is potentially age-independent (Kasimatis *et al.*, 2018; Fig. S1B). However, given

the experiment size and high reproductive variance, a difference between genotypes of 71%

would be required to achieve 80% power to detect a significant difference. Thus, within-

generation assays likely are not sufficiently sensitive to capture the nuanced fitness effects

suggested by the trend means, despite the high abundance of these proteins.

We turned to experimental evolution to overcome these power limitations of within-

generation experimental design. Specifically, we competed the wildtype allele against the

deletion allele in an obligate outcrossing *C. elegans* population by evolving ten replicate

populations of 1,000 individuals from a base population that was heterozygous at the NSPF locus

in an otherwise homogeneous genetic background. After 20 generations, we documented a

significant overall decrease in the frequency of the deletion allele across replicates (Model 1: z-

value = 3.3, p = 0.001; Model 2: z-value = 2.3, p = 0.02). Additionally, the change in frequency

exceeded that expected by neutral processes alone in seven of the ten replicates (Fig. 1). These

results support the within generation fecundity trends and indicate a direct net fitness cost when

the NSPF genes are removed, such that the wildtype allele confers an approximately 0.1% fitness

advantage over the deletion allele (Model 1: *s* = 0.001; Model 2: *s* = 0.0008). Together, the

favoring of the wildtype allele coupled with the fecundity do not provide evidence of a cost of

the NSPF proteins during mating. Rather, they provide a net fitness effect that is congruent with

being beneficial to at least one sex.

NSPF proteins localize to the spicules in adult males

We characterized the functional location of NSPF proteins during fertilization using whole-worm

immunostaining of a His-tagged *nspf-1* transgene re-inserted into the deletion allele background

(PX814). We scored the presence the fluorescent antibody bound to NSPF-1::6xHis in six

anatomical regions: head, distal gonad, proximal gonad, vas deferens, cloaca, and tail (Fig. 2A).

Throughout male development, we found that NSPF presence and abundance was correlated

with male reproductive maturity (Fig. 2B; Fig S2). During larval stage 3 (L3), the gonad is still

developing and no staining was present, which supports our previous finding that NSPF

production is linked with sperm production (File S4). Sperm production begins during the L4

stage and correspondingly L4 males had staining in both the distal (n = 17 of 126) and proximal

gonad (n = 1 of 126; Fig. 3A). Interestingly, 43% of L4 males had staining in the tail region (n =

55 of 126) specifically overlapping with the developing spicules. These patterns are significantly

different from control males (χ^2 = 175.5, df = 3, p < 0.0001), of which only one individual

showed non-specific staining along the cuticle in the proximal gonad region (Fig. 3B).

In adult males, NSPF proteins localized solely to the spicules (Fig. 3C). Abundance

appeared highest on days 1 and 2 of adulthood with 100% of males (n = 100 and 150,

respectively) showing staining in the tail region (Fig. 2B). Again, these patterns were

significantly different from the off-target staining or spicule auto-fluorescence observed in

control males during day 1 (n = 13 of 99 stained; χ^2 = 149.5, df = 1, p < 0.0001) and day 2 (n =

45 of 150 stained; $\chi^2 = 158.5$, df = 1, p < 0.0001) of adulthood (Fig. 3D). Older males, however,

had a decreased signal of NSPF proteins (n = 59 of 106), though still significantly different from

control males ($\chi^2 = 39.4$, df = 1, p < 0.0001). This pattern of protein abundance throughout

development suggests that NSPF production is tightly correlated with spermatogenesis and peak

reproductive output and mating activity during early adulthood.

While immunostaining of whole adult worms did not identify NSPF protein within the

gonad, dissected testes of adult males confirmed localization of NSPF within spermatids (Fig.

S3). The punctate patterning around the nucleus overlaps with the location of MOs, which

supports the proteomic characterization of this gene family. Thus, whole worm immunostaining

likely lacked the sensitivity to visualize staining within small sperm subcellular structures.

Males transfer NSPF proteins to the female uterus during mating

To test our hypothesis that NSPF proteins form a constituent of seminal fluid, we examined

whether males transferred NSPF proteins during mating. Specifically, control pseudo-females

(JK574) were mated with His-tagged males (VX300) for 24 hours. Of the pseudo-females

without eggs in utero, 38% (n = 8 of 21) had a clear signal of staining around the vulva and in the

uterine space (Fig. 4A). This signal was significantly different from that of control females

mated to control males ($\chi^2 = 9.7$, df = 2, p = 0.008; Fig. 4B). Only 12% (n = 15 of 129) of

pseudo-females with eggs in utero showed a clear signal of staining, which was not significantly

different from pseudo-females mated to control males ($\chi^2 = 0.53$, df = 1, p = 0.47). However, in

these females the NSPF proteins appear to be in the uterus lumen, which suggests that eggs may

mask the fluorescent signal (Fig. 4C). These results indicate that males transfer NSPF proteins

during ejaculation and support our hypothesis that MOs contribute proteins to seminal fluid.

NSPF proteins localize to the uterus of unmated hermaphrodites

To determine if self-sperm NSPF patterns differed from male-derived sperm patterns, we

characterized NSPF localization in His-tagged hermaphrodites (VX300) raised in the presence of

males throughout development. Again, L3 individuals showed no signal of staining (File S4). L4

hermaphrodites, however, had little to no staining and did not significantly differ from control

females (p = 1; Fig. 2C) despite mRNA expression of *nspf-1* during this developmental stage

(Harris *et al.*, 2019). The non-overlapping mRNA and protein localization patterns are likely to

be an artifact of our whole-animal immunostaining procedure. Adult hermaphrodites all showed

the same vulva and uterus localization pattern of staining as we observed in mated, control

females (Fig. 2C; Fig. S2). On days 1 and 2 of adulthood 54% (n = 54 of 100) and 41% (n = 61

of 150) of hermaphrodites, respectively, showed staining, which was significantly more than

control females (Day 1: χ^2 = 49.9, df = 1, p < 0.0001; Day 2: χ^2 = 22.6, df = 1, p < 0.0001). On

day 8 of adulthood, 67% (n = 90 of 134) of hermaphrodites showed the same significant staining

pattern (χ^2 = 147.2, df = 6, p < 0.0001; Fig. 2C). Because self-sperm are depleted by day 8 late in

adulthood (Hirsh *et al.*, 1976), these data further support the male transfer of NSPF proteins to

the uterus. Across all developmental time points, there was an unexpected absence of signal in

the spermathecae, likely due to experimental constraints.

To distinguish whether the early adulthood signal of NSPF staining was caused by self-

fertilization or mating, we examined a male-less population of His-tagged hermaphrodites on day

1 of adulthood prior to the start of egg laying. If the NSPF proteins were costly to female

reproduction, then we would expect self-sperm derived NSPF to be lost or significantly reduced.

We found that 59% (n = 59 of 100) of hermaphrodites showed the same vulva and uterus pattern

of staining (Fig. 4D) despite no opportunity for male transfer of NSPF proteins. This frequency

was not significantly different from the number of day 1 hermaphrodites raised in the presence of

males (p = 0.57). These results suggest that the uterine localization of the NSPF proteins –

whether secreted by self-sperm or male sperm – likely reflects their site of activity and not

simply a byproduct of the point of transfer during mating. Moreover, they reinforce the benefit of

NSPF proteins to reproductive fitness and imply that this benefit is directed toward female

reproduction.

368   **DISCUSSION**

Seminal fluid is a complex bioactive mixture whose components are implicated in a variety of

reproductive, behavioral and physiological functions and thus likely evolves under multiple

selective pressures. By linking the genes creating seminal fluid proteins with their evolutionary

history, effect on fitness, and phenotypic contribution we can understand the evolutionary forces

driving seminal fluid evolution through its impacts on reproductive function. We took an

integrative approach to connect the highly abundant MO NSPF proteins across the genotype,

phenotype, fitness landscape. We confirmed that the NSPF genes are indeed a component of

seminal fluid that is transferred to females, which demonstrates that MO fusion provides a novel

source of seminal fluid proteins. Moreover, the localization of NSPF proteins differ from that of

mRNA transcripts, highlighting the nested layers between genotype space and phenotype space

that can often be overlooked. Finally, we quantified the small, but significant contribution made

to reproductive fitness by the presence of NSPF proteins, suggestive of a fitness benefit

potentially to females. Together this work demonstrates the power of *Caenorhabditis* to

disentangle beneficial versus antagonistic effects of seminal fluid proteins and expand our

perspective of seminal fluid protein function.

The manipulable hermaphroditic mating system of *C. elegans*, relative to study systems

like *Drosophila*, offers a novel means of distinguishing sexual conflict from sexual cooperation

between male-derived sperm and self-sperm interactions (Schärer *et al.*, 2015). We capitalized

on this advantage to show that NSPF production is not only retained during the lineage transition

to self-fertilization, but that the proteins localize to the same uterus region in unmated

hermaphrodites as mated females (Fig. 4). This pattern supports the conserved molecular

evolution of the *nspf* gene sequences and implies that the function of the sperm-dependent NSPF

proteins within the uterus is mating-independent. Importantly, the conserved expression in

hermaphrodites indicates that these proteins are not involved in sexual conflict, as there would be

strong selection favoring their loss (Yin *et al.*, 2018). Instead, these genes appear beneficial to

fitness in a female-biased manner, as supported by fitness assays. In particular, if NSPF genes

provided an advantage to males to the detriment of females, we would expect females mated

with deletion males to trend toward higher reproductive success than females mated with

wildtype males. However, our assay and previous assays support the opposite of such a scenario,

suggesting NSPF promotes sexually-concordant reproductive fitness, as shown more

convincingly during experimental evolution.

A common challenge in evolutionary biology is to detect the fitness effects of variants at

a single gene (Barrett & Hoekstra, 2011). Our experimental evolution design to compete the

wildtype and deletion alleles in a highly-replicated fashion allowed us to compound small effect

sizes over tens of generations. The nematode model system allows for this approach to be done

at a larger scale and over a longer time periods than similar studies (Wong & Rundle, 2013),

which, when coupled with genomics, provides greater power to detect small or fluctuating

selection coefficients. The overall trend across replicates was a decrease in the frequency of the

deletion allele beyond the neutral expectation (Fig. 2). The estimated selection coefficient,

despite being small, would be powerful enough to prevent *nspf* gene loss over time in nature. The

frequency of the deletion allele did not continually decrease in every replicate at each sampling

time point, however, which likely reflected a combination of variance and stochasticity during

mating along with sampling error and the relatively small selection coefficient. The fluctuation

of the deletion allele could also reflect the environment conditions in the lab which do not

perfectly replicate nature, where the NSPF proteins may provide a greater or more consistent

benefit in populations with larger effective sizes. This approach of allelic competition in

experimental evolution (Gray & Cutter, 2014; Teotonio *et al.*, 2017) may represent a promising

avenue for future seminal fluid protein studies, and for genetic variants of other phenotypes,

where small fitness effect sizes have proved challenging to characterize.

The localization of NSPF protein to the spicule (Fig. 3) and uterus (Fig. 4) was

unexpected given that the NSPFs were previously characterized as being associated with sperm

MOs (Kasimatis *et al.*, 2018) and that mRNA expression data indicated that all three *nspf* genes

are expressed in the gonad only (Kruse *et al.*, 2016). Staining of dissected testes provided the

necessary sensitivity to confirm that NSPF proteins localize to spermatid MOs. Our whole-worm

staining approach did, however, identify additional regions of localization that may result from

the anatomical difference in the location of inactivated versus activated sperm. The activation

process occurs as sperm pass from a male to a female/hermaphrodite while the spicules are

everted in the vulva, thus creating a direct link between MO fusion and the spicules. This overlap

in timing, however, does not explain the spicule signal seen in L4 males, which warrants further

work. In hermaphrodites, the parsimonious hypothesis is that NSPF proteins passively diffuse

from the spermathecae, where activated sperm reside, to the uterus. The location of NSPF

proteins within the uterus is similar to other proteins that males transfer, which originate in

newly identified extracellular vesicles (Nikonorova *et al.*, 2022). While the NSPFs do not appear

in these vesicles, it is possible that this alternative vesicle type also contributes to seminal fluid

proteins, supporting our finding that nematodes employ a distinct mechanism for secreting

seminal fluid proteins than either mammals or *Drosophila* (McGraw *et al.*, 2015).

Our data motivate multiple hypotheses for future testing to narrow down the specific

beneficial function of the NSPF proteins. First, the NSPF proteins could interact with a male-

female signaling network that promotes egg laying. While the interacting partner for NSPF

proteins remains unknown, it is possible that NSPFs bind to muscle proteins that form the uterus

walls and are involved in egg laying behavior. Under this hypothesis, localization to the spicules is puzzling. However, previous work has shown the females pressurize the gonad to push some of the ejaculate mass and spicules out of the vulva (Barker, 1994; Kleemann & Basolo, 2007). This process could create a residue on the spicule cuticle as a byproduct, though we cannot rule out a functional relationship between the NSPF proteins and spicule cuticle.

Alternatively, the NSPF proteins could impact the uterine environment. Specifically, seminal fluid of some species contains anti-microbial factors (Samakovlis *et al.*, 1991; Gao & Zhu, 2016). Given the density of protein right above the vulva, one hypothesis is that the NSPF proteins play an anti-microbial role to protect the uterus from bacterial infection. Previous work has shown that the uterus is exposed to pathogenic infection from the external environment during mating and egg laying (Muir & Tan, 2008). The male spicules are presumably also exposed during eversion, and thus any anti-microbial activity of the NSPF residue on the spicules might serve to prevent bacterial infection when the spicules retract. Biochemically, the alpha-helix structure of NSPF proteins lends plausibility to this hypothesis, as such secondary structures can confer generic bacterial lysing properties (Huan *et al.*, 2020). Future work testing the fitness effects of the NSPF deletion allele in worms raised in more pathogenic bacteria is warranted.

Our work demonstrates the complex but subtle dynamics of seminal fluid proteins uncovered when using integrative approaches. The NSPF seminal fluid proteins, in particular, exhibit sexually-concordant rather than sexually-antagonistic fitness effects. We also showed how *Caenorhabditis* nematodes are an excellent system for expanding our knowledge of post-insemination interactions not only for their technical benefits, but also for their novel sperm components and potential for making mating system contrasts. Connecting the layers of the genotype-phenotype-fitness map for seminal fluid proteins will help us uncover the genetics of fertility and the evolution of reproductive success.

Acknowledgements: We thank the Phillips lab for sharing plasmids pMS4 and pZCS16 and strain PX814. We thank Zachary Stevenson for transgenic advice, Jack Hu and Mathias Renaud for immunostaining advice, and Stephen Banse, Eric Haig and the Cutter and Claycomb labs for their helpful discussion.

Funding: This research was supported by a Natural Sciences and Engineering Research Council

of Canada (NSERC) Banting Postdoctoral Fellowship to KRK and NSERC Discovery Grant

Program funding to ADC and LR. The funders had no role in study design, data collection and

analysis, decision to publish, or preparation of the manuscript.

Author Contributions: KRK, ADC, and LR devised the project. KRK created strains VX194

and VX300. KRK performed the fecundity assay and experimental evolution. CR performed the

immunostaining assays with assistance from KRK. KRK analyzed the data. KRK wrote the

manuscript with the support of the other authors.

Data accessibility

The oligonucleotides used in this study are available as Table S1. The fertility data are available

in File S1. The raw amplicon data will be made publicly available at NCBI SRA upon

publication. The processed amplicon reads are available in File S2. The immunostaining data are

available in File S3 and the complete protocol is available in File S4. All scripts are available via

the Cutter Lab GitHub repository NSPF (https://github.com/Cutterlab/NSPF). Worm strains

JK574 and VX300 are available from the *Caenorhabditis* Genetics Center. Strains PX814 and

VX194 are available from the Cutter lab upon request.

References

Albritton, S.E., Kranz, A.-L., Rao, P., Kramer, M., Dieterich, C. & Ercan, S. (2014) Sex-biased

gene expression and evolution of the x chromosome in nematodes. *Genetics*, **197**, 865–883.

Arnqvist, G. & Rowe, L. (2005) *Sexual conflict*. Princeton University Press.

Baldi, C., Cho, S. & Ellis, R.E. (2009) Mutations in two independent pathways are sufficient to

create hermaphroditic nematodes. *Science*, **326**, 1002–1005.

Barker, D.M. (1994) Copulatory plugs and paternity assurance in the nematode *Caenorhabditis*

elegans. *Animal Behaviour*, **48**, 147–156.

Barrett, R.D.H. & Hoekstra, H.E. (2011) Molecular spandrels: tests of adaptation at the genetic

level. *Nature Reviews Genetics*, **12**, 767–780.

Begun, D.J., Whitley, P., Todd, B.L., Waldrip-Dail, H.M. & Clark, A.G. (2000) Molecular
population genetics of male accessory gland proteins in *Drosophila*. *Genetics*, **156**, 1879–
1888.

Brenner, S. (1974) The genetics of *Caenorhabditis elegans*. *Genetics*, **77**, 71–94.

Chapman, T. (2011) Seminal fluid-mediated fitness traits in *Drosophila*. *Heredity*, **87**, 511–521.

Cirera, S. & Aguade, M. (1997) Evolutionary history of the sex-peptide (Acp70A) gene region in
Drosophila melanogaster. *Genetics*, **147**, 189–197.

Clark, N.L., Aagaard, J.E. & Swanson, W.J. (2006) Evolution of reproductive proteins from
animals and plants. *Reproduction*, **131**, 11–22.

Dickinson, D.J., Pani, A.M., Heppert, J.K., Higgins, C.D. & 2015. (2015) Streamlined genome
engineering with a self-excising drug selection cassette. *Genetics*, **200**, 1035–1049.

Ellis, R.E. & Stanfield, G.M. (2014) The regulation of spermatogenesis and sperm function in
nematodes. *Seminars in Cell and Developmental Biology*, **29**, 17–30.

Gao, B. & Zhu, S. (2016) The drosomycin multigene family: three-disulfide variants from
Drosophila takahashii possess antibacterial activity. *Nature Reviews Genetics*, **6**, 32175.

Gray, J.C. & Cutter, A.D. (2014) Mainstreaming *Caenorhabditis elegans* in experimental
evolution. *Proceedings of the Royal Society B: Biological Sciences*, **281**, 20133055.

Hansen, J.M., Chavez, D.R. & Stanfield, G.M. (2015) COMP-1 promotes competitive advantage
of nematode sperm. *eLife*, 4:e05423.

Harris, T.W., Arnaboldi, V., Cain, S., Chan, J., Chen, W.J., Cho, J., *et al.* (2019) WormBase: a
modern model organism information resource. *Nucleic Acids Research*, **29**, 82.

Hirsh, D., Oppenheim, D. & Klass, M. (1976) Development of the reproductive system of
Caenorhabditis elegans. *Developmental Biology*, **49**, 200–219.

Hopkins, B.R. & Perry, J.C. (2022) The evolution of sex peptide: sexual conflict, cooperation,
and coevolution. *Biological Reviews*, **97**, 1426–1448.

Huan, Y., Kong, Q., Mou, H. & Yi, H. (2020) Antimicrobial peptides: classification, design,
application and research progress in multiple fields. *Frontiers in Microbiology*, **11**, 582779.

Kasimatis, K.R., Moerdyk-Schauwecker, M.J., Timmermeyer, N. & Phillips, P.C. (2018)
Proteomic and evolutionary analyses of sperm activation identify uncharacterized genes in
Caenorhabditis nematodes. *BMC Genomics*, **19**, 593.

Kenyon, C., Chang, J., Gensch, E., Rudner, A. & Tabtiang, R. (1993) A *C. elegans* mutant that
lives twice as long as wild type. *Nature*, **366**, 461–464.

Kleemann, G.A. & Basolo, A.L. (2007) Facultative decrease in mating resistance in
hermaphroditic *Caenorhabditis elegans* with self-sperm depletion. *Animal Behaviour*, **74**,
1339–1347.

Kruse, F., Junker, J.P., Oudenaarden, A. van & Bakkers, J. (2016) Tomo-seq: A method to obtain
genome-wide expression data with spatial resolution. *Methods in cell biology*, **135**, 299–307.

Lewontin, R.C. (1974) *The genetic basis of evolutionary change*. Columbia University Press.

McDonough, C.E., Whittington, E., Pitnick, S. & Dorus, S. (2016) Proteomics of reproductive
systems: Towards a molecular understanding of postmating, prezygotic reproductive barriers.
Journal of Proteomics, **135**, 26–37.

McGraw, L.A., Suarez, S.S. & Wolfner, M.F. (2015) On a matter of seminal importance.
BioEssays, **37**, 142–147.

Muir, R.E. & Tan, M.-W. (2008) Virulence of *Leucobacter chromiireducens* subsp. *solipictus* to
Caenorhabditis elegans: characterization of a novel host-pathogen interaction. *Applied and*
Environmental Microbiology, **74**, 4185–4198.

Nikonorova, I.A., Wang, J., Cope, A.L., Tilton, P.E., Power, K.M., Walsh, J.D., *et al.* (2022)
Isolation, profiling, and tracking of extracellular vesicle cargo in *Caenorhabditis elegans*.
Current Biology, **32**, 1–13.

Perry, J.C., Sirot, L. & Wigby, S. (2013) The seminal symphony: how to compose an ejaculate.
Trends in Ecology & Evolution, **28**, 414–422.

Poiani, A. (2006) Complexity of seminal fluid: a review. *Behavioral Ecology and Sociobiology*,
60, 289–310.

Rogers, A.B., Cormier, K.S. & Fox, J.G. (2006) Thiol-reactive compounds prevent nonspecific
antibody binding in immunohistochemistry. *Laboratory Investigation*, **86**, 526–533.

Samakovlis, C., Kylsten, P., Kimbrell, D.A., Engström, A. & Hultmark, D. (1991) The andropin
gene and its product, a male-specific antibacterial peptide in *Drosophila melanogaster*. *The*
EMBO Journal, **10**, 163–169.

Schärer, L., Janicke, T. & Ramm, S.A. (2015) Sexual conflict in hermaphrodites. *Cold Spring*
Harbor Perspectives in Biology, **7**, a017673.

Shakes, D.C., Wu, J., Sadler, P.L., LaPrade, K., Moore, L.L., Noritake, A., *et al.* (2009)
Spermatogenesis-specific features of the meiotic program in *Caenorhabditis elegans*. *PLoS*
Genetics, **5**, e1000611.

Sirot, L.K., Wong, A., Chapman, T. & Wolfner, M.F. (2015) Sexual conflict and seminal fluid
proteins: a dynamic landscape of sexual interactions. *Cold Spring Harbor Perspectives in*
Biology, **7**, a017533.

Smith, J.R. & Stanfield, G.M. (2011) TRY-5 Is a sperm-activating protease in *Caenorhabditis*
elegans seminal fluid. *PLoS Genetics*, **7,** e1002375.

Swanson, W.J. & Vacquier, V.D. (2002) Reproductive protein evolution. *Annual Review of*
Ecology and Systematics, **33**, 161–179.

R Core Team. (2020) *R: A language and environment for statistical computing*. Foundation for
Statistical Computing.

Teotonio, H., Estes, S., Phillips, P.C. & Baer, C.F. (2017) Experimental evolution with
Caenorhabditis nematodes. *Genetics*, **206**, 691–716.

Ting, J.J., Woodruff, G.C., Leung, G., Shin, N.-R., Cutter, A.D. & Haag, E.S. (2014) Intense
sperm-mediated sexual conflict promotes reproductive isolation in *Caenorhabditis*
nematodes. *PLoS Biology*, **12**, e1001915.

Wilburn, D.B. & Swanson, W.J. (2016) From molecules to mating: Rapid evolution and
biochemical studies of reproductive proteins. *Journal of Proteomics*, **135**, 12–25.

Wong, A. & Rundle, H. (2013) Selection on the *Drosophila* seminal fluid protein Acp62F.
Ecology and Evolution, **3**, 1942–1950.

Yin, D., Schwarz, E.M., Thomas, C.G., Felde, R.L., Korf, I.F., Cutter, A.D., *et al.* (2018) Rapid
genome shrinkage in a self-fertile nematode reveals sperm competition proteins. *Science*,
359, 55–61.

Figures

Figure 1. Changes in the frequency of the deletion allele over time estimated using

amplicon sequencing data. All ten replicates started with 50% deletion allele (dashed line). The

shaded gray region represents the expected allele frequency range under genetic drift and random

mating estimated by simulation. There was a significant decrease in the frequency of the deletion

allele over time (Model 1: s = 0.001, z-value = 3.3, p < 0.001; Model 2: s = 0.0008, z-value =

2.3, p < 0.05).

Figure 2. Quantification of antibody staining of 6x His-tagged NSPF-1. A) Diagram of the

anatomical regions scored for presence of fluorescence in hermaphrodites/females and males. **B)**

The percent of control males (JK574, triangle) and His-tagged males (VX300, circle) with

fluorescence in the tail region during larval stage 4 (L4), day 1 of adulthood (D1), day 2 of

adulthood (D2), and day 8 of adulthood (D8). **C)** The percent of control females (JK574,

triangle) and His-tagged hermaphrodites (VX300, circle) with fluorescence in the vulva region

during larval stage 4 (L4), day 1 of adulthood (D1), day 2 of adulthood (D2), and day 8 of

adulthood (D8). Immunostaining counts across all anatomical regions is shown in Fig. S2.

Figure 3. NSPF-1 localizes the spicules of adult males. A) Protein localized to the gonad of L4

His-tagged males. **B)** L4 control males showed off-target staining, especially in the cuticle, but

not in the spicules. **C)** In adult His-tagged males, protein localized to the spicule structure in the

male tail region. **D)** In adult control males, low levels of autofluorescence are detected in the

spicules. However, the signal is markedly different than see in His-tagged males. In all images,

the arrow points to the spicule.

Figure 4. NSPF-1 localizes to the vulva and uterus of mated females and adult hermaphrodites. A) Control pseudo-females mated to His-tagged males showed evidence of protein localization in the vulva and uterus, especially when eggs were not yet present in the uterus. **B)** Control pseudo-females mated to control males showed no evidence of staining. **C)** Control pseudo-females mated with eggs present in the uterus after mating with His-tagged males also had protein present in the uterus, though the signal was not always clear. However, when present the proteins seemed to localize to the lumen and potentially along the surface of vulva muscle cells, as shown in the merged image. **C)** His-tagged adult hermaphrodites who had never been mated by a male showed the same protein localization in the uterus. In all images, the arrow points to the vulva.

Supplemental Files

Fig S1. Single male fecundity assays. A) The total fecundity of wildtype males with the NSPF genes (gray) and deletion males with the NSPF genes removed (orange) at day 1 of adulthood and day 5 of adulthood. There was no significant difference in reproductive success between male types. Day 1 data were replotted from Kasimatis et al. (2018). **B)** The proportion of progeny coming from wildtype males (gray) or deletion males (orange) versus a wildtype competitor male. There was no significant difference in sperm competitive ability between male types (Kasimatis *et al.*, 2018). Competitive fecundity success data were replotted from Kasimatis et al. (2018).

Fig S2. Immunostaining quantification development. A) Count data showing the percent of control males (JK574, triangle) and His-tagged males (VX300, circle) with fluorescence in each anatomical region during larval stage 4 (L4, orange), day 1 of adulthood (D1, green), day 2 of adulthood (D2, blue), and day 8 of adulthood (D8, pink). **C)** Count data showing the percent of control females (JK574, triangle) and His-tagged hermaphrodites (VX300, circle) with fluorescence in each anatomical region during larval stage 4 (L4, orange), day 1 of adulthood (D1, green), day 2 of adulthood (D2, blue), and day 8 of adulthood (D8, pink).

Fig S3. NSPF-1 localizes the MOs of spermatids in adult males. A) Protein (green) localized in a punctate pattern around the nucleus (blue) of spermatids in the dissected testes of adult His-tagged males. **B)** Protein localization within a single spermatid (outlined) of an adult His-tagged male. **C)** Protein localization within multiple spermatids (arrows) of an adult His-tagged male.

Table S1. Primers used in this study.
Table S2. Strains used in this study.

File S1. Late life fecundity data.
File S2. Processed amplicon reads.
File S3. Complete immunostaining protocols and buffer recipes.
File S4. Immunostaining data.