Title: Parvalbumin Neurons and Cortical Coding of Dynamic Stimuli: A Network Model

Jian Carlo Nocon[1,2,3,4], Isaac Paul Boyd[1,2,3,4], Howard Gritton[5,6], Xue Han[1,2,3,4], and Kamal
Sen[1,2,3,4,*]

[1]Neurophotonics Center, Boston University, Boston, Massachusetts, United States of America
02215,
[2]Center for Systems Neuroscience, Boston University, Boston, Massachusetts, United States of
America, 02215
[3]Hearing Research Center, Boston University, Boston, Massachusetts, United States of America,
02215
[4]Department of Biomedical Engineering, Boston University, Boston, Massachusetts, United
States of America, 02215
[5]Department of Comparative Biosciences, University of Illinois, Urbana, Illinois, United States
of America, 61820
[6]Department of Bioengineering, University of Illinois, Urbana, Illinois, United States of
America, 61820

[*]Corresponding Author: kamalsen@bu.edu

Abstract

Cortical circuits feature both excitatory and inhibitory cells that underlie the encoding of dynamic sensory stimuli, e.g., speech, music, odors, and natural scenes. While previous studies have shown that inhibition plays an important role in shaping the neural code, how excitatory and inhibitory cells coordinate to enhance encoding of temporally dynamic stimuli is not fully understood. Recent experimental recordings in mouse auditory cortex have shown that optogenetic suppression of parvalbumin neurons results in a decrease of neural discriminability of dynamic stimuli. Here, we present a multilayer model of a cortical circuit that mechanistically explains these results. The model is based on parvalbumin neurons which respond to both stimulus onsets and offsets, as observed experimentally, and incorporates characteristic short-term synaptic plasticity profiles of excitatory and parvalbumin neurons. We reveal that by tuning the relative strengths of onset and offset inputs to parvalbumin neurons, the model generates different regimes of coding dominated by rapid firing rate modulations or spike timing. Moreover, the model replicates the experimentally observed reduction in neural discrimination performance during optogenetic suppression of parvalbumin neurons. These results suggest that distinct onset and offset inputs to parvalbumin neurons enhance cortical discriminability of dynamic stimuli by encoding distinct temporal features, enhancing temporal coding, and reducing cortical noise.

New & Noteworthy

Here we propose a model for the mechanisms that underlie neuron responses in the auditory cortex. This study focuses on single channel tuning of simple interactions between neurons. Using physiologically relevant parameters that underly parvalbumin and excitatory neurons, in the proposed artificial network, we show that we can recreate observed results in live studies.

Key Words

Auditory cortex, cortical circuits, complex scene analysis, neural coding, parvalbumin

Introduction

Temporal processing of dynamic stimuli is critical for human perception of behaviorally relevant stimuli, e.g., speech, music, and natural scenes. While auditory cortex (ACx) is thought to contribute to such processing, the underlying cortical mechanisms remain poorly understood. In a recent study (1), we investigated the processing of dynamic stimuli in mouse ACx, quantifying neural discriminability of dynamic sounds in ACx, and found that optogenetic suppression of parvalbumin (PV) neurons degraded discrimination performance, and specifically temporal coding, in ACx. In this study, we present a network model to address how PV neurons shape cortical discrimination performance.

PV neurons in ACx have been found to respond to both onset and offset in sounds (2, 3), suggesting they are specialized for rapid encoding of stimulus boundaries (4). Additionally, previous models have shown that the short-term synaptic plasticity (STP) profiles of inhibitory interneurons are critical for temporal context-dependent modulation (5). Here, we implemented a multi-layer cortical network model with excitatory neurons, and PV neurons responding to both onset- and offsets to explain the experimental observations in (1). The model reveals that PV

66 neurons, and synaptic depression from PV to excitatory synapses are critical in sculpting cortical
67 responses to dynamic stimuli.
68
69 **Materials and Methods**
70
71   **Circuit structure overview.** The model network is composed of three layers: the first
72 layer representing pre-cortical input (IC, input cell), and the second (intermediate) and third
73 (output) layer both representing ACx. Both cortical layers consist of excitatory cells and
74 inhibitory parvalbumin-expressing (PV) cells.
75
Architecture. The input layer consists of one input channel, based on previous
experimental and modeling studies (6, 7). While previous experimental studies consisted of four
spatial locations, our focus in this study was to dissect the effects of "within-channel" inhibition-
mediated by PV neurons. Thus, we simplified the model architecture to focus on a single
channel, which contains two excitatory subchannels that respond to different temporal features,
i.e., onsets and offsets, of auditory stimuli. In each layer, both excitatory subchannels converge
on a parvalbumin neuron, which then inhibits the excitatory cells in the next layer. Our goals
were to match the discrimination performance and firing rate of excitatory neurons recorded in
mouse ACx (1).
85
86   **Stimuli.** For simulations of responses from (8), target stimuli consisted of two white
87 noise tokens modulated by human speech-shaped envelopes with a sampling frequency of 200
88 kHz.
89
90   **Input spike generation via spectro-temporal receptive fields.** Input responses were
91 generated from spectro-temporal receptive fields (STRFs). The STRFs for the input layer were
92 generated using the STRFlab Matlab toolbox (9) and modeled as the product of independent
93 spectral and temporal kernels:

$$STRF(t, f) = F(t) * G(f)$$

94   The temporal kernel was modeled as the difference of two linear functions (10, 11):

$$F(t) = e^{-\frac{t}{\alpha}} \left[\frac{C_1 \left(\frac{t}{\alpha}\right)^{n_1}}{n_1!} - \frac{C_2 \left(\frac{t}{\alpha}\right)^{n_2}}{n_2!} \right] \#(1)$$

C_1 and C_2 control the weights of the excitatory and inhibitory regions of the temporal kernel,
respectively. C_1 was fixed at 1, while C_2 was fixed at 0.88 to allow for a broader spiking profile.
α, which controls the timescale of the kernel, was fixed at 9.7 ms, while n_1 was fixed at 5 and n_2
was fixed at 8. Using this kernel allowed us to capture the temporal profile of STRFs measured
in the mouse medial geniculate body (12).
The spectral kernel of the STRF was modeled as a 2D Gabor filter (13):

$$G(f) = e^{-\left[\frac{(f-f_0)}{\sqrt{2}\sigma_f}\right]^2} \cos\left(2\pi * \Omega_f * (f - f_0)\right) \#(2)$$

The frequency range is determined by the best frequency f_0, the spectral bandwidth σ_f, and
modulation frequency Ω_f. Respectively, these parameters were fixed at 4300 Hz, 2000 Hz, and
50 μs to generate a broadband STRF for all simulations.

The convolution of the STRF and stimulus spectrograms generated a firing rate profile,
which was used to drive a Poisson spike generator. Refractoriness was modeled using a
sigmoidal recovery function (14):

$$w(\Delta) = \begin{cases} \dfrac{(\Delta - t_a)^g}{(\Delta - t_a)^g + t_r^g} & for\ \Delta > t_a \\ 0 & for\ \Delta \leq t_a \end{cases} \#(3)$$

107   Absolute refractory period (τ_a) was fixed at 1 ms, the relative refractory period (τ_r) was fixed at 1
108   ms, and the timescale of the recovery function (γ) was fixed at 2 ms. For each of the two target
109   stimuli, 10 trials were generated, each using a novel random token for Poisson spike generation.
110   To model responses to offsets, the convolution between the STRF and stimuli was
111   inverted such that the minimum and maximum values were switched:

$$FR_{off}(t) = -FR_{on}(t) + \max(FR_{on(t)}) \#(4)$$

112   A similar scheme has been used in previous models to approximate offset responses in auditory
113   cortex (15). To prevent evoked spiking at offset cells before stimulus playback, FR_{off} was set to
114   zero for all times before its first zero-crossing. Finally, FR_{off} was thresholded such that all
115   negative values were set to zero.
116
117   **Model structure and parameters.** We utilized the DynaSim toolbox (16) to implement
118   the model of mouse ACx. Neurons were modeled as leaky integrate-and-fire units with
119   membrane voltage V:

$$\frac{dV}{dt} = \frac{[(E_L - V) - Ri_{syn} + RI_{tonic}]}{RC} \#(5)$$

where E_L is the equilibrium potential, R is the membrane resistance, i_{syn} is the synaptic input
current, I_{tonic} is applied tonic current, and C is the membrane capacitance. Universal parameters,
such as capacitance and threshold voltage, are given in Table 1. Membrane resistances were
varied between cell types to approximate the differences in integration time constants (Table 2),
with PV cells having a faster membrane time constant than that of excitatory cells (5).
Additionally, resting membrane potentials for PV cells were set higher than those of excitatory
cells to ensure more rapid firing. When membrane voltage passed threshold, voltage was shunted
to V_{reset} for an absolute refractory period t_{ref}. In excitatory cells, spike-rate adaptation was also
implemented (17):

$$\frac{dg_{sra}}{dt} = -\frac{g_{sra}}{\tau_{sra}} \#(6)$$

$$g_{sra} \rightarrow g_{sra} + \Delta g_{sra} \#(7)$$

After a post-synaptic spike, adaptation conductance increases by Δg_{sra} and recovers to zero with
recovery time τ_{sra} (18). We found that implementing refractoriness and spike-rate adaptation in
this manner allowed the model to match the firing rates modulations in experimental data.
To simulate network noise, the output excitatory cell received a Poisson-like noise input
with a refractory period of 1 ms and a firing rate of 8 Hz, which matched the average
spontaneous activity in experimental data. To decrease simulation time, network noise was not
modeled using a separate unit; rather, tokens for Poisson spike times were generated before
integration of cell voltage.
137
Synaptic connections. Throughout simulations, certain synaptic weights were fixed.
Within the circuit, the synaptic connectivity between excitatory cells (IC->E_{Int}, E_{Int}->E_{Out}) was

kept constant. Additionally, the synaptic conductance of the inputs to PV cells (IC->PV$_{Int}$ and
E$_{Int}$->PV$_{Out}$) was fixed to ensure that a PV neuron would fire around 3 ms before an excitatory
cell receiving the same unitary input to approximate spike-time latencies observed in mouse
auditory cortex (19). Additionally, all synapses featured short-term synaptic depression, based on
previous models (20).
Post-synaptic currents (PSCs) were modeled using the difference of two decaying
exponential functions with time constants τ_1 and τ_2, with $\tau_1 > \tau_2$:

$$s(t) = B\left(e^{-\frac{t}{\tau_1}} - e^{-\frac{t}{\tau_2}}\right)u(t) \#(8)$$

147   where B is a normalization constant such that s = 1 at maximum, and u(t) is the unit step. For
148   simulations, s was rewritten as two ordinary differential equations:

$$\frac{ds}{dt} = \frac{\left(\frac{\tau_2}{\tau_1}\right)^{\frac{\tau_1}{\tau_2-\tau_1}} * x - s}{\tau_2} \#(9)$$

$$\frac{dx}{dt} = -\frac{x}{\tau_1} \#(10)$$

149   where x defines spike inputs to the post-synaptic cell. When a pre-synaptic spike is detected, x is
150   increased by synaptic strength P. P is then decreased by a fraction (f$_P$) of its current value (20):

$$x \rightarrow x + P \#(11)$$

$$P \rightarrow (1 - f_P)P \#(12)$$

151   In the absence of pre-synaptic spikes, P recovers to 1 with time constant τ_P:

$$\frac{dP}{dt} = -\frac{(1-P)}{\tau_P} \#(13)$$

152   The full equation for post-synaptic currents is given as:

$$i_{syn}(V, i) = g_{syn} * s * (V - E_{syn}) * netcon \#(14)$$

where g$_{syn}$ is synaptic conductance; E$_{syn}$ is reversal potential; and *netcon* is a binary matrix of
network connections between cell populations, with rows representing sources and columns
representing targets.
Values of τ_1, τ_2, and E$_{syn}$ were based on recorded excitatory and inhibitory PSCs in
mouse cortex (19) and kept constant throughout all simulations (21). Table 3 outlines the
parameters for all synaptic connections.
159
Biological rationale. STRFs at the IC were based on experimentally recorded data in
mouse auditory neurons (12). Here, we start with the assumption that the model contains two
excitatory subchannels that capture different features of auditory stimuli, with one responding to
sound onsets and the other responding to sound offsets consistent with past studies that have
reported parallel processing of onset and offset-responding cells in mouse auditory cortex (3).
We model PV neurons as having both onset and offset components, the most frequently observed
finding experimentally (2). Previous studies have shown the existence of both excitatory and
inhibitory neurons within cortex (22–24), and other studies have found that PV cells encode gaps
between bursts of white noise in auditory cortex (4).
169
Calculating discrimination performance using spike distance measures. Neural
discrimination performance was calculated using a template-matching approach from previous
experimental and modeling studies (6, 7). Briefly, spike trains were classified to one of the two

target stimuli based on whose template, one from each stimulus, yielded a smaller spike distance.
For each simulation set of 20 trials, 100 iterations of template matching were done, in which one
of the 10 spike trains for each target was chosen as a template, and all remaining trials were
matched to each template to determine target identity. All possible pairs of templates were used
across the 100 iterations to calculate an average value of performance. To ensure that
discrimination performance was only based on spiking during stimulus playback, spiking
between the simulation times of 300 ms and 3300 ms was used for analysis. This approach
utilized spike train distances as inputs, and several measures were used, each of which capture
different features of responses: SPIKE-distance; ISI-distance; rate-independent (RI) SPIKE-
distance. Previous analysis also utilized spike count distance, the difference in number of spikes
between trains, but discriminability based on this measure was close to chance for most data.
Thus, spike count distance-based performance was excluded from this study.
SPIKE-distance (25, 26) calculates the dissimilarity between two spike trains based on
differences in spike timing and instantaneous firing rate without additional parameters. For one
spike train in a pair, the instantaneous spike timing difference at time t is:

$$S_1(t) = \frac{\Delta t_P^{(1)}(t)x_F^{(1)} + \Delta t_F^{(1)}(t)x_P^{(1)}}{x_{ISI}^{(1)}(t)}, t_P^{(1)} \leq t \leq t_F^{(1)} \#(15)$$

where Δt_P represents the distance between the preceding spike from train 1 ($t_P^{(1)}$) and the nearest
spike from train 2, Δt_F represents the distance between the following spike from train 1 ($t_F^{(1)}$) and
the nearest spike from train 2, x_F is the absolute difference between t and $t_F^{(1)}$, and x_P is the
absolute difference between t and $t_P^{(1)}$. To calculate $S_2(t)$, the spike timing difference from the
view of the other train, all spike times and ISIs are replaced with the relevant values in train 2.
The pairwise instantaneous difference between the two trains is calculated as:

$$S''(t) = \frac{S_1(t) + S_2(t)}{2\langle x_{ISI}^1(t), x_{ISI}^2(t)\rangle} \#(16)$$

194   Finally, $S_1(t)$ and $S_2(t)$ are locally weighted by their instantaneous spike rates to account
195   for differences in firing rate, and the total distance between the two spike trains is the integral of
196   the profile:

$$S(t) = \frac{S_1(t)x_{ISI}^2(t) + S_2(t)x_{ISI}^1(t)}{2\langle x_{ISI}^1(t), x_{ISI}^2(t)\rangle^2} \#(17)$$

$$D_S = \frac{1}{T}\int_0^T S(t)dt \#(18)$$

197   ISI-distance calculates the dissimilarity between two spike trains based on differences in
198   instantaneous rate synchrony. For a given time point:

$$I(t) = \frac{\left|x_{ISI}^{(1)}(t) - x_{ISI}^{(2)}(t)\right|}{\max\left(x_{ISI}^{(1)}(t), x_{ISI}^{(2)}(t)\right)} \#(19)$$

This profile is then integrated along the spike train length to give a distance value, with values of
0 obtained for either identical spike trains or pairs with the same constant firing rate and a global
phase shift difference.
The RI-SPIKE-distance is rate-independent, as it does not take differences in local firing
rate between the two spike trains into account. From SPIKE-distance calculations, the final step
of weighing $S_1(t)$ and $S_2(t)$ by their instantaneous spike rates is skipped, yielding:

$$S_{1,2}^{RI}(t) = \frac{S_1(t) + S_2(t)}{2\langle x_{ISI}^1(t), x_{ISI}^2(t)\rangle} \#(20)$$

Like the other measures, the dissimilarity profile is integrated to give a distance value, with a
value of 0 obtained for two identical spike trains. cSPIKE was used to calculate SPIKE-, ISI-,
and RI-SPIKE distances between all possible spike train pairs (27).

Parameter searches for synaptic depression strengths. To determine how the
dynamics of synaptic depression affect spiking activity within the circuit, f_P was varied for each
synapse type (E->E, PV->E, and E->PV). For all simulations, all three distance-based
performance values at the output excitatory unit were calculated, along with the average firing
rate during stimulus playback. The values of f_P that yielded firing and discriminability values
approximating those seen in experimental data were then chosen for the rest of the simulations in
this study.

Optogenetic suppression of PV neurons. To replicate the experimental effect of
optogenetic suppression of PV cells, all PV neurons received a hyperpolarizing current of 0.03
nA, and the mean firing rate of network noise inputs to the output excitatory unit was increased
to 30 Hz to match the rise in spontaneous activity during suppression. The first effect was
modeled to replicate the effect of increased spiking across all cells within experimentally
recorded data, while the second effect captured the downstream effects of increased noise via
network perturbation. To account for the variability from Poisson-like noise at output, we ran 30
simulations of the same set of 20 trials (10 per target stimulus) for each condition.

Results

Mouse ACx units exhibit high neural discrimination of temporally dynamic stimuli.
To better understand cortical coding of complex scenes in a mouse model amenable to circuit
manipulation using genetic tools, we recently utilized a cocktail party-like experimental
paradigm (6) while recording from neurons in ACx (Figure 1A) (1). Specifically, we recorded
responses to spatially distributed sound mixtures to determine how competing sound sources
influence cortical coding of stimuli. The recording configuration consisted of 4 speakers
arranged in horizontal space around the mouse at 4 locations on the azimuthal plane: directly in
front (0º), two contralateral (90º and 45º) and one ipsilateral (-90º) to the right auditory cortex
recording area. Two target stimuli were used, each consisting of a white noise token modulated
by human speech-shaped envelopes from a speech corpus (28).
Figure 1B illustrates responses from an example single-unit cell during target-only
conditions. Here, the spike rasters and peri-stimulus time histograms (PSTHs) follow the
amplitude envelope of each target stimulus. To quantify the discriminability between the two
target stimuli, we utilized a template matching-based classifier (see: Materials and Methods,
'Calculating discrimination performance using spike distance measures'). The input to this
classifier was one of several spike train distance measures: SPIKE-distance, which measures
differences in spike timing and firing rate modulation between trains (27); inter-spike interval
(ISI)-distance, which only measures differences in firing rate modulation; rate-independent (RI)-
SPIKE-distance, which measures differences in spike timing while being independent of firing
rate differences. We focused on a subset of single-unit neurons that exhibited high SPIKE-
distance-based performance (above 70%) at one or more locations in the spatial grid that we

refer to as hotspots. Across the population of hotspots, we found that SPIKE-distance-based
performance was highest, followed by ISI-distance-based performance and then RI-SPIKE-
distance-based performance (Figure 1C).
To investigate the role of PV interneurons in auditory discrimination performance, we
compared discrimination performance at these hotspots with and without optogenetic
suppression of PV neurons in ACx. Upon optogenetic suppression, we found that performance
significantly decreased for all distance measures. These results indicate that optogenetic
suppression disrupts both timing and rate-based coding within ACx.
We organized hotspots based on the order of SPIKE-distance-, ISI-distance, and RI-
SPIKE-distance-based performances. In hotspots where firing rate-based (i.e., ISI-distance)
performance was lowest (Figure 2A), we found that spike timing-based (i.e., RI-SPIKE-distance)
performance primarily drove the high SPIKE-distance-based performance. When ACx was
optogenetically suppressed, both the overall and spike timing-based performance was reduced,
while firing rate-based performance remained unchanged. Hotspots where firing rate-based
distance was second highest (Figure 2B) or highest (Figure 2C) indicate values where high
SPIKE-distance-based performance was driven by firing rate modulations. Upon optogenetic
suppression, all three performance measures were degraded at these hotspots. Across all three
types of hotspots, we found those with higher rate-based performances tended to have higher
firing rates during stimulus playback, which increased during optogenetic suppression (Figure
2D).
Because the average SPIKE-distance-based performance was similar across all types, we
calculated the ratio of spike timing-based performance to firing rate-based performance for each
hotspot (Figure 2E). This measure allowed us to determine the extent to which SPIKE-distance-
based performance was based on either spike timing or firing rate modulation, with higher ratios
emphasizing the former.
274
Network Model: Distinct synaptic depression profiles across excitatory and PV
inhibitory synapses support high discrimination performance. We created a multi-layer
network model to replicate temporal responses to dynamic stimuli in mouse ACx neurons
(Figure 3A). The input layer consists of an onset cell and offset cell with Poisson firing with
absolute and relative refractory periods of 1ms. Within the cortical intermediate and output
layers, the excitatory cell from the preceding layer synapses onto a feed-forward inhibitory
circuit consisting of one excitatory cell (E) and one inhibitory PV cell (red) (Figure 3B). Table 1
shows the universal simulation parameters, while Table 2 shows the parameters for all cells in
this model.
In all synapses, short-term plasticity (STP) was implemented as previously described
(20). The model parameters are consistent with previous modeling studies in auditory cortex (5).
Specifically, all synapses in this circuit featured depression, the strength of which was mildest
between excitatory cells, followed by excitatory inputs to PV units, with the strongest depression
in synapses from PV to excitatory units (Figure 3C). Recovery time constants for each synapse
were chosen to match the firing rate modulations in experimental data. Table 3 shows the
parameters for all synapses in this circuit.
To investigate how changes in depression strengths affected model responses, we varied
the synaptic depression strength (f_P) in each synapse type (E->E, E->PV, and PV->E), ranging
from zero ($f_P = 0$) to maximal depression ($f_P = 1$). We first varied the depression strength
between excitatory cells (Figure 4A). We found that weak synaptic depression was necessary for

high firing rates and performance values at the output excitatory cell. When synaptic depression
strength increased, performance dropped towards chance values and firing at output dropped to
the network noise rate of 8 Hz.
When we varied the synaptic depression strength at excitatory inputs to PV cells (Figure
4B), we found that all performance values did not greatly change at higher depression strengths,
while firing rate greatly increased.
Finally, when we varied the synaptic depression strength of PV inhibition on excitatory
cells (Figure 4C), we found that SPIKE-distance-based performance plateaued at higher values
of f_P, while ISI-distance-based performance increased, and RI-SPIKE-distance-based
performance decreased. Additionally, firing rate steadily increased with synaptic depression
strength. From these simulations, we chose depression strengths, i.e., f_P values that were
consistent with the firing rates and performances that approximated the experimental single unit
data from mouse ACx, as marked by the horizontal dashed lines in Figure 4. Thus, these results
revealed that obtaining relatively high discrimination performance and firing rates consistent
with the data requires relatively strong depression in PV->E synapses, moderate depression in E-
>PV synapses and mild depression in E->E synapses.

Relative strengths of onset and offset component in PV neurons in model explains
diversity of discrimination profiles observed experimentally. To model the diversity of rate-
modulation hotspots (low, medium, and high) identified in the data, we next varied the strength
between the onset and offset components preceding PV neurons and the PV neurons in both
cortical layers while keeping all other circuit parameters constant. For hotspots with lower ratios
of spike timing to firing rate-based performances, the offset component of PV neurons was
reduced relative to the onset component. (Figure 5A).
Figure 5B illustrates the model's response to one of the two target stimuli shown in
Figure 1B. Throughout the ascending circuit, the firing rate pattern from each input is
maintained. In the model, output units exhibit periods of firing rate modulation where spiking is
close to zero during the quiet periods of the target stimulus.
We also found that this circuit effectively captures the pattern of firing rate-based versus
spike timing-based performance in high rate-based performance hotspots observed
experimentally (Figure 5C). For the simulations shown in 5B, ISI-distance-based performance
was higher than SPIKE-distance- and RI-SPIKE-distance-based performance was consistent with
the patterns observed in the high rate-based performance hotspots in the data (Figure 2C).
The model also captures the relative spike latencies of each cell type within cortex
(Figure 5D). Given the same input from IC, the onset component of PV neurons fires first, which
inhibits the excitatory PSC from the same spike input at E. These results are consistent with the
differences in receptive fields between excitatory and PV neurons in mouse ACx (19). Due to
synaptic depression, later spikes from the PV cell have a much weaker inhibitory effect on the
excitatory cell, thus allowing the E cell to fire during this period of weakened inhibition. The
descending edge of excitatory firing is then shaped by inhibition from the offset component in
PV neurons, which sharpens firing rate modulations (Figure 5E).
The parameter space underlying the relative offset to PV strength in neurons versus input
gain (Figure 5F), shows the distinction between the timing-based regime and the rate-based
regime. The graph displays the approximate ratio between the timing-based regime and the rate-
based regime (RI-SPIKE performance/ISI performance). A relatively high input gain and low
offset component results in a rate-modulation dominated regime of coding (blue region).

To capture hotspots with higher ratios of spike timing to firing rate-based performances,
we ran simulations where the offset component of PV neurons was higher than the onset
component (Figure 6A). Within experimental data, we found that these hotspots exhibited
responses where the noise floor firing rate during stimulus playback was lower than the
spontaneous firing rate. Thus, we hypothesized that increased inhibition during offset periods
facilitates higher spike timing-based performance. Figure 6B shows the responses to one of the
two target stimuli across all cells within the circuit. The output excitatory cell exhibited an
average firing rate of 20 Hz during stimulus playback, which approximates values seen in mouse
ACx. For these simulations, SPIKE-distance- and RI-SPIKE-distance-based performances were
almost equal, with ISI-distance-based performance being the lowest (Figure 6C), which agrees
with the trends observed in the low rate-based performance hotspots in the data (Figure 2A).
Voltage traces and firing rate modulation patterns at output neurons retain the same sharpening
mechanism seen in simulations of hotspots with high firing rate-based performances (Figures
6D-E). Figure 6F illustrates the range of parameter values for the spike timing-dominated regime
(highlighted region, 0.035 µS $(IC_{off}\text{->}PV)_{gsyn}$, 0.1 input gain), with relatively low input gain and
a relatively high offset component. Thus, the model reflects the diversity of firing rate-based
performance seen in experimental data and reveals that altering the relative onset and offset
components of PV neurons effectively captures the broad range of coding profiles seen in mouse
ACx.
360
Model explains experimentally observed effects of optogenetic suppression of PV
neurons. We next employed the model to simulate the effects of reduced PV signaling and
compared the results to empirical data collected in mouse ACx during optogenetic suppression of
PV cells. For the optogenetic condition, we inhibited PV neurons in all layers with a
hyperpolarizing current to model the elevated spontaneous firing rate observed in single unit
recordings (1), we increased the mean firing rate of the Poisson-like network noise to simulate
the increased excitability in the cortical network (Figure 7A).
In agreement with experimental results, we found that discrimination performance values
at the output unit decreased with optogenetic suppression for all spike distance measures (Figure
7B). Control simulations were found to approximate the firing rate profiles found within mouse
ACx, where the firing rate profile followed the temporal envelope of both target stimuli and
showed very little firing during the quiet periods of both targets (Figure 7C, top). During
simulations where PV neurons were suppressed, we saw a decrease in firing in all PV cells and a
slight increase in firing at both cortical excitatory units (Figure 7C, bottom). Notably, we found
that the noise floor was larger in optogenetic suppression trials than in the control condition
during stimulus playback (Figure 7D).
When we simulated optogenetic suppression for low firing rate populations that show
enhanced spike timing-based performance, the model replicated the experimental results across
all three measures of temporal and rate dependent discriminability. Specifically, optogenetic
suppression led to a decrease in both SPIKE-distance- and RI-SPIKE-distance-based
performances, while ISI-distance-based performance remained unchanged (Figure 8A), which
agrees with the data in Figure 2A. Like the simulations from the firing rate-based hotspot,
responses during the control condition were found to approximate those within mouse ACx
(Figure 8B, top). During optogenetic simulations, we saw the same increase in the noise floor
during stimulus playback as in the rate-based hotspot simulations (Figure 8B, bottom, and Figure

8C). Thus, the three-layer circuit model captures the experimentally observed reduction in neural
discriminability with optogenetic suppression of PV neurons in mouse ACx.
388
389 **Discussion**
390
Diverse neural discrimination performance profiles in ACx. In previous experimental
study examining the effects of optogenetically suppressing PV cells in mouse ACx, we found
that performance was degraded in terms of both spike timing, firing rate modulations, and a
combination of the two (1). In this study we further analyzed the degree to which high neural
discriminability was determined by either spike timing or firing rate modulations within each
hotspot. We then set out to determine if we could construct a model that effectively captures the
diversity of rate and temporal coding components seen in single unit data. In addition, we
focused on revealing the properties by which PV neurons could influence discriminability across
a range of hotspots present in mouse ACx.
400
Network model explains diversity of profiles. Previous computational studies of the
mouse auditory cortex primarily focused on modeling responses to transient stimuli such as tones
and white noise bursts (5, 29). Here, we sought to replicate responses to longer, and temporally
dynamic stimuli in mouse ACx. using experimentally recorded spectral temporal receptive fields
in mouse MGB (12). This input is then temporally sharpened by intra-cortical processing via
inhibition from PV interneurons.
One critical finding of the model is that the profile of synaptic depression parameters in
distinct synapses is critical for attaining high neural discriminability in auditory cortex.
Similarly, synaptic depression at inputs to PV cells and their inhibitory effect on excitatory cells
were both found to facilitate rapid modulations in firing rate within the circuit. Specifically, as
the thalamic input was processed in cortical layers, firing rate peaks were maintained, and offset
components of PV neurons enhanced suppression of the troughs in firing rate, thereby enhancing
firing rate modulations.
414
**Potential role of onset and offset components of PV interneurons in generating
diverse discrimination performance profiles and responses to stimulus envelopes.** We
demonstrated that the model circuit was able to capture the diversity of discriminability profiles
within auditory cortex by changing the relative strengths of onset and offset components of PV
neurons. When onset components were relatively high compared to offset, excitatory units
exhibited more prominent rate modulation dominated coding, i.e., low spike timing to firing rate-
based performance ratios. When offset components of PV neural responses were relatively
stronger compared to onset, the model captured the remaining set of hotspots with spike timing
dominated coding (high spike timing to firing rate-based performance ratios). Previous studies
have found that PV neurons are implicated in encoding gaps in noise within auditory cortex
which suggests an important contributions to spike timing (4). A critical feature in our model is
that PV neurons have both onset and offset components, based on experimental observations
from (2).These components may arise from distinct parallel onset and offset inputs originating in
the thalamus (3). Our model further elucidates the contribution of offset component in PV
neurons in enhancing temporal coding of dynamic stimuli.
430

Model limitations. The model in this study is a simplified version of cortical circuits, as
it only includes feed-forward circuits between excitatory and PV units. It does not account for
any recurrent inhibitory mechanisms between the two neuron types, nor does it include any other
interneuron type, such as somatostatin-expressing (SOM) or vasoactive intestinal peptide-
expressing (VIP) neurons, which might further facilitate the high neural discriminability seen in
cortex. Additionally, the model utilized single synapses over a population model in which
multiple inputs converge onto neurons. These simplifications allowed us to better dissect which
components of the cortical network are integral for capturing the variety of hotspots seen in
experimental data. The current results suggest an important role of onset and offset components
in PV neurons to temporally sharpen cortical responses and improve cortical discrimination
performance. Future models should incorporate other interneuron types to capture additional
aspects of cortical processing. For example, SOM neurons may be important to capture "cross-
channel" surround inhibition, and VIP neurons may be important for capturing top-down effects
on cortical processing (30).
In the model, the responses of offset neurons were modeled phenomenologically similar
to a previous study on auditory thalamic responses in mice (15). We found that this
approximation produced responses like the responses of offset neurons in the data. Other studies
have hypothesized that responses to stimulus offsets along auditory pathways are mediated by
hyperpolarization-activated current, amplified by downstream mechanisms, or inherited from
upstream regions (31, 32). Future studies should explore these mechanistic models of offset-
responding neurons.
Finally, while the model utilizes STP between all synapses, the parameters chosen to
model the temporal profile and strength of STP were selected to match the rapid firing rate
modulations noted in the experimental data. The recovery times and depression strengths that
were chosen for the circuit approximate the relative strengths of synaptic depression found
within cortex, with mild depression between excitatory inputs to relay cells and strong
depression in inhibitory inputs to excitatory cells (5). With these parameters, the model captured
both the neural discrimination performance profiles and mean firing rate values seen within the
experimental data.

Key model predictions and future directions. In this study, we presented a
computational model describing how PV-driven inhibition transforms thalamic inputs across
cortical layers to facilitates high neural discrimination performance within auditory cortex. The
model is based on physiological data in mice and posits that 1) cortical processing refines
stimulus encoding at the thalamic level as spiking propagates downstream, 2) spiking in PV
neurons are more phasic in time when compared to responses in excitatory neurons, 3) PV
neurons with both onset and offset components sharpen spiking activity in excitatory neurons,
thus leading to increased neural discriminability performance within cortex.
In this model, we focused on dissecting the rich temporal dynamics of "within-channel"
inhibition mediated by PV neurons, to replicate experimental data from clean trials in which only
a single stimulus was present. We did not model "cross-channel" interactions across spatial
channels, which may be critical for modeling the emergence of hotspots at different spatial
configurations. Based on the architecture of previous models (7, 30), future models of mouse
ACx should incorporate multiple spatial channels, with cross channel "surround" inhibition. The
analysis found that suppression of PV neurons reduced performance across all configurations of
target and masker location (1); thus, we predict that other interneurons types, e.g., SOM and VIP

neurons, may play an important role in determining the emergence of hotspots at different spatial configurations.

More generally, the modeling paradigm developed in this study may be useful in investigating the roles of different cortical interneuron types in complex scene analysis. The experimental data that were used to create and validate this model involved the optogenetic suppression of PV interneurons within mouse ACx. Future experiments with optogenetic manipulations or opto-tagging of other interneuron types within cortex, such as SOM or VIP neurons can be used to expand the model circuit. Ultimately, an understanding of the contributions of the diverse populations of interneurons in cortex, will be necessary to reveal cortical circuit mechanisms underlying complex scene analysis.

Grants

The research in this study was supported by the National Institutes of Health (#1R34NS111742-01) and the National Science Foundation (#IIS-1835270 and #2319321).

References

1.   **Nocon JC**, **Gritton HJ**, **James NM**, **Mount RA**, **Qu Z**, **Han X**, **Sen K**. Parvalbumin neurons enhance temporal coding and reduce cortical noise in complex auditory scenes. *Commun Biol* 6: 751, 2023. doi: 10.1038/s42003-023-05126-0.

2.   **Olsen T**, **Hasenstaub A**. Offset responses in the auditory cortex show unique history dependence. *J Neurosci* 42: 7370–85, 2022. doi: 10.1523/JNEUROSCI.0494-22.2022.

3.   **Liu J**, **Whiteway MR**, **Sheikhattar A**, **Butts DA**, **Babadi B**, **Kanold PO**. Parallel Processing of Sound Dynamics across Mouse Auditory Cortex via Spatially Patterned Thalamic Inputs and Distinct Areal Intracortical Circuits. *Cell Rep* 27: 872-885 e7, 2019. doi: 10.1016/j.celrep.2019.03.069.

4.   **Keller CH**, **Kaylegian K**, **Wehr M**. Gap encoding by parvalbumin-expressing interneurons in auditory cortex. *J Neurophysiol* 120: 105–114, 2018. doi: 10.1152/jn.00911.2017.

5.   **Seay MJ**, **Natan RG**, **Geffen MN**, **Buonomano DV**. Differential Short-Term Plasticity of PV and SST Neurons Accounts for Adaptation and Facilitation of Cortical Neurons to Auditory Tones. *J Neurosci* 40: 9224–9235, 2020. doi: 10.1523/JNEUROSCI.0686-20.2020.

6.   **Maddox RK**, **Billimoria CP**, **Perrone BP**, **Shinn-Cunningham BG**, **Sen K**. Competing sound sources reveal spatial effects in cortical processing. *PLoS Biol* 10: e1001319, 2012. doi: 10.1371/journal.pbio.1001319.

7.   **Dong J**, **Colburn HS**, **Sen K**. Cortical Transformation of Spatial Processing for Solving the Cocktail Party Problem: A Computational Model(1,2,3). *eNeuro* 3, 2016. doi: 10.1523/ENEURO.0086-15.2015.

8.   **Nocon JC**, **Gritton HJ**, **James NM**, **Mount RA**, **Qu Z**, **Han X**, **Sen K**. Parvalbumin neurons enhance temporal coding and reduce cortical noise in complex auditory scenes in mice. 2023.

9.   **Theunissen FE**, **David SV**, **Singh NC**, **Hsu A**, **Vinje WE**, **Gallant JL**. Estimating spatio-temporal receptive fields of auditory and visual neurons from their responses to natural stimuli. *Network* 12: 289–316, 2001.

10.   **Adelson EH**, **Bergen JR**. Spatiotemporal energy models for the perception of motion. *J Opt Soc Am A* 2: 284–99, 1985. doi: 10.1364/josaa.2.000284.

11.   **Dayan P**, **Abbott LF**. Space-Time Receptive Fields. In: *Theoretical neuroscience*, edited
by Sejnowski TJ, Poggio T. Cambridge, MA: The MIT Press, 2001, p. 68–69.
12.   **Lohse M**, **Bajo VM**, **King AJ**, **Willmore BDB**. Neural circuits underlying auditory
contrast gain control and their perceptual implications. *Nat Commun* 11: 324, 2020. doi:
10.1038/s41467-019-14163-5.
13.   **Qiu A**, **Schreiner CE**, **Escabi MA**. Gabor analysis of auditory midbrain receptive fields:
spectro-temporal and binaural composition. *J Neurophysiol* 90: 456–76, 2003. doi:
10.1152/jn.00851.2002.
14.   **Schaette R**, **Gollisch T**, **Herz AV**. Spike-train variability of auditory neurons in vivo:
dynamic responses follow predictions from constant stimuli. *J Neurophysiol* 93: 3270–81, 2005.
doi: 10.1152/jn.00758.2004.
15.   **Anderson LA**, **Linden JF**. Mind the Gap: Two Dissociable Mechanisms of Temporal
Processing in the Auditory System. *J Neurosci* 36: 1977–95, 2016. doi:
10.1523/JNEUROSCI.1652-15.2016.
16.   **Sherfey JS**, **Soplata AE**, **Ardid S**, **Roberts EA**, **Stanley DA**, **Pittman-Polletta BR**,
Kopell NJ. DynaSim: A MATLAB Toolbox for Neural Modeling and Simulation. *Front*
Neuroinform 12: 10, 2018. doi: 10.3389/fninf.2018.00010.
17.   **Fanselow EE**, **Connors BW**. The roles of somatostatin-expressing (GIN) and fast-
spiking inhibitory interneurons in UP-DOWN states of mouse neocortex. *J Neurophysiol* 104:
596–606, 2010. doi: 10.1152/jn.00206.2010.
18.   **Dayan P**, **Abbott LF**. Spike-Rate Adaptation and Refractoriness. In: *Theoretical*
neuroscience, edited by Sejnowski TJ, Poggio T. Cambridge, MA: The MIT Press, 2001, p. 165–
166.
19.   **Li LY**, **Xiong XR**, **Ibrahim LA**, **Yuan W**, **Tao HW**, **Zhang LI**. Differential Receptive
Field Properties of Parvalbumin and Somatostatin Inhibitory Neurons in Mouse Auditory Cortex.
Cereb Cortex 25: 1782–91, 2015. doi: 10.1093/cercor/bht417.
20.   **Dayan P**, **Abbott LF**. Release Probability and Short-Term Plasticity. In: *Theoretical*
neuroscience, edited by Sejnowski TJ, Poggio T. Cambridge, MA: The MIT Press, 2001, p. 184–
187.
21.   **Udakis M**, **Pedrosa V**, **Chamberlain SEL**, **Clopath C**, **Mellor JR**. Interneuron-specific
plasticity at parvalbumin and somatostatin inhibitory synapses onto CA1 pyramidal neurons
shapes hippocampal output. *Nat Commun* 11: 4395, 2020. doi: 10.1038/s41467-020-18074-8.
22.   **Pfeffer CK**, **Xue M**, **He M**, **Huang ZJ**, **Scanziani M**. Inhibition of inhibition in visual
cortex: the logic of connections between molecularly distinct interneurons. *Nat Neurosci* 16:
1068–76, 2013. doi: 10.1038/nn.3446.
23.   **Li LY**, **Ji XY**, **Liang F**, **Li YT**, **Xiao Z**, **Tao HW**, **Zhang LI**. A feedforward inhibitory
circuit mediates lateral refinement of sensory representation in upper layer 2/3 of mouse primary
auditory cortex. *J Neurosci* 34: 13670–83, 2014. doi: 10.1523/JNEUROSCI.1516-14.2014.
24.   **Xue M**, **Atallah BV**, **Scanziani M**. Equalizing excitation-inhibition ratios across visual
cortical neurons. *Nature* 511: 596–600, 2014. doi: 10.1038/nature13321.
25.   **Kreuz T**, **Chicharro D**, **Houghton C**, **Andrzejak RG**, **Mormann F**. Monitoring spike
train synchrony. *J Neurophysiol* 109: 1457–72, 2013. doi: 10.1152/jn.00873.2012.
26.   **Satuvuori E**, **Mulansky M**, **Daffertshofer A**, **Kreuz T**. Using spike train distances to
identify the most discriminative neuronal subpopulation. *J Neurosci Methods* 308: 354–365,
2018. doi: 10.1016/j.jneumeth.2018.09.008.

27.   **Satuvuori E**, **Kreuz T**. Which spike train distance is most suitable for distinguishing rate
and temporal coding? *J Neurosci Methods* 299: 22–33, 2018. doi:
10.1016/j.jneumeth.2018.02.009.
28.   **Rothauser EH**, **Chapman WD**, **Guttman N**, **Nordby KS**, **Silbiger HR**, **Urbanek GE**,
Weinstock M. I.E.E.E. recommended practice for speech quality measurements. *IEEE Trans
Audio Electroacoust* 17: 225–246, 1969. doi: 10.1109/TAU.1969.1162058.
29.   **Moore AK**, **Weible AP**, **Balmer TS**, **Trussell LO**, **Wehr M**. Rapid Rebalancing of
Excitation and Inhibition by Cortical Circuitry. *Neuron* 97: 1341-1355 e6, 2018. doi:
10.1016/j.neuron.2018.01.045.
30.   **Chou KF**, **Sen K**. AIM: A network model of attention in auditory cortex. *PLoS Comput
Biol* 17: e1009356, 2021. doi: 10.1371/journal.pcbi.1009356.
31.   **Kopp-Scheinpflug C**, **Tozer AJ**, **Robinson SW**, **Tempel BL**, **Hennig MH**, **Forsythe
ID**. The sound of silence: ionic mechanisms encoding sound termination. *Neuron* 71: 911–25,
2011. doi: 10.1016/j.neuron.2011.06.028.
32.   **Solyga M**, **Barkat TR**. Emergence and function of cortical offset responses in sound
termination detection. *Elife* 10, 2021. doi: 10.7554/eLife.72240.

584
585

586   **Acknowledgements:** We thank Michael Economo, Alberto Cruz-Martín, Monty Escabí, and
587   Kristina Penikis for helpful discussions and comments on the manuscript.
588
589   **Funding:** The work in this manuscript was funded by the National Science Foundation (#IIS-
590   1835270) and the National Institutes of Health and the National Institute of Neurological
591   Diseases and Stroke (#1R34NS111742-01).

592
593   **Conflicts of Interest:** The authors of this paper declare no conflicts of interest.
594

 Figures and Tables

597
Figure 1. Experimental data from mouse ACx. (A) Diagram of experimental setup and spatial
location of target stimuli. During trials, one of two target stimuli consisting of white noise
modulated by a human speech-shaped envelope was played from one of the four speakers along
the azimuth in horizontal space. **(B)** Responses to spatial configuration shown in A where the
two targets originated at 90°, from an example single unit in mouse ACx. Responses followed
the amplitude envelope of both targets, resulting in high SPIKE-distance-based neural
discrimination performance (92%) at this configuration. **(C)** Average discrimination performance
using different spike distance metrics across the population of 23 high-performing single units (n
= 49 hotspots during clean trials). Paired t-tests found significant differences between control and
optogenetic trials for all spike distance-based performances (SPIKE-distance: p = 8e-10; ISI-
distance: p = 3e-08; RI-SPIKE-distance: p = 1e-08).
609

610
Figure 2. Hotspots in single-unit experimental data reflect a broad range of timing and
rate-based contributions to performance. Hotspots in single-unit data were sorted by
comparing firing rate modulation-based (ISI-distance) performance to the other two measures
(SPIKE-distance, firing rate modulation and spike timing; RI-SPIKE-distance, spike timing
only). **(A)** Hotspots where ISI-distance-based performance was the worst out of all three
performance measures (n = 9), as indicated by the horizontal dashed line. **(B)** Hotspots where
ISI-distance-based performance was higher than RI-SPIKE-distance-based performance but not
SPIKE-distance-based performance (n = 15). **(C)** Hotspots where ISI-distance performance was
higher than the other two performance values (n = 25). **(D)** Auditory stimulus evoked average
firing rates during sound alone and PV optogenetic inhibition for each of the three hotspot types.
(E) Histogram showing all 49 hotspots from single-units, sorted by the ratio between RI-SPIKE-
distance-based performance and ISI-distance-based performance, with a ratio of 1 indicated by
the dashed line. Ratios lower than 1 indicate hotspots where high neural discrimination
performance is primarily driven by modulations in firing rate, rather than spike timing.
625

Figure 3. Computational model of auditory cortex layers and thalamic input. (A) Three-layer ACx model. The input stage of the model consists of a Poisson-like thalamic neuron with input from the inferior colliculus (IC) with both onset and offset type inputs. Firing rate of onset units is based on the convolution of an MGB STRF with stimulus spectrograms. Offset responses were generated by inverting the convolution between the STRF and stimulus and adding the absolute maximum value of the resulting firing rate trace. Both the intermediate (Int) and output (Out) layers exhibit feed-forward inhibitory circuits mediated by PV cells (red). PV neurons have both onset and offset components in their responses, the most frequent pattern observed experimentally (2). The excitatory cell in the output layer exhibited Poisson noise to simulate network noise in ACx. **(B)** Target 1 waveform with example FR trace from STRF convolution for onset- (red) and offset-responding (green) units. **(C)** Example traces of synaptic plasticity. Left: mild synaptic depression between excitatory cells. Right: synaptic depression in excitatory inputs to PV cells. Bottom: strong synaptic depression at inhibitory inputs to excitatory cells.

C [nF]	0.1
V_{thresh} [mV]	-47
I_{tonic} [nA]	0
Simulation time [ms]	3500
Integration time step [ms]	0.1
IC input gain	0.165

642 **Table 1. Universal DynaSim parameters.**

Unit	R (MΩ)	Δg_{sra} (nS/spk)	τ_{sra} (ms)	t_{ref} (ms)	E_L (mV)	V_{reset} (mV)
IC	200	0	--	1	-65	-54
E	200	0.3	100	1	-65	-54
PV	100	0	--	0.5	-57	-52

643   **Table 2. Unit parameters.** DynaSim parameters that differed between units.

				Nocon et al. (2023)		
Connection	τ_1 (ms)	τ_2 (ms)	E_{syn} (mV)	g_{syn} (nS)	f_P	τ_P (ms)
E->E	0.7	1.5	0	20	0.1	30
N->E$_{Out}$	0.7	1.5	0	15	0	--
On->PV	0.1	1	0	20	0.2	80
Off->PV	0.1	1	0	5	0	80
PV->E$_{On}$ PV->E$_{Off}$	1	4.5	-80	25	0.5	120

644 **Table 3. Synaptic parameters.** DynaSim parameters that differed between synapses. N
645 represents simulated network noise input from a Poisson unit in the output layer to model
646 network noise in ACx.
647

Figure 4. Effects of synaptic depression strengths on output discriminability and firing.
Simulations for a rate-based hotspot in output excitatory cells, where synaptic depression
strengths (f_P) were varied while keeping all other parameters constant. **(A)** Synaptic depression
strength between excitatory cells versus discriminability measures (top) and firing rate (bottom).
Simulations where $f_{P,E->E} = 0.1$ (indicated by the vertical dashed line) leads to SPIKE-distance-
based neural discriminability and firing rate values that approximate those of experimental data
in hotspots with high firing rate-based performances, both of which are indicated by horizontal
dashed lines in the upper and lower panels. **(B)** Synaptic depression strength between excitatory
inputs to PV cells versus discriminability measures (top) and firing rate (bottom). Simulations
where $f_{P,E->PV} = 0.2$ results in performance and firing rate values that approximate those of
experimental data. **(C)** Synaptic depression strength of inhibitory inputs to excitatory cells versus
discriminability measures and firing rate. Simulations where $f_{P,PV->E} = 0.5$ resulted in
performance and firing rate values that approximate those in hotspots with high firing rate-based
performance.

664
Figure 5. Hotspots with high firing rate-based performance can be modeled using PV
neurons with a relatively high onset component, (A) To simulate hotspots where firing rate-
based performance was higher than spike timing-based performance, inhibition strength from
offset-sensitive PV units (On->PV, 20 µS) was reduced relative to that of onset-sensitive PV
units (Off->PV, 5 µS). **(B)** Responses during the first second of one target stimulus (black
waveform, bottom) from each cell, with rasters plotted in gray and the corresponding PSTH
overlaid in the same colors as those shown in Figure 3A. To better show firing rate patterns in all
cells, PSTHs were smoothed using a moving-average filter spanning 3 samples. Scale bars at
bottom-right corner indicates vertical scale of PSTHs and horizontal time scale. Average firing
rate at the output excitatory cell was 32 Hz during stimulus playback. **(C)** Performance measures
of simulations shown in A. Here, ISI-distance-based performance is higher than RI-SPIKE-
distance-based performance, indicating that the high SPIKE-distance-based performance is
dominated by rapid changes in firing rate modulation. **(D)** Example voltage traces from output
layer neurons. Onset-sensitive PV neurons (top) fire first from IC inputs, which then leads to
inhibition of E cells (middle), thereby sculpting the ascending edge of firing rate modulations in
excitatory cells. The descending edge is sculpted by offset-sensitive PV cells (bottom). Scale
bars measure 20mV and 10ms. **(E)** Zoomed-in smoothed PSTHs (bin size 20ms) of output layer
cells during the first half second of sound onset. PSTHs of PV cells indicate how inhibition
shapes the firing rate patterns in excitatory cells. **(F)** Example parameter search showing
approximate RI-SPIKE/ISI performance values testing IC input gain vs (IC_{Off}->PV)g_{syn}. Outlined
section highlights ISI dominant region in parameter space (black outline). The color plot shows 7
by 7 zero-padded averaging convolution of a single parameter sweep (average of 49 cells per cell
shown). Values were first subtracted by the grid mean and then the mean was reintroduced after
convolution to reduce noise from zero padding.

Figure 6. Hotspots with high spike timing-based performance can be modeled using PV neurons with a relatively high offset component. (A) To simulate hotspots where spike timing-based performance was higher than firing rate-based performance, inhibition strength from offset-sensitive PV units (w_{Off}, green) was elevated relative to onset-sensitive PV units (w_{On}, red). **(B)** Responses during the first second of one target stimulus (black waveform, bottom) from each cell, with rasters plotted in gray and the corresponding smoothed PSTH overlaid in the same colors as those shown in Figure 3A. Scale bars at bottom-right corner indicates vertical scale of PSTHs and horizontal time scale. Average firing rate at the output excitatory cell was 20 Hz during stimulus playback. **(C)** Performance measures of simulations shown in A. Here, ISI-distance-based performance is lower than RI-SPIKE-distance-based performance, indicating that the high SPIKE-distance-based performance is dominated by spike timing comparisons between trains. **(D)** Example voltage traces from output layer neurons. Onset-sensitive PV neurons (top) fire first from IC inputs, which then leads to inhibition of E cells (middle), thereby sculpting the ascending edge of firing rate modulations in excitatory cells. The descending edge is sculpted by offset-sensitive PV cells (bottom). Scale bars measure 20mV and 10ms. **(E)** Zoomed-in smoothed PSTHs (bin size 20ms) of output layer cells during the first half second of sound onset. **(F)** E Example parameter search showing approximate RI-SPIKE/ISI performance values testing IC input gain vs (IC_{Off}->PV)g_{syn}. Outlined section highlights RI-SPIKE dominant region in parameter space (black outline). Convolution method is the same as described in Figure 5F.

709
Figure 7. Simulations of responses during optogenetic suppression for a hotspot with high
firing rate-based performance. For all simulations used in this figure, synaptic connectivity
strengths were the same as in Figure 5. **(A)** Diagram showing changes to circuit during
optogenetic suppression simulations. All PV cells are suppressed with a hyperpolarizing current
of 0.03 nA, and the network noise input to the output excitatory cell is increased from 8 Hz to 30
Hz. **(B)** Average performance values during control and optogenetic conditions across 30 sets of
simulations per condition, with error bars representing ± 1 standard deviation. The significance
between laser and control trials for all distance metrics (p < 0.001) was calculated using
Wilcoxon rank sum test. **(C)** Example rasters and PSTHs to each target stimulus during one
control and one optogenetic simulation. **(D)** Inset showing zoomed-in portion of the response to
target stimulus #1, between 0.6 and 1.8s after sound onset, as outlined in C. Responses during
optogenetic trials show a higher noise floor, compared to the control condition.

722
Figure 8. Simulations of responses during optogenetic suppression in a hotspot with high
spike timing-based performance. For all simulations used in this Figure, synaptic connectivity
strengths were the same as in Figure 6. **(A)** Average performance values during control and
optogenetic trials across 30 sets of simulations per condition, with error bars representing $\pm$ 1
standard deviation. The significance between laser and control trials for SPIKE and RI-SPIKE
distance ($p < 0.001$) and ISI distance ($p < 0.01$) was calculated using Wilcoxon rank sum test.
(B) Example rasters and PSTHs to each target stimulus during one control and one optogenetic
simulation. **(C)** Inset showing zoomed-in portion of the response to target stimulus #2, between
2.45s and 2.75s after sound onset, as outlined in B. Responses during optogenetic trials show a
higher noise floor, compared to the control condition.
733