Differential Inhibitory Responses to Temporal Features Enhance Cortical Coding of Dynamic Stimuli: A Network Model

Jian Carlo Nocon[1,2,3,4], Howard Gritton[5,6], Xue Han[1,2,3,4], and Kamal Sen[1,2,3,4,*]

[1]Neurophotonics Center, Boston University, Boston, Massachusetts, United States of America 02215,
[2]Center for Systems Neuroscience, Boston University, Boston, Massachusetts, United States of America, 02215
[3]Hearing Research Center, Boston University, Boston, Massachusetts, United States of America, 02215
[4]Department of Biomedical Engineering, Boston University, Boston, Massachusetts, United States of America, 02215
[5]Department of Comparative Biosciences, University of Illinois, Urbana, Illinois, United States of America, 61820
[6]Department of Bioengineering, University of Illinois, Urbana, Illinois, United States of America, 61820

[*]Corresponding Author: kamalsen@bu.edu

Abstract

Cortical circuits feature both excitatory and inhibitory cells that underlie the encoding of dynamic sensory stimuli, e.g., speech, music, odors, and natural scenes. While previous studies have shown that inhibition plays an important role in shaping the neural code, little is known about how excitatory and inhibitory cells coordinate to enhance encoding of temporally dynamic stimuli. Recent experimental recordings in mouse auditory cortex (ACx) have shown that optogenetic suppression of parvalbumin (PV) neurons results in a decrease of neural discriminability between dynamic stimuli with speech envelope modulations. Here, we present a multilayer model of a cortical circuit that mechanistically explains these results. The model incorporates characteristic short-term synaptic plasticity (STP) profiles of excitatory and PV cells. Crucially, the model performance is based on populations of PV cells that separately respond to stimulus onsets and offsets. We reveal that by tuning the relative strengths of inhibition from onset- and offset-responding PV cells, the cortical network model captures the broad range of neural discriminability profiles in cortical single-unit data, with varying contributions from rapid firing rate modulations and spike timing. The model also replicates and explains the experimentally observed reduction in neural discrimination performance during optogenetic suppression of PV neurons. These results suggest that distinct populations of PV neurons enhance cortical discriminability of dynamic stimuli by encoding distinct temporal features, enhancing temporal coding, and reducing cortical noise.

Introduction

Temporal processing of dynamic stimuli is critical for human perception of behaviorally relevant stimuli, e.g., speech, music, and natural scenes. Although auditory cortex (ACx) is thought to contribute to such processing, the underlying cortical mechanisms remain poorly understood. In a recent study (Nocon et al., 2022), we investigated the processing of dynamic stimuli in mouse ACx, quantifying neural discriminability of dynamic sounds in ACx, and found that optogenetic suppression of parvalbumin (PV) neurons degraded discrimination performance, and specifically temporal coding, in ACx. How do PV neurons shape cortical discrimination performance and temporal coding? In this study we present a network model to address this question.

PV neurons in ACx have been found to encode onset or offset responses (Liu et al., 2019), suggesting they are specialized for rapid encoding of stimulus boundaries (Keller et al., 2018). Additionally, previous models have shown that the short-term synaptic plasticity (STP) profiles of inhibitory interneurons are critical for temporal context-dependent modulation (Seay et al., 2020). Here, we implemented a multi-layer cortical network model with excitatory neurons, and both onset- and offset-responding PV neurons to explain the experimental observations in (Nocon et al., 2022). The model reveals that both onset and offset-responding PV neurons, and synaptic depression from PV to excitatory synapses are critical in sculpting cortical responses to dynamic stimuli. Moreover, offset PV neurons strongly contribute to the temporal coding component in cortical discrimination. The model identifies key candidate cortical mechanisms that support temporal processing of behaviorally relevant dynamic stimuli, e.g., speech and music.

Materials and Methods

Circuit structure overview

The model network is composed of three layers: the first layer representing pre-cortical input (IC, input cell), and the second intermediate and third output layer both representing ACx. Both cortical layers consist of excitatory cells and inhibitory parvalbumin-expressing (PV) cells.

Architecture

The input layer consists of one spatial input channel that correspond to center-preferring neurons, based on previous experimental and modeling studies (Maddox et al., 2012; Dong et al., 2016). While previous experimental studies consisted of four spatial locations, this study's focus was on dissecting the effects of "within-channel" inhibition-mediated by PV neurons. Thus, we simplified the model architecture to focus on one spatially tuned channel, which contains two subchannels that respond to different temporal features of auditory stimuli. The first goal was to match the discrimination performance and firing rate of excitatory neurons recorded in mouse ACx. The second goal was to capture the variety of spike timing-based and firing rate-based discriminability performances experimentally observed in ACx.

Generating input spikes

Stimuli

For all simulations, target stimuli consisted of two white noise tokens modulated by human speech-shaped envelopes with a sampling frequency of 195312 Hz. These same stimuli were used for previous experiments on complex scene analysis in mouse ACx (Nocon et al., 2022).

Spectro-temporal receptive fields

Input responses were generated from spectro-temporal receptive fields (STRFs). The STRFs for the input layer were generated using the STRFlab Matlab toolbox (Theunissen et al., 2001) and modeled as the product of independent spectral and temporal kernels:

$$STRF(t, f) = F(t) * G(f)$$

The temporal kernel was modeled as the difference of two linear functions (Adelson and Bergen, 1985; Dayan and Abbott, 2001c):

$$F(t) = e^{-\frac{t}{\alpha}} \left[\frac{C_1 \left(\frac{t}{\alpha}\right)^{n_1}}{n_1!} - \frac{C_2 \left(\frac{t}{\alpha}\right)^{n_2}}{n_2!} \right]$$

C_1 and C_2 control the weights of the excitatory and inhibitory regions of the temporal kernel, respectively. C_1 was fixed at 1, while C_2 was fixed at 0.88 to allow for a broader spiking profile. α, which controls the timescale of the kernel, was fixed at 9.7 ms, while n_1 was fixed at 5 and n_2 was fixed at 7. Using this kernel allowed us to capture the temporal profile of STRFs measured in the mouse medial geniculate body (Lohse et al., 2020).

The spectral kernel of the STRF was modeled as a 2D Gabor filter (Qiu et al., 2003):

$$G(f) = e^{-\left[\frac{(f-f_0)}{\sqrt{2}\sigma_f}\right]^2} \cos\left(2\pi * \Omega_f * (f - f_0)\right)$$

The frequency range is determined by the best frequency f_0, the spectral bandwidth σ_f, and
modulation frequency Ω_f. Respectively, these parameters were fixed at 4300 Hz, 2000 Hz, and
50 μs to generate a broadband STRF for all simulations.
The convolution of the STRF and stimulus spectrograms generated a firing rate profile,
which was used to drive a Poisson spike generator. Refractoriness was modeled using a
sigmoidal recovery function (Schaette et al., 2005):

$$w(\Delta) = \begin{cases} \dfrac{(\Delta - \tau_a)^\gamma}{(\Delta - \tau_a)^\gamma + \tau_r^\gamma} & for\ \Delta > \tau_a \\ 0 & for\ \Delta \le \tau_a \end{cases}$$

108   Absolute refractory period (τ_a) was fixed at 1 ms, the relative refractory period (τ_r) was fixed at 1
109   ms, and the timescale of the recovery function (γ) was fixed at 2 ms. For each of the two target
110   stimuli, 10 trials were generated, each using a novel random token for Poisson spike generation.
111   To model responses to offsets, the convolution between the STRF and stimuli was
112   inverted such that the minimum and maximum values were switched:

$$FR_{off}(t) = -FR_{on}(t) + \max\left(FR_{on(t)}\right)$$

113   A similar scheme has been used in previous models to approximate offset responses in auditory
114   cortex (Anderson and Linden, 2016). To prevent evoked spiking at offset cells before stimulus
115   playback, FR_{off} was set to zero for all times before its first zero-crossing. Finally, FR_{off} was
116   thresholded such that all negative values were set to zero.
117

Model structure and parameters
Model neurons
120   We utilized the DynaSim toolbox (Sherfey et al., 2018) to implement the model of mouse
121   ACx. Neurons were modeled as leaky integrate-and-fire units with membrane voltage V:

$$\frac{dV}{dt} = \frac{\left[(E_L - V) - Ri_{syn} + RI_{tonic}\right]}{RC}$$

where E_L is the equilibrium potential, R is the membrane resistance, i_{syn} is the synaptic input
current, I_{tonic} is applied tonic current, and C is the membrane capacitance. Universal parameters,
such as capacitance and threshold voltage, are given in Table 1. Membrane resistances were
varied between cell types to approximate the differences in integration time constants (Table 2),
with PV cells having a faster membrane time constant than that of excitatory cells (Seay et al.,
2020). Additionally, resting membrane potentials for PV cells were set higher than those of
excitatory cells to ensure more rapid firing. When membrane voltage passed threshold, voltage
was shunted to V_{reset} for an absolute refractory period t_{ref}. In excitatory cells, spike-rate
adaptation was also implemented (Fanselow and Connors, 2010):

$$\frac{dg_{sra}}{dt} = -\frac{g_{sra}}{\tau_{sra}}$$

$$g_{sra} \to g_{sra} + \Delta g_{sra}$$

After a post-synaptic spike, adaptation conductance increases by Δg_{sra} and recovers to zero with
recovery time τ_{sra} (Dayan and Abbott, 2001b). We found that implementing refractoriness and
spike-rate adaptation in this manner allowed the model to match the firing rates modulations in
experimental data.
To simulate network noise, the output excitatory cell received a Poisson-like noise input
with a refractory period of 1 ms and a firing rate of 8 Hz, which matched the average
spontaneous activity in experimental data. To decrease simulation time, network noise was not

modeled using a separate unit; rather, tokens for Poisson spike times were generated before
integration of cell voltage.
140

Synaptic connections

Throughout simulations, certain synaptic weights were fixed. Within the circuit, the
synaptic connectivity between excitatory cells (IC->E_{Int}, E_{Int}->E_{Out}) was kept constant.
Additionally, the synaptic conductance of the inputs to PV cells (IC->PV_{Int} and E_{Int}->PV_{Out}) was
fixed to ensure that a PV neuron would fire around 3 ms before an excitatory cell receiving the
same unitary input to approximate spike-time latencies observed in mouse auditory cortex (Li et
al., 2015). Additionally, all synapses featured short-term synaptic depression, based on previous
models (Dayan and Abbott, 2001a; Seay et al., 2020).
Post-synaptic currents (PSCs) were modeled using the difference of two decaying
exponential functions with time constants τ_1 and τ_2, with $\tau_1 > \tau_2$:

$$s(t) = B\left(e^{-\frac{t}{\tau_1}} - e^{-\frac{t}{\tau_2}}\right)u(t)$$

151 where B is a normalization constant such that s = 1 at maximum, and u(t) is the unit step. For
152 simulations, s was rewritten as two ordinary differential equations:

$$\frac{ds}{dt} = \frac{\left(\frac{\tau_2}{\tau_1}\right)^{\frac{\tau_1}{\tau_2-\tau_1}} * x - s}{\tau_2}$$
$$\frac{dx}{dt} = -\frac{x}{\tau_1}$$

153 where x defines spike inputs to the post-synaptic cell. When a pre-synaptic spike is detected, x is
154 increased by synaptic strength P. P is then decreased by a fraction (f_P) of its current value (Dayan
155 and Abbott, 2001a):

$$x \rightarrow x + P$$
$$P \rightarrow (1 - f_P)P$$

156 In the absence of pre-synaptic spikes, P recovers to 1 with time constant τ_P:

$$\frac{dP}{dt} = -\frac{(1-P)}{\tau_P}$$

157 The full equation for post-synaptic currents is given as:

$$i_{syn}(V, i) = g_{syn} * s * \left(V - E_{syn}\right) * netcon$$

where g_{syn} is synaptic conductance; E_{syn} is reversal potential; and *netcon* is a binary matrix of
network connections between cell populations, with rows representing sources and columns
representing targets.
Values of τ_1, τ_2, and E_{syn} were based on recorded excitatory and inhibitory PSCs in
mouse cortex (Li et al., 2015) and kept constant throughout all simulations (Udakis et al., 2020).
Table 3 outlines the parameters for all synaptic connections.
164

Biological rationale

STRFs at the IC were based on experimentally recorded data in mouse auditory neurons
(Lohse et al., 2020). Here, we start with the assumption that one spatially tuned channel contains
two subchannels that capture different features of auditory stimuli, with one responding to sound
onsets and the other responding to sound offsets. Past studies have reported parallel processing
of onset and offset-responding cells in mouse auditory cortex (Liu et al., 2019). We also assume

that these channels are independent of each other before entering ACx, and that these
interactions between onset and offset channels are mediated via PV cells. Previous studies have
shown the existence of both excitatory and inhibitory neurons within cortex (Pfeffer et al., 2013;
Li et al., 2014; Xue et al., 2014), and other studies have found that PV cells encode gaps between
bursts of white noise in auditory cortex (Keller et al., 2018).
176

Data analysis

Calculating discrimination performance using spike distance measures

Neural discrimination performance was calculated using a template-matching approach
from previous experimental and modeling studies (Maddox et al., 2012; Dong et al., 2016).
Briefly, spike trains were classified to one of the two target stimuli based on whose template, one
from each stimulus, yielded a smaller spike distance. For each simulation set of 20 trials, 100
iterations of template matching were done, in which one of the 10 spike trains for each target was
chosen as a template, and all remaining trials were matched to each template to determine target
identity. All possible pairs of templates were used across the 100 iterations to calculate an
average value of performance. To ensure that discrimination performance was only based on
spiking during stimulus playback, spiking between the simulation times of 300 ms and 3300 ms
was used for analysis. This approach utilized spike train distances as inputs, and several
measures were used, each of which capture different features of responses: SPIKE-distance; ISI-
distance; rate-independent (RI) SPIKE-distance. Previous analysis also utilized spike count
distance, the difference in number of spikes between trains, but discriminability based on this
measure was close to chance for most data. Thus, spike count distance-based performance was
excluded from this study.
SPIKE-distance (Kreuz et al., 2013; Satuvuori et al., 2018) calculates the dissimilarity
between two spike trains based on differences in spike timing and instantaneous firing rate
without additional parameters. For one spike train in a pair, the instantaneous spike timing
difference at time t is:

$$S_1(t) = \frac{\Delta t_P^{(1)}(t)x_F^{(1)} + \Delta t_F^{(1)}(t)x_P^{(1)}}{x_{ISI}^{(1)}(t)}, t_P^{(1)} \le t \le t_F^{(1)}$$

where Δt_P represents the distance between the preceding spike from train 1 ($t_P^{(1)}$) and the nearest
spike from train 2, Δt_F represents the distance between the following spike from train 1 ($t_F^{(1)}$) and
the nearest spike from train 2, x_F is the absolute difference between t and $t_F^{(1)}$, and x_P is the
absolute difference between t and $t_P^{(1)}$. To calculate $S_2(t)$, the spike timing difference from the
view of the other train, all spike times and ISIs are replaced with the relevant values in train 2.
The pairwise instantaneous difference between the two trains is calculated as:

$$S''(t) = \frac{S_1(t) + S_2(t)}{2\langle x_{ISI}^1(t), x_{ISI}^2(t)\rangle}$$

204 Finally, $S_1(t)$ and $S_2(t)$ are locally weighted by their instantaneous spike rates to account
205 for differences in firing rate, and the total distance between the two spike trains is the integral of
206 the profile:

$$S(t) = \frac{S_1(t)x_{ISI}^2(t) + S_2(t)x_{ISI}^1(t)}{2\langle x_{ISI}^1(t), x_{ISI}^2(t)\rangle^2}$$

$$D_S = \frac{1}{T}\int_0^T S(t)dt$$

ISI-distance calculates the dissimilarity between two spike trains based on differences in instantaneous rate synchrony. For a given time point:

$$I(t) = \frac{\left| x_{ISI}^{(1)}(t) - x_{ISI}^{(2)}(t) \right|}{\max\left(x_{ISI}^{(1)}(t), x_{ISI}^{(2)}(t) \right)}$$

This profile is then integrated along the spike train length to give a distance value, with values of 0 obtained for either identical spike trains or pairs with the same constant firing rate and a global phase shift difference.

The RI-SPIKE-distance is rate-independent, as it does not take differences in local firing rate between the two spike trains into account. From SPIKE-distance calculations, the final step of weighing $S_1(t)$ and $S_2(t)$ by their instantaneous spike rates is skipped, yielding:

$$S_{1,2}^{RI}(t) = \frac{S_1(t) + S_2(t)}{2\langle x_{ISI}^1(t), x_{ISI}^2(t) \rangle}$$

Like the other measures, the dissimilarity profile is integrated to give a distance value, with a value of 0 obtained for two identical spike trains. cSPIKE was used to calculate SPIKE-, ISI-, and RI-SPIKE distances between all possible spike train pairs (Satuvuori and Kreuz, 2018).

Simulation types

Parameter searches for synaptic depression strengths

To determine how the dynamics of synaptic depression affect spiking activity within the circuit, f_P was varied for each synapse type (E->E, PV->E, and E->PV). For all simulations, all three distance-based performance values at the output excitatory unit were calculated, along with the average firing rate during stimulus playback. The values of f_P that yielded firing and discriminability values approximating those seen in experimental data were then chosen for the rest of the simulations in this study.

Optogenetic suppression of PV neurons

To replicate the experimental effect of optogenetic suppression of PV cells, all PV neurons received a hyperpolarizing current of 0.03 nA, and the mean firing rate of network noise inputs to the output excitatory unit was increased to 12 Hz to match the rise in spontaneous activity during suppression. The first effect was modeled to replicate the effect of increased spiking across all cells within experimentally recorded data, while the second effect captured the downstream effects of increased noise via network perturbation. To account for the variability from Poisson-like noise at output, we ran 5 simulations of the same set of 20 trials (10 per target stimulus) for each condition.

Results

Mouse ACx units exhibit high neural discrimination of temporally dynamic stimuli

To better understand cortical coding of complex scenes in a mouse model amenable to circuit manipulation using genetic tools, we recently utilized a cocktail party-like experimental paradigm (Maddox et al., 2012) while recording from neurons in ACx (Figure 1A) (Nocon et al.,

2022). Specifically, we recorded responses to spatially distributed sound mixtures to determine
how competing sound sources influence cortical coding of stimuli. The recording configuration
consisted of 4 speakers arranged in horizontal space around the mouse at 4 locations on the
azimuthal plane: directly in front (0º), two contralateral (90º and 45º) and one ipsilateral (-90º) to
the right auditory cortex recording area. Two target stimuli were used, each consisting of a white
noise token modulated by human speech-shaped envelopes from a speech corpus (Rothauser,
1969).
Figure 1B illustrates responses from an example single-unit cell during target-only
conditions. Here, the spike rasters and peri-stimulus time histograms (PSTHs) follow the
amplitude envelope of each target stimulus. To quantify the discriminability between the two
target stimuli, we utilized a template matching-based classifier (see: Materials and Methods,
'Calculating discrimination performance using spike distance measures'). The input to this
classifier was one of several spike train distance measures: SPIKE-distance, which measures
differences in spike timing and firing rate modulation between trains (Satuvuori and Kreuz,
2018); inter-spike interval (ISI)-distance, which only measures differences in firing rate
modulation; rate-independent (RI)-SPIKE-distance, which measures differences in spike timing
while being independent of firing rate differences. We focused on a subset of single-unit neurons
that exhibited high SPIKE-distance-based performance (above 70%) at one or more locations in
the spatial grid that we refer to as hotspots. Across the population of hotspots, we found that
SPIKE-distance-based performance was highest, followed by ISI-distance-based performance
and then RI-SPIKE-distance-based performance (Figure 1C).
To investigate the role of PV interneurons in auditory discrimination performance, we
compared discrimination performance at these hotspots with and without optogenetic
suppression of PV neurons in ACx. Upon optogenetic suppression, we found that performance
significantly decreased for all distance measures. These results indicate that optogenetic
suppression disrupts both timing and rate-based coding within ACx.
We organized hotspots based on the order of SPIKE-distance-, ISI-distance, and RI-
SPIKE-distance-based performances. In hotspots where firing rate-based (i.e., ISI-distance)
performance was lowest (Figure 2A), we found that spike timing-based (i.e., RI-SPIKE-distance)
performance primarily drove the high SPIKE-distance-based performance. When ACx was
optogenetically suppressed, both the overall and spike timing-based performance was reduced,
while firing rate-based performance remained unchanged. Hotspots where firing rate-based
distance was second highest (Figure 2B) or highest (Figure 2C) indicate values where high
SPIKE-distance-based performance was driven by firing rate modulations. Upon optogenetic
suppression, all three performance measures were degraded at these hotspots. Across all three
types of hotspots, we found those with higher rate-based performances tended to have higher
firing rates during stimulus playback, which increased during optogenetic suppression (Figure
2D).
Because the average SPIKE-distance-based performance was similar across all types, we
calculated the ratio of spike timing-based performance to firing rate-based performance for each
hotspot (Figure 2E). This measure allowed us to determine the extent to which SPIKE-distance-
based performance was based on either spike timing or firing rate modulation, with higher ratios
emphasizing the former.
289

Network Model: Distinct synaptic depression profiles across excitatory and PV inhibitory synapses support high discrimination performance

We created a multi-layer network model to replicate temporal responses to dynamic stimuli in mouse ACx neurons (Figure 3A). The input layer consists of a single Poisson-like thalamic cell simulating pre-cortical input (IC) that exhibits absolute and relative refractory periods of 1 ms. Within the cortical intermediate and output layers, the excitatory cell from the preceding layer synapses onto a feed-forward inhibitory circuit consisting of one relay cell (E) and one onset-responding inhibitory PV cell (red) and one PV cell that fires during offsets in sound stimuli (green) (Figure 3B). Table 1 shows the universal simulation parameters, while Table 2 shows the parameters for all cells in this model.

In all synapses, short-term plasticity (STP) was implemented as previously described (Dayan and Abbott, 2001a). The model is also consistent with previous modeling studies (Seay et al., 2020), in that all synapses in this circuit featured depression, the strength of which was mildest between excitatory cells, followed by excitatory inputs to PV units, with the strongest depression in synapses from PV to excitatory units (Figure 3C). Recovery time constants for each synapse were chosen to match the rapid firing rate modulations in experimental data. Table 3 shows the parameters for all synapses in this circuit.

To investigate how changes in depression strengths affected model responses, we varied the synaptic depression strength (f_P) in each synapse type (E->E, E->PV, and PV->E), ranging from zero ($f_P = 0$) to maximal depression ($f_P = 1$). We first varied the depression strength between excitatory cells (Figure 4A). We found that weak synaptic depression was necessary for high firing rates and performance values at the output excitatory cell. When synaptic depression strength increased, performance dropped towards chance values and firing at output dropped to the network noise rate of 8 Hz.

When we varied the synaptic depression strength at excitatory inputs to PV cells (Figure 4B), we found that all performance values did not greatly change at higher depression strengths, while firing rate greatly increased.

Finally, when we varied the synaptic depression strength of PV inhibition on excitatory cells (Figure 4C), we found that SPIKE-distance-based performance plateaued at higher values of f_P, while ISI-distance-based performance increased, and RI-SPIKE-distance-based performance decreased. Additionally, firing rate steadily increased with synaptic depression strength. From these simulations, we chose depression strengths, i.e., f_P values that were consistent with the firing rates and performances that approximated the experimental single unit data from mouse ACx, as marked by the horizontal dashed lines in Figure 4. Thus, these results revealed that obtaining relatively high discrimination performance and firing rates consistent with the data requires relatively strong depression in PV->E synapses, moderate depression in E->PV synapses and mild depression in E->E synapses.

Relative strengths of onset and offset PV populations in model explains diversity of discrimination profiles observed experimentally

To model the diversity of rate-modulation hotspots (low, medium, and high) identified in the data, we next varied the inhibitory strength of offset PV neurons in both cortical layers while keeping all other circuit parameters constant. For hotspots with lower ratios of spike timing to

firing rate-based performances, the inhibitory signaling currents from offset PV neurons was
reduced relative to that from onset PV neurons (Figure 5A).
Figure 5B illustrates the model's response to one of the two target stimuli shown in
Figure 1B. Throughout the ascending circuit, the firing rate pattern from each input is
maintained. In the model, output units exhibit periods of firing rate modulation where spiking is
close to zero during the quiet periods of the target stimulus. This range of rate modulation at the
output contributes to its high SPIKE-distance-based discrimination performance of 90% between
both target stimuli.
We also found that this circuit effectively captures the balance between firing rate-based
versus spike timing-based performance in hotspots (Figure 5C). For the simulations shown in
5B, ISI-distance-based performance was higher than SPIKE-distance- and RI-SPIKE-distance-
based performance was consistent with the patterns observed in the data (Figure 2C).
The model also captures the relative spike latencies of each cell within cortex (Figure
5D). Given the same input from IC, the onset-responding PV neuron fires first, which inhibits the
excitatory PSC from the same spike input at E. These results are consistent with the differences
in receptive fields between excitatory and PV neurons in mouse ACx (Li et al., 2015). Due to
synaptic depression, later spikes from the onset PV cell have a much weaker inhibitory effect on
the excitatory cell, thus allowing the E cell to fire during this period of weakened inhibition. The
descending edge of excitatory firing is then shaped by inhibition from the offset PV cell, which
sharpens firing rate modulations. This sharpening is clearly apparent when the PSTHs of all
output units are compared (Figure 5E).
To capture hotspots with higher ratios of spike timing to firing rate-based performances,
we ran simulations where inhibition strength from offset PV neurons was higher than that from
onset PVs (Figure 6A). Within experimental data, we found that these hotspots exhibited
responses where the noise floor firing rate during stimulus playback was lower than the
spontaneous firing rate. Thus, we hypothesized that increased inhibition during offset periods
facilitates higher spike timing-based performance. Figure 6B shows the responses to one of the
two target stimuli across all cells within the circuit. The output excitatory cell exhibited a firing
rate of 20 Hz during stimulus playback, which approximates values seen in mouse ACx. For
these simulations, SPIKE-distance- and RI-SPIKE-distance-based performances were almost
equal, with ISI-distance-based performance being the lowest (Figure 6C), which agrees with the
trends observed in the data. Voltage traces and firing rate modulation patterns at output neurons
retain the same sharpening mechanism seen in simulations of hotspots with high firing rate-based
performances (Figures 6D-E). Thus, the model reflects the diversity of firing rate-based
performance seen in experimental data and reveals that altering the relative connectivity
strengths of onset and offset PV neurons effectively captures the broad range of neural
discriminability profiles seen in mouse ACx.
371

Model explains experimentally observed effects of optogenetic suppression of PV neurons

We next employed the model to simulate the effects of reduced PV signaling and
compared the results to empirical data collected in mouse ACx during optogenetic suppression of
PV cells. For the optogenetic condition, we inhibited PV neurons in all layers with a
hyperpolarizing current of 0.03 nA. To model the elevated spontaneous firing rate observed in
single unit recordings (Nocon et al., 2022), we also simulated this by increasing the mean firing
rate of the Poisson-like network noise (Figure 7A).

In agreement with experimental results, we found that discrimination performance values
at the output unit decreased with optogenetic suppression for all spike distance measures (Figure
7B). Control simulations were found to approximate the firing rate profiles found within mouse
ACx, where the firing rate profile followed the temporal envelope of both target stimuli and
showed very little firing during the quiet periods of both targets (Figure 7C, top). During
simulations where PV neurons were suppressed, we saw a decrease in firing in all PV cells and a
slight increase in firing at both cortical excitatory units (Figure 7C, bottom). Notably, we found
that the noise floor was larger in optogenetic suppression trials than in the control condition
during stimulus playback (Figure 7D).
When we simulated optogenetic suppression for low firing rate populations that show
enhanced spike timing-based performance, the model replicated the experimental results across
all three measures of temporal and rate dependent discriminability. Specifically, optogenetic
suppression led to a decrease in both SPIKE-distance- and RI-SPIKE-distance-based
performances, while ISI-distance-based performance remained unchanged (Figure 8A), which
agrees with the data in Figure 2A. Like the simulations from the firing rate-based hotspot,
responses during the control condition were found to approximate those within mouse ACx
(Figure 8B, top). During optogenetic simulations, we saw the same increase in the noise floor
during stimulus playback as in the rate-based hotspot simulations (Figure 8B, bottom, and Figure
8C). Thus, the three-layer circuit model captures the experimentally observed reduction in neural
discriminability with optogenetic suppression of PV neurons in mouse ACx.
400

Discussion

Diverse neural discrimination performance profiles in ACx

In previous experimental study examining the effects of optogenetically suppressing PV
cells in mouse ACx, we found that performance was degraded in terms of both spike timing,
firing rate modulations, and a combination of the two (Nocon et al., 2022). In this study we
further analyzed the degree to which high neural discriminability was determined by either spike
timing or firing rate modulations within each hotspot. We then set out to determine if we could
construct a model that effectively captures the diversity of rate and temporal coding components
seen in single unit data. In addition, we focused on revealing the properties by which PV neurons
could influence discriminability across a range of hotspots present in mouse ACx.

Network model explains diversity of profiles

Previous computational studies of the mouse auditory cortex primarily focused on
modeling responses to transient stimuli such as tones and white noise bursts (Moore et al., 2018;
Seay et al., 2020). Here, we sought to replicate responses to longer, and temporally dynamic
stimuli in mouse ACx. Previously, we proposed a computational model of cortical responses to
target songs within songbirds, which showed sparser spiking activity compared to recordings in
mice (Maddox et al., 2012; Dong et al., 2016). Inputs in the songbird model were based on
STRFs recorded in zebra finch auditory thalamus (Amin et al., 2010), which exhibited a much
sharper temporal profile than those recorded in mouse MGB (Lohse et al., 2020). The model
follows the higher stimulus evoked firing rate observed in mouse auditory thalamus, which is
then temporally sharpened by intra-cortical processing via inhibition from PV interneurons.

One critical finding of the model is that the profile of synaptic depression parameters in distinct synapses is critical for attaining high neural discriminability in auditory cortex. Similarly, synaptic depression at inputs to PV cells and their inhibitory effect on excitatory cells were both found to facilitate rapid modulations in firing rate within the circuit. Specifically, as the thalamic input was processed in cortical layers, firing rate peaks were maintained, and offset-responding PV units enhanced suppression of the troughs in firing rate, thereby enhancing firing rate modulations.

Potential role of onset and offset-responding PV interneurons in generating diverse discrimination performance profiles

We demonstrated that the model circuit was able to capture the diversity of discriminability profiles within auditory cortex by changing the ratio between inhibition strength from onset PV cells and offset PV cells. When inhibition from onset cells was larger than that of offset cells, excitatory units exhibited higher firing rate-based performance. When inhibition from offset PV neurons was stronger, the model captured the remaining set of hotspots with high spike timing to firing rate-based performance ratios. Previous studies have found that PV neurons are implicated in encoding gaps in noise within auditory cortex suggesting an important contributions to spike timing (Keller et al., 2018). The circuit, which features sub-populations of PV neurons that separately respond to sound onsets or offsets, arising from distinct parallel networks that originate in thalamic regions (Liu et al., 2019), further elucidates the contribution of offset responding PV neurons to enhance temporal coding of dynamic stimuli.

Model limitations

The model in this study is a simplified version of cortical circuits. It only includes feed-forward circuits between excitatory and PV units. It does not account for any recurrent inhibitory mechanisms between the two neuron types, nor does it include any other interneuron type, such as somatostatin-expressing (Sst) or vasoactive intestinal peptide-expressing (VIP) neurons, which might further facilitate the high neural discriminability seen in cortex. Additionally, the model utilized single synapses over a population model in which multiple inputs converge onto neurons. These simplifications allowed us to better dissect which components of the cortical network are integral for capturing the variety of hotspots seen in experimental data. The current results suggest an important role of phasic PV inhibition during both sound onsets and offsets to temporally sharpen cortical responses and improve cortical discrimination performance. Future models should incorporate other interneuron types to capture additional aspects of cortical processing. For example, Sst neurons may be important to capture "cross-channel" surround inhibition, and VIP neurons may be important for capturing top-down effects on cortical processing (Chou and Sen, 2021).

In the model, the responses of offset neurons were modeled phenomenologically by inverting the convolutions between the STRF and stimulus waveforms, such that the maximum onset firing rate becomes zero at offset cells and the minimum firing rates become the maxima. A similar approach was used in a previous study on auditory thalamic responses in mice (Anderson and Linden, 2016), and we found that this approximation matched the responses of offset cells seen in the data. Other studies have hypothesized that responses to stimulus offsets along auditory pathways are mediated by hyperpolarization-activated current, amplified by

downstream mechanisms, or inherited from upstream regions (Kopp-Scheinpflug et al., 2011;
Solyga and Barkat, 2021). Future studies should explore mechanistic models of offset neurons.
Finally, while the model utilizes STP between all synapses, the parameters chosen to
model the temporal profile and strength of STP were selected to match the rapid firing rate
modulations noted in the experimental data. The recovery times and depression strengths that
were chosen for the circuit approximate the relative strengths of synaptic depression found
within cortex, with mild depression between excitatory inputs to relay cells and strong
depression in inhibitory inputs to excitatory cells (Seay et al., 2020). With these parameters, the
model captured both the neural discrimination performance profiles and mean firing rate values
seen within the experimental data.
478

Key model predictions and future directions

In this study, we presented a computational model describing how PV-driven inhibition
transforms thalamic inputs across cortical layers to facilitate high neural discrimination
performance within auditory cortex. The computational model is based on physiological data in
mice and posits that 1) cortical processing refines stimulus encoding at the thalamic level as
spiking propagates downstream, 2) spiking in PV neurons are more phasic in time when
compared to responses in excitatory neurons, 3) PV neurons that are differentially tuned to
temporal features in auditory stimuli sharpen spiking activity in excitatory neurons, thus leading
to increased neural discriminability performance within cortex.
487

In this model, we focused on dissecting the rich temporal dynamics of "within-channel"
inhibition mediated by PV neurons, to replicate experimental data from clean trials in which only
a single stimulus was present. We did not model "cross-channel" interactions across spatial
channels, which may be critical for modeling the emergence of hotspots at different spatial
configurations. Based on the architecture of previous models (Dong et al., 2016; Chou and Sen,
2021), future models of mouse ACx should incorporate multiple spatial channels, with cross
channel "surround" inhibition. The analysis found that suppression of PV neurons reduced
performance across all configurations of target and masker location (Nocon et al., 2022); thus,
we predict that other interneurons types, e.g., Sst and VIP neurons, may play an important role in
determining the emergence of hotspots at different spatial configurations.

More generally, the modeling paradigm developed in this study may be useful in
investigating the roles of different cortical interneuron types in complex scene analysis. The
experimental data that were used to create and validate this model involved the optogenetic
suppression of PV interneurons within mouse ACx. Future experiments with optogenetic
manipulations or opto-tagging of other interneuron types within cortex, such as Sst or VIP
neurons can be used to expand the model circuit. Ultimately, an understanding of the
contributions of the diverse populations of interneurons in cortex, will be necessary to reveal
cortical circuit mechanisms underlying complex scene analysis.
506

Acknowledgements

508 We thank Michael Economo, Alberto Cruz-Martín, and Monty Escabi for helpful discussions
509 and comments on the manuscript. The work in this manuscript was funded by NSF grant #IIS-
510 1835270 and NIH/NINDS grant #1R34NS111742-01.

References

Adelson EH, Bergen JR (1985) Spatiotemporal energy models for the perception of motion. J
Opt Soc Am A 2:284-299.
Amin N, Gill P, Theunissen FE (2010) Role of the zebra finch auditory thalamus in generating
complex representations for natural sounds. J Neurophysiol 104:784-798.
Anderson LA, Linden JF (2016) Mind the Gap: Two Dissociable Mechanisms of Temporal
Processing in the Auditory System. J Neurosci 36:1977-1995.
Chou KF, Sen K (2021) AIM: A network model of attention in auditory cortex. PLoS Comput
Biol 17:e1009356.
Dayan P, Abbott LF (2001a) Release probability and short-term plasticity. In: Theoretical
neuroscience (Sejnowski TJ, Poggio T, eds), pp 184-187. Cambridge, MA: The MIT
Press.
Dayan P, Abbott LF (2001b) Spike-rate adaptation. In: Theoretical neuroscience (Sejnowski TJ,
Poggio T, eds), pp 165-166. Cambridge, MA: The MIT Press.
Dayan P, Abbott LF (2001c) Space-Time Receptive Fields. In: Theoretical neuroscience
(Sejnowski TJ, Poggio T, eds), pp 68-69. Cambridge, MA: The MIT Press.
Dong J, Colburn HS, Sen K (2016) Cortical Transformation of Spatial Processing for Solving the
Cocktail Party Problem: A Computational Model(1,2,3). eNeuro 3.
Fanselow EE, Connors BW (2010) The roles of somatostatin-expressing (GIN) and fast-spiking
inhibitory interneurons in UP-DOWN states of mouse neocortex. J Neurophysiol
104:596-606.
Keller CH, Kaylegian K, Wehr M (2018) Gap encoding by parvalbumin-expressing interneurons
in auditory cortex. J Neurophysiol 120:105-114.
Kopp-Scheinpflug C, Tozer AJ, Robinson SW, Tempel BL, Hennig MH, Forsythe ID (2011) The
sound of silence: ionic mechanisms encoding sound termination. Neuron 71:911-925.
Kreuz T, Chicharro D, Houghton C, Andrzejak RG, Mormann F (2013) Monitoring spike train
synchrony. J Neurophysiol 109:1457-1472.
Li LY, Xiong XR, Ibrahim LA, Yuan W, Tao HW, Zhang LI (2015) Differential Receptive Field
Properties of Parvalbumin and Somatostatin Inhibitory Neurons in Mouse Auditory
Cortex. Cereb Cortex 25:1782-1791.
Li LY, Ji XY, Liang F, Li YT, Xiao Z, Tao HW, Zhang LI (2014) A feedforward inhibitory
circuit mediates lateral refinement of sensory representation in upper layer 2/3 of mouse
primary auditory cortex. J Neurosci 34:13670-13683.
Liu J, Whiteway MR, Sheikhattar A, Butts DA, Babadi B, Kanold PO (2019) Parallel Processing
of Sound Dynamics across Mouse Auditory Cortex via Spatially Patterned Thalamic
Inputs and Distinct Areal Intracortical Circuits. Cell Rep 27:872-885 e877.
Lohse M, Bajo VM, King AJ, Willmore BDB (2020) Neural circuits underlying auditory contrast
gain control and their perceptual implications. Nat Commun 11:324.
Maddox RK, Billimoria CP, Perrone BP, Shinn-Cunningham BG, Sen K (2012) Competing
sound sources reveal spatial effects in cortical processing. PLoS Biol 10:e1001319.
Moore AK, Weible AP, Balmer TS, Trussell LO, Wehr M (2018) Rapid Rebalancing of
Excitation and Inhibition by Cortical Circuitry. Neuron 97:1341-1355 e1346.
Nocon JC, Gritton HJ, James NM, Han X, Sen K (2022) Parvalbumin neurons, temporal coding,
and cortical noise in complex scene analysis. bioRxiv.
Pfeffer CK, Xue M, He M, Huang ZJ, Scanziani M (2013) Inhibition of inhibition in visual
cortex: the logic of connections between molecularly distinct interneurons. Nat Neurosci
16:1068-1076.

Qiu A, Schreiner CE, Escabi MA (2003) Gabor analysis of auditory midbrain receptive fields:
spectro-temporal and binaural composition. J Neurophysiol 90:456-476.
Rothauser EH, Chapman, W. D., Guttman, N., Nordby, K. S., Silbiger, H. R., Urbanek, G. E., &
Weinstock, M. (1969) I.E.E.E. recommended practice for speech quality measurements.
IEEE Trans Audio Electroacoust 17:225-246.
Satuvuori E, Kreuz T (2018) Which spike train distance is most suitable for distinguishing rate
and temporal coding? J Neurosci Methods 299:22-33.
Satuvuori E, Mulansky M, Daffertshofer A, Kreuz T (2018) Using spike train distances to
identify the most discriminative neuronal subpopulation. J Neurosci Methods 308:354-
365.
Schaette R, Gollisch T, Herz AV (2005) Spike-train variability of auditory neurons in vivo:
dynamic responses follow predictions from constant stimuli. J Neurophysiol 93:3270-
3281.
Seay MJ, Natan RG, Geffen MN, Buonomano DV (2020) Differential Short-Term Plasticity of
PV and SST Neurons Accounts for Adaptation and Facilitation of Cortical Neurons to
Auditory Tones. J Neurosci 40:9224-9235.
Sherfey JS, Soplata AE, Ardid S, Roberts EA, Stanley DA, Pittman-Polletta BR, Kopell NJ
(2018) DynaSim: A MATLAB Toolbox for Neural Modeling and Simulation. Front
Neuroinform 12:10.
Solyga M, Barkat TR (2021) Emergence and function of cortical offset responses in sound
termination detection. Elife 10.
Theunissen FE, David SV, Singh NC, Hsu A, Vinje WE, Gallant JL (2001) Estimating spatio-
temporal receptive fields of auditory and visual neurons from their responses to natural
stimuli. Network 12:289-316.
Udakis M, Pedrosa V, Chamberlain SEL, Clopath C, Mellor JR (2020) Interneuron-specific
plasticity at parvalbumin and somatostatin inhibitory synapses onto CA1 pyramidal
neurons shapes hippocampal output. Nat Commun 11:4395.
Xue M, Atallah BV, Scanziani M (2014) Equalizing excitation-inhibition ratios across visual
cortical neurons. Nature 511:596-600.
588
589

Figure 1. Experimental data from mouse ACx. (A) Diagram of experimental setup and spatial location of target stimuli. During trials, one of two target stimuli consisting of white noise modulated by a human speech-shaped envelope was played from one of the four speakers along the azimuth in horizontal space. **(B)** Responses to spatial configuration shown in A where the two targets originated at 90°, from an example single unit in mouse ACx. Responses followed the amplitude envelope of both targets, resulting in high SPIKE-distance-based neural discrimination performance (92%) at this configuration. **(C)** Average discrimination performance using different spike distance metrics across the population of 24 high-performing single units (n = 50 hotspots during clean trials). Paired t-tests found significant differences between control and optogenetic trials for all spike distance-based performances (SPIKE-distance: $p = 8e\text{-}10$; ISI-distance: $p = 3e\text{-}08$; RI-SPIKE-distance: $p = 1e\text{-}08$).

Figure 2. Hotspots in single-unit experimental data reflect a broad range of timing and
rate-based contributions to performance. Hotspots in single-unit data were sorted by
comparing firing rate modulation-based (ISI-distance) performance to the other two measures
(SPIKE-distance, firing rate modulation and spike timing; RI-SPIKE-distance, spike timing
only). **(A)** Hotspots where ISI-distance-based performance was the worst out of all three
performance measures (n = 9), as indicated by the horizontal dashed line. **(B)** Hotspots where
ISI-distance-based performance was higher than RI-SPIKE-distance-based performance but not
SPIKE-distance-based performance (n = 16). **(C)** Hotspots where ISI-distance performance was
higher than the other two performance values (n = 25). **(D)** Auditory stimulus evoked average
firing rates during sound alone and PV optogenetic inhibition for each of the three hotspot types.
(E) Histogram showing all 50 hotspots from single-units, sorted by the ratio between RI-SPIKE-
distance-based performance and ISI-distance-based performance, with a ratio of 1 indicated by
the dashed line. Ratios lower than 1 indicate hotspots where high neural discrimination
performance is primarily driven by modulations in firing rate, rather than spike timing.
616

Figure 3. Computational model of auditory cortex layers and thalamic input. (A) Three-layer ACx model. The input stage of the model consists of a Poisson-like thalamic neuron with input from the inferior colliculus (IC) with a firing rate based on the convolution of an MGB STRF with stimulus spectrograms. Both the intermediate **(Int)** and output **(Out)** layers exhibit feed-forward inhibitory circuits mediated by PV cells (red). Excitatory cells also received inhibition from offset-sensitive PV cells (green), which were driven by excitatory cells that respond to stimulus offsets (not shown). Offset responses were generated by inverting the convolution between the STRF and stimulus and adding the absolute maximum value of the resulting firing rate trace, and columns of offset-responding excitatory cells have identical structure as onset-responding columns shown in A. The excitatory cell in the output layer exhibited Poisson noise to simulate network noise in ACx. **(B)** Target 1 waveform with example FR trace from STRF convolution for onset- (red) and offset-responding (green) units. Scale bar measures 50 Hz and 250 ms. **(C)** Example traces of synaptic plasticity. Left: mild synaptic depression between excitatory cells. Middle: synaptic depression in excitatory inputs to PV cells. Right: strong synaptic depression at inhibitory inputs to excitatory cells.

C [nF]	0.1
V_{thresh} [mV]	-47
I_{tonic} [nA]	0
Simulation time [ms]	3500
Integration time step [ms]	0.1

633 **Table 1. Universal DynaSim parameters.**

Unit	R (MΩ)	Δg_{sra} (nS/spk)	τ_{sra} (ms)	t_{ref} (ms)	E_L (mV)	V_{reset} (mV)
IC	200	0	--	1	-65	-54
E	200	0.3	100	1	-65	-54
PV	100	0	--	0.5	-57	-52

634   **Table 2. Unit parameters.** DynaSim parameters that differed between units.

Connection	g_{syn} (nS)	τ_1 (ms)	τ_2 (ms)	E_{syn} (mV)	f_D	τ_D (ms)
E->E	20	0.7	1.5	0	0.1	30
N->E_{Out}	15	0.7	1.5	0	0	--
IC->PV_{Int} E_{Int}->PV_{Out}	32	0.1	1	0	0.2	80
PV->E	25	1	4.5	-80	0.5	120

Table 3. Synaptic parameters. DynaSim parameters that differed between synapses. N represents simulated network noise from Poisson units in the output layer that were not explicitly modeled as a separate population.

Figure 4. Effects of synaptic depression strengths on output discriminability and firing.
Simulations for a rate-based hotspot in output excitatory cells, where synaptic depression
strengths (f_P) were varied while keeping all other parameters constant. **(A)** Synaptic depression
strength between excitatory cells versus discriminability measures (top) and firing rate (bottom).
Simulations where $f_{P,E->E} = 0.1$ (indicated by the vertical dashed line) leads to SPIKE-distance-
based neural discriminability and firing rate values that approximate those of experimental data
in hotspots with high firing rate-based performances, both of which are indicated by horizontal
dashed lines. **(B)** Synaptic depression strength between excitatory inputs to PV cells versus
discriminability measures (top) and firing rate (bottom). Simulations where $f_{P,E->PV} = 0.2$ results
in performance and firing rate values that approximate those of experimental data. **(C)** Synaptic
depression strength of inhibitory inputs to excitatory cells versus discriminability measures and
firing rate. Simulations where $f_{P,PV->E} = 0.5$ resulted in performance and firing rate values that
approximate those in hotspots with high firing rate-based performance.

Figure 5. Circuit simulations of a hotspot with high firing rate-based performance are facilitated by high onset inhibition. (**A**) To simulate hotspots where firing rate-based performance was higher than spike timing-based performance, inhibition strength from offset-sensitive PV units (w_{Off}, green) was reduced relative to that of onset-sensitive PV units (w_{On}, red). (**B**) Responses during the first second of one target stimulus (black waveform, bottom) from each cell, with rasters plotted in gray and the corresponding PSTH overlaid in the same colors as those shown in Figure 3A. To better show firing rate patterns in all cells, PSTHs were smoothed using a moving-average filter spanning 3 samples. Scale bars at bottom-right corner indicates vertical scale of PSTHs and horizontal time scale. Average firing rate at the output excitatory cell was 32 Hz during stimulus playback. (**C**) Performance measures of simulations shown in A. Here, ISI-distance-based performance is higher than RI-SPIKE-distance-based performance, indicating that the high SPIKE-distance-based performance is dominated by rapid changes in firing rate modulation. (**D**) Example voltage traces from output layer neurons. Onset-sensitive PV neurons (top) fire first from IC inputs, which then leads to inhibition of E cells (middle), thereby sculpting the ascending edge of firing rate modulations in excitatory cells. The descending edge is sculpted by offset-sensitive PV cells (bottom). Scale bars measure 20mV and 10ms. (**E**) Zoomed-in smoothed PSTHs (bin size 20ms) of output layer cells during the first half second of sound onset. PSTHs of PV cells indicate how inhibition shapes the firing rate patterns in excitatory cells.

Figure 6. Circuit simulations of a hotspot with high spike timing-based performance are facilitated by high offset inhibition. (A) To simulate hotspots where spike timing-based performance was higher than firing rate-based performance, inhibition strength from offset-sensitive PV units (w_{Off}, green) was elevated relative to onset-sensitive PV units (w_{On}, red). **(B)** Responses during the first second of one target stimulus (black waveform, bottom) from each cell, with rasters plotted in gray and the corresponding smoothed PSTH overlaid in the same colors as those shown in Figure 3A. Scale bars at bottom-right corner indicates vertical scale of PSTHs and horizontal time scale. Average firing rate at the output excitatory cell was 20 Hz during stimulus playback. **(C)** Performance measures of simulations shown in A. Here, ISI-distance-based performance is higher than RI-SPIKE-distance-based performance, indicating that the high SPIKE-distance-based performance is dominated by rapid changes in firing rate modulation. **(D)** Example voltage traces from output layer neurons. Onset-sensitive PV neurons (top) fire first from IC inputs, which then leads to inhibition of E cells (middle), thereby sculpting the ascending edge of firing rate modulations in excitatory cells. The descending edge is sculpted by offset-sensitive PV cells (bottom). Scale bars measure 20mV and 10ms. **(E)** Zoomed-in smoothed PSTHs (bin size 20ms) of output layer cells during the first half second of sound onset.

Figure 7. Simulations of responses during optogenetic suppression for a hotspot with high
firing rate-based performance. For all simulations used in this figure, synaptic connectivity
strengths were the same as in Figure 5. **(A)** Diagram showing changes to circuit during
optogenetic suppression simulations. All PV cells are suppressed with a hyperpolarizing current
of 0.03 nA, and the network noise input to the output excitatory cell is increased from 8 Hz to 12
Hz. **(B)** Average performance values during control and optogenetic conditions across 5 sets of
simulations per condition, with error bars representing $\pm$ s.e.m. **(C)** Example rasters and PSTHs
to each target stimulus during one control and one optogenetic simulation. **(D)** Inset showing
zoomed-in portion of the response to target stimulus #1, between 0.6 and 1.8s after sound onset,
as outlined in C. Responses during optogenetic trials show a higher noise floor, compared to the
control or non-laser condition.
701

Figure 8. Simulations of responses during optogenetic suppression in a hotspot with high
spike timing-based performance. For all simulations used in this Figure, synaptic connectivity
strengths were the same as in Figure 6. **(A)** Average performance values during control and
optogenetic trials across 5 sets of simulations per condition, with error bars representing ± s.e.m.
(B) Example rasters and PSTHs to each target stimulus during one control and one optogenetic
simulation. **(C)** Inset showing zoomed-in portion of the response to target stimulus #2, between
2.45 and 2.75s after sound onset, as outlined in B. Responses during optogenetic trials show a
higher noise floor, compared to the control or non-laser condition.
710

A Low rate-based performance hotspots

B Medium rate-based performance hotspots

C High rate-based performance hotspots

D Firing rate changes during optogenetic suppression

E Timing-based versus rate-based control performances

A

Out

E

w_{On} w_{Off} Offset

PV   PV

Int

E

w_{On} w_{Off} Offset

PV   PV

IC

Onset

B

Target 1

Onset FR

Offset FR

C

Mild depression

E->E

Depression

E->PV

Strong depression

PV->E

A High firing rate-based performance hotspots

g_{SYN} (μS)

B

	PV_{On}	E	PV_{Off}
Out			
Int	PV_{On}	E	PV_{Off}
IC			

75 Hz
250 ms

C Output E performance

Performance

100%
90%
80%
70%
60%
50%

SPIKE ISI RI-SPIKE

Spike distance measure

D

Out PV_{On}
Out E
Out PV_{Off}

E

Output spike count

— PV_{On}
— E
— PV_{Off}

400 650 900
Time (ms)