1   **Title:** Neotropical birds and mammals show divergent behaviour responses to human pressure

3   **Keywords:** *Activity patterns, dial activity, vertebrates, landscape of fear, nocturnality, diurnality*

5   **Authors:** Pablo Jose Negret*[1,2,3], Mathew Scott Luskin[2,4], Bibiana Gomez-Valencia[5], Angelica Diaz-
6   Pulido[5], Luis Hernando Romero[5], Adriana Restrepo[5], Julie G. Zaehringer[3], Jose Manuel Ochoa-
7   Quintero[4,5], Calebe Pereira Mendes[7]
8

Affiliations:

[1]The University of Queensland, School of Earth and Environmental Sciences, Qld 4072, Australia

[2]The University of Queensland, Centre for Biodiversity and Conservation Science, Qld 4072, Australia

[3] University of Bern, Wyss Academy for Nature, Centre for Development and Environment, Institute
of Geography, Bern, Switzerland

[4]The University of Queensland, School of Biological Sciences, Qld 4072, Australia

[5]Instituto de Investigación de Recursos Biológicos Alexander Von Humboldt, Bogotá D.C., Colombia

[6]Universidade Federal do Mato Grosso do Sul, Campo Grande 79070-900, Brazil

[7]Nanyang Technological University, Asian School of the Environment, Singapore

Corresponding authors mailing address:

*Pablo Jose Negret

Email: p.negret@uq.edu.au

Word count: 5686

Acknowledgments:

We would like to acknowledge the people from the Maron Ecology & Conservation Policy Lab and

the Ecological Cascades Lab for helpful discussion on the interpretation of the results and to C.

Rojano, L. Miranda y R. Ávila (Fundación Cunaguaro), A. Lopera, F. Hernández, R. Fernández y R.

Rodríguez (Corporación Gaica), Á. E. Cáceres-Gómez, N. Ferro-Muñoz. F. Cáceres-Gómez (Biotica

Consultores) for the field work, and S. Pérez for the revision of the bird species lists. We thank

Ecopetrol S.A. and Instituto de Investigación de Recursos Biológicos Alexander Von Humboldt for

financing the study within the framework of the FIBRAS Agreement.

Neotropical birds and mammals show divergent behaviour responses to human pressure

Abstract:

Human presence and habitat disturbance (together 'human pressure' hereafter) can generate a deep

fear in animals and this can influence their behaviour. Altered animal behaviour, such as shifts in diel

activity patterns, affect many species and species interactions, which can induce changes in individual

fitness, species-level population persistence, evolutionary dynamics, and ecosystem-level biodiversity.

However, whether dial activity behavioural responses to human pressure are consistent among key

functional groups has been poorly studied. For example, while medium to large mammal species tend

to become more nocturnal in areas with high human pressure, it's unclear if sympatric/co-occurring

birds display similar or opposite patterns. This is an important knowledge gap because synchronous

or opposing guild-level shifts can shape consequences for food-web dynamics (predation and

competition), stability of interaction networks and ecosystem functioning. Here we used information

from camera trapping along a gradient of human pressure in the Colombian Llanos region to assess

diel activity changes in birds and mammals. We found that the diel activity of over 45% of the bird

and 50% of the mammals assessed significantly changed where there was higher human pressures,

with mammals becoming more nocturnal and birds more diurnal. The opposing behavioural

responses to humans among vertebrate functional groups has significant repercussions for the fields

of community ecology, including intraguild predation and competition, and may be a significant

ecosystem-level conservation consideration.

Introduction:

The extent of human pressure has increased dramatically in the last 20 years with over two thirds of the global land area under moderate or intense human pressure (Williams et al., 2020b). The related habitat changes and specific anthropic pressures (e.g. hunting) have vastly altered wildlife communities including extirpations, extinctions, and changing ecosystem functioning (Dirzo et al., 2014; Haddad et al., 2015). These direct effects are also accompanied by the indirect effects of humans on the behaviour of remaining wildlife, which poses a significant threat to species fitness and species interactions as well as overall biodiversity and ecosystem functioning (Gaynor et al., 2018; Sih et al., 2011; Zanette and Clinchy, 2019).

Human presence and habitat disturbance (together 'human pressure' hereafter) can generate a deep fear in animals and this "landscape of fear" can influence their diel activity patterns and use of space (Gallagher et al., 2017; Laundre et al., 2010; Zanette and Clinchy, 2019). Medium and large mammals tend to avoid humans by increasing their nocturnality (Gaynor et al., 2018; Mendes et al., 2020) and reducing their home range closer to urban environments (O'Donnell and Delbarco-Trillo, 2020). However, a recent study found scarce evidence of activity pattern shifts for mammals in areas with high human pressure (oil palm plantations) in the Colombian Llanos (Pardo et al., 2021). Further investigation is needed since these variations in wildlife responses to humans are not well understood but they could have immense repercussions for species interactions such as predation and competition, which can restructure entire food webs, energy flows and ecosystem functioning.

Bird behavioural responses to humans may differ from mammals but our understanding is more limited, especially in the tropics (Bonier et al., 2007; Chace and Walsh, 2006; Zanette and Clinchy, 2019). Chilean forest birds shifted their daily activity peaks from early morning in old-growth forests, to noon in plantations, and to the afternoon in logged forests (Fontúrbel et al., 2021) and forest birds in Hawaii showed increased diurnality in fragmented landscapes (Smetzer et al., 2022). One difference underlying variation in vertebrate functional group responses could be related to harvest rates, which are often predicted by larger body size and ease of capture (less arboreality or flight)

(Dirzo et al., 2014). In a global review of hunting in tropical forests, birds and mammals both showed

significant declines in hunting-accessible areas, although 25% less for birds than mammals (Benítez-

López et al., 2017). There is also evidence that birds need to forage more in fragmented habitats,

which could increase their diurnality (Redpath, 1995; Saunders, 1980). Fearful animals near humans

may perceive safety differently depending on their ability to hide versus escape, with arboreal birds

preferring higher daytime visibility and ease of escape via flight, while terrestrial mammals may view

nocturnal stealthy foraging as safer (Gallagher et al., 2017; Zanette and Clinchy, 2019).

Understanding bird behavioural responses to human pressure and how this responses may differ from

those of mammals is an important knowledge gap because synchronous or opposing guild-level shifts

among mammals and birds can shape consequences for food-web dynamics and ecosystem

functioning (Zanette and Clinchy, 2019).

Colombia, one of the most biodiverse countries is affected by high levels of habitat degradation and

human pressures have doubled in the last 50 years (Correa Ayram et al., 2020). However, there is

limited evidence if there are changes to wildlife behaviour do to this increase in human pressure. We

used camera trapping along a gradient of human pressure in the Colombian Llanos to measure

changes in daily activity patterns of birds and mammals. For medium and large birds and mammals,

we expected to observe shifts towards nocturnality for species susceptible to poaching and hunting

and no changes for small species. Finally, we expect to see congruent responses from birds and

mammals with similar ecological characteristics, such as game species, and species with similar body

size and diet.

Methods:

Survey

To detect possible shifts in the activity patterns of birds and mammals, we used a large dataset of

camera-traps (Bushnell Core DS No Glow) to record the activity period of bird and mammal species

in areas with different levels of human pressure. The study area is located in the Llanos region in

Colombia. The sampling occurred from November 2020 to February of 2021 in three areas with

different levels of human pressure (Fig 1). The region is highly biodiverse but also under increased

pressure from agricultural expansion (Williams et al., 2020a). The areas are all immersed in a very

diversified landscape with different grades of human pressure within and among the areas, ranging

from isolated forest patches near human settlements and roads to continuous forest, including areas of

direct influence of hydrocarbon exploitations with proximity to roads and urban settlements, as well

as areas of forest, morichal and natural savannah or pastures for livestock with scattered trees and

bushes.

We sampled each of the three sites with 367 camera stations located along a gradient of Human

Footprint Index -HFI- (Correa Ayram et al., 2020); 186 in the Meta Foothills site (22-85 HFI), 138 in

the Casanare Foothills site (22-91 HFI) and 43 in the Tillavá River site (8-45 HFI). Each station

received one camera for a period between 16 and 96 days. Each camera trap station was placed no less

than one km away from any other camera. The camera-traps were placed around 50 cm above the

ground, recording photos in response to the activation of a passive infrared sensor. All images were

stored and processed in the Wildlife Insights platform and specialists verified the animal

identifications (Diaz-Pulido et al., 2022a, 2022b, 2022c). We used 30 minutes as a minimum time

allowed between two records for the same species at the same camera, as a safety measure to avoid

repeated records of the same individual. This effort resulted on a total of 16,939 camera/days and 165

species recorded; 120 birds and 45 mammals (SI Table 1).

With the data obtained from camera-trapping, we classified the records in two groups based on the

time of sunset and sunrise. Records taken when the sun was up in the sky (i.e. after sunrise and before

sunset) were considered "day" records, while records taken after the sunset or before sunrise were

considered "night" records. Since the exact timing of sunrise slightly varies along the study area

according to latitude and date, we determined each camera station's longitude, latitude, date and time

and determined the exact timing of sunrise and sunset from the U.S naval observatory solar calendar

(U.S. Naval Observatory, 2022). The records for each species in each sampling station were classified

in day/night. Species with less than 5 captures in either the day or night were excluded from the

analysis as the number of observations for these species was not enough to allow for statistical

comparison. Then, using the day/night records for each one of the camera trap stations as a binomial

variable (this is, every record as a Bernoulli trial), we evaluated the effect of human pressure on 11

bird and 14 mammal species, which had enough data to perform the analysis across the study area

(Table 1).

Human pressure proxy

We used the recently developed Colombian Legacy-adjusted Human Footprint Index (LHFI) for

2018, which is the most recent product of its kind for Colombia. It is temporally consistent, and has a

more precise spatial resolution (300 m) than any other national (Etter et al., 2011) or global human

footprint map (Venter et al., 2016; Williams et al., 2020b). This index measures the level of human

pressure on ecosystems and provides a baseline for monitoring human pressure on biodiversity. It was

generated by the Humboldt institute (Correa Ayram et al., 2019) using the same methodology as for

the one developed for the year 2015 by Correa Ayram et al., (2020). The LHFI is estimated using

seven primary variables: land use type, rural population density, distance to roads, distance to

settlements, fragmentation of natural vegetation, the biomass relative to natural potential, and time of

intervention on ecosystems in years. All seven primary variables were estimated at 2018 and re-scaled

between 0 and 5 to reflect their relative contribution to human impact and transformation. Values of 0

indicated a null contribution, while values of 5 indicated a very high contribution. Then the variables

were added, and the final index was normalized between 0 and 100. See Correa Ayram et al., (2020)

for a complete description of the variables and the methodology. The Human Footprint Index -HFI-

(Correa Ayram et al., 2020) for each camera station was recorded.

Data analysis

We used logistic regressions to evaluate the effect of human pressure on birds' and mammals' daily

activity patterns, with the day/night records for each species within each camera station as a binomial

response variable and the LHFI as the explanatory variable. To determine the best spatial scale for

each species, four logistic models were created using the average LHFI from buffers with radius of

250, 500, 750 and 1000 m around each camera trap station, which were compared by the Akaike

Information Criterion corrected for small sample sizes (AICc) (Burnham and Anderson, 2018). For

each species, the model with smallest AICc value was used for the logistic regression, while the

others were excluded. Since the model selection was used only to select the best spatial scale for each

species, a null model was not required (Mendes et al., 2020). For the species which presented

responses to human pressure, we also calculated their activity shift, here described as the difference in

the model predicted diurnality between the most preserved and most disturbed areas in which the

species were recorded. Finally, the activity period of all species was also plotted for visual inspection

(Fig. 2), and circular metrics of diel activity, such as mean direction and Rho (i.e. a measurement of

circular mean and dispersion, respectively) were calculated and reported (SI Table 2). The species'

diet, body size, activity pattern (diurnal, nocturnal or cathemeral) and whether it was a game species

or not was determined based on both local expert knowledge and literature (Emmons and Feer, 1997;

Hilty and Brown, 2001; IUCN, 2022; Myers et al., 2022; Ocampo et al., 2021). All analysis were

performed in R version 4.2.1 (R Core Team, 2022), using the packages "lme4" (Bates et al., 2015),

"circular" (Agostinelli and Lund, 2022), and ggplot2 (Wickham, 2016).

Results:

We found evidence of significant activity shifts in response to human pressure for seven out of 14

species of mammals (50%) and five of the 11 species of birds (45%) that had enough records to be

assessed. For mammals, six became more nocturnal in disturbed areas while one more diurnal (Table.

1, Fig. 3a & 4a), while for birds four became more diurnal while one became more nocturnal (Table.

1, Fig. 3b & 4b). The average increase in nocturnality for mammals was 9.5% while the average

increase in diurnality for birds was 7%. However, we must clarify that by "becoming more nocturnal"

or "becoming more diurnal", we did not mean that a species shifted its peak of activity from a period

to another. Instead, it means that although the species main activity is still occurring during its natural

activity period (i.e. night for nocturnal species and day for diurnal species), the proportions of the

day/night records have changed (Mendes et al., 2020).

The bird species which responded to human pressure by becoming more diurnal were the swainson's

thrush (*Catharus ustulatus*), the little tinamou (*Crypturellus soui*), the white-tipped dove (*Leptotila*

verreauxi) and the amazonian motmot (*Momotus momota*). The pauraque (*Nyctidromus albicollis*)

was the only bird species that responded to human pressure by becoming more nocturnal. Other six

bird species did not present detectable activity shifts in response to human pressure. The grey-necked

wood rail (*Aramides cajaneus*), the pectoral sparrow (*Arremon taciturnus*), the cinereous tinamou

(*Crypturellus cinereus*), the grey-fronted dove (*Leptotila rufaxilla*), the black-faced antbird

(*Myrmoborus myotherinus*) and the southern lapwing (*Vanellus chilensis*).

In relation to mammals, the species which responded to human pressure by becoming more nocturnal

were the paca (*Cuniculus paca*), the agouti (*Dasyprocta fuliginosa*), the nine-banded armadillo

(*Dasypus novemcinctus*), the common opossum (*Didelphis marsupialis*), the grey four-eyed opossum

(*Philander opossum*) and the collared anteater (*Tamandua tetradactyla*). Three of these species

(agouti, paca and armadillo) are heavily hunted or persecuted. The tayra (*Eira barbara*) was the only

mammal species that responded to human pressure by becoming more diurnal. Other seven species,

the crab-eating fox (*Cerdocyon thous*), the ocelot (*Leopardus pardalis*), the red brocket (*Mazama*

americana), the giant anteater (*Myrmecophaga tridactyla*), the ring-tailed coati (*Nasua nasua*), the

white-tailed deer (*Odocoileus virginianus*) and the collared peccary (*Dicotyles tajacu*), did not present

detectable activity shifts in response to human pressure.

Discussion:

We found evidence of significant activity shifts in response to human pressure for 45% of the bird

species assessed, with four of five shifts towards more diurnal activity patterns. The four species

exhibiting diurnal shifts were already diurnal and one is a game species (little tinamou; *Crypturellus*

soui) and the pauraque (*Nyctidromus albicollis*) shifted from cathemeral in more intact areas towards

nocturnality. The increases in diurnality may be a response to increased foraging required to counter

the decreases in food availability in degraded habitats [e.g. due to logging, edges, fragmentation

(Fontúrbel et al., 2021; Redpath, 1995; Saunders, 1980; Smetzer et al., 2022)]. Degraded habitats may

also influence vertebrates activity patterns because they have higher variation in microclimate, which

can affect other environmental factors such as sunlight (Bennie et al., 2014; Ewers and Banks-Leite,

2013; Fontúrbel et al., 2021; Schmidt et al., 2017), insect prey (Checa et al., 2014; Outhwaite et al.,

2022) and other key resources like flowering, nectar production and fruit ripening (Jackson, 1966;

Vitasse et al., 2021).

Three of the six bird species that didn't show a statistically significant shift in daily activity patterns

(black-faced antbird, southern lapwing and pectoral sparrow) have few captures. The grey-necked

wood rail and the grey-fronted dove had non-significant shifts toward diurnality with increased human

pressure. The cinereous tinamou had a non-significant increase in nocturnality with human influence,

opposite to the significant shift found in the little tinamou. These two tinamous are sympatric,

morphologically similar, and closely related ground-dwelling birds, which may generate a pressure

for temporal niche partitioning (Dias et al., 2016; Negret et al., 2015b) that may be reinforced in

degraded habitats with lower food availability.

In relation to mammals, we found that 50% of the species assessed showed activity shifts in response

to human pressure. These results align with the findings of other studies that show changes in activity

patterns for mammals species with increased human pressure (Bennie et al., 2014; Gaynor et al.,

2018; Mendes et al., 2020). From the seven mammal species which had a significant activity shift,

four are nocturnal and three of those are also game species. All of them had shifts towards a more

nocturnal behaviour. From the three diurnal species which presented activity pattern shifts, the agouti

and the Common opossum, which are hunted, became more nocturnal when in disturbed areas, while

the tayra became more diurnal in areas with higher human pressure. The Tayra is a generalist species

that is able to subsist on second growth forest and agricultural mosaics without forest cover, and may

subsidize their diet taking advantage of domesticated animals such as chicken (Meneguetti et al.,

2014; Michalski and Peres, 2005). These characteristics could explain its shift towards diurnality in

areas where there are anthropogenic sources of food that can be opportunistically exploited during

daytime.

Four of the seven mammal species that didn't show a statistically significant shift in daily activity

patterns (ocelot, ring tailed coati, white-tailed deer and collared peccary), had a low number of

records which could have influenced the explanatory power of those models. The giant anteater, a

species that is occasionally killed or captured and that is classified as Vulnerable (Bertassoni and

Abba, 2014; Emmons and Feer, 1997) had an increase in nocturnality in areas with higher human

pressure as was the case for the nocturnal mammal species that had significant changes in daily

activity patterns. The crab-eating fox and red brocket deer had an increase in diurnality in areas with

higher human pressure. These two species can be found in degraded landscapes and have generalist

diets (Emmons and Feer, 1997), which suggests that they might be less affected by human pressure

than other species.

The effect of human pressure on vertebrate species goes beyond impacting species richness and

abundance. Our results show that increased human pressure can have a marked impact in the activity

pattern of multiple vertebrate groups. Particularly, we found that 45% of the bird species and 50% of

the mammal species assessed had shifts in their activity pattern with increases in human pressure.

Bird species had a general shift towards diurnality, while mammal species towards nocturnality,

revealing that different vertebrate groups can react distinctly to increased human presence and

activity. There is a need for more research on the impact of human pressure on vertebrates' behaviour,

as such changes could result in marked shifts away from natural patterns, with consequences for

fitness, population persistence, community interactions, and evolution (Chace and Walsh, 2006;

Gaynor et al., 2018; Lin et al., 2012). The availability of camera trap information is increasing

exponentially, but usually the only information reported is in relation to species' community

composition and population densities (Antunes et al., 2022; Negret et al., 2015a). Our analysis

highlights the fact that many behavioural characteristics, such as species' daily activity, may vary

between species, individuals and sites and that this variation can have impacts on the ecology and

conservation of these species. Future studies should focus on the impact of human pressure on the

behavioural patterns of vertebrates (Gaynor et al., 2018; Sih et al., 2011), especially for birds and

other groups for which information in this regard is scarce (Bonier et al., 2007; Chace and Walsh,

2006; Fontúrbel et al., 2021; Smetzer et al., 2022) and for regions that are underrepresented. Our

findings are a step in this direction.

References:

Agostinelli, C., Lund, U., 2022. R package "circular": Circular Statistics (version 0.4-95).

Antunes, A.C., Montanarin, A., Gräbin, D.M., Monteiro, E.C. dos S., de Pinho, F.F., Alvarenga, G.C.,

Ahumada, J., Wallace, R.B., Ramalho, E.E., Barnett, A.P.A., Bager, A., Costa Lopes, A.M.,

Keuroghlian, A., Giroux, A., Herrera, A.M., Almeida Correa, A.P., Meiga, A.Y., Jácomo, A.T.

de A., Barros Barban, A., Antunes, A., Coelho, A.G. de A., Camilo, A.R., Nunes, A.V., Gomes,

A.C. dos S.M., Silva Zanzini, A.C., Castro, A.B., Desbiez, A.L.J., Figueiredo, A., Thoisy, B.,

Gauzens, B., Oliveira, B.T., Lima, C.A., Peres, C.A., Durigan, C.C., Brocardo, C.R., Rosa, C.A.,

Zárate☐Castañeda, C., Monteza☐Moreno, C.M., Carnicer, C., Trinca, C.T., Polli, D.J., Ferraz,

D. da S., Lane, D.F., Gomes da Rocha, D., Barcelos, D.C., Auz, D., Rosa, D.C.P., Silva, D.A.,

Silvério, D.V., Eaton, D.P., Nakano☐Oliveira, E., Venticinque, E., Carvalho Junior, E.,

Mendonça, E.N., Vieira, E.M., Isasi☐Catalá, E., Fischer, E., Castro, E.P., Oliveira, E.G., Melo,

F.R., Lima Muniz, F., Rohe, F., Baccaro, F.B., Michalski, F., Paim, F.P., Santos, F., Anaguano,

F., Palmeira, F.B.L., Reis, F. da S., Aguiar☐Silva, F.H., Batista, G. de A., Zapata☐Ríos, G.,

Forero☐Medina, G., De Souza Ferreira Neto, G., Alves, G.B., Ayala, G., Pedersoli, G.H.P., El

Bizri, H.R., Alves do Prado, H., Mozerle, H.B., Costa, H.C.M., Lima, I.J., Palacios, J., Assis, J.

de R., Boubli, J.P., Metzger, J.P., Teixeira, J.V., Miranda, J.M.D., Polisar, J., Salvador, J.,

Borges☐Almeida, K., Didier, K., Dayane de Lima Pereira, K., Torralvo, K., Gajapersad, K.,

Silveira, L., Maioli, L.U., Maracahipes☐Santos, L., Valenzuela, L., Benavalli, L., Fletcher, L.,

Paolucci, L.N., Zanzini, L.P., Zago da Silva, L., Ribeiro Rodrigues, L.C., Benchimol, M.,

Oliveira, M.A., Lima, M., Basto da Silva, M., Santos Junior, M.A., Viscarra, M., Cohn☐Haft,

M., Abrahams, M.I., Benedetti, M.A., Marmontel, M., Hirt, M.R., Tôrres, N.M., Cruz Junior,

O.F., Alvarez☐Loayza, P., Jansen, P., Prist, P.R., Brando, P.M., Bernardes Perônico, P., Leite,

R. do N., Rabelo, R.M., Sollmann, R., Beltrão☐Mendes, R., Ferreira, R.A.F., Coutinho, R.,

Oliveira, R. da C., Ilha, R., Hilário, R.R., Pires, R.A.P., Sampaio, R., Silva Moreira, R.,

Botero☐Arias, R., Vasquez Martinez, R., Nóbrega, R.A. de A., Fadini, R.F., Morato, R.G.,

Carneiro, R.L., Almeida, R.P.S., Ramos, R.M., Schaub, R., Dornas, R., Cueva, R., Rolim, S.,

Laurindo, S., Espinosa, S., Fernandes, T.N., Sanaiotti, T.M., Alvim, T.H.G., Dornas, T.T., Piña,

T.E.N., Caetano Andrade, V.L., Santiago, W.T.V., Magnusson, W.E., Campos, Z., Ribeiro,

M.C., 2022. AMAZONIA CAMTRAP: A dataset of mammal, bird, and reptile species recorded

with camera traps in the Amazon forest. Ecology e3738. https://doi.org/10.1002/ecy.3738

Bates, D., Mächler, M., Bolker, B., Walker, S., 2015. Fitting Linear Mixed-Effects Models Using

lme4. J. Stat. Softw. 67. https://doi.org/10.18637/jss.v067.i01

Benítez-López, A., Alkemade, R., Schipper, A.M., Ingram, D.J., Verweij, P.A., Eikelboom, J.A.J.,

Huijbregts, M.A.J., 2017. The impact of hunting on tropical mammal and bird populations.

Science (80-.). 356, 180–183. https://doi.org/10.1126/science.aaj1891

Bennie, J.J., Duffy, J.P., Inger, R., Gaston, K.J., 2014. Biogeography of time partitioning in mammals.

Proc. Natl. Acad. Sci. U. S. A. 111, 13727–13732. https://doi.org/10.1073/pnas.1216063110

Bertassoni, M.F., Abba, A.M., 2014. Myrmecophaga tridactyla [WWW Document]. IUCN Red List

Threat. Species. URL https://dx.doi.org/10.2305/IUCN.UK.2014-1.RLTS.T14224A47441961.en

(accessed 5.23.22).

Bonier, F., Martin, P.R., Wingfield, J.C., 2007. Urban birds have broader environmental tolerance.

Biol. Lett. 3, 670–673. https://doi.org/10.1098/rsbl.2007.0349

Burnham, K.P., Anderson, D.R., 2018. Model Selection and Multimodel inference: A Practical

Information-Theoretic Approach, 2nd ed, Sociological Methods and Research.

Chace, J.F., Walsh, J.J., 2006. Urban effects on native avifauna: A review. Landsc. Urban Plan. 74,

46–69. https://doi.org/10.1016/j.landurbplan.2004.08.007

Checa, M.F., Rodriguez, J., Willmott, K.R., Liger, B., 2014. Microclimate variability significantly

affects the composition, abundance and phenology of butterfly communities in a highly

threatened neotropical dry forest. Florida Entomol. 97, 1–13.

https://doi.org/10.1896/054.097.0101

Correa Ayram, C.A., Diaz-Timote, J., Etter, A., Ramirez, W., Corzo, G., 2019. Human Space

Footprint Index in Colombia, years 2015 and 2018 [WWW Document]. Alexander von

Humboldt Biol. Resour. Res. Inst. URL

http://geonetwork.humboldt.org.co/geonetwork/srv/spa/catalog.search;jsessionid=E36E885D872

F105C5F20E73670B68E29?node=srv#/metadata/3f37fa6b-5290-4399-9ea3-eaafcd0b2fbe

Correa Ayram, C.A., Etter, A., Díaz-Timoté, J., Rodríguez Buriticá, S., Ramírez, W., Corzo, G., 2020.

Spatiotemporal evaluation of the human footprint in Colombia: Four decades of anthropic

impact in highly biodiverse ecosystems. Ecol. Indic. 117, 106630.

https://doi.org/10.1016/j.ecolind.2020.106630

Dias, L.C.S., Bernardo, C.S.S., Srbek-Araujo, A.C., 2016. Daily and seasonal activity patterns of the

Solitary Tinamou (Tinamus solitarius) in the Atlantic Forest of Southeastern Brazil. Wilson J.

Ornithol. 128, 885–894. https://doi.org/10.1676/15-157.1

Diaz-Pulido, A., Restrepo, A., Rodríguez, R., Hernández, O.F., Lopera, A., Rosero, Y., Moreno, G.,

2022a. Piedemonte Meta Ecopetrol. Bogotá. Colombia.

Diaz-Pulido, A., Restrepo, A., Rodríguez, R., Hernández, O.F., Lopera, A., Rosero, Y., Moreno, G.,

2022b. Río Tillavá Ecopetrol. Bogota, Colombia.

Diaz-Pulido, A., Restrepo, A., Rojano, C., Miranda, L., Ávila, R., 2022c. Piedemonte Casanare

Ecopetrol. Bogota, Colombia.

Dirzo, R., Young, H.S., Galetti, M., Ceballos, G., Isaac, N.J.B., Collen, B., 2014. Defaunation in the

Anthropocene. Science (80-.). 345, 401–406. https://doi.org/10.1126/science.1251817

Emmons, L.H., Feer, F., 1997. Neotropical Rainforest Mammals: A Field Guide, Second Edi. ed.

University of Chicago Press.

Etter, A., McAlpine, C.A., Seabrook, L., Wilson, K.A., 2011. Incorporating temporality and

biophysical vulnerability to quantify the human spatial footprint on ecosystems. Biol. Conserv.

144, 1585–1594. https://doi.org/10.1016/j.biocon.2011.02.004

Ewers, R.M., Banks-Leite, C., 2013. Fragmentation Impairs the Microclimate Buffering Effect of

Tropical Forests. PLoS One 8. https://doi.org/10.1371/journal.pone.0058093

Fontúrbel, F.E., Orellana, J.I., Rodríguez-Gómez, G.B., Tabilo, C.A., Castaño-Villa, G.J., 2021.

Habitat disturbance can alter forest understory bird activity patterns: A regional-scale

assessment with camera-traps. For. Ecol. Manage. 479, 118618.

https://doi.org/10.1016/j.foreco.2020.118618

Gallagher, A.J., Creel, S., Wilson, R.P., Cooke, S.J., 2017. Energy Landscapes and the Landscape of

Fear. Trends Ecol. Evol. 32, 88–96. https://doi.org/10.1016/j.tree.2016.10.010

Gaynor, K.M., Hojnowski, C.E., Carter, N.H., Brashares, J.S., 2018. The influence of human

disturbance on wildlife nocturnality. Science (80-.). 360, 1232–1235.

https://doi.org/10.1126/science.aar7121

Haddad, N.M., Brudvig, L.A., Clobert, J., Davies, K.F., Gonzalez, A., Holt, R.D., Lovejoy, T.E.,

Sexton, J.O., Austin, M.P., Collins, C.D., Cook, W.M., Damschen, E.I., Ewers, R.M., Foster,

B.L., Jenkins, C.N., King, A.J., Laurance, W.F., Levey, D.J., Margules, C.R., Melbourne, B.A.,

Nicholls, A.O., Orrock, J.L., Song, D.X., Townshend, J.R., 2015. Habitat fragmentation and its

lasting impact on Earth's ecosystems. Sci. Adv. 1, 1–10. https://doi.org/10.1126/sciadv.1500052

Hilty, S.L., Brown, W.L., 2001. Guía de las Aves de Colombia.

IUCN, 2022. The IUCN Red List of Threatened Species [WWW Document]. URL

http://www.iucnredlist.org/

Jackson, M.T., 1966. Effects of Microclimate on Spring Flowering Phenology. Ecology 167–174.

Laundre, J.W., Hernandez, L., Ripple, W.J., 2010. The Landscape of Fear: Ecological Implications of

Being Afraid. Open Ecol. J. 3, 1–7. https://doi.org/10.2174/1874213001003030001

Lin, T., Coppack, T., Lin, Q.X., Kulemeyer, C., Schmidt, A., Behm, H., Luo, T., 2012. Does avian

flight initiation distance indicate tolerance towards urban disturbance? Ecol. Indic. 15, 30–35.

https://doi.org/10.1016/j.ecolind.2011.09.018

Mendes, C.P., Carreira, D., Pedrosa, F., Beca, G., Lautenschlager, L., Akkawi, P., Bercê, W., Ferraz,

K.M.P.M.B., Galetti, M., 2020. Landscape of human fear in Neotropical rainforest mammals.

Biol. Conserv. 241, 108257. https://doi.org/10.1016/j.biocon.2019.108257

Meneguetti, D.U. de O., Zan, R.A., Ramos, L.J., 2014. First report of occurance running over of

tayras (Eira barbara Linnaeus, 1758 - carnivore, mustelidae) in the Jamari valley, Rondônia,

western Amazon. Colégio Apl. da Univ. Fed. do Acre 1, 190–194.

Michalski, F., Peres, C.A., 2005. Anthropogenic determinants of primate and carnivore local

extinctions in a fragmented forest landscape of southern Amazonia. Biol. Conserv. 124, 383–

396. https://doi.org/10.1016/j.biocon.2005.01.045

Myers, P., Espinosa, R.., Parr, C.S., Jones, T., Hammond, G.S., Dewey, T.A., 2022. The Animal

Diversity Web [WWW Document]. URL https://animaldiversity.org

Negret, P.J., Garzon, O., Stevenson, P.R., 2015a. First preliminary inventory of Non-Flying Mammals

of the Alto Fragua Indi-Wasi National Park, Colombia. Mammology Notes 2, 32–35.

Negret, P.J., Garzón, O., Stevenson, P.R., Laverde-, O., Gar On, O., 2015b. New ecological

information for the Black Tinamou (Tinamus osgoodi hershkovitzi). Auk 132, 533–539.

https://doi.org/10.1642/AUK-14-116.1

O'Donnell, K., Delbarco-Trillo, J., 2020. Changes in the home range sizes of terrestrial vertebrates in

response to urban disturbance: a meta-analysis. J. Urban Ecol. 6, 1–8.

https://doi.org/10.1093/jue/juaa014

Ocampo, D., Borja-Acosta, K.G., Lozano-Flórez, J., Cifuentes-Acevedo, S., Arbeláez-Cortés, E.,

Bayly, N.J., Caguazango, Á., Coral-Jaramillo, B., Cueva, D., Forero, F., Gómez, J.P., Gómez,

C., Loaiza-Muñoz, M.A., Londoño, G.A., Losada-Prado, S., Pérez-Peña, S., Ramírez-Chaves,

H.E., Rodríguez-Posada, M.E., Sanabria-Mejía, J., Sánchez-Martínez, M., Serrano-Cardozo,

V.H., Sierra-Buitrago, M.S., Soto-Patiño, J., Acevedo-Charry, O., 2021. Body mass data set for

1,317 bird and 270 mammal species from Colombia. Ecology 102, 13473147.

https://doi.org/10.1002/ecy.3273

Outhwaite, C.L., McCann, P., Newbold, T., 2022. Agriculture and climate change are reshaping insect

biodiversity worldwide. Nature 605. https://doi.org/10.1038/s41586-022-04644-x

Pardo, L.E., Edwards, W., Campbell, M.J., Gómez-Valencia, B., Clements, G.R., Laurance, W.F.,

2021. Effects of oil palm and human presence on activity patterns of terrestrial mammals in the

Colombian Llanos. Mamm. Biol. 101, 775–789. https://doi.org/10.1007/s42991-021-00153-y

R Core Team, 2022. R: A Language and Environment for Statistical Computing. R Foundation for

Statistical Computing.

Redpath, S.M., 1995. Impact of habitat fragmentation on activity and hunting behavior in the tawny

owl, Strix aluco. Behav. Ecol. 6, 410–413. https://doi.org/10.1093/beheco/6.4.410

Saunders, D.A., 1980. Food and movements of the short-billed form of the white-tailed black

cockatoo. Wildl. Res. 7, 257–269. https://doi.org/10.1071/WR9800257

Schmidt, M., Jochheim, H., Kersebaum, K.C., Lischeid, G., Nendel, C., 2017. Gradients of

microclimate, carbon and nitrogen in transition zones of fragmented landscapes – a review.

Agric. For. Meteorol. 232, 659–671. https://doi.org/10.1016/j.agrformet.2016.10.022

Sih, A., Ferrari, M.C.O., Harris, D.J., 2011. Evolution and behavioural responses to human-induced

rapid environmental change. Evol. Appl. 4, 367–387. https://doi.org/10.1111/j.1752-

4571.2010.00166.x

Smetzer, J.R., Paxton, K.L., Hart, P.J., Paxton, E.H., 2022. Activity patterns of Hawaiian forest birds

in a fragmented and continuous landscape. J. Avian Biol. 1–15.

https://doi.org/10.1111/jav.02905

U.S. Naval Observatory, 2022. Astronomical Applications Department [WWW Document]. URL

https://aa.usno.navy.mil/data/RS_OneYear

Venter, O., Sanderson, E.W., Magrach, A., Allan, J.R., Beher, J., Jones, K.R., Possingham, H.P.,

Laurance, W.F., Wood, P., Fekete, B.M., Levy, M.A., Watson, J.E.M., 2016. Sixteen years of

change in the global terrestrial human footprint and implications for biodiversity conservation.

Nat. Commun. 7, 1–11. https://doi.org/10.1038/ncomms12558

Vitasse, Y., Baumgarten, F., Zohner, C.M., Kaewthongrach, R., Fu, Y.H., Walde, M.G., Moser, B.,

2021. Impact of microclimatic conditions and resource availability on spring and autumn

phenology of temperate tree seedlings. New Phytol. 232, 537–550.

https://doi.org/10.1111/nph.17606

Wickham, H., 2016. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag, New York.

Springer, New York.

Williams, B.A., Grantham, H.S., Watson, J.E.M., Alvarez, S.J., Simmonds, J.S., Rogéliz, C.A., Da

Silva, M., Forero-Medina, G., Etter, A., Nogales, J., Walschburger, T., Hyman, G., Beyer, H.L.,

2020a. Minimising the loss of biodiversity and ecosystem services in an intact landscape under

risk of rapid agricultural development. Environ. Res. Lett. 15. https://doi.org/10.1088/1748-

9326/ab5ff7

Williams, B.A., Venter, O., Allan, J.R., Atkinson, S.C., Rehbein, J.A., Ward, M., Di Marco, M.,

Grantham, H.S., Ervin, J., Goetz, S.J., Hansen, A.J., Jantz, P., Pillay, R., Rodríguez-Buriticá, S.,

Supples, C., Virnig, A.L.S., Watson, J.E.M., 2020b. Change in Terrestrial Human Footprint

Drives Continued Loss of Intact Ecosystems. One Earth 3, 371–382.

https://doi.org/10.1016/j.oneear.2020.08.009

Zanette, L.Y., Clinchy, M., 2019. Ecology of fear. Curr. Biol. 29, R309–R313.

https://doi.org/10.1016/j.cub.2019.02.042

Figures and Tables:

Table 1. Main results of the regression analysis, testing the effect of human pressure on the diurnality of each mammal and bird species assessed. Significant
results are in bold.

Class	Species	Common name	Spatial Scale	Estimate	P value	Z value	Sample Size (Cameras)	N. Records
Mammals	*Cerdocyon thous*	Crab-eating fox	500	0.03	0.244	1.164	56	293
	Cuniculus paca	**Paca**	500	-0.07	**<<0.01**	-6.156	107	621
	Dasyprocta fuliginosa	**Agouti**	1000	-0.04	**<<0.01**	-3.838	71	371
	Dasypus novemcinctus	**Nine-banded armadillo**	500	-0.03	**0.046**	-1.999	137	715
	Didelphis marsupialis	**Common opossum**	500	-0.10	**<<0.01**	-7.977	171	1030
	Eira barbara	**Tayra**	500	0.04	**0.006**	2.767	73	193
	Leopardus pardalis	Ocelot	1000	0.00	0.895	0.132	60	83
	Mazama americana	Red brocket	500	0.02	0.330	0.974	41	117
	Myrmecophaga tridactyla	Giant anteater	1000	-0.01	0.324	-0.986	102	283
	Nasua nasua	Ring tailed coati	1000	0.04	0.295	1.048	19	29
	Odocoileus virginianus	White-tailed deer	500	-0.02	0.149	-1.442	35	76
	Pecari tajacu	Collared peccary	250	0.01	0.693	0.395	33	78
	Philander opossum	**Gray four-eyed opossum**	1000	-0.08	**<<0.01**	-5.035	58	479
	Tamandua tetradactyla	**Collared anteater**	250	-0.03	**<<0.01**	-3.868	164	362
Birds	*Aramides cajaneus*	Grey-necked wood rail	1000	0.01	0.115	1.578	114	1044
	Arremon taciturnus	Pectoral sparrow	250	-0.02	0.240	-1.176	23	77
	Catharus ustulatus	**Swainson's thrush**	500	0.10	**<<0.01**	4.608	51	546
	Crypturellus cinereus	Cinereous tinamou	250	-0.01	0.260	-1.127	109	428
	Crypturellus soui	**Little tinamou**	250	0.07	**0.003**	2.948	48	188
	Leptotila rufaxilla	Grey-fronted dove	1000	0.02	0.239	1.176	124	1451
	Leptotila verreauxi	**White-tipped dove**	1000	0.09	**<<0.01**	4.331	87	647
	Momotus momota	**Amazonian motmot**	500	0.05	**0.011**	2.535	56	176
	Myrmoborus myotherinus	Black-faced antbird	500	0.10	0.416	0.813	6	20
	Nyctidromus albicollis	**Pauraque**	250	-0.05	**<<0.01**	-4.963	48	318
	Vanellus chilensis	Southern lapwing	1000	-0.94	0.317	-1.000	4	17

Figure 1. (a) In red the location of the study areas in Colombia. In grey the llanos region. Location of the three sites where cameras were deployed in the study area: **(b)** Meta Foothills. **(c)** Casanare Foothills. **(d)** Tillavá River. Camera trap stations are depicted by purple circles. The level of human pressure in the study area is depicted using the Legacy-adjusted Human Footprint Index (LHFI) for 2018. Red colours depict areas of high human pressure while blue colours areas of low human pressure.

487 a)

Figure 2. Circadian activity of **(a)** seven mammals and **(b)** five bird species in three areas in the
Colombian llanos. White bars represent camaras deployed in more preserved sited, and dark grey bars
represent camaras deployed in more disturbed areas. Light grey is the overlap in the circadian activity
between the more preserved and more disturbed camara sites. For this figure, the location where a
camara was deployed was considered disturbed when it presented a Human Footprint index above the
mean, considering only the sited where camaras where deployed that had recordings of the species.
The species depicted are the ones that presented statistically significant changes in diurnality based on

a logistic regression. Mammal and bird silhouettes were obtained from the PhyloPic public domain
database (http://phylopic.org/).

Figure 3. Shifts in diurnality of **(a)** seven mammals and **(b)** five bird species in response to human pressure in three sites in the Colombian llanos, from the most preserved sites (black point) to the most disturbed sites (arrow point) in which each species was recorded. Arrows to the right represent shifts toward diurnality, while arrows to the left represent shifts toward nocturnality. The species depicted are the ones that presented statistically significant changes in diurnality based on a logistic regression. Mammal and bird silhouettes were obtained from the PhyloPic public domain database (http://phylopic.org/).

561 **a)**

Figure 4. Changes in diurnal activity for **(a)** seven mammals and **(b)** five bird species with increases
in human pressure. The human pressure (x-axis) was measured using the Legacy-adjusted Human
Footprint Index (LHFI) for 2018 of each site as a proxy (Correa Ayram et al., 2019). The species
depicted are the ones that presented statistically significant changes in diurnality based on a logistic
regression. Mammal and bird silhouettes were obtained from the PhyloPic public domain database
(http://phylopic.org/).
586

   Supplementary material

   SI Table 1. Mammal, bird and reptile species recorded with the camera traps in each of the three sites
   (Casanare, Meta and Tillava) in the Llanos region in Colombia.

Species	Class	Sites	Number of Records
Aramides cajaneus	Aves	Casanare	279
Arremon axillaris	Aves	Casanare	26
Arremon brunneinucha	Aves	Casanare	22
Arremon taciturnus	Aves	Casanare	2
Atlapetes semirufus	Aves	Casanare	1
Bubulcus ibis	Aves	Casanare	4
Buteogallus schistaceus	Aves	Casanare	1
Cacicus cela	Aves	Casanare	2
Cantorchilus leucotis	Aves	Casanare	2
Catharus aurantiirostris	Aves	Casanare	5
Catharus dryas	Aves	Casanare	18
Catharus minimus	Aves	Casanare	2
Catharus ustulatus	Aves	Casanare	187
Ceratopipra erythrocephala	Aves	Casanare	3
Chamaepetes goudotii	Aves	Casanare	1
Chamaeza campanisona	Aves	Casanare	26
Chamaeza turdina	Aves	Casanare	8
Coereba flaveola	Aves	Casanare	1
Crotophaga ani	Aves	Casanare	1
Crotophaga major	Aves	Casanare	11
Crypturellus cinereus	Aves	Casanare	61
Crypturellus soui	Aves	Casanare	127
Cyanocorax violaceus	Aves	Casanare	25
Cyanocorax yncas	Aves	Casanare	3
Dendrocincla fuliginosa	Aves	Casanare	1
Diglossa cyanea	Aves	Casanare	1
Dromococcyx pavoninus	Aves	Casanare	1
Eurypyga helias	Aves	Casanare	7
Gallus gallus domesticus	Aves	Casanare	71

Leptotila rufaxilla	Aves	Casanare	1232
Leptotila verreauxi	Aves	Casanare	580
Malacoptila mystacalis	Aves	Casanare	2
Mesembrinibis cayennensis	Aves	Casanare	3
Micrastur semitorquatus	Aves	Casanare	1
Mitu tomentosum	Aves	Casanare	11
Momotus momota	Aves	Casanare	131
Myioborus miniatus	Aves	Casanare	1
Myrmeciza longipes	Aves	Casanare	5
Myrmoborus leucophrys	Aves	Casanare	7
Nycticorax nycticorax	Aves	Casanare	2
Nyctidromus albicollis	Aves	Casanare	212
Odontophorus gujanensis	Aves	Casanare	1
Odontophorus strophium	Aves	Casanare	1
Ortalis ruficauda	Aves	Casanare	95
Pachyramphus rufus	Aves	Casanare	1
Patagioenas speciosa	Aves	Casanare	1
Penelope argyrotis	Aves	Casanare	1
Penelope jacquacu	Aves	Casanare	2
Pheugopedius rutilus	Aves	Casanare	1
Phimosus infuscatus	Aves	Casanare	2
Piaya cayana	Aves	Casanare	3
Piranga rubra	Aves	Casanare	1
Psarocolius decumanus	Aves	Casanare	16
Pteroglossus castanotis	Aves	Casanare	1
Quiscalus lugubris	Aves	Casanare	3
Rupornis magnirostris	Aves	Casanare	8
Spinus psaltria	Aves	Casanare	1
Tangara palmarum	Aves	Casanare	1
Taraba major	Aves	Casanare	2
Thryophilus rufalbus	Aves	Casanare	4
Tinamus guttatus	Aves	Casanare	1
Tinamus tao	Aves	Casanare	14
Turdus albicollis	Aves	Casanare	2

Turdus debilis	Aves	Casanare	1
Turdus ignobilis	Aves	Casanare	1
Turdus leucomelas	Aves	Casanare	95
Turdus nudigenis	Aves	Casanare	14
Zentrygon linearis	Aves	Casanare	170
Bos taurus	Mammalia	Casanare	207
Caluromys lanatus	Mammalia	Casanare	3
Canis familiaris	Mammalia	Casanare	192
Cerdocyon thous	Mammalia	Casanare	81
Coendou prehensilis	Mammalia	Casanare	12
Cuniculus paca	Mammalia	Casanare	388
Dasyprocta fuliginosa	Mammalia	Casanare	235
Dasypus novemcinctus	Mammalia	Casanare	425
Didelphis marsupialis	Mammalia	Casanare	482
Eira barbara	Mammalia	Casanare	124
Equus caballus	Mammalia	Casanare	21
Felis catus	Mammalia	Casanare	96
Galictis vittata	Mammalia	Casanare	6
Herpailurus yagouaroundi	Mammalia	Casanare	10
Homo sapiens	Mammalia	Casanare	259
Leopardus pardalis	Mammalia	Casanare	46
Leopardus wiedii	Mammalia	Casanare	7
Lontra longicaudis	Mammalia	Casanare	1
Marmosa robinsoni	Mammalia	Casanare	61
Mazama americana	Mammalia	Casanare	117
Metachirus nudicaudatus	Mammalia	Casanare	90
Myrmecophaga tridactyla	Mammalia	Casanare	82
Nasua nasua	Mammalia	Casanare	14
Odocoileus virginianus	Mammalia	Casanare	41
Pecari tajacu	Mammalia	Casanare	62
Procyon cancrivorus	Mammalia	Casanare	7
Proechimys canicollis	Mammalia	Casanare	1
Proechimys oconnelli	Mammalia	Casanare	6
Puma concolor	Mammalia	Casanare	5

Sapajus apella	Mammalia	Casanare	15
Sciurus granatensis	Mammalia	Casanare	63
Tamandua tetradactyla	Mammalia	Casanare	178
Urocyon cinereoargenteus	Mammalia	Casanare	1
Ameiva ameiva	Reptilia	Casanare	4
Iguana iguana	Reptilia	Casanare	2
Anabacerthia striaticollis	Aves	Meta	1
Anas platyrhynchos domesticus	Aves	Meta	4
Aramides cajaneus	Aves	Meta	762
Ardea alba	Aves	Meta	1
Arremon taciturnus	Aves	Meta	75
Arremonops conirostris	Aves	Meta	9
Athene cunicularia	Aves	Meta	3
Basileuterus culicivorus	Aves	Meta	9
Bubulcus ibis	Aves	Meta	80
Burhinus bistriatus	Aves	Meta	35
Buteogallus meridionalis	Aves	Meta	2
Cantorchilus leucotis	Aves	Meta	12
Caracara cheriway	Aves	Meta	6
Catharus ustulatus	Aves	Meta	359
Cercomacroides tyrannina	Aves	Meta	1
Chamaeza campanisona	Aves	Meta	1
Chordeiles pusillus	Aves	Meta	1
Cochlearius cochlearius	Aves	Meta	1
Coragyps atratus	Aves	Meta	3
Crotophaga ani	Aves	Meta	8
Crotophaga major	Aves	Meta	19
Crypturellus cinereus	Aves	Meta	327
Crypturellus soui	Aves	Meta	60
Cyanocorax violaceus	Aves	Meta	43
Dendrocincla fuliginosa	Aves	Meta	18
Dysithamnus mentalis	Aves	Meta	1
Egretta caerulea	Aves	Meta	1
Eudocimus ruber	Aves	Meta	1

Eurypyga helias	Aves	Meta	1
Florisuga mellivora	Aves	Meta	1
Gallus gallus	Aves	Meta	408
Geothlypis philadelphia	Aves	Meta	1
Geotrygon montana	Aves	Meta	34
Gymnomystax mexicanus	Aves	Meta	10
Leptotila rufaxilla	Aves	Meta	216
Leptotila verreauxi	Aves	Meta	65
Manacus manacus	Aves	Meta	2
Megascops choliba	Aves	Meta	3
Meleagris gallopavo	Aves	Meta	1
Mesembrinibis cayennensis	Aves	Meta	89
Microcerculus marginatus	Aves	Meta	1
Milvago chimachima	Aves	Meta	4
Mitu tomentosum	Aves	Meta	2
Momotus momota	Aves	Meta	37
Myiothlypis fulvicauda	Aves	Meta	1
Myrmoborus leucophrys	Aves	Meta	8
Myrmoborus myotherinus	Aves	Meta	20
Myrmophylax atrothorax	Aves	Meta	7
Numida meleagris	Aves	Meta	3
Nyctidromus albicollis	Aves	Meta	105
Odontophorus gujanensis	Aves	Meta	9
Ortalis guttata	Aves	Meta	30
Penelope argyrotis	Aves	Meta	1
Penelope jacquacu	Aves	Meta	41
Phaethornis atrimentalis	Aves	Meta	3
Phaethornis griseogularis	Aves	Meta	3
Pheugopedius rutilus	Aves	Meta	2
Phimosus infuscatus	Aves	Meta	11
Piaya cayana	Aves	Meta	3
Pitangus sulphuratus	Aves	Meta	3
Psarocolius angustifrons	Aves	Meta	2
Psarocolius decumanus	Aves	Meta	9

Ramphocelus carbo	Aves	Meta	4
Rupornis magnirostris	Aves	Meta	31
Saltator maximus	Aves	Meta	11
Setophaga ruticilla	Aves	Meta	2
Sporophila intermedia	Aves	Meta	2
Sturnella magna	Aves	Meta	1
Synallaxis albescens	Aves	Meta	1
Syrigma sibilatrix	Aves	Meta	17
Tachyphonus rufus	Aves	Meta	1
Tangara episcopus	Aves	Meta	1
Tersina viridis	Aves	Meta	1
Tigrisoma lineatum	Aves	Meta	2
Troglodytes aedon	Aves	Meta	2
Turdus ignobilis	Aves	Meta	2
Turdus leucomelas	Aves	Meta	38
Turdus nudigenis	Aves	Meta	19
Vanellus chilensis	Aves	Meta	17
Veniliornis passerinus	Aves	Meta	1
Zentrygon linearis	Aves	Meta	37
Bos taurus	Mammalia	Meta	764
Cabassous unicinctus	Mammalia	Meta	9
Caluromys lanatus	Mammalia	Meta	3
Canis familiaris	Mammalia	Meta	176
Cerdocyon thous	Mammalia	Meta	201
Choloepus didactylus	Mammalia	Meta	3
Coendou prehensilis	Mammalia	Meta	6
Cuniculus paca	Mammalia	Meta	173
Dasyprocta fuliginosa	Mammalia	Meta	28
Dasypus kappleri	Mammalia	Meta	9
Dasypus novemcinctus	Mammalia	Meta	288
Didelphis marsupialis	Mammalia	Meta	523
Eira barbara	Mammalia	Meta	65
Equus caballus	Mammalia	Meta	212
Equus mulus	Mammalia	Meta	11

Felis catus	Mammalia	Meta	68
Galictis vittata	Mammalia	Meta	1
Herpailurus yagouaroundi	Mammalia	Meta	2
Homo sapiens	Mammalia	Meta	386
Leopardus pardalis	Mammalia	Meta	30
Leopardus wiedii	Mammalia	Meta	3
Myrmecophaga tridactyla	Mammalia	Meta	198
Nasua nasua	Mammalia	Meta	15
Odocoileus virginianus	Mammalia	Meta	6
Philander opossum	Mammalia	Meta	479
Plecturocebus ornatus	Mammalia	Meta	3
Priodontes maximus	Mammalia	Meta	1
Procyon cancrivorus	Mammalia	Meta	14
Puma concolor	Mammalia	Meta	3
Saimiri sciureus	Mammalia	Meta	73
Sapajus apella	Mammalia	Meta	37
Sciurus granatensis	Mammalia	Meta	90
Sciurus igniventris	Mammalia	Meta	6
Tamandua tetradactyla	Mammalia	Meta	156
Ameiva ameiva	Reptilia	Meta	2
Iguana iguana	Reptilia	Meta	15
Aramides cajaneus	Aves	Tillava	3
Buteogallus meridionalis	Aves	Tillava	1
Caracara cheriway	Aves	Tillava	3
Catharus minimus	Aves	Tillava	1
Crypturellus cinereus	Aves	Tillava	40
Crypturellus soui	Aves	Tillava	1
Dendroplex picus	Aves	Tillava	1
Dryocopus lineatus	Aves	Tillava	1
Geotrygon montana	Aves	Tillava	3
Leptotila rufaxilla	Aves	Tillava	3
Leptotila verreauxi	Aves	Tillava	2
Momotus momota	Aves	Tillava	8
Nyctidromus albicollis	Aves	Tillava	1

Phaethornis griseogularis	Aves	Tillava	1
Ramphocelus carbo	Aves	Tillava	1
Rupornis magnirostris	Aves	Tillava	3
Saltator maximus	Aves	Tillava	1
Tinamus guttatus	Aves	Tillava	2
Tinamus major	Aves	Tillava	1
Turdus albicollis	Aves	Tillava	1
Bos taurus	Mammalia	Tillava	20
Canis familiaris	Mammalia	Tillava	1
Cerdocyon thous	Mammalia	Tillava	11
Cuniculus paca	Mammalia	Tillava	60
Dasyprocta fuliginosa	Mammalia	Tillava	108
Dasypus novemcinctus	Mammalia	Tillava	2
Didelphis marsupialis	Mammalia	Tillava	25
Eira barbara	Mammalia	Tillava	4
Equus caballus	Mammalia	Tillava	1
Homo sapiens	Mammalia	Tillava	57
Hydrochoerus hydrochaeris	Mammalia	Tillava	1
Leopardus pardalis	Mammalia	Tillava	7
Marmosa robinsoni	Mammalia	Tillava	4
Myrmecophaga tridactyla	Mammalia	Tillava	2
Odocoileus hemionus	Mammalia	Tillava	1
Odocoileus virginianus	Mammalia	Tillava	29
Pecari tajacu	Mammalia	Tillava	16
Priodontes maximus	Mammalia	Tillava	4
Puma concolor	Mammalia	Tillava	3
Sapajus apella	Mammalia	Tillava	3
Sciurus granatensis	Mammalia	Tillava	2
Tamandua tetradactyla	Mammalia	Tillava	28
Tapirus terrestris	Mammalia	Tillava	7

Class	Species	Common name	Diet	Body size in kg (Animal Diversity Web 2022)	Body size in kg (Emmons 1997)	Body size in kg (Ocampo et al. 2021)	Game Species	Activity	Nocturnal records (%)	Mean Activity Time	Rho	N. Records	N. Sites
Mammals	*Cerdocyon thous*	Crab-eating fox	Omnivore	5 - 8	3.6 - 7.9	3.45	No	N	88.1	23:43	0.46	293	56
	Cuniculus paca	**Paca**	Herbivore	7 - 12	5 - 13	6.119	Yes	N	93.56	23:16	0.65	621	107
	Dasyprocta fuliginosa	**Agouti**	Herbivore	1.3 - 4	3.5 - 6	3.048	Yes	D	13.48	12:50	0.46	371	71
	Dasypus novemcinctus	**Nine-banded armadillo**	Omnivore	3.6 - 7.7	2.7 - 6.3	4.73	Yes	N	95.94	23:15	0.69	715	137
	Didelphis marsupialis	**Common opossum**	Omnivore	1.5	0.6 - 1.6	0.232	Yes**	D	96.8	23:45	0.62	1030	171
	Eira barbara	**Tayra**	Omnivore	3 - 6	2.7 - 7		No	D	11.4	11:37	0.53	193	73
	Leopardus pardalis	Ocelot	Carnivore	8.5 - 16	8 - 9	13.5	Yes*	N	79.52	00:04	0.4	83	60
	Mazama americana	Red brocket	Herbivore	20 - 55	24 - 48		Yes	C	40.17	11:49	0.09	117	41
	Myrmecophaga tridactyla	Giant anteater	Insectivore	18 - 39	22 - 39		Yes**	C	70.57	23:36	0.36	283	102
	Nasua nasua	Ring tailed coati	Omnivore	3 - 6	3 - 7.2	2.825	No	D	17.24	13:37	0.41	29	19
	Odocoileus virginianus	White-tailed deer	Herbivore	57 - 137	30 - 50		Yes	C	31.58	10:27	0.23	76	35
	Pecari tajacu	Collared peccary	Omnivore	15 - 25	17 - 35		Yes	D	15.38	11:09	0.53	78	33
	Philander opossum	**Gray four-eyed opossum**	Omnivore	0.2 - 0.7	0.2 - 0.6	0.405	No	N	94.36	23:12	0.58	479	58
	Tamandua tetradactyla	**Collared anteater**	Insectivore	4.5	3.6 - 8.4		No	N	87.02	00:28	0.56	362	164
Birds	*Aramides cajaneus*	Grey-necked wood rail	Omnivore			0.42	No	D	11.3	11:40	0.5	1044	114
	Arremon taciturnus	Pectoral sparrow	Omnivore			0.03	No	D	12.99	11:41	0.46	77	23
	Catharus ustulatus	**Swainson's thrush**	Omnivore			0.03	No	D	5.13	11:21	0.61	546	51
	Crypturellus cinereus	Cinereous tinamou	Omnivore			0.49	Yes**	D	5.37	11:06	0.55	428	109
	Crypturellus soui	**Little tinamou**	Omnivore			0.20	Yes**	D	5.32	11:17	0.52	188	48
	Leptotila rufaxilla	Grey-fronted dove	Herbivore			0.14	No	D	2.34	11:25	0.65	1451	124
	Leptotila verreauxi	**White-tipped dove**	Herbivore			0.13	No	D	2.01	11:05	0.68	647	87
	Momotus momota	**Amazonian motmot**	Omnivore			0.11	No	D	2.84	11:30	0.61	176	56
	Myrmoborus myotherinus	Black-faced antbird	Insectivore			0.02	No	D	50	13:38	0.17	20	6
	Nyctidromus albicollis	**Pauraque**	Insectivore			0.05	No	C	75.79	19:49	0.23	318	48
	Vanellus chilensis	Southern lapwing	Omnivore			0.26	No	D	29.41	13:26	0.32	17	4

599