Genetic risk for Multiple Sclerosis originated in Pastoralist Steppe populations

Authors

William Barrie[1]§, Yaoling Yang[2,3]§, Kathrine E. Attfield[4]§, Evan Irving-Pease[5]§, Gabriele Scorrano[5]§, Lise Torp Jensen[6]§, Angelos P. Armen[4], Evangelos Antonios Dimopoulos[7], Aaron Stern[8], Alba Refoyo-Martinez[5], Abigail Ramsøe[5], Charleen Gaunitz[5], Fabrice Demeter[5], Marie Louise S. Jørkov[9], Stig Bermann Møller[10], Bente Springborg[10], Lutz Klassen[11], Inger Marie Hyldgård[11], Niels Wickmann[12], Lasse Vinner[5], Thorfinn Sand Korneliussen[5], Martin Sikora[5], Kristian Kristiansen[5,13], Santiago Rodriguez[3], Rasmus Nielsen[5,8], Astrid K. N. Iversen[4]@, Daniel J. Lawson[2,3]*@, Lars Fugger[4,14]*@, Eske Willerslev[1,5]*@

Affiliations

[1]Department of Zoology, University of Cambridge, Cambridge, UK,
[2]Department of Statistical Sciences, School of Mathematics, University of Bristol, Bristol, UK,
[3]Integrative Epidemiology Unit, Population Health Sciences, University of Bristol, Bristol, UK,
[4]Oxford Centre for Neuroinflammation, Nuffield Department of Clinical Neurosciences, John Radcliffe Hospital, University of Oxford, Oxford, UK
[5]Lundbeck Foundation GeoGenetics Centre, Globe Institute, University of Copenhagen, Copenhagen, Denmark,
[6]Department of Clinical Medicine, Aarhus University Hospital, Aarhus, Denmark
[7]Pathogen Genomics and Evolution Group, Department of Veterinary Medicine, University of Cambridge, Cambridge, UK
[8]Department of Integrative Biology, University of California, Berkeley
[9]Laboratory of Biological Anthropology, Department of Forensic Medicine, University of Copenhagen, Denmark
[10]Ålborg Historiske Museum, Nordjyske Museer, Vang Mark 25, 9380 Vestbjerg, Denmark
[11]Museum Østdanmark - Djursland og Randers. Stemannsgade 2, DK-8900 Randers C, Denmark
[12]Museum Vestsjælland, Forten 10, 4300 Holbæk, Denmark
[13]Department of Historical Studies, University of Gothenburg, SE-41255, Gothenburg, Sweden
[14]MRC Human Immunology Unit, John Radcliffe Hospital, University of Oxford, Oxford, UK

* Corresponding authors; email: Dan.Lawson@bristol.ac.uk, lars.fugger@ndcn.ox.ac.uk, ew482@cam.ac.uk
§ Joint first authors
@ Joint last authors

SUMMARY

Multiple sclerosis (MS) is a modern neuro-inflammatory and -degenerative disease, which is most prevalent in Northern Europe. Whilst it is known that inherited risk to MS is located within or within close proximity to immune genes it is unknown when, where and how this genetic risk originated. By using the largest ancient genome dataset from the Stone Age, along with new Medieval and post-

Medieval genomes, we show that many of the genetic risk variants for MS rose to higher frequency
among pastoralists located on the Pontic Steppe, and were brought into Europe by the Yamnaya-
related migration approximately 5,000 years ago. We further show that these MS-associated
immunogenetic variants underwent positive selection both within the Steppe population, and later in
Europe, likely driven by pathogenic challenges coinciding with dietary and lifestyle environmental
changes. This study highlights the critical importance of this period as a determinant of modern
immune responses and its subsequent impact on the risk of developing MS in a changing
environment.
52
<u>INTRODUCTION</u>
Multiple sclerosis (MS) is an autoimmune disease of the brain and spinal cord that currently affects
more than 2.5 million people worldwide. The prevalence varies markedly with ethnicity and
geographical location, with the highest prevalence observed in Europe (142.81 per 100.000 people),
and Northern Europeans being particularly susceptible to developing the disease[1]. The origins and
reasons for the geographical variation are poorly understood, yet such biases may hold important
clues as to why the prevalence of autoimmune diseases, including MS, has continued to rise during
the last 50 years.
61
62   While still elusive, MS etiology is thought to involve gene-gene and gene-environmental interactions.
63   Accumulating evidence suggests that exogenous triggers initiate a cascade of events involving a
64   multitude of cells and immune pathways in genetically vulnerable individuals, which may ultimately
65   lead to MS neuropathology[2].
66
Genome-wide association studies have identified 233 commonly occurring genetic variants that are
associated with MS; 32 variants are located in the HLA region and 201 outside the HLA region[3]. The
strongest MS associations are found in the HLA region with the most prominent of these, HLA-
DRB1*15:01, conferring an approximately three-fold increase in the risk of MS. Collectively, genetic
factors are estimated to explain approximately 30% of the overall disease risk, while environmental
and lifestyle factors are considered the major contributors to MS. Such determinants may include
geographically varying exposures like infections and low sun exposure/vitamin D deficiency. For
instance, while infection with Epstein-Barr virus frequently occurs in childhood and usually is
symptomless, delayed infection into early adulthood, as typically observed in countries with high
standards of hygiene, is associated with a 32-fold increased risk of MS[4,5]. Lifestyle factors associated
with increased MS risk such as smoking, obesity during adolescence, and nutrition/gut health also
vary geographically[6]. Dysregulations including autoimmunity in modern immune systems could also
result from the absence of ancient immunological triggers to which humans have evolutionarily
adapted, for instance by disturbing the delicate balance of pro- and anti-inflammatory pathways[7].
81
European ancestry has been postulated to explain part of the global difference in MS prevalence
globally in admixed populations[8]. Specifically, cases in African Americans exhibit increased
European ancestry in the HLA region compared to controls, with European haplotypes conferring
more MS risk for most HLA alleles, including HLA-DRB1*15:01. Conversely, Asian American cases
have decreased European ancestry in the HLA region compared to controls. Although Ancient
European ancestry and MS risk in Europe are known to be geographically structured (Figure 1a-b),
the effect of ancestry variation within Europe on MS prevalence is unknown.
89
Modern ancestry is viewed as a mixture of genetic ancestries derived from ancient populations, who
can be distinguished by their subsistence lifestyle: Western Hunter-Gatherers (WHG), Eastern

Hunter-Gatherers (EHG), Caucasus Hunter-Gatherers (CHG), Anatolian Farmers, and Steppe
Pastoralists (Figure 1c-d). Using the largest ancient genome dataset from the Stone Age, presented in
the accompanying study 'Population Genomics of Stone Age Eurasia'[9], coupled with new Medieval
and post-Medieval genomes, we quantified modern European ancestry with respect to these ancient
ancestries to identify signals of lifestyle-specific evolution. Then we determined whether the variants
associated with an increased risk for MS have undergone positive selection. We asked when selection
occurred and whether the targets of selection were specific to diet and lifestyle. Finally, we examined
the environmental conditions that may have caused selection for risk variants, including human
subsistence practice and exposure to pathogens.
101
102

Figure 1: Population history of Europe is associated with modern-day distribution of MS.
a) Modern-day geographical distribution of MS in Europe. Prevalence data for MS (cases per
100,000) was obtained from [1]. b) Steppe ancestry in modern samples as estimated by [10]. c-d) A model
of European prehistory[11] onto which our reference samples have been projected using NNLS (see
Methods), and the same data represented spatially. Chronologically, Western Hunter-Gatherers
(WHG) and Eastern Hunter-Gatherers (EHG) were largely replaced by Anatolian Farmers amid
demographic changes during the "Neolithic transition" around 9,000 years ago. Later migrations
during the Bronze Age about 5,000 years ago brought a roughly equal Steppe ancestry component
from the Pontic-Caspian Steppe to Europe, an ancestry descended from the EHG from the Middle
Don River region and Caucasus Hunter-Gatherers (CHG)[9]. Steppe ancestry has been associated with
the Yamnaya culture and then with the expansion westwards of the Corded Ware Complex and Bell
Beaker culture, and the eastwards expansion in the form of the Afanasievo culture[12,10]. Samples are
vertical bars representing their "admixture estimate" estimated by NNLS (methods) from six

ancestries: EHG (green), WHG (pink), CHG (yellow), Farmer (blue), Steppe (cyan) or an Outgroup
(represented by ancient Africans, red). Important population expansions are shown as growing bars
and "recent" (post-Bronze age) non-reference admixed populations are shown for the Denmark time-
transect (see Supplementary Figure 1.1 for details).
123
<u>RESULTS</u>
We obtained local ancestry (i.e. ancestry at specific loci) labels for ~410,000 self-identified "white
British" individuals in the UK Biobank[13], using a reference panel of 318 ancient DNA (aDNA)
samples[9] (Figure 1; Supplementary Figure 1.1) from the Mesolithic and Neolithic, including Steppe
pastoralists. Comparing the ancestry at each labelled single nucleotide polymorphism (SNP,
n=549,323) to genome-wide ancestry in the UK Biobank provided a "local ancestry anomaly score"
(Methods), for which two regions stood out as having undergone the most significant ancestry-
specific evolution in this period: LCT/MCM6, regulating lactase persistence[14], and the HLA region
(Figure 2, top).
133
To determine whether this evolution of the HLA region has subsequently impacted diseases that are
strongly associated with risk alleles found within this region, we investigated the history of variants
associated with two HLA-associated autoimmune diseases, multiple sclerosis (MS) and rheumatoid
arthritis (RA), using the largest ancient genome dataset from the Stone Age coupled with 86 new
Medieval and post-Medieval genomes from Denmark (Supplementary Figure 1.1, Supplementary
Note 1, ST1). Alongside modern data, with our newly published genomes we have an almost
complete transect from approximately 10,000 years ago to the present.
141
The allele frequencies of SNPs conferring the highest risk for MS (all in the HLA class II region) in
our ancient groups show striking patterns. In particular the tag SNP (rs3135388-T) for HLA-
DRB1*15:01, the largest risk factor for MS, first appeared in an Italian Neolithic individual (sampleId
R3 from Grotta Continenza, C14 dated to between 5,836-5,723 BCE, coverage 4.05X) and rapidly
increased in frequency around the time of the emergence of the Yamnaya culture around 5,300 years
ago in Steppe and Steppe-derived populations (Figure 2). From risk allele frequencies of individuals
in the UK Biobank born in, and of a 'typical ancestral background' for, each country[9], we found
HLA-DRB1*15:01 frequency peaks in modern populations of Finland, Sweden and Iceland, and in
ancient populations with high Steppe ancestry (Figure 2, inset).
151

Figure 2. Areas of unusual local ancestry in the genome, and ancient and modern frequencies of DRB1*15:01.

*a): Local Ancestry Anomaly Score measuring the difference between the local ancestry and the genome-wide average (capped at -log10(p)=20; see Methods). b) HLA-DRB1*15:01 frequencies in ancient and modern (inset) populations; this is the highest effect variant for MS risk (calculated using rs3135388 tag SNP). For the ancient data, for each ancestry (CHG, EHG, WHG, Farmer, Steppe) the five populations with the highest amount of that ancestry are coloured and labelled. DRB1*15:01 was present before the Steppe expansion, but rose to high frequency during the Yamnaya formation (shaded red). The geographical distribution of DRB1*15:01 frequency in modern populations in the UK Biobank is also shown (inset).*

To investigate the risk of a particular ancestry at all MS-associated fine-mapped loci[3] present in the UK Biobank imputed dataset (n=205/233, see methods), we used the local ancestry dataset to calculate a risk ratio (see Methods: Weighted Average Prevalence) for each ancestry. For MS, Steppe ancestry has the highest risk ratio in nearly all HLA SNPs, while Farmer and 'Outgroup' ancestry (represented by ancient Africans) are often the most protective (Figure 3, top), meaning a Steppe-derived haplotype at these positions confers MS risk.

(a) Risk ratio for all MS−associated SNPs based on WAP

Figure 3: Associations between local ancestry and MS in a modern population.

a) Risk ratio of SNPs for MS based on weighted average prevalence (WAP; see Methods), when

decomposed by inferred ancestry. Each ancestry is assigned a mean and confidence interval based on

bootstrap resampling, for each chromosome (faded where non-significant). The three HLA regions

are split from the rest of chromosome 6, and SNPs with risk ratio >1.2 or <0.8 are annotated. b-c)

Genome-wide Ancestral Risk Scores (ARS, see Methods) for MS. Confidence intervals are estimated

by either bootstrapping over individuals (b, which can be interpreted as testing power to reject a null

of no association between MS and ancestry) and bootstrapping over SNPs (c, which can be

interpreted as testing whether ancestry is associated with MS genome-wide).

Having shown that some ancestries carry higher risk, we calculated an aggregate risk for each

ancestry across the same SNPs using a new statistic, the Ancestral Risk Score (ARS). ARS is

computed in a large modern sample with local ancestry labels, estimating the relative risk for a

modern individual consisting of entirely one ancestry, mitigating the effects of low aDNA sample
numbers and bias[15], and being robust to intervening drift and selection. We used effect size estimates
from previous association studies, under an additive model, with confidence intervals obtained via an
accelerated bootstrap[16] (Supplementary Note 4). In the ARS for MS (Figure 3 bottom), Steppe
ancestry had a large and significant risk, followed by WHG, CHG and EHG; Neolithic Farmer and
Outgroup ancestry had the lowest ARS (Figure 3). Therefore Steppe ancestry is contributing the most
risk for MS overall. We tested for a genome-wide association by resampling loci, and found that
Steppe risk is much reduced but still clearly exceeds Farmer, a pattern which holds even when
excluding SNPs on the HLA (Supplementary Note 4, Figure S4.1).
195
The fact that all but two MS-associated HLA SNPs confer risk within Steppe ancestry implies that
this risk has a common evolutionary history. We therefore investigated whether ancestry was
important for prediction using three types of association study in the UK Biobank for disease-
associated SNPs, controlling for age, sex and the first 18 PCs. The first of these is a regular SNP-
based association as conducted in GWAS. The second uses local ancestry probabilities instead of
genotype values (Supplementary Note 3). The third is based on Haplotype trend regression (HTR)
which is used to detect interactions between SNPs[17] by treating each haplotype's probability as a
feature from which to predict a trait, instead of using SNPs as in a regular GWAS. We developed a
new method called Haplotype Trend Regression with eXtra flexibility (HTRX, Supplementary Note
5) that searches for haplotype patterns that include single SNPs and non-contiguous haplotypes. To
prevent overfitting, we reported out-of-sample variation explained, and showed by simulation (see
Supplementary Figure 4.4) that HTRX predicts the same variance as regular GWAS when interactions
are absent, but explains more variance when the interaction strength increases.
209
Although our cohort of self-identified "white British" individuals is relatively under-powered with
respect to MS (cases=1,949; controls=398,049; prevalence=0.487%), MS was associated with Steppe
and Farmer ancestry ($p<1e-10$) in the HLA region (Supplementary Figure 4.1). In 3 out of 4 main LD
blocks within the HLA (class I, two subregions of class II determined by LD blocks at 32.41-32.68Mb
and 33.04-33.08Mb, and class III), local ancestry explains significantly more variation in total than
SNP variation (Figure 4; measured by average out-of-sample McFadden's R^2 for logistic regression,
see Methods). While increased ancestry performance over GWAS can be explained by tagging of
SNPs outside the region, increased HTRX performance over GWAS quantifies the total effect of a
haplotype, including rare SNPs and epistasis. Across the entire HLA region, haplotypes explain at
least 17% more out-of-sample variation than GWAS (2.90%, compared to 2.48%). Interaction signals
are also observed within class I, within class II, and between class I and III.
221

Figure 4: MS association in the HLA.

Comparison of variance explained in MS within the UK Biobank, for all fine-mapped HLA SNPs with an independent contribution[3]. The plots compare GWAS (treating SNPs as having independent effect), local ancestry at those SNPs, and HTRX (haplotypes) after accounting for covariates (Methods). a) is for fine-mapped MS-associated SNPs in the HLA. b) is HLA class I and -III, c) is HLA class II, d) is HLA class I, e) is HLA class III, f) and g) are subregions of HLA class II chosen from LD. HTRX has small "up-arrows" where these are lower bounds (Methods). h) Genetic correlations in the HLA region at our time-depth from Ancestry-based LD (LDA, see Methods) and Supplementary Figure 6.5 for LD.

This interaction risk can be attributed to particular ancestries: for example, multiple haplotypes at the 32.41-32.68Mb region are Steppe-associated and have high MS odds ratios. We further tested whether co-occurring ancestries at each loci were associated with MS (Methods; Supplementary Figure 4.2), but found no evidence that risk was associated with anything other than Steppe ancestry.

Having established that Steppe ancestry contributes most of the HLA-associated risk for MS, we
investigated evidence for polygenic selection on the disease-associated variants using two methods.
Firstly, we used a novel chromosome painting technique based on inference of a sample's nearest
neighbours in the marginal trees of an ARG that contains labelled individuals (Irving-Pease et al.,
submitted). The resulting ancestral path labels, for haplotypes in both ancient and modern individuals,
allowed us to infer allele frequency trajectories for risk associated variants, while controlling for
changes in admixture proportions through time. These paths extend backwards from the present day to
approximately 15,000 years ago, and are labelled with the unique population that a path travels
through. We stress that the path labels are not representative of a continuous population, but represent
a path backwards in time that encompasses that ancestry. For example, the CHG path originates in
Caucasus hunter-gatherers, before merging with EHG to form the Steppe population, and then merges
with other ancestries in later European populations (Figure 1).
250
251   Because not all fine-mapped SNPs had ancestral path labels (missing OR=10.4%) and due to the
252   difficulty in accurately inferring HLA alleles in ancient samples[18], we LD-pruned genome-wide
253   significant summary statistics from the same study[3] for which we did have ancestry path labels (n=62,
254   see methods). This allowed us to test for polygenic selection across disease-associated variants using
255   CLUES[19] and PALM[20].
256
For MS, we found evidence that disease risk was selectively increased when considering all ancestries
collectively (p=5.06e-05; ω=0.0029), between 5,000-2,000 years ago (Figure 5). Conditioning on each
of the four long-term ancestral paths (CHG, EHG, WHG and ANA), we found a statistically
significant signal of selection in CHG (p=6.45e-3; ω=0.009). None of the other ancestral paths
reached nominal significance, although ANA (p=0.0743; ω=0.011) and EHG (p=0.064; ω=0.0045)
paths were close. Again, it is likely that the selection occurred in the pastoralist population of the
Steppe, as that population consists of approximately half CHG ancestry[11] (Figure 1). The SNP driving
the largest change in genetic risk over time was rs3129934, in both the pan-ancestry (p=9.52e-06;
s=0.017) and CHG (p=0.019; s=0.008) analyses, which tags the HLA-DRB1*15:01 haplotype[21]. We
also tested three other alleles that tag the HLA-DRB1*15:01 haplotype (rs3129889, rs3135388 and
rs3135391) for evidence of selection, and found that the ancestry stratified signal was consistently
strongest in CHG (Figure 5). None of the four tag SNPs were detected on either the EHG or WHG
backgrounds, indicating that the HLA-DRB1*15:01 haplotype likely originated in the basal
population ancestral to both ANA and CHG.
271

272
Figure 5: Evidence for selection on MS-associated SNPs.
a) Stacked line plot of the pan-ancestry PALM analysis for MS, showing the contribution of alleles to
disease risk over time. Individual SNPs are stacked, with their trajectories polarised to show the
frequency of the positive risk allele and weighted by their scaled effect size: when a given SNP bar
becomes wider over time the risk allele has increased in frequency, and vice versa. SNPs are sorted
by their marginal p-value and direction of effect, with selected SNPs that increase risk plotted on top.
SNPs are also coloured by their marginal p-values, and significant SNPs are shown in yellow. The y-
axis shows the scaled polygenic risk score (PRS), which ranges from 0 to 1, representing the
maximum possible additive genetic risk in a population.
b) Maximum likelihood trajectories for four SNPs tagging DRB1*15:01. The background is shaded
for the approximate time period in which the ancestry existed as an actual population. None of the
tagging alleles are present on the EHG or WHG ancestral paths.

Our second selection measure introduces a new statistic, Linkage Disequilibrium of Ancestry (LDA).
LDA is the correlation between ancestries across SNPs, measuring whether recombination events
between ancestries are high compared to recombination within ancestries. From this we constructed

289   an "LDA score" using the fine-mapped SNPs, which is the total amount of genome in LDA with a
290   given SNP. A high LDA score indicates that the haplotype inherited from the reference population is
291   longer than expected, while a low score indicates that the haplotype is shorter than expected (i.e.
292   underwent more recombination). For example, the LCT/MCM6 region exhibits a high LDA score
293   (Supplementary Figure 6.4), as expected from a relatively recent selective sweep[22].
294

The HLA has significantly *lower* LDA scores than the rest of chromosome 6 (Supplementary Figure
6.4). We simulated the LDA score under selection (Supplementary Figure 6.1; Methods), which
showed that when SNP frequencies are increasing in the most recent population, single locus selection
cannot explain this signal (Supplementary Figure 6.2-3). Instead, different loci in LD must have
independently reached high frequency in different ancestral populations that admixed, with selection
favouring haplotypes of mixed ancestry over single-ancestry haplotypes. Although multi-SNP
selection has been modelled[23], the interaction with prior population structure is less explored and is
important for the HLA, justifying a new term, "recombinant favouring selection".
303

The HLA region contains the highest "Outgroup" ancestry anywhere on the genome (Figure 6),
reflecting high nucleotide diversity. Unlike other measures of balancing selection such as Fst (Figure
6), LDA describes excess ancestry LD from specific, dated populations and therefore need not be
correlated with them. For the HLA class II region, the selection measures all line up (LDA score, Fst,
pi), but for class I, the LDA score has an additional non-diverse minimum at 30.8Mb, implying that
here the genome is ancestrally diverse but genetically strongly constrained. The LDA score is thus
informative about the type of selection being detected, and whether it has been subject to change.
311

Figure 6: Signatures of selection at the HLA locus showing different regions of the HLA (coloured bar) and locations of MS-associated SNPs (vertical lines, coloured by the variance explained by 6 ancestries). *a): Whole Chromosome 6 "local ancestry" decomposition by genetic position. b). HLA "local ancestry" decomposition. c): LDA score; low values are indicative of selection for multiple linked loci, while high values indicate positive selection. d): pi scores (nucleotide diversity) for CEU (Northern and Western European ancestry). MS-associated SNPs fall in highly diverse regions of the HLA. e): Fst scores (divergence between two populations) for CEU vs YRI(Yoruba); locally higher scores indicate regions that have undergone differential selection between the two populations.*

Because MS would not have conferred a fitness advantage on ancient individuals, it is likely that this selection was driven by traits with shared genetic architecture, of which increased risk for MS in the present is a consequence. We therefore looked at LD-pruned MS-associated SNPs that showed statistically significant evidence for selection using CLUES (n=26) and which also had a genome-

wide significant trait association (p < 5e-8) in any of the 4,359 traits from the UK Biobank[13] (UK
Biobank Neale Lab, Round 2: http://www.nealelab.is/uk-biobank/). We found that many selected
SNPs are also associated with celiac disease (n=15), white blood cell/neutrophil count (n=15/n=15),
hypothyroidism (n=14) and haemoglobin concentration (n=14) (Supplementary Figure 7.1). This
raised the possibility that the selection had increased risk for both MS and celiac disease, and when
we tested celiac disease for polygenic selection, we found significant evidence for positive selection,
increasing genetic risk (p=9.65e-3; ω=0.846, Supplementary Note 6).

Because the UK Biobank is underpowered with respect to many traits and diseases, we also undertook
a manual literature search (see methods) for all SNPs that reached genome-wide significance for
association with MS in the summary stats (i.e., not LD-pruned, as independence is not required) and
which showed statistically significant evidence for selection using CLUES (n=94). We found that
most of the alleles under positive selection are associated with protective effects against specific
pathogens (virus, bacteria, fungi and parasites) and/or infectious diseases within one or several
ancestral paths (disease or pathogen associated/total selected in ancestry path: pan-ancestry 36/44;
ANA 24/31; CHG 25/29; EHG 27/35; WHG 9/10, Supplementary Note 8, ST13, Supplementary
Figure 8.1), although we note that GWAS data for many infectious diseases are not available. We
observed that the selected alleles had protective associations with several chronic viruses (EBV, VZV,
HSV, and CMV) and to viruses or diseases not associated with transmission in small hunter-gatherer
groups (e.g., measles, mumps, influenza, whooping cough). Moreover, many selected alleles
conferred a reduction of risk of parasites, of skin and subcutaneous tissue, gastrointestinal, respiratory,
urinary tract, and sexually transmitted infections, or of pathogens associated with these or other
infections (e.g., malaria, toxoplasmosis, entamoeba histolytica, clostridium difficile, tuberculosis,
streptococcus pyrogenes, and chlamydia) (Supplementary Note 8, ST13, Supplementary Figure 8.1).

We contrasted these findings for MS with results for RA, a common inflammatory HLA class II-
associated disease that primarily affects the joints causing pain, swelling and stiffness[24], which shows
a strikingly different ancestry risk profile. HLA-DRB1*04:01 is the largest genetic risk factor for RA;
in the CLUES analysis, the tag SNP for this allele (rs660895) displayed evidence of continuous
negative selection until approximately 3,000 years ago (p=4.63e-4, Supplementary Figure 5.1). We
found that WHG and EHG ancestries often confer the most risk at SNPs associated with RA (Relative
Risk ratio of RA-associated SNPs based on WAP, see Methods); and these ancestries have
contributed the greatest risk for RA on aggregate, reflected in a higher ARS for these ancestries
(Supplementary Note 4), while Steppe and Outgroup ancestry have the lowest scores (Supplementary
Figure 3.1). These results were recapitulated in the local ancestry GWAS (Supplementary Note 3).

We found that RA-associated SNPs have undergone negative polygenic selection (p=7.93e-3,
Supplementary Figure 5.1) over the last approximately 15,000 years; when this is decomposed by
ancestry path, we found significant evidence for negative selection in both the CHG (p=3.09e-5) and
ANA (p=1.20e-3) ancestry paths. We found no evidence for negative selection in the EHG and WHG
paths, although both show a trend of increasing risk, and EHG nears significance (p=0.0842).

These results demonstrate that genetic risk for RA was higher in the distant past, in contrast to MS,
with RA-associated risk variants present at higher frequencies in European hunter-gatherer
populations before the arrival of agriculture. In order to understand what caused the high risk in
hunter-gatherer populations and subsequent negative selection, we again undertook a manual
literature search for pleiotropic effects of SNPs associated with RA. Because the number of SNPs that
reached genome-wide significance in the GWAS study and also showed statistically significant

evidence for directional selection was large, we only analysed LD-pruned SNPs (n=42). We found
that the majority of selected SNPs were associated with protection against distinct pathogens and/or
infectious diseases across all paths (disease or pathogen associated/total selected in ancestry path:
pan-ancestry 9/13; ANA 10/13; CHG 8/11; EHG 10/16; WHG 10/12). We found that selected RA-
risk alleles were often linked to the same pathogens or diseases as in the MS analysis, although the
number of protective associations to distinct pathogens were fewer (Supplementary Note 8, ST14,
Supplementary Figure 8.1).

DISCUSSION
The last 10,000 years have seen some of the most extreme global experiments in lifestyle with the
emergence of farming in some regions and a pastoral lifestyle in others. While 5,000 years ago farmer
ancestry predominated across Europe, a relatively diverged ancestry arrived with the Steppe
migrations around this time. We have shown that this ancestry contributes the most genetic risk for
MS today, and that these variants were the result of positive selection coinciding with the emergence
of a pastoralist lifestyle on the Pontic-Caspian Steppe, and continued selection in the subsequent
admixed post-Stone Age populations in Europe. This ultimately created a legacy of heterogeneity in
MS risk observed across Europe today. These results address the long-standing debate around the
north-south gradient in MS prevalence in Europe, and suggest that the Steppe ancestry gradient in
modern populations - specifically at the HLA region - across the continent causes this phenomenon in
combination with environmental factors. Furthermore, while epistasis between MS-associated variants
in the HLA region has been demonstrated before[25, 26, 27, 28], we have shown that accounting for this
explains 17% more variance than independent SNPs effects alone. Many of the haplotypes carrying
these risk alleles have ancestry-specific origins, which could be exploited for individual risk
prediction and may offer a pathway from ancestry associations into a mechanistic understanding of
MS risk. We have contrasted these findings with results for rheumatoid arthritis (RA), another HLA
class II associated chronic inflammatory disease, and found that the genetic risk for RA exhibits a
contrasting pattern: genetic risk was highest in Stone Age hunter-gatherer ancestry and decreased over
time.

Our interpretation of this history is that co-evolution between pathogens and their human hosts has
resulted in massive and divergent ancestry-specific selection on immune response genes according to
lifestyle and environment, driven by a range of pathogenic drivers, and "recombinant favouring
selection" after these populations merged. The Late Neolithic and Early Bronze Age was a time of
massively increased infectious diseases in human populations, due to increased population density as
well as contact with, and consumption of, domesticated animals. Many diseases trace their origins to
this period, such as tuberculosis (TB) caused by the intracellular bacteria *Mycobacterium tuberculosis*
or *Mycobacterium bovis*[29, 30], bubonic plague caused by *Yersinia pestis*[31, 32, 33], herpes simplex virus[34],
and chickenpox caused by varicella-zoster virus[35], and we have shown that many of the MS- and RA-
associated variants under selection confer resistance to a range of infectious diseases and pathogens
(Supplementary Note 8). For example, HLA-DRB1*15:01 is associated with protection against TB[36]
and increased risk for lepromatous leprosy[37]. However, we are underpowered to detect specific
associations beyond this hypothesis due to poor knowledge of the distribution and diversity of past
diseases, poor preservation of endogenous pathogens in the archaeological record, and a lack of well-
powered GWAS studies for many infectious diseases.

A pattern that repeatedly appears is that of lifestyle change driving changes in risk and phenotypic
outcomes. We have shown that in the past environmental changes driven by lifestyle innovation
inadvertently drove an increase in genetic risk for MS. Today, with increasing prevalence of MS cases

observed over the last five decades[38, 39]), we again observe a striking correlation with changes in our
environment, including lifestyle choices and improved hygiene, which no longer favours this previous
genetic architecture. Instead, the fine balance of genetically-driven cells within the immune system,
which are needed to combat a broad repertoire of pathogens without harming self-tissue, has been met
with new challenges, including a potential absence of requirement. For example, while a population of
immune cells, T helper 1 (Th1), direct strong cellular immune responses against intracellular
pathogens, T helper 2 (Th2) cells mediate humoral immune responses against extracellular bacteria
and parasites and further have the capacity to guide the restoring of homeostasis,thus preventing
damage of the infected tissue via immune-regulatory cytokines. We have shown that the majority of
selected MS-associated SNPs are associated with protection against a wide range of pathogens,
consistent with strong but balanced Th1/Th2 immunity in the Bronze age, where a diversification of
pathogens likely took place. In contrast, although MS pathogenesis is complex and multicellular of
nature, CD4+ Th cells, in particular IFN-ɣ producing Th1 cells and IL-17-producing Th17 cells play a
key role in disease development[2]. The skewed Th1/Th2 balance observed in MS may partly result
from the developed world's increased sanitation, which has led to drastically reduced burden of
parasites, which the immune system had evolved to efficiently combat[40]. In the case of RA, the
exposure of Hunter Gatherer populations to the respiratory or gastrointestinal pathogens linked to
triggering RA[41] was likely low. The new pathogenic challenges associated with agriculture, animal
domestication, pastoralism, and higher population densities might have substantially increased the risk
of developing RA in genetically predisposed individuals, resulting in negative selection. If true, this
would present a parallel between RA in the Bronze Age and MS today, in which lifestyle changes
have exposed previously favourable genetic variants as autoimmune disease risks.
445
More broadly, it is clear that this was a critical period in human history during which highly
genetically and culturally divergent populations evolved and eventually mixed. These separate
histories dictate the genetic risk and prevalence of several autoimmune diseases today. Surprisingly,
the emergence of the pastoralist Steppe lifestyle may have had an impact on immune response as great
as or greater than the emergence of farming during the Neolithic transition, commonly held to be the
greatest lifestyle change in human history.
452
453   DATA AVAILABILITY
454   All collapsed and paired-end sequence data for novel samples sequenced in this study will be made
455   publicly available on the European Nucleotide Archive, together with trimmed sequence alignment
456   map files, aligned using human build GRCh37. Previously published ancient genomic data used in
457   this study are detailed in ST15, and are all already publicly available.
458
CODE AVAILABILITY
The modified version of CLUES used in this study is available from https://github.com/standard-
aaron/clues. The pipeline and conda environment necessary to replicate the analysis of allele
frequency trajectories and polygenic selection in Supplementary Note 6 are available on Github at
https://github.com/ekirving/ms_paper. The code to create Ancestry Anomaly scores based on
Chromosome painting is on Github at https://github.com/danjlawson/ms_paper. The codes to
compute LDA and LDA score are available on Github at
https://github.com/YaolingYang/LDAandLDAscore. The codes to implement HTRX and its
simulation are on Github at https://github.com/YaolingYang/HTRX. The codes to implement ARS
calculation are on Github at https://github.com/will-camb/ms_paper.
469

ACKNOWLEDGEMENTS
We extend our thanks to all the former and current staff at the Lundbeck Foundation GeoGenetics
Centre and the GeoGenetics Sequencing Core, and to colleagues across the many institutions detailed
below. We are particularly grateful to Maria Madrona, Lærke Hansen and Julie Bitz-Thorsen for
laboratory assistance; and to Julie Hansen, Sandra Mularczyk, Katja Thorø Michler, Emilie Neerup
Nielsen for their help with sampling, and to Line Olsen as project manager for the Lundbeck
Foundation GeoGenetics Centre project. We thank UK Biobank Ltd. for access to the UK Biobank
genomic resource. We are thankful to Illumina Inc. for collaboration. E.W. thanks St. John's College,
Cambridge, for providing a stimulating environment of discussion and learning.
479
AUTHOR CONTRIBUTIONS
W.B., Y.Y., K.E.A., E.K.I-P, G.S., and L.T.J. contributed equally to this work.
A.I., D.J.L., L.F., and E.W. led the study.
W.B., A.R-M., L.F., R.N., and E.W. conceptualised the study.
R.N., K.K., L.F., and E.W. acquired funding for research.
A.R., C.G., F.D., M.L.S.J., S.B.M., B.S., L.K., I.M.H., N.W., L.V., and T.S.K., were involved in
sample collection and processing
W.B., Y.Y., E.K.I-P, A.S., S.R., and D.J.L. were involved in developing and applying methodology.
W.B., Y.Y., E.K.I-P, G.S., A.A., A.R., E.A.D., M.S., S.R., A.I., and D.J.L. undertook formal analyses
of data.
W.B., Y.Y., K.E.A., E.K.I-P, and L.T.J., A.I., L.F., and E.W. drafted the main text (W.B. led this).
W.B., Y.Y., E.K.I-P, G.S., L.T.J., E.A.D., A.S., F.D., M.L.S.J., S.B.M., B.S., L.K., I.M.H., N.W.,
L.V., A.I., and D.J.L. drafted supplementary notes and materials.
W.B., Y.Y., K.E.A., E.K.I-P, L.T.J., A.A., K.K., R.N., A.I., D.J.L., L.F., and E.W. were involved in
reviewing drafts and editing.
All co-authors read, commented on, and agreed upon the submitted manuscript.
496
497
498 COMPETING INTERESTS
499 The authors declare no competing interests
500

501 Supplementary Figures

502
503

Supplementary Figure 1.1. Ancient sample PCA, map, ancestry proportions through time for
samples in Denmark. (1) PC1 vs PC2 of the filtered Western Eurasian ancient samples included in
this study. Black circled points are Danish Medieval and post-Medieval samples published here for
the first time. Major component ancestry locations are labelled. (2) Map of ancient filtered Western
Eurasian ancient samples included in this study (3a) Map of reference data and time transect of
Denmark as in Figure 1. (3b) More recent ancient data (samples <4,200 years ago) not used as
reference, showing the clines of the main ancestry components from (3a).

513
514 **Supplementary Figure 1.2. Modern prevalences of RA (left) and CD (right).**
515 Modern-day geographical distribution of RA and CD prevalence in Europe. Prevalence data for RA
516 (cases per 100,000) was obtained from [42]. For CD, the seroprevalence (%) is based on the presence of
517 transglutaminase and/or endomysial autoantibodies; data were obtained from [43].
518

519
Supplementary Figure 1.3 Association between genome-wide Steppe ancestry, MS prevalence
and DRB1*15:01 frequency in modern populations in the UK Biobank.

Supplementary Figure 2.1. Ancient and modern prevalences of HLA-DRB1*04:01 (rs3817964)
and HLA-DQ2.5 (rs2187668).
Top and middle: Ancient distributions of HLA-DRB1*04:01, the largest genetic risk factor in RA,
and HLA-DQ2.5, the largest genetic risk factor in CD. Average frequency across all populations is
shown (blue line, 10 time bins) as well as the Bronze Age (red shading).
Bottom: Modern distribution of HLA-DRB1*04:01 (left) and HLA-DQ2.5 (right) in populations in
the UK Biobank. NB the tag SNPs may be less effective at tagging these types in non-European
populations, so we urge caution in interpretation - especially in African populations.
533
534

535
Supplementary Figure 3.1: Associations between local ancestry and RA in a modern population.
a) Risk ratio of SNPs for RA based on weighted average prevalence (WAP; see Methods), when
decomposed by inferred ancestry. Each ancestry is assigned a mean and confidence interval based on
bootstrap resampling, for each chromosome (faded where non-significant). SNPs with risk ratio >1.15
or <0.85 are annotated. b-c) Genome-wide Ancestral Risk Scores (ARS, see Methods) for RA.
Confidence intervals are estimated by either bootstrapping over individuals (b, which can be
interpreted as testing power to reject a null of no association between RA and ancestry) and
bootstrapping over SNPs (c, which can be interpreted as testing whether ancestry is associated with
RA genome-wide).
545

Supplementary Figure 3.2: Associations between local ancestry and CD in a modern population.
a) Risk ratio of SNPs for CD based on weighted average prevalence (WAP; see Methods), when decomposed by inferred ancestry. Each ancestry is assigned a mean and confidence interval based on bootstrap resampling, for each chromosome (faded where non-significant). SNPs with risk ratio >1.15 or <0.85 are annotated. b-c) Genome-wide Ancestral Risk Scores (ARS, see Methods) for CD. Confidence intervals are estimated by either bootstrapping over individuals (b, which can be interpreted as testing power to reject a null of no association between CD and ancestry) and bootstrapping over SNPs (c, which can be interpreted as testing whether ancestry is associated with CD genome-wide).

559
560 **Supplementary Figure 4.1: Association with MS risk at externally ascertained SNPs, for (top)**
561 **ancestry, and (bottom) SNPs**.
562 Due to the UK Biobank being less powered (having fewer cases) than the Case-Control study from
563 which these SNPs were found, the only statistically significant association is in the HLA.
564

Supplementary Figure 4.2: Comparison between MS-risk and local ancestry for 3 example

SNPs.

In the HLA class II region, all SNPs share a pattern in which high Steppe ancestry is associated with

high MS-risk. The risk decreases monotonically and is not present in the Steppe precursor populations

(Hunter Gathers), but is with the admixed Bronze-age European populations (Steppe + Farmer).

572
573
574 **Supplementary Figure 4.3: Decomposition of individuals ancestry at MS risk SNPs in terms of**
575 **(left) the ancestry of those SNPs alone, or (right) the Weighted average prevalence of MS in**
576 **each ancestry after "logit" transformation.**
577

Supplementary Figure 4.4: Simulation study with four SNPs showing the boxplots of out-of-

sample variance (with the red line representing the average) explained by HTRX compared to

GWAS, HTR and the true model.

The total variance explained by HTRX is the same as SNP and bigger than HTR when there are no

interactions. When interaction (with subtitle "int") exists, HTRX significantly outperforms GWAS

and HTR. In all situations, HTRX works similarly to the truth.

a) Rheumatoid arthritis (r^2 < 0.05; window 250 kb) (n = 153) | All ancestries | ω = -0.005 | se = 0.0021 | z = -2.655 | p = 0.00793

b)   rs660895 | chr6:32577380 | Gene(s): HLA-DRB1 - HLA-DQA1 | A/G

logLR = 12.26 | p = 4.63e-04 | epoch = 0-529 | s = -0.00710

587
Supplementary Figure 5.1: Evidence for selection on RA-associated SNPs.
a) Stacked line plot of the pan-ancestry PALM analysis for RA, showing the contribution of alleles to
disease risk over time. Individual SNPs are stacked, with their trajectories polarised to show the

frequency of the positive risk allele and weighted by their scaled effect size: when a given SNP bar
becomes wider over time the risk allele has increased in frequency, and vice versa. SNPs are sorted by
their marginal p-value and direction of effect, with selected SNPs that increase risk plotted on top.
SNPs are also coloured by their marginal p-values, and significant SNPs are shown in yellow. The y-
axis shows the scaled polygenic risk score (PRS), which ranges from 0 to 1, representing the
maximum possible additive genetic risk in a population.
b) Posterior likelihood trajectory for rs660895, tagging HLA-DRB1*04:01, inferred by CLUES.
598

599
600   **Supplementary Figure 6.1: Simulating Low LDA score**.
601   Left: A simulated history in which a single population splits into two ("Steppe" and "Farmer") after
602   2200 generations and experiences positive selection on different loci (m_1 in P_1 and m_2 in P_2). After
603   2900 generations the populations merge ("Europeans") but selection continues on *both* loci.
604

Supplementary Figure 6.2: LDAS simulation with positive or balancing selection in the modern population. The left two columns show simulations with a single variant satisfying the observed constraint that modern-day frequencies are not decreasing (i.e. not negative selection). The right column shows simulations with two variants, also obeying this constraint. The model for simulating 2 loci is the same as in Supplementary Figure 6.1, and that for 1 locus is in the top right of this plot (which differs only in the location of the selected variant in the separated populations).

613
614 **Supplementary Figure 6.3: LDAS simulation with single locus negatively selected in the modern**
615 **population.** In two cases this generates a low LDAS score, which requires recent negative selection
616 (which is ruled out for HLA by the observed frequency trend). In this case, one ancestry dominates
617 the region and recombination to the other conveys risk. The model used is in the top right of
618 Supplementary Figure 6.2.
619

620
621
622 **Supplementary Figure 6.4: LDAS on chromosome 6 and 2.** LDA score is a) high in the
623 LCT/MCM6 region while is b) low in the HLA region.

624
625 **Supplementary Figure 6.5: Pairwise Linkage Disequilibrium (LD) plot (measured by D') for all**
626 **the MS-associated SNPs on chromosome 6.**

627
628
629

 Supplementary Figure 7.1 Allele frequency plots for positively selected MS-associated SNPs that
are also associated with other phenotypes in the UK Biobank. Traits 1-5.
SNPs are shown with their maximum likelihood trajectories, and polarised by the direction of their
effect on the marginal UK Biobank trait (i.e. showing the 'risk' allele). Phenotypes are ordered
according to the number of common SNPs, non-significant SNPs are shown with partial transparency,
portions of the trajectory with low posterior density are cropped off, and the background is shaded for
the approximate time period in which the ancestry existed as an actual population.

Supplementary Figure 7.2 Allele frequency plots for positively selected MS-associated SNPs that
are also associated with other phenotypes in the UK Biobank. Traits 6-10.
SNPs are shown with their maximum likelihood trajectories, and polarised by the direction of their
effect on the marginal UK Biobank trait (i.e. showing the 'risk' allele). Phenotypes are ordered
according to the number of common SNPs, non-significant SNPs are shown with partial transparency,
portions of the trajectory with low posterior density are cropped off, and the background is shaded for
the approximate time period in which the ancestry existed as an actual population. Note that many
phenotypes are underpowered in the UKBiobank GWAS, hence why MS appears as just the joint 7th
in this list.

Supplementary Figure 7.3 Allele frequency plots for positively selected MS-associated SNPs that are also associated with other phenotypes in the UK Biobank. Traits 11-15.

SNPs are shown with their maximum likelihood trajectories, and polarised by the direction of their effect on the marginal UK Biobank trait (i.e. showing the 'risk' allele). Phenotypes are ordered according to the number of common SNPs, non-significant SNPs are shown with partial transparency, portions of the trajectory with low posterior density are cropped off, and the background is shaded for the approximate time period in which the ancestry existed as an actual population.

The number of protective links to pathogens or infectious diseases within the MS- and RA-risk SNPs

■ RA (n=42) ■ MS (n=94)

Supplementary Figure 8.1 The number of protective associations with pathogens or infectious
diseases for the MA- and RA-associated selected SNPs
The number of protective associations to specific pathogens and/or diseases associated with the MS-
and RA-SNPs that showed statistically significant evidence for selection using CLUES. One SNP can
have a link to more than one pathogen and/or disease (see ST13 and ST14 for details on each SNP).
Fifteen and twelve SNPs had no detectable links to any pathogen or infectious disease in the MS and
RA SNP sets, respectively.
666
667

<u>METHODS</u>
Data Generation
Overview
In order to examine variants associated with phenotypes backwards in time, we assembled a large
ancient DNA dataset. Here we present new genomic data from 86 ancient individuals from Medieval
and post-Medieval periods from Denmark (Supplementary Figure 1, Supplementary Note 1, ST1).
The samples range in age from around the XIth to the XVIIIth century. We extracted ancient DNA
from tooth cementum or petrous bone and shotgun sequenced the 86 genomes to a depth of genomic
coverage ranging from 0.02 X to 1.6 X (mean = 0.39 X and median = 0.27 X). The genomes of the
new 86 individuals were imputed using the 1,000 Genomes phased data as a reference panel by an
imputation method designed for low coverage genomes (GLIMPSE[44]), and we also imputed 1,664
ancient genomes presented in the accompanying study 'Population Genomics of Stone Age Eurasia'[9].
Depending on the specific data quality requirements for the downstream analyses, we filtered out
samples with poor coverage, variant sites with low MAF and with low imputation quality (average

genotype probability < 0.98). Our dataset of ancient individuals span approximately 15,000 years
across Eurasia (Supplementary Figure 1).
684
Ancient data DNA extraction and library preparation
Laboratory work was conducted in the dedicated ancient DNA clean-room facilities at the Lundbeck
Foundation GeoGenetics Centre (Globe Institute, University of Copenhagen). A total of 86 Medieval
and post-Medieval human samples from Denmark (ST2) were processed using semi-automated
procedures. Each sample was processed in parallel. For each extract non USER-treated and USER-
treated (NEB) libraries were built[45]. All libraries were sequenced on the NovaSeq6000 instrument at
the GeoGenetics Sequencing Core, Copenhagen, using S4 200 cycles kits version 1.5. A more
detailed description of DNA extraction and library preparation can be found in Supplementary Note 1.
693
Basic bioinformatics
The sequencing data was demultiplexed using the Illumina software BCL Convert
(https://emea.support.illumina.com/sequencing/sequencing_software/bcl-convert.html, Illumina Inc.) .
Adapter sequences were trimmed and overlapping reads were collapsed using AdapterRemoval
(2.2.4[46]). Single-end collapsed reads of at least 30bp and paired-end reads were mapped to the human
reference genome build 37 using bwa (0.7.17[47]) with seeding disabled to allow for higher sensitivity.
Paired- and single-end reads for each library and lane were merged, and duplicates were marked using
Picard MarkDuplicates (2.18.26, http://picard.sourceforge.net) with a pixel distance of 12000. Read
depth and coverage were determined using samtools (1.10[48]) with the all sites used in the calculation
(-a). Data was then merged to sample level and duplicates were marked again.
704
DNA authentication
In order to determine the authenticity of the ancient reads, post-mortem DNA damage patterns were
quantified using mapDamage2.0[49]. Next, two different methods were used to estimate the levels of
contamination. Firstly, we applied ContamMix in order to quantify the fraction of exogenous reads in
the mitochondrial reads by comparing the mtDNA consensus genome to possible contaminant
genomes[50]. The consensus was constructed using an in-house perl script that used sites with at least 5x
coverage, and bases were only called if observed in at least 70% of reads covering the site. Lastly, we
applied ANGSD (0.931[51]) to estimate nuclear contamination by quantifying heterozygosity
on the X chromosome in males. Both contamination estimates only used filtered reads
with a base quality of ≥20 and mapping quality of ≥30.
715
Imputation
We combined the 86 newly sequenced Medieval and post-Medieval Danish individuals with 1,664
previously published ancient genomes[9]. We then excluded individuals showing:
contamination (more than 5%); low autosomal coverage (less than 0.1 X); low
genome−wide average imputation genotype probability (less than 0.98), and we chose
the best quality sample in a close relative pair (first or second degree relative). A total
of 1,557 individuals passed all filters, and were used in downstream analyses. We
restricted the analysis to SNPs with imputation INFO score ≥ 0.5 and MAF ≥ 0.05.
724
725   **Kinship analysis and uniparental haplogroup inferences**
726   READ[52] was used to detect the degree of relatedness between pairs of individuals.

727 The mtDNA haplogroups of the Medieval and post-Medieval individuals were assigned using
728 HaploGrep2[53]. Y chromosome haplogroup assignment was inferred following the workflow already
729 published[54]. More details can be found in Supplementary Note 2.
730
Population genetic analyses
We used principal component analysis (PCA) (Supplementary Figure 1.1) to investigate the overall
population structure of the dataset. We used plink[55], excluding SNPs with minor allele frequency
(MAF) < 0.05 in the imputed panel. Based on 1,210 ancient western Eurasia imputed genomes, the
Medieval and post-Medieval samples cluster very close to each other, displaying a relatively low
genetic variability and situated within the genetic variability observed in the post-Bronze Age western
Eurasian populations.
738
We used three methods to estimate ancestry components in our ancient samples: model-based
clustering (ADMIXTURE[56]) (Supplementary Note 1, Figure S1.1) on a subset of 826,248 SNPs;
qpAdm[57] (Supplementary Note 1 Figure S1.2 and Table S1.1) with a reference panel of three genetic
ancestries (WHG, Neolithic Farmer, and Steppe) on the same 826,248 SNPs. We performed qpAdm
applying the option "allsnps: YES" and a set of 7 outgroups was used as "right populations":
Siberia_UpperPaleolithic_UstIshim, Siberia_UpperPaleolithic_Yana,
Russia_UpperPaleolithic_Sunghir, Switzerland_Mesolithic, Iran_Neolithic, Siberia_Neolithic,
USA_Beringia. We set a minimum threshold of 100,000 SNPs and only results with $p > 0.05$ only
have been considered. Finally we ran chromosome painting[58] using a panel of 7 ancestries (as on the
UK Biobank). We ran chromosome painting on all ancient individuals not in the reference panel,
using a reference panel of ancient donors grouped into populations to represent specific ancestries:
western hunter-gatherer (WHG), eastern hunter-gatherer (EHG), Caucasus hunter-gatherer (CHG),
Neolithic Farmer, Yamnaya, African and EastAsian (method described in [9] Supplementary Note 3h).
Painting followed the pipeline of [59] based on GLOBETROTTER[60], with admixture proportions
estimated using Non-Negative Least squares. We also painted individuals born in Denmark of a
typical ancestry based on density-based clustering of the first 18 PCs[9]. This generated both local
ancestry probabilities and genome-wide ancestry fractions for each painted individual. The reference
panel used for chromosome painting was designed to capture the various components of European
ancestry only, and so we urge caution in interpreting these results for non-European samples.
758
This dataset provides the opportunity to study the population history of Denmark from the Mesolithic
to the post-Medieval period, covering around 10,000 years, which can be considered a typical
Northern European population. Our results clearly demonstrate the impact of previously described
demographic events, including the influx of Neolithic Farmer ancestry ~9,000 years ago and Steppe
ancestry ~5,000 years ago[12, 10]. We highlight genetic continuity from the Bronze Age to the post-
Medieval period (Supplementary Note 1 Figure S1.1), although *qpAdm* detected a small increase in
the Neolithic Farmer component during the Viking Age (Supplementary Note 1 Figure S1.2 and
Table S1.1), while the Medieval period marked a time of increased genetic diversity, likely reflecting
increased mobility across Europe. This genetic continuity is further confirmed by the haplogroups
identified in the uniparental genetic makers (Supplementary Note 2). Together, these results suggest
that after the Bronze Age Steppe migration there was no other major gene flow into Denmark from
populations with significantly different Neolithic and Bronze Age ancestry compositions, and
therefore no changes in these ancestry components in the Danish population.
772

Local ancestry
We used two estimates of local ancestry from [9]: (1) first coalescent labels generated by running
Chromopainter[58] on all "White British" individuals in the UK Biobank, using the same reference
panel described above. Henceforth 'local ancestry'. (2) Ancestry path labels in GBR, FIN and TSI
1000G populations[61]) and 1015 ancient genomes generated using a neural network to assign ancestry
paths based on a sample's nearest neighbours at the first five informative nodes of a marginal tree
sequence, where an informative node is defined as one which has at least one leaf from the reference
set of ancient samples described above ([9] Supplementary Note S3i). Henceforth 'ancestry path labels'.
781
SNP associations
We aimed to generate SNP associations from previous studies for each phenotype in a consistent
approach. To generate a list of SNPs associated with multiple sclerosis (MS), rheumatoid arthritis
(RA) and celiac disease (CD), we used two approaches: in the first, we downloaded fine-mapped
SNPs from previous association studies. For each fine-mapped SNP, if the SNP did not have an
ancestry path label, we found the SNP in highest LD that did, with a minimum threshold of $r^2 \geq 0.7$
in the GBR, FIN and TSI 1000G populations using LDLinkR[62]. The final SNPs used for each
phenotype can be found in ST4 (MS), ST5 (RA), and ST6 (CD).
790
791 For MS, we used data from [3]. For non-MHC SNPs, we used the 'discovery' SNPs with P(joined) and
792 OR(joined) generated in the replication phase. For MHC variants, we searched the literature for the
793 reported HLA alleles and amino-acid polymorphisms (ST3). In total, we generated 205 SNPs which
794 were either fine-mapped or in high LD with a fine-mapped SNP (15 MHC, 190 non-MHC).
795
796 For RA, we downloaded 57 genome-wide significant non-MHC SNPs for seropositive RA in
797 Europeans[63]. We retrieved MHC associations separately ([64], with associated ORs and p-values from
798 [65]). In total, we generated 51 SNPs which were either fine-mapped or in high LD with a fine-mapped
799 SNP (3 MHC, 48 non-MHC).
800
801 For CD, we retrieved non-MHC SNPs from [66]. We used MHC SNPs from [67], with associated ORs and
802 p-values from [68]. In total, this generated 32 SNPs which were either fine-mapped or in high LD with a
803 fine-mapped SNP (3 MHC, 29 non-MHC).
804
Secondly, because we could not always find tag SNPs for fine-mapped SNPs that were present in our
ancestry path labels dataset, we found that we were losing significant signal from the HLA, therefore
we generated a second set of SNP associations. We downloaded full summary statistics for each
disease (MS: [3]; RA: [69], CD: http://www.nealelab.is/uk-biobank/), restricted to sites present in the
ancestry path labels dataset, and ran Plink's (PLINK v1.90b4.4[70]) clump method (parameters: --
clump-p1 5e-8 --clump-r2 0.05 --clump-kb 250 as in [71] using LD in the GBR, FIN and TSI 1000G
populations[61] to extract genome-wide significant independent SNPs.
812
813 In the main text we report results for the first set of SNPs ('fine-mapped') for analyses involving local
814 ancestry in modern data, and the second set of SNPs ('pruned') for analyses involving polygenic
815 measures of selection (CLUES/PALM).
816
REGIONS OF UNUSUAL ANCESTRY AND GENE ENRICHMENT
To assess which regions of ancestry were unusual, we converted the ancestry estimates to a Z-score.
Specifically, we let $A(i, j, j)$ denote the probability of the kth ancestry ($k = 1, \ldots, K$) at the jth SNP
($j = 1, \ldots, J$) of a chromosome for the ith individual ($i = 1, \ldots, N$). We then computed the mean

painting for each SNP, $A(j,k) = \frac{1}{N}\sum_{i=1}^{N} \quad A(i,j,k)$. From this we estimated a scale parameter μ_k

and deviation parameter σ_k using a block-median approach. Specifically we partitioned the genome

into 0.5Mb regions, and within each, computed the mean and standard deviation of the ancestry. The

parameter estimates are then the median values over the whole genome. We then computed an

anomaly score for each SNP for each ancestry $Z(j,k) = (A(j,k) - \mu_k)/\sigma_k$.

To arrive at an anomaly score for each SNP aggregated over all ancestries, we also had to account for

correlations in the ancestry paintings. Instead of scaling each ancestry deviation $A^*(j,k) = A(j,k) - $

μ_k by its standard deviation, we instead "whitened" them, i.e. rotated the data to have an independent

signal. Let $C = A^{*T}A^*$ by a $K \times K$ covariance matrix, and let $C^{-1} = UDV^T$ be the Singular Value

Decomposition. Then $W = UD^{1/2}$ is the whitening matrix from which $Z = A^*W$ are normally

distributed with covariance matrix diag(1) under the null that A^* is normally distributed with mean 0

and unknown covariance Σ. The "ancestry anomaly score" test statistic for each SNP is $t(j) = $

$\sum_{k=1}^{K} \quad Z(j,k)^2$, which is Chi-squared distributed with K degrees of freedom under the null, and we

reported p-values from this.

To test for gene enrichment we formed a list of all SNPs reaching genome-wide significance ($p < $

5^{-8}) and using the R package *gprofiler2*[72] converted these to a unique list of genes. We then used *gost*

to perform an enrichment test for each GO term, for which we used default p-value correction via the

g:Profiler SCS method. This is an empirical correction based on performing random lookups of the

same number of genes under the null, to control the error rate and ensure that 95% of reported

categories (at p=0.05) are correct.

ALLELE FREQUENCY PLOTS OVER TIME

To investigate how effect allele frequencies have changed over time, we extracted high effect alleles

for each phenotype from the ancient data. We excluded all non-Eurasian samples, grouped them by

'groupLabel', excluded any group with fewer than 4 samples, and coloured points by ancestry

proportion according to genome-wide NNLS based on chromosome painting (above).

CLUSTER ANALYSIS

In order to understand whether risk-conferring haplotypes evolved in the Steppe population, or in a

pre- or post-dating population in which Steppe ancestry is high, we used k-means clustering on the

dosage of each ancestry for each selected significant SNP and investigated the dosage distribution of

clusters with significantly higher MS prevalence. For the target SNPs, the Elbow method[73]. suggested

selecting around 5-7 clusters, of which we chose 6. After performing the k-means cluster analysis, we

calculated the average probability for each ancestry for case individuals. Furthermore, we calculated

the prevalence of MS in each cluster, and performed a one-sample t-test to investigate whether it

differs from the overall MS prevalence (0.487%). This tests whether particular combinations of

ancestry are associated with the phenotype at a SNP. Clusters with high MS risk-ratios have high

Steppe components (Supplementary Figure 4.2), leading to the conclusion that Steppe ancestry alone

is driving this signal.

WEIGHTED AVERAGE PREVALENCE

In order to quantify the risk of each ancestry for each SNP, we calculated the weighted average

prevalence (WAP) for each ancestry based on the result of k-means clustering (above).

For the jth SNP, let $P_{jkm} = n_{jm}\underline{P}_{jkm}$ denote the sum of the kth ancestry probabilities of all the
individuals in the mth cluster ($k, m = 1,\ldots,6$), where n_{jm} is the cluster size of the mth cluster. Let
π_{jm} denote the prevalence of MS in the mth cluster, the weighted average prevalence for the kth
ancestry is defined as:

871
$$\underline{\pi}_{jk} = \frac{P_{jkm}\pi_{jm}}{\sum_{m=1}^{6} P_{jkm}},$$

where P_{jkm} is defined as the weight for each cluster.

For each ancestry, WAP measures the association of that ancestry with MS risk across all clusters. To
make a clear comparison, we calculated the risk ratio (compared to the overall MS prevalence) for
each ancestry at each SNP, and assigned a mean and confidence interval for the risk ratios of each
ancestry at each chromosome (Figure3, Supplementary Figure 3.1 and 3.2).

PCA/UMAP OF WAP/AVERAGE DOSAGE
We performed principal component analysis (PCA) on the average ancestry probability and WAP at
each MS-associated SNP (Supplementary Figure 4.3). The former shows that all of the HLA SNPs
except three from HLA class II and III have much larger Outgroup components compared with the
others. The latter analysis indicates a strong association between Steppe and MS risk. Also, Outgroup
ancestry at rs10914539 from chromosome 1 exceptionally reduces the incidence of MS, while
Outgroup ancestry at rs771767 (chromosome 3) and rs137956 (chromosome 22) significantly boosts
MS risk.

ANCESTRAL RISK SCORES
Following methods developed in Irving-Pease et al. (*submitted*), we calculated the effect allele
painting frequency for a given ancestry $f_{\{anc,i\}}$ for SNP i using the formula:

891
$$f_{\{anc,i\}} = \frac{\sum_{j}^{M_{effect}} Painting\ certainty_{\{j,i,anc\}}}{\sum_{j}^{M_{alt}} Painting\ certainty_{\{j,i,anc\}} + \sum_{j}^{M_{effect}} Painting\ certainty_{\{j,i,anc\}}},$$

where there are M_{effect} individuals homozygous for the effect allele, M_{alt} individuals homozygous
for the other allele, and $\sum_{j}^{M_{effect}} Painting\ certainty_{\{j,i,anc\}}$ is the sum of the painting
probabilities for that ancestry anc in individuals homozygous for the effect allele at SNP i. This can
be interpreted as an estimate of an ancestral contribution to effect allele frequency in a modern
population. Per-SNP painting frequencies can be found in ST4, ST5, and ST6.

To calculate the ancestral risk score (ARS) we summed over all I pruned SNPs in an additive model:

899
$$ARS_{anc} = \sum_{i}^{I} f_{\{anc,i\}} * beta_i.$$

We then ran a transformation step as in [74]. To obtain 95% confidence intervals, we ran an accelerated
bootstrap over loci, which accounts for the skew of data to better estimate confidence intervals [75].

LOCAL ANCESTRY AND GENOTYPE GWAS
We used the UK Biobank to fit GWAS models for local ancestry values and genotype values
separately, using only SNPs known to be associated with the phenotype ('fine-mapped' SNPs). We

used the following phenotype codes for each phenotype: MS: Data-Field 131043; RA: Data-Field
131849 (seropositive); CD: Data-Field 21068.

Let Y_i denote the phenotype status for the ith individual ($i = 1,\ldots,399998$), which takes value 1 for a
case and 0 for control, and let $\pi_i = Pr(Y_i = 1)$ denote the probability that this individual has the
event. Let X_{ijk} denote the kth ancestry probability ($k = 1,\ldots,K$) for the jth SNP ($j = 1,\ldots,205$) of
the ith individual. C_{ic} is the cth predictor ($c = 1,\ldots,N_c$) for the ith individual. We used the following
logistic regression model for GWAS, which assumes the effects of alleles are additive:

915
$$Y_i \sim Bin(1, \pi_i); \ log(\frac{\pi_i}{1 - \pi_i}) = \sum_{k=1}^{K} \beta_{jk} X_{ijk} + \sum_{c=1}^{N_c} \gamma_c C_{ic}.$$

We used N_c=20 predictors in the GWAS models, including sex, age and the first 18 PCs, which are
sufficient to capture most of the population structure in the UK Biobank[76].

First, we built the model with $K = 1$. By using only one ancestry probability in each model, we aimed
to find the statistical significance of each SNP under each ancestry. Then, we built the model with
$K = 5$, i.e. using all 6 local ancestry probabilities which sum to 1. We calculated the variance
explained by each SNP by summing up the variance explained by X_{ijk}(k=1,...,5).

We considered fitting the multivariate models by using all the SNPs as covariates. However, the
dataset only contains 1,982 cases. Even though only one ancestry is included, the multivariate model
contains 191 predictors, which could result in overfitting problems. Therefore, the GWAS models are
preferred over multivariate models.

We also fitted a logistic regression model for GWAS using the genotype data as follows:

931
$$Y_i \sim Bin(1, \pi_i); \ log(\frac{\pi_i}{1 - \pi_i}) = \beta_j X_{ij} + \sum_{c=1}^{N_c} \gamma_c C_{ic},$$

where $X_{ij} \in \{0,1,2\}$ denotes the number of copies of the reference allele of the jth SNP ($j = $
$1,\ldots,205$) that the ith individual has, and C_{ic} ($c = 1,\ldots,N_c$) denotes the covariates including age, sex
and first 18 PCs for the ith individual, where N_c=20. Due to the UK Biobank being underpowered
compared to the Case-Control study from which these SNPs were found, the only statistically
significant (at $p < 10^{-5}$) association is in the HLA class II tagging HLA-DRB1*15:01.

COMPARISON OF GWAS MODELS USING PAINTING AND GENOTYPE DATA
We compared the variance explained by SNPs from the GWAS model using the painting data (all 6
local ancestry probabilities) with that from GWAS model using the genotype data. McFadden's
pseudo R squared measure[77] is widely used for estimating the variance explained by the logistic
regression models. McFadden's pseudo R squared is defined as

943
$$R^2 = 1 - \frac{ln(L_M)}{lm(L_0)},$$

944   where L_M and 0 are the likelihoods for the fitted and the null model, respectively. Taking overfitting
945   into account, we propose the adjusted McFadden's pseudo R squared by penalizing the number of
946   predictors:

947
$$Adjusted \ R^2 = 1 - \frac{ln(L_M)/(N - k)}{ln(L_0)/(N - 1)},$$

948   where N is the sample size and k is the number of predictors.
949
950   Specifically, $R^2(SNPs)$ is calculated as the extra variance in addition to sex, age and 18 PCs that can
951   be explained by SNPs:
952 $$R^2(SNPs) = R^2(sex + age + 18\,PCs + SNPs) - R^2(sex + age + 18\,PCs).$$
953
Notably, two SNPs stand out for explaining much larger variance than others when fitting the GWAS
model using the genotype data, but overall more SNPs from GWAS painting explain more than 0.1%
variance, which indicates the painting data are probably more efficient for estimating the effect sizes
of SNPs and detecting significant SNPs. Also, some SNPs from GWAS models using painting data
explain almost the same amount of variance, suggesting that these SNPs consist of very similar
ancestries.
960
HAPLOTYPE TREND REGRESSION WITH eXtra FLEXIBILITY (HTRX)
We propose Haplotype Trend Regression with eXtra flexibility (HTRX) which searches for haplotype
patterns that include single SNPs and non-contiguous haplotypes. HTRX is an association between a
template of n SNPs and a phenotype. A template gives a value for each SNP taking values of '0' or
'1', reflecting whether the reference allele of each SNP is present or absent, or an 'X' meaning either
value is allowed. For example, haplotype '1X0' corresponds to a 3-SNP haplotype where the first
SNP is the alternative allele and the third SNP is the reference allele, while the second SNP can be
either the reference or the alternative allele. Therefore, haplotype '1X0' is essentially only a 2-SNP
haplotype.
970
971   To examine the association between a haplotype and a binary phenotype, we replace the genotype
972   term with a haplotype from the standard GWAS model:
973 $$Y_i \sim Bin(1, \pi_i); \; log\left(\frac{\pi_i}{1-\pi_i}\right) = \beta_j H_{ij} + \sum_{c=1}^{N_c} \gamma_c \, C_{ic},$$
974   where H_{ij} denotes the jth haplotype probability for the ith individual:

$$H_{ij} = \begin{cases} 1 & \text{if } i\text{th individual has haplotype } j \text{ in both genomes,} \\ 1/2 & \text{if } i\text{th individual has haplotype } j \text{ in one of the two genomes,} \\ 0 & \text{otherwise.} \end{cases}$$

975
976
977   HTRX can identify gene-gene interactions, and is superior to HTR not only because it can extract
978   combinations of significant SNPs within a region, leading to improved predictive performance, but
979   the haplotypes are more interpretable as multi-SNP haplotypes are only reported when they lead to
980   increased predictive performance.
981
982   **HTRX Model selection procedure for shorter haplotypes**
983   Fitting HTRX models directly on the whole dataset can lead to significant overfitting, especially when
984   the number of SNPs increases. When overfitting occurs, the models experience poorer predictive
985   accuracy against unseen data. Further, HTRX introduces an enormous model space which much be
986   searched.
987
988   To address these problems, we implement a two-step procedure.
989

990   Step 1: select **candidate** models. This is to address the model search problem, and is chosen to obtain
991   a set of models more diverse than traditional bootstrap resampling (Efron, 1979[78]).
992
993   (1) Randomly sample a subset (50%) of data. Specifically, when the outcome is binary, stratified
994   sampling is used to ensure the subset has approximately the same proportion of cases and controls as
995   the whole data;
996
997   (2) Start from a model with fixed covariates (18 PCs, sex and age), and perform forward regression on
998   the subset, i.e. iteratively choose a feature (in addition to the fixed covariates) to add whose inclusion
999   enables the model to explain the largest variance, and select s models with the lowest Bayesian
1000   Information Criteria (BIC)[79] to enter the candidate model pool;
1001
1002   (3) repeat (1)-(2) B times, and select all the different models in the candidate model pool as the
1003   candidate models.
1004
1005   Step 2: select the best model using 10-fold cross-validation.
1006
1007   (1) Randomly split the whole data into 10 groups with approximately equal sizes, using stratified
1008   sampling when the outcome is binary;
1009
1010   (2) In each of the 10 folds, use a different group as the test dataset, and take the remaining groups as
1011   the training dataset. Then, fit all the candidate models on the training dataset, and use these fitted
1012   models to compute the additional variance explained by features (out-of-sample R^2) in the test
1013   dataset. Finally, select the candidate model with the biggest average out-of-sample R^2 as the best
1014   model.
1015
1016   **HTRX Model selection procedure for longer haplotypes (Cumulative HTRX)**
1017   Longer haplotypes are important for discovering interactions. However, there are $3^k - 1$ haplotypes
1018   in HTRX if the region contains k SNPs, making it unrealistic for regions with large numbers of SNPs.
1019   To address this issue, we proposed cumulative HTRX to control the number of haplotypes, which is
1020   also a two-step procedure.
1021
1022   Step 1: extend haplotypes and select candidate models.
1023
1024   (1) Randomly sample a subset (50%) of data, use stratified sampling when the outcome is binary. This
1025   subset is used for all the analysis in (2) and (3);
1026
(2) Start with L randomly chosen SNPs from the entire k SNPs, and keep the top M haplotypes that
are chosen from the forward regression. Then add another SNP to the M haplotypes to create $3M + 2$
haplotypes. There are $3M$ haplotypes obtained by adding `0', `1' or `X' to the previous M haplotypes,
as well as 2 bases of the added SNP, i.e. `XX...X0' and `XX...X1' (as `X' was implicitly used in the
previous step). The top M haplotypes from them are then selected using forward regression. Repeat
this process until obtaining M haplotypes which include $k - 1$ SNPs;
1033
1034   (3) Add the last SNP to create $3M + 2$ haplotypes. Afterwards, start from a model with fixed
1035   covariates (18 PCs, sex and age), perform forward regression on the training set, and select s models
1036   with the lowest BIC to enter the candidate model pool;

1037

1038 (4) repeat (1)-(3) B times, and select all the different models in the candidate model pool as the
1039 candidate models.

1040

1041 Step 2: select the best model using 10-fold cross-validation, as described in "**HTRX Model selection**
1042 **procedure for shorter haplotypes**".

1043

1044 We note that because the search procedure in Step 1(2) may miss some highly predictive haplotypes,
1045 cumulative HTRX acts as a lower bound on the variance explainable by HTRX.

1046

As a model criticism, only common and highly predictive haplotypes (i.e. those with the greatest
adjusted R^2) are correctly identified, but the increased complexity of the search space of HTRX leads
to haplotype subsets that are not significant on their own but are significant when interacting with
other haplotype subsets being missed. This issue would be eased if we increase all the parameters s, l,
M and B but with higher computational cost, or improve the search by optimizing the order of adding
SNPs. This leads to a decreased certainty that the exact haplotypes proposed are `correct', but together
reinforces the inference that interaction is extremely important.

1054

Simulation for HTRX
To investigate how the total variance explained by HTRX compare to GWAS and HTR, we used a
simulation study comparing:
(1) linear models (denoted by "lm") and generalized linear models with a logit link-function (denoted
by "glm");
(2) models with or without actual interaction effects;
(3) models with or without rare SNPs (frequency smaller than 5%);
(4) remove or retain rare haplotypes when rare SNPs exist.

1063

We started from creating the genotypes for 4 different SNPs G_{ijq} ($i = 1,\ldots,100,000$ denotes the
index of individuals, $j = 1("1XXX"), 2("X1XX"), 3("XX1X") \ and \ 4("XXX1")$ represents the index
of SNPs, and $q = 1,2$ for two genomes as individuals are diploid). If no rare SNPs were included, we
sampled the frequency F_j of these 4 SNPs from 5% to 95%; otherwise, we sampled the frequency of
the first 2 SNPs from 2% to 5% (in practice, we obtained $F_1 = 2.8\%$ and $F_2 = 3.1\%$ under our seed)
while the last 2 SNPs from 5% to 95%. For the ith individual, we sampled $G_{ijq} \sim Bin(1, F_j)$ for the qth
genome of the jth SNP, and took the average value of two genomes as the genotype for the jth SNP of
the ith individual: $G_{ij} = \frac{G_{ij1} + G_{ij2}}{2}$. Based on the genotype data, we obtained the haplotype data for
each individual, and we considered removing haplotypes rarer than 0.1% or not when rare SNPs were
generated. In addition, we sampled 20 fixed covariates (including sex, age and 18 PCs) C_{ic} where $c =$
$1,\ldots,20$ from UK Biobank for 100000 individuals.

1075

1076 Next, we sampled the effect sizes of SNPs β_{G_j} and covariates β_{C_c}, and standardize them by their
1077 standard deviations: $\beta_{G_i} \sim \frac{U(-1,1)}{sd(G_j)}$ and $\beta_{C_c} \sim \frac{U(-1,1)}{sd(C_c)}$ for each fixed j and c, respectively. When
1078 interaction exists, we created a fixed effect size for haplotype "11XX" as twice the average absolute
1079 SNP effects: $\beta_{H_1} = \frac{1}{2}\sum_{j=1}^{4} \ \ |\beta_{G_j}|$ where H_1 refers to "11XX"; otherwise, $H_1 = 0$. Note that $F_{H_1} =$
1080 0.09% when rare SNPs are included.

1081

Finally, we sampled the outcome based on the outcome score (for the ith individual)

$$O_i = \sum_{c=1}^{20} \quad \beta_c C_{ic} + \gamma(\sum_{j=1}^{4} \quad \beta_{G_j} G_{ij} + \beta_{H_1} H_1) + e_i + w,$$

where γ is the effect scale of SNPs and haplotype "11XX", $e_i \sim N(0,0.1)$ is the random error and w is

a fixed intercept term. For linear models, the outcome $Y_i = O_i$; while for generalized linear models,

we sampled the outcome from binomial distribution: $Y_i \sim Bin(1, \pi_i)$, where $\pi_i = \frac{e^{O_i}}{1+e^{O_i}}$ is the

probability that the i th individual has the case.

1088

1089 As the simulation is intended to compare the variance explained by HTRX, HTR and SNPs (GWAS)

1090 in addition to fixed covariates, we tripled the effect sizes of SNPs and haplotype "11XX" (if

1091 interaction exists) by setting $\gamma = 3$. In "glm", to ensure a reasonable case prevalence (e.g. below 5%),

1092 we set $w = -7$, which was also applied in "lm".

1093

We applied the procedure described in " **HTRX Model selection procedure for shorter haplotypes**"

for HTRX, HTR and GWAS, and visualized the distribution of the out-of-sample R^2 for each of the

best models selected by each method in Supplementary Figure 4.4. In both "lm" and "glm", HTRX

has equal predictive performance as the true model. It performs as well as GWAS when the

interaction effects is absent, explains more variance when an interaction is present, and is significantly

more explanatory than HTR. When rare SNPs are included, the only effective interaction term is rare.

In this case the difference between GWAS and HTRX becomes smaller as expected, and removing the

rare haplotypes hardly reduces the performance of HTRX.

1102

1103 In conclusion, we demonstrate through simulation that our HTRX implementation a) searches

1104 haplotype space effectively, and b) protects against overfitting. This makes it a superior approach

1105 compared to HTR and GWAS to integration SNP effects with gene-gene interaction. Its robustness

1106 also retains when there are rare effective SNPs and haplotypes.

1107

POLYGENIC SELECTION TEST

We inferred allele frequency trajectories and selection coefficients for a set of LD-puned genome-

wide significant trait associated variants using a modified version of the software CLUES[19]. To

account for population structure in our samples, we applied a novel chromosome painting technique

based on inference of a sample's nearest neighbours in the marginal trees of an ARG that contains

labelled individuals (Irving-Pease et al., *submitted*). We ran CLUES using a time-series of imputed

aDNA genotype probabilities obtained from 1,015 ancient West Eurasian samples that passed all

quality control filters. We produced four additional models for each trait associated variant, by

conditioning the analysis on one of the four ancestral path labels from our chromosome painting

model: either Western hunter-gatherers (WHG), Eastern hunter-gatherers (EHG), Caucasus hunter-

gatherers (CHG), or Anatolian farmers (ANA). We then inferred polygenic selection gradients (ω)

and p-values for each of MS, CD and RA, in all ancestral paths, using the software PALM[20]. Full

methods and results can be found in Supplementary Note 6.

1121

1122 **ANCESTRY LINKAGE DISEQUILIBRIUM (LDA) AND ANCESTRY LINKAGE**

1123 **DISEQUILIBRIUM SCORE (LDAS)**

1124 In population genetics, linkage disequilibrium (LD) is defined as the non-random association of

1125 alleles at different loci in a given population[80]. We propose an ancestry linkage disequilibrium (LDA)

1126 approach to measure the association of ancestries between SNPs.

1127

1128 Let $A(i, j, k)$ denote the probability of the kth ancestry ($k = 1, \ldots, K$) at the jth SNP ($j = 1, \ldots, J$) of a
1129 chromosome for the ith individual ($i = 1, \ldots, N$).

1130

1131 We define the distance between SNP l and m as the average L_2 norm between ancestries at those
1132 SNPs. Specifically we compute the L_2 norm for the ith genome as

1133 $$D_i(l, m) = ||A(i, l, \cdot) - A(i, m, \cdot)||_2 = \sqrt{\frac{1}{K}\sum_{k=1}^{K} \quad (A(i, l, k) - A(i, m, k))^2}.$$

1134

1135 Then we compute the distance between SNP l and m by averaging $D_i(l, m)$:

1136 $$D(l, m) = \frac{1}{N}\sum_{i=1}^{N} \quad D_i(l, m).$$

1137

We define $D^*(l, m)$ as the theoretical distance between SNP l and m if there were no linkage
disequilibrium of ancestry (LDA) between them. $D^*(l, m)$ is estimated by

$$D^*(l, m) \approx \frac{1}{N}\sum_{i=1}^{N} \quad ||A(i^*, l, \cdot) - A(i, m, \cdot)||_2,$$

where $i^* \in \{1, \ldots, N\}$ are re-sampled without replacement at SNP l. Using the empirical distribution
of ancestry probabilities accounts for variability in both the average ancestry and its distribution
across SNPs. Ancestry assignment can be very precise in regions of the genome where our reference
panel matches our data, and uncertain in others where we only have distant relatives of the underlying
populations.

1146

1147 The LDA between SNP l and m is a similarity, defined in terms of the negative distance $-D(l, m)$
1148 normalized by the expected value $D^*(l, m)$ under no LD, as:

1149 $$LDA(l, m) = \frac{D^*(l, m) - D(l, m)}{D^*(l, m)}.$$

1150

1151 LDA therefore takes an expected value 0 when haplotypes are randomly assigned at different SNPs,
1152 and positive values when the ancestries of haplotypes are correlated.

1153

LDA is a pairwise quantity. To arrive at a per-SNP property, we define the LDA score (LDAS) of
SNP j as the total LDA of this SNP with the rest of the genome, i.e. the integral of the LDA for that
SNP. Because this quantity decreases to zero as we move away from the target SNP, this is in practice
computed within an XcM-window (we use $X = 5$ as LDA is approximately zero outside this region in
our data) on both sides of the SNP. Note that we measure this quantity in terms of the genetic
distance, and therefore LDAS is measuring the length of ancestry-specific haplotypes compared to
individual-level recombination rates.

1161

1162 As a technical note, when the SNPs approach either end of the chromosome, they no longer have a
1163 complete window, which results in a smaller LDAS. This would be appropriate for measuring total
1164 ancestry correlations, but to make LDAS useful for detecting anomalous SNPs, we use the LDAS of
1165 the symmetric side of the SNP to estimate the LDAS within the non-existent window.

1166

$$LDAS(j; X) = \begin{cases} \int_{gd(j)-X}^{gd(j)+X} LDA(j, l) \, dgd & \text{if } X \le gd(j) \le tg - X, \\ \int_{0}^{gd(j)+X} LDA(j, l) \, dgd + \int_{2gd(j)}^{gd(j)+X} LDA(j, l) \, dgd & \text{if } gd(j) < X, \\ \int_{gd(j)-X}^{tg} LDA(j, l) \, dgd + \int_{gd(j)-X}^{2gd(j)-tg} LDA(j, l) \, dgd & \text{if } gd(j) > tg - X. \end{cases}$$

1167
1168

1169   where $gd(l)$ is the genetic distance (i.e. position in cM) of SNP l, and tg is the total genetic distance

1170   of a chromosome. We also assume the LDA on either end of the chromosome equals the LDA of the

1171   SNP closest to the end: $LDA(j, gd = 0) = LDA(j, l_{min(gd)})$ and $LDA(j, gd = td) =$

1172 $LDA(j, l_{max(gd)})$, where gd is the genetic distance, $l_{min(gd)}$ and $l_{max(gd)}$ are the indexes of the SNP

1173   with the smallest and largest genetic distance, respectively.

1174

1175   The integral $\int_{gd(j)-X}^{gd(j)+X} LDA(j,l)dgd$ is computed assuming linear interpolation of the LDA score

1176   between adjacent SNPs.

1177

1178   LDA thus quantifies the correlations between the ancestry of two SNPs, measuring the proportion of

1179   individuals who have experienced a recombination leading to a change in ancestry, relative to the

1180   genome-wide baseline. The LDA score is the total amount of genome in LDA with each SNP

1181   (measured in recombination map distance).

1182

SIMULATION FOR SELECTION: LDA

An ancient population P_0 evolved for 2200 generations before splitting into two sub-populations P_1

(Steppe) and P_2 (Farmer). After evolving 400 generations, we added mutation "m_1" and "m_2" at the

different locus in P_1 and P_2. Both added mutations were then positively selected in the following 300

generations, after which they merged to P_3, where both added mutations experienced strong positive

selection for 20 generations. Finally, we sampled 1000 individuals from P_3 to compute their ancestry

proportions of P_1 and P_2 using the "chromosome painting" technique, and calculated the LDA score

of the simulated chromosome positions.

1191

1192   The above describes the simulation in Supplementary Figure 6.1.

1193

We investigated balancing selection at 2 loci as well. The balancing selection in P_1 and P_2 ensured the

mutated allele reaches around 50% frequency, while positive selection made the mutated allele

become almost the only allele. In P_3, if m_1 or m_2 was positively selected, its frequency reached more

than 80% regardless of whether the allele experienced balancing or positive selection in P_1 or P_2,

because we set a strong positive selection. If m_1 or m_2 was balancing selected in P_3, its frequency

slightly increased, e.g. if m_1 underwent balancing selection in P_1, it had 25% frequency when P_3 was

created, and the frequency reached around 37.5% after 20 generations of balancing selection in P_3.

1201

1202   The results (Supplementary Figure 6.2) show that positive selection in P_3 resulted in low LDA scores

1203   around the selected locus, if this allele was not uncommon (i.e. had 50% (balancing selection) or

1204   100% frequency (positive selection) in subpopulation P_1 or P_2). Note that the balancing selection in

1205 P_1 or P_2 worked the same as "weak positive selection", because m_1 and m_2 were rare when they first

1206   occurred, which were positively selected until 50% frequency.

1207

1208   We also performed simulations for selection at a single locus (Supplementary Figure 6.2&6.3).

1209

1210   Stage 1: We added a mutation m_1 in the 1600 generation in P_0, which then underwent balancing

1211   selection until generation 2200, when P_0 split into P_1 and P_2, where the frequency of m_1 was around

1212   50%.

1213

1214   Stage 2: Then we explored different combinations of positive, balancing and negative selection of m_1
1215   in P_1 and P_2. the frequency of m_1 reached 80%, 50% and 20% when it was positively, balancing or
1216   negatively selected, respectively, until generation 2899. Here we sampled 20 individuals each in P_1
1217   and P_2 as the ancient samples.
1218
1219   Stage 3: Then P_1 and P_2 merged into P_3 in generation 2900. In P_3, for each combination of selection
1220   in Stage 2, we simulated positive, balancing and negative selection for m_1. The selection lasted for 20
1221   generations, and then we sampled 4000 individuals from P_3 as the modern population.
1222
1223   Results: when m_1 was positively selected in only one of P_1 and P_2, and it experienced negative
1224   selection in P_3, the LDA scores around the locus of m_1 were low. Otherwise, no abnormal LDA
1225   scores were found at m_1.
1226
1227
1228

REFERENCES

1. Walton, C. *et al.* Rising prevalence of multiple sclerosis worldwide: Insights from the Atlas of MS, third edition. *Mult. Scler. J.* **26**, 1816–1821 (2020).

2. Attfield, K. E., Jensen, L. T., Kaufmann, M., Friese, M. A. & Fugger, L. The immunology of multiple sclerosis. *Nat. Rev. Immunol.* (2022) doi:10.1038/s41577-022-00718-z.

3. International Multiple Sclerosis Genetics Consortium *et al.* Multiple sclerosis genomic map implicates peripheral immune cells and microglia in susceptibility. *Science* **365**, eaav7188 (2019).

4. Bjornevik, K. *et al.* Longitudinal analysis reveals high prevalence of Epstein-Barr virus associated with multiple sclerosis. *Science* **375**, 296–301 (2022).

5. Lanz, T. V. *et al.* Clonally expanded B cells in multiple sclerosis bind EBV EBNA1 and GlialCAM. *Nature* **603**, 321–327 (2022).

6. Olsson, T., Barcellos, L. F. & Alfredsson, L. Interactions between genetic, lifestyle and environmental risk factors for multiple sclerosis. *Nat. Rev. Neurol.* **13**, 25–36 (2017).

7. Benton, M. L. *et al.* The influence of evolutionary history on human health and disease. *Nat. Rev. Genet.* **22**, 269–283 (2021).

8. Chi, C. *et al.* Admixture mapping reveals evidence of differential multiple sclerosis risk by genetic ancestry. *PLOS Genet.* **15**, e1007808 (2019).

9. Allentoft, M. E. *et al. Population Genomics of Stone Age Eurasia.* http://biorxiv.org/lookup/doi/10.1101/2022.05.04.490594 (2022) doi:10.1101/2022.05.04.490594.

10. Haak, W. *et al.* Massive migration from the steppe was a source for Indo-European languages in Europe. *Nature* **522**, 207–211 (2015).

11. Jones, E. R. *et al.* Upper Palaeolithic genomes reveal deep roots of modern Eurasians. *Nat. Commun.* **6**, 8912 (2015).

12. Allentoft, M. E. *et al.* Population genomics of Bronze Age Eurasia. *Nature* **522**, 167–172 (2015).

13. Bycroft, C. *et al.* The UK Biobank resource with deep phenotyping and genomic data. *Nature* **562**, 203–209 (2018).

14. Itan, Y., Powell, A., Beaumont, M. A., Burger, J. & Thomas, M. G. The Origins of Lactase

Persistence in Europe. *PLoS Comput. Biol.* **5**, e1000491 (2009).

15.   Dehasque, M. *et al.* Inference of natural selection from ancient DNA. *Evol. Lett.* **4**, 94–108

(2020).

16.   Efron, B. Better Bootstrap Confidence Intervals. *J. Am. Stat. Assoc.* **82**, 171–185 (1987).

17.   Zaykin, D. V. *et al.* Testing Association of Statistically Inferred Haplotypes with Discrete and

Continuous Traits in Samples of Unrelated Individuals. *Hum. Hered.* **53**, 79–91 (2002).

18.   Thuesen, N. H., Klausen, M. S., Gopalakrishnan, S., Trolle, T. & Renaud, G. *Benchmarking*

freely available human leukocyte antigen typing algorithms across varying genes, coverages and

typing resolutions. http://biorxiv.org/lookup/doi/10.1101/2022.06.28.497888 (2022)

doi:10.1101/2022.06.28.497888.

19.   Stern, A. J., Wilton, P. R. & Nielsen, R. An approximate full-likelihood method for inferring

selection and allele frequency trajectories from DNA sequence data. *PLOS Genet.* **15**, e1008384

(2019).

20.   Stern, A. J., Speidel, L., Zaitlen, N. A. & Nielsen, R. Disentangling selection on genetically

correlated polygenic traits via whole-genome genealogies. *Am. J. Hum. Genet.* **108**, 219–239

(2021).

21.   Comabella, M. *et al.* Identification of a Novel Risk Locus for Multiple Sclerosis at 13q31.3

by a Pooled Genome-Wide Scan of 500,000 Single Nucleotide Polymorphisms. *PLoS ONE* **3**,

e3490 (2008).

22.   Bersaglieri, T. *et al.* Genetic Signatures of Strong Recent Positive Selection at the Lactase

Gene. *Am. J. Hum. Genet.* **74**, 1111–1120 (2004).

23.   He, Z., Dai, X., Beaumont, M. & Yu, F. Detecting and Quantifying Natural Selection at Two

Linked Loci from Time Series Data of Allele Frequencies with Forward-in-Time Simulations.

Genetics **216**, 521–541 (2020).

24.   Fugger, L., Jensen, L. T. & Rossjohn, J. Challenges, Progress, and Prospects of Developing

Therapies to Treat Autoimmune Diseases. *Cell* **181**, 63–80 (2020).

25.   Gregersen, J. W. *et al.* Functional epistasis on a common MHC haplotype associated with

multiple sclerosis. *Nature* **443**, 574–577 (2006).

26.   Wang, J. H. *et al.* Modeling the cumulative genetic risk for multiple sclerosis from genome-

wide association data. *Genome Med.* **3**, 3 (2011).

27.   Cotsapas, C. & Mitrovic, M. Genome-wide association studies of multiple sclerosis. *Clin.*

Transl. Immunol. **7**, e1018 (2018).

28.   Slim, L., Chatelain, C., Foucauld, H. de & Azencott, C.-A. A systematic analysis of gene–

gene interaction in multiple sclerosis. *BMC Med. Genomics* **15**, 100 (2022).

29.   Bos, K. I. *et al.* Pre-Columbian mycobacterial genomes reveal seals as a source of New World

human tuberculosis. *Nature* **514**, 494–497 (2014).

30.   Sabin, S. *et al.* A seventeenth-century Mycobacterium tuberculosis genome supports a

Neolithic emergence of the Mycobacterium tuberculosis complex. *Genome Biol.* **21**, 201 (2020).

31.   Rasmussen, S. *et al.* Early Divergent Strains of Yersinia pestis in Eurasia 5,000 Years Ago.

Cell **163**, 571–582 (2015).

32.   Spyrou, M. A. *et al.* Analysis of 3800-year-old Yersinia pestis genomes suggests Bronze Age

origin for bubonic plague. *Nat. Commun.* **9**, 2234 (2018).

33.   Rascovan, N. *et al.* Emergence and Spread of Basal Lineages of Yersinia pestis during the

Neolithic Decline. *Cell* **176**, 295-305.e10 (2019).

34.   Guellil, M. *et al.* Ancient herpes simplex 1 genomes reveal recent viral structure in Eurasia.

Sci. Adv. **8**, eabo4435.

35.   Pontremoli, C., Forni, D., Clerici, M., Cagliani, R. & Sironi, M. Possible European Origin of

Circulating Varicella Zoster Virus Strains. *J. Infect. Dis.* jiz227 (2019) doi:10.1093/infdis/jiz227.

36.   Tian, C. *et al.* Genome-wide association and HLA region fine-mapping studies identify

susceptibility loci for multiple common infections. *Nat. Commun.* **8**, 599 (2017).

37.   Krause-Kyora, B. *et al.* Ancient DNA study reveals HLA susceptibility locus for leprosy in

medieval Europeans. *Nat. Commun.* **9**, 1569 (2018).

38.   Wallin, M. T. *et al.* The prevalence of MS in the United States: A population-based estimate

using health claims data. *Neurology* **92**, e1029–e1040 (2019).

39.   Feigin, V. L. *et al.* Global, regional, and national burden of neurological disorders, 1990–

2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet Neurol.* **18**, 459–

480 (2019).

40.   Fleming, J. & Fabry, Z. The hygiene hypothesis and multiple sclerosis. *Ann. Neurol.* **61**, 85–

89 (2007).

41.   Joo, Y. B., Lim, Y.-H., Kim, K.-J., Park, K.-S. & Park, Y.-J. Respiratory viral infections and

the risk of rheumatoid arthritis. *Arthritis Res. Ther.* **21**, 199 (2019).

42.   Safiri, S. *et al.* Global, regional and national burden of rheumatoid arthritis 1990–2017: a

systematic analysis of the Global Burden of Disease study 2017. *Ann. Rheum. Dis.* **78**, 1463–1471

(2019).

43.   Lindfors, K. *et al.* Coeliac disease. *Nat. Rev. Dis. Primer* **5**, 3 (2019).

44.   Rubinacci, S., Ribeiro, D. M., Hofmeister, R. J. & Delaneau, O. Efficient phasing and

imputation of low-coverage sequencing data using large reference panels. *Nat. Genet.* **53**, 120–126

(2021).

45.   Meyer, M. & Kircher, M. Illumina Sequencing Library Preparation for Highly Multiplexed

Target Capture and Sequencing. *Cold Spring Harb. Protoc.* **2010**, pdb.prot5448 (2010).

46.   Schubert, M., Lindgreen, S. & Orlando, L. AdapterRemoval v2: rapid adapter trimming,

identification, and read merging. *BMC Res. Notes* **9**, 88 (2016).

47.   Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler transform.

Bioinformatics **25**, 1754–1760 (2009).

48.   Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–

2079 (2009).

49.   Jónsson, H., Ginolhac, A., Schubert, M., Johnson, P. L. F. & Orlando, L. mapDamage2.0: fast

approximate Bayesian estimates of ancient DNA damage parameters. *Bioinformatics* **29**, 1682–

1684 (2013).

50.   Fu, Q. *et al.* A Revised Timescale for Human Evolution Based on Ancient Mitochondrial

Genomes. *Curr. Biol.* **23**, 553–559 (2013).

51.   Korneliussen, T. S., Albrechtsen, A. & Nielsen, R. ANGSD: Analysis of Next Generation

Sequencing Data. *BMC Bioinformatics* **15**, 356 (2014).

52.   Monroy Kuhn, J. M., Jakobsson, M. & Günther, T. Estimating genetic kin relationships in

1342   prehistoric populations. *PLOS ONE* **13**, e0195491 (2018).

1343 53. Weissensteiner, H. *et al.* HaploGrep 2: mitochondrial haplogroup classification in the era of

1344   high-throughput sequencing. *Nucleic Acids Res.* **44**, W58–W63 (2016).

1345 54. Scorrano, G., Yediay, F. E., Pinotti, T., Feizabadifarahani, M. & Kristiansen, K. The genetic

1346   and cultural impact of the Steppe migration into Europe. *Ann. Hum. Biol.* **48**, 223–233 (2021).

1347 55. Purcell, S. *et al.* PLINK: A Tool Set for Whole-Genome Association and Population-Based

1348   Linkage Analyses. *Am. J. Hum. Genet.* **81**, 559–575 (2007).

1349 56. Shringarpure, S. S., Bustamante, C. D., Lange, K. & Alexander, D. H. Efficient analysis of

1350   large datasets and sex bias with ADMIXTURE. *BMC Bioinformatics* **17**, 218 (2016).

1351 57. Patterson, N. *et al.* Ancient Admixture in Human History. *Genetics* **192**, 1065–1093 (2012).

1352 58. Lawson, D. J., Hellenthal, G., Myers, S. & Falush, D. Inference of Population Structure using

1353   Dense Haplotype Data. *PLoS Genet.* **8**, e1002453 (2012).

1354 59. Margaryan, A. *et al.* Population genomics of the Viking world. *Nature* **585**, 390–396 (2020).

1355 60. Hellenthal, G. *et al.* A Genetic Atlas of Human Admixture History. *Science* **343**, 747–751

1356   (2014).

1357 61. The 1000 Genomes Project Consortium *et al.* A global reference for human genetic variation.

1358   *Nature* **526**, 68–74 (2015).

1359 62. Myers, T. A., Chanock, S. J. & Machiela, M. J. LDlinkR: An R Package for Rapidly

1360   Calculating Linkage Disequilibrium Statistics in Diverse Populations. *Front. Genet.* **11**, 157

1361   (2020).

1362 63. Ishigaki, K. *et al.* *Trans-ancestry genome-wide association study identifies novel genetic*

1363   *mechanisms in rheumatoid arthritis*. http://medrxiv.org/lookup/doi/10.1101/2021.12.01.21267132

1364   (2021) doi:10.1101/2021.12.01.21267132.

1365 64. Alekseyenko, A. V. *et al.* Causal graph-based analysis of genome-wide association data in

1366   rheumatoid arthritis. *Biol. Direct* **6**, 25 (2011).

1367 65. Raychaudhuri, S. *et al.* Five amino acids in three HLA proteins explain most of the

1368   association between MHC and seropositive rheumatoid arthritis. *Nat. Genet.* **44**, 291–296 (2012).

1369 66. Spanish Consortium on the Genetics of Coeliac Disease (CEGEC) *et al.* Dense genotyping

identifies and localizes multiple common and rare variant association signals in celiac disease. *Nat.*

Genet. **43**, 1193–1201 (2011).

67.   Monsuur, A. J. *et al.* Effective Detection of Human Leukocyte Antigen Risk Alleles in Celiac

Disease Using Tag Single Nucleotide Polymorphisms. *PLoS ONE* **3**, e2270 (2008).

68.   Gutierrez-Achury, J. *et al.* Fine mapping in the MHC region accounts for 18% additional

genetic risk for celiac disease. *Nat. Genet.* **47**, 577–578 (2015).

69.   the RACI consortium *et al.* Genetics of rheumatoid arthritis contributes to biology and drug

discovery. *Nature* **506**, 376–381 (2014).

70.   Chang, C. C. *et al.* Second-generation PLINK: rising to the challenge of larger and richer

datasets. *GigaScience* **4**, 7 (2015).

71.   Ju, D. & Mathieson, I. The evolution of skin pigmentation-associated variation in West

Eurasia. *Proc. Natl. Acad. Sci.* **118**, e2009227118 (2021).

72.   Kolberg, L., Raudvere, U., Kuzmin, I., Vilo, J. & Peterson, H. gprofiler2 -- an R package for

gene list functional enrichment analysis and namespace conversion toolset g:Profiler.

F1000Research **9**, ELIXIR-709 (2020).

73.   Thorndike, R. L. Who belongs in the family? *Psychometrika* **18**, 267–276 (1953).

74.   Berg, J. J. & Coop, G. A Population Genetic Signal of Polygenic Adaptation. *PLoS Genet.* **10**,

e1004412 (2014).

75.   Frangos, C. C. & Schucany, W. R. Jackknife estimation of the bootstrap acceleration

constant. *Comput. Stat. Data Anal.* **9**, 271–281 (1990).

76.   Sarmanova, A., Morris, T. & Lawson, D. J. *Population stratification in GWAS meta-analysis*

should be standardized to the best available reference datasets.

http://biorxiv.org/lookup/doi/10.1101/2020.09.03.281568 (2020) doi:10.1101/2020.09.03.281568.

77.   McFadden, D. Conditional logit analysis of qualitative choice behavior. (1973).

78.   Efron, B. Bootstrap Methods: Another Look at the Jackknife. *Ann. Stat.* **7**, 1–26 (1979).

79.   Kass, R. E. & Wasserman, L. A Reference Bayesian Test for Nested Hypotheses and its

Relationship to the Schwarz Criterion. *J. Am. Stat. Assoc.* **90**, 928–934 (1995).

80.   Slatkin, M. Linkage disequilibrium — understanding the evolutionary past and mapping the

medical future. *Nat. Rev. Genet.* **9**, 477–485 (2008).

1399