1

Reduced mitochondrial transcription sensitizes acute myeloid leukemia cells to BCL-2 inhibition

Short title: Inhibition of mtDNA expression to treat leukemia

Authors

Laleh S. Arabanian[1], Jenni Adamsson[2], Anke Unger[4], Raffaella Di Lucrezia[4], Tim Bergbrede[4], Arghavan Ashouri[1], Erik Larsson[1], Peter Nussbaumer[4], Bert M. Klebl[4,5], Lars Palmqvist[2,3,*], Claes M. Gustafsson[1,3*,†]

[*]Corresponding authors: lars.palmqvist@clinchem.gu.se and claes.gustafsson@medkem.gu.se

[†]Lead Contact

Affiliations

[1] Department of Medical Biochemistry and Cell Biology, University of Gothenburg, Gothenburg, Sweden

[2] Department of Laboratory Medicine, University of Gothenburg, Gothenburg, Sweden.

[3] Department of Clinical Chemistry, Sahlgrenska University Hospital, Gothenburg, Sweden

[4] Lead Discovery Center GmbH, Otto-Hahn-Str. 15, D-44227 Dortmund, Germany

[5] The Norwegian College of Fishery Science, UiT The Arctic University of Norway, PO Box 6050 Langnes, N-9037 Tromsø, Norway

Competing interests:

CMG is a scientific co-founder of Pretzel Therapeutics Inc. Together with RD, TB, AU, PN, BMK who are employees of LDC, CMG is also co-inventor of the patent application WO 2019/057821. Medicinal chemistry and pharmacology part of this work was financed by the Max-Planck Gesellschaft e.V. under the framework agreement between Max-Planck and LDC.

Abstract

Overcoming drug-resistance and the subsequent relapse that often occurs with monotherapy

is crucial in the treatment of acute myeloid leukemia. We here demonstrate that therapy-

resistant leukemia initiating cells can be targeted using a novel inhibitor of mitochondrial

transcription (IMT). The compound inhibits mitochondrial RNA polymerase activity and

sensitizes the resistant population to the induction of apoptosis. *In vitro* studies on acute

myeloid leukemia cells demonstrate that IMT prevents cell proliferation, and together with a

selective BCL-2 inhibitor, venetoclax, induces apoptosis and suppress oxidative

phosphorylation (OXPHOS) synergistically. AML mouse models treated with IMT in

combination with venetoclax show prolonged survival in venetoclax-resistant models. Our

findings suggest that certain therapy-resistant leukemia cell populations display a unique

dependency on mitochondrial transcription and can be targeted with IMT.

Introduction

Acute myeloid leukemia (AML) comprises a devastating and heterogeneous group of diseases, for which patient survival rates are dismal despite several new treatment options. Current cares are still mainly based on unselective chemotherapy but for most patients also with addition of more directed treatments after patient stratification based on genetic aberrations found at diagnosis. However, these strategies lead to cures in only a fraction of AML patients due to therapy resistance developed by the leukemic cells leading to relapse. Curative treatment usually needs allogenic stem cell transplantation, which is not applicable in many cases, highlighting the critical need to develop different approaches. The anti-apoptotic molecule BCL-2 has been a therapeutic target in lymphoid and myeloid leukemia, as in many cancers overexpression of BCL-2 results in dysregulation of apoptosis (1). ABT-199 (venetoclax), is a specific BH3-mimetic that causes apoptosis by preventing BCL-2 from binding pro-apoptotic BAX and BAK1 proteins (2). However, complete responses occur in just ~20% of AML patients treated with Venetoclax (3). In addition, many patients relapse, with regrowth initiated by therapy-resistant leukemic clones, suggesting that single agent administration is unlikely to yield lasting responses in most cases. Venetoclax is therefore used in combination with conventional therapies. Together with hypomethylating agents, Venetoclax is now standard of care for patients deemed unfit for intensive induction therapy (4).

Resistance to venetoclax may result from overexpression of the anti-apoptotic proteins MCL1 or BCL2L1 (5), or through changes in levels of other anti-apoptotic molecules (6). Aberrant pro-survival regulation can also be achieved by inactivation of TP53 function, which controls many aspects of apoptosis (6, 7). MCL1-dependent resistance to venetoclax can be overcome when the compound is combined with the MDM2 inhibitor RG7388, which causes p53

activation. A combination of BCL-2 inhibition and TP53 activation may thus provide an

interesting therapeutic approach for treatment of AML (8).

Other combination approaches that promote apoptosis may also be of interest. The majority

of ATP required for leukemia stem cell (LSC) function is generated by oxidative

phosphorylation (OXPHOS) (9, 10). Interestingly, venetoclax-resistant leukemic cells display

increased levels of OXPHOS and are protected from mitochondrial stress (7). These data

suggest approaches to specifically target OXPHOS in leukemia may be an attractive option.

In support of this notion, inhibition of mitochondrial protein translation by antibiotics that

reduce OXPHOS capacity results in a selective cytotoxic effect on leukemic cell when used

in combination with tyrosine kinase inhibitors (11). Furthermore, LSCs from *de novo* AML

patients display a unique dependency on amino acid metabolism to fuel OXPHOS. When

venetoclax is combined with the hypomethylating agent azacytidine, the two compounds act

in concert to inhibit amino acid uptake, leading to a drop in OXPHOS and thus cell death

(12). More recently, AML therapy using a combination of cytarabine (AraC) and venetoclax

has highlighted the central role of mitochondrial adaptation during treatment (10, 13),

indicating the need for a mitochondrial directed combination therapy for AML. Such a

therapy may help to overcome existing treatment resistance.

Recently we developed novel, first-in-class inhibitors of mitochondrial transcription (IMTs)

(14). IMT treatment specifically targets mitochondrial RNA polymerase (POLRMT) and

reduce transcription of mitochondrial DNA, which codes for thirteen key components of the

OXPHOS machinery, including subunits of respiratory complexes I, III, IV, and the ATP

synthase. Consequently, IMT treatment impairs OXPHOS system biogenesis in a dose-

dependent manner and decreases cell viability in various tumor cells. Surprisingly, several

weeks of per oral IMT treatment is well tolerated in mice and does not cause toxicity in

normal tissues, despite inducing a strong anti-tumor response in human cancer xenografts

(14).

Here we examine AML cell lines for their sensitivity for treatment with IMT. We find that a

combination of IMT and venetoclax displays a synergistic on cell growth and induction of

apoptosis. We demonstrated the efficacy of this approach in leukemia mouse models

generated by human AML cell lines as well as primary samples from AML patients.

Results

Synergistic effect of IMT in combination with venetoclax on AML cell lines

We were curious whether our inhibitors of mitochondrial transcription (IMTs) (14) could be effective against AML cells. The coumarin class of IMTs was optimized by medicinal chemistry to generate lead IMT compounds with improved metabolic stability in mice, such as LDC204857, herein simply referred to as IMT (Fig 1A, Fig. S1 and Table S1). IMT is a close analog of previously described IMT1B (Fig. S1) with a highly similar biochemical and cellular inhibition profile (14). It shows an extended half-life *in vivo* due to its increased metabolic stability (Table S1). Thus, by using IMT we could consistently reduce the mouse dose to 50 mg/kg per oral once daily, which is very well tolerated and does not exert any adverse effects in mice.

Using four AML cell lines with different expression of TP53 (MV4-11 and MOLM-13 expressing wild type TP53; THP-1 and KG-1 expressing mutated TP53), we monitored the effects of increasing concentrations of IMT on cellular growth using a luminescence-based cell viability assay. Only marginal effects were observed after a 72-h period of exposure to IMT, but all cell lines displayed dose-dependent antileukemic activity after 144 h of treatment (Fig. 1B), in line with our previous observations (14). We also monitored venetoclax sensitivity by assessing cell viability, and noted that, in line with published work (15), a dose-dependent reduction in cell growth occurs after three days and becomes more pronounced after 6 days (Fig. 1B).

To then investigate whether IMTs show a synergistic effect when combined with venetoclax, we conducted a combination study in AML cell lines (Fig. 1C). After six days of treatment, we found a clear synergistic effect on cell proliferation in MV4-11 and MOLM-13 cell lines,

which express wild type TP53, as determined by combination analysis (Bliss-independence

model (16)). Treatment of THP-1 and KG-1 cell lines, which harbor mutated TP53, does not

show a synergistic effect even when using higher concentrations of both IMT and venetoclax

(Fig. 1C, additional treatment data are shown in Fig. S2, Bliss value calculations for both

time points are summarized Fig. 1 - Source Data 1.

Combination of IMT and venetoclax synergistically induces apoptosis and loss of

mitochondrial membrane potential in MV4-11 and MOLM-13 cells.

To investigate the factors involved in the inhibition of cell growth with IMT and venetoclax,

AML cell lines were treated with single agents (IMT or venetoclax) or a combination of the

two (Fig. 2A), and the stages of apoptosis were monitored using Annexin V and PI staining.

No detectable increase in apoptosis was induced by IMT exposure alone in any AML cell

line. Venetoclax treatment resulted in both early and late apoptosis induction in MV4-11 and

MOLM-13 cell lines, and employing a combination of IMT and venetoclax significantly

enhanced apoptosis induction in these wild-type cell lines. On the other hand, combination

treatment did not induce apoptosis in THP-1 and KG-1 mutated cell lines (Fig. 2A).

One distinctive feature of apoptosis is the disruption of mitochondrial function, which can be

measured as loss of mitochondrial membrane potential due to membrane depolarization. As a

combination of IMT and venetoclax could synergistically induce apoptosis, we were curious

whether combination drug treatment thus affects mitochondrial membrane potential. We

observed that venetoclax-sensitive cell lines (MV4-11 and MOLM-13) showed a dramatic

loss of mitochondrial membrane potential when treated with both venetoclax and IMT (Fig.

2B). On the other hand, mitochondrial membrane potential in THP-1 cells was not affected

by single nor by combination treatment. Interestingly, although KG-1 was venetoclax

insensitive, the combination treatment significantly enhanced loss of mitochondrial

membrane potential (Fig. 2B). These data suggest that a combination of IMT and venetoclax

is cytotoxic and synergistically induces cell death in sensitive cells, presumably through

disruption of mitochondrial function.

159 Combination of IMT and venetoclax significantly suppresses OXPHOS in AML cells.

We next studied whether combination treatment with IMT and venetoclax affects OXPHOS

in AML cells. OXPHOS was evaluated by determining oxygen consumption rate (OCR) (Fig.

3A), which was measured first as basal respiration and then oligomycin (OMY)-induced leak

respiration (ATP-linked respiration) after the addition of oligomycin to inhibit ATP synthase.

Stepwise carbonyl cyanide m-chlorophenyl hydrazone (CCCP) titrations were subsequently

carried out to reach the electron transport chain respiratory capacity state. Finally, antimycin

a (AMA) was added to inhibit complex I/III of the electron transport chain and to terminate

respiration, allowing for the determination and correction of residual oxygen consumption,

which is indicative of non-mitochondrial oxygen consumption (Fig. 3A). We noted a

tendency to inhibition of OXPHOS respiration in venetoclax-sensitive MOLM-13 and MV4-

11 cells when combination treatment was used (Fig. 3B). In comparison, OXPHOS levels in

venetoclax-resistant THP-1 cells were unaffected in the presence of single compound or

combination treatments, confirming that the inhibition we observed is not a general effect of

drug treatment (Fig. 3B). In addition, there was no conclusive OCR data obtained for KG-1

cell line.

In normal hematopoietic cells, glycolysis is upregulated when OXPHOS is blocked (17). In

contrast to normal hematopoietic cells, high energy turnover AML cells rely predominantly

on OXPHOS due to poorly regulated metabolic adaptation. To verify the reliance on

OXPHOS, AML cells were grown in growth medium supplemented with either glucose or

galactose (Fig. 3C). Two ATP molecules are produced when cells use glucose for glycolysis,

but if galactose is used instead, no ATP is generated (18). We observed no significant

decrease in the level of ATP production (Fig. 3C) nor in cell counts (Fig. S3A) in the absence

of glucose, indicating that glycolysis is not a major source of energy in AML cells. To test

whether AML cells use glycolysis under treatment conditions, OCR was measured in the

presence or absence of a high concentration of glucose (Fig. S3B). OCR measured for basal

respiration was not significantly different with excess amounts of glucose, again indicating

the inability of AML cells to switch to glycolysis when OXPHOS is disrupted.

To assess whether AML cells could rely on glycolysis for ATP generation when OxPhos was

inhibited by IMT, we cultured AML cells in medium with galactose instead of glucose, a

condition that limits glycolysis-dependent ATP production. Even in these glycolytic

conditions, we observed minimal ATP production and cell viability, suggesting that AML

cells are poorly adapted to switch to glycolysis when mitochondrial function is impaired. This

finding underscores the high dependence of AML cells on OxPhos for energy production and

supports the proposed mechanism of IMT efficacy, as it appears to exploit the inability of

AML cells to compensate through glycolysis when mitochondrial transcription is inhibited.

Combination of IMT and venetoclax synergistically delays leukemia initiation in cell-

derived leukemia mouse models.

Following the confirmed synergy between IMT and venetoclax in AML cell lines, we

proceeded to test the dual inhibition of mitochondrial transcription and BCL-2 in an AML

cell-derived xenograft (CDX) transplantation mouse model. First, the pharmacokinetic profile

of IMT was determined in plasma, liver, and bone marrow (BM) of mice, 1, 8, and 24 h post-

administration (Fig. S4A). As described above (Fig. S1), the IMT mouse PK profile was superior to IMT1B. IMT was detected in all tissues at all time points when administered alone. To determine the concentration of IMT and venetoclax when administered in combination, plasma samples were then collected 8–11 h post-administration of single or combination formulations and the concentrations of IMT and/or venetoclax in samples were measured (Fig. S4B). Values obtained were all within the effective range, as combination treatment did not affect the individual compound concentrations (14). For xenograft experiments, NBSGW mice were intravenously transplanted with MV4-11 AML cells (Fig. 4A). Two weeks after transplantation, engraftment of AML cells was confirmed (evaluated by gating on hCD45+ population by flow cytometry in peripheral blood). Treatment then began with vehicle, 50 mg/kg IMT, 100 mg/kg venetoclax, or a combination of IMT (50 mg/kg) and venetoclax (100 mg/kg) (Fig. 4A), dosed as such based on previously published protocols (14, 18). The efficacy of treatments was determined by measuring the proportion of transplanted AML cells in peripheral blood and by survival analysis (Fig. 4B). In all treatment groups, the hCD45+ population dropped immediately after treatment, and vehicle- and venetoclax-treated mice then exhibited a rapid rebound. In contrast, in IMT-treated mice the increase in leukemic cell proportion was delayed until eight weeks post-transplantation, and in the combination-treated group for to up to 20 weeks (Fig. 4B). These observations correlate with results from survival analysis (Fig. 4C). IMT treatment of MV4-11 transplanted mice significantly delayed the development of leukemia, inducing a 37% increase in median overall survival (57.5 → 79 days). Development of leukemia was further delayed by combination treatment to 131% (57.5 → 133 days), and the group contained two mice that did not show any disease symptoms after 200 days (post-transplantation) when they had to be euthanized according to ethical permissions. Moreover, leukemic cells were not detectable in the BM of surviving mice at the time of termination. All other mice showed

symptoms of leukemia and of these, all but one combination-treated mouse had an increased

percentage of leukemic cells in BM (Fig. 4D). These results confirm that treatment with a

combination of IMT and venetoclax significantly prolongs leukemia progression in mice.

Moreover, the general condition of mice such as behavior, fur, mobility and weight, assessed

before (week 2) and after treatment (week 5), were not altered in the IMT-, venetoclax-, or

combination-treatment groups when compared to the vehicle-treated group (weights shown in

Fig. S5B), suggesting that the treatments were well tolerated. As with single IMT therapy

(14), treatment with combination therapy did not display any evidence of organ toxicity or

drug-associated death in the mice (Fig. S5C, D).

IMT and venetoclax synergistically decrease the tumor burden and increase the life

span in an AML-PDX mouse model resistant to venetoclax.

The CDX mouse model demonstrated IMT could sensitize AML cell lines to BCL-2

inhibitors *in vivo*. We next analyzed the effects of combination treatment on a more refined *in*

vivo model by generating xenografts using AML patient-derived cells (AML PDX). We first

tested an AML-PDX mouse model with resistance to venetoclax (AML sample harboring

TP53 mutation). Two weeks post-transplantation with AML PDX cells, NSGS mice were

orally treated with IMT, venetoclax, or a combination of the two for 21 days (Fig. 5A). The

efficacy of the treatments was determined based on the proportion of human AML cells

(determined by hCD45+ population). Six out of 10 mice in each group were euthanized

immediately after the last compound administration for end-of-treatment analysis. We found

that the combination of IMT and venetoclax resulted in significant inhibition of tumor burden

in blood, BM, and spleen of treated mice (Fig. 5B). By examining histology sections of BM

and spleen, we found that tissues from combination-treated animals contained less

engraftment of leukemic blasts than single- and vehicle-treated mice (Fig. 5C). Accordingly,

the frequencies of hCD45+ cells in the BM and spleens of the combination-treated group

were largely reduced in comparison to other groups (Fig. 5D), and engrafted tumor cells in

combination-treated PDX mice showed a lower proliferating percentage in comparison to

single or vehicle treated groups (shown by KI67+/hCD45+ immunohistochemistry staining;

Fig. 5E). This demonstrates a synergy between IMT and venetoclax in reducing tumor

burden.

To further analyze the cellular effects of the IMT-venetoclax combination treatment, we

analyzed transcriptomes of purified viable AML blasts isolated from PDX models at end of

treatment (Fig. 6A). We confirmed that combination treatment had a more pronounced effect

on gene transcription as compared to single treatments. In agreement with its more

pronounced effect on tumor burden, combination treatment suppressed expression of several

genes involved in leukemogenesis, including MX1, KCTD12, ITGA and RGL1 (Table S2)

(19). Functional annotation and enrichment analysis were performed using the DAVID

bioinformatics resources (Database for Annotation, Visualization, and Integrated Discovery)

to identify significantly enriched gene ontology (GO) terms and pathways associated with the

gene set. The gene list in Table S2 points to a potentially integrated function of the gene set

in 1- Innate immune involvement, 2- Mitochondrial Function and Immune Signaling and 3-

Signaling and Trafficking. The presence of annotations related to innate immune response,

extracellular region, and transmembrane receptors suggests that the genes may be involved in

the initial immune defense, recognizing pathogens, and activating immune cells. The

indication of mitochondrial function and immune signaling approves that mitochondria are

known not only for energy production but also for their emerging role in immune signaling,

such as by producing reactive oxygen species (ROS) in response to infection. For signaling

and trafficking, the presence of terms like "GTP binding," "endosome," and "calcium"

highlights intracellular signaling, possibly in relation to immune activation or cellular responses The presence of terms like "GTP binding," "endosome," and "calcium" highlights intracellular signaling, possibly in relation to immune activation or cellular responses. We also analyzed survival by monitoring the mice for disease progression and morbidity. Peripheral blood engraftments were analyzed at regular intervals until the appearance of disease symptoms (Fig. 6B and Fig. S6A). The leukemia burden immediately prior to euthanization demonstrated that the cytoreductive effect of the combination treatment persists (Fig. 6B). However, blood count (determined by RBCs and HGB levels) did not vary significantly between experimental groups, nor did the magnitude of splenomegaly or liver weight vary at the time of euthanization (Fig. S6B–D). Interestingly, mice treated with a combination of IMT and venetoclax succumbed to death significantly later than other treatment groups, with a median survival of 112 days vs 96, 99, and 102 days for vehicle-, IMT-, and venetoclax-treated groups, respectively (Fig. 6**C**). Combination treatment thus increases the life span of venetoclax-resistant AML mouse models, demonstrating that IMT can sensitize AML cells towards other therapies.

IMT does not affect tumor burden or life span in an AML-PDX mouse model sensitive to venetoclax.

For comparison, we also monitored effects of combination treatment on a PDX model based on AML cells sensitive to venetoclax treatment (AML cells harboring FLT3-ITD, NMP1 and DNMT3A). As expected, we observed that venetoclax treatment significantly reduced the tumor burden in the blood and BM of mice. Combination treatment with IMT did not add to this effect (Fig. 7A). By examining histology sections, we found that both BM and spleen tissues from combination- and venetoclax-treated animals contained reduced proportion of leukemic blasts compared to IMT- and vehicle-treated mice (Fig. 7B). White blood counts

(WBCs), red blood counts (RBCs), and hemoglobin (HGB) levels were not significantly

different across experimental groups (Fig. S7A). The total body and liver weight of PDX

mice were very similar in all groups (Fig. S7B, D), but both venetoclax and combination

treatments reduced splenomegaly (Fig. S7C).

The remaining mice were used for survival analysis, which involved regular blood count and

chimerism analysis until the appearance of disease symptoms (Fig. 7C). Both venetoclax and

IMT caused a modest increase in life span, but combination treatment did not offer an

additional effect (Fig. 7D).

Discussion

In recent years, the genetic complexity and heterogeneity of leukemia cell populations has become apparent (20, 21) and developing effective strategies to target human leukemia initiating cells has thus proven to be a challenging task. Currently, there is considerable interest in therapies, such as venetoclax, that can restore normal apoptosis of cancer cells by directly interacting with the BCL-2 family of proteins. These are effective for some leukemias, but many patients eventually become refractory to treatment and ultimately succumb to their disease (22, 23). It has become increasingly important to determine which biological properties are relatively consistent in leukemic cell populations irrespective of genetic makeup (tumor population). Recent observations indicate that several cancer cell types, including AML, are reliant on OXPHOS for metabolism (24), making inhibition of mitochondrial function a promising therapeutic strategy for AML (12, 25-28). Therapy-resistant leukemic cells are more reliant on oxidative respiration than glycolysis for energy generation, suggesting that the OXPHOS dependency and lack of normal glycolytic activity in AML cells may be characteristic of cancers (29). Understanding resistance mechanisms will allow for the design of strategies to counteract drug resistance or impede resistance from developing. In AMLs with aberrant expression of BCL-2, venetoclax can reduce OXPHOS (9) and trigger cell death. However, venetoclax resistance can occur in patients treated with the BCL-2 inhibitor and in some subtypes of AML patients with poor prognoses (30).

Previous work has indicated that AML cells display increased mitochondrial biogenesis and inhibition of mitochondrial translation can be used as a strategy for human AML (18, 27). In our work, we target another process involved in mitochondrial gene expression, i.e., transcription. IMT treatment targets POLRMT, which has two main functions in mitochondria: it transcribes mtDNA encoded genes and provides the primers required to

initiate mtDNA replication. Inhibition of POLRMT can thus deliver a double punch, both

reducing mtDNA copy number and inhibit OXPHOS biogenesis. As demonstrated here

inhibition of POLRMT can overcome venetoclax therapy resistance in AML. These

observations are in line with recent reports of the importance of mitochondrial function in

leukemia cell survival (31, 32). By developing a clinically relevant and well-tolerated

approach to deliver IMTs in immunodeficient mice transplanted with AML cells, we were

able to reduce the tumor burden in all (cell- and patient-derived) xenografted mice treated

with a combination of IMT and venetoclax. This demonstrates that reducing POLRMT

activity results in successful targeting of leukemic cells. Our *in vitro* investigations confirmed

that the combination of IMT and venetoclax is necessary to significantly decrease ATP levels

and consequently induce apoptosis. In addition, efficacy studies of AML PDX models

showed that single compounds are not sufficient to eradicate AML cells, but that the

combination regimen reduced AML tumor burden and prolonged survival in all treated mice.

These data indicate the advantage of both downregulating POLRMT activity in combination

with increasing apoptosis and highlights the importance of rational combination therapies

(33, 34). IMT offers potential advantages over other OXPHOS-targeting therapies due to its

lower toxicity, as observed in preclinical models. While other strategies, such as

mitochondrial translation inhibition or electron transport chain inhibitors, effectively target

OXPHOS in AML, they may come with higher toxicity risks. IMT's selective targeting of

mitochondrial transcription appears to minimize adverse effects, making it a promising

candidate for use in combination regimens with BCL-2 inhibitors like venetoclax. This

approach could improve therapeutic outcomes while maintaining tolerability, addressing a

significant challenge in AML treatment.

In our study, we observed that the combination of IMT and venetoclax did not show

synergistic effects in TP53 mutant AML cells, particularly in the THP-1 cell line. Although

statistical analyses indicated potential interaction, the biological impact was minimal in TP53

mutant cells, contrasting with the robust synergy observed in TP53 wild-type cells like MV4-

11 and MOLM-13. This suggests that TP53 mutation status may influence the efficacy of this

combination therapy, highlighting a potential limitation in targeting mitochondrial

transcription to potentiate BCL-2 inhibition in TP53 mutant AML context. These findings

underscore the need for broad mutational profiling when designing combination therapies for

AML.

In summary, we have introduced a new therapeutic strategy to target therapy-resistant

leukemia with OXPHOS-dependent properties. Despite several limitations with the design of

pre-clinical cancer models and the intensity and duration of treatments, we demonstrate here

that the combined inhibition of BCL-2 and mitochondrial transcription can target therapy-

resistant AMLs. This combination treatment succeeds as IMT primes cell vulnerability to

apoptosis induction. Consequently, our findings may be of value for the future development

of better clinical strategies for the treatment of AML.

Acknowledgments

The authors would like to thank the staff at the Department of Experimental Biomedicine at

the University of Gothenburg, namely, Jenny Nyman, Claudia Engwall, Tova Johansson and

Hanna Brissman for tremendous assistance with mouse experiments. They would also like to

thank Dr Rebekah Hailes for editing the manuscript and to acknowledge Lenka Palova-

Jelinkova, Klara Danova, and Pavla Jestrabova from SOTIO Biotech (Prague, Czech

Republic) for valuable input on *in vitro* and *in vivo* studies. The authors acknowledge support

from the National Genomics Infrastructure in Stockholm funded by Science for Life

Laboratory, the Knut and Alice Wallenberg Foundation and the Swedish Research Council,

and SNIC/Uppsala Multidisciplinary Center for Advanced Computational Science for

assistance with massively parallel sequencing and access to the UPPMAX computational

infrastructure. This research is supported by Swedish Research Council (2017-01257:CMG),

Swedish Cancer Foundation (20 1291 PjF:CMG, and 20 0925 PjF:LP), Knut and Alice

Wallenberg Foundation (2017.0080: CMG), The Swedish state under the agreement between

the Swedish government and the county councils, the ALF agreement (ALFGBG-

966275:CMG and ALFGBG-965206: LP), and Assar Gabrielsson Foundation for cancer

research grant (LSA)

Materials and Methods

Cell lines and culture conditions

All cell lines (MV4-11, MOLM13, THP-1, KG-1, OCI-AML3, and OCI-AML2) were

purchased from DSMZ (Braunschweig, Germany) and cultured in RPMI-1640 (Sigma

Aldrich) supplemented with 10–20% heat inactivated fetal bovine serum (Gibco) and 1%

penicillin/streptomycin (Invitrogen). Cells were maintained in a humidified incubator at 37

°C and 5% CO_2. RPMI-1640 without glucose was used for *in vitro* treatment experiments and

supplemented by physiological concentrations (5 mM) of glucose (D-Glucose, Sigma) or

galactose (D-galactose, Sigma).

Generation of *in vivo* AML mouse model

Mice were purchased from Jackson laboratories (USA) and maintained in a pathogen-free facility at the University of Gothenburg. All animal experiments were conducted in accordance with guidelines approved by the research animal ethics committee at the University of Gothenburg. For AML cell line xenograft (CDX) experiments, 6–8-week-old, male NBSGW mice (NOD.Cg-Kit^{W-41J} Tyr$^+$ Prkdc^{scid} Il2rg^{tm1Wjl}/ThomJ, Jackson laboratories) were used. Mice were transplanted intravenously with ~1 × 10^6 MV4-11 cells. For patient-derived xenograft (PDX) experiments, 6–8-week-old, male NSGS mice (NOD.Cg-Prkdc^{scid} Il2rg^{tm1Wjl} Tg (CMV-IL3,CSF2,KITLG)1Eav/MloySzJ, Jackson laboratories) were sub-lethally irradiated with 1 Gy radiation. AML PDX splenocytes were purchased from Jackson laboratories (id: J000106134*, male, 58 years old, AML, M4, positive for FLT3-ITD, DNMT3A, and NPM1mutaions; and id: 000106566*, male, 48 years old, AML no classification specified, positive for TP53 mutation). To obtain adequate number of PDX cells for the scale of the efficacy experiments, the splenocytes were initially expanded in vivo. Mice were transplanted intravenously with ~1 × 10^6 splenocytes from AML PDX mice containing more than 80% patient AML cells.

Human leukemia cell frequency assessment by flow cytometry

The percentage of human leukemic cells (MV4-11 cell lines and AML PDX cells) in mouse peripheral blood, BM, and spleen was determined by flow cytometry. Mice were bled at indicated time points after transplantation. Prior to staining, peripheral red blood cells were lysed using ammonium chloride for 8–10 min at room temperature. Cells were stained with antibodies, washed, and submitted for flow-cytometric analysis using a FACS Aria (BD Biosciences, USA) or CytoFlex instrument (Beckman Coulter, California, USA). At least 10,000 events were acquired, and the percentage was calculated relative to gated viable mononuclear cells (MNCs). To discriminate between human and mouse cells, cells were

stained with antibodies against human CD45 (clone HI30, BD Biosciences) and mouse CD45

(clone 2D1, BD Biosciences). Engraftment was deemed positive if human CD45+ cells

represented >0.1% of the viable MNCs. 7AAD (BD Biosciences) served as viability marker.

Preparation of compounds for *in vivo* drug administration

IMT was optimized based on the structure activity and property relationship of the coumarin

mtRNAPol inhibitor class by LDC (Dortmund, Germany). Venetoclax was obtained from

MedChemExpress (Hölzel Diagnostika Handels GmbH, Germany). Following the

development of a suitable vehicle for the formulation of IMT and venetoclax, as well as the

combination of both compounds, the test articles were formulated using 0.5% aqueous

methylcellulose solution (Sigma-Aldrich, Germany). After randomization, and two weeks

post-transplantation or with confirmation of engraftment, mice were treated with vehicle

(0.5% methylcellulose), 50 mg/kg IMT, 100 mg/kg venetoclax, or a combination (100 mg/kg

venetoclax + 50 mg/kg IMT) by oral administration daily for 21 days.

Cell proliferation assay

Cells were seeded out ca 20 h prior to treatment. For dose response experiments, cells were

treated with a wide range of concentrations of single compounds. Cell proliferation assays

were performed every 24 h from 72–168 h, where cell viability/ATP production was

determined using CellTiter-Glo (Promega).

Apoptosis assay and Mitochondrial potential assay

Apoptosis was evaluated by Annexin V and PI staining and mitochondrial membrane

potential was evaluated by lipophilic cationic dye JC-1. Cells were treated with IC50

concentrations of IMT, venetoclax, or combination of both, analyzed using an advanced

image cytometer (Nucleocounter NucleoCounter's accompanying software NucleoView).

High-resolution respirometry

AML cell lines were treated with DMSO, IMT, venetoclax, or combination of IMT and

venetoclax for 3–4 days. The mitochondrial respiratory capacity (oxygen consumption rate)

was measured using high resolution respirometry (O2K, Oroboros, Innsbruck, Austria). A

substrate-uncoupler-inhibitor titration (SUIT) protocol was used to measure specific features

of mitochondrial respiration. Respiratory capacities were tested in a sequence of coupling

states: basal respiration; LEAK respiration using 5 mM oligomycin (Sigma); and non-

coupled electron transfer state (ET state) where the optimum carbonyl cyanide 3-

chlorophenylhydrazone (CCCP, Sigma) concentration for maximum flux was determined by

titration (max. 3 µM CCCP). Mitochondrial respiration was inhibited using 2.5 µM antimycin

(Sigma). Where stated, excess glucose (D-Glucose, Merck) was added at a final

concentration of 20 mM. Instrumental and oxygen background fluxes were calibrated as a

function of oxygen concentration and subtracted from the total volume-specific oxygen flux.

Calibration and data analysis was performed using Datlab v.6.1 software (Oroboros

Instruments).

Pharmacokinetics analysis

484   IMT and venetoclax were extracted from plasma and tissues by protein precipitation using

485   acetonitrile. Tissue samples were homogenized using double the (sample) volume of PBS,

486   and bone marrow samples were suspended in 40 µL of PBS prior to extraction with

487   acetonitrile. Samples were analyzed by liquid chromatography tandem-mass spectrometry

488   using a Prominence UFLC system (Shimadzu) coupled to a Qtrap 5500 instrument

(ABSciex). Test articles were separated on a C18 column using a gradient elution with an acetonitrile/water mixture containing 0.1% formic acid as the mobile phase. Chromatographic conditions and mass spectrometry parameters were optimized for each test article prior to sample analysis. Concentrations of each test article were calculated by means of a standard curve.

mRNA extraction and transcriptome sequencing of AML PDX cells

For RNAseq experiment BM samples were FACS sorted using BD FACS Aria fusion (BD Biosciences). Human AML cells were isolated using human CD45 antibody. 7-AAD (BD Biosciences) was used for dead cell exclusion. RNA was extracted using RNeasy Micro kit (Qiagen). RNAseq library preparation and RNA-sequencing was done using the SMARTer Total Stranded RNA pico kit. RNA library was prepared at National Genomics Infrastructure (NGI) to be sequenced on the NovaSeq S6000 on 0.5 lanes of one S4 flowcell.

Bioinformatics analysis

Analysis was performed in R (www.r-project.org, accessed on 10 May 2022) using DESeq2, clusterProfiler and ReactomePA packages. Prefiltering to keep genes that have at least 10 reads total was performed before differential expression analysis was done with DESeq2. Visualization of differentially expressed genes was done with EnhancedVolcano to create volcano plots and gridExtra for creating tables. The *p*-value cutoff was set to 0.1 to allow for more genes to be found in EP (End-of-treatment) group comparisons and the Res/Sen (Resistant/Sensitive) group comparisons. Upregulated and downregulated genes were plotted and colored according to Fold Change (FC) greater than 1.5 or -1.5 and the *p*-value cutoff 0.1.

Functional annotation and enrichment analysis were performed using the DAVID

bioinformatics resources (Database for Annotation, Visualization, and Integrated Discovery)

to identify significantly enriched gene ontology (GO) terms and pathways associated with the

gene set (35).

Statistical Analysis

Unless otherwise noted, data were summarized using mean ± standard error of the mean (±

SEM). Per-group sample sizes presented in figures and results are reported from ≥3 separate

experiments, unless stated otherwise. Survival distributions were estimated using the Kaplan–

Meier method, and group comparisons were conducted using the log-rank test. Data were

analyzed using GraphPad Prism 9.0. Combination analysis was summarized using the Bliss-

independent model (16). The formula $Y_{ab,P} = Y_a + Y_b - Y_a Y_b$ was used where drug A at dose

a inhibits Y_a percent of cell growth, and drug B at dose b inhibits Y_b percent of cell growth.

The predicted inhibitory effect of combined percentage was estimated by $Y_{ab,P}$. The observed

combined percentage inhibition $Y_{ab,O}$ was then compared with $Y_{ab,P}$, and if $Y_{ab,O} > Y_{ab,P}$, the

combination treatment was thought to be more efficacious (synergy). If $Y_{ab,O} = Y_{ab,P}$ the

combination treatment was thought to be additive.

Author contributions:

Conceptualization (LSA, RD, TB, AU, PN, BMK)

Methodology (TB, AU, PN, BMK)

Investigation (LSA, JA, CN)

Supervision (BMK, LP, CG)

Writing—original draft (LSA)

Writing—review & editing (LSA, JA, AU, RD, TB, PN, BMK, LP, CMG)

Competing interests:

CMG is a scientific co-founder of Pretzel Therapeutics Inc. Together with RD, TB, AU, PN,

BMK who are employees of LDC, CMG is also co-inventor of the patent application WO

2019/057821. The Medicinal chemistry and pharmacology part of this work was financed by

the Max-Planck Gesellschaft e.V. under the framework agreement between Max-Planck and

LDC.

Data and materials availability:

All data needed to evaluate the conclusions in the paper are present in the paper and/or the

Supplementary Materials. RNA-seq datasets have been deposited in the GSE repository

(GSE213976). Additional data related to this paper may be requested from the authors.

References

1. Guieze R, Liu VM, Rosebrock D, Jourdain AA, Hernandez-Sanchez M, Martinez Zurita A, et al. Mitochondrial Reprogramming Underlies Resistance to BCL-2 Inhibition in Lymphoid Malignancies. Cancer Cell. 2019;36(4):369-84 e13.

2. Souers AJ, Leverson JD, Boghaert ER, Ackler SL, Catron ND, Chen J, et al. ABT-199, a potent and selective BCL-2 inhibitor, achieves antitumor activity while sparing platelets. Nat Med. 2013;19(2):202-8.

3. Konopleva M, Pollyea DA, Potluri J, Chyla B, Hogdal L, Busman T, et al. Efficacy and Biological Correlates of Response in a Phase II Study of Venetoclax Monotherapy in Patients with Acute Myelogenous Leukemia. Cancer Discov. 2016;6(10):1106-17.

4. Roberts AW, Wei AH, Huang DCS. BCL2 and MCL1 inhibitors for hematologic malignancies. Blood. 2021;138(13):1120-36.

5. Ramsey HE, Fischer MA, Lee T, Gorska AE, Arrate MP, Fuller L, et al. A Novel MCL1 Inhibitor Combined with Venetoclax Rescues Venetoclax-Resistant Acute Myelogenous Leukemia. Cancer Discov. 2018;8(12):1566-81.

6. Choi JH, Bogenberger JM, Tibes R. Targeting Apoptosis in Acute Myeloid Leukemia: Current Status and Future Directions of BCL-2 Inhibition with Venetoclax and Beyond. Target Oncol. 2020;15(2):147-62.

7. Nechiporuk T, Kurtz SE, Nikolova O, Liu T, Jones CL, D'Alessandro A, et al. The TP53 Apoptotic Network Is a Primary Mediator of Resistance to BCL2 Inhibition in AML Cells. Cancer Discov. 2019;9(7):910-25.

8. Pan R, Ruvolo V, Mu H, Leverson JD, Nichols G, Reed JC, et al. Synthetic Lethality of Combined Bcl-2 Inhibition and p53 Activation in AML: Mechanisms and Superior Antileukemic Efficacy. Cancer Cell. 2017;32(6):748-60 e6.

9. Lagadinou ED, Sach A, Callahan K, Rossi RM, Neering SJ, Minhajuddin M, et al. BCL-2 inhibition targets oxidative phosphorylation and selectively eradicates quiescent human leukemia stem cells. Cell Stem Cell. 2013;12(3):329-41.

10. Farge T, Saland E, de Toni F, Aroua N, Hosseini M, Perry R, et al. Chemotherapy-Resistant Human Acute Myeloid Leukemia Cells Are Not Enriched for Leukemic Stem Cells but Require Oxidative Metabolism. Cancer Discov. 2017;7(7):716-35.

11. Kuntz EM, Baquero P, Michie AM, Dunn K, Tardito S, Holyoake TL, et al. Targeting mitochondrial oxidative phosphorylation eradicates therapy-resistant chronic myeloid leukemia stem cells. Nat Med. 2017;23(10):1234-40.

12. Jones CL, Stevens BM, D'Alessandro A, Reisz JA, Culp-Hill R, Nemkov T, et al. Inhibition of Amino Acid Metabolism Selectively Targets Human Leukemia Stem Cells. Cancer Cell. 2018;34(5):724-40 e4.

13. Henkenius K, Greene BH, Barckhausen C, Hartmann R, Marken M, Kaiser T, et al. Maintenance of cellular respiration indicates drug resistance in acute myeloid leukemia. Leuk Res. 2017;62:56-63.

14. Bonekamp NA, Peter B, Hillen HS, Felser A, Bergbrede T, Choidas A, et al. Small-molecule inhibitors of human mitochondrial DNA transcription. Nature. 2020;588(7839):712-6.

15. Jonas BA, Pollyea DA. How we use venetoclax with hypomethylating agents for the treatment of newly diagnosed patients with acute myeloid leukemia. Leukemia. 2019;33(12):2795-804.

16. Zhao W, Sachsenmeier K, Zhang L, Sult E, Hollingsworth RE, Yang H. A New Bliss Independence Model to Analyze Drug Combination Data. J Biomol Screen. 2014;19(5):817-21.

17. Chapuis N, Poulain L, Birsen R, Tamburini J, Bouscary D. Rationale for Targeting Deregulated
Metabolic Pathways as a Therapeutic Strategy in Acute Myeloid Leukemia. Front Oncol.
2019;9:405.

18. Sharon D, Cathelin S, Mirali S, Di Trani JM, Yanofsky DJ, Keon KA, et al. Inhibition of mitochondrial
translation overcomes venetoclax resistance in AML through activation of the integrated stress
response. Sci Transl Med. 2019;11(516).

19. Ebert BL, Pretz J, Bosco J, Chang CY, Tamayo P, Galili N, et al. Identification of RPS14 as a 5q-
syndrome gene by RNA interference screen. Nature. 2008;451(7176):335-9.

20. Patel JP, Gonen M, Figueroa ME, Fernandez H, Sun Z, Racevskis J, et al. Prognostic relevance of
integrated genetic profiling in acute myeloid leukemia. N Engl J Med. 2012;366(12):1079-89.

21. Testa U. Leukemia stem cells. Ann Hematol. 2011;90(3):245-71.

22. Savona MR, Rathmell JC. Mitochondrial Homeostasis in AML and Gasping for Response in
Resistance to BCL2 Blockade. Cancer Discov. 2019;9(7):831-3.

23. Lin KH, Winter PS, Xie A, Roth C, Martz CA, Stein EM, et al. Targeting MCL-1/BCL-XL Forestalls the
Acquisition of Resistance to ABT-199 in Acute Myeloid Leukemia. Sci Rep. 2016;6:27696.

24. Basak NP, Banerjee S. Mitochondrial dependency in progression of acute myeloid leukemia.
Mitochondrion. 2015;21:41-8.

25. Pollyea DA, Stevens BM, Jones CL, Winters A, Pei S, Minhajuddin M, et al. Venetoclax with
azacitidine disrupts energy metabolism and targets leukemia stem cells in patients with acute
myeloid leukemia. Nat Med. 2018;24(12):1859-66.

26. Cole A, Wang Z, Coyaud E, Voisin V, Gronda M, Jitkova Y, et al. Inhibition of the Mitochondrial
Protease ClpP as a Therapeutic Strategy for Human Acute Myeloid Leukemia. Cancer Cell.
2015;27(6):864-76.

27. Skrtic M, Sriskanthadevan S, Jhas B, Gebbia M, Wang X, Wang Z, et al. Inhibition of mitochondrial
translation as a therapeutic strategy for human acute myeloid leukemia. Cancer Cell.
2011;20(5):674-88.

28. Molina JR, Sun Y, Protopopova M, Gera S, Bandi M, Bristow C, et al. An inhibitor of oxidative
phosphorylation exploits cancer vulnerability. Nat Med. 2018;24(7):1036-46.

29. Kreitz J, Schonfeld C, Seibert M, Stolp V, Alshamleh I, Oellerich T, et al. Metabolic Plasticity of
Acute Myeloid Leukemia. Cells. 2019;8(8).

30. Saliba AN, John AJ, Kaufmann SH. Resistance to venetoclax and hypomethylating agents in acute
myeloid leukemia. Cancer Drug Resist. 2021;4:125-42.

31. Baccelli I, Gareau Y, Lehnertz B, Gingras S, Spinella JF, Corneau S, et al. Mubritinib Targets the
Electron Transport Chain Complex I and Reveals the Landscape of OXPHOS Dependency in Acute
Myeloid Leukemia. Cancer Cell. 2019;36(1):84-99 e8.

32. Bosc C, Selak MA, Sarry JE. Resistance Is Futile: Targeting Mitochondrial Energetics and
Metabolism to Overcome Drug Resistance in Cancer Treatment. Cell Metab. 2017;26(5):705-7.

33. Hosseini M, Rezvani HR, Aroua N, Bosc C, Farge T, Saland E, et al. Targeting Myeloperoxidase
Disrupts Mitochondrial Redox Balance and Overcomes Cytarabine Resistance in Human Acute
Myeloid Leukemia. Cancer Res. 2019;79(20):5191-203.

34. Jin S, Cojocari D, Purkal JJ, Popovic R, Talaty NN, Xiao Y, et al. 5-Azacitidine Induces NOXA to
Prime AML Cells for Venetoclax-Mediated Apoptosis. Clin Cancer Res. 2020;26(13):3371-83.

35.Huang da W, Sherman BT, Lempicki RA. Systematic and integrative analysis of large gene lists
using DAVID bioinformatics resources. Nat Protoc. 2009;4(1):44-57.

    Figure legends.

    Figure 1. IMT+venetoclax combination treatment synergistically inhibits cell

    viability/growth of AML cell lines *in vitro*.

    (**A**) Chemical structure of IMT (LDC204857) (**B**) AML cell lines were treated with single

    agents IMT or venetoclax at a range of concentrations (0.1–1000 nM and 0.01–100 nM,

    respectively), and assessed by cell viability tests after three and six days. (**C**) Synergism

    analysis of AML cell lines treated with a combination of IMT (0.05, 0.1, 0.2 μM for MV4-11

    and MOLM-13, and 0.1, 0.5, and 1 μM for THP-1 and KG-1) and venetoclax (Vex; 1, 2, 4

    nM for MV4-11 and MOLM-13, and 1, 5, 10 nM for THP-1 and KG-1) for six days. 'S'

    represents synergy of the combination treatment, evaluated using the Bliss independent

    model. Proliferation percentages are normalized to DMSO treated. All results are from three

    independent experiments.

    Figure 2. Combination treatment significantly enhances apoptosis induction in AML

    cell lines.

    AML cell lines were treated with IMT (500 nM) and venetoclax (5 nM) for 72–96 h. (**A**)

    Apoptosis was determined by Annexin V/PI staining of the cells and analyzed by flow

    cytometry. (**B**) Loss of mitochondrial membrane potential was calculated as the percentage of

    depolarized (apoptotic) cells out of the total cell population. Error bars represent standard

    error of the mean where * represents $P < 0.05$, **$P < 0.01$, and ***$p < 0.001$.

Figure 3. Combination of IMT and venetoclax dramatically decreases OXPHOS (in a

glycolysis-independent manner) in sensitive cell lines.

(**A**) Representative graph of an OXPHOS measurement using mitostress compounds:

oligomycin (OMY), carbonyl cyanide m-chlorophenyl hydrazone (CCCP), and antimycin

(AMA). (**B**) AML cell lines (MV4-11, MOLM-13, and THP-1) were treated with either IMT

(500 nM), venetoclax (5 nM), or a combination of both for four days. DMSO was added to

control wells. Oxygen consumption rate (OCR) was measured with an Oroboros O2K

instrument at physiological glucose levels (5 mM) using mitostress compounds. (**C**) AML

cell lines were cultured in growth medium supplemented with either 5 mM glucose or

galactose, and cell viability tests (ATP measurements) were performed after 24 and 48 h. All

results are from at least three independent experiments. Error bars show ± SEM. Statistical

tests on all comparisons in (C) showed non-significant differences.

Figure 4. Combination of IMT and venetoclax significantly prolongs the survival of

mice with leukemia.

(**A**) Outline of CDX efficacy experiment: NBSGW mice were transplanted with MV4-11

human leukemia cells, then treated with either vehicle, IMT, venetoclax, or a combination of

IMT and venetoclax (n=10 per treatment group). Peripheral blood was taken before treatment

(week 2) and after treatment at regular intervals until disease progression was observed. (**B**)

Analysis of proportion of human CD45+ cells in mice peripheral blood in all treatment

groups at indicated time points, determined by flow cytometry. (**C**) Kaplan-Meier analysis of

survival. Statistical significance was calculated using log-rank (Mantle-Cox) test. ****p <

0.0001, Error bars represent ± SEM. (**D**) Analysis of proportion of human CD45+ cells in the

bone marrow of all treatment groups (four mice per group compared to two surviving mice)

at time of euthanization.

Figure 5. Synergistic effect of (IMT and venetoclax) combination treatment on tumor burden and survival of AML-PDX transplanted mice.

(**A**) Experimental outline: NSGS mice were transplanted with primary AML splenocytes from PDX mice and were then treated with either vehicle, IMT, venetoclax, or a combination of IMT and venetoclax (n=10 per treatment group). After the final dosing, three or four mice were euthanized (end-of-treatment group) and the rest (n=6) were maintained for survival analysis. (**B**) Percentage of viable human CD45-positive (hCD45+) cells in peripheral blood, BM, and spleen of treated mice determined by flow cytometry. (**C**) Immunohistochemistry analysis of mouse BM and spleen sections stained with anti-human CD45 (brown) and counterstained with DAB (blue). (**D**) Quantitative analysis of hCD45+ cells on immunohistochemistry sections of BM and spleen sections of treated mice. (**E**) Proliferating tumor cells (CD45) are shown by quantification of Ki67 positive cells in immunohistochemistry-stained BM sections at end-of-treatment (results from at least three technical replicates and two biological replicates in each treatment group).

Figure 6. Synergistic effect of (IMT and venetoclax) combination treatment on gene transcription and survival of AML-PDX transplanted mice.

The experimental outline was shown in figure 5A, but n=10 per treatment group. After the final dosing, three or four mice were euthanized (end-of-treatment group) and the rest (n=6) were maintained for survival analysis. (**A**) Gene expression profiling by RNA-seq of purified AML blasts isolated from PDX models treated with venetoclax, IMT, or a combination thereof. Indicated expression changes are relative to a vehicle treated group. Significantly enriched genes are indicated in red ($q < 0.05$). (**B**) Peripheral blood engraftment analysis of survival group. (**C**) Survival analysis of treated mice shown by Kaplan-Meier analysis (n=6/treatment group). *$P < 0.05$, **$P < 0.01$, and ***$p < 0.001$.

Figure 7. Leukemia progression induced by primary AML cells is inhibited by venetoclax treatment.

As in Fig. 5A, efficacy experiment outline: NSGS mice were transplanted with primary AML splenocytes from PDX mice and were then treated with either vehicle, IMT, venetoclax, or a combination of IMT and venetoclax (n=12 per treatment group). 8–12 h after the last dosing, half of the mice (n=6/treatment group) were euthanized (end-of-treatment group) and the other half were maintained for survival analysis. (**A**) Percentage of viable human CD45-positive (hCD45+) cells in peripheral blood and bone marrow of treated mice (n=6 for each treatment group), determined by flow cytometry. (**B**) Immunohistochemistry analysis of hCD45+ cells on bone marrow and spleen tissue sections of treated mice (images are representative from at least two mice per group). Scale bars represent 100 μM. (**C**) Peripheral blood engraftment analysis of survival group. (**D**) Mice treated with single or a combination of compounds showed prolonged survival in comparison to the vehicle-treated group. Kaplan-Meier analysis shows the relative survival curve. $*P < 0.05$, $**P < 0.01$, and $***p < 0.001$.

Figure 1

a

LDC204857

b

Day 3

Cell growth (% of control) vs IMT (nM)

- THP-1
- KG-1
- MV4-11
- MOLM-13

Day 3

Cell growth (% of control) vs Venetoclax (nM)

- THP-1 — TP53-mut
- KG-1 — TP53-mut
- MV4-11 — TP53-wt
- MOLM-13 — TP53-wt

Day 6

Cell growth (% of control) vs IMT (nM)

- THP-1
- KG-1
- MV4-11
- MOLM-13

Day 6

Cell growth (% of control) vs Venetoclax (nM)

- THP-1 — TP53-mut
- KG-1 — TP53-mut
- MV4-11 — TP53-wt
- MOLM-13 — TP53-wt

c

MV4-11 day 6

% proliferation (normalized)

- 0.2 µM IMT
- 0.1 µM IMT
- 0.05 µM IMT
- DMSO

DMSO, 1 nM Vex, 2 nM Vex, 4 nM Vex

MOLM-13 day 6

% proliferation (normalized)

- 0.2 µM IMT
- 0.1 µM IMT
- 0.05 µM IMT
- DMSO

DMSO, 1 nM Vex, 2 nM Vex, 4 nM Vex

THP-1 day6

% proliferation (normalized)

- 1 µM IMT
- 0.5 µM IMT
- 0.1 µM IMT
- DMSO

DMSO, 1 nM Vex, 5 nM Vex, 10 nM Vex

KG-1 day 6

% proliferation (normalized)

- 1 µM IMT
- 0.5 µM IMT
- 0.1 µM IMT
- DMSO

DMSO, 1 nM Vex, 5 nM Vex, 10 nM Vex

Figure 2

a

b

Figure 3

Figure 4

a

b

c

d

Figure 6

A

B

Human CD45+ cells (%) in peripheral blood survival analysis

***

Vehicle Venetoclax IMT Combi

C

Dosing period (21d)

% survival

Days post-transplantation

Median Survival (days)

Vehicle (n=6)   96
IMT (n=6)   99
Venetoclax (n=6)   102
Combi (n=6)   112

***
**
***

Figure 7

B   Anti-hCD45   Nucleus