1

Reduced mitochondrial transcription sensitizes acute myeloid leukemia cells to BCL-2 inhibition

Short title: Inhibition of mtDNA expression to treat leukemia

Authors

Laleh S. Arabanian[1], Jenni Adamsson[2], Anke Unger[4], Raffaella Di Lucrezia[4], Tim Bergbrede[4], Arghavan Ashouri[1], Erik Larsson[1], Peter Nussbaumer[4], Bert M. Klebl[4,5], Lars Palmqvist[2,3,*], Claes M. Gustafsson[1,3*,†]

Affiliations

[1] Department of Medical Biochemistry and Cell Biology, University of Gothenburg, Gothenburg, Sweden

[2] Department of Laboratory Medicine, University of Gothenburg, Gothenburg, Sweden.

[3] Department of Clinical Chemistry, Sahlgrenska University Hospital, Gothenburg, Sweden

[4] Lead Discovery Center GmbH, Otto-Hahn-Str. 15, D-44227 Dortmund, Germany

[5] The Norwegian College of Fishery Science, UiT The Arctic University of Norway, PO Box 6050 Langnes, N-9037 Tromsø, Norway

*Corresponding authors: lars.palmqvist@clinchem.gu.se and claes.gustafsson@medkem.gu.se

†Lead Contact

Abstract

Overcoming drug-resistance and the subsequent relapse that often occurs with monotherapy is

crucial in the treatment of acute myeloid leukemia. We here demonstrate that therapy-resistant

leukemia initiating cells can be targeted using a novel inhibitor of mitochondrial transcription

(IMT). The compound inhibits mitochondrial RNA polymerase activity and sensitizes the

resistant population to the induction of apoptosis. *In vitro* studies on acute myeloid leukemia

cells demonstrate that IMT prevents cell proliferation, and together with a selective BCL-2

inhibitor, venetoclax, induce apoptosis and suppress oxidative phosphorylation (OXPHOS)

synergistically. AML mouse models treated with IMT in combination with venetoclax show

prolonged survival in venetoclax-resistant models. Our findings suggest that certain therapy-

resistant leukemia cell populations display a unique dependency on mitochondrial transcription

and can be targeted with IMT.

Teaser

Leukemia cells resistant to standard treatment are sensitive to a recently developed inhibitor of

mitochondrial gene expression.

Introduction

Acute myeloid leukemia (AML) comprises a devastating and heterogeneous group of diseases, for which patient survival rates have remained stagnant over the last several decades. There is no one standard AML treatment as it depends on the individual situation of each patient. Current cares are focused on allogenic stem cell transplantation as well as patient stratification and therapy selection based on genetic aberrations. These lead to cures in only a fraction of AML patients, highlighting the critical need to develop different approaches. The anti-apoptotic molecule BCL-2 has been a therapeutic target in lymphoid and myeloid leukemia, as in many cancers overexpression of BCL-2 results in dysregulation of apoptosis (*1*). ABT-199 (venetoclax), is a specific BH3-mimetic that causes apoptosis by preventing BCL-2 from binding pro-apoptotic BAX and BAK1 proteins (*2*). However, complete responses occur in just ~20% of AML patients treated with Venetoclax (*3*). In addition, many patients relapse, with regrowth initiated by therapy-resistant leukemic clones, suggesting that single agent administration is unlikely to yield lasting responses in most cases. Venetoclax is therefore used in combination with conventional therapies. Together with hypomethylating agents, Venetoclax is now standard of care for patients deemed unfit for intensive induction therapy (*4*).

Resistance to venetoclax may result from overexpression of the anti-apoptotic proteins MCL1 or BCL2L1 (*5*), or through changes in levels of other anti-apoptotic molecules (*6*). Aberrant pro-survival regulation can also be achieved by inactivation of TP53 function, which controls many aspects of apoptosis (*6, 7*). MCL1-dependent resistance to venetoclax can be overcome when the compound is combined with the MDM2 inhibitor RG7388, which causes p53 activation. A combination of BCL-2 inhibition and TP53 activation may thus provide an interesting therapeutic approach for treatment of AML (*8*).

Other combination approaches that promote apoptosis may also be of interest. The majority of

ATP required for leukemia stem cell (LSC) function is generated by oxidative phosphorylation

(OXPHOS) (*9, 10*). Interestingly, venetoclax-resistant leukemic cells display increased levels

of OXPHOS and are protected from mitochondrial stress (*7*). These data suggest approaches

to specifically target OXPHOS in leukemia may be an attractive option. In support of this

notion, inhibition of mitochondrial protein translation by antibiotics that reduce OXPHOS

capacity results in a selective cytotoxic effect on leukemic cell when used in combination with

thyrosine kinase inhibitors (*11*). Furthermore, LSCs from *de novo* AML patients display a

unique dependency on amino acid metabolism to fuel OXPHOS. When venetoclax is combined

with the hypomethylating agent azacytidine, the two compounds act in concert to inhibit amino

acid uptake, leading to a drop in OXPHOS and thus cell death (*12*). More recently, AML

therapy using a combination of cytarabine (AraC) and venetoclax has highlighted the central

role of mitochondrial adaptation during treatment (*10, 13*), indicating the need for a

mitochondrial directed combination therapy for AML. Such a therapy may help to overcome

existing treatment resistance.

Recently we developed novel, first-in-class inhibitors of mitochondrial transcription (IMTs)

(*14*). IMT treatment specifically targets mitochondrial RNA polymerase (POLRMT) and

reduce transcription of mitochondrial DNA, which codes for thirteen key components of the

OXPHOS machinery, including subunits of respiratory complexes I, III, IV, and the ATP

synthase. As a consequence, IMT treatment impairs OXPHOS system biogenesis in a dose-

dependent manner, and decreases cell viability in various tumor cells. Surprisingly, several

weeks of per oral IMT treatment is well tolerated in mice and does not cause toxicity in normal

tissues, despite inducing a strong anti-tumor response in human cancer xenografts (*14*).

Here we examine AML cell lines for their sensitivity for treatment with IMT. We find that a

combination of IMT and venetoclax displays a synergistic on cell growth and induction of

apoptosis. We demonstrate that the efficacy of this approach in leukemia mouse models

generated by human AML cell lines as well as primary samples from AML patients.

Results

Synergistic effect of IMT in combination with venetoclax on AML cell lines

We were curious whether our inhibitors of mitochondrial transcription (IMTs) (*14*) could be effective against AML cells. The coumarin class of IMTs was optimized by medicinal chemistry to generate lead IMT compounds with improved metabolic stability in mice, such as LDC204857, herein simply referred to as IMT (Fig 1A, Fig. S1 and Table S1). IMT is a close analog of previously described IMT1B (Fig. S1) with a highly similar biochemical and cellular inhibition profile (*14*). It shows an extended half-life *in vivo* due to its increased metabolic stability (Table S1). Thus, by using IMT we could consistently reduce the mouse dose to 50 mg/kg per oral once daily, which is very well tolerated and does not exert any adverse effects in mice.

Using four AML cell lines with different expression of TP53 (MV4-11 and MOLM-13 expressing wild type TP53; THP-1 and KG-1 expressing mutated TP53), we monitored the effects of increasing concentrations of IMT on cellular growth using a luminescence-based cell viability assay. Only marginal effects were observed after a 72-h period of exposure to IMT, but all cell lines displayed dose-dependent antileukemic activity after 144 h of treatment (Fig. 1B), in line with our previous observations (*14*). We also monitored venetoclax sensitivity by assessing cell viability, and noted that, in line with published work (*15*), a dose-dependent reduction in cell growth occurs after three days and becomes more pronounced after 6 days (Fig. 1B).

To then investigate whether IMTs show a synergistic effect when combined with venetoclax, we conducted a combination study in AML cell lines (Fig. 1C). After six days of treatment, we found a clear synergistic effect on cell proliferation in MV4-11 and MOLM-13 cell lines, which

express wild type TP53, as determined by combination analysis (Bliss-independence model

(*16*)). Treatment of THP-1 and KG-1 cell lines, which harbor mutated TP53, only revealed a

synergistic effect when using higher concentrations of both IMT and venetoclax (Fig. 1C,

additional treatment data are shown in Fig. S2).

Combination of IMT and venetoclax synergistically induces apoptosis and loss of

mitochondrial membrane potential in MV4-11 and MOLM-13 cells

To investigate the factors involved in the inhibition of cell growth with IMT and venetoclax,

AML cell lines were treated with single agents (IMT or venetoclax) or a combination of the

two (Fig. 2A), and the stages of apoptosis were monitored using Annexin V and PI staining.

No detectable increase in apoptosis was induced by IMT exposure alone in any AML cell line.

Venetoclax treatment resulted in both early and late apoptosis induction in MV4-11 and

MOLM-13 cell lines, and employing a combination of IMT and venetoclax significantly

enhanced apoptosis induction in these wild type cell lines. On the other hand, combination

treatment did not induce apoptosis in THP-1 and KG-1 mutated cell lines (Fig. 2A).

One distinctive feature of apoptosis is the disruption of mitochondrial function, which can be

measured as loss of mitochondrial membrane potential due to membrane depolarization. As a

combination of IMT and venetoclax could synergistically induce apoptosis, we were curious

whether combination drug treatment thus affects mitochondrial membrane potential. We

observed that venetoclax-sensitive cell lines (MV4-11 and MOLM-13) showed a dramatic loss

of mitochondrial membrane potential when treated with both venetoclax and IMT (Fig. 2B).

On the other hand, mitochondrial membrane potential in THP-1 cells was not affected by single

nor by combination treatment. Interestingly, although KG-1 was venetoclax insensitive, the

combination treatment significantly enhanced loss of mitochondrial membrane potential (Fig.

2B). These data suggest that a combination of IMT and venetoclax is cytotoxic and

synergistically induces cell death in sensitive cells, presumably through disruption of

mitochondrial function.

Combination of IMT and venetoclax significantly suppresses OXPHOS in AML cells

We next studied whether combination treatment with IMT and venetoclax affects OXPHOS in

AML cells. OXPHOS was evaluated by determining oxygen consumption rate (OCR) (Fig.

3A), which was measured first as basal respiration and then oligomycin (OMY)-induced leak

respiration (ATP-linked respiration) after the addition of oligomycin to inhibit ATP synthase.

Stepwise carbonyl cyanide m-chlorophenyl hydrazone (CCCP) titrations were subsequently

carried out to reach the electron transport chain respiratory capacity state. Finally, antimycin a

(AMA) was added to inhibit complex I/III of the electron transport chain and to terminate

respiration, allowing for the determination and correction of residual oxygen consumption,

which is indicative of non-mitochondrial oxygen consumption (Fig. 3A). We noted that

combination treatment significantly inhibits OXPHOS respiration in venetoclax-sensitive

MOLM-13 and MV4-11 cells (Fig. 3B). In comparison, OXPHOS levels in venetoclax-

resistant THP-1 cells were unaffected in the presence of single compound or combination

treatments, confirming that the inhibition we observed is not a general effect of drug treatment

(Fig. 3B).

In normal hematopoietic cells, glycolysis is upregulated when OXPHOS is blocked (*17*). In

contrast to normal hematopoietic cells, high energy turnover AML cells rely predominantly on

OXPHOS due to poorly regulated metabolic adaptation. To verify the reliance on OXPHOS,

AML cells were grown in growth medium supplemented with either glucose or galactose (Fig.

3C). Two ATP molecules are produced when cells use glucose for glycolysis, but if galactose

is used instead, no ATP is generated (*18*). We observed no significant decrease in the level of

ATP production (Fig. 3C) nor in cell counts (Fig. S3A) in the absence of glucose, indicating

that glycolysis is not a major source of energy in AML cells. To test whether AML cells use

glycolysis under treatment conditions, OCR was measured in the presence or absence of a high

concentration of glucose (Fig. S3B). OCR measured for basal respiration was not significantly

different with excess amounts of glucose, again indicating the inability of AML cells to switch

to glycolysis when OXPHOS is disrupted.

Combination of IMT and venetoclax synergistically delays leukemia initiation in cell-derived leukemia mouse models

Following the confirmed synergy between IMT and venetoclax in AML cell lines, we

proceeded to test the dual inhibition of mitochondrial transcription and BCL-2 in an AML cell-

derived xenograft (CDX) transplantation mouse model. First, the pharmacokinetic profile of

IMT was determined in plasma, liver, and bone marrow (BM) of mice, 1, 8, and 24 h post-

administration (Fig. S4A). As described above (Fig. S1), the IMT mouse PK profile was

superior to IMT1B. IMT was detected in all tissues at all time points when administered alone.

To determine the concentration of IMT and venetoclax when administered in combination,

plasma samples were then collected 8–11 h post-administration of single or combination

formulations and the concentrations of IMT and/or venetoclax in samples were measured (Fig.

S4B). Values obtained were all within the effective range, as combination treatment did not

affect the individual compound concentrations (*14*). For xenograft experiments, NBSGW mice

were intravenously transplanted with MV4-11 AML cells (Fig. 4A). Two weeks after

transplantation, engraftment of AML cells was confirmed (evaluated by gating on hCD45+

population by flow cytometry in peripheral blood). Treatment then began with vehicle, 50

mg/kg IMT, 100 mg/kg venetoclax, or a combination of IMT (50 mg/kg) and venetoclax (100

mg/kg) (Fig. 4A), dosed as such based on previously published protocols (*14, 18*). The efficacy of treatments was determined by measuring the proportion of transplanted AML cells in peripheral blood and by survival analysis (Fig. 4B). In all treatment groups, the hCD45+ population dropped immediately after treatment, and vehicle- and venetoclax-treated mice then exhibited a rapid rebound. In contrast, in IMT-treated mice the increase in leukemic cell proportion was delayed until eight weeks post-transplantation, and in the combination-treated group for to up to 20 weeks (Fig. 4B). These observations correlate with results from survival analysis (Fig. 4C). IMT treatment of MV4-11 transplanted mice significantly delayed the development of leukemia, inducing a 37% increase in median overall survival (57.5 → 79 days). Development of leukemia was further delayed by combination treatment to 131% (57.5 → 133 days), and the group contained two mice that did not show any disease symptoms after 200 days (post-transplantation) when they had to be euthanized according to ethical permissions. Moreover, leukemic cells were not detectable in the BM of surviving mice at the time of termination. All other mice showed symptoms of leukemia and of these, all but one combination-treated mouse had an increased percentage of leukemic cells in BM (Fig. 4D). These results confirm that treatment with a combination of IMT and venetoclax significantly prolongs leukemia progression in mice. Moreover, the general condition of mice such as behavior, fur, mobility and weight, assessed before (week 2) and after treatment (week 5), were not altered in the IMT-, venetoclax-, or combination-treatment groups when compared to the vehicle-treated group (weights shown in Fig. S5B), suggesting that the treatments were well tolerated. As with single IMT therapy (*14*), treatment with combination therapy did not display any evidence of organ toxicity or drug-associated death in the mice (Fig. S5C, D).

IMT and venetoclax synergistically decrease the tumor burden and increase the life span in an AML-PDX mouse model resistant to venetoclax.

The CDX mouse model demonstrated IMT could sensitize AML cell lines to BCL-2 inhibitors

in vivo. We next analyzed effects of combination treatment on a more refined *in vivo* model by

generating xenografts using AML patient-derived cells (AML PDX). We first tested an AML-

PDX mouse model with resistance to venetoclax (AML sample harboring TP53 mutation).

Two weeks post-transplantation with AML PDX cells, NSGS mice were orally treated with

IMT, venetoclax, or a combination of the two for 21 days (Fig. 5A). The efficacy of the

treatments was determined based on the proportion of human AML cells (determined by

hCD45+ population). Six out of 10 mice in each group were euthanized immediately after the

last compound administration for end-of-treatment analysis. We found that the combination of

IMT and venetoclax resulted in significant inhibition of tumor burden in blood, BM, and spleen

of treated mice (Fig. 5B). By examining histology sections of BM and spleen, we found that

tissues from combination-treated animals contained less engraftment of leukemic blasts than

single- and vehicle-treated mice (Fig. 5C). Accordingly, the frequencies of hCD45+ cells in

the BM and spleens of the combination-treated group were largely reduced in comparison to

other groups (Fig. 5D), and engrafted tumor cells in combination-treated PDX mice showed a

lower proliferating percentage in comparison to single or vehicle treated groups (shown by

KI67+/hCD45+ immunohistochemistry staining; Fig. 5E). This demonstrates a synergy

between IMT and venetoclax in reducing tumor burden.

To further analyze the cellular effects of the IMT-venetoclax combination treatment, we

analyzed transcriptomes of purified viable AML blasts isolated from PDX models at end of

treatment (Fig. 6A). We confirmed that combination treatment had a more pronounced effect

on gene transcription as compared to single treatments. In agreement with its more pronounced

effect on tumor burden, combination treatment suppressed expression of a number of genes

involved in leukemogenesis, including MX1, KCTD12, ITGA and RGL1 (Table S2) (*19*).

We also analyzed survival by monitoring the mice for disease progression and morbidity.

Peripheral blood engraftments were analyzed at regular intervals until the appearance of

disease symptoms (Fig. 6B and Fig. S6A). The leukemia burden immediately prior to

euthanization demonstrated that the cytoreductive effect of the combination treatment persists

(Fig. 6B). However, blood count (determined by RBCs and HGB levels) did not vary

significantly between experimental groups, nor did the magnitude of splenomegaly or liver

weight vary at the time of euthanization (Fig. S6B–D). Interestingly, mice treated with a

combination of IMT and venetoclax succumbed to death significantly later than other treatment

groups, with a median survival of 112 days vs 96, 99, and 102 days for vehicle-, IMT-, and

venetoclax-treated groups, respectively (Fig. 6**C**). Combination treatment thus increases the

life span of venetoclax-resistant AML mouse models, demonstrating that IMT can sensitize

AML cells towards other therapies.

IMT does not affect tumor burden or life span in an AML-PDX mouse model sensitive to

venetoclax.

For comparison, we also monitored effects of combination treatment on a PDX model based

on AML cells sensitive to venetoclax treatment (AML cells harboring FLT3-ITD, NMP1 and

DNMT3A). As expected, we observed that venetoclax treatment significantly reduced the

tumor burden in the blood and BM of mice. Combination treatment with IMT did not add to

this effect (Fig. 7A). By examining histology sections, we found that both BM and spleen

tissues from combination- and venetoclax-treated animals contained reduced proportion of

leukemic blasts compared to IMT- and vehicle-treated mice (Fig. 7B). White blood counts

(WBCs), red blood counts (RBCs), and hemoglobin (HGB) levels were not significantly

different across experimental groups (Fig. S7A). Total body and liver weight of PDX mice

were very similar in all groups (Fig. S7B, D), but both venetoclax and combination treatments

reduced splenomegaly (Fig. S7C).

The remaining half of mice were used for survival analysis, which involved regular blood count

and chimerism analysis until the appearance of disease symptoms (Fig. 7C). Both venetoclax

and IMT caused a modest increase in life span, but combination treatment did not offer an

additional effect (Fig. 7D).

Discussion

In recent years, the genetic complexity and heterogeneity of leukemia cell populations has become apparent (*20, 21*) and developing effective strategies to target human leukemia initiating cells has thus proven to be a challenging task. Currently, there is considerable interest in therapies, such as venetoclax, that can restore normal apoptosis of cancer cells by directly interacting with the BCL-2 family of proteins. These are effective for some leukemias, but many patients eventually become refractory to treatment and ultimately succumb to their disease (*22, 23*). It has become increasingly important to determine which biological properties are relatively consistent in LSC populations irrespective of genetic makeup (tumor population). Recent observations indicate that several cancer cell types, including LSCs, are reliant on OXPHOS for metabolism (*24*), making inhibition of mitochondrial function a promising therapeutic strategy for AML (*12, 25-28*). Therapy-resistant LSCs are more reliant on oxidative respiration than glycolysis for energy generation, suggesting that the OXPHOS dependency and lack of normal glycolytic activity in LSCs may be characteristic of cancers (*29*). Understanding resistance mechanisms will allow for the design of strategies to counteract drug resistance or impede resistance from developing. In AMLs with aberrant expression of BCL-2, venetoclax can reduce OXPHOS (*9*) and trigger cell death. However, venetoclax resistance can occur in patients treated with the BCL-2 inhibitor and in some subtypes of AML patients with poor prognoses (*30*).

Previous work has indicated that AML cells display increased mitochondrial biogenesis and inhibition of mitochondrial translation can be used as a strategy for human AML (*18, 27*). In our work, we target another process involved in mitochondrial gene expression, i.e., transcription. IMT treatment targets POLRMT, which has two main functions in mitochondria: it transcribes mtDNA encoded genes and provides the primers required to initiate mtDNA

replication. Inhibition of POLRMT can thus deliver a double punch, both reducing mtDNA

copy number and inhibit OXPHOS biogenesis. As demonstrated here inhibition of POLRMT

can overcome venetoclax therapy resistance in AML. These observations are in line with recent

reports of the importance of mitochondrial function in leukemia cell survival (*31, 32*). By

developing a clinically relevant and well-tolerated approach to deliver IMTs in

immunodeficient mice transplanted with AML cells, we were able to reduce the tumor burden

in all (cell- and patient-derived) xenografted mice treated with a combination of IMT and

venetoclax. This demonstrates that reducing POLRMT activity results in successful targeting

of leukemic cells. Our *in vitro* investigations confirmed that the combination of IMT and

venetoclax is necessary to significantly decrease ATP levels and consequently induce

apoptosis. In addition, efficacy studies of AML PDX models showed that single compounds

are not sufficient to eradicate AML cells, but that the combination regimen reduced AML

tumor burden and prolonged survival in all treated mice. These data indicate the advantage of

both downregulating POLRMT activity in combination with increasing apoptosis and

highlights the importance of rational combination therapies (*33, 34*).

In summary, we have introduced a new therapeutic strategy to target therapy-resistant leukemia

with OXPHOS-dependent properties. Despite several limitations with the design of pre-clinical

cancer models and the intensity and duration of treatments, we demonstrate here that the

combined inhibition of BCL-2 and mitochondrial transcription can target therapy-resistant

AMLs. This combination treatment succeeds as IMT primes cell vulnerability to apoptosis

induction. Consequently, our findings may be of value for the future development of better

clinical strategies for the treatment of AML.

Materials and Methods

Cell lines and culture conditions

All cell lines (MV4-11, MOLM13, THP-1, KG-1, OCI-AML3, and OCI-AML2) were purchased from DSMZ (Braunschweig, Germany) and cultured in RPMI-1640 (Sigma Aldrich) supplemented with 10–20% heat inactivated fetal bovine serum (Gibco) and 1% penicillin/streptomycin (Invitrogen). Cells were maintained in a humidified incubator at 37 °C and 5% CO_2. RPMI-1640 without glucose was used for *in vitro* treatment experiments and supplemented by physiological concentrations (5 mM) of glucose (D-Glucose, Sigma) or galactose (D-galactose, Sigma).

Generation of *in vivo* AML mouse model

Mice were purchased from Jackson laboratories (USA) and maintained in a pathogen-free facility at the University of Gothenburg. All animal experiments were conducted in accordance with guidelines approved by the research animal ethics committee at the University of Gothenburg. For AML cell line xenograft (CDX) experiments, 6–8-week-old, male NBSGW mice (NOD.Cg-Kit^{W-41J} Tyr$^+$ Prkdc^{scid} Il2rg^{tm1Wjl}/ThomJ, Jackson laboratories) were used. Mice were transplanted intravenously with $\sim 1 \times 10^6$ MV4-11 cells. For patient-derived xenograft (PDX) experiments, 6–8-week-old, male NSGS mice (NOD.Cg-Prkdc^{scid} Il2rg^{tm1Wjl} Tg(CMV-IL3,CSF2,KITLG)1Eav/MloySzJ, Jackson laboratories) were sub-lethally irradiated with 1 Gy radiation. AML PDX splenocytes were purchased from Jackson laboratories (id: J000106134*, Positive for FLT3-ITD, DNMT3A, and NPM1mutaions; and id: 000106566*, Positive for TP53 mutation). Mice were transplanted intravenously with $\sim 1 \times 10^6$ splenocytes from AML PDX mice containing more than 80% patient AML cells.

Human leukemia cell frequency assessment by flow cytometry

The percentage of human leukemic cells (MV4-11 cell lines and AML PDX cells) in mouse

peripheral blood, BM, and spleen was determined by flow cytometry. Mice were bled at

indicated time points after transplantation. Prior to staining, peripheral red blood cells were

lyzed using ammonium chloride for 8–10 min at room temperature. Cells were stained with

antibodies, washed, and submitted for flow-cytometric analysis using a FACS Aria (BD

Biosciences, USA) or CytoFlex instrument (Beckman Coulter, California, USA). At least

10,000 events were acquired, and the percentage was calculated relative to gated viable

mononuclear cells (MNCs). To discriminate between human and mouse cells, cells were

stained with antibodies against human CD45 (clone HI30, BD Biosciences) and mouse CD45

(clone 2D1, BD Biosciences). Engraftment was deemed positive if human CD45+ cells

represented >0.1% of the viable MNCs. 7AAD (BD Biosciences) served as viability marker.

Preparation of compounds for *in vivo* drug administration

IMT was optimized based on the structure activity and property relationship of the coumarin

mtRNAPol inhibitor class by LDC (Dortmund, Germany). Venetoclax was obtained from

MedChemExpress (Hölzel Diagnostika Handels GmbH, Germany). Following the

development of a suitable vehicle for the formulation of IMT and venetoclax, as well as the

combination of both compounds, the test articles were formulated using 0.5% aqueous

methylcellulose solution (Sigma-Aldrich, Germany). After randomization, and two weeks

post-transplantation or with confirmation of engraftment, mice were treated with vehicle (0.5%

methylcellulose), 50 mg/kg IMT, 100 mg/kg venetoclax, or a combination (100 mg/kg

venetoclax + 50 mg/kg IMT) by oral administration daily for 21 days.

Cell proliferation assay

Cells were seeded out ca 20 h prior to treatment. For dose response experiments, cells were

treated with a wide range of concentrations of single compounds. Cell proliferation assays were

performed every 24 h from 72–168 h, where cell viability/ATP production was determined

using CellTiter-Glo (Promega).

Apoptosis assay and Mitochondrial potential assay

Apoptosis was evaluated by Annexin V and PI staining and mitochondrial membrane potential

was evaluated by lipophilic cationic dye JC-1. Cells were treated with IC50 concentrations of

IMT, venetoclax, or combination of both, analyzed using an advanced image cytometer

(Nucleocounter NucleoCounter's accompanying software NucleoView).

High-resolution respirometry

AML cell lines were treated with DMSO, IMT, venetoclax, or combination of IMT and

venetoclax for 3–4 days. The mitochondrial respiratory capacity (oxygen consumption rate)

was measured using high resolution respirometry (O2K, Oroboros, Innsbruck, Austria). A

substrate-uncoupler-inhibitor titration (SUIT) protocol was used to measure specific features

of mitochondrial respiration. Respiratory capacities were tested in a sequence of coupling

states: basal respiration; LEAK respiration using 5 mM oligomycin (Sigma); and non-coupled

electron transfer state (ET state) where the optimum carbonyl cyanide 3-

chlorophenylhydrazone (CCCP, Sigma) concentration for maximum flux was determined by

titration (max. 3 μM CCCP). Mitochondrial respiration was inhibited using 2.5 μM antimycin

(Sigma). Where stated, excess glucose (D-Glucose, Merck) was added at a final concentration

of 20 mM. Instrumental and oxygen background fluxes were calibrated as a function of oxygen

concentration and subtracted from the total volume-specific oxygen flux. Calibration and data

analysis was performed using Datlab v.6.1 software (Oroboros Instruments).

Pharmacokinetics analysis

IMT and venetoclax were extracted from plasma and tissues by protein precipitation using acetonitrile. Tissue samples were homogenized using double the (sample) volume of PBS, and bone marrow samples were suspended in 40 µL of PBS prior to extraction with acetonitrile. Samples were analyzed by liquid chromatography tandem-mass spectrometry using a Prominence UFLC system (Shimadzu) coupled to a Qtrap 5500 instrument (ABSciex). Test articles were separated on a C18 column using a gradient elution with an acetonitrile/water mixture containing 0.1% formic acid as the mobile phase. Chromatographic conditions and mass spectrometry parameters were optimized for each test article prior to sample analysis. Concentrations of each test article were calculated by means of a standard curve.

mRNA extraction and transcriptome sequencing of AML PDX cells

For RNAseq experiment BM samples were FACS sorted using BD FACS Aria fusion (BD Biosciences). Human AML cells were isolated using human CD45 antibody. 7-AAD (BD Biosciences) was used for dead cell exclusion. RNA was extracted using RNeasy Micro kit (Qiagen). RNAseq library preparation and RNA-sequencing was done using the SMARTer Total Stranded RNA pico kit. RNA library was prepared at National Genomics Infrastructure (NGI) to be sequenced on the NovaSeq S6000 on 0.5 lanes of one S4 flowcell.

Bioinformatics analysis

Analysis was performed in R (www.r-project.org, accessed on 10 May 2022) using DESeq2, clusterProfiler and ReactomePA packages. Prefiltering to keep genes that have at least 10 reads total was performed before differential expression analysis was done with DESeq2. Visualization of differentially expressed genes was done with EnhancedVolcano to create

434   volcano plots and gridExtra for creating tables. The *p*-value cutoff was set to 0.1 to allow for

435   more genes to be found in EP (End-of-treatment) group comparisons and the Res/Sen

436   (Resistant/Sensitive) group comparisons. Upregulated and downregulated genes were plotted

437   and colored according to Fold Change (FC) greater than 1.5 or -1.5 and the *p*-value cutoff 0.1.

Statistical Analysis

Unless otherwise noted, data were summarized using mean ± standard error of the mean (±

SEM). Per-group sample sizes presented in figures and results are reported from ≥3 separate

experiments, unless stated otherwise. Survival distributions were estimated using the Kaplan–

Meier method, and group comparisons were conducted using the log-rank test. Data were

analyzed using GraphPad Prism 9.0. Combination analysis was summarized using the Bliss-

independent model (*16*). The formula $Y_{ab,P} = Y_a + Y_b - Y_a Y_b$ was used where drug A at dose a

inhibits Y_a percent of cell growth, and drug B at dose b inhibits Y_b percent of cell growth. The

predicted inhibitory effect of combined percentage was estimated by $Y_{ab,P}$. The observed

combined percentage inhibition $Y_{ab,O}$ was then compared with $Y_{ab,P}$, and if $Y_{ab,O} > Y_{ab,P}$, the

combination treatment was thought to be more efficacious (synergy). If $Y_{ab,O} = Y_{ab,P}$ the

combination treatment was thought to be additive.

References

1. R. Guieze, V. M. Liu, D. Rosebrock, A. A. Jourdain, M. Hernandez-Sanchez, A. Martinez Zurita, J. Sun, E. Ten Hacken, K. Baranowski, P. A. Thompson, J. M. Heo, Z. Cartun, O. Aygun, J. B. Iorgulescu, W. Zhang, G. Notarangelo, D. Livitz, S. Li, M. S. Davids, A. Biran, S. M. Fernandes, J. R. Brown, A. Lako, Z. B. Ciantra, M. A. Lawlor, D. B. Keskin, N. D. Udeshi, W. G. Wierda, K. J. Livak, A. G. Letai, D. Neuberg, J. W. Harper, S. A. Carr, F. Piccioni, C. J. Ott, I. Leshchiner, C. M. Johannessen, J. Doench, V. K. Mootha, G. Getz, C. J. Wu, Mitochondrial Reprogramming Underlies Resistance to BCL-2 Inhibition in Lymphoid Malignancies. *Cancer Cell* **36**, 369-384 e313 (2019).

2. A. J. Souers, J. D. Leverson, E. R. Boghaert, S. L. Ackler, N. D. Catron, J. Chen, B. D. Dayton, H. Ding, S. H. Enschede, W. J. Fairbrother, D. C. Huang, S. G. Hymowitz, S. Jin, S. L. Khaw, P. J. Kovar, L. T. Lam, J. Lee, H. L. Maecker, K. C. Marsh, K. D. Mason, M. J. Mitten, P. M. Nimmer, A. Oleksijew, C. H. Park, C. M. Park, D. C. Phillips, A. W. Roberts, D. Sampath, J. F. Seymour, M. L. Smith, G. M. Sullivan, S. K. Tahir, C. Tse, M. D. Wendt, Y. Xiao, J. C. Xue, H. Zhang, R. A. Humerickhouse, S. H. Rosenberg, S. W. Elmore, ABT-199, a potent and selective BCL-2 inhibitor, achieves antitumor activity while sparing platelets. *Nat Med* **19**, 202-208 (2013).

3. M. Konopleva, D. A. Pollyea, J. Potluri, B. Chyla, L. Hogdal, T. Busman, E. McKeegan, A. H. Salem, M. Zhu, J. L. Ricker, W. Blum, C. D. DiNardo, T. Kadia, M. Dunbar, R. Kirby, N. Falotico, J. Leverson, R. Humerickhouse, M. Mabry, R. Stone, H. Kantarjian, A. Letai, Efficacy and Biological Correlates of Response in a Phase II Study of Venetoclax Monotherapy in Patients with Acute Myelogenous Leukemia. *Cancer Discov* **6**, 1106-1117 (2016).

4. A. W. Roberts, A. H. Wei, D. C. S. Huang, BCL2 and MCL1 inhibitors for hematologic malignancies. *Blood* **138**, 1120-1136 (2021).

5. H. E. Ramsey, M. A. Fischer, T. Lee, A. E. Gorska, M. P. Arrate, L. Fuller, K. L. Boyd, S. A. Strickland, J. Sensintaffar, L. J. Hogdal, G. D. Ayers, E. T. Olejniczak, S. W. Fesik, M. R. Savona, A Novel MCL1 Inhibitor Combined with Venetoclax Rescues Venetoclax-Resistant Acute Myelogenous Leukemia. *Cancer Discov* **8**, 1566-1581 (2018).

6. J. H. Choi, J. M. Bogenberger, R. Tibes, Targeting Apoptosis in Acute Myeloid Leukemia: Current Status and Future Directions of BCL-2 Inhibition with Venetoclax and Beyond. *Target Oncol* **15**, 147-162 (2020).

7. T. Nechiporuk, S. E. Kurtz, O. Nikolova, T. Liu, C. L. Jones, A. D'Alessandro, R. Culp-Hill, A. d'Almeida, S. K. Joshi, M. Rosenberg, C. E. Tognon, A. V. Danilov, B. J. Druker, B. H. Chang, S. K. McWeeney, J. W. Tyner, The TP53 Apoptotic Network Is a Primary Mediator of Resistance to BCL2 Inhibition in AML Cells. *Cancer Discov* **9**, 910-925 (2019).

8. R. Pan, V. Ruvolo, H. Mu, J. D. Leverson, G. Nichols, J. C. Reed, M. Konopleva, M. Andreeff, Synthetic Lethality of Combined Bcl-2 Inhibition and p53 Activation in AML: Mechanisms and Superior Antileukemic Efficacy. *Cancer Cell* **32**, 748-760 e746 (2017).

9. E. D. Lagadinou, A. Sach, K. Callahan, R. M. Rossi, S. J. Neering, M. Minhajuddin, J. M. Ashton, S. Pei, V. Grose, K. M. O'Dwyer, J. L. Liesveld, P. S. Brookes, M. W. Becker, C. T. Jordan, BCL-2 inhibition targets oxidative phosphorylation and selectively eradicates quiescent human leukemia stem cells. *Cell Stem Cell* **12**, 329-341 (2013).

10. T. Farge, E. Saland, F. de Toni, N. Aroua, M. Hosseini, R. Perry, C. Bosc, M. Sugita, L. Stuani, M. Fraisse, S. Scotland, C. Larrue, H. Boutzen, V. Feliu, M. L. Nicolau-Travers, S. Cassant-Sourdy, N. Broin, M. David, N. Serhan, A. Sarry, S. Tavitian, T. Kaoma, L. Vallar, J. Iacovoni, L. K. Linares, C. Montersino, R. Castellano, E. Griessinger, Y. Collette, O. Duchamp, Y. Barreira, P. Hirsch, T. Palama, L. Gales, F. Delhommeau, B. H. Garmy-Susini, J. C. Portais, F. Vergez, M. Selak, G. Danet-Desnoyers, M. Carroll, C. Recher, J. E. Sarry,

Chemotherapy-Resistant Human Acute Myeloid Leukemia Cells Are Not Enriched for Leukemic Stem Cells but Require Oxidative Metabolism. *Cancer Discov* **7**, 716-735 (2017).

11. E. M. Kuntz, P. Baquero, A. M. Michie, K. Dunn, S. Tardito, T. L. Holyoake, G. V. Helgason, E. Gottlieb, Targeting mitochondrial oxidative phosphorylation eradicates therapy-resistant chronic myeloid leukemia stem cells. *Nat Med* **23**, 1234-1240 (2017).

12. C. L. Jones, B. M. Stevens, A. D'Alessandro, J. A. Reisz, R. Culp-Hill, T. Nemkov, S. Pei, N. Khan, B. Adane, H. Ye, A. Krug, D. Reinhold, C. Smith, J. DeGregori, D. A. Pollyea, C. T. Jordan, Inhibition of Amino Acid Metabolism Selectively Targets Human Leukemia Stem Cells. *Cancer Cell* **34**, 724-740 e724 (2018).

13. K. Henkenius, B. H. Greene, C. Barckhausen, R. Hartmann, M. Marken, T. Kaiser, M. Rehberger, S. K. Metzelder, W. J. Parak, A. Neubauer, C. Brendel, E. Mack, Maintenance of cellular respiration indicates drug resistance in acute myeloid leukemia. *Leuk Res* **62**, 56-63 (2017).

14. N. A. Bonekamp, B. Peter, H. S. Hillen, A. Felser, T. Bergbrede, A. Choidas, M. Horn, A. Unger, R. Di Lucrezia, I. Atanassov, X. Li, U. Koch, S. Menninger, J. Boros, P. Habenberger, P. Giavalisco, P. Cramer, M. S. Denzel, P. Nussbaumer, B. Klebl, M. Falkenberg, C. M. Gustafsson, N. G. Larsson, Small-molecule inhibitors of human mitochondrial DNA transcription. *Nature* **588**, 712-716 (2020).

15. B. A. Jonas, D. A. Pollyea, How we use venetoclax with hypomethylating agents for the treatment of newly diagnosed patients with acute myeloid leukemia. *Leukemia* **33**, 2795-2804 (2019).

16. W. Zhao, K. Sachsenmeier, L. Zhang, E. Sult, R. E. Hollingsworth, H. Yang, A New Bliss Independence Model to Analyze Drug Combination Data. *J Biomol Screen* **19**, 817-821 (2014).

17. N. Chapuis, L. Poulain, R. Birsen, J. Tamburini, D. Bouscary, Rationale for Targeting Deregulated Metabolic Pathways as a Therapeutic Strategy in Acute Myeloid Leukemia. *Front Oncol* **9**, 405 (2019).

18. D. Sharon, S. Cathelin, S. Mirali, J. M. Di Trani, D. J. Yanofsky, K. A. Keon, J. L. Rubinstein, A. D. Schimmer, T. Ketela, S. M. Chan, Inhibition of mitochondrial translation overcomes venetoclax resistance in AML through activation of the integrated stress response. *Sci Transl Med* **11**, (2019).

19. B. L. Ebert, J. Pretz, J. Bosco, C. Y. Chang, P. Tamayo, N. Galili, A. Raza, D. E. Root, E. Attar, S. R. Ellis, T. R. Golub, Identification of RPS14 as a 5q- syndrome gene by RNA interference screen. *Nature* **451**, 335-339 (2008).

20. J. P. Patel, M. Gonen, M. E. Figueroa, H. Fernandez, Z. Sun, J. Racevskis, P. Van Vlierberghe, I. Dolgalev, S. Thomas, O. Aminova, K. Huberman, J. Cheng, A. Viale, N. D. Socci, A. Heguy, A. Cherry, G. Vance, R. R. Higgins, R. P. Ketterling, R. E. Gallagher, M. Litzow, M. R. van den Brink, H. M. Lazarus, J. M. Rowe, S. Luger, A. Ferrando, E. Paietta, M. S. Tallman, A. Melnick, O. Abdel-Wahab, R. L. Levine, Prognostic relevance of integrated genetic profiling in acute myeloid leukemia. *N Engl J Med* **366**, 1079-1089 (2012).

21. U. Testa, Leukemia stem cells. *Ann Hematol* **90**, 245-271 (2011).

22. M. R. Savona, J. C. Rathmell, Mitochondrial Homeostasis in AML and Gasping for Response in Resistance to BCL2 Blockade. *Cancer Discov* **9**, 831-833 (2019).

23. K. H. Lin, P. S. Winter, A. Xie, C. Roth, C. A. Martz, E. M. Stein, G. R. Anderson, J. P. Tingley, K. C. Wood, Targeting MCL-1/BCL-XL Forestalls the Acquisition of Resistance to ABT-199 in Acute Myeloid Leukemia. *Sci Rep* **6**, 27696 (2016).

24. N. P. Basak, S. Banerjee, Mitochondrial dependency in progression of acute myeloid leukemia. *Mitochondrion* **21**, 41-48 (2015).

25.   D. A. Pollyea, B. M. Stevens, C. L. Jones, A. Winters, S. Pei, M. Minhajuddin, A.
D'Alessandro, R. Culp-Hill, K. A. Riemondy, A. E. Gillen, J. R. Hesselberth, D. Abbott, D.
Schatz, J. A. Gutman, E. Purev, C. Smith, C. T. Jordan, Venetoclax with azacitidine disrupts
energy metabolism and targets leukemia stem cells in patients with acute myeloid leukemia.
Nat Med **24**, 1859-1866 (2018).

26.   A. Cole, Z. Wang, E. Coyaud, V. Voisin, M. Gronda, Y. Jitkova, R. Mattson, R. Hurren, S.
Babovic, N. Maclean, I. Restall, X. Wang, D. V. Jeyaraju, M. A. Sukhai, S. Prabha, S. Bashir,
A. Ramakrishnan, E. Leung, Y. H. Qia, N. Zhang, K. R. Combes, T. Ketela, F. Lin, W. A.
Houry, A. Aman, R. Al-Awar, W. Zheng, E. Wienholds, C. J. Xu, J. Dick, J. C. Wang, J.
Moffat, M. D. Minden, C. J. Eaves, G. D. Bader, Z. Hao, S. M. Kornblau, B. Raught, A. D.
Schimmer, Inhibition of the Mitochondrial Protease ClpP as a Therapeutic Strategy for
Human Acute Myeloid Leukemia. *Cancer Cell* **27**, 864-876 (2015).

27.   M. Skrtic, S. Sriskanthadevan, B. Jhas, M. Gebbia, X. Wang, Z. Wang, R. Hurren, Y. Jitkova,
M. Gronda, N. Maclean, C. K. Lai, Y. Eberhard, J. Bartoszko, P. Spagnuolo, A. C. Rutledge,
A. Datti, T. Ketela, J. Moffat, B. H. Robinson, J. H. Cameron, J. Wrana, C. J. Eaves, M. D.
Minden, J. C. Wang, J. E. Dick, K. Humphries, C. Nislow, G. Giaever, A. D. Schimmer,
Inhibition of mitochondrial translation as a therapeutic strategy for human acute myeloid
leukemia. *Cancer Cell* **20**, 674-688 (2011).

28.   J. R. Molina, Y. Sun, M. Protopopova, S. Gera, M. Bandi, C. Bristow, T. McAfoos, P.
Morlacchi, J. Ackroyd, A. A. Agip, G. Al-Atrash, J. Asara, J. Bardenhagen, C. C. Carrillo, C.
Carroll, E. Chang, S. Ciurea, J. B. Cross, B. Czako, A. Deem, N. Daver, J. F. de Groot, J. W.
Dong, N. Feng, G. Gao, J. Gay, M. G. Do, J. Greer, V. Giuliani, J. Han, L. Han, V. K. Henry,
J. Hirst, S. Huang, Y. Jiang, Z. Kang, T. Khor, S. Konoplev, Y. H. Lin, G. Liu, A. Lodi, T.
Lofton, H. Ma, M. Mahendra, P. Matre, R. Mullinax, M. Peoples, A. Petrocchi, J. Rodriguez-
Canale, R. Serreli, T. Shi, M. Smith, Y. Tabe, J. Theroff, S. Tiziani, Q. Xu, Q. Zhang, F.
Muller, R. A. DePinho, C. Toniatti, G. F. Draetta, T. P. Heffernan, M. Konopleva, P. Jones,
M. E. Di Francesco, J. R. Marszalek, An inhibitor of oxidative phosphorylation exploits
cancer vulnerability. *Nat Med* **24**, 1036-1046 (2018).

29.   J. Kreitz, C. Schonfeld, M. Seibert, V. Stolp, I. Alshamleh, T. Oellerich, B. Steffen, H.
Schwalbe, F. Schnutgen, N. Kurrle, H. Serve, Metabolic Plasticity of Acute Myeloid
Leukemia. *Cells* **8**, (2019).

30.   A. N. Saliba, A. J. John, S. H. Kaufmann, Resistance to venetoclax and hypomethylating
agents in acute myeloid leukemia. *Cancer Drug Resist* **4**, 125-142 (2021).

31.   I. Baccelli, Y. Gareau, B. Lehnertz, S. Gingras, J. F. Spinella, S. Corneau, N. Mayotte, S.
Girard, M. Frechette, V. Blouin-Chagnon, K. Leveille, I. Boivin, T. MacRae, J. Krosl, C.
Thiollier, V. P. Lavallee, E. Kanshin, T. Bertomeu, J. Coulombe-Huntington, C. St-Denis, M.
E. Bordeleau, G. Boucher, P. P. Roux, S. Lemieux, M. Tyers, P. Thibault, J. Hebert, A.
Marinier, G. Sauvageau, Mubritinib Targets the Electron Transport Chain Complex I and
Reveals the Landscape of OXPHOS Dependency in Acute Myeloid Leukemia. *Cancer Cell*
36, 84-99 e88 (2019).

32.   C. Bosc, M. A. Selak, J. E. Sarry, Resistance Is Futile: Targeting Mitochondrial Energetics
and Metabolism to Overcome Drug Resistance in Cancer Treatment. *Cell Metab* **26**, 705-707
(2017).

33.   M. Hosseini, H. R. Rezvani, N. Aroua, C. Bosc, T. Farge, E. Saland, V. Guyonnet-Duperat, S.
Zaghdoudi, L. Jarrou, C. Larrue, M. Sabatier, P. L. Mouchel, M. Gotanegre, M. Piechaczyk,
G. Bossis, C. Recher, J. E. Sarry, Targeting Myeloperoxidase Disrupts Mitochondrial Redox
Balance and Overcomes Cytarabine Resistance in Human Acute Myeloid Leukemia. *Cancer*
Res **79**, 5191-5203 (2019).

34.   S. Jin, D. Cojocari, J. J. Purkal, R. Popovic, N. N. Talaty, Y. Xiao, L. R. Solomon, E. R.
Boghaert, J. D. Leverson, D. C. Phillips, 5-Azacitidine Induces NOXA to Prime AML Cells
for Venetoclax-Mediated Apoptosis. *Clin Cancer Res* **26**, 3371-3383 (2020).

Acknowledgments

General:

The authors would like to thank the staff at the Department of Experimental Biomedicine at

the University of Gothenburg, namely, Jenny Nyman, Claudia Engwall, Tova Johansson and

Hanna Brissman for tremendous assistance with mouse experiments. They would also like to

thank Dr Rebekah Hailes for editing the manuscript and to acknowledge Lenka Palova-

Jelinkova, Klara Danova, and Pavla Jestrabova from SOTIO Biotech (Prague, Czech Republic)

for valuable input on *in vitro* and *in vivo* studies. The authors acknowledge support from the

National Genomics Infrastructure in Stockholm funded by Science for Life Laboratory, the

Knut and Alice Wallenberg Foundation and the Swedish Research Council, and SNIC/Uppsala

Multidisciplinary Center for Advanced Computational Science for assistance with massively

parallel sequencing and access to the UPPMAX computational infrastructure.

Funding:

Swedish Research Council grant 2017-01257 (CMG)

Swedish Cancer Foundation grant 20 1291 PjF (CMG)

Swedish Cancer Foundation grant 20 0925 PjF (LP)

Knut and Alice Wallenberg Foundation grant 2017.0080 (CMG)

The Swedish state under the agreement between the Swedish government and the county

councils, the ALF agreement grant ALFGBG-966275 (CMG)

The Swedish state under the agreement between the Swedish government and the county

councils, the ALF agreement grant ALFGBG-965206 (LP)

Assar Gabrielsson Foundation for cancer research grant (LSA)

Author contributions:

Conceptualization (LSA, RD, TB, AU, PN, BMK)

Methodology (TB, AU, PN, BMK)

Investigation (LSA, JA, CN)

Supervision (BMK, LP, CG)

Writing—original draft (LSA)

Writing—review & editing (LSA, JA, AU, RD, TB, PN, BMK, LP, CMG)

Competing interests:

CMG is a scientific co-founder of Pretzel Therapeutics Inc. Together with RD, TB, AU, PN,

BMK who are employees of LDC, CMG is also co-inventor of the patent application WO

2019/057821. Medicinal chemistry and pharmacology part of this work was financed by the

Max-Planck Gesellschaft e.V. under the framework agreement between Max-Planck and LDC.

Data and materials availability:

All data needed to evaluate the conclusions in the paper are present in the paper and/or the

Supplementary Materials. RNA-seq datasets have been deposited in the GSE repository

(GSE213976). Additional data related to this paper may be requested from the authors.

Figure 1

Figure 1. IMT+venetoclax combination treatment synergistically inhibits cell

viability/growth of AML cell lines *in vitro*.

(**A**) Chemical structure of IMT (LDC204857) (**B**) AML cell lines were treated with single

agents IMT or venetoclax at a range of concentrations (0.1–1000 nM and 0.01–100 nM,

respectively), and assessed by cell viability tests after three and six days. (**C**) Synergism

analysis of AML cell lines treated with a combination of IMT (0.05, 0.1, 0.2 μM for MV4-11

and MOLM-13, and 0.1, 0.5, and 1 μM for THP-1 and KG-1) and venetoclax (Vex; 1, 2, 4 nM

for MV4-11 and MOLM-13, and 1, 5, 10 nM for THP-1 and KG-1) for six days. 'S' represents

synergy of the combination treatment, evaluated using the Bliss independent model.

Proliferation percentages are normalized to DMSO treated. All results are from three

independent experiments.

Figure 2

A

B

Figure 2. Combination treatment significantly enhances apoptosis induction in AML cell

lines.

AML cell lines were treated with IMT (500 nM) and venetoclax (5 nM) for 72–96 h. (**A**)

Apoptosis was determined by Annexin V/PI staining of the cells and analyzed by flow

cytometry. (**B**) Loss of mitochondrial membrane potential was calculated as the percentage of

depolarized (apoptotic) cells out of the total cell population. Error bars represent standard error

of the mean where * represents $P < 0.05$, **$P < 0.01$, and ***$p < 0.001$.

Figure 3

Figure 3. Combination of IMT and venetoclax dramatically decreases OXPHOS (in a

glycolysis-independent manner) in sensitive cell lines.

(**A**) Representative graph of an OXPHOS measurement using mitostress compounds:

oligomycin (OMY), carbonyl cyanide m-chlorophenyl hydrazone (CCCP), and antimycin

(AMA). (**B**) AML cell lines (MV4-11, MOLM-13, and THP-1) were treated with either IMT

(500 nM), venetoclax (5 nM), or a combination of both for four days. DMSO was added to

control wells. Oxygen consumption rate (OCR) was measured with an Oroboros O2K

instrument at physiological glucose levels (5 mM) using mitostress compounds. (**C**) AML cell

lines were cultured in growth medium supplemented with either 5 mM glucose or galactose,

and cell viability tests (ATP measurements) were performed after 24 and 48 h. All results are

from at least three independent experiments. Error bars show ± SEM. Statistical tests on all

comparisons in (C) showed non-significant differences.

Figure 4

A

B

C

D

Figure 4. Combination of IMT and venetoclax significantly prolongs the survival of mice

with leukemia.

(**A**) Outline of CDX efficacy experiment: NBSGW mice were transplanted with MV4-11

human leukemia cells, then treated with either vehicle, IMT, venetoclax, or a combination of

IMT and venetoclax (n=10 per treatment group). Peripheral blood was taken before treatment

(week 2) and after treatment at regular intervals until disease progression was observed. (**B**)

Analysis of proportion of human CD45+ cells in mice peripheral blood in all treatment groups

at indicated time points, determined by flow cytometry. (**C**) Kaplan-Meier analysis of survival.

Statistical significance was calculated using log-rank (Mantle-Cox) test. ****$p < 0.0001$, Error

bars represent ± SEM. (**D**) Analysis of proportion of human CD45+ cells in the bone marrow

of all treatment groups at time of euthanization.

Figure 5

Anti-hCD45 Nucleus

Figure 5. Synergistic effect of (IMT and venetoclax) combination treatment on tumor

burden and survival of AML-PDX transplanted mice.

(**A**) Experimental outline: NSGS mice were transplanted with primary AML splenocytes from

PDX mice and were then treated with either vehicle, IMT, venetoclax, or a combination of IMT

and venetoclax (n=10 per treatment group). After the final dosing, three or four mice were

euthanized (end-of-treatment group) and the rest (n=6) were maintained for survival analysis.

(**B**) Percentage of viable human CD45-positive (hCD45+) cells in peripheral blood, BM, and

spleen of treated mice determined by flow cytometry. (**C**) Immunohistochemistry analysis of

mouse BM and spleen sections stained with anti-human CD45 (brown) and counterstained with

DAB (blue). (**D**) Quantitative analysis of hCD45+ cells on immunohistochemistry sections of

BM and spleen sections of treated mice. (**E**) Proliferating tumor cells (CD45) are shown by

quantification of Ki67 positive cells in immunohistochemistry-stained BM sections at end-of-

treatment (results from at least three technical replicates and two biological replicates in each

treatment group).

A

B

Figure 6. Synergistic effect of (IMT and venetoclax) combination treatment on gene

transcription and survival of AML-PDX transplanted mice.

The experimental outline was shown in figure 5A, but n=10 per treatment group. After the final

dosing, three or four mice were euthanized (end-of-treatment group) and the rest (n=6) were

maintained for survival analysis. (**A**) Gene expression profiling by RNA-seq of purified AML

blasts isolated from PDX models treated with venetoclax, IMT, or a combination thereof.

Indicated expression changes are relative to a vehicle treated group. Significantly enriched

genes are indicated in red (q < 0.05). (**B**) Peripheral blood engraftment analysis of survival

group. (**C**) Survival analysis of treated mice shown by Kaplan-Meier analysis (n=6/treatment

group). *P < 0.05, **P < 0.01, and ***p < 0.001.

A

B

C

D

Figure 7. Leukemia progression induced by primary AML cells is inhibited by

venetoclax treatment.

As in Fig. 5A, efficacy experiment outline: NSGS mice were transplanted with primary AML

splenocytes from PDX mice and were then treated with either vehicle, IMT, venetoclax, or a

combination of IMT and venetoclax (n=12 per treatment group). 8–12 h after the last dosing,

half of the mice (n=6/treatment group) were euthanized (end-of-treatment group) and the

other half were maintained for survival analysis. (**A**) Percentage of viable human CD45-

positive (hCD45+) cells in peripheral blood and bone marrow of treated mice (n=6 for each

treatment group), determined by flow cytometry. (**B**) Immunohistochemistry analysis of

hCD45+ cells on bone marrow and spleen tissue sections of treated mice (images are

representative from at least two mice per group). Scale bars represent 100 μM. (C) Peripheral

blood engraftment analysis of survival group. (**D**) Mice treated with single or a combination

of compounds showed prolonged survival in comparison to the vehicle-treated group.

Kaplan-Meier analysis shows the relative survival curve. *P < 0.05, **P < 0.01, and ***p <

0.001.