A pan-variant mRNA-LNP T cell vaccine protects HLA transgenic mice from mortality after infection with SARS-CoV-2 Beta

Brandon Carter[1,10,+], Pinghan Huang[2,+], Ge Liu[1,10], Yuejin Liang[2], Paulo J.C. Lin[5], Bi-Hung Peng[3], Lindsay McKay[6], Alexander Dimitrakakis[1,10], Jason Hsu[3], Vivian Tat[4], Panatda Saenkham-Huntsinger[2], Jinjin Chen[7], Clarety Kaseke[8], Gaurav D. Gaiha[8,9], Qiaobing Xu[7], Anthony Griffiths[6], Ying K. Tam[5], Chien-Te K. Tseng[2,3,4,*], David K. Gifford[1,10,11,*]

1. MIT Computer Science and Artificial Intelligence Laboratory, Cambridge, MA, USA
2. Department of Microbiology and Immunology, The University of Texas Medical Branch, Galveston, TX, USA
3. Department of Neuroscience, Cell Biology, and Anatomy, The University of Texas Medical Branch, Galveston, TX, USA
4. Department of Pathology, The University of Texas Medical Branch, Galveston, TX, USA
5. Acuitas Therapeutics, Vancouver, BC, Canada
6. Boston University's National Emerging Infectious Diseases Laboratories, Boston, MA, USA
7. Department of Biomedical Engineering, Tufts University, Medford, MA, USA
8. Ragon Institute of MGH, MIT, and Harvard, Cambridge, MA, USA
9. Division of Gastroenterology, Massachusetts General Hospital, Boston, MA, USA
10. MIT Electrical Engineering and Computer Science, Cambridge, MA, USA
11. MIT Biological Engineering, Cambridge, MA, USA

* Correspondence to: sktseng@utmb.edu, gifford@mit.edu
+ These authors contributed equally to this work.

Abstract

Clinically licensed COVID-19 vaccines ameliorate viral infection by inducing vaccinee production of neutralizing antibodies that bind to the SARS-CoV-2 Spike protein to inhibit viral cellular entry (Walsh et al., 2020; Baden et al., 2021), however the clinical effectiveness of these vaccines is transitory as viral variants arise that escape antibody neutralization (Tregoning et al., 2021; Willett et al., 2022). Vaccines that solely rely upon a T cell response to combat viral infection could be transformational because they can be based on highly conserved short peptide epitopes that hold the potential for pan-variant immunity, but a mRNA-LNP T cell vaccine has not been shown to be sufficient for effective antiviral prophylaxis. Here we show that a mRNA-LNP vaccine based on highly conserved short peptide epitopes activates a CD8+ and CD4+ T cell response that prevents mortality in HLA-A*02:01 transgenic mice infected with the SARS-CoV-2 Beta variant of concern (B.1.351). In mice vaccinated with the T cell vaccine, 24% of the nucleated cells in lung were CD8+ T cells on day 7 post infection. This was 5.5 times more CD8+ T cell infiltration of the lungs in response to infection compared to the Pfizer-BioNTech

Comirnaty® vaccine. Between days 2 and 7 post infection, the number of $CD8^+$ T cells in the

lung increased in mice vaccinated with the T cell vaccine and decreased in mice vaccinated with

Comirnaty®. The T cell vaccine did not produce neutralizing antibodies, and thus our results

demonstrate that SARS-CoV-2 viral infection can be controlled by a T cell response alone. Our

results suggest that further study is merited for pan-variant T cell vaccines, and that T cell

vaccines may be relevant for individuals that cannot produce neutralizing antibodies or to help

mitigate Long COVID.

Main

Current strategies for COVID-19 vaccine design utilize one or more SARS-CoV-2 Spike protein

subunits to primarily activate the humoral arm of the adaptive immune response to produce

neutralizing antibodies to the Spike receptor binding domain (RBD) (Walsh et al., 2020; Baden

et al., 2021).   Vaccination to produce neutralizing antibodies is a natural objective, as

neutralizing antibodies present an effective barrier to the viral infection of permissive cells by

binding to the RBD and thus blocking cellular entry via the ACE2 receptor.   However, the

strategy of focusing on Spike as the sole vaccine target has proven problematic as Spike rapidly

evolves to produce structural variants that evade antibody-based acquired immunity from

vaccination or infection with previous viral variants (Tregoning et al., 2021; Willett et al., 2022).

Compared to the original Wuhan variant, novel viral variants are arising that are more

contagious (Zhang et al., 2021), and infectious to a broader range of host species (Shuai et al.,

2021).   Thus, vaccine designers are pursuing a stream of novel Spike variant vaccines.

Multivalent Spike vaccines and bivalent booster vaccines provide protection against multiple

known variants of concern (VOC) of SARS-CoV-2 but are not necessarily protective against

unknown future variants (Martinez et al., 2021).   Mosaic RBD nanoparticles that display

disparate SARS-CoV RBDs have been found to produce effective neutralizing antibodies

against both SARS-CoV-1 and SARS-CoV-2 (Cohen et al., 2022), but the robustness of mosaic

RBD protection against possible future Spike mutations depends upon conserved Spike

structural epitopes. In contrast, the vaccine approach we present here depends upon

conserved T cell epitopes drawn from the entire viral proteome for protection against future

variants.

We show in this study that an immunogenic T cell vaccine with 11 conserved short MHC

displayed epitopes ("MIT-T-COVID vaccine") provides effective prophylaxis against the onset of

SARS-CoV-2-induced morbidity and mortality caused by SARS-CoV-2 Beta infection in

transgenic mice carrying human HLA-A*02:01. We demonstrate the MIT-T-COVID vaccine

causes significant infiltration of CD8$^+$ and CD4$^+$ T lymphocytes in the lungs. Multiple designs for

T cell vaccines for SARS-CoV-2 have been proposed (Liu et al., 2020; Nathan et al., 2021;

Heitmann et al., 2022; Pardieck et al., 2022) and are in clinical trials (NCT05113862,

NCT0488536, NCT05069623, NCT04954469), but identifying the mechanisms behind the

efficacy of pure T cell vaccines remains an open question.   Substantial literature suggests that

T cell responses are integral to the adaptive immunity to COVID-19 (Moss, 2022).   For

example, a study that ablated the B cell compartment of the immune system in Spike vaccinated

mice found that CD8$^+$ T cells alone can control viral infection (Israelso et al., 2021). Pardieck et

al. (2022) found that vaccination with a single CD8$^+$ T cell epitope conferred protection against

mortality from the Leiden-0008/2020 SARS-CoV-2 variant in K18-hACE2 transgenic mice, but

unlike our study, required three doses for efficacy, did not engage a CD4$^+$ T cell response, and

did not identify significant T cell infiltration of the lungs. In addition to specific antibody

responses, COVID-19 vaccinees and convalescent patients possess SARS-CoV-2 specific

CD8$^+$ and CD4$^+$ T cells, suggesting the contribution of the T cell compartment to the adaptive

immunity to COVID-19 (Sekine et al., 2020), and clinical findings have revealed vaccine-induced

T cell responses in B cell-deficient patients (Shree, 2022).   It has also been reported that

vaccination by WA (Wuhan) Spike in a mouse model failed to produce antibodies fully capable

of neutralizing the SA (Beta) variant of CoV-2, yet immunized mice were protected against Beta

strain challenge, leading to the conclusion that T cell responses were sufficient to control Beta

SARS-CoV-2 (Kingstad-Bakke et al., 2022).   In addition, vaccination with T cell epitope-rich

Nucleocapsid protein produced specific T cell responses thought to be causally associated with

viral control (Matchett et al., 2021).   Intranasal vaccination of mice with SARS-CoV-1

Nucleocapsid followed by challenge with 10^4 Plaque-forming units of SARS-CoV-1 prevented

mortality in 75% of the mice (Zhao et al, 2016). Combined Spike and Nucleocapsid vaccination

improved viral control compared to Spike vaccination alone in preclinical models, while CD8$^+$ T

cell depletion demonstrated the role of CD8$^+$ T cells in viral control and protection from weight

loss (Hajnik et al., 2022).

Vaccine Design

We designed a COVID-19 T cell vaccine using T cell epitopes that are unchanged over 19

presently known SARS-CoV-2 variants of concern (VOC) (Methods).   Since T cell epitopes can

originate from any part of the viral proteome, they can be drawn from portions of the proteome

that are likely to be evolutionarily stable.   The prediction of stable epitopes can be

accomplished by historical analysis of thousands of viral variants (Liu et al., 2020), structural

analysis (Nathan et al., 2021), or the functional analysis of mutations lethal to the virus.

Beginning with stable epitopes as candidates, our T cell vaccine design proceeds by vaccine

epitope selection that optimizes population coverage where every vaccinated individual is

predicted to experience on average multiple immunogenic peptide-HLA hits (Liu et al., 2021; Liu

et al., 2022).

The MIT-T-COVID vaccine realizes an *n-times* coverage objective by encoding multiple

epitopes for each target MHC class I and II diplotype to (1) expand diverse sets of T cell

clonotypes to fight viral infection, (2) accommodate variations in epitope immunogenicity

between vaccinees, and (3) reduce the chances that viral evolution will lead to immune system

escape (Liu et al., 2022).   The MIT-T-COVID vaccines consists of eight MHC class I epitopes

and three MHC class II epitopes (Figure 1A, Methods, Supplemental Table 1).   The MHC class

I and II vaccine peptides are encoded into a single mRNA construct for delivery with the same

Acuitas LNP delivery platform that is used by the Pfizer-BioNTech Comirnaty® vaccine (Figure

1A).   The eight MIT-T-COVID MHC class I epitopes are the HLA-A*02:01 subset of the MHC

class I *de novo* MIRA only vaccine design of Liu et al. (2021) that used combinatorial

optimization to select vaccine epitopes to maximize *n-times* population coverage over HLA

haplotype frequencies (Liu et al., 2022).   For inclusion in the assembled construct, the eight

MHC class I vaccine peptides were randomly shuffled, and alternate peptides were flanked with

five additional amino acids at each terminus as originally flanked in the SARS-CoV-2 proteome.

Selected epitopes were flanked to test if flanking enhanced or impaired epitope presentation.

The three MHC class II epitopes were selected by considering all SARS-CoV-2 proteome

windows of length 13-25 and selecting conserved peptides that are predicted to be displayed by

the mouse MHC class II allele H-2-IAb (Methods).   The mRNA construct encodes a secretion

signal sequence at the N-terminus and an MHC class I trafficking signal (MITD) at the C-

terminus (Kreiter et al., 2008). Peptide sequences are joined by non-immunogenic

glycine/serine linkers (Sahin et al., 2017). The construct also included control peptides for HLA-

A*02:01 and H-2-IAb (CMV pp65: NLVPMVATV for HLA-A*02:01 and Human CD74:

KPVSKMRMATPLLMQAL for H-2-IAb) (Vita et al., 2018).

Challenge Study

We vaccinated with three test articles: a negative control injection (PBS), the Pfizer-BioNTech

Comirnaty® vaccine (wastage that was refrozen and then thawed), and the MIT-T-COVID

vaccine (Figure 1B). We vaccinated HLA-A*02:01 human transgenic mice with these three

vaccines, vaccinating 11 age-matched male mice with each vaccine. Comirnaty® or MIT-T-

COVID vaccinations contained of 10 µg of mRNA. The mice were vaccinated at Day 0 and

boosted at Day 21 (Methods). At Day 35 three mice from each group were sacrificed for

immunogenicity studies, and at Day 37 the remaining eight mice were challenged intranasally

(I.N.) with 5 x 10^4 $TCID_{50}$ of the SARS-CoV-2 B.1.351 variant. Challenged mice were subjected

to daily monitoring for the onset of morbidity (i.e., weight changes and other signs of illness) and

any mortality. At Day 39 (2 days post infection) three mice were taken from each group and

sacrificed to determine viral burdens and to perform lung histopathology. At Day 44 (7 days

post infection) the remaining mice were sacrificed to determine viral burdens and to perform

lung histopathology.

Immunogenicity of vaccines

Vaccination by the MIT-T-COVID vaccine expanded CD8$^+$ and CD4$^+$ T cells that expressed

interferon gamma (IFN-γ) or tumor necrosis factor alpha (TNF-α) when queried by vaccine

epitopes (Figures 2A-B). The observed variability of immunogenicity of MIT-T-COVID epitopes

in convalescent COVID patients (Supplemental Table 1) and in the present study supports our

use of multiple epitopes per MHC diplotype for *n-times* coverage. Vaccination by Comirnaty®

produced no significant T cell responses to vaccine epitopes when compared to PBS. CD8$^+$ T

cells that are activated by the CD8-4 epitope (YLQPRTFLL, Spike 269-277) are expanded in

animals vaccinated with the MIT-T-COVID vaccine (IFN-γ 1.316% ± 0.529% of CD8$^+$ T cells, *P*

= 0.0087 vs. PBS; TNF-α 0.380% ± 0.086% of CD8$^+$ T cells, *P* = 0.0149 vs. PBS; Figure 2A).

Similarly, the CD8-8 epitope (TVYSHLLLV, ORF3a 89-97) activated CD8$^+$ T cells that are

expanded by the MIT-T-COVID vaccine (IFN-γ 0.560% ± 0.021% of CD8$^+$ T cells, *P* = 0.001 vs.

PBS; TNF-α 0.250% ± 0.125% of CD8$^+$ T cells, *P* = 0.015 vs. PBS) and the CD4-2 epitope

(RFYFYTSKTTVASLIN, ORF1ab 1421-1436) activated CD4$^+$ T cells are expanded by the MIT-

T-COVID vaccine (TNF-α 0.078% ± 0.027%, *P* = 0.058 vs. PBS, *P* = 0.010 vs. Comirnaty®,

IFN-γ not significant, Figure 2B). The lack of HLA-A*02:01 transgenic animal response to

certain CD8$^+$ epitopes that were immunogenic in patients (Supplemental Table 1) is consistent

with past studies of transgenic mouse models (Kotturi et al., 2009) and the variability of

immunogenicity of epitopes between in-bred mice (Croft et al., 2019).

T cells that lack IFN-γ responses can produce effective immune mediators (Nakiboneka et al.,

2019). We found that the fraction of CD4$^+$ T cells that are FoxP3$^+$ were not expanded in

Comirnaty® or MIT-T-COVID vaccinated animals, suggesting that fraction of T regulatory cells

that could induce tolerance were not expanded by vaccination (Figure 2C).

Vaccine efficacy against the onset of morbidity and mortality and the viral loads in

challenged mice

Upon viral challenge, both PBS and MIT-T-COVID vaccinated animals exhibited a more than

10% weight loss by Day 3 (PBS mean weight reduction 11.386% ± 1.688%, MIT mean weight

reduction 10.851% ± 0.641%), with the MIT-T-COVID vaccinated animals beginning to recover

from Day 4 onward (Figure 3A). The weight phenotype was mirrored by the clinical score

phenotype, with PBS animals not recovering and MIT-T-COVID vaccinated animals improving

from Day 5 onward (Figure 3B, Methods). The Comirnaty® vaccine protected animals from

significant weight loss and poor clinical scores.

When the yields of infectious progeny virus were measured at 2 days post infection (dpi), we

noted that mice immunized with MIT-T-COVID vaccine had 6.706 ± 0.076 log10 TCID$_{50}$/g,

compared to 7.258 ± 0.367 log10 TCID$_{50}$/g in the PBS control, representing a moderate 3.6-fold

reduction in viral replication (P = 0.046; Figure 3D). In contrast, mice immunized with

Comirnaty® had infectious viral titers that were below the detection limits at 2 dpi. Infectious

viral titer was not significant for all test articles at 7 dpi (Figure 3D). Lung viral mRNA levels

measured by qPCR followed the same trends as infectious virus progeny in the lungs, with

Comirnaty® showing no significant levels. Despite a reduced content of total and sub-genomic

viral RNAs associated with MIT-T-COVID vaccine samples as assessed by qPCR, the reduction

was insignificant, compared to those of PBS control (Supplemental Figure 1).

All the Comirnaty® and MIT-T-COVID vaccine animals survived to 7 dpi, when the study was

terminated, with the difference in clinical scores of these two vaccine groups becoming

insignificant (Figure 3C). In contrast, four of five PBS control animals had been euthanized

because of weight loss > 20%.   The survival of all five animals vaccinated with Comirnaty®

and the MIT-T-COVID vaccine was significant vs. PBS (P = 0.0053, logrank test).

We noted that mice immunization with the Comirnaty® vaccine elicited substantial specific

antibodies capable of completely neutralizing Beta and WA-1 variants of CoV-2 with 100%

neutralizing titers (NT_{100}) of 896 ± 350 and fold 2048 ± 1887, respectively (Figure 3E). The

higher neutralizing titer against WA-1 is expected as it is variant matched to Comirnaty®.   As

expected for a peptide-based vaccine, no neutralizing titer could be readily detected in the

serum from mice immunized with MIT-T-COVID vaccine or PBS.   Total specific IgG/IgM

antibodies were also measured by ELISA against a cell lysate prepared from WA-1-infected

Vero E6 cells.   Comirnaty® has a log10 IgG/IgM titer of 4.915 ± 0.213, while the log10 titer

produced by PBS was 2.452 ± 0.321 and the MIT-T-COVID vaccine produced a log10 titer of

2.266 ± 0.291 (Figure 3F).   The low but detectable titers in PBS and MIT-T-COVID vaccine-

immunized mice may represent an early IgM response to viral infection.

All lung samples were subjected to immunohistochemistry (IHC) staining for SARS-CoV-2 spike

protein (Supplemental Figure 2). We found specimens vaccinated with PBS exhibited extensive

staining indicative of viral infection throughout the epithelium of both the bronchioles and the

alveolar sacs, with the viral infection appearing more intense at 2 dpi. Although viral infection is

significantly reduced by 7 dpi, viral antigen was still readily detectable throughout alveoli. In

comparison, specimens vaccinated with the MIT-T-COVID vaccine exhibited similarly extensive

viral infection at 2 dpi throughout the bronchiolar and alveolar epithelia, albeit somewhat

reduced in intensity. However, by 7 dpi, viral infection was significantly reduced in both extent

and intensity, with brown puncta being detected only in a few alveoli scattered throughout the

tissue. Contrasted with both PBS and MIT-T-COVID vaccinated specimens, the Comirnaty®-

vaccinated specimens exhibited significantly reduced viral infections at both 2 and 7 dpi. Apart

from a single area at 7 dpi (see Supplemental Figure 3), viral antigen was undetected at both

timepoints in Comirnaty® vaccinated animals.

Paraffin-embedded and H&E-stained lung specimens of differentially immunized mice,

harvested at 2 and 7 dpi, were subjected to histopathological examination (Supplemental Figure

4). We found that at 7 dpi mice immunized with MIT-T-COVID vaccine exhibited extensive

lymphocytic infiltrations in perivascular regions and spaces from around bronchi, bronchioles, to

alveoli. Fewer infiltrations were found in mice immunized with either Comirnaty® or PBS.

Additionally, these infiltrations only localized at perivascular regions around bronchi and large

bronchioles. Despite the less intensive and localized inflammatory infiltrates, we also noted

widespread congestion, hemorrhage, and few foci of thromboembolism were exclusively

observed within the lungs of Comirnaty®-immunized mice but not others (Supplemental Figure

4). We also noted the lung histopathology was milder but the same pattern at 2 dpi than those of

7 dpi (data not shown).

Lung specimens were subjected to IHC staining for CD8$^+$ and CD4$^+$ cells at both 2 and 7 dpi

(Figure 4; Supplemental Figures 5-7). At 2 dpi, we found a significant increase in CD8$^+$ T cells

infiltrating the lungs in mice immunized with MIT-T-COVID (12.6% ± 5.91% of all nucleated cells

were CD8$^+$) or Comirnaty® (12.4% ± 2.35%) compared to mice immunized with PBS (PBS

mean 4.48% ± 1.99%; P = 0.044 vs. MIT-T-COVID, P = 0.050 vs. Comirnaty®; Figure 4A).

At 7 dpi, we found a significant increase in CD8$^+$ T cells infiltrating the lungs in mice immunized

with MIT-T-COVID (24.0% ± 5.21% of all nucleated cells were CD8$^+$) compared to mice

immunized with Comirnaty® (Comirnaty® mean 4.35% ± 2.20%, P = 0.001) or PBS (PBS mean

7.32% ± 4.88%, P = 0.001; Figure 4A). We observed a significant increase in CD8$^+$ T cells

infiltrating the lungs in mice immunized with MIT-T-COVID between 2 and 7 dpi (P = 0.0039),

and a significant decrease in CD8$^+$ T cells infiltrating the lungs in mice immunized with

Comirnaty® between 2 and 7 dpi (P = 0.044; Figure 4A). At 7 dpi, we also found a significant

increase in CD4$^+$ T cells in filtrating the lungs in mice immunized with MIT-T-COVID (7.09% ±

4.05%) compared to mice immunized with PBS (PBS mean 0.41% ± 0.39%; P = 0.0062; Figure

4B).

Discussion

Existing Spike vaccines produce a T cell response that is thought to be important for vaccine

effectiveness and durability (Moss, 2022).   However, the T cell response induced by a Spike

vaccine may be less effective in promoting durable immune responses than the T cell response

induced by a pure T cell vaccine. Spike based T cell responses may contain epitopes that are

subject to evolutionary change (de Silva et al., 2021; Redd et al., 2021), or Spike epitopes may

not be as well presented or immunogenic as epitopes drawn from a more diverse set of SARS-

CoV-2 proteins.   Thus, T cell augmentation strategies for Spike vaccines (Liu et al., 2021) and

pure T cell vaccines will require further exploration to unravel the hierarchy of immune

responses to their components, and how to structure a vaccine for optimal prophylaxis.   In

patients with impaired antibody responses, T cell vaccines would eliminate the burden of non-
immunogenic B cell epitopes (Benjamini et al., 2022) and provide immune protection, at least to
a certain extent, against infection.
284
285 The marginal IgG/IgM antibody titers that were not neutralizing elicited by mice immunized with
286 the MIT-T-COVID vaccine indicates that the protective mechanism of the vaccine was likely
287 based on a T cell response to the virus. The reduction in viral titer on day 2 shows that T cell
288 responses were present early in infection.   In the absence of neutralizing antibodies these
289 responses were sufficient to rescue the vaccinated mice from the onset of mortality.
290
We expect that a T cell vaccine would provide prophylaxis similar to what we have described
against future SARS-CoV variants and strains that conserve the vaccine's epitopes.   We chose
Beta (B.1.351) for our challenge study as it has a severe phenotype in a wild-type mouse
background (Shuai et al., 2021).   Transgenic human ACE2 (hACE2) mice exhibit encephalitis
post COVID infection that does not represent human pathology (Kumari et al., 2021) and thus
may be a poor model for T cell vaccine effectiveness. Instead, we chose to evaluate the HLA-
A*02:01 component of our vaccine design in a HLA-A*02:01 transgenic animal model given the
predominance of HLA-A02 in the human population (Ellis et al., 2000).
299
The MIT-T-COVID vaccine induced a response where circulating T cells migrated rapidly and
efficiently into the lung upon viral challenge, as evidenced by more intense and widespread
lymphocytic infiltrations than other groups. Further, the degree of CD8$^+$ T cell infiltration of the
lungs significantly increased in mice vaccinated with the MIT-T-COVID vaccine between days 2
and 7 post infection, and significantly decreased in mice vaccinated with the Comirnaty®
vaccine. The lack of this T cell response in Comirnaty® vaccinated animals merits further study.
306
Another finding is that widespread congestion, hemorrhage, and few foci of thromboembolism
were exclusively observed within the lungs of Comirnaty®-immunized mice. Although this
finding is unrelated to the MIT-T-COVID vaccine, it suggests the possibility of immunization-
induced side-effects or immunopathology of mRNA-based CoV-2 vaccines. The exact
mechanism of pulmonary embolism in this case remains unknown for now and should be
explored.
313

Cytotoxic T cells (CD8$^+$ T cells) have the ability to kill virally infected cells, and thus T cell

vaccines might provide long-term immunity and be effective therapeutics for Long COVID.

Post-acute sequelae of COVID-19 (PASC, "Long COVID") causes persistent symptoms in ~10%

of people past twelve weeks of COVID infection (WHO, 2021).   Long COVID has been

associated with the continued presence of Spike in the blood, suggesting that a tissue based

viral reservoir remains in Long COVID patients (Swank et al., 2022).

Our results suggest that if one goal of a vaccine is protection against novel viral strains it may

be appropriate to develop vaccines that permit symptomatic infection but protect against severe

illness.   Current regulatory criteria for licensing vaccines are solely based upon the prevention

of symptomatic illness for vaccine matched viral strains.   Further research on T cell vaccines

may reveal novel vaccine designs that prevent severe illness while providing protection against

viral variants as an additional tool in the fight against global pandemics.

A

```
MRVTAPRTLILLLSGALALTETWAGS G G S G G G G S G G YLYALVYFL G G S G G G G S G G RSKNPLLYDANYFLCWHTN
G G S G G G G S G G FVDGVPFVV G G S G G G G S G G AYYVGYLQPRTFLLKYNEN G G S G G G G S G G
FLNRFTTTL G G S G G G G S G G RLTKYTMADLVYALRHFDE G G S G G G G S G G SIIAYTMSL G G S G G G G S G G
LLLFVTVYSHLLLVAAGLE G G S G G G G S G G ATSRTLSYY G G S G G G G S G G KTFPPTEPK G G S G G G G S G G
EEIAIILASFSASTSAFVETVKGLDY G G S G G G G S G G KSILSPLYAFASEAARVVRSIFSRTL G G S G G G G S G G
VDYGARFYFYTSKTTVASLINTLNDLGGSGGGGSGGNLVPMVATVGGSGGGGSGGKPVSKMRMATPLLMQALGGSLGGGG
SGIVGIVAGLAVLAVVVIGAVVATVMCRRKSSGGKGGSYSQAASSDSAQGSDVSLTA
```

331 **B**

335 **Figure 1.** (A) Assembled vaccine construct containing a secretion signal sequence (red),

336 peptides (bold) joined by non-immunogenic glycine/serine linkers, and an MHC class I trafficking

337 signal (blue). (B) Study design.

A

B

C

Figure 2. Vaccine immunogenicity. (A) CD8[+] T cell responses, (B) CD4[+] T cell responses, (C) FoxP3[+] T Regulatory cells as a percentage of all CD4[+] cells. The CD8 pool includes MHC class I peptides CD8-1—CD8-8 (Supplemental Table 1). The CD4 pool includes MHC class II peptides CD4-1 and CD4-2. The Pep pool includes all query peptides in Supplemental Table 1. Error bars indicate the standard deviation around each mean. *P* values were computed by one-way ANOVA with Tukey's test. *$P < 0.05$, **$P < 0.01$.

Figure 3. Study phenotypic data, lung viral titer, and vaccine antibody responses. (A) Weights vs. days post infection, (B) clinical scores vs. days post infection, (C) Kaplan-Meier mortality curve (mortality at 80% weight loss), (D) lung viral titer, (E) maximum serum dilution that provided 100% neutralization of viral infection *in vitro*, (F) IgG/IgM titer measured by ELISA against cell lysate infected with WA-1 SARS-CoV-2. Dotted lines in (D)-(F) indicate assay limits of detection. Error bars indicate the standard deviation around each mean. *P* values were computed by one-way ANOVA with Tukey's test except (C) *P* values were computed using the logrank test. **P* < 0.05 and ***P* < 0.01.

Figure 4. Lung immunohistochemistry for CD8⁺ and CD4⁺ cells. Counts of (A) CD8⁺ and (B) CD4⁺ T cells in expressed as a percentage of all nucleated cells visible in each field from lung tissue. Example CD8⁺ stain images at 7 dpi for (C) MIT-T-COVID, (D) Pfizer/BNT, and (E) PBS vaccinated animals. Lung samples were subjected to IHC staining for CD8 (brown) with hematoxylin counterstain (blue). Images were taken at 10x magnification. Red outlines in (C)-(E) indicate CD8⁺ cells identified and counted by CellProfiler software (Methods). Error bars indicate the standard deviation around each mean. P values were computed by two-way ANOVA with Tukey's test. *$P < 0.05$ and **$P < 0.01$. See also Supplemental Figures 5-7.

Methods

Mice

A total of 33 male HLA-A*02:01 human transgenic mice (Taconic 9659) were used that were between 12 and 16 weeks old when they received their initial vaccination.   These CB6F1 background mice carry the MHC class I alleles HLA-A*02:01, H-2-Kb, H-2-Db, H-2-Kd, and H-2-Dd, and H-2-Ld. The mice carry the MHC class II alleles H-2-IAd, H-2-IEd, and H-2-IAb.

Tissue culture and virus

Vero E6 cells (ATCC, CRL:1586) were grown in minimum essential medium (EMEM, Gibco) supplemented with penicillin (100 units/mL), streptomycin (100 µg/mL) and 10% fetal bovine serum (FBS). Two strains of SARS-CoV-2 were used in this study, SARS-CoV-2 (US_WA-1/2020 isolate) and Beta (B.1.351/SA, Strain: hCoV-19/USA/MD-HP01542/2021). Both viruses were propagated and quantified in Vero E6 cells and stored at -80 °C until needed.

MIT-T-COVID vaccine design

MHC Class I epitopes. Eight peptides from SARS-CoV-2 were selected as a subset of the MHC class I *de novo* MIRA only vaccine design of Liu et al., 2021.   We filtered this set of 36 peptides to the 8 peptides predicted to be displayed by HLA-A*02:01 by a combined MIRA and machine learning model of peptide-HLA immunogenicity (Liu et al., 2021). The combined model predicts which HLA molecule displayed a peptide that was observed to be immunogenic in a MIRA experiment, and uses machine learning predictions of peptide display for HLA alleles not observed or peptides not tested in MIRA data. Thus, all eight MHC class I peptides in our vaccine were previously observed to be immunogenic in data from convalescent COVID-19 patients (Snyder et al., 2021). We further validated that all peptides are predicted to bind HLA-A*02:01 with high (less than or equal to 50 nM) affinity using the NetMHCpan-4.1 (Reynisson et al., 2020) and MHCflurry 2.0 (O'Donnell et al., 2020) machine learning models. For inclusion in the assembled construct, the eight vaccine peptides were randomly shuffled, and alternate peptides were flanked with five additional amino acids at each terminus as originally flanked in the SARS-CoV-2 proteome.   The CD4-2 vaccine epitope produced a CD8$^+$ response, and this may be a consequence of a contained MHC class I epitope, with candidates including

SLINTLNDL (HLA-A*02:01 predicted binding 55 nM), ASLINTLNDL (H-2-Db predicted binding

289 nM), FYTSKTTVASL (H-2-Kd predicted binding 293 nM), FYFYTSKTTV (H-2-Kd predicted

binding 40 nM), and YFYTSKTTV (H-2-Kd predicted binding 56 nM).

<u>MHC Class II epitopes</u>. Three peptides from SARS-CoV-2 were optimized for predicted binding

to H-2-IAb. We scored all SARS-CoV-2 peptides of length 13-25 using the sliding window

approach of Liu et al. (2020) and a machine learning ensemble that outputs the mean predicted

binding affinity (IC50) of NetMHCIIpan-4.0 (Reynisson et al., 2020) and PUFFIN (Zeng et al.,

2019). We selected the top three peptides by predicted binding affinity using a greedy selection

strategy with a minimum edit distance constraint of 5 between peptides to avoid selecting

overlapping windows. All three peptides were flanked with an additional five amino acids per

terminus from the SARS-CoV-2 proteome.

The start and end amino acid positions of each vaccine peptide in its origin gene is shown in

Supplemental Table 1. For SARS-CoV-2, peptides are aligned to reference proteins in UniProt

(UniProt Consortium, 2019) (UniProt IDs: P0DTC2 (S), P0DTC3 (ORF3a), P0DTC5 (M),

P0DTC9 (N), P0DTD1 (ORF1ab)).   All of the epitopes are conserved over these variants of

concern: Alpha (B.1.1.7), Beta (B.1.351), Gamma (P.1), Delta (21A, 21J, B.1.617.2), Kappa

(B.1.617.1), Epsilon (B.1.427, B.1.429), Iota (B.1.526), Lambda (C.37), Mu (B.1.621) , Omicron

(BA.1, BA.2, BA.4, B.5, BA.2.12.1, BA.2.75), EU1 (B.1.177).

MIT-T-COVID vaccine formulation

Codon optimization for mouse expression of the MIT-T-COVID vaccine construct was performed

using the IDT Codon Optimization Tool (Integrated DNA Technologies). The resulting nucleic

acid sequence is provided in Supplemental Table 2. RNA was synthesized by TriLink

BioTechnologies as a modified mRNA transcript with full substitution of 5-Methoxy-U, capped

(Cap 1) using CleanCap® AG and polyadenylated (120A). RNA containing lipid nanoparticles

were prepared as previously described (Pardi et al., 2017). Briefly, an ethanolic solution of

ALC-0315 (Patent WO2017075531), cholesterol, distearoylphosphatidylcholine (DSPC), and 2-

[(polyethylene glycol)-2000] N,N ditetradecylacetamide (ALC-0159, Patent Application

US14732218) was rapidly mixed with an solution of RNA in citrate buffer at pH 4.0 (composition

described in Patent WO2018081480). Physical properties of the LNP such as size and

polydispersity were assessed by Malvern Zetasizer and encapsulation efficiency by Ribogreen
assay (Life Technologies).
499
500   Animal vaccination and virus challenge
501
Thirty-three HLA-A*02:01 human transgenic mice were randomly divided into three groups and
immunized twice at three-week intervals with vehicle (PBS/300 mM sucrose), 10 µg of
Comirnaty® vaccine or 10 µg of MIT-T-COVID vaccine. All vaccines were administered as a
50 µl intramuscular injection. The Comirnaty® vaccine was wastage vaccine that was diluted for
human administration (0.9% NaCl diluent), and remaining unusable wastage vaccine in vials
was flash frozen at -80 °C.   This wastage vaccine was later thawed and immediately
administered without dilution (50 ul is 10 ug of mRNA). No Comirnaty® was used that could
have been administered to humans.   Since Comirnaty® was thawed twice, our results may not
be representative of its best performance.   The MIT-T-COVID vaccine was diluted to 10 µg in
50 µL with PBS with 300 mM sucrose and then administered. At two weeks post booster
immunization, eight mice of each group were challenged with 5×10^4 $TCID_{50}$/60 µL of SARS-
CoV-2 (B.1.351/SA, Strain: hCoV-19/USA/MD-HP01542/2021) via intranasal (IN) route. Mice
were weighted daily and clinically observed at least once daily and scored based on a 1–4
grading system that describes the clinical wellbeing. Three mice in each group were euthanized
at 2 dpi for assessing viral loads and histopathology of the lung. The remaining five mice were
continued monitored for weight changes, other signs of clinical illness, and mortality (if any) for
up to 7 dpi before euthanasia for assessing antibody responses within the blood and viral loads
and histopathology of the lung. Animal studies were conducted at Galveston National
Laboratory at University of Texas Medical Branch at Galveston, Texas, based on a protocol
approved by the Institutional Animal Care and Use Committee at UTMB at Galveston.
522
523   Assessment of mortality and morbidity
524
525   Differentially immunized and challenged mice were monitored at least once each day for the
526   morbidity and mortality and assigned the clinical scores based on the following: 1: Healthy, 2:
527   ruffled fur, lethargic. 3: hunched posture, orbital tightening, increased respiratory rate, and/or >
528   15% weight loss, and 4: dyspnea and/or cyanosis, reluctance to move when stimulated or >
529   20% weight loss.
530

Immunogenicity measurements

At 14 days post booster immunization, three mice of each group were sacrificed for harvesting

splenocytes in 2ml R10 medium (RPMI, 10% FBS, 1%P/S, 10 mM HEPES). Briefly, spleens

were homogenized and subjected to filtration onto 40 μm cell strainers, followed by a wash of

strainers with 10 mL PBS and centrifuged at 500 g for 5 min at 4 °C. Cell pellet was

resuspended by using 2 mL of red blood cell lysis buffer (Thermo Fisher Scientific) for 2-3 min,

followed by a supplement of 20 mL PBS. Resulting cell suspensions were centrifuged at 500 g

for 5 min, resuspended in 2 mL R10 medium before counting the numbers under a microscope.

For a brief in vitro stimulation, aliquots of 10^6 cells were incubated with indicated peptide at a

final concentration of 1 μg/mL in each well of a 96-well plate. Golgiplug (5 μg/mL, BD

Bioscience) was added into the culture at 1 hr post stimulation and followed by an additional 4

hrs incubation. For cell surface staining, cultured splenocytes were resuspended in 40 μL FACS

buffer containing fluorochrome-conjugated antibodies and incubated 1 hr at 4 °C followed by cell

fixation using Cytofix/Cytoperm buffer (BD Bioscience) for 20 min at 4 °C. Cells were further

incubated with fluorochrome-conjugated cytokine antibodies overnight. The next day, cells were

washed twice using 1 × Perm/Wash buffer and resuspended in 300 μL FACS buffer for analysis.

For Foxp3 staining, cells were fixed and permeabilized by Foxp3/Transcription Factor Staining

Buffer Set (Thermo Fisher Scientific) according to the instruction. Fixable Viability Dye

eFluor506 (Thermo Fisher Scientific) was also used in all sample staining to exclude dead cells

from our data analysis. The fluorochrome-conjugated anti-mouse antibodies were listed as

below: FITC-conjugated CD3 (17A2, Biolegend), efluor450-conjugated CD4 (GK1.5,

eBioscience), PE-Cy7-conjugated CD8 (53-6.7, eBioscience), PE-conjugated IFN-γ (XMG1.2,

eBioscience), PerCP-eFlour710- conjugated TNF-α (MP6-XV22, eBioscience). For the Foxp3

staining, Pacific Blue-conjugated CD4 (GK1.5), FITC-conjugated CD25 (PC61), Percp-Cy5.5-

conjugated CTLA4 (UC-10-4B9), PE-conjugated Foxp3 (FJK-16s). Cell acquisition was

performed a BD LSR Fortessa and data were analyzed via FlowJo software version 10 (BD

Bioscience).

Viral titer assay

For virus quantitation, the frozen lung specimens were weighed before homogenization in

PBS/2% FBS solution using the TissueLyser (Qiagen), as previously described (Tseng et al.,

2007). The homogenates were centrifuged to remove cellular debris. Cell debris-free

homogenates were used to quantifying infectious viruses in the standard Vero E6 cell-based

infectivity assays in 96-well microtiter plates, as we routinely used in the lab (Tseng et al.,

2012). Titers of virus were expressed as 50% tissue culture infectious dose per gram of tissue

$(TCID_{50}/g)$.

Antibody neutralization assay

Sera of mice collected at 7 dpi were used for measuring specific antibody responses. Briefly,

sera were heat-inactivated (56 °C) for 30 min, were stored at -80 °C until needed. For

determining the SARS-CoV-2 neutralizing antibody titers, serially two-fold (starting from 1:40)

and duplicate dilutions of heat-inactivated sera were incubated with 100 TCID of SARS-CoV-2

(US_WA-1/2020 isolate) or Beta (B.1.351/SA, Strain: hCoV-19/USA/MD-HP01542/2021) at 37

°C for 1 h before transferring into designated wells of confluent Vero E6 cells grown in 96-well

microtiter plates. Vero E6 cells cultured with medium with or without virus were included as

positive and negative controls, respectively. After incubation at 37 °C for 3 days, individual wells

were observed under the microscopy for the status of virus-induced formation of cytopathic

effect. The 100% neutralizing titers (NT_{100}) of sera were expressed as the lowest dilution folds

capable of completely preventing the formation of viral infection-induced cytopathic effect in

100% of the wells.

Serum IgG/IgM response by ELISA

ELISA was applied to verified serum IgG and IgM responses with SARS-CoV-2 (US_WA-1/2020

isolate) infected Vero-E6 cell lysate. In brief, SARS-CoV-2 infected Vero-E6 cell lysate was

coated on 96-well plate (Corning) at 1 µg/well in PBS for overnight at 4 °C. The plates were

blocked using 1% BSA/PBST for 1 h at room temperature. The 5-fold serial-dilution serum from

each mouse (starting at 1:100) was then added into antigen-coated plates and incubated for 1 h

at 37 °C. The plates were then washed three times with PBST (PBS/0.1% Tween-20) followed

by incubation with 100 µL of anti-mouse IgG and IgM HRP conjugated secondary antibody

(Jackson immunoresearch) (1:2000) for 1 h at 37 °C. After three times wash using PBST, 100

µL of ABTS substrate (Seracare) was added to the plates and incubated for 30 min in the dark.

After stopped by adding 1% SDS. Then absorbance at 405 nm (OD 405 nm) was measured

with the plate reader (Molecular Devices) and analyzed with GraphPad Prism Version 9.1.2.

RNA extraction and quantitative RT-PCR

Lung tissues were weighted and homogenized in 1 mL of Trizol reagent (Invitrogen) using

TissueLyser (Qiagen). The RNA was then extracted using Direct-zol RNA miniprep kits (Zymo

research) according to the manufacturer's instructions. 500 ng total RNA was then applied to

cDNA synthesis using iScript cDNA Synthesis kit (Biorad) according to the manufacturer's

instructions. The viral genomic RNA, subgenomic RNA and mice 18s rRNA has then been

amplified using iQ SYBR green supermix (Biorad) and performed using CFX96 real time system

(Biorad). The samples were run in duplicate using the following conditions: 95 °C for 3 min then

45 cycles of 95 °C for 15 s and 58 °C for 30 s. The level of expression was then normalized with

18s rRNA and calculated using $2^{-\Delta Ct}$ method, as we have previously described (Tseng et al.,

2007; Agrawal et al., 2015)

The primer set for SARS-CoV-2 RNA amplification is nCoV-F

(ACAGGTACGTTAATAGTTAATAGCGT) and nCoV-R (ATATTGCAGCAGTACGCACACA).

SgLeadSARS2-F (CGATCTCTTGTAGATCTGTTCTC) and nCoV-R

(ATATTGCAGCAGTACGCACACA) were used for subgenomic SARS-CoV-2 RNA

amplification. Mouse 18s rRNA was served as the internal control and amplified using 18s-

F(GGACCAGAGCGAAAGCATTTGCC) and 18s-R (TCAATCTCGGGTGGCTGAACGC).

Immunohistochemistry

All slides were prepared by the Histopathology Core (UTMB) into 5 μm paraffin-embedded

sections for immunohistochemistry (IHC). IHC staining and analysis were performed by UTMB

according to previously published protocols (Tseng et al., 2007; Yoshikawa et al., 2009). In

brief, a standard IHC sequential incubation staining protocol was followed to detect the SARS-

CoV-2 spike (S) protein, CD4[+] cells, or CD8[+] cells using a rabbit-raised anti-SARS-CoV-2 S

protein antibody (1:5000 dilution, ab272504, Abcam plc, Cambridge UK), a rabbit monoclonal

anti-CD4 antibody (1:250 dilution, ab183685, ab183685, Abcam plc, Cambridge, UK), and a

rabbit monoclonal anti-CD8 antibody (1:500 dilution, ab217344, Abcam plc, Cambridge, UK)

followed by peroxidase-conjugated secondary antibody and 3,3'-Diaminobenzidine (DAB)

substrate kit (MP-7802, Vector Laboratories, Burlingame, CA). Slides were counterstained with

hematoxylin (MHS16-500ML, Sigma-Aldrich Inc., St. Louis, MO) and antigen expression was

examined under 10X and 40X magnifications using an Olympus IX71 microscope.

CD4[+] and CD8[+] cell counts were quantified using CellProfiler 4.2.4 (Stirling et al., 2021). The

image analysis pipeline included the following modules: (1) UnmixColors (for each of DAB and

hematoxylin stains), (2) ImageMath (applied to DAB stain images only; subtract 0.5 from all

intensity values and set values less than 0 equal to 0), and (3) IdentifyPrimaryObjects (require

object diameter 20-60 pixels [DAB stain] or 15-60 pixels [hematoxylin stain]; Global threshold

strategy; Otsu thresholding method with two-class thresholding, threshold smoothing scale

1.3488, threshold correction factor 1.0, lower threshold bound 0.0, upper threshold bound 1.0,

no log transform before thresholding; Shape method to distinguish clumped objects and draw

dividing lines between clumped objects; automatically calculate size of smoothing filter for

declumping; automatically calculate minimum allowed distance between local maxima; speed

up by using lower-resolution image to find local maxima; fill holes in identified objects after

declumping only; continue handling objects if excessive number of objects identified).

Statistical Analysis

One-way ANOVA tests were performed in Python using the SciPy package (Virtanen et al.,

2020). Two-way ANOVA and Tukey's tests were performed in Python using the statsmodels

package (Seabold and Perktold, 2010). Logrank tests were performed using GraphPad Prism.

Acknowledgements

This work was supported in part by Schmidt Futures and a C3.ai grant to DG. We benefited

from thoughtful comments from Michael Birnbaum.

Declaration of Interests

DG and BC are founders of Think Therapeutics. PL and YT are employees of Acuitas

Therapeutics.

Supplemental Data

MHC Class	Vaccine Peptide	Query Peptide	Query ID	Organism	Gene	Vaccine Peptide Start-End	Query Peptide Start-End	MIRA	MCMT
1	YLYALVYFL	YLYALVYFL	CD8-1	SARS-CoV-2	ORF3a	107-115	107-115	0.86	0.19
1	RSKNPLLYDANYFLCWHTN	LLYDANYFL	CD8-2	SARS-CoV-2	ORF3a	134-152	139-147	0.70	0.44
1	FVDGVPFVV	FVDGVPFVV	CD8-3	SARS-CoV-2	ORF1ab	4726-4734	4726-4734	0.68	
1	AYYVGYLQPRTFLLKYNEN	YLQPRTFLL	CD8-4	SARS-CoV-2	S	264-282	269-277	0.66	0.62
1	FLNRFTTTL	FLNRFTTTL	CD8-5	SARS-CoV-2	ORF1ab	3482-3490	3482-3490	0.53	
1	RLTKYTMADLVYALRHFDE	TMADLVYAL	CD8-6	SARS-CoV-2	ORF1ab	4510-4528	4515-4523	0.44	
1	SIIAYTMSL	SIIAYTMSL	CD8-7	SARS-CoV-2	S	691-699	691-699	0.59	
1	LLLFVTVYSHLLLVAAGLE	TVYSHLLLV	CD8-8	SARS-CoV-2	ORF3a	84-102	89-97	0.61	
1	ATSRTLSYY	ATSRTLSYY	CD8(-)-1	SARS-CoV-2	M	171-179	171-179		
1	KTFPPTEPK	KTFPPTEPK	CD8(-)-2	SARS-CoV-2	N	361-369	361-369		
1	NLVPMVATV	NLVPMVATV	A0201 (+)	CMV	pp65	495-503	495-503		
2	KSILSPLYAFASEAARVVRSIFSRTL	PLYAFASEAARVVRSI	CD4-1	SARS-CoV-2	ORF1ab	527-552	532-547		
2	VDYGARFYFYTSKTTVASLINTLNDL	RFYFYTSKTTVASLIN	CD4-2	SARS-Cov-2	ORF1ab	1416-1441	1421-1436		
2	EEIAIILASFSASTSAFVETVKGLDY	–	Synthesis failed	SARS-CoV-2	ORF1ab	471-496	-		
2	KPVSKMRMATPLLMQAL	KPVSKMRMATPLLMQAL	IAb (+)	Homo sapiens	CD74	102-118	102-118		

Supplemental Table 1. MIT-T-COVID vaccine peptides, query peptides (vaccine immunogenicity), peptide origin, and appearance probability in HLA-A*02:01 convalescent COVID-19 patients in Snyder et al. (2020) (MIRA column) and Kared et al. (2021) (MCMT). Selected peptides were tested by other studies for their immunogenicity in convalescent COVID-19 patients whose HLA type included HLA-A*02:01. The study by Snyder et al. (2020) included 80 HLA-A*02:01 convalescent COVID-19 patients and tested peptides individually or in small pools with the Multiplex Identification of T-cell Receptor Antigen Specificity (MIRA) assay. Query peptides were first filtered to only consider those with predicted HLA-A*02:01 binding affinity less than or equal to 25 nM. The MIRA fraction in Supplemental Table 1 is the number of individuals positive for a pool containing a query peptide divided by 80. The Kared et al. (2021) study evaluated 16 HLA-A*02:01 convalescent COVID-19 patient by mass cytometry–

678   based multiplexed tetramer (MCMT) staining.   The MCMT fraction in Supplemental Table 1 is

679   the number of individuals positive for a query peptide divided by 16.

```
ATG AGG GTC ACA GCT CCT CGG ACC TTG ATC CTC CTT TTG TCT GGT GCT CTT
GCA CTG ACT GAG ACT TGG GCC GGG TCA GGA GGC AGT GGA GGA GGA GGA TCC
GGG GGT TAT TTG TAT GCT CTG GTT TAT TTT CTG GGC GGG TCC GGA GGC GGT
GGC TCT GGC GGG AGG TCC AAG AAT CCA CTT CTC TAC GAC GCA AAC TAT TTC
TTG TGT TGG CAC ACC AAT GGG GGG AGC GGT GGC GGA GGA AGC GGT GGG TTC
GTG GAC GGA GTT CCC TTT GTT GTT GGT GGG TCA GGC GGA GGA GGC TCT GGC
GGG GCT TAC TAT GTA GGG TAC CTG CAG CCC CGA ACA TTC CTT TTG AAA TAC
AAC GAG AAC GGT GGA TCC GGT GGG GGA GGA AGT GGA GGG TTT CTG AAT AGA
TTC ACC ACC ACT CTG GGA GGT TCT GGC GGC GGG GGT TCT GGT GGA CGG CTG
ACT AAA TAC ACA ATG GCC GAT CTT GTT TAC GCA TTG CGG CAT TTT GAT GAG
GGA GGC AGT GGC GGA GGG GGA TCC GGC GGC AGC ATA ATA GCT TAC ACC ATG
TCA CTG GGA GGG AGC GGA GGG GGC GGG AGC GGC GGT TTG CTC CTT TTC GTG
ACA GTG TAT AGC CAT CTC CTT CTG GTG GCA GCT GGC CTT GAA GGG GGG AGC
GGT GGA GGA GGT AGC GGT GGC GCC ACT TCT AGG ACA TTG AGT TAC TAT GGG
GGC AGC GGA GGA GGA GGT TCT GGA GGC AAG ACC TTC CCC CCT ACA GAG CCC
AAG GGA GGT TCC GGC GGC GGG GGC AGT GGT GGG GAA GAG ATC GCC ATT ATC
TTG GCT TCC TTT AGT GCT TCA ACA AGC GCT TTT GTA GAG ACC GTA AAG GGC
CTC GAT TAT GGA GGT TCA GGG GGA GGA GGC TCA GGT GGG AAG TCA ATA CTG
TCT CCT CTT TAT GCA TTT GCA TCA GAG GCT GCA AGA GTT GTC CGA TCC ATT TTT
TCT CGC ACT CTT GGG GGG TCC GGC GGA GGC GGG TCT GGC GGT GTG GAT TAT
GGT GCT AGG TTT TAT TTT TAC ACT TCC AAA ACC ACT GTT GCC TCT CTC ATA AAT
ACC CTC AAT GAC CTG GGA GGG TCT GGT GGC GGG GGG AGT GGC GGG AAT CTT
GTT CCT ATG GTT GCA ACA GTA GGA GGT TCC GGA GGG GGT GGC AGC GGA GGA
AAG CCA GTG TCC AAG ATG AGA ATG GCA ACC CCT TTG CTG ATG CAG GCC CTG
GGT GGC AGT CTG GGA GGT GGT GGC TCC GGC ATC GTA GGT ATA GTC GCC GGA
CTT GCA GTT TTG GCC GTG GTA GTG ATA GGC GCA GTT GTT GCC ACC GTT ATG
TGC CGA CGA AAG AGT TCA GGC GGC AAG GGT GGT TCT TAC TCC CAA GCT GCA
AGC TCC GAC TCC GCT CAG GGG AGT GAT GTT AGC TTG ACT GCA TGA
```

682   **Supplemental Table 2.** Nucleic acid sequence for assembled vaccine construct of Figure 1A.

684
685
686

687
688 **Supplemental Figure 1.** (A) Lung viral RNA level, and (B) lung subgenomic viral RNA level.
689 Error bars indicate the standard deviation around each mean.
690

2 DPI

7 DPI

a.
PBS

b.
MIT

c.
Pfizer
BNT

Supplemental Figure 2. All lung samples were subjected to IHC staining for SARS-CoV-2
spike protein (brown) with hematoxylin counterstain (blue), representative images of which are
shown here. Inset images taken at 40X magnification. Black arrowheads indicate selected areas
of viral infection. (a) Specimens vaccinated with PBS exhibited extensive staining indicative of
viral infection throughout the epithelium of both the bronchioles and the alveolar sacs, with the
viral infection appearing more intense at 2 days post-infection (dpi, left) than at 7 dpi (right).
Although viral infection is significantly reduced by 7 dpi, viral antigen was still readily detectable
throughout alveoli. (b) In comparison, specimens vaccinated with the MIT-T-COVID vaccine
exhibited similarly extensive viral infection at 2 dpi (left) throughout the bronchiolar and alveolar
epithelia, albeit somewhat reduced in intensity. However, by 7 dpi (right), viral infection was
significantly reduced in both extent and intensity, with brown puncta being detected only in a few
alveoli scattered throughout the tissue. (c) Contrasted with both PBS and MIT-T-COVID
vaccinated specimens, the Pfizer/BNT-vaccinated specimens exhibited significantly reduced
viral infections at both 2 (left) and 7 dpi (right). With the exception of a single area at 7 dpi (see
Supplemental Figure 3), viral antigen was undetected at both timepoints.

Supplemental Figure 3. Sole region of detectable SARS-CoV-2 spike antigen in alveoli of

Comirnaty® vaccinated lung specimens at 7 days post-infection. Black arrowhead indicates

infected cell.

712
713
Supplemental Figure 4. Lung histopathology.   Lungs of mice vaccinated with MIT-T-COVID

(A) or Pfizer/BNT (B) are compared with those with PBS (C). At 7 dpi, the MIT-T-COVID

vaccinated group showed extensive lymphocytic infiltrations in perivascular regions and spaces

around bronchi, bronchioles, and alveoli. There are fewer infiltrations found in the Pfizer/BNT or

PBS groups and they are only localized at perivascular regions around bronchi and large

bronchioles. There is widespread congestion along with hemorrhage and few foci of

thromboembolism (arrow in D) seen in the Pfizer/BNT group but not others. Bar = 80 μm in A, B

and C; Bar = 40 μm in D.

Supplemental Figure 5. Lung immunohistochemistry for CD4⁺ cells at 7 dpi. Example CD4⁺ stain images for (A) MIT-T-COVID, (B) Pfizer/BNT, and (C) PBS vaccinated animals. Lung samples were subjected to IHC staining for CD4 (brown) with hematoxylin counterstain (blue). Images were taken at 10X magnification. Red outlines indicate CD4⁺ cells identified and counted by CellProfiler software (Methods). See also Figure 4 and Supplemental Figures 6-7.

Supplemental Figure 6. Lung immunohistochemistry for CD8$^+$ and CD4$^+$ cells at 2 dpi.

Example CD8$^+$ stain images for (A) MIT-T-COVID, (B) Pfizer/BNT, and (C) PBS vaccinated

animals. Example CD4$^+$ stain images for (D) MIT-T-COVID, (E) Pfizer/BNT, and (F) PBS

vaccinated animals. Lung samples were subjected to IHC staining for CD4 (brown) with

hematoxylin counterstain (blue). Images were taken at 10X magnification. Red outlines

indicate cells identified and counted by CellProfiler software (Methods). See also Figure 4,

Supplemental Figure 5, and Supplemental Figure 7.

Supplemental Figure 7. Complete set of cell counts in lung immunohistochemistry images for

CD8$^+$ and CD4$^+$ cells at 2 and 7 dpi. Lung samples were subjected to IHC staining for CD4 or

CD8 (brown) with hematoxylin counterstain (blue). Images were taken at 10X magnification.

Red outlines indicate cells identified by CellProfiler software (Methods). See also Figure 4 and

Supplemental Figures 5-6.

References

1. Agrawal AS, Garron T, Tao X, Peng BH, Wakamiya M, Chan TS, Couch RB, Tseng CT. Generation of a transgenic mouse model of Middle East respiratory syndrome coronavirus infection and disease. J Virol. 2015 Apr;89(7):3659-70. doi: 10.1128/JVI.03427-14. Epub 2015 Jan 14. PMID: 25589660; PMCID: PMC4403411.

2. Baden LR, El Sahly HM, Essink B, Kotloff K, Frey S, Novak R, Diemert D, Spector SA, Rouphael N, Creech CB, McGettigan J, Khetan S, Segall N, Solis J, Brosz A, Fierro C, Schwartz H, Neuzil K, Corey L, Gilbert P, Janes H, Follmann D, Marovich M, Mascola J, Polakowski L, Ledgerwood J, Graham BS, Bennett H, Pajon R, Knightly C, Leav B, Deng W, Zhou H, Han S, Ivarsson M, Miller J, Zaks T; COVE Study Group. Efficacy and Safety of the mRNA-1273 SARS-CoV-2 Vaccine. N Engl J Med. 2021 Feb 4;384(5):403-416. doi: 10.1056/NEJMoa2035389. Epub 2020 Dec 30. PMID: 33378609; PMCID: PMC7787219.

3. Benjamini O, Rokach L, Itchaki G, Braester A, Shvidel L, Goldschmidt N, Shapira S, Dally N, Avigdor A, Rahav G, Lustig Y, Ben David SS, Fineman R, Paz A, Bairey O, Polliack A, Levy I, Tadmor T. Safety and efficacy of the BNT162b mRNA COVID-19 vaccine in patients with chronic lymphocytic leukemia. Haematologica. 2022 Mar 1;107(3):625-634. doi: 10.3324/haematol.2021.279196. PMID: 34320789; PMCID: PMC8883569.

4. Cohen AA, van Doremalen N, Greaney AJ, Andersen H, Sharma A, Starr TN, Keeffe JR, Fan C, Schulz JE, Gnanapragasam PNP, Kakutani LM, West AP Jr, Saturday G, Lee YE, Gao H, Jette CA, Lewis MG, Tan TK, Townsend AR, Bloom JD, Munster VJ, Bjorkman PJ. Mosaic RBD nanoparticles protect against challenge by diverse sarbecoviruses in animal models. Science. 2022 Aug 5;377(6606):eabq0839. doi: 10.1126/science.abq0839. Epub 2022 Aug 5. PMID: 35857620; PMCID: PMC9273039.

5. Croft NP, Smith SA, Pickering J, Sidney J, Peters B, Faridi P, Witney MJ, Sebastian P, Flesch IEA, Heading SL, Sette A, La Gruta NL, Purcell AW, Tscharke DC. Most viral peptides displayed by class I MHC on infected cells are immunogenic. Proc Natl Acad

Sci U S A. 2019 Feb 19;116(8):3112-3117. doi: 10.1073/pnas.1815239116. Epub 2019

Feb 4. PMID: 30718433; PMCID: PMC6386720.

6. de Silva TI, Liu G, Lindsey BB, Dong D, Moore SC, Hsu NS, Shah D, Wellington D,

Mentzer AJ, Angyal A, Brown R, Parker MD, Ying Z, Yao X, Turtle L, Dunachie S;

COVID-19 Genomics UK (COG-UK) Consortium, Maini MK, Ogg G, Knight JC;

ISARIC4C Investigators, Peng Y, Rowland-Jones SL, Dong T. The impact of viral

mutations on recognition by SARS-CoV-2 specific T cells. iScience. 2021 Nov

19;24(11):103353. doi: 10.1016/j.isci.2021.103353. Epub 2021 Oct 28. PMID: 34729465;

PMCID: PMC8552693.

7. Ellis JM, Henson V, Slack R, Ng J, Hartzman RJ, Katovich Hurley C. Frequencies of

HLA-A2 alleles in five U.S. population groups. Predominance Of A*02011 and

identification of HLA-A*0231. Hum Immunol. 2000 Mar;61(3):334-40. doi:

10.1016/s0198-8859(99)00155-x. PMID: 10689125.

8. Geers D, Shamier MC, Bogers S, den Hartog G, Gommers L, Nieuwkoop NN, Schmitz

KS, Rijsbergen LC, van Osch JAT, Dijkhuizen E, Smits G, Comvalius A, van Mourik D,

Caniels TG, van Gils MJ, Sanders RW, Oude Munnink BB, Molenkamp R, de Jager HJ,

Haagmans BL, de Swart RL, Koopmans MPG, van Binnendijk RS, de Vries RD,

GeurtsvanKessel CH. SARS-CoV-2 variants of concern partially escape humoral but not

T-cell responses in COVID-19 convalescent donors and vaccinees. Sci Immunol. 2021

May 25;6(59):eabj1750. doi: 10.1126/sciimmunol.abj1750. PMID: 34035118; PMCID:

PMC9268159.

9. Hajnik RL, Plante JA, Liang Y, Alameh MG, Tang J, Bonam SR, Zhong C, Adam A,

Scharton D, Rafael GH, Liu Y, Hazell NC, Sun J, Soong L, Shi PY, Wang T, Walker DH,

Sun J, Weissman D, Weaver SC, Plante KS, Hu H. Dual spike and nucleocapsid mRNA

vaccination confer protection against SARS-CoV-2 Omicron and Delta variants in

preclinical models. Sci Transl Med. 2022 Sep 14;14(662):eabq1945. doi:

10.1126/scitranslmed.abq1945. Epub 2022 Sep 14. PMID: 36103514.

10. Heitmann JS, Bilich T, Tandler C, Nelde A, Maringer Y, Marconato M, Reusch J, Jäger

S, Denk M, Richter M, Anton L, Weber LM, Roerden M, Bauer J, Rieth J, Wacker M,

Hörber S, Peter A, Meisner C, Fischer I, Löffler MW, Karbach J, Jäger E, Klein R,

Rammensee HG, Salih HR, Walz JS. A COVID-19 peptide vaccine for the induction of SARS-CoV-2 T cell immunity. Nature. 2022 Jan;601(7894):617-622. doi: 10.1038/s41586-021-04232-5. Epub 2021 Nov 23. PMID: 34814158; PMCID: PMC8791831.

11. Israelow B, Mao T, Klein J, Song E, Menasche B, Omer SB, Iwasaki A. Adaptive immune determinants of viral clearance and protection in mouse models of SARS-CoV-2. Sci Immunol. 2021 Oct 15;6(64):eabl4509. doi: 10.1126/sciimmunol.abl4509. Epub 2021 Sep 2. PMID: 34623900; PMCID: PMC9047536.

12. Kared H, Redd AD, Bloch EM, Bonny TS, Sumatoh H, Kairi F, Carbajo D, Abel B, Newell EW, Bettinotti MP, Benner SE, Patel EU, Littlefield K, Laeyendecker O, Shoham S, Sullivan D, Casadevall A, Pekosz A, Nardin A, Fehlings M, Tobian AA, Quinn TC. SARS-CoV-2-specific CD8+ T cell responses in convalescent COVID-19 individuals. J Clin Invest. 2021 Mar 1;131(5):e145476. doi: 10.1172/JCI145476. PMID: 33427749; PMCID: PMC7919723.

13. Kingstad-Bakke B, Lee W, Chandrasekar SS, Gasper DJ, Salas-Quinchucua C, Cleven T, Sullivan JA, Talaat A, Osorio JE, Suresh M. Vaccine-induced systemic and mucosal T cell immunity to SARS-CoV-2 viral variants. Proc Natl Acad Sci U S A. 2022 May 17;119(20):e2118312119. doi: 10.1073/pnas.2118312119. Epub 2022 May 13. PMID: 35561224; PMCID: PMC9171754.

14. Kotturi MF, Assarsson E, Peters B, Grey H, Oseroff C, Pasquetto V, Sette A. Of mice and humans: how good are HLA transgenic mice as a model of human immune responses? Immunome Res. 2009 Jun 17;5:3. doi: 10.1186/1745-7580-5-3. PMID: 19534819; PMCID: PMC2702351.

15. Kreiter S, Selmi A, Diken M, Sebastian M, Osterloh P, Schild H, Huber C, Türeci O, Sahin U. Increased antigen presentation efficiency by coupling antigens to MHC class I trafficking signals. J Immunol. 2008 Jan 1;180(1):309-18. doi: 10.4049/jimmunol.180.1.309. Erratum in: J Immunol. 2012 Sep 1;189(5):2682. PMID: 18097032.

888   16. Kumari P, Rothan HA, Natekar JP, Stone S, Pathak H, Strate PG, Arora K, Brinton MA,
889   Kumar M. Neuroinvasion and Encephalitis Following Intranasal Inoculation of SARS-
890   CoV-2 in K18-hACE2 Mice. Viruses. 2021 Jan 19;13(1):132. doi: 10.3390/v13010132.
891   PMID: 33477869; PMCID: PMC7832889.
892
893   17. Liu G, Carter B, Bricken T, Jain S, Viard M, Carrington M, Gifford DK. Computationally
894   Optimized SARS-CoV-2 MHC Class I and II Vaccine Formulations Predicted to Target
895   Human Haplotype Distributions. Cell Syst. 2020 Aug 26;11(2):131-144.e6. doi:
896   10.1016/j.cels.2020.06.009. Epub 2020 Jul 27. PMID: 32721383; PMCID: PMC7384425.
897
898   18. Liu G, Carter B, Gifford DK. Predicted Cellular Immunity Population Coverage Gaps for
899   SARS-CoV-2 Subunit Vaccines and Their Augmentation by Compact Peptide Sets. Cell
900   Syst. 2021 Jan 20;12(1):102-107.e4. doi: 10.1016/j.cels.2020.11.010. Epub 2020 Nov
901   27. PMID: 33321075; PMCID: PMC7691134.
902
903   19. Liu, G., Carter B., Dimitrakakis A., Gifford D. K. Maximum n-times Coverage for Vaccine
904   Design, International Conference on Learning Representations, ICLR (2022),
905   https://arxiv.org/abs/2101.10902
906
907   20. Martinez DR, Schäfer A, Leist SR, De la Cruz G, West A, Atochina-Vasserman EN,
908   Lindesmith LC, Pardi N, Parks R, Barr M, Li D, Yount B, Saunders KO, Weissman D,
909   Haynes BF, Montgomery SA, Baric RS. Chimeric spike mRNA vaccines protect against
910   Sarbecovirus challenge in mice. Science. 2021 Aug 27;373(6558):991-998. doi:
911   10.1126/science.abi4506. Epub 2021 Jun 22. PMID: 34214046; PMCID: PMC8899822.
912
913   21. Matchett WE, Joag V, Stolley JM, Shepherd FK, Quarnstrom CF, Mickelson CK,
914   Wijeyesinghe S, Soerens AG, Becker S, Thiede JM, Weyu E, O'Flanagan S, Walter JA,
915   Vu MN, Menachery VD, Bold TD, Vezys V, Jenkins MK, Langlois RA, Masopust D.
916   Nucleocapsid vaccine elicits spike-independent SARS-CoV-2 protective immunity. J
917   Immunol. 2021 Jun 30;: PMID: 33948591; PMCID: PMC8095198.
918
919   22. Moss P. The T cell immune response against SARS-CoV-2. Nat Immunol. 2022
920   Feb;23(2):186-193. doi: 10.1038/s41590-021-01122-w. Epub 2022 Feb 1. PMID:
921   35105982.

23. Nakiboneka R, Mugaba S, Auma BO, Kintu C, Lindan C, Nanteza MB, Kaleebu P, Serwanga J. Interferon gamma (IFN-γ) negative CD4+ and CD8+ T-cells can produce immune mediators in response to viral antigens. Vaccine. 2019 Jan 3;37(1):113-122. doi: 10.1016/j.vaccine.2018.11.024. Epub 2018 Nov 17. PMID: 30459072; PMCID: PMC6290111.

24. Nathan A, Rossin EJ, Kaseke C, Park RJ, Khatri A, Koundakjian D, Urbach JM, Singh NK, Bashirova A, Tano-Menka R, Senjobe F, Waring MT, Piechocka-Trocha A, Garcia-Beltran WF, Iafrate AJ, Naranbhai V, Carrington M, Walker BD, Gaiha GD. Structure-guided T cell vaccine design for SARS-CoV-2 variants and sarbecoviruses. Cell. 2021 Aug 19;184(17):4401-4413.e10. doi: 10.1016/j.cell.2021.06.029. Epub 2021 Jun 30. PMID: 34265281; PMCID: PMC8241654.

25. O'Donnell TJ, Rubinsteyn A, Laserson U. MHCflurry 2.0: Improved Pan-Allele Prediction of MHC Class I-Presented Peptides by Incorporating Antigen Processing. Cell Syst. 2020 Jul 22;11(1):42-48.e7. doi: 10.1016/j.cels.2020.06.010. Epub 2020 Jul 14. Erratum in: Cell Syst. 2020 Oct 21;11(4):418-419. PMID: 32711842.

26. Pardi N, Hogan MJ, Pelc RS, Muramatsu H, Andersen H, DeMaso CR, Dowd KA, Sutherland LL, Scearce RM, Parks R, Wagner W, Granados A, Greenhouse J, Walker M, Willis E, Yu JS, McGee CE, Sempowski GD, Mui BL, Tam YK, Huang YJ, Vanlandingham D, Holmes VM, Balachandran H, Sahu S, Lifton M, Higgs S, Hensley SE, Madden TD, Hope MJ, Karikó K, Santra S, Graham BS, Lewis MG, Pierson TC, Haynes BF, Weissman D. Zika virus protection by a single low-dose nucleoside-modified mRNA vaccination. Nature. 2017 Mar 9;543(7644):248-251. doi: 10.1038/nature21428. Epub 2017 Feb 2. PMID: 28151488; PMCID: PMC5344708.

27. Pardieck IN, van der Sluis TC, van der Gracht ETI, Veerkamp DMB, Behr FM, van Duikeren S, Beyrend G, Rip J, Nadafi R, Beyranvand Nejad E, Mülling N, Brasem DJ, Camps MGM, Myeni SK, Bredenbeek PJ, Kikkert M, Kim Y, Cicin-Sain L, Abdelaal T, van Gisbergen KPJM, Franken KLMC, Drijfhout JW, Melief CJM, Zondag GCM, Ossendorp F, Arens R. A third vaccination with a single T cell epitope confers protection

955   in a murine model of SARS-CoV-2 infection. Nat Commun. 2022 Jul 8;13(1):3966. doi:
956   10.1038/s41467-022-31721-6. PMID: 35803932; PMCID: PMC9267705.
957

28. Redd AD, Nardin A, Kared H, Bloch EM, Pekosz A, Laeyendecker O, Abel B, Fehlings
M, Quinn TC, Tobian AA. CD8+ T cell responses in COVID-19 convalescent individuals
target conserved epitopes from multiple prominent SARS-CoV-2 circulating variants.
medRxiv [Preprint]. 2021 Feb 12:2021.02.11.21251585. doi:
10.1101/2021.02.11.21251585. Update in: Open Forum Infect Dis. 2021 Mar
30;8(7):ofab143. PMID: 33594378; PMCID: PMC7885937.
964

965   29. Reynisson B, Alvarez B, Paul S, Peters B, Nielsen M. NetMHCpan-4.1 and
966   NetMHCIIpan-4.0: improved predictions of MHC antigen presentation by concurrent
967   motif deconvolution and integration of MS MHC eluted ligand data. Nucleic Acids Res.
968   2020 Jul 2;48(W1):W449-W454. doi: 10.1093/nar/gkaa379. PMID: 32406916; PMCID:
969   PMC7319546.
970

30. Sahin U, Derhovanessian E, Miller M, Kloke BP, Simon P, Löwer M, Bukur V, Tadmor
AD, Luxemburger U, Schrörs B, Omokoko T, Vormehr M, Albrecht C, Paruzynski A,
Kuhn AN, Buck J, Heesch S, Schreeb KH, Müller F, Ortseifer I, Vogler I, Godehardt E,
Attig S, Rae R, Breitkreuz A, Tolliver C, Suchan M, Martic G, Hohberger A, Sorn P,
Diekmann J, Ciesla J, Waksmann O, Brück AK, Witt M, Zillgen M, Rothermel A,
Kasemann B, Langer D, Bolte S, Diken M, Kreiter S, Nemecek R, Gebhardt C, Grabbe
S, Höller C, Utikal J, Huber C, Loquai C, Türeci Ö. Personalized RNA mutanome
vaccines mobilize poly-specific therapeutic immunity against cancer. Nature. 2017 Jul
13;547(7662):222-226. doi: 10.1038/nature23003. Epub 2017 Jul 5. PMID: 28678784.
980

981   31. Seabold, S., Perktold, J. Statsmodels: Econometric and statistical modeling with python.
982   Proceedings of the 9th Python in Science Conference. 2010; 57(61).
983

32. Sekine T, Perez-Potti A, Rivera-Ballesteros O, Strålin K, Gorin JB, Olsson A, Llewellyn-
Lacey S, Kamal H, Bogdanovic G, Muschiol S, Wullimann DJ, Kammann T, Emgård J,
Parrot T, Folkesson E; Karolinska COVID-19 Study Group, Rooyackers O, Eriksson LI,
Henter JI, Sönnerborg A, Allander T, Albert J, Nielsen M, Klingström J, Gredmark-Russ
S, Björkström NK, Sandberg JK, Price DA, Ljunggren HG, Aleman S, Buggert M. Robust

T Cell Immunity in Convalescent Individuals with Asymptomatic or Mild COVID-19. Cell.

2020 Oct 1;183(1):158-168.e14. doi: 10.1016/j.cell.2020.08.017. Epub 2020 Aug 14.

PMID: 32979941; PMCID: PMC7427556.

33. Shuai H, Chan JF, Yuen TT, Yoon C, Hu JC, Wen L, Hu B, Yang D, Wang Y, Hou Y,

Huang X, Chai Y, Chan CC, Poon VK, Lu L, Zhang RQ, Chan WM, Ip JD, Chu AW, Hu

YF, Cai JP, Chan KH, Zhou J, Sridhar S, Zhang BZ, Yuan S, Zhang AJ, Huang JD, To

KK, Yuen KY, Chu H. Emerging SARS-CoV-2 variants expand species tropism to

murines. EBioMedicine. 2021 Nov;73:103643. doi: 10.1016/j.ebiom.2021.103643. Epub

2021 Oct 21. PMID: 34689086; PMCID: PMC8530107.

34. Snyder TM, Gittelman RM, Klinger M, May DH, Osborne EJ, Taniguchi R, Zahid HJ,

Kaplan IM, Dines JN, Noakes MT, Pandya R, Chen X, Elasady S, Svejnoha E, Ebert P,

Pesesky MW, De Almeida P, O'Donnell H, DeGottardi Q, Keitany G, Lu J, Vong A,

Elyanow R, Fields P, Greissl J, Baldo L, Semprini S, Cerchione C, Nicolini F, Mazza M,

Delmonte OM, Dobbs K, Laguna-Goya R, Carreño-Tarragona G, Barrio S, Imberti L,

Sottini A, Quiros-Roldan E, Rossi C, Biondi A, Bettini LR, D'Angio M, Bonfanti P,

Tompkins MF, Alba C, Dalgard C, Sambri V, Martinelli G, Goldman JD, Heath JR, Su

HC, Notarangelo LD, Paz-Artal E, Martinez-Lopez J, Carlson JM, Robins HS. Magnitude

and Dynamics of the T-Cell Response to SARS-CoV-2 Infection at Both Individual and

Population Levels. medRxiv [Preprint]. 2020 Sep 17:2020.07.31.20165647. doi:

10.1101/2020.07.31.20165647. PMID: 32793919; PMCID: PMC7418734.

1011

1012   35. Stirling DR, Swain-Bowden MJ, Lucas AM, Carpenter AE, Cimini BA, Goodman A.

1013   CellProfiler 4: improvements in speed, utility and usability. BMC Bioinformatics. 2021

1014   Sep 10;22(1):433. doi: 10.1186/s12859-021-04344-9. PMID: 34507520; PMCID:

1015   PMC8431850.

1016

1017   36. Swank Z, Senussi Y, Manickas-Hill Z, Yu XG, Li JZ, Alter G, Walt DR. Persistent

1018   circulating SARS-CoV-2 spike is associated with post-acute COVID-19 sequelae. Clin

1019   Infect Dis. 2022 Sep 2:ciac722. doi: 10.1093/cid/ciac722. Epub ahead of print. PMID:

1020   36052466.

1021

1022   37. Tregoning JS, Flight KE, Higham SL, Wang Z, Pierce BF. Progress of the COVID-19
1023   vaccine effort: viruses, vaccines and variants versus efficacy, effectiveness and escape.
1024   Nat Rev Immunol. 2021 Oct;21(10):626-636. doi: 10.1038/s41577-021-00592-1. Epub
1025   2021 Aug 9. PMID: 34373623; PMCID: PMC8351583.

1026

1027   38. Tseng CTK, Huang C, Newman P, Wang N, Narayanan K, Watts DM, Makino S,
1028   Packard M, Zaki SR, Chan TS, and Peters CJ. 2007. SARS Coronavirus Infection
1029   of Transgenic Mice bearing the Human Angiotensin Converting Enzyme 2 (hACE2) Virus
1030   Receptor. *JVI* 81: 1162-1173.

1031
39. Tseng CT, Sbrana E, Iwata-Yoshikawa N, Newman PC, Garron T, Atmar RL, Peters CJ,
Couch RB. Immunization with SARS coronavirus vaccines leads to pulmonary
immunopathology on challenge with the SARS virus. PLoS One. 2012;7(4):e35421. doi:
10.1371/journal.pone.0035421. Epub 2012 Apr 20. Erratum in: PLoS One. 2012;7(8).
doi:10.1371/annotation/2965cfae-b77d-4014-8b7b-236e01a35492. PMID: 22536382;
PMCID: PMC3335060.

1038

1039   40. UniProt Consortium. UniProt: a worldwide hub of protein knowledge. Nucleic Acids Res.
1040   2019 Jan 8;47(D1):D506-D515. doi: 10.1093/nar/gky1049. PMID: 30395287; PMCID:
1041   PMC6323992.

1042

41. Virtanen P, Gommers R, Oliphant TE, Haberland M, Reddy T, Cournapeau D, Burovski
E, Peterson P, Weckesser W, Bright J, van der Walt SJ, Brett M, Wilson J, Millman KJ,
Mayorov N, Nelson ARJ, Jones E, Kern R, Larson E, Carey CJ, Polat İ, Feng Y, Moore
EW, VanderPlas J, Laxalde D, Perktold J, Cimrman R, Henriksen I, Quintero EA, Harris
CR, Archibald AM, Ribeiro AH, Pedregosa F, van Mulbregt P; SciPy 1.0 Contributors.
SciPy 1.0: fundamental algorithms for scientific computing in Python. Nat Methods. 2020
Mar;17(3):261-272. doi: 10.1038/s41592-019-0686-2. Epub 2020 Feb 3. Erratum in: Nat
Methods. 2020 Feb 24;: PMID: 32015543; PMCID: PMC7056644.

1051
1052   42. Vita R, Mahajan S, Overton JA, Dhanda SK, Martini S, Cantrell JR, Wheeler DK, Sette
1053   A, Peters B. The Immune Epitope Database (IEDB): 2018 update. Nucleic Acids Res.
1054   2018 Oct 24. doi: 10.1093/nar/gky1006. [Epub ahead of print] PubMed PMID: 30357391.

1055

43. Walsh EE, Frenck RW Jr, Falsey AR, Kitchin N, Absalon J, Gurtman A, Lockhart S,
Neuzil K, Mulligan MJ, Bailey R, Swanson KA, Li P, Koury K, Kalina W, Cooper D,
Fontes-Garfias C, Shi PY, Türeci Ö, Tompkins KR, Lyke KE, Raabe V, Dormitzer PR,
Jansen KU, Şahin U, Gruber WC. Safety and Immunogenicity of Two RNA-Based Covid-
19 Vaccine Candidates. N Engl J Med. 2020 Dec 17;383(25):2439-2450. doi:
10.1056/NEJMoa2027906. Epub 2020 Oct 14. PMID: 33053279; PMCID: PMC7583697.
1062

44. Willett BJ, Grove J, MacLean OA, Wilkie C, De Lorenzo G, Furnon W, Cantoni D, Scott
S, Logan N, Ashraf S, Manali M, Szemiel A, Cowton V, Vink E, Harvey WT, Davis C,
Asamaphan P, Smollett K, Tong L, Orton R, Hughes J, Holland P, Silva V, Pascall DJ,
Puxty K, da Silva Filipe A, Yebra G, Shaaban S, Holden MTG, Pinto RM, Gunson R,
Templeton K, Murcia PR, Patel AH, Klenerman P, Dunachie S; PITCH Consortium;
COVID-19 Genomics UK (COG-UK) Consortium, Haughney J, Robertson DL, Palmarini
M, Ray S, Thomson EC. SARS-CoV-2 Omicron is an immune escape variant with an
altered cell entry pathway. Nat Microbiol. 2022 Aug;7(8):1161-1179. doi:
10.1038/s41564-022-01143-7. Epub 2022 Jul 7. PMID: 35798890; PMCID:
PMC9352574.
1073

1074   45. World Health Organization (WHO). In the wake of the pandemic: preparing for Long
1075   COVID (2021). Policy Brief 39. Accessed August 14, 2022.
1076   https://apps.who.int/iris/bitstream/handle/10665/339629/Policy-brief-39-1997-8073-
1077   eng.pdf
1078

1079   46. Zeng H, Gifford DK. Quantification of Uncertainty in Peptide-MHC Binding Prediction
1080   Improves High-Affinity Peptide Selection for Therapeutic Design. Cell Syst. 2019 Aug
1081   28;9(2):159-166.e3. doi: 10.1016/j.cels.2019.05.004. Epub 2019 Jun 5. PMID:
1082   31176619; PMCID: PMC6715517.
1083

1084   47. Zhang X, Wu S, Wu B, Yang Q, Chen A, Li Y, Zhang Y, Pan T, Zhang H, He X. SARS-
1085   CoV-2 Omicron strain exhibits potent capabilities for immune evasion and viral entrance.
1086   Signal Transduct Target Ther. 2021 Dec 17;6(1):430. doi: 10.1038/s41392-021-00852-5.
1087   PMID: 34921135; PMCID: PMC8678971.
1088

1089   48. Zhao J, Zhao J, Mangalam AK, Channappanavar R, Fett C, Meyerholz DK, Agnihothram

1090   S, Baric RS, David CS, Perlman S. Airway Memory CD4(+) T Cells Mediate Protective

1091   Immunity against Emerging Respiratory Coronaviruses. Immunity. 2016 Jun

1092   21;44(6):1379-91. doi: 10.1016/j.immuni.2016.05.006. Epub 2016 Jun 7. PMID:

1093   27287409; PMCID: PMC4917442.