Selection leads to remarkable variability in the outcomes of hybridization across replicate hybrid zones

S. Eryn McFarlane[1,2], Joshua P. Jahner[1], Dorothea Lindtke[3], C. Alex Buerkle[1], Elizabeth G. Mandeville[4,5]

[1] Department of Botany, University of Wyoming, Laramie, Wyoming, USA
[2] Department of Biology, York University, Toronto, Canada
[3] Agroscope, Animal GenoPhenomics, Posieux, Switzerland
[4] Department of Integrative Biology, University of Guelph, Guelph, Ontario, Canada
[5] Biology Department, Northern Michigan University, Marquette, Michigan, USA

ORCID
S. Eryn McFarlane   https://orcid.org/0000-0002-0706-458X
Joshua P. Jahner   https://orcid.org/0000-0001-8121-6783
Dorothea Lindtke   https://orcid.org/0000-0003-1339-1410
C. Alex Buerkle   https://orcid.org/0000-0003-4222-8858
Elizabeth G. Mandeville   https://orcid.org/0000-0002-9461-9140

Correspondence:
S. Eryn McFarlane, Department of Biology, York University, Toronto, Canada.
Email: emcfar@yorku.ca

Keywords: *ancestry, conservation, drift, incompatibility, introgression, secondary contact, speciation*

Running head: Variable outcomes of hybridization

1 Abstract

Hybrid zones have been held up as an opportunity to see speciation in action, as windows into the evolutionary process. When hybrid zones are replicated, it is assumed that if the same genetic incompatibilities are maintaining reproductive isolation across all instances of secondary contact, those incompatibilities should be identifiable by consistent patterns in the genome. In contrast, changes in allele frequencies due to genetic drift should be idiosyncratic for each hybrid zone. To test this assumption, we simulated 20 replicates of each of 12 hybrid zone scenarios with varied genetic incompatibilities, rates of migration, and selection. We found remarkable variability in the outcomes of hybridization in replicate hybrid zones, particularly with Bateson-Dobzhansky-Muller incompatibilities and strong selection. We found substantial differences among replicates in the overall genomic composition of individuals, including admixture proportions, inter-specific ancestry complement, and number of ancestry junctions. Additionally, we found substantial variation in genomic clines among replicates at focal loci, regardless of locus-specific selection. From this, we conclude that processes other than selection are responsible for some consistent outcomes of hybridization, whereas selection on incompatibilities can lead to genomically widespread and highly variable outcomes. We use these results to highlight the challenge of mapping between pattern and process in hybrid zones and call attention to how selection against incompatibilities will commonly lead to unpredictable, variable outcomes. Our hope is that this study informs future research on replicate hybrid zones and encourages further development of statistical techniques, theoretical models, and exploration of additional axes of variation to understand the maintenance of reproductive isolation.

20 Introduction

"Same thing you do when you cross an elephant and a rhino. El-eph-I-no." — Ted Lasso

Following and during speciation, the local or global persistence of novel species will commonly be tested through range shifts and secondary contact between diverged lineages. Species will be more likely to persist in the face of potential interspecific gene flow if hybrid fitness is low due to incompatible alleles or genotypes (Coyne & Orr 2004). For some groups, we have good descriptions of the genetics of incompatibilities and evolutionary models for their origin (Fishman & Sweigart 2018, Larson *et al.* 2018, Maheshwari & Barbash 2011, Orr 1995, Powell *et al.* 2020). Likewise, we have well-developed expressions for equilibrium clinal variation following secondary contact (Barton 1979, 1983, Barton & Hewitt 1985, Barton & Gale 1993, Endler 1977). Together these have formed an expectation that we can use empirical observations of gene exchange and ancestry variation in hybrid zones to identify exceptional loci. Specifically, we can use hybrid zones to look for genetic loci associated with reproductive isolation and variation in fitness between hybrids and parental type individuals. For example, geographic and genomic cline methods identify loci with deviations in allelic ancestry or state relative to a genome wide distribution of ancestry (Gompert & Buerkle 2012, Szymura & Barton 1986). There is growing interest in using contrasts of multiple hybrid zones to build our understanding of their consistent and idiosyncratic features (e.g. Simon *et al.* 2021, Westram *et al.* 2021). This mirrors diverse research in evolutionary biology that uses replicates or multiple instances of evolution to reveal parallel or concordant evolution and to implicate predictable and potentially deterministic features of evolution (Elmer *et al.* 2014, Ferris *et al.* 2021, Tenaillon *et al.* 2016).

In hybrid zones, outcomes of secondary contact and the composition of hybrid populations will be driven by drift, gene flow, selection, and attributes of the genome such as linkage and recombination rates. It is an open question what level of consistency among replicate hybrid populations we should expect with different levels of selection against recombinant genotypes. Would high consistency among replicates indicate strong selection and effective reproductive isolation, or would consistency among replicates instead be a hallmark of neutral, clinal variation? We might expect consistency to be indicative of natural selection, particularly directional selection (Pardo-Diaz *et al.* 2012, Whitney *et al.* 2006). Indeed, a number of studies have suggested some measure of locus-specific consistency as a way to determine candidate regions for incompatibility loci (e.g. Kingston *et al.* 2017, Gagnaire *et al.* 2013, Caballero-Villalobos *et al.* 2021, Larson *et al.* 2014). However, the fitnesses of hybrid genotypes that contribute to intrinsic post-zygotic reproductive isolation do not involve directional selection, but are predicted to be multilocus, epistatic incompatibilities (Bateson 1909, Dobzhansky 1937, Muller 1942). Rather than deterministic and consistent shifts in allele frequency,

the evolutionary dynamics associated with these fitness matrices include unstable equilibria and frequency dependence that could lead to high variance in outcomes, despite strong selection.

In a world in which the evolutionary processes governing hybridization generate predictable outcomes, one could map between empirical observations and evolutionary processes. Specifically, hybrid zone characteristics (e.g. genetic composition and ancestry of individuals, number of hybrids, time since contact) would allow inference of the different demographic conditions, including selection and gene flow. Moreover, replicate hybrid zones could be used to understand how these evolutionary processes vary across space, time, and ecological variation (Rieseberg *et al.* 1999, Buerkle & Rieseberg 2001, Harrison & Larson 2016, Westram *et al.* 2021). Consistency or parallelism of loci with exceptional differentiation or introgression between populations (via F_{ST}, and geographic or genomic clines) has been suggested as evidence for either loci associated with adaptive introgression or genomic regions associated with reproductive isolation (Langdon *et al.* 2022, Matute *et al.* 2020, Rieseberg *et al.* 1999, Simon *et al.* 2020, 2021, Westram *et al.* 2021). Explicitly stated, this predicts that consistent selection against a hybrid genotype, regardless of environment, will lead to detectable consistency across replicate hybrid zones, and this consistency is a hallmark of selection. Further, this assumes that replicate hybrid zones between the same species pairs will behave the same way (i.e. similar to replicated lab populations) across time or space. But the question remains: if hybrid zones are indeed windows into evolution (Harrison 1990), do different scenes viewed through different windows imply differences in process? If instead the outcomes of hybridization are highly idiosyncratic or capricious (*sensu* Lewontin 1966), potentially due to stochasticity following initial contact in different hybrid zones, the ability to link patterns to processes will be limited no matter how many replicate hybrid zones are sampled.

To resolve expectations for consistency and contingency in evolutionary replicates, we simulated replicate outcomes of hybridization, varying selection, migration, and genetic architecture of incompatibilities. We have deliberately simulated only simple genetic architectures, specifically two or four locus incompatibilities, rather than tackling polygenic selection. Specifically, we simulated either an inter-genomic (hereafter 'BDMI', i.e. a Bateson-Dobzhansky-Muller incompatibility) or intra-genomic incompatibility (hereafter 'pathway'). BDMIs are theorized as the result of different alleles arising at two (or more) loci in diverged lineages, where an incompatibility arises when the lineages hybridize, and heterozygous genotypes have a negative epistatic interaction, resulting in lower fitness in F_1, F_2, or both categories of individuals (Bateson 1909, Dobzhansky 1936, Muller 1942). For example, (Powell *et al.* 2020) found a two locus incompatibility that contributes to melanoma in *Xiphophorus maculatus* and *X. hellerii* hybrids. In contrast, pathway incompatibilities have been associated with *cis* regulation of gene expression, where incompatibilities can result in disregulated expression in hybrids (Landry *et al.* 2007).

We are particularly interested in strictly intrinsic incompatibilities (i.e., hybrids have lower fitness without ecological, or extrinsic selection (Coughlan & Matute 2020)). Intrinsic genetic incompatibilities are expected to be the final step of reproductive isolation, preventing gene flow between species even if there is no longer prezygotic isolation, such as in the case of range shifts or translocations (Coughlan & Matute 2020). There is an expectation that intrinsic incompatibilities could be more general across taxa than mechanisms leading to prezygotic isolation (Xiong & Mallet 2022). Once we understand how intrinsic incompatibilities in consistent environments might result in variable outcomes across sites, we can develop theoretical expectations for more complex processes. For example, there are not yet theoretical expectations for how ecological interactions with genetic incompatibilities (Thompson *et al.* 2022), 'coupling' of rates of introgression across the genome (Barton 1983, Barton & De Cara 2009), or 'hybridization load' (Moran *et al.* 2021) might behave across replicate hybrid zones, or, eventually even across taxa. Theoretical understanding of the variability of intrinsic incompatibilities across replicate hybrid zones is needed as a baseline before adding the additional complications of ecological selection.

Many simulations we use here were originally performed in association with Lindtke & Buerkle (2015), but were not previously used to quantify consistency in outcomes of hybridization. Lindtke & Buerkle (2015) demonstrated that different genetic architectures causing reproductive isolation resulted in variation in introgression and the effectiveness of reproductive isolation. In contrast, we examined variation among replicates to 1) quantify the degree of variability across replicate hybrid zones with identical evolutionary conditions (genomic architecture and fitness consequences of incompatibilities, population size, migration); 2) determine how different evolutionary conditions affect consistency across replicate hybrid zones; and 3) contrast variability of genotypes at specific loci, particularly those underlying the incompatibilities between species. Finally, we examined how consistency among populations increased with unequal parental populations and

to what extent fthis ratio between parental populations made it easier to pinpoint effects of selection on locus-specific introgression. We hypothesize that the effects of drift at both the population and locus-specific levels make it difficult to detect selection. We predict that even when there is consistency among replicate hybrid populations, as is expected when there parental population sizes are unequal, the consistency is not due to strong selection at either the individual or locus-specific level, but rather due to drift leading to the loss of rare alleles.

Methods

Simulation framework

Following Lindtke & Buerkle (2015) and Gompert *et al.* (2012), we simulated diploid whole genomes (2N = 20) using the software dfuse (https://www.uwyo.edu/buerkle/software/dfuse/, https://doi.org/10.5061/dryad.0506g). A total of 510 loci were simulated per individual, with 51 equally spaced loci per chromosome. Individuals were arranged in a chain of 11 demes, with infinitely sized populations of different species at either end of the chain of demes and 150 individuals in the intermediate demes. Thus, the meta populations act as sink populations, where there can be admixture from the infinite parental source populations (Fig. 1, Lindtke & Buerkle 2015)). Simulations of 11 demes allowed for population structure, with individuals in the spatially central deme unlikely to have direct parentage involving the two species (F1 or backcrossed hybrids). This is consistent with a 'classical' hybrid zone, rather than a 'mosaic' hybrid zone (Rand & Harrison 1989). We also explored additional, complementary analyses for a 3 deme case (as in Lindtke & Buerkle 2015) to verify results were robust to different deme structures.

To determine the degree of variability among replicate hybrid zones across a broad range of evolutionary conditions, we considered 20 independent replicates of 12 different scenarios that varied by migration rate, strength of selection, and type of incompatibility. Migration rate was set as $m/2$, where m was either 0.01 or 0.2 and could occur in a stepwise fashion between demes (except from deme 2 to 1 and from deme 10 to 11, where migration was not allowed). Viability selection (c) against individuals carrying mismatched alleles in either a BDMI or pathway incompatibility was either 0, 0.2, or 0.9. For all BDMI scenarios, two loci on the first chromosome (the fourth and sixth loci on the first chromosome: 1.4, 1.6) were specified as the causal variants, whereas two loci on both the first and second chromosomes (1.2, 1.4, 2.2, 2.4) were specified for all pathway scenarios (see Fig. 1 for genotypic fitness values). Incompatibility loci positions were chosen arbitrarily while maintaining the same distance between markers (Lindtke & Buerkle 2015).

Individual genotypes, admixture proportion (q), inter-specific ancestry (Q_{12}), and the number of junctions between ancestry blocks were all recorded for each individual every 10 generations (from 10–100). The latter three statistics describe the ancestry of hybrid individuals: the admixture proportion (q) is the proportion of ancestry in an individual from one of the two parental species, the inter-specific ancestry (Q_{12}) is the proportion of an individual's diploid genome where one copy comes from parental population 1 and the other parental population 2, (Fig. S1; Gompert *et al.* 2014, Lindtke *et al.* 2014, Shastry *et al.* 2021), and the number of junctions describes the extent of recombination between ancestry from the parental genomes. With more generations of hybridization and recombination, inter-specific ancestry (Q_{12}) declines and the number of junctions increases. We chose these metrics to describe our hybrid individuals, as the distributions of q, Q_{12}, and junctions have been used to infer the processes underlying hybrid zones (Jiggins & Mallet 2000, Schumer *et al.* 2017). A more comprehensive description of the simulations can be found in Lindtke & Buerkle (2015).

Quantifying variability across replicate hybrid zones

Our primary goal was to determine under which scenarios replicate hybrid zones are consistent with one another. For all analyses, we considered only the central deme where the proportion of hybrids is typically the highest (deme 6; results not shown). Additionally, we focused on the outcomes from generations 10 and 100. As our primary metric of variability, we calculated Fisher's F statistics from analysis of variance (ANOVA) in R v4.0.5 (R Core Team 2021) to quantify the ratio of among versus within replicate variance for a given simulation scenario. Larger F statistics correspond to a greater variance among replicates relative to variation within replicates. We also report the R^2, the proportion of variance explained by replicate.

¹⁵⁴ We first focused on the variability of individual level parameters by calculating ANOVA F statistics based
¹⁵⁵ on q, Q_{12}, and the number of transitions (junctions) in ancestry along chromosomes. Next, we asked to
¹⁵⁶ what extent specific genetic loci respond similarly across replicates. To do this, we calculated ANOVA F
¹⁵⁷ statistics based on the genotypes of three specific loci: a locus directly under selection (locus 1.4); a locus
¹⁵⁸ on the same chromosome as selected loci (locus 1.10); and a physically unlinked locus (3.4). In addition
¹⁵⁹ to ANOVA on genotypes, we performed ANOVA on minor ancestry. We set the response variable as either
¹⁶⁰ the locus specific genotype (0, 0.5 or 1), if mean q for the replicate was < 0.5, or, if $q > 0.5$ then we had
¹⁶¹ the response variable as 1-genotype. This ensures that a minor ancestry score of 1 meant an individual had
¹⁶² two copies of the rarer ancestry at the replicate level. As a complementary approach to assess genotypic
¹⁶³ variability among individuals and replicates, we used principal component analysis (PCA) in R based on raw
¹⁶⁴ genotypes from all 510 loci. For each replicate, we calculated a 90% confidence ellipse of the first two PCs
¹⁶⁵ using the *ordiellipse* function in vegan v2.6-2 (Oksanen *et al.* 2022). Finally, we examined the variability
¹⁶⁶ in the genetic composition of hybrids using using multinomial regression that reflects the probabilities of
¹⁶⁷ homozygous and heterozygous genotypes given an individual's genome-wide admixture proportion (Gompert
¹⁶⁸ & Buerkle 2010). We consider any significant effect of replicate identified by ANOVA to be evidence of
¹⁶⁹ variability, although we recognize that statistically significant differences could arise through variation that
¹⁷⁰ is not large in magnitude.

Unequal parental populations

¹⁷² Because we expected rare alleles to go to fixation quickly in small populations, we focused our simulations on
¹⁷³ scenarios with equal proportions of each parental type. However, given that many instances of anthropogenic
¹⁷⁴ hybridization are related to conservation translocations or range expansions, we also wanted to ask whether
¹⁷⁵ selection could lead to consistency when there are unequal parental population sizes. We predict that unequal
¹⁷⁶ parental populations could lead to less variation among replicates, as it is expected that more common alleles
¹⁷⁷ go to fixation according to their frequency (Kimura 1962). We simulated 3 deme chains, as above, for m=0.01
¹⁷⁸ or m=0.2, c=0.9, and a BDMI. However, instead of 150 individuals per deme, we varied the population sizes
¹⁷⁹ to manipulate the parental population sizes, giving ratios of 100:1, 10:1, 5:1, 2:1 and 1:1 (Table S4). we then
¹⁸⁰ followed the same protocol as described above to estimate the F statistics and R^2 to determine to what extent
¹⁸¹ the proportion of locus-specific variation explained by replicate was different for each of the three focal loci
¹⁸² (1.4, 1.10 and 3.4, as described above).

Results

¹⁸⁴ We examined 12 scenarios that varied with respect to the degree of gene flow ($m = 0.01$ versus 0.2), strength
¹⁸⁵ of selection ($c = 0$ versus 0.2 versus 0.9), and genetic architecture (BDMI versus pathway). In general,
¹⁸⁶ proportionally more variation in any of the response variables (q, Q_{12}, or number of junctions) was associated
¹⁸⁷ with differences among replicates (i.e. had larger ANOVA F statistics) when selection was high, gene flow
¹⁸⁸ was low, and when incompatibilities were due to a BDMI (Fig. 2). At generation 10, there was significant
¹⁸⁹ variation explained by replicate in number of junctions in all 12 scenarios (F range: 3.26–146, R^2 range:
¹⁹⁰ 0.020–0.482), in admixture proportion (q) for all 12 scenarios (F range: 2.16–1148, R^2 range: 0.014–0.880),
¹⁹¹ and in inter-specific ancestry (Q_{12}) for 12 scenarios (F range: 0.72–153, R^2 range: 0.005–0.494) (Table S1).
¹⁹² For the three deme structure, we also found substantial among replicate variation in q when selection was
¹⁹³ high and migration was low (Fig. S2), but the high selection, high migration scenario was more consistent
¹⁹⁴ than in the 11 deme structure. At generation 100, replicates differed in the number of junctions for all 12
¹⁹⁵ scenarios (F range: 17.76–1183, R^2 range: 0.079–0.883), in q for all 12 scenarios (F range: 7.24–3094, R^2
¹⁹⁶ range: 0.044–0.952), and in Q_{12} for 11 scenarios (F range: 1.07–433, R^2 range: 0.007–0.734) (Table S1).
¹⁹⁷ For those scenarios with the largest F statistics for q and Q_{12}, extreme variation was not isolated to the
¹⁹⁸ chromosome with incompatible loci, but instead extended throughout the genome (Fig. 3; note that Q_{12} was
¹⁹⁹ less variable in the three deme structure, Fig. S3). Accordingly, in these scenarios the genetic composition of
²⁰⁰ hybrids varied greatly across replicates for key characteristics such as the proportion of F2 hybrids and the
²⁰¹ direction of backcrossing (Fig. 4), although there was less variation among replicates in the three deme case
²⁰² (Fig. S4). Across scenarios, log-transformed F statistics for q and Q_{12} were highly correlated with one another
²⁰³ in generation 10 (r = 0.874; Fig. S5) and generation 100 (r = 0.933; Fig. S6). Further, a replicate's median

q, and to a lesser extent Q_{12}, in generation 10 were often correlated with its median value in generation 100, especially when selection was strong ($c = 0.9$) (Figs. S7, S8).

Each of the three focal loci (1.4: locus under selection; 1.10: locus physically linked to causal locus; 3.4: physically unlinked locus) had different proportions of variation explained by replicate, depending on the simulation scenario. Replicate was a significant predictor of variation in all scenarios for each of the 3 loci (Table S2). The ratio of variance explained by replicate in locus 1.4 ranged from F = 2.7 to 233, where there was little variation explained due to replicate in the BDMI scenario with high migration and moderate selection ($m = 0.2$, $c = 0.2$, F = 2.7, P < 0.001, $R^2 = 0.017$). In contrast, in the BDMI scenario with low migration and high selection, there was huge proportion of variation in genotype explained by replicates (F = 233.1, P < 0.001, $R^2 = 0.598$). Similar patterns across scenarios were found for the other two loci, suggesting that the patterns extended across the genome. We found that using minor ancestry instead of genotype recovered a similar pattern, where replicate explained substantial variation in nearly all scenarios (Table S3). The PCA based on the genotypes from all 510 loci failed to differentiate the replicates from generation 10, except for the scenarios involving a BDMI with $m = 0.01$ and $c = 0.9$ (Fig. S9). At generation 100, however, PCA was able to distinguish among replicates for all low gene flow ($m = 0.01$) scenarios, but replicates were qualitatively identical for all high gene flow ($m = 0.2$) scenarios (Fig. S10). Finally, there was substantial variation in the multinomial regressions of ancestry each of the three focal loci in both the 11 deme scenario (Figs. 5, S11, S12, S13) and the three deme scenario (Fig. S14).

We found that differences in parental population size affected the proportion of variation explained by replicate, where replicate hybrid zones tend to be more consistent when parental populations are unequal (Fig. 6). This was true for both low (m=0.01) and high (m=0.2) migration scenarios, although there was generally more consistency in high migration scenarios. Comparing locus 1.4, 1.10 and 3.4, it was clear that while consistency changed with the ratio between parental populations, the changes were similar regardless of locus-specific selection. So while consistency changed depending on the ratio between parental populations, this was not consistency due to selection.

Discussion

For many of the evolutionary scenarios we modeled, there was remarkable variation in the outcomes of hybridization across replicates (Fig. 2) despite their identical evolutionary and demographic conditions. Notably, both genome-wide ancestry composition of individuals and genotypes at specific loci varied extensively across replicates. Strong selection on the same loci led to increased genotypic variation among replicates, in contrast to naïve expectations that selection would lead to more consistent outcomes. Moreover, this result extended across the entire genome with pronounced intra- and inter-chromosomal linkage disequilibrium (Fig. 3). Even loci that were not on the same chromosome as the causal loci for incompatibilities were affected by the strong viability selection. This lack of a direct causal link between selection strength and outcomes of hybridization suggests caution when tying patterns of consistency to evolutionary processes in replicate hybrid zones. Similar outcomes of hybridization may occur in the presence or absence of strong selection. Indeed, we found the least variability explained by among replicate differences when selection was weak, suggesting that consistency could indicate a lack of reproductive isolation. Variability in outcomes could arise from many causes. Our results highlight a prominent role for stochasticity in the composition of hybrid zones (Blount et al. 2008), particularly when selection is strong, meaning that chance events in early generations of hybridization lead to evolution proceeding in dramatically different directions (Gould 1989). Below, we discuss these results in the context of both speciation research and conservation genetics.

Replicate hybrid zones in speciation genetics

When the geographic ranges of species collide in secondary contact, outcomes of hybridization have been expected to be similar when mechanisms of reproductive isolation are the same. Further, previous work has shown that selection on individual loci can lead these loci to deviate in ancestry from neutral loci elsewhere in the genome (Gompert & Buerkle 2011), though attributing deviation at exceptional loci to drift or selection is fraught (Gompert & Buerkle 2011, McFarlane et al. 2021), as many processes can produce this pattern. These assumptions underlie the basic premise of replicate hybrid zone research: similarities across replicates should be indicative of shared isolating barriers, independent of the effects of genetic drift at each individual

population, thereby giving insight into the nature of reproductive isolation between a pair of species (Langdon et al. 2022, Matute et al. 2020, Rieseberg et al. 1999, Simon et al. 2020, 2021, Westram et al. 2021). Despite this expectation, many of our simulations highlight remarkable variation in individual-level parameters (Figs. 2, 4), genotypes (Fig. 3), and introgression (Fig. 5) across replicates. For example, more than 80% of the variation in q is explained by among replicate variation when migration is rare and selection against BDMIs is strong (Table S1). When reproductive isolation is strong, one might anticipate consistency, where strong selection might shore up reproductive isolation. But if there are variable outcomes, then reproductive isolation could collapse in some places but not others, under identical conditions. These unstable equilibria resulting from frequency dependent selection and underdominance (in the context of multi-locus epistasis) in our simulations suggest that a substantial amount of stochasticity could result when reproductive isolation is tested in secondary contact.

In contrast to the idea that strong selection could lead to consistency across replicates, we found that BDMIs combined with strong selection lead to remarkable variation among replicates. Since underdominance created by BDMIs can lead to alternative homozygous genotypic states with full fitness, alternative alleles can rise to high frequency in our different simulation replicates (Fig. 1). This was reflected by a bimodal distribution of q scores among replicates particularly when selection on BDMIs was high (Fig. 2). We found that the stronger the barriers to gene flow at specific loci, the more variable were the outcomes under identical conditions, and the more variation was explained by among replicate differences, and thus, the more unstable the allele frequencies are across replicates. Further, any drift will lead to alternative homozygotes being favoured due to frequency dependent selection (Ayala & Campbell 1974, Heino et al. 1998, Gompert et al. 2012). This means that small perturbations to identical systems due to stochasticity can result in drastically different outcomes (Fig. 1). The unstable equilibrium dynamics predominate in the first few generations after secondary contact, but eventually steady migration will bring hybrid zones back to migration-drift balance. In the meantime, selection for either parental homozygote has set the trajectory of the system for later generations. This is why late generations were well predicted by early generations (Figs. S7, S8). This is consistent with previous work that has demonstrated the sensitivity of evolutionary trajectories to starting conditions (Simões et al. 2008, Losos 2011) and the historical sequence of events (i.e. historicity; Lewontin 1966); while the replicates in our simulations have identical conditions, stochasticity and drift lead to differences in early generations that are perpetuated.

One noteworthy difference between our simulations and some empirical studies of replicate hybrid zones is that many of our simulations had equal numbers of individuals between admixing populations. Since we have the same N_e for each parental population, this could lead to the bimodality of high frequency alleles in the BDMIs between replicates; either pair of alleles is equally fit, and equally likely, so which ever becomes more frequent due to drift is favored by selection (Fig. 1). We chose to model equal starting numbers of individuals from admixing populations because the outcomes of hybridization (i.e. which pair of alleles go to fixation) should be more easily be predicted based on starting allele frequencies when there is asymmetry (Kimura 1962). Thus, there could be increased consistency, but it would not be because of selection. We tested whether changing the starting ratios of the parental populations affected consistency among replicates, and we found that there was generally more consistency when there was more asymmetry (Fig. 6). However, we have also shown that this increase in consistency at particular loci was independent of selection, and was likely due to drift. If selection led to more consistency across replicates than drift did, which would allow consistency to be used to infer selection, this would be most likely be seen in the 1:1 scenarios. In empirical studies with asymmetry, we expect that observed consistency would likely be due to drift, not selection, and that in these cases it would be even more difficult to point to selection as the underlying mechanism.

Our conception of speciation typically includes a set of phenotypes and underlying genotypes that contribute to reproductive isolation between diverged taxa. However, the results of our simulation study suggest that if taxa hybridize and make fertile and viable progeny in multiple instances of secondary contact, a single, consistent set of loci contributing to reproductive isolation would lead to variable evolutionary dynamics of hybrids and gene flow. That variability will make it more challenging to map the genes associated with reproductive isolation, even if there is a single, consistent genetic architecture. Indeed, we have previously noted that genetic drift can lead to patterns in the genome that can mimic the predicted outcomes of selection in secondary contact (Gompert & Buerkle 2011, McFarlane et al. 2021). Beyond the challenge of mapping pattern to process, the variation in the effectiveness of reproductive isolation even under (unrealistically) constant conditions suggests that the nature of isolation and gene flow between species is more complex than

we have typically appreciated, or than would be suggested by a simple interpretation of 'speciation genetics'. For these reasons, we suggest that understanding the variation in effectiveness of incompatibilities, where secondary contact occurs in more than one location (or time point), is an interesting avenue for future research. Additionally, coupling between barrier effects, including loci involved in pre- or postzygotic isolation, has been predicted to increase consistency across clines within a hybrid zone (Barton 1983, Butlin & Smadja 2018). While we have not considered coupling directly here, we predict that this strengthening of reproductive isolation could increase variability across replicate hybrid zones, as the potential for stochasticity acting on multiple traits would likely increase the variability, not decrease it. While coupling would likely take more than the 10 generations we have focused on here to occur, it would be an interesting question to determine the extent to which coupling affected consistency across replicate hybrid zones.

Implications for conservation research

In the realm of conservation, where replicate hybrid zones are studied because of the potential negative consequences of hybridization between endemic and introduced species (Allendorf *et al.* 2001, Mandeville *et al.* 2017, Todesco *et al.* 2016, Wolf *et al.* 2001), our findings are somewhat challenging. Given finite budgets for conservation, the ability to generalize from one setting with potentially hybridizing individuals to other unsampled settings (i.e. out-of-sample prediction) would be extremely useful. However, except in specific circumstances (e.g. when selection against hybrids is known to be low), we anticipate substantial variation among hybrid zones even under identical demographic and evolutionary processes. This means that to assess the likely extent of hybridization in each population, for example to protect individuals in those settings with the least hybridization, each locality might need to be sampled. Since this pattern is driven by genetic drift, the variation among localities will be even higher with small effective population sizes, as would be expected if one species has been introduced to the range of another. The predictability of evolution is negatively (but not linearly) related to effective population size (Szendro *et al.* 2013), so when considering small, introduced or endangered populations, practitioners must assume that drift and the corresponding stochasticity in the composition of populations are even stronger. One optimistic point worth highlighting is the consistency found across time within replicate hybrid zones. If conditions do not change over time, the distribution of admixture proportions in generation 100 is highly correlated with the distribution in generation 10 (Figs. S7, S8). Practitioners will need to weigh the need for temporal replication to 'check in' relative to the need for spatial replication, based on our finding that every new hybrid zone could have a different distribution than those already sampled. However, one caveat is that this temporal stability is based on a system with constant conditions, which is fairly unrealistic.

Caveats

One difference between our study and some empirical studies is the use of minor ancestry at haplotypes, rather than admixture proportion, to determine to what extent there are regions of the genome with depleted variation, suggesting incompatibilities (e.g. Langdon *et al.* 2022, Schumer *et al.* 2018). While we explored this for specific loci (Table S3), we are conscious that polarizing by minor parent assumes that backcrossing in either direction has the same consequences, regardless of the ancestry of the genetic background. Given that most of our simulations assumed the same parental population size and equal contributions to admixture (i.e. there was no asymmetry), the population determined to be the minor ancestor varied among replicates (Fig. 1). When we compared minor ancestry at specific loci, we found a very low correlation between (arbitrarily chosen) replicate 1 and 2 and significant variation across replicates in nearly all scenarios (Table S3), suggesting that using minor ancestry rather than genotype does not substantially decrease the variation between replicates.

A second key difference between our simulations and the expectations in empirical systems is the heterogeneous recombination landscape that has been demonstrated to affect speciation (Nachman & Payseur 2012). Our simulations have a homogenous recombination landscape based on a Poisson process (Lindtke & Buerkle 2015). This homogenous recombination means that recombination hotspots and cool spots, which could lead to purging of even beneficial alleles in empirical systems (Moran *et al.* 2021), have not been simulated here. This also means that we would not have found a correlation between recombination and minor ancestry as in (Langdon *et al.* 2022), although we also did not find a correlation of minor ancestry between replicate

populations. Future work could simulate such recombination landscapes and ask how this affects variation in outcomes across replicate hybrid zones.

A third caveat is that we have not simulated adaptive introgression, where an allele from one population introgresses through the hybrid zone into the other population due to directional selection. We expect that some of the same problems we have demonstrated will apply in that circumstance as well, even if underdominance does not drive variation as we expect it does for BDMIs. Some empirical studies have found evidence of adaptive introgression from introduced species, using replicate hybrid zones (e.g., Bay *et al.* 2019, Calfee *et al.* 2020), but few have explicitly ruled out genetic drift as a mechanism, and selection is regularly implicated even when hybrid zones do not show concordant patterns (but see Nouhaud *et al.* 2022). Given that adaptive introgression is only expected to be possible when recombination is quite high, allowing a beneficial allele to be separated from a neutral or deleterious haplotype (Moran *et al.* 2021, Uecker *et al.* 2015), the *a priori* expectation must be that introgression is commonly due to non-adaptive forces (*sensu* Barrett & Hoekstra 2011).

Finally, more generally, our results are illustrative of variability that arises in a particular set of circumstances and are not meant to be a perfect reflection of the most likely population structure and fitnesses in secondary contact for a pair of taxa in nature. While we have perturbed these simulations to examine robustness, such as considering both an 11 deme and 3 deme population structure (Lindtke & Buerkle 2015; Figs. S2, S3), simulations can never cover all possible biological variability. The expectations for any natural setting should be informed by knowledge of the system, which will include an appreciation of the complexities due to chance in finite populations, unstable equilibrium allele frequencies (and ancestries), and the dynamics of gene flow between populations with potentially unequal and variable size through time.

Strategies for analyses of replicate hybrid zones

Even when there is substantial variability across hybrid zones, there is still the opportunity to capitalize on replication when available. This might be particularly fruitful when the genetic architecture of incompatibilities is expected to be polygenic or selection is less strong, as such cases have less variation explained by among replicate differences (Table S1). Given that polygenic architectures are likely responsible for many cases of reproductive isolation (Barton 2022), we are encouraged by our results. By accounting statistically for this variability across replicates (rather than assuming all replicates are comparable), one can examine both global and local patterns. For example, some studies have used replicate hybrid zones to ask how ecological variation affects distributions of admixture (e.g. Chatfield *et al.* 2010, Culumber *et al.* 2012, Mandeville *et al.* 2019, 2021, Velo-Antón *et al.* 2021), a question that assumes variation across replicates. Here, we have used ANOVA to estimate variance due to replicate, but we have not incorporated other variables (e.g. environmental variability) into our analyses. ANOVAs are excellent for our balanced simulations, but might not be the appropriate method for studies that want to account for variability among replicates and also ask how some other variable (e.g. human activity, temporal variation) affects hybridization outcomes. Some studies of replicate hybrid zones have used PCA or similar approaches to assess to what extent there were substantial differences among replicates, and then using pairwise correlations between studies, to determine to what extent there was concurrence between admixture distributions, allele frequencies, or cline parameters (e.g. Westram *et al.* 2021). We found that PCAs did not differentiate among replicates well in the 10th generation after admixture (Fig. S9), suggesting that PCAs are not sufficient to confirm similarity among replicates.

One could build more sophisticated hierarchical models beyond ANOVA, but the usual rules about sample size for hierarchical models will apply. For example, Calfee *et al.* (2020) fit a classic logistic cline model to inferred genome-wide individual ancestry proportions, which fit a cline specific to two replicate instances of the escape of *Apis mellifera scutellata* and subsequent hybridization with *Apis mellifera*. They were thus able to make inferences about the similarities and differences of these two hybrid zones, while accounting for between replicate variation. By using a hierarchical model, which allows for differences among populations in genome wide distributions, but assumes that these distributions are all pulled from a global population, there is an opportunity to compare clines at the same locus across populations, without assuming that the populations are perfectly comparable. This means that different taxa and systems will be more fruitful for understanding causes of different outcomes of hybridization, such as multiply introduced species like trout (Mandeville *et al.* 2019), mussels (Simon *et al.* 2020), and organisms that are amenable to laboratory settings

where there can be substantial replication (e.g. *Saccharomyces* sp., Dunn *et al.* 2013; *Tigriopus californicus*, Pritchard & Edmands 2013). There are opportunities to use replicate hybrid zones to shed light on both local and global patterns of hybridization and introgression, while being cautious to not assume that replicates will be predictive of other replicates or unsampled sites.

One challenge when studying hybridization across spatial replicates is to think about how independent replicate sites are. At one end of the spectrum, 'replicates' may not be independent at all for study systems that have high levels of gene flow over the spatial scale of sampling. In contrast, some hybrid zones may be truly independent, such as instances of secondary contact isolated from one another in a mosaic hybrid zone or by a substantial distance. For example, bighorn sheep have been reintroduced to several isolated mountain ranges across North America using founders from multiple source ancestries (Jahner *et al.* 2019). More often, studies may involve multiple transects of secondary contact between diverged lineages, as in studies of house mice in Europe (Bímová *et al.* 2011, Payseur *et al.* 2004, Teeter *et al.* 2010) or *Gryllus* crickets in North America (Larson *et al.* 2014). Replicated hybrid zones may also have complex spatial structure, and replicates may be hierarchically nested, as is the case with replicating across fish hybrid zones in small streams within a larger river basin, and then also replicating across basins (e.g. *Catostomus* sucker hybridization or rainbow × cutthroat trout hybridization; Mandeville *et al.* 2017, Muhlfeld *et al.* 2017). Geographic distribution of hybridizing species, as well as the dispersal mode and capacity of study organisms, are likely to be quite important in defining independence of replicate sites, and may influence decisions about statistical treatment of comparisons. We emphasize that studies of replicate hybrid zones at all levels of independence are likely to be informative, but clearly defining the independence of replicates is an important goal.

Conclusion

This research was motivated by our own ambitions to use replicate hybrid zones to ask basic and applied questions in our study systems. Indeed, we have written about the promise of replicate hybrid zones in our own papers (Buerkle & Lexer 2008, Jahner *et al.* 2021, Lindtke *et al.* 2012, Mandeville *et al.* 2017, McFarlane *et al.* 2021), so we were fairly confident that we could link the outcomes of hybridization to the evolutionary processes that generated them, at least for some scenarios. As it turns out, early generation stochasticity can dramatically alter the outcomes of hybridization across hybrid zones, leading to extreme variability across replicates with identical starting conditions. Moreover, the degree of variability is amplified with increasing strength of selection on incompatibilities, suggesting that the 'simplest scenarios' will be the most difficult in which to tie pattern to process. Despite these results, we believe the future of replicate hybrid zone research remains promising, particularly if potential reasons for variation among hybrid zones are considered appropriately. We hope this study will encourage 1) the development of new statistical models that explicitly account for variability across hybrid zones and use this variability to better understand evolutionary and demographic causes of reproductive isolation; 2) future modeling exercises that examine the countless other important axes of variability that were not explored here; and 3) new empirical studies of replicate hybrid zones that acknowledge the potential lack of identifiability of evolutionary processes based on our observed outcomes of secondary contact.

Acknowledgements

We are grateful to Cassandre Pyne, Amanda Meuser, Amy Pitura, Teaghan Frauley, Chris Nice, and Zach Gompert for previous discussions of this manuscript. We are also thankful for the comments of three thoughtful reviewers. This research was supported by the Modelscape Consortium with funding from NSF (OIA-2019528). Analyses were supported by the University of Wyoming's Advanced Research Computing Center and its Beartooth Computing Environment, Intel x86_64 cluster (https://doi.org/10.15786/M2FY47).

Data Accessibility Statement

The original source code for the simulation software `dfuse` can be found at the Dryad Digital Repository associated with Lindtke & Buerkle (2015) (doi: 10.5061/dryad.0506g). All code and input files for analyses

can be found at on Zenado https://zenodo.org/account/settings/github/repository/erynmcfarlane/predict_hybrids, (McFarlane *et al.* 2022).

Benefit-Sharing Statement

Benefits from this research accrue from the sharing of our data and results on public databases as described above.

Author contributions

SEM and EGM had the initial idea to think carefully about the utility of replicate hybrid zones in evolution and conservation. DL and SEM ran all of the simulations. SEM, JPJ, and CAB went down a deep, statistical rabbit hole before ultimately deciding to use ANOVA to analyze everything. SEM wrote the initial draft of the manuscript, and all authors contributed to subsequent revisions.

References

Allendorf F, Leary R, Spruell P, Wenburg J (2001) The problems with hybrids: setting conservation guidelines. *Trends in Ecology & Evolution*, **16**, 613–622.

Ayala FJ, Campbell CA (1974) Frequency-dependent selection. *Annual review of Ecology and systematics*, pp. 115–138.

Barrett RD, Hoekstra HE (2011) Molecular spandrels: tests of adaptation at the genetic level. *Nature Reviews Genetics*, **12**, 767–780.

Barton NH (1979) The dynamics of hybrid zones. *Heredity*, **43**, 341–359.

Barton NH (1983) Multilocus clines. *Evolution*, **37**, 454–471.

Barton NH (2022) The "new synthesis". *Proceedings of the National Academy of Sciences*, **119**, e2122147119.

Barton NH, De Cara MAR (2009) The evolution of strong reproductive isolation. *Evolution*, **63**, 1171–1190.

Barton NH, Gale KS (1993) Genetic analysis of hybrid zones. In: *Hybrid Zones and the Evolutionary Process* (ed. Harrison RG), pp. 13–45, Oxford University Press, New York.

Barton NH, Hewitt GM (1985) Analysis of hybrid zones. *Annual Review of Ecology and Systematics*, **16**, 113–148.

Bateson W (1909) Heredity and variation in modern lights. In: *Darwin and modern science* (ed. Seward AC), pp. 85–101, Cambridge University Press, Cambridge, UK.

Bay RA, Taylor EB, Schluter D (2019) Parallel introgression and selection on introduced alleles in a native species. *Molecular Ecology*, **28**, 2802–2813.

Bímová BV, Macholán M, Baird SJ, *et al.* (2011) Reinforcement selection acting on the european house mouse hybrid zone. *Molecular Ecology*, **20**, 2403–2424.

Blount ZD, Borland CZ, Lenski RE (2008) Historical contingency and the evolution of a key innovation in an experimental population of *Escherichia coli*. *Proceedings of the National Academy of Sciences*, **105**, 7899–7906.

Buerkle CA, Lexer C (2008) Admixture as the basis for genetic mapping. *Trends in Ecology & Evolution*, **23**, 686–694.

Buerkle CA, Rieseberg LH (2001) Low intraspecific variation for genomic isolation between hybridizing sunflower species. *Evolution*, **55**, 684–691.

Butlin RK, Smadja CM (2018) Coupling, reinforcement, and speciation. *The American Naturalist*, **191**, 155–172.

Caballero-Villalobos L, Silva-Arias GA, Turchetto C, *et al.* (2021) Neutral and adaptive genomic variation in hybrid zones of two ecologically diverged petunia species (solanaceae). *Botanical Journal of the Linnean Society*, **196**, 100–122.

Calfee E, Agra MN, Palacio MA, Ramírez SR, Coop G (2020) Selection and hybridization shaped the rapid spread of african honey bee ancestry in the americas. *PLoS genetics*, **16**, e1009038.

Chatfield MWH, Kozak K, Fitzpatrick B, Tucker P (2010) Patterns of differential introgression in a salamander hybrid zone: inferences from genetic data and ecological niche modelling. *Molecular Ecology*, **19**, 4265–4282.

Coughlan JM, Matute DR (2020) The importance of intrinsic postzygotic barriers throughout the speciation process. *Philosophical Transactions of the Royal Society B*, **375**, 20190533.

Coyne JA, Orr HA (2004) *Speciation*. Sinauer Associates, Sunderland, Massachusetts.

Culumber Z, Shepard D, Coleman S, Rosenthal G, Tobler M (2012) Physiological adaptation along environmental gradients and replicated hybrid zone structure in swordtails (teleostei: Xiphophorus). *Journal of evolutionary biology*, **25**, 1800–1814.

Dobzhansky T (1936) Studies on hybrid sterility. ii. localization of sterility factors in *Drosophila pseudoobscura* hybrids. *Genetics*, **21**, 113–135.

Dobzhansky TH (1937) *Genetics and the Origin of Species*. Columbia University Press, New York.

Dunn B, Paulish T, Stanbery A, *et al.* (2013) Recurrent rearrangement during adaptive evolution in an interspecific yeast hybrid suggests a model for rapid introgression. *PLoS genetics*, **9**, e1003366.

Elmer KR, Fan S, Kusche H, *et al.* (2014) Parallel evolution of Nicaraguan crater lake cichlid fishes via non-parallel routes. *Nature Communications*, **5**, 1–8.

Endler JA (1977) *Geographic Variation, Speciation, and Clines*. Princeton University Press, Princeton, NJ.

Ferris KG, Chavez AS, Suzuki TA, *et al.* (2021) The genomics of rapid climatic adaptation and parallel evolution in North American house mice. *PLoS Genetics*, **17**, e1009495.

Fishman L, Sweigart AL (2018) When two rights make a wrong: the evolutionary genetics of plant hybrid incompatibilities. *Annual Review of Plant Biology*, **69**, 707–731.

Gagnaire PA, Pavey SA, Normandeau E, Bernatchez L (2013) The genetic architecture of reproductive isolation during speciation-with-gene-flow in lake whitefish species pairs assessed by rad sequencing. *Evolution*, **67**, 2483–2497.

Gompert Z, Buerkle CA (2010) *introgress*: a software package for mapping components of isolation in hybrids. *Molecular Ecology Resources*, **10**, 378–384.

Gompert Z, Buerkle CA (2011) Bayesian estimation of genomic clines. *Molecular Ecology*, **20**, 2111–2127.

Gompert Z, Buerkle CA (2012) bgc: Software for Bayesian estimation of genomic clines. *Molecular Ecology Resources*, **12**, 1168–1176.

Gompert Z, Lucas LK, Buerkle CA, Forister ML, Fordyce JA, Nice CC (2014) Admixture and the organization of genetic diversity in a butterfly species complex revealed through common and rare genetic variants. *Molecular Ecology*, **23**, 4555–4573.

Gompert Z, Parchman TL, Buerkle CA (2012) Genomics of isolation in hybrids. *Philosophical Transactions of the Royal Society B: Biological Sciences*, **367**, 439–450.

Gould SJ (1989) *Wonderful Life: The Burgess Shale and the Nature of History*. W. W. Norton & Company, Inc., New York, NY, USA.

Harrison RG (1990) Hybrid zones: windows on evolutionary process. *Oxford Surveys in Evolutionary Biology*, **7**, 69–128.

Harrison RG, Larson EL (2016) Heterogeneous genome divergence, differential introgression, and the origin and structure of hybrid zones. *Molecular ecology*, **25**, 2454–2466.

Heino M, Metz JA, Kaitala V (1998) The enigma of frequency-dependent selection. *Trends in Ecology & Evolution*, **13**, 367–370.

Jahner JP, Matocq MD, Malaney JL, *et al.* (2019) The genetic legacy of 50 years of desert bighorn sheep translocations. *Evolutionary Applications*, **12**, 198–213.

Jahner JP, Parchman TL, Matocq MD (2021) Multigenerational backcrossing and introgression between two woodrat species at an abrupt ecological transition. *Molecular Ecology*, **30**, 4245–4258.

Jiggins CD, Mallet J (2000) Bimodal hybrid zones and speciation. *Trends in Ecology & Evolution*, **15**, 250–255.

Kimura M (1962) On the probability of fixation of mutant genes in a population. *Genetics*, **47**, 713–719.

Kingston SE, Parchman TL, Gompert Z, Buerkle C, Braun MJ (2017) Heterogeneity and concordance in locus-specific differentiation and introgression between species of towhees. *Journal of evolutionary biology*, **30**, 474–485.

Landry CR, Hartl DL, Ranz JM (2007) Genome clashes in hybrids: insights from gene expression. *Heredity*, **99**, 483–493.

Langdon QK, Powell DL, Kim B, *et al.* (2022) Predictability and parallelism in the contemporary evolution of hybrid genomes. *PLoS Genetics*, **18**, e1009914.

Larson EL, Vanderpool D, Sarver BA, *et al.* (2018) The evolution of polymorphic hybrid incompatibilities in house mice. *Genetics*, **209**, 845–859.

Larson EL, White TA, Ross CL, Harrison RG (2014) Gene flow and the maintenance of species boundaries. *Molecular ecology*, **23**, 1668–1678.

Lewontin RC (1966) Is nature probable or capricious? *BioScience*, **16**, 25–27.

Lindtke D, Buerkle CA (2015) The genetic architecture of hybrid incompatibilities and their effect on barriers to introgression in secondary contact. *Evolution*, **69**, 1987–2004.

Lindtke D, Buerkle CA, Barbará T, *et al.* (2012) Recombinant hybrids retain heterozygosity at many loci: new insights into the genomics of reproductive isolation in *Populus*. *Molecular Ecology*, **21**, 5042–5058.

Lindtke D, Gompert Z, Lexer C, Buerkle CA (2014) Unexpected ancestry of *Populus* seedlings from a hybrid zone implies a large role for postzygotic selection in the maintenance of species. *Molecular ecology*, **23**, 4316–4330.

Losos JB (2011) Convergence, adaptation, and constraint. *Evolution: International Journal of Organic Evolution*, **65**, 1827–1840.

Maheshwari S, Barbash DA (2011) The genetics of hybrid incompatibilities. *Annual Review of Genetics*, **45**, 331–355.

Mandeville EG, Hall RO, Buerkle CA (2021) Ecological outcomes of hybridization vary extensively in catostomus fishes. *bioRxiv*.

Mandeville EG, Parchman TL, Thompson KG, *et al.* (2017) Inconsistent reproductive isolation revealed by interactions between *Catostomus* fish species. *Evolution Letters*, **1**, 255–268.

Mandeville EG, Walters AW, Nordberg BJ, Higgins KH, Burckhardt JC, Wagner CE (2019) Variable hybridization outcomes in trout are predicted by historical fish stocking and environmental context. *Molecular ecology*, **28**, 3738–3755.

Matute DR, Comeault AA, Earley E, *et al.* (2020) Rapid and predictable evolution of admixed populations between two drosophila species pairs. *Genetics*, **214**, 211–230.

McFarlane SE, Jahner JP, Lindtke D, Buerkle CA, Mandeville EG (2022) Dataset for 'selection leads to remarkable variability in the outcomes of hybridization across replicate hybrid zones'. Data available at Zenado, https://zenodo.org/account/settings/github/repository/erynmcfarlane/predict_hybrids.

McFarlane SE, Senn HV, Smith SL, Pemberton JM (2021) Locus-specific introgression in young hybrid swarms: drift may dominate selection. *Molecular Ecology*, **30**, 2104–2115.

Moran BM, Payne C, Langdon Q, Powell DL, Brandvain Y, Schumer M (2021) The genomic consequences of hybridization. *Elife*, **10**, e69016.

Muhlfeld CC, Kovach RP, Al-Chokhachy R, *et al.* (2017) Legacy introductions and climatic variation explain spatiotemporal patterns of invasive hybridization in a native trout. *Global Change Biology*, **23**, 4663–4674.

Muller HJ (1942) Isolating mechanisms, evolution, and temperature. *Biological Symposia*, **6**, 71–125.

Nachman MW, Payseur BA (2012) Recombination rate variation and speciation: theoretical predictions and empirical results from rabbits and mice. *Philosophical Transactions of the Royal Society B: Biological Sciences*, **367**, 409–421.

Nouhaud P, Martin SH, Portinha B, Sousa VC, Kulmuni J (2022) Rapid and predictable genome evolution across three hybrid ant populations. *Plos Biology*, **20**, e3001914.

Oksanen J, Simpson GL, Blanchet FG, *et al.* (2022) *vegan: Community Ecology Package. R package version 2.6-2*. https://CRAN.R-project.org/package=vegan.

Orr H (1995) The population-genetics of speciation - the evolution of hybrid incompatibilities. *Genetics*, **139**, 1805–1813.

Pardo-Diaz C, Salazar C, Baxter SW, *et al.* (2012) Adaptive introgression across species boundaries in heliconius butterflies. *PLoS genetics*, **8**, e1002752.

Payseur BA, Krenz JG, Nachman MW (2004) Differential patterns of introgression across the X chromosome in a hybrid zone between two species of house mice. *Evolution*, **58**, 2064–2078.

Powell DL, García-Olazábal M, Keegan M, *et al.* (2020) Natural hybridization reveals incompatible alleles that cause melanoma in swordtail fish. *Science*, **368**, 731–736.

Pritchard VL, Edmands S (2013) The genomic trajectory of hybrid swarms: outcomes of repeated crosses between populations of tigriopus californicus. *Evolution: International Journal of Organic Evolution*, **67**, 774–791.

R Core Team (2021) *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria.

Rand DM, Harrison RG (1989) Ecological genetics of a mosaic hybrid zone: mitochondrial, nuclear, and reproductive differentiation of crickets by soil type. *Evolution*, **43**, 432–449.

Rieseberg LH, Whitton J, Gardner K (1999) Hybrid zones and the genetic architecture of a barrier to gene flow between two sunflower species. *Genetics*, **152**, 713–727.

Schumer M, Powell DL, Delclós PJ, *et al.* (2017) Assortative mating and persistent reproductive isolation in hybrids. *Proceedings of the National Academy of Sciences*, **114**, 10936–10941.

Schumer M, Xu C, Powell DL, *et al.* (2018) Natural selection interacts with recombination to shape the evolution of hybrid genomes. *Science*, **360**, 656–660.

Shastry V, Adams PE, Lindtke D, *et al.* (2021) Model-based genotype and ancestry estimation for potential hybrids with mixed-ploidy. *Molecular Ecology Resources*, **21**, 1434–1451.

Simões P, Santos J, Fragata I, Mueller LD, Rose MR, Matos M (2008) How repeatable is adaptive evolution? the role of geographical origin and founder effects in laboratory adaptation. *Evolution: International Journal of Organic Evolution*, **62**, 1817–1829.

Simon A, Arbiol C, Nielsen EE, *et al.* (2020) Replicated anthropogenic hybridisations reveal parallel patterns of admixture in marine mussels. *Evolutionary Applications*, **13**, 575–599.

Simon A, Fraïsse C, El Ayari T, *et al.* (2021) How do species barriers decay? concordance and local introgression in mosaic hybrid zones of mussels. *Journal of Evolutionary Biology*, **34**, 208–223.

Szendro IG, Franke J, de Visser JAG, Krug J (2013) Predictability of evolution depends nonmonotonically on population size. *Proceedings of the National Academy of Sciences*, **110**, 571–576.

Szymura JM, Barton NH (1986) Genetic analysis of a hybrid zone between the fire-bellied toads, *Bombina bombina* and *B. variegata*, near Cracow in southern Poland. *Evolution*, **40**, 1141–1159.

Teeter KC, Thibodeau LM, Gompert Z, Buerkle CA, Nachman MW, Tucker PK (2010) The variable genomic architecture of isolation between hybridizing species of house mouse. *Evolution*, **64**, 472–485.

Tenaillon O, Barrick JE, Ribeck N, *et al.* (2016) Tempo and mode of genome evolution in a 50,000-generation experiment. *Nature*, **536**, 165–170.

Thompson KA, Peichel CL, Rennison DJ, *et al.* (2022) Analysis of ancestry heterozygosity suggests that hybrid incompatibilities in threespine stickleback are environment dependent. *PLoS biology*, **20**, e3001469.

Todesco M, Pascual MA, Owens GL, *et al.* (2016) Hybridization and extinction. *Evolutionary applications*, **9**, 892–908.

Uecker H, Setter D, Hermisson J (2015) Adaptive gene introgression after secondary contact. *Journal of mathematical biology*, **70**, 1523–1580.

Velo-Antón G, Lourenço A, Galán P, Nicieza A, Tarroso P (2021) Landscape resistance constrains hybridization across contact zones in a reproductively and morphologically polymorphic salamander. *Scientific reports*, **11**, 1–16.

Westram AM, Faria R, Johannesson K, Butlin R (2021) Using replicate hybrid zones to understand the genomic basis of adaptive divergence. *Molecular Ecology*, **30**, 3797–3814.

Whitney KD, Randell RA, Rieseberg LH (2006) Adaptive introgression of herbivore resistance traits in the weedy sunflower *Helianthus annuus*. *American Naturalist*, **167**, 794–807.

Wolf DE, Takebayashi N, Rieseberg LH (2001) Predicting the risk of extinction through hybridization. *Conservation Biology*, **15**, 1039–1053.

Xiong T, Mallet J (2022) On the impermanence of species: The collapse of genetic incompatibilities in hybridizing populations. *Evolution*, **76**, 2498–2512.

Figures

Figure 1: (A) Analyses were based on the outcomes of hybridization in either 11 deme or 3 deme simulations. Genotype fitness matrices are shown for the (B) BDMI and (C) pathway genetic architectures. Levels of selection (c) were set to either 0, 0.2, or 0.9. (D) The number of individuals from generation 10 having each genotypic combination for the two BDMI loci (1.4 and 1.6) is displayed for each of the twenty replicates from deme 6 of the BDMI, migration (m) = 0.01, selection (c) = 0.9 scenario.

Figure 2: Selection leads to remarkable variation in the outcomes of hybridization across replicate hybrid zones. The distribution of admixture proportion (q) from generation 10 is shown for 20 replicates from the 12 scenarios that varied by migration (m), selection (c) and incompatibility (BDMI versus pathway). Colors correspond to different replicates and match those in other figures. See Fig. S6 for results from generation 100.

Figure 3: Locus-specific variation in average (among individuals in a replicate) admixture proportion (q, panel A) and inter-specific ancestry (Q_{12}, panel B) for replicates (different colours), six combinations (columns) of migration ($m = 0.01$ or 0.2) and selection ($c = 0$, 0.2 or 0.9), and two types of incompatibilities (BDMI or pathways, in rows). Colors correspond to different replicates and match those in other figures. The black line is the average across all replicates, as in Lindtke & Buerkle (2015). The plots are for simulations after ten generations. Chromosome 1 has loci with BDMI or pathway incompatibilities, chromosome 2 has pathway incompatibility loci, while chromosome 3 has only neutral loci, and is representative of the rest of the genome (chromosomes 4–10).

Figure 4: Individual ancestry varies greatly across the twenty replicates from the BDMI, migration (m) = 0.01, selection (c) = 0.9 simulations (generation 10, deme 6). Paired estimates of admixture proportion (q) and inter-specific ancestry (Q_{12}) can be used to classify individuals as parentals ($q = 0$; $Q_{12} = 0$ or 1), F1 hybrids ($q = 0.5$; $Q_{12}= 1$), F2 hybrids (on average $q = 0.5$, $Q_{12} = 0.5$), or backcrosses (residing on the solid black lines) (Gompert *et al.* 2014, Lindtke *et al.* 2014, Shastry *et al.* 2021). Colors correspond to different replicates and match those in other figures.

Figure 5: Probability of each AA (homozygous; solid line) and Aa (codominant heterozygote; dashed line) for each of three simulated SNPs in the BDMI scenario where migration is low (m=0.01), and selection is high (c=0.9) after 10 generations. Colors correspond to different replicates and match those in other figures. This multinomial regression shows the probability of either a homozygous or heterozygous genotype for a given admixture proportion q.

Figure 6: F statistics and R^2 for each locus 1.4 (under selection), locus 1.10 (in linkage with causal incompatibility loci) and locus 3.4 (neutral) from 20 replicates of BDMIs with strong selection (c=0.9), both low and high gene flow across demes (m=0.01 in red and m=0.2 in blue). We varied the ratio of asymmetry between the parental populations from 100:1 to 1:1 (Table S4). We found that asymmetry affects the consistency across replicates, but that the effect of asymmetry is essentially the same across loci, suggesting that consistency is not indicative of locus specific selection, even when consistency is relatively high.